Pigs are highly susceptible to but do not transmit mink-derived highly pathogenic avian influenza virus H5N1 clade 2.3.4.4b

Taeyong Kwon[1], Jessie D. Trujillo[1], Mariano Carossino[2], Eu Lim Lyoo[1], Chester D. McDowell[1], Konner Cool[1], Franco S. Matias-Ferreyra[1], Trushar Jeevan[3], Igor Morozov[1], Natasha N. Gaudreault[1], Udeni B.R. Balasuriya[2], Richard J. Webby[3], Nikolaus Osterrieder[1,4], Juergen A. Richt[1, *]

1 Department of Diagnostic Medicine/Pathobiology, College of Veterinary Medicine, Kansas State University, Manhattan, KS, USA

2 Louisiana Animal Disease Diagnostic Laboratory and Department of Pathobiological Sciences, School of Veterinary Medicine, Louisiana State University, Baton Rouge, LA, USA

3 Department of Infectious Diseases, St. Jude Children's Research Hospital, Memphis, TN, USA

4 Institut für Virologie, Freie Universität Berlin, Berlin, Germany

 * Corresponding author: Dr. Juergen A. Richt; e-mail: jricht@ksu.edu

Abstract

Rapid evolution of highly pathogenic avian influenza viruses (HPAIVs) is driven by antigenic drift but

also by reassortment, which might result in robust replication in and transmission to mammals. Recently,

spillover of clade 2.3.4.4b HPAIV to mammals including humans, and their transmission between

mammal species has been reported. This study aimed to evaluate the pathogenicity and transmissibility of

a mink-derived clade 2.3.4.4b H5N1 HPAIV isolate from Spain in pigs. Experimental infection caused

moderate to severe interstitial pneumonia with necrotizing bronchiolitis with high titers of virus present in

the lower respiratory tract of principal-infected pigs and 100% seroconversion. Principal-infected pigs

shed limited amount of virus through the nasal and oral cavities, and importantly, there was no

transmission to contact sentinel pigs. Notably, critical mammalian-like mutations such as PB2-E627K and

HA-Q222L emerged at low frequencies in clinical samples and tissues derived from principal-infected

pigs. It is concluded that pigs are highly susceptible to infection with the mink-derived clade 2.3.4.4b

H5N1 HPAIV and provide a favorable environment for HPAI viruses to acquire mammalian-like

adaptations. This work is critical for further risk assessment on the ability of the newly emerging H5N1

clade 2.3.4.4b HPAIVs to cross the species barriers into mammals.

Introduction

Influenza A viruses (IAV) are members of the *Orthomyxoviridae*, family that encompasses single-stranded, negative-sense, segmented RNA viruses. IAVs are classified into different subtypes based on the two major surface proteins: hemagglutinin (HA) and neuraminidase (NA). Wild aquatic birds are natural hosts for most IAV types, except for H17N10 and H18N11, which were first isolated from bats and seem endemic in multiple bat species [1,2]. Highly pathogenic avian influenza (HPAI) viruses arise from the evolution of low pathogenic avian influenza (LPAI) viruses by insertion of a stretch of basic amino acids into the HA cleavage site, which enables efficient HA activation by ubiquitous cellular, furin-like proteases; they cause systemic, lethal infections in poultry [3].

Since the first detection of HPAI H5 subtype viruses with the isolation of the A/Goose/Guangdong/1/1996 (GsGD) H5N1 virus in southern China in 1996, this lineage has genetically evolved and spread worldwide primarily by migratory birds [4]. Continuous evolution of the GsGD-like H5 HPAI viruses lead to the emergence of genetically distinct clades, and reassortment with other clades of HPAI viruses or local LPAI viruses has resulted in remarkable genetic diversification through the acquisition of novel gene segments [5]. Eventually, HPAI H5N1 clade 2 viruses became dominant and their descendants have been responsible for subsequent HPAI outbreaks. One remarkable HPAI outbreak in North America happened in 2014/2015, when Asian-origin HPAI H5 clade 2.3.4.4 viruses spread to North America via the Pacific flyway (Alaska, Western Canada/USA) with migratory wild birds and underwent reassortment with local North American LPAIVs, resulting in HPAI H5 viruses with multiple NA subtypes, including H5N1, H5N2, H5N5, H5N6, and H5N8, collectively called H5Nx viruses [6]. The geographic expansion of the H5Nx HPAI viruses via wild birds and increased virus molecular evolution resulted in the emergence and dominance of H5 subtype clade 2.3.4.4b viruses which contributed to the H5N8 outbreaks in 2016/17 and 2020/21 and the H5N1 outbreaks in 2021/22 [7]. By 2021/22, H5N1 clade 2.3.4.4b infections had caused massive HPAI outbreaks in Europe with subsequent spread to North America via Iceland and the trans-Atlantic flyways [8].

The normal distribution of the receptors for avian and mammalian IAVs in the human respiratory tract

usually prevents efficient direct transmission of avian influenza viruses to humans, since the respective

avian-like receptor (sialic acid linked to galactose by an alpha-2,3 linkage) is found only on cells in the

lower respiratory tract region [9]. Nonetheless, in certain instances the human-animal interface allows avian

influenza viruses to cross species barriers and infect humans. Human infection with avian influenza

viruses occurs sporadically and results in mild to severe illness with a wide range of clinical symptoms.

Specifically, it has been reported that human infections with avian H5 and H7 subtype viruses can cause

severe illness in humans with case fatality rates of 52% and 39%, respectively [10]. Human infections with

HPAI H5N1 viruses have been attributed to direct exposure to infected poultry or infected wild birds.

Importantly, the current epidemiological data do not provide evidence for sustained human-to-human

transmission of HPAI H5 viruses [11]. Since the beginning of 2020, human infections with clade 2.3.4.4b

H5N1 have been documented in China, Spain, UK, USA, and Vietnam, and all cases were linked to close

contact with infected poultry [12]. In addition, the ability of the clade 2.3.4.4b H5N1 viruses to cross the

species barrier is clearly supported by its detection in a variety of mammalian species such as terrestrial

and marine mammals [12]. In October 2022, an outbreak of the clade 2.3.4.4b H5N1 HPAI virus in farmed

mink in Spain was reported, with clear evidence of sustained mink-to-mink transmission [13], raising

concerns about the adaptation of the clade 2.2.4.4b H5N1 virus to mammals. Therefore, in order to

determine the risk of the mink-derived clade 2.3.4.4b H5N1 virus isolate, this study evaluated its

virulence and transmissibility in pigs, which are natural hosts for IAVs and the "mixing vessel" for the

reassortment and establishment of potentially pandemic influenza viruses.

Materials and Methods

Cells and virus

Madin-Darby canine kidney (MDCK) cells were maintained in Dulbecco's Modified Eagle Medium

(DMEM; Corning, Manassas, VA, USA) supplemented with 5% fetal bovine serum (FBS; R&D systems,

Flower Branch, GA, USA) and a 1% antibiotic-antimycotic solution (Gibco, Grand Island, NY, USA).

The mink-derived clade 2.3.4.4b H5N1 isolate, A/Mink/Spain/3691-8_22VIR10586-10/2022, was kindly

provided by Francesco Bonfante and Isabella Monne from the Istituto Zooprofilattico Sperimentale delle

Venezie, Legnaro, Italy, and Monserrat Agüero and Azucena Sánchez from the Laboratorio Central

Veterinario (LCV), Ministry of Agriculture, Fisheries and Food, Madrid, Spain, via Dr. Richard Webby

from St. Jude Childrens' Research Hospital, Memphis, TN.

Pig infection experiment

Fifteen 4-week-old piglets were obtained from a high health status pig herd at Kansas State University.

The pigs were separated into two groups: a principal-infected group and a sentinel group that were housed

in separate pens. Nine pigs were challenged with a total dose of 2.2×10^7 TCID$_{50}$ in 4 mL of DMEM: 1

mL orally, 1mL intranasally, and 2 mL intratracheally (see Figure 1A). On day 2 after infection, principal-

infected and sentinel pigs were co-mingled and co-housed in a single pen. Nasal swab samples were

collected in 2 mL of DMEM containing 1X antibiotic/antimycotic solution at -1, 1 to 14, 17, and 21 days

post-challenge (DPC), oropharyngeal swabs on days -1, 1, 3, 5, 7, 10, 14, 17, and 21 DPC. Three of the

principal-infected pigs each were euthanized and necropsied at 3 and 5 DPC, and the remaining 3

principal and 6 sentinel pigs were euthanized and necropsied at 21 DPC. At necropsy, gross pathology

evaluation was performed (see below) and bronchoalveolar lavage fluid (BALF) and tissue samples were

collected for virological and histopathological evaluation.

Virus isolation

MDCK cells were used to isolate the virus from clinical samples, which were filtered through a 0.45 μm

syringe filter except for BALF. Ten-fold serial dilutions of samples were prepared and transferred on

confluent MDCK cells after washing with PBS. The plates were subjected to immunofluorescence assay

(IFA) on day 2. Briefly, the cells were fixed with ice-cold methanol and incubated with the HB65

monoclonal antibody targeting the IAV nucleoprotein [14]. After washing, the cells were incubated with

anti-mouse conjugated with Alexa Fluor 488 (Invitrogen, Carlsbad, CA, USA). Virus titers were

calculated using the Reed-Muench method [15].

RNA extraction and M gene-specific RT-qPCR

Viral RNA was extracted using a magnetic bead-based automated extraction system (Taco Mini and total

nucleic acid extraction kit, Gene Reach, USA) [16]. Briefly, the swabs, BALF samples and 10% tissue

homogenates were mixed with an equal volume of the RLT buffer (Qiagen, Germantown, MD, USA).

Two hundreds microliter of the sample lysate and 200 µL isopropanol was added into the extraction plate.

RT-qPCR was performed using the previously published primers and the modified probe (A to G at 66) in

qScript XLT 1-Step RT-qPCR ToughMix (QuantaBio, Beverly, MA, USA) [17].

Serology

Virus neutralization tests were performed according to a previously established protocol [18]. Briefly, equal

volumes of heat-treated serum (50 µL) and 100 $TCID_{50}$/50 µL of virus were mixed and incubated at 37°C

for 1 hour. The mixture was transferred onto pre-washed MDCK cells. On day 2, IFA was performed to

determine 50% inhibition of virus growth. In addition, commercially available enzyme-linked

immunosorbent assay (ELISA) kits were used to evaluate the antibody responses according to the

manufacturers' instruction: (1) ID Screen® Influenza A Nucleoprotein Swine Indirect, Innovative

Diagnostics, France, (2) ID Screen® Influenza A Antibody Competition Multi-Species, Innovative

Diagnostics, France, and (3) IDEXX AI MultiS-Screen Ab Test, IDEXX, USA.

Gross pathology and histopathology

Post-mortem examinations were conducted at 3, 5, and 21 DPC. Macroscopic pathology was scored *in*

toto, and the percentage of lung lesions calculated based on the previous published method protocols [19].

Tissue samples were fixed in 10% neutral buffered formalin, processed, and stained with hematoxylin and

eosin (H&E). Microscopic pathology was evaluated based on microanatomy of the lung using previously

described scoring scale with minor modifications [20]. Each lung section was scored from 0 to 4 scale (0=

absent, 1=minimal, 2=mild, 3=moderate, and 4=severe) for six separate criteria typically associated with

influenza A virus infections in pigs: (i) epithelial necrosis, attenuation or disruption; (ii) airway exudate;

(iii) percentage of airways with inflammation; (iv) peribronchiolar and perivascular lymphocytic

inflammation; (v) alveolar exudate; (vi) alveolar septal inflammation. For scoring the percentage of

airways with inflammation (iii): 0%=0, up to 10% involvement =1, 10–39% involvement =2, 40–69%

involvement =3 and, greater than 70% involvement =4.

Influenza A virus-specific immunohistochemistry (IHC)

Immunohistochemistry (IHC) for detection of Influenza virus A H5N1 nucleoprotein (NP) antigen was

performed on the automated BOND RXm platform and the Polymer Refine Red Detection kit (Leica

Biosystems, Buffalo Grove, IL). Following automated deparaffinization, four-micron formalin-fixed,

paraffin-embedded tissue sections on positively charged Superfrost® Plus slides (VWR, Radnor, PA)

were subjected to automated heat-induced epitope retrieval (HIER) using a ready-to-use EDTA-based

retrieval solution (pH 9.0, Leica Biosystems) at 100 °C for 20 min. Subsequently, tissue sections were

incubated with the primary antibody (rabbit polyclonal anti-Influenza A virus NP [ThermoFisher

Scientific, PA5-32242] diluted 1:500 in Primary Antibody diluent [Leica Biosystems]) for 30 min at

ambient temperature followed by a polymer-labeled goat anti-rabbit IgG coupled with alkaline

phosphatase (30 min). Fast Red was used as the chromogen (15 min), and counterstaining was performed

with hematoxylin for 5 min. Slides were dried in a 60 °C oven for 30 min and mounted with a permanent

mounting medium (Micromount®, Leica Biosystems). Lung sections from a pig experimentally infected

with swine influenza virus A/swine/Texas/4199-2/1998 H3N2 were used as a positive assay control.

Influenza A virus H5N1 mink-specific RNAscope® in situ hybridization.

For RNAscope® in situ hybridization (ISH), an anti-sense probe targeting the NP and the hemagglutinin

(HA [H5 subtype]) of influenza virus A/Mink/Spain/3691-8_22VIR10586-10/2022 clade 2.3.4.4b

(GISAID # EPI_ISL_15878539) were designed (Advanced Cell Diagnostics (ACD), Newark, CA, USA).

Sections of formalin-fixed paraffin-embedded tissues were generated as indicated above, and the

RNAscope® ISH assay was performed using the RNAscope 2.5 LSx Reagent Kit (ACD) on the

automated BOND RXm platform (Leica Biosystems, Buffalo Grove, IL, USA). Following automated

baking and deparaffinization, tissue sections were subjected to heat-induced epitope retrieval (HIER)

using an EDTA-based solution (pH 9.0; Leica Biosystems) at 100°C for 15 min, protease digestion using

the RNAscope® 2.5 LSx Protease for 15 min at 40 °C, and incubation with a ready-to-use hydrogen

peroxide solution for 10 min at room temperature. Slides were incubated with each probe mixture for 2 h

at 40 °C, and the signal was amplified using a specific set of amplifiers (AMP1 through AMP6) as

recommended by the manufacturer. The signal was detected using a Fast-Red solution for 10 min at room

temperature. Slides were counterstained with a ready-to-use hematoxylin for 5 min, followed by five

washes with 1X BOND Wash Solution (Leica Biosystems). Slides were finally rinsed in deionized water,

dried in a 60 °C oven for 30 min, and mounted with Ecomount® (Biocare, Concord, CA, USA). Lung

sections from a pig experimentally infected with swine influenza virus A/swine/Texas/4199-2/1998 H3N2

were used as a positive assay control for the NP-specific probe, while lung sections from this study were

used in conjunction with the former to validate the HA-specific probe.

Next-generation sequencing (NGS)

The whole genome sequence of the mink-derived H5N1 clade 2.3.4.4b virus was determined using the

Illumina MiSeq sequencing platform (Illumina, San Diego, CA, USA). Briefly, viral RNA was extracted

from the challenge virus inoculum and virus-positive clinical samples using the QIAamp viral RNA mini

kit (Qiagen, Germantown, MD, USA) according to the manufacturer's instructions. Viral gene segments

were amplified using SuperScript™ III One-Step RT-PCR System with Platinum™ Taq DNA Polymerase

(Thermo Fisher Scientific, Waltham, WA, USA) with the Opti and Uni universal influenza primer sets [21,22]

or with SuperScript™ IV First-Strand Synthesis System and Platinum™ SuperFi™ DNA Polymerase

(Thermo Fisher Scientific, Waltham, WA, USA) with previously published segment specific primer sets

[23]. All samples were normalized to 100−500 ng of DNA prior to library preparations. Sequencing libraries

were prepared using the Illumina DNA Prep kit (Illumina, San Diego, CA). Libraries were sequenced

using pair-end chemistry on the Illumina MiSeq platform with the Miseq v2 Reagent kit (300 cycles).

Sequencing reads were demultiplexed and parsed into individual FASTQ files and imported into CLC

Genomics Workbench version 22.0.1 (Qiagen, Germantown, MD, USA) for analysis. Reads were

trimmed to remove primer sequences and filtered to remove low quality and short reads. The trimmed

reads were mapped to the reference sequences (GISAID accession numbers: EPI2220590 to

EPI2220597). Following read mapping, all samples were run through the low frequency variant caller

module within CLC Genomic Workbench with a frequency cutoff greater than 2%.

Results

Infection of pigs with the mink-derived clade 2.3.4.4b H5N1

This study aimed to elucidate the pathogenicity and transmissibility of the mink-derived clade 2.3.4.4b

H5N1 virus in pigs. At 24 hours post-challenge, all principal-infected pigs became lethargic, and five of

nine pigs were febrile with average temperatures of 41 °C. However, the average temperature decreased

to 39.5 °C at 2 DPC and pigs were healthy with no obvious clinical signs throughout the rest of the

observation period (Supplementary Figure 1).

Infectious virus was isolated from nasal (6/9 animals positive) and oropharyngeal swabs (4/9 positive)

from principal infected pigs and the titers ranged from 4.46×10^1 to 1×10^3 TCID$_{50}$/mL at 1 DPC (Table

1). At 3 DPC, one nasal and one oropharyngeal swab were virus positive, with titers of 2.15×10^3 and

4.64×10^2 TCID$_{50}$/mL, respectively. Infectious virus was still present in the nasal swab at 5 DPC ($2.15 \times$

10^2 TCID$_{50}$/mL). RT-qPCR results showed that all principal-infected animals were positive on at least one

time point from 1–5 DPC in either the nasal or oropharyngeal swabs (Supplementary Figure 2A and 2B).

All swabs collected from sentinel pigs were negative by virus isolation and RT-qPCR throughout the

entire observation period.

To determine virus replication and distribution in tissues, virus titers were determined in BALF and

various tissue samples. All BALF samples at 3 and 5 DPC were positive with virus titers of 2.15×10^3 to

4.64×10^4 TCID$_{50}$/mL at 3 DPC and 4.64×10^2 to 3.16×10^3 TCID$_{50}$/mL at 5 DPC (Figure 1B).

Infectious virus could also be isolated from tracheas (3/6), bronchi (2/6), lung tissues (5/6),

tracheobronchial lymph nodes (1/6), and tonsils (1/6) of six principal-infected pigs that were euthanized

at 3 (n=3 pigs) and 5 (n=3 pigs) DPC (Figure 1C and 1D). Interestingly, viral RNA was detectable in the

ethmoturbinates of all three animals sacrificed at 3 DPC (Supplementary Figure 2C). In addition, viral

RNA was present in the nasal turbinate of one pig at 3 DPC; this pig shed infectious virus from the nasal

cavity at 1 and 3 DPC. In contrast, the brain and olfactory bulb at 3 and 5 DPC were negative as assessed

by virus isolation and RT-qPCR (Supplementary Figure 2C and 2D). As expected, BALF and tissue

samples from both principal-infected and sentinel animals at 21 DPC were negative by both virus

isolation and RT-qPCR.

To determine the antibody responses in principal-infected and sentinel pigs, virus neutralization tests

(VNTs) and enzyme-link immunosorbent assays (ELISAs) were performed. Neutralizing antibodies were

detectable in all three principal-infected animals starting at 7 DPC and all principals were sero-positive at

14 and 21 DPC (Figure 1E), with neutralization titers ranging from 1:10 to 1:160. The NP-based indirect

ELISA revealed that two out of three principal-infected animals seroconverted at 7 DPC, with the levels

of antibodies in all three principal-infected pigs against NP increasing by 14 and 21 DPC (Figure 1F).

Two different NP-based competitive ELISAs confirmed the seroconversion of the three principal-infected

pigs at the 14 and 21 DPC (Supplementary Figure 3A and 3B).

Macroscopic and microscopic pathology

At 3 DPC, moderate to severe, multifocal to coalescing interstitial pneumonia with marked congestion

and edema affected all lobes were noted in 2/3 euthanized animals, while the remaining animal had mild,

multifocal pneumonia with congestion predominately the right lung and left caudal lung lobes

(Supplementary Figure 4A–4C). Pulmonary lesions consisted of moderate to severe non-suppurative

interstitial pneumonia with necrotizing bronchitis/bronchiolitis affecting multiple bronchopulmonary

segments, predominantly the pulmonary parenchyma adjacent to terminal bronchioles. Affected

bronchopulmonary segments were characterized by degeneration, attenuation and necrosis of the

bronchiolar epithelium and intraluminal cell debris and degenerate neutrophils, as well as peribronchiolar

and perivascular histiocytic, neutrophilic and lymphocytic inflammation locally extending to and

expanding alveolar septa. Smaller bronchi and bronchioles were most commonly affected, and necrotizing

alveolitis was characterized by alveolar spaces that contain inflammatory cells, occasional fibrin and

necrotic cell debris; the epithelial lining was segmentally to partially denuded or lined by swollen

degenerate epithelium (Figure 3A). These microscopic changes correlated with intracytoplasmic and/or

intranuclear viral NP antigen frequently within bronchial/bronchiolar epithelial cells, and less frequently

within pneumocytes and alveolar macrophages (Figure 3B). Viral NP (as determined by NP-specific IHC)

and H5-specific M or HA RNA (as determined by RNAscope® ISH) were predominantly detected within

airway epithelia and intraluminal necrotic cell debris, and sporadically in inflammatory cells within the

bronchiolar lamina propria (Figure 4A and 4B). In 1/3 pigs (#839), there was localized suppurative and

erosive rhinitis affecting the olfactory region (Supplementary Figure 5A). Within the affected region, viral

antigen was detected in sporadic foci of the olfactory neuroepithelium (Supplementary Figure 5B). Larger

airways (segmental bronchi and trachea) were affected by mild, multifocal neutrophilic to

lymphohistiocytic bronchitis and tracheitis, with frequent bronchial but rare tracheal epithelial cells

containing viral antigen (Supplementary Figure 5C and 5D).

By 5 DPC, multifocal to coalescing areas of pulmonary consolidation were observed on all pigs

(Supplementary Figure 4D–4F). In 2/3 pigs, focally extensive pneumonia was accompanied by large air-

filled bullae formation in the left dorsal lobes. Microscopically, alterations were similar to but more

extensive than those at 3 DPC with intense inflammation affecting larger regions of the pulmonary

parenchyma, with frequent bronchial/bronchiolar involvement and locally extensive alveolar collapse and

consolidation (Figure 3C). Viral NP antigen and, less frequently, NP- and HA-specific RNA were detected

in alveolar macrophages and intraluminal cell debris (Figure 3D and 4C–4D). In addition, numerous

individuals and small clusters of epithelia on medium to large airways contain viral antigen. Microscopic

changes in the trachea were mild and characterized by neutrophilic and lymphohistiocytic tracheitis with

sporadic epithelial cell degeneration/necrosis and rare intracytoplasmic viral antigen (Supplementary

Figure 5E and 5F). Multifocal, mild suppurative to erosive rhinitis affecting rostral turbinates was

identified, although in lieu of intralesional viral antigen. On 21 DPC, multiple foci of mild

lymphohistiocytic interstitial pneumonia were noted, accompanied by bronchiolar epithelial hyperplasia

(repair) with very rare bronchiolar epithelial cells or proprial mononuclear cells containing viral antigen

(Figure 3E and 3F).

Interestingly, Influenza A virus NP-positive antigen presenting cells were common within

tracheobronchial lymph nodes, most abundant at 3 DPC (with an average of 10-20 cells per 200X field)

and gradually reducing in numbers through 21 DPC (Supplementary Figure 6). No other significant gross

or histologic lesions, or viral antigen were identified in other organs examined.

Emergence of mammalian-like mutations in H5N1-infected pigs

The whole-length genome of the H5N1 virus inoculum, two nasal swabs, five oropharyngeal swabs and

six BALF samples were genetically characterized to determine genetic evolution of the H5N1 virus after

replication for up to 5 days in pigs. Our results show that the original mammalian-like PB2 T271A

mutation present in the inoculum remained stable with a frequency below 2% with no evidence of a

change in mutational frequency after virus replication for several days pigs (Table 2 and Supplementary

Table 2). Amino acid substitutions with a significant frequency (>2%) were found throughout all eight

viral segments in clinical samples from H5N1-infected pigs; these mutations were below a frequency of

2% in the virus inoculum (Supplementary Table 2). NGS analysis revealed that oropharyngeal swab #517

on 1 DPC had the mammalian-like K526R mutation of PB2 at a frequency of approximately 3% (Table

2). Interestingly, the critical mammalian adaptation mutations, PB2-E627K and PB2-E627V, were

identified in the oropharyngeal swab of pig #571 (E627K; 1 DPC) , the BALF sample of pig #839

(E627K; 3 DPC), in the oropharyngeal swab of pig #94 (E627V; 1 DPC) and the BALF sample of pig

#213 (E627V; 5 DPC) at a frequency of approximately 5%, 5%, 7% or 2%, respectively (see Table 2;

Supplementary Table 1). Furthermore, the oropharyngeal swab of pig #94 (1 DPC) had the Q222L

adaptation mutation in the HA gene at the frequency of 2.1%.

Discussion

Rapid evolution of IAV is driven by two mechanisms: (1) the introduction of point mutations by the error-

prone RNA polymerase, also called antigenic drift; and (2) by acquisition of one or more RNA segments

from another influenza virus, a process called reassortment or antigenic shift. Molecular evolution allows

IAVs to cross species barriers and acquire the ability to infect a wide range of hosts from avian species to

swine to humans. This broad host tropism provides opportunities for IAVs to gain distinct geno- and

phenotypes, to attain pandemic potential, and to evade pre-existing immunity in humans [24]. Specific host-

virus-environment characteristics accelerate the emergence of novel IAVs with pandemic potential. For

pandemic preparedness, it is critical to assess the potential pandemic risk of newly emerged IAVs; the

evaluation of certain virus characteristics is mainly done by *in vitro* and sometimes also *in vivo*

experiments [25]. However, it is essential to assess the pathogenesis and transmissibility of IAVs in target

animal models. Pigs are natural hosts for IAVs, and it is known that certain IAVs can be efficiently

transmitted between pigs. Importantly, interspecies transmissions between human and pigs and *vice versa*

are not uncommon and are facilitated by an extensive human-pig interface in various agricultural settings

[26]. Furthermore, pigs express receptors specific for both, mammalian and avian IAVs, providing

opportunities for co-infections with these viruses and the subsequent generation of novel reassortant

influenza viruses [27]. Recently, interspecies transmission of clade 2.3.4.4b H5N1 virus to mink and

sustained mink-to-mink transmission was reported in Spain, posing a significant concern to public health

[13]. Therefore, it was crucial to evaluate the pathogenicity and transmissibility of the recently emerged

clade 2.3.4.4b H5N1 virus isolated from diseased mink in domestic pigs.

One of key findings in this study was that experimental infection of pigs with the mink-derived clade

2.3.4.4b H5N1 virus resulted in productive virus replication and seroconversion in 100% of the principal-

infected pigs. Virus titers in BALF samples of principal-infected animals reached $10^{3.3}$ to $10^{4.7}$ TCID$_{50}$/mL

at 3 DPC and declined to $10^{2.7}$ to $10^{3.5}$ TCID$_{50}$/mL at 5 DPC. In addition, the virus was found in other

respiratory tissues, such as trachea, bronchi, as well as lymphoid tissues, such as tracheobronchial lymph

nodes and tonsils. Moreover, gross and histological evaluation of various tissues demonstrated that the

infection caused acute multifocal to coalescing necrotizing broncho-interstitial pneumonia at 3 and 5

DPC, with residual mild interstitial pneumonia still present at 21 DPC. Based on the microscopic lesion

distribution, virus titers and viral antigen/RNA tissue expression, it is clear that the lesions and

distribution of this mink-derived clade 2.3.4.4b H5N1 virus mostly involved the lower respiratory tract

(i.e., pulmonary parenchyma) with only mild alterations on the upper respiratory tract. Previous studies

demonstrated virus replication of avian-derived H5Nx clade 2.3.4.4 HPAI viruses from North America in

the lower respiratory tract of pigs upon experimental infection [28]. In the latter study, infection of pigs with

the avian-origin H5Nx clade 2.3.4.4 viruses, isolated in 2014 and 2015 in the U.S, led to viral RNA

detection in 80-100% of the animals and successful virus isolation in 60-100% of the BALF samples at 3

and 5 DPC [28]. Consistent with virus detection in the majority of animals, 60-100% of infected animals

seroconverted by 21 DPC (28). In addition, in a very recent study, low susceptibility of pigs against

experimental infection with an avian-derived H5N1 clade 2.3.4.4b virus, isolated from chickens in

Germany in 2022, was reported [29]. This chicken H5N1 clade 2.3.4.4b isolate lacked any mammalian-

adaptive mutations. Nasal and alimentary exposure of pigs to this avian-derived H5N21 clade 2.3.4.4b

virus only resulted in marginal virus replication, without inducing any clinical signs or pathological

changes. Only 1 out of 8 pigs seroconverted, indicating a rather high resistance of pigs to infection with

the avian-derived derived H5N21 clade 2.3.4.4b virus without mammalian-like adaptations [(29)].

Experimental infection of pigs with LPAI viruses without mammalian-like adaptations were also

previously performed, but none of the experimental infections with LPAI H5 viruses resulted in a

productive infection [30]. We conclude from our present study with the mink-derived clade 2.3.4.4b H5N1

HPAI virus, that this virus is not only able to efficiently infect mink, but also pigs. The mink-derived

H5N1 clade 2.3.4.4b virus efficiently replicated in the lower respiratory tract of pigs inducing sizeable

lung lesions; in addition, nasal and oral shedding was observed on early days after infection although less

efficient than reported for swine-derived influenza viruses [31,32]. However, it is important to note that the

mink-derived H5N1 clade 2.3.4.4b virus was unable to spread from principal-infected to co-mingled in-

contact sentinel animals; this was most likely due to the limited amount of virus shedding by the nasal

and oral cavities in principal-infected pigs. Interestingly, neither infectious virus, viral RNA nor

histopathological changes were present in tissues from the central nervous system of principal-infected

pigs in our study. In recent spillover events, wild mammals infected with the clade 2.3.4.4b H5N1 virus

primarily exhibited neurological symptoms with histopathological changes in brains as well as viral RNA

detection in CNS tissues [33-35]. Similarly, the mink infected with the H5N1 clade 2.3.4.4b virus used in this

study also showed neurological manifestations, such as ataxia and tremors [13]. We could not see any

neurological clinical signs in principal-infected pigs, indicating that the pathogenicity of the H5N1 clade

2.3.4.4b viruses could vary in different mammalian host species.

Earlier studies showed that experimental infection of pigs with avian-origin LPAI H5 subtype viruses

isolated between 1980 and 2005 led to nasal shedding of infectious virus of up to $10^{5.5}$ egg infectious

doses (EID)$_{50}$/mL and $10^{5.3}$ EID$_{50}$/100 mg [36,37]. In addition, it was shown that experimental infection of

pigs with human and avian-origin HPAI H5N1 viruses isolated in early (1997) and later (2004) in the

emergence of the HPAIV H5N1 resulted in the isolation of infectious virus in nasal swabs/nasal tracts

over the course of infection, but no pig-to-pig transmission [38,39]. When pigs were infected with the index

human H5N1 HPAIV strain (HK156-97) and an initial chicken H5N1 HPAIV isolate (CHK258-97) by the

oral and nasal routes, both the human and chicken H5N1 viruses replicated well in the nasal tract of pigs,

with the chicken isolate to higher titers than the human virus. Importantly, neither the human nor the

chicken H5N1 virus transmitted to contact pigs in the same pen. Thus, pigs support the replication of the

index human and an early chicken H5N1 virus to modest titers, and there was no detectable transmission

to contact animals [38]. Similarly, experimental studies on the replication and transmissibility of 2004 avian

H5N1 HPAI viruses associated with human infections in Vietnam to pigs revealed that H5N1 all viruses

tested replicated in the swine respiratory tract but none were transmitted to contact pigs (39).

Epidemiological studies showed limited serologic evidence (8/3175 pigs positive; 0.25%) of exposure to

the HPAI H5N1 viruses in Vietnamese pigs [39]. These findings indicate that pigs can be infected with the

2004 Vietnamese HPAI H5N1 viruses but that these viruses are not readily transmitted between pigs

under experimental conditions [39]. In contrast to the above described work, no viral RNA was detected in

nasal swabs of pigs infected with avian-derived clade 2.3.4.4 H5Nx viruses in 2014 [28], and only limited

detection of viral RNA but failure to isolate infectious virus was reported when pigs were infected with

an avian-derived clade 2.3.4.4b H5N1 virus without mammalian-like adaptations [29]. Collectively, these

results suggest that some isolates of HPAI H5 subtype viruses can replicate efficiently in the upper

respiratory tract of pigs, with limited replication in the lower respiratory tract. There is no transmission of

these viruses to co-mingled contact sentinels.

Through the continuous evolution of the HPAI H5Nx viruses over two decades, the

A/Goose/Guangdong/1/1996 H5N1 lineage (GsGD) may have acquired increased fitness in avian species

resulting in efficient spread across the globe. However, the continued adaptation of the H5N1 clade

2.3.4.4b virus to avian species may, in turn, contribute to a loss of fitness for mammalian species, which

resulted in its reduced replication capacity in the upper respiratory tract of pigs. Despite the shedding of

infectious virus in nasal and oropharyngeal secretions from principal-infected pigs, there was no evidence

of transmission of the mink-derived HPAI H5N1 clade 2.3.4.4b virus to sentinel pigs under our

experimental conditions. This result was consistent with previous findings indicating a lack of pig-to-pig

transmission for H5- or H7-subtype HPAIVs, regardless of the presence of infectious virus in nasal and/or

oral excretions [28,29,38,39]. The lack of transmissibility of the mink-derived H5N1 clade 2.3.4.4b isolate in

pigs is supported by the absence of seroconversion in the co-mingled sentinel animal; also, this virus did

not transmitted to the 11 farm workers that had been in close contact with the H5N1-infected minks at the

site of the outbreak [13]. Furthermore, a recent study showed that two other mink-derived H5N1 viruses

which differ by several amino acids in the PB2, PB1, PA, NA, NS1 and NS2 proteins, did not support

transmission to contact animals (neighboring cages) through respiratory droplets in the ferret model [40].

Importantly, it has been previously documented that H5 subtype viruses can acquire the ability to transmit

efficiently among pigs through reassortment. Lee *et al.* [41] isolated two H5N2 viruses from pigs showing

typical clinical symptoms of influenza-like illness; one was an avian-origin virus, the other one was an

avian-swine reassortant virus with PB2, PA, NP and M genes derived from swine influenza viruses. Upon

experimental infection of pigs, both viruses could be isolated from nasal swabs, but pig-to-pig

transmission occurred only in the group infected with the reassortant H5N2 virus [41].

One of the important findings is that the H5N1 viruses inoculated into pigs acquired mammalian-like

mutations/adaptations in the course of several days of pig infection, which potentially results in increased

fitness of the H5N1 virus in mammalian hosts. There is a total of four mink-derived H5N1 isolates from

the infected Spanish farm which were sequenced; all of them possessed the PB2 T271A mutation. The

PB2 T271A mutation is known to lead to enhanced polymerase activity in mammalian cells and might

have contributed to cross-species transmission from avian species to mammals including mink [13,42]. Our

results show that the T271A mutation was stable with no evidence of a change in the frequency of the

mutation after replication of the virus for several days in pigs. However, minor variants possessing the

mammalian-like E627K mutation in PB2 emerged in oropharyngeal swab (pig #714) and BALF (pig

#839) samples obtained from principal-infected pigs, even though they were found at low frequencies

(4.7% and 5.2%, respectively). The E627K mutation is a key determinant for mammalian adaptation of

avian influenza viruses; its introduction into the genome of an avian influenza virus enables efficient

replication of the avian-origin polymerase complex in mammalian cells [43]. It is well known that H5N1

viruses harboring the PB2 E627K mutation have increased virulence in mice; it also contributes to air-

borne virus transmission in ferrets and contact transmission in guinea pigs [44-48]. Similarly, an amino acid

substitution from glutamic acid (E) to valine (V) at position 627 of the PB2 gene (E627V) was shown to

increase viral replication in mammalian cells and virulence in mice models [49]; an E627V mutation was

detected as a minor virus population in the oropharyngeal swab of pig #94 and BALF sample of pig #213.

Furthermore, the oropharyngeal swab of gig #517 possessed a PB2 K526R mutation at the frequency of

3.1%; the K526R mutation was shown to enhance the effect of the PB2 E627K on avian influenza virus

replication in mammalian cells and also their virulence in mice [50]. Another key mutation is the Q222L

(according to the H5 numbering system) in the HA gene which is equivalent to a Q226L mutation using

the H3 numbering system. This mutation was found in oropharyngeal swab of pig #94 at 1 DPC. Amino

acid (AA) position 222 in the HA is involved in the binding of HA to its sialic acid receptor and plays a

critical role in determining the host range of IAVs. The HA of mammalian IAVs binds to α2,6-linked

sialic acid, whereas the HA of avian IAVs binds to α2,3-linked sialic acid. Several amino acids in the HA

receptor binding site are responsible for switching sialic acid specificity; however, the Q222L and G224S

AA positions in the H5 numbering system, which are equivalent to AA positions Q226L and G228S in the

H3 numbering system, are crucial for increased α2,6-sialic acid receptor preference in different subtypes

of avian IAVs; they are located in the receptor binding domain and interact directly with the host's sialic

acid receptor [51]. In addition, the Q222L mutation was previously described to contribute to the emergence

of transmissible H5N1 viruses via aerosols after serial passage of H5N1 in ferrets [45,46]. Despite the critical

role of these mutations in pathogenicity and transmissibility of IAVs in mammalian hosts, it still remains

unclear how the H5N1 clade 2.3.4.4b virus underwent its molecular evolution, and eventually acquired

critical mammalian-like mutations. In addition to the role of pigs in the ecology of IAVs as the "mixing

vessel", the emergence of mammalian-adapted H5N1 viruses in pigs is close to reality; therefore, pigs

may serve as the "hotspot" to introduce mammalian-like mutations into avian influenza viruses although

the frequencies of the mammalian-like mutations described here are rather low.

Spillover of H5N1 HPAI clade 2.3.4.4b viruses to mammals and their sustained transmission between

mammals has been reported [13,52]. The majority of mammals infected by the HPAI H5N1 clade 2.3.4.4b

virus are carnivores, and the consumption of infected dead wild birds is presumably the basis for these

spillover events [53]. In contrast to the HPAI H5N1 clade 2.3.4.4b virus infections in wild avians and wild

mammals, outbreaks in farmed avian and mammal species pose a much greater risk to humans, due to its

close proximity to occupational workers and -in the case of mammalian species - the favorable

environment for the acquisition of mammalian-like adaptations. Therefore, evaluating the

infectivity/susceptibility and transmissibility of the emerging HPAI viruses in well-established

mammalian animal models such as swine and ferrets is crucial to determine their pandemic potential. The

present study demonstrated that infection of pigs with the mink-derived clade 2.3.4.4b H5N1 virus led to

a productive viral replication in the respiratory tracts of all principal-infected pigs including

seroconversion; however, virus shedding from principal-infected pigs was not high and/or frequent

enough for an efficient transmission to co-mingled sentinel pigs. Notably, key mammalian-like

mutations were found in clinical and/or tissue samples derived from principal-infected pigs. In

conclusion, the results obtained in the present study suggest that the mink-derived H5N1 clade 2.3.4.4b

virus exhibited increased infectivity in pigs when compared to avian-origin H5N1 clade 2.3.4.4b viruses,

however, did not transmit to co-mingled sentinel animals; therefore, this virus would still be placed in a

moderate risk group in terms of transmission ability to humans.

References

1   Tong, S. *et al.* A distinct lineage of influenza A virus from bats. *Proc Natl Acad Sci U S A* **109**,
450   4269-4274 (2012). https://doi.org/10.1073/pnas.1116200109
451   2   Tong, S. *et al.* New world bats harbor diverse influenza A viruses. *PLoS Pathog* **9**, e1003657
452   (2013). https://doi.org/10.1371/journal.ppat.1003657

3   Vey, M. *et al.* Hemagglutinin activation of pathogenic avian influenza viruses of serotype H7
requires the protease recognition motif R-X-K/R-R. *Virology* **188**, 408-413 (1992).
https://doi.org:10.1016/0042-6822(92)90775-k

4   Xu, X., Subbarao, Cox, N. J. & Guo, Y. Genetic characterization of the pathogenic influenza
A/Goose/Guangdong/1/96 (H5N1) virus: similarity of its hemagglutinin gene to those of H5N1
viruses from the 1997 outbreaks in Hong Kong. *Virology* **261**, 15-19 (1999).
https://doi.org:10.1006/viro.1999.9820

5   WHO/OIE/FAO H5N1 Evolution Working Group. Toward a unified nomenclature system for highly
pathogenic avian influenza virus (H5N1). *Emerg Infect Dis* **14**, e1 (2008).
https://doi.org:10.3201/eid1407.071681

6   World Health Organization. *Evolution of the influenza A(H5) haemagglutinin: WHO/OIE/FAO H5*
Working Group reports a new clade designated 2.3.4.4,
<https://www.who.int/publications/m/item/evolution-of-the-influenza-a(h5)-haemagglutinin-
who-oie-fao-h5-working-group-reports-a-new-clade-designated-2.3.4.4> (2015).

7   Pohlmann, A. *et al.* Has Epizootic Become Enzootic? Evidence for a Fundamental Change in the
Infection Dynamics of Highly Pathogenic Avian Influenza in Europe, 2021. *mBio* **13**, e0060922
(2022). https://doi.org:10.1128/mbio.00609-22

8   Bevins, S. N. *et al.* Intercontinental Movement of Highly Pathogenic Avian Influenza A(H5N1)
Clade 2.3.4.4 Virus to the United States, 2021. *Emerg Infect Dis* **28**, 1006-1011 (2022).
https://doi.org:10.3201/eid2805.220318

9   Shinya, K. *et al.* Avian flu: influenza virus receptors in the human airway. *Nature* **440**, 435-436
(2006). https://doi.org:10.1038/440435a

10   World Health Organization.   (2023).

11   Harfoot, R. & Webby, R. J. H5 influenza, a global update. *J Microbiol* **55**, 196-203 (2017).
https://doi.org:10.1007/s12275-017-7062-7

12   World Health Organization.   (2022).

13   Aguero, M. *et al.* Highly pathogenic avian influenza A(H5N1) virus infection in farmed minks,
Spain, October 2022. *Euro Surveill* **28** (2023). https://doi.org:10.2807/1560-
7917.ES.2023.28.3.2300001

14   Ma, W. *et al.* 2009 pandemic H1N1 influenza virus causes disease and upregulation of genes
related to inflammatory and immune responses, cell death, and lipid metabolism in pigs. *J Virol*
85, 11626-11637 (2011). https://doi.org:10.1128/JVI.05705-11

15   Reed, L. J. & Muench, H. A Simple Method of Estimating Fifty Per Cent Endpoints12. *American*
Journal of Epidemiology **27**, 493-497 (1938). https://doi.org:10.1093/oxfordjournals.aje.a118408

16   Gaudreault, N. N. *et al.* SARS-CoV-2 infection, disease and transmission in domestic cats. *Emerg*
Microbes Infect **9**, 2322–2332 (2020). https://doi.org:10.1080/22221751.2020.1833687

17   Sponseller, B. A. *et al.* Influenza A pandemic (H1N1) 2009 virus infection in domestic cat. *Emerg*
Infect Dis **16**, 534-537 (2010). https://doi.org:10.3201/eid1603.091737

18   Gauger, P. C. & Vincent, A. L. Serum Virus Neutralization Assay for Detection and Quantitation of
Serum Neutralizing Antibodies to Influenza A Virus in Swine. *Methods Mol Biol* **2123**, 321-333
(2020). https://doi.org:10.1007/978-1-0716-0346-8_23

19   Artiaga, B. L. *et al.* Evaluating alpha-galactosylceramide as an adjuvant for live attenuated
influenza vaccines in pigs. *Anim Dis* **2**, 19 (2022). https://doi.org:10.1186/s44149-022-00051-x

20   Lee, J. *et al.* Effects of PB1-F2 on the pathogenicity of H1N1 swine influenza virus in mice and
pigs. *J Gen Virol* **98**, 31-42 (2017). https://doi.org:10.1099/jgv.0.000695

21   Lee, D. H. Complete Genome Sequencing of Influenza A Viruses Using Next-Generation
Sequencing. *Methods Mol Biol* **2123**, 69-79 (2020). https://doi.org:10.1007/978-1-0716-0346-
8_6

Zhou, B. & Wentworth, D. E. Influenza A virus molecular virology techniques. *Methods Mol Biol* **865**, 175-192 (2012). https://doi.org:10.1007/978-1-61779-621-0_11

Hoffmann, E., Stech, J., Guan, Y., Webster, R. G. & Perez, D. R. Universal primer set for the full-length amplification of all influenza A viruses. *Arch Virol* **146**, 2275-2289 (2001). https://doi.org:10.1007/s007050170002

Mena, I. *et al.* Origins of the 2009 H1N1 influenza pandemic in swine in Mexico. *Elife* **5** (2016). https://doi.org:10.7554/eLife.16777

World Health Organization. Tool for Influenza Pandemic Risk Assessment (TIPRA). (2020).

Rajao, D. S. & Vincent, A. L. Swine as a model for influenza A virus infection and immunity. *ILAR J* **56**, 44-52 (2015). https://doi.org:10.1093/ilar/ilv002

Ito, T. *et al.* Molecular basis for the generation in pigs of influenza A viruses with pandemic potential. *J Virol* **72**, 7367-7373 (1998). https://doi.org:10.1128/JVI.72.9.7367-7373.1998

Kaplan, B. S., Torchetti, M. K., Lager, K. M., Webby, R. J. & Vincent, A. L. Absence of clinical disease and contact transmission of HPAI H5NX clade 2.3.4.4 from North America in experimentally infected pigs. *Influenza Other Respir Viruses* **11**, 464-470 (2017). https://doi.org:10.1111/irv.12463

Graaf, A. *et al.* Low Susceptibility of Pigs against Experimental Infection with HPAI Virus H5N1 Clade 2.3.4.4b. *Emerg Infect Dis* **29**, 1492-1495 (2023). https://doi.org:10.3201/eid2907.230296

Balzli, C. *et al.* Susceptibility of swine to H5 and H7 low pathogenic avian influenza viruses. *Influenza Other Respir Viruses* **10**, 346-352 (2016). https://doi.org:10.1111/irv.12386

Richt, J. A. *et al.* Pathogenic and antigenic properties of phylogenetically distinct reassortant H3N2 swine influenza viruses cocirculating in the United States. *J Clin Microbiol* **41**, 3198-3205 (2003). https://doi.org:10.1128/JCM.41.7.3198-3205.2003

Ma, W. *et al.* Identification of H2N3 influenza A viruses from swine in the United States. *Proc Natl Acad Sci U S A* **104**, 20949-20954 (2007). https://doi.org:10.1073/pnas.0710286104

Elsmo, E. J. *et al.* Pathology of natural infection with highly pathogenic avian influenza virus (H5N1) clade 2.3.4.4b in wild terrestrial mammals in the United States in 2022. *bioRxiv* (2023). https://doi.org:10.1101/2023.03.10.532068

Thorsson, E. *et al.* Highly Pathogenic Avian Influenza A(H5N1) Virus in a Harbor Porpoise, Sweden. *Emerg Infect Dis* **29**, 852-855 (2023). https://doi.org:10.3201/eid2904.221426

Bordes, L. *et al.* Highly Pathogenic Avian Influenza H5N1 Virus Infections in Wild Red Foxes (Vulpes vulpes) Show Neurotropism and Adaptive Virus Mutations. *Microbiol Spectr* **11**, e0286722 (2023). https://doi.org:10.1128/spectrum.02867-22

Kida, H. *et al.* Potential for transmission of avian influenza viruses to pigs. *J Gen Virol* **75 (Pt 9)**, 2183-2188 (1994). https://doi.org:10.1099/0022-1317-75-9-2183

De Vleeschauwer, A., Van Poucke, S., Braeckmans, D., Van Doorsselaere, J. & Van Reeth, K. Efficient transmission of swine-adapted but not wholly avian influenza viruses among pigs and from pigs to ferrets. *J Infect Dis* **200**, 1884-1892 (2009). https://doi.org:10.1086/648475

Shortridge, K. F. *et al.* Characterization of avian H5N1 influenza viruses from poultry in Hong Kong. *Virology* **252**, 331-342 (1998). https://doi.org:10.1006/viro.1998.9488

Choi, Y. K. *et al.* Studies of H5N1 influenza virus infection of pigs by using viruses isolated in Vietnam and Thailand in 2004. *J Virol* **79**, 10821-10825 (2005). https://doi.org:10.1128/JVI.79.16.10821-10825.2005

Maemura, T. *et al.* Characterization of highly pathogenic clade 2.3.4.4b H5N1 mink influenza viruses. *EBioMedicine* **97**, 104827 (2023). https://doi.org:10.1016/j.ebiom.2023.104827

Lee, J. H. *et al.* Isolation and genetic characterization of H5N2 influenza viruses from pigs in Korea. *J Virol* **83**, 4205-4215 (2009). https://doi.org:10.1128/JVI.02403-09

42 Bussey, K. A., Bousse, T. L., Desmet, E. A., Kim, B. & Takimoto, T. PB2 residue 271 plays a key role
in enhanced polymerase activity of influenza A viruses in mammalian host cells. *J Virol* **84**, 4395-
4406 (2010). https://doi.org:10.1128/JVI.02642-09

43 Neumann, G., Chen, H., Gao, G. F., Shu, Y. & Kawaoka, Y. H5N1 influenza viruses: outbreaks and
biological properties. *Cell Res* **20**, 51-61 (2010). https://doi.org:10.1038/cr.2009.124

44 Kim, J. H. *et al.* Role of host-specific amino acids in the pathogenicity of avian H5N1 influenza
viruses in mice. *Journal of General Virology* **91**, 1284-1289 (2009).
https://doi.org:10.1099/vir.0.018143-0

45 Herfst, S. *et al.* Airborne transmission of influenza A/H5N1 virus between ferrets. *Science* **336**,
1534-1541 (2012). https://doi.org:10.1126/science.1213362

46 Linster, M. *et al.* Identification, characterization, and natural selection of mutations driving
airborne transmission of A/H5N1 virus. *Cell* **157**, 329-339 (2014).
https://doi.org:10.1016/j.cell.2014.02.040

47 Hatta, M., Gao, P., Halfmann, P. & Kawaoka, Y. Molecular Basis for High Virulence of Hong Kong
H5N1 Influenza A Viruses. *Science* **293**, 1840-1842 (2001).
https://doi.org:10.1126/science.1062882

48 Steel, J., Lowen, A. C., Mubareka, S. & Palese, P. Transmission of influenza virus in a mammalian
host is increased by PB2 amino acids 627K or 627E/701N. *PLoS Pathog* **5**, e1000252 (2009).
https://doi.org:10.1371/journal.ppat.1000252

49 Taft, A. S. *et al.* Identification of mammalian-adapting mutations in the polymerase complex of
an avian H5N1 influenza virus. *Nat Commun* **6**, 7491 (2015).
https://doi.org:10.1038/ncomms8491

50 Song, W. *et al.* The K526R substitution in viral protein PB2 enhances the effects of E627K on
influenza virus replication. *Nat Commun* **5**, 5509 (2014). https://doi.org:10.1038/ncomms6509

51 Stevens, J. *et al.* Structure and receptor specificity of the hemagglutinin from an H5N1 influenza
virus. *Science* **312**, 404-410 (2006). https://doi.org:10.1126/science.1124513

52 Lindh, E. *et al.* Highly pathogenic avian influenza A(H5N1) virus infection on multiple fur farms in
the South and Central Ostrobothnia regions of Finland, July 2023. *Euro Surveill* **28** (2023).
https://doi.org:10.2807/1560-7917.ES.2023.28.31.2300400

53 Adlhoch, C. *et al.* Avian influenza overview April - June 2023. *EFSA J* **21**, e08191 (2023).
https://doi.org:10.2903/j.efsa.2023.8191

Acknowledgments

We gratefully thank Yonghai Li, Patricia Assato, Michelle Zajac, Isaac Fitz for technical assistance, and

Innovative Diagnostics for providing ELISA kits. Funding for this study was provided through grants

from the NIAID supported Center of Excellence for Influenza Research and Response (CEIRR) under

contract number 75N93021C00016, the National Bio and Agro-Defense Facility (NBAF) Transition Fund

from the State of Kansas, the MCB Core of the Center on Emerging and Zoonotic Infectious Diseases

(CEZID) of the National Institutes of General Medical Sciences under award number P20GM130448, the

NIAID Centers of Excellence for Influenza Research and Surveillance (CEIRS) under contract number

HHSN 272201400006C.

Author contributions

J.A.R and N.O. conceptualized the research project. T.K., J.D.T., M.C., E.L.L., C.D.M., K.C., F.S.M.,

T.J., I.M., N.N.G., U.B.R.B., N.O., R.J.W. and J.A.R acquired the data.

Competing interests

The J.A.R. laboratory received support from Tonix Pharmaceuticals, Genus *plc*, Xing Technologies, and

Zoetis, outside of the reported work. J.A.R. is inventor on patents and patent applications on the use of

antivirals and vaccines for the treatment and prevention of virus infections, owned by Kansas State

University. The other authors declare no competing interests.

Figure legions

Figure 1. Infection with a mink-derived clade 2.3.4.4b H5N1 isolate in pigs. **A** The pigs (n=9) were orally, intra-nasally, and intra-tracheally infected with 2.2×10^7 TCID$_{50}$ of A/Mink/Spain/3691-8_22VIR10586-10/2022 and sentinel pigs (n=6) were introduced in a same pen at 2 days post-challenge (DPC). Swab and serum samples were collected throughout the study. Principal infected pigs were euthanized and necropsied at 3 and 5 DPC (n=3 per time point) and the remaining three principals and 6 sentinels were euthanized and necropsied at 21 DPC. **B** Infectious virus titers in bronchoalveolar lavage fluids at 3 and 5 DPC and **C and D** infectious virus titers in tissues at (C) 3 DPC and (D) 5 DPC. **E and F** Antibody responses were evaluated in three principal pigs at -1, 7, 14, and 21 DPC. Virus neutralization (VN) titers (E) were a reciprocal of the highest serum dilution inhibiting replication of the homologous H5N1 virus. ELISA antibody titers were calculated in a swine indirect ELISA kit (ID Screen® Influenza A Nucleoprotein Swine Indirect, Innovative Diagnostics, France) where S/P ratios were calculated from O.D. values and then converted to antibody titers according to the manufacturers' instruction. Dash lines represent the limit of detection of each assay (the limit of detection of each assay (4.64×10^1 TCID$_{50}$/mL for BALF titration, 2.92×10^1 TCID$_{50}$/mL for tissue titration, 1:10 for VN assay, and 1053 for ELISA antibody titer). Empty circle represents negative results in each assay.

Figure 2. Macroscopic and microscopic lung scores in a mink-derived clade 2.3.4.4b H5N1 virus-infected pigs. Infected pigs were euthanized at 3, 5, and 21 DPC for pathological evaluation. Macroscopic lung scores were determined by the percentage of the surface area showing IAV-induced pneumonia. Microscopic scores were calculated by the average of six different criteria (see Materials and Methods)

Figure 3. Histological features and tropism of mink-derived H5N1 clade 2.3.4.4b influenza virus in pigs. Histologically, pulmonary lesions were characterized by necrotizing to lymphohistiocytic

bronchointerstitial pneumonia. At 3 DPC (A and B), multifocal bronchopulmonary segments were

characterized by bronchiolar epithelial cell necrosis (A, inset 1) and associated areas of pulmonary

parenchyma with expanded alveolar septa infiltrated by lymphocytes and histiocytes and alveolar

epithelial necrosis (A, inset 2). Influenza A virus NP intracytoplasmic antigen was detected within

bronchiolar epithelial cells (B, asterisk), pneumocytes and alveolar macrophages (B, inset). Intranuclear

viral antigen was typically detected within infected epithelial cells. Similar but more extensive and severe

histological changes were noted at 5 DPC (C and D), with similar alveolar (C, inset 1) and bronchiolar (C,

inset 2) alterations. Influenza A virus NP antigen was more abundant but had a similar cellular distribution

compared to 3 DPC (D and insets 1 and 2). At 21 DPC, the airway epithelia had mostly repaired but

multifocal areas of interstitial inflammation persisted (E, inset). Only sporadic epithelial cells contained

viral antigen at this timepoint (F and insets).

Figure 4. Detection of mink-derived H5N1 clade 2.3.4.4b influenza virus RNA via in situ hybridization

(ISH). Probes specific to the NP and HA (H5) genes were designed. Viral RNA was predominantly

detected within bronchiolar epithelial cells and intraluminal cellular debris at 3 (A and B) and 5 DPC (C

and D). RNAscope® ISH, Fast Red, total magnification 200X.

644   **Figure 1**

646   **Figure 2.**

Figure 3.

651   **Figure 4.**

652 **Table 1.** Virus shedding in pigs infected with the mink-derived clade 2.3.4.4b H5N1 virus.

Days post-challenge (DPC)			1	2	3	4	5	6 to 21[a]
Nasal swab	Principal	# of positive / # of animals	6/9	0/9	1/9	0/6	1/6	0/3
		Virus titer ($TCID_{50}$/mL)	4.6×10^1 to 1×10^3	Neg	2.15×10^3	Neg	2.15×10^2	Neg
	Sentinel	# of positive / # of animals	0/6	0/6	0/6	0/6	0/6	0/6
Oropharyngeal swab	Principal	# of positive / # of animals	4/9	N/A[b]	1/9	N/A[b]	0/6	0/3
		Virus titer ($TCID_{50}$/mL)	1×10^2 to 4.6×10^2	N/A[b]	4.6×10^2	N/A[b]	Neg	Neg
	Sentinel	# of positive / # of animals	0/6	N/A[b]	0/6	N/A[b]	0/6	0/6

653 [a]Nasal swabs were collected at 6 to 14, 17 and 21 DPC and oropharyngeal swabs were at 7, 10, 14, 17,

654 and 21 DPC. [b]N/A: Samples were not collected.

Table 2. Mammalian-like adaptive mutations after replication of a mink-derived clade 2.3.4.4b H5N1 in pigs.

Viral protein	Amino acid substitution	Inoculum	Nasal swab, #517, 1 DPC	Nasal swab, #517, 3 DPC	OP swab, #94, 1 DPC	OP swab, #517, 1 DPC	OP swab, #571, 1 DPC	OP swab, #839, 1 DPC	OP swab, #634, 3 DPC	BALF, #517, 3 DPC	BALF, #575, 3 DPC	BALF, #839, 3 DPC	BALF, #94, 5 DPC	BALF, #213, 5 DPC	BALF, #714, 5 DPC
PB2 [a]	K526R	< 2%	< 2%	< 2%	< 2%	**3.14%**	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%
	E627K	< 2%	< 2%	< 2%	< 2%	< 2%	**4.71%**	< 2%	< 2%	< 2%	< 2%	**5.19%**	< 2%	< 2%	< 2%
	E627V	< 2%	< 2%	< 2%	**6.96%**	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	**2.02%**	< 2%
HA	Q222L	< 2%	< 2%	< 2%	**2.10%**	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%	< 2%

[a] Amino acid substitution with a frequency >2% was not found at position of 271 of PB2 in all samples tested.

Supplementary Figure 1. Daily body temperature of principal infected and sentinels in this study. The value is expressed as the mean ± SEM. The dash line indicates 40.5 °C, which is considered fever in this study.

Supplementary Figure 2. Viral RNA detection in (A) nasal swab and (B) oropharyngeal swab of H5N1-infected pigs (n=9, at 1, 2, and 3 days post-challenge (DPC); and n=6, at 4 and 5 DPC). The Cq values of positive swabs were shown in the graphs. The distribution of viral RNA in tissues of H5N1-infected pigs at (C) 3 DPC and (D) 5 DPC. Empty circles represent the single well positive in duplicates.

Supplementary Figure 3. Antibody responses in H5N1-infected pigs employing two competitive

ELISAs: (A) ID Screen® Influenza A Antibody Competition Multi-Species and (B) IDEXX AI MultiS-

Screen Ab Test. Empty and grey circles represent negative and suspicious.

3 DPC   **5 DPC**   **21 DPC**

Supplementary Figure 4. Gross changes in the lungs of a mink-derived clade 2.3.4.4b H5N1 virus-infected pigs at 3, 5 and 21 DPC. At 3 DPC, two pigs (A and B) had moderate to severe, multifocal to coalescing interstitial pneumonia with congestion and edema affecting all lobes (A and B), while the third animal (C) had mild, multifocal pneumonia with congestion affecting predominately the right lung and left caudal lung lobes (C). At 5 DPC (D–F), coalescing areas of pulmonary consolidation with congestion and edema were noted in all pigs. The right middle and caudal lobes were most commonly affected (D–F) with occasional bullae (asterisk). At 21 DPC, only a few areas of pulmonary consolidation were noted (G–I).

Supplementary Figure 5. Histological and immunohistochemical findings in the upper respiratory tract of pigs infected with a mink-derived clade 2.3.4.4b H5N1 virus. In a single animal, a focal area of the olfactory epithelium is mildly infiltrated by few lymphocytes (A, arrows) that also mildly infiltrate the subjacent lamina propria (A, arrowhead) at 3 DPC. Within this area, olfactory epithelia contain abundant influenza NP antigen (B, intracytoplasmic and intranuclear). At 3 and 5 DPC, there is mild lymphohistiocytic and neutrophilic tracheitis (C and E) with lymphocytes and histiocytes within the lamina propria and sporadic neutrophils and lymphocytes transmigrating through the lining epithelium (arrows). Sporadic epithelial cells contain intracytoplasmic and intranuclear influenza NP antigen (D and F, arrows). H&E (A, C, E) and IHC (Fast Red, B, D and F), total magnification 200X.

Supplementary Figure 6. Immunohistochemical findings in the tracheobronchial lymph nodes of pigs infected with a mink-derived clade 2.3.4.4b H5N1 virus. At 3 DPC (A), there are numerous influenza NP antigen-positive cells with dendritic projections of their cytoplasm (consistent with antigen presenting cells). The number of immune-positive cells gradually declines at 5 and 21 DPC (B and C, respectively). IHC, Fast Red, total magnification 100X.

Supplementary Table 2. Frequencies of mutations of a mink-derived clade 2.3.4.4b H5N1 in pigs. Blank represents less than 2%.

Viral protein	Amino acid substitution	Inoculum	Nasal swab, #517, 1 DPC	Nasal swab, #517, 3 DPC	OP swab, #94, 1 DPC	OP swab, #517, 1 DPC	OP swab, #571, 1 DPC	OP swab, #839, 1 DPC	OP swab, #634, 3 DPC	BALF, #517, 3 DPC	BALF, #575, 3 DPC	BALF, #839, 3 DPC	BALF, #94, 5 DPC	BALF, #213, 5 DPC	BALF, #714, 5 DPC
PB2	R70W														3.35
	K80R								9.67						
	R101G											3.47			
	P104T							3.25							
	N137K		6.95												
	K140E								2.70						
	V167I												2.57		
	P169S											2.25			
	E180G														9.86
	E188G					2.47									
	D195N		3.28		7.01				3.43						
	V220I												2.49		
	T238A											2.28			
	C239Y													3.71	
	M243K	2.68													
	T245A			3.64											
	R269G														8.89
	S279L						2.12								
	S286N						4.70								
	L317S													2.52	
	N348S												3.04		
	E396K										2.22				
	F420L														2.80
	M431T													2.06	
	H437R			3.91											
	K440R				7.25										
	I463M							3.04							
	S474P												3.00		
	M483K	2.43													
	S497G	2.67													
	K526R					3.14									
	D611N											4.03	2.35		
	T612A					6.26	2.11				2.37		3.95		
	L618I			3.83											
	E627K						4.71					5.19			
	E627V				6.96									2.02	
	R641G								4.29						
	M645V		2.37		4.09	2.80	2.76	5.20			2.57			2.08	20.07
	Y658C	2.50													
	E681K												2.80		
	G682E										3.30				
	L691S											2.24			

Protein	Mutation																
	R703K		4.87			2.04											
	S709G					2.11											
	A717T								2.31								
	K721R										4.53						
	D740N				6.17												
	A750S																3.19
PB1	K11E							10.49									
	T36A																3.32
	M40T																2.49
	Q48K										2.06						
	A91T					3.89											
	G101W										3.43						
	T123A												2.01				
	Y129C										2.65						
	P138T									3.10							
	M171V							2.10									
	M174T									4.14							
	V191I										3.18						
	R192K				3.45												
	K207R												3.19				
	K209R										5.92						
	A221T													4.85			
	I248V												4.47				
	G250R								3.02								
	S261G							2.51									
	V273I					2.21						3.28					
	E277G			4.14													
	N292D											2.84					
	Q313R				7.01												
	P315S								4.91								
	A320T							2.79									
	I322T												2.49				
	V336A											4.64					
	A340S	2.63															
	A349V							2.78									
	D377N								2.22								
	K387E	2.50															
	K387R		3.80														
	L424P												2.06				
	I450R											3.53					
	D464G							5.79									
	I475V			4.10													
	M546V																2.11
	A547V										2.17						
	L598P									3.69							
	P607S							2.39									
	L611S											3.11					
	L628P			4.88													

Protein	Mutation	1	2	3	4	5	6	7	8	9	10	11	12	13	14
	N641S												2.92		
	V644A														3.09
	V660I	31.83	60.36	78.21	33.58	19.61	31.99	19.32	16.78	22.83	15.91	16.95	4.59	13.76	10.24
	C692R			10.62											
	I711V								3.32						
	K745R				2.18										
PA	F4L												2.17		
	F9L												3.40		
	K19R							2.02							
	F46S				2.72							3.63			
	S65P								5.15						
	V100A													5.69	
	K113R							3.37							
	F117S				3.65										
	K139R				2.80										
	H146Q				5.08										
	F150L					3.52						15.63			
	Y161C										2.27				
	T162A		18.72												
	E198K		2.09												
	K245E			25.03											
	L246P			6.31											
	N254D														2.13
	R266Q												4.00		
	D272N		2.66												
	K318R												3.48		
	V323A										2.25				
	N331D		2.12												
	Q340K								3.46						
	D352N														9.70
	K358R												5.97		
	D380Y				2.14										
	D394N														9.01
	R401K											3.36			
	L425F		4.06	63.88				2.32							
	E427K					4.25									
	A433T													3.97	
	M441R										5.50				
	K461R												4.38		
	F480L					3.86									
	M485I												3.17		
	N499T												5.48		
	H510R														12.57
	E538K	5.95								4.13			6.50	6.06	
	E538G		3.34	59.10											
	G578S														12.95
	L589R								2.50						
	E599G											2.08			

Gene	Mutation													
	E623G			2.73										
	S632P							5.48						
	S632F								3.63					
	V645A						2.16							
	F681L		13.38											
	E688K						3.47							
	N696D												4.89	
	L711F												4.70	
HA	Q2R				2.13									
	E14G		2.62											
	V16I										3.79			
	H28Y	8.19												
	V57I													2.31
	D68G						5.25							
	V80A						6.22							
	L101M	3.39												
	A140V									2.29	3.69			
	S142F	4.12												
	I151V												6.59	
	K152E												3.86	
	D155N												6.92	
	I174M										2.01			
	N183D						5.81							
	L190I											8.85		
	T202R										4.49			
	Q207Stop	2.23												
	Q222L		2.10											
	M289V	2.98												
	C302R										2.57			
	Y305F						3.23							
	K310R						3.40							
	V312I	22.45					30.67							
	L322P										2.70			
	R327K									2.45				
	G341E												4.43	
	M374I	15.49		3.76			68.45					2.52	8.03	
	M413I												6.69	
	D419Y							2.70						
	V481A											2.09		
	N483S						2.19							
	I510V												9.45	3.12
NP	I57V	5.04												
	G86E		5.33											
	V105A						2.10							
	R156H												8.67	
	A181V													7.39
	D290N					3.17								
	F338S	3.75												

	S344T												3.47	2.20			
	S402P														5.72		
	V408I														6.53		
NA	T10I			4.89				5.28									
	H36R														3.28		
	N73I	2.16															
	Q78L	3.87															
	S82L									2.48							
	V83A					2.05											
	N104S			2.94													
	L134P											2.51					
	L140P	2.39															
	P162L													11.73			
	A178D										3.61						
	I188V										2.51						
	G236S																15.58
	S237P									2.44							
	V264A											2.29					
	V304I									2.75							
	Y353Stop																2.05
	E372K		11.12														
	D376G													2.24			
	T381N													4.87			
	M396I											5.12					
M1	V15I										5.21						
	A25T													4.37			
	M43I	3.16															
	I59V		2.62														
	L103S			2.03													
	G220R																5.97
M2	S20N											2.36					
	L40S		3.05														
	T65K																2.79
	R77Q																2.31
	S82N								2.17								
NS1	G70D										2.00						
	M79V				2.23	2.86	2.81							15.39			3.88
	D189Y														19.15		
	Q220R				3.87				2.73	2.14				2.35	3.22		2.06
	L222R			2.18													
	Stop231W				2.13	4.64	3.00	2.58	3.44	2.17				17.38	3.74		6.73
	N29S					3.23											
NEP	M31I														19.15		
	R63G				3.87				2.73	2.14				2.35	3.22		2.06
	W65G			2.18													
	E74K													2.23			
	E74G				2.13	4.64	3.00	2.58	3.44	2.17				17.38	3.74		6.73
	K88M			2.28													