Knockdown of PHOX2B in the retrotrapezoid nucleus reduces the central CO$_2$ chemoreflex in

rats.

Silvia Cardani [* 1,2], Tara A. Janes[* 1,2], William Betzner[1§], Silvia Pagliardini[# 1,2,3]

[1]Department of Physiology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton AB

T6G2E1 Canada,

[2]Women and Children's Health Research Institute, University of Alberta, Edmonton AB T6G2E1

Canada.

[3]Neuroscience and Mental Health Institute, University of Alberta, Edmonton AB T6G2E1 Canada.

*SC and TAJ contributed equally to this work

[§] Current address: Departments of Clinical Neurosciences and Community Health Sciences,
Hotchkiss Brain Institute, University of Calgary Cumming School of Medicine, Calgary AB T2N 4N1

[#]Corresponding author: Silvia Pagliardini, Department of Physiology, Faculty of Medicine &

Dentistry, University of Alberta, 3020F Katz Group- Rexall Centre

Email: silviap@ualberta.ca.

Acknowledgments/ Author contributions

Concept and experimental design, S.P. and S.C.; Data collection, S.C., W.B. and S.P.; Data analysis,

S.C. and T.A.J.; Manuscript preparation, S.C., T.A.J. and S.P.

S.C. is supported by the Canadian Lung Association BaO Fellowship. T.A.J. is supported by the

Canadian Institutes for Health Research (CIHR) Postdoctoral Fellowships. S.P. is supported by a

Women and Children's Health Research Institute Innovation Grant and CIHR Project scheme grant.

ABSTRACT

PHOX2B is a transcription factor essential for the development of the autonomic nervous system. Heterozygous mutations in the PHOX2B coding region are responsible for the occurrence of Congenital Central Hypoventilation Syndrome (CCHS), a rare neurological disorder characterised by inadequate chemosensitivity and life-threatening sleep-related hypoventilation. Animal studies suggest that chemoreflex defects are caused in part by the improper development or function of PHOX2B expressing neurons in the retrotrapezoid nucleus (RTN), a central hub for CO_2 chemosensitivity.

Although the function of PHOX2B in rodents during development is well established, its role in the adult respiratory network remains unknown. In this study, we investigated whether reduction in PHOX2B expression in chemosensitive neuromedin-B (NMB) expressing neurons in the RTN altered respiratory function. Four weeks following local RTN injection of a lentiviral vector expressing the short hairpin RNA (shRNA) targeting *Phox2b* mRNA, a reduction of PHOX2B expression was observed in *Nmb* neurons compared to both naïve rats and rats injected with the non-target shRNA. PHOX2B knockdown did not affect breathing in room air or under hypoxia, but ventilation was significantly impaired during hypercapnia. PHOX2B knockdown did not alter *Nmb* expression but reduced the expression of both *Task2* and *Gpr4*, two CO_2 sensors in the RTN. We conclude that PHOX2B in the adult brain has an important role in CO_2 chemoreception and reduced PHOX2B expression in CCHS beyond the developmental period may contribute to the impaired central chemoreflex function.

INTRODUCTION

The Paired Like Homeobox 2B (PHOX2B) is a transcription factor essential for embryonic differentiation and for the maintenance of the neuronal phenotype of different classes of neurons in both the central and peripheral nervous systems (Brunet & Pattyn, 2002; Pattyn et al., 1997, 1999, 2000; Stanke et al., 1999; Tiveron et al., 1996). In recent years, the PHOX2B gene has been extensively investigated, not only for its neurodevelopmental role, but also because its heterozygous mutation leads to Congenital Central Hypoventilation Syndrome (CCHS, OMIM 209880; Amiel et al., 2003), Hirschsprung's disease (HSCR; OMIM 142623; Berry-Kravis et al., 2006; Trochet et al., 2005), and neuroblastoma (Bourdeaut et al., 2005; Van Limpt et al., 2004). CCHS is a rare (1:200,000 births in France (Trang et al., 2005) and 1:148,000 births in Japan (Shimokaze et al., 2015) disorder characterised by a failure to respond to both hypercapnia and hypoxia (Amiel et al., 2003; Di Lascio et al., 2021; Weese-Mayer et al., 2017). The most frequent PHOX2B mutation is an expansion of a polyalanine repeat, ranging from +5 to +13 residues on the C-terminal of PHOX2B, an area that regulates DNA binding affinity and protein solubility (Amiel et al., 2003; Di Lascio et al., 2016; Weese-Mayer et al., 2003). CCHS symptoms present with varying degrees of severity depending on the expansion of the mutation (Berry-Kravis et al., 2006; Di Lascio et al., 2018; Matera, 2004). In most mild cases, patients display normal ventilation during wakefulness, hypoventilation during sleep leading to hypercarbia and hypoxemia, as well as a lack of sensitivity and discomfort to these challenges. In the most severe of cases, hypoventilation may also be present while awake.

Respiratory disturbances resulting from the PHOX2B mutations appear to stem from its widespread expression in respiratory-related neurons both during development and into adulthood. Although PHOX2B is absent in respiratory rhythmogenic neurons of the preBötzinger

complex, it is highly expressed in adult neurons that are responsible for CO_2 and O_2 chemoreflexes

(Kang et al., 2007). Among those are the neurons of the retrotrapezoid nucleus (RTN), the locus

coeruleus, the carotid bodies and the nucleus of the solitary tract that, together with the dorsal

motor nucleus of the vagus nerve and the area postrema, make up to the dorsal vagal complex, a

structure that provides the major integrative centre for the mammalian autonomic nervous

system (Cutsforth-Gregory & Benarroch, 2017; Guyenet et al., 2019; Kang et al., 2007; Zoccal et al.,

2014). Previous work by our group and others specifically investigated the role of PHOX2B-

expressing RTN neurons by either selective lesions or inactivation of neuronal activity of these

neurons (Abbott et al., 2011; Basting et al., 2015; Marina et al., 2010; Nattie & Li, 1994; Souza et

al., 2018, 2023; Janes et al., 2024) and showed that RTN neurons have a key role in central CO_2

chemoreception.

Defects observed in the CO_2-chemoreflex of CCHS transgenic rodent models are primarily caused

by the improper development or function of PHOX2B-expressing neurons in the RTN (Dubreuil et

al., 2008; Ramanantsoa et al., 2011), although anatomo-pathological studies in CCHS patients

suggest that the respiratory impairment may be caused by a more widespread brain dysfunction

(Harper et al., 2015; Nobuta et al., 2015). Transgenic mice born with PHOX2B knockout fail to

develop the RTN, the locus coeruleus (Pattyn et al., 1999) and catecholaminergic groups in A1, A2,

A5 and A7 (Pattyn et al., 2000), as well as neurons of the nucleus of the solitary tract and the

carotid bodies. Mice harbouring heterozygous PHOX2B knockout develop apparently normal but

have depressed responses to hypoxia and hypercapnia (Dauger, 2003) and sleep disordered

breathing in the perinatal period (Durand et al., 2005). Furthermore, the expression of the most

common heterozygous PHOX2B mutation found in CCHS patients (PHOX2B +7Ala expansion) in

mice resulted in a near complete loss of RTN neurons, impaired CO_2 response and consequent

death in the newborn period (Dubreuil et al., 2008; Madani et al., 2021; Ramanantsoa et al., 2011;

Takakura et al., 2008). More selective expression of the mutant PHOX2B in the RTN neurons

allowed for survival in the neonatal period and partial recovery of the CO_2 response in adulthood

(Ramanantsoa et al., 2011).

Multiple potential cellular mechanisms have been implicated in the etiology of CCHS, from loss of

PHOX2B function in development due to heterozygous mutant PHOX2B expression, to gain of

function of the mutated protein that may alter transcriptional activity in neurons, as well as

aggregate formation and premature cell death (Bachetti et al., 2005; Di Lascio et al., 2013;

Dubreuil et al., 2008; Durand et al., 2005; Goridis et al., 2010; Trochet et al., 2005). While its role

during embryonic neurodevelopment is established, PHOX2B is still expressed in selected neuronal

populations in the adult brain (Kang et al., 2007; Stornetta et al., 2006) and its function, with the

exception of the contribution to the maintenance of the noradrenergic phenotype (Fan et al.,

2011), is not yet fully understood. Considering the fundamental role of PHOX2B in CCHS

pathogenesis and its expression in key structures for CO_2 chemoreflex responses, it is essential to

understand how both wild-type and mutant PHOX2B proteins impact CO_2 homeostasis and

ventilation not only across development, but also in adulthood.

In order to have a better understanding of the role of PHOX2B in the CO_2 homeostatic processes

we used a non-replicating lentivirus vector of two short-hairpin RNA (shRNA) clones targeting

selectively *Phox2b* mRNA to knockdown the expression of PHOX2B in the RTN of adult rats and

tested ventilation and chemoreflex responses. In parallel, we also determined whether

knockdown of PHOX2B in adult RTN neurons negatively affected cell survival. Finally, we sought to

provide a mechanistic link between PHOX2B expression and the chemosensitive properties of RTN

neurons, which have been attributed to two proton sensors, the proton-activated G protein-

coupled receptor (GPR4) and the proton-modulated potassium channel (TASK-2) (Gestreau et al.,

2010; Guyenet et al., 2016; Kumar et al., 2015). The expression of these sensors is partially

overlapping in RTN neurons and the genetic deletion of either impairs the central respiratory

chemoreflex with maximal impairment when both sensors are eliminated (Gestreau et al., 2010;

Guyenet et al., 2016; Kumar et al., 2015).

Our results indicate that progressive knockdown of PHOX2B in RTN causes a reduction in the

chemoreflex response that was proportional to the decrease in the fraction of *Nmb*/PHOX2B

expressing RTN neurons. This effect was associated with a reduction in the expression of *Gpr4* and

Task2 mRNA in *Nmb* expressing cells in which PHOX2B was knocked down, suggesting a role for

PHOX2B in the transcription of the proton sensors in RTN neurons.

RESULTS

To directly investigate the role of PHOX2B in RTN neurons, we performed selective knockdown of

this protein in adult rats through local injection of a non-replicating lentiviral vector of two short-

hairpin RNA (shRNA) clones targeting two different sequences of the *Phox2b* mRNA (n=25) or non-

target shRNA as control (NT-shRNA; n=23). Initial experiments utilized 200 nl/injection at each of 4

stereotaxic coordinates (1×10^9 VP/ml; n=14). Four weeks post injection, we assessed ventilatory

parameters in room air, hypercapnia, and hypoxia. Interestingly, a small but significant impairment

in the CO_2 chemoreflex was observed not only in PHOX2B-shRNA but also in NT-shRNA rats.

Specifically, room air breathing was not affected by RTN injection, but allometric V_E during 7% CO_2

was blunted in both NT-shRNA and PHOX2B-shRNA rats when compared to pre-surgical baseline (-

21% and -24%, respectively; Supp. Fig. 1A). Analysis of the air convection requirement ratio (CRR;

V_E/VO_2) confirmed that chemoreflex impairment during 7% CO_2 exposure was of central origin and

not due to lowered metabolism (data not shown). Given that CO_2 chemoreflexes were impaired in

both NT-shRNA and PHOX2B-shRNA rats, and histological analysis showed significant reduction of

Nmb^+/PHOX2B$^+$ RTN cells in NT-shRNA and PHOX2B-shRNA rats (33% and 19% cells remaining,

respectively; Suppl. Fig. 1), we concluded that the volume and titre of the shRNA viral constructs

triggered off-target effects (Mcbride et al., 2008; Van Gestel et al., 2014) that cause cell death in

RTN neurons.

PHOX2B expression is modestly reduced two weeks following infection without respiratory

impairment.

Due to the off-target effects obtained with large volume injections, we reduced injection volume

of both PHOX2B shRNA (n=17) and NT-shRNA viruses (n=17) (4x100nl/injection; 1x10^9 VP/ml).

Two weeks post-surgery we assessed respiratory function during exposure to room air,

hypercapnia and hypoxia. Overall, we observed little change in respiration at this time point.

During room air, as well as in 5% and 7.2% CO_2, frequency (f_R), was reduced in every experimental

group (naïve, NT-shRNA, PHOX2B-shRNA) compared to baseline (Air: p=0.014; 5% CO_2: p< 0.001;

7.2% CO_2 p< 0.001; Fig. 1A) but no effect on treatment was observed, suggesting that these

changes were likely due to the increase in age and body weight during the experiment (naïve:

+32.2%; NT-shRNA: +27.8%; PHOX2B-shRNA: +28.6% body weight). Tidal volume was reduced in

the PHOX2B-shRNA rats at 7.2% CO_2 compared to both naïve rats (naive: 9.6 ± 1.2 ml·kg^{-1};

PHOX2B-shRNA: 8.5 ± 0.9 ml·kg^{-1}; -11%; p= 0.022) and baseline conditions (baseline shRNA: 9.6 ±

1.3 ml·kg^{-1}; -11%; p< 0.001; Fig. 1B) but it was not different from NT-shRNA.

Despite similar changes in body weight across treatment groups, allometric ventilation (V_E ALLO)

was increased in naïve rats compared to the other treatment groups in room air (naïve: 38.4 ± 4.5

ml·min^{-1}; NT-shRNA: 34.7 ± 2.8 ml·min^{-1}; PHOX2B-shRNA: 34.0 ± 4.5 ml·min^{-1}; p=0.005; Fig. 1C), but

no changes in V_E ALLO were observed in hypercapnia (7.2% CO_2) across treatments (naïve : 91.5 ±

13.2 ml·min^{-1}; NT-shRNA: 84.4 ± 13.7 ml·min^{-1}; PHOX2B-shRNA: 80.5 ± 17.2 ml·min^{-1}). Oxygen

consumption (VO_2 ALLO) was not different between experimental conditions or treatment groups

(Fig. 1D), and V_E/VO_2 ALLO indicated that ventilation was matched with metabolism for each

treatment across all gas compositions (naïve: 7.7 ± 1.3; NT-shRNA: 7.2 ± 1.3; PHOX2B-shRNA: 6.9 ±

2.0; Fig. 1E;). These results suggest that despite a reduction in hypercapnic V_T in the PHOX2B-

shRNA rats, overall ventilation was not impaired. Since we observed differences in room air V_E

ALLO between treatment groups, we normalized these data to show the hypercapnic ventilatory

response (HCVR, i.e., absolute change from room air; Fig. 1F). The HCVR was not affected by the

PHOX2B shRNA treatment two weeks post-injection (naïve: 53.1 ± 13.4 ml·min^{-1}; NT-shRNA: 49.6 ±

13.2 ml·min^{-1}; PHOX2B-shRNA: 46.5 ± 14.8 ml·min^{-1}).

To determine the extent of PHOX2B knockdown in RTN neurons we combined RNAScope® and

immunohistochemistry assays to quantify RTN neurons expressing *Nmb* and PHOX2B (Fig. 2B). We

reasoned that by assessing the fraction of RTN neurons that express only *Nmb* (compared to

Nmb$^+$/PHOX2B$^+$), we would be able to determine the level of PHOX2B knockdown in our

experiments. Two weeks post viral injection, we observed a significant decrease (15.2%) in the

total number of *Nmb*$^+$ RTN neurons (i.e., *Nmb*$^+$/PHOX2B$^+$ plus *Nmb*$^+$/ PHOX2B$^-$) in NT-shRNA rats

compared to naïve rats (naïve; 337.6 ± 11.95; n=4; NT-shRNA: 286.0 ± 10.44; n=4; p<0.001; Fig.

2C). The total number of *Nmb*$^+$ RTN neurons in PHOX2B-shRNA ratswas reduced compared to

naïve rats (PHOX2B-ShRNA 218.8 ± 7.8 n=4; -35.2%; p<0.001) and compared to NT-shRNA (-23.5%;

p<0.001).

In order to assess PHOX2B knockdown, we then quantified the number of neurons expressing

either *Nmb* and PHOX2B (*Nmb*$^+$/PHOX2B$^+$) or *Nmb* only (*Nmb*$^+$/PHOX2B$^-$). The number of

Nmb^+/PHOX2B$^+$ neurons in PHOX2B-shRNA rat was reduced by 36.1 % compared to NT-shRNA and

by 46.2% compared to naïve rats (Nmb^+/PHOX2B$^+$ NT-shRNA: 232.3 ± 11.6 vs naïve: 275.8 ± 17.1;

vs PHOX2B-shRNA: 148.3 ± 12.6; p<0.05 and 0.001, respectively; Fig. 2C,D). On the contrary, the

fraction of Nmb^+/PHOX2B$^-$ neurons in PHOX2B-shRNA rat was not different compared to naïve and

NT-shRNA rats (Nmb^+/PHOX2B$^-$ NT-shRNA: 53.7 +- 8.6; vs naïve: 61.8 +-13.9; vs PHOX2B-shRNA:

70.5 +- 8.7 cells; Fig. 2C,E). These results thus indicate that despite modest loss of Nmb^+ neurons in

NT-shRNA and PHOX2B-shRNA treated rats and nearly 50% reduction of Nmb^+/PHOX2B$^+$ in

PHOX2B-shRNA rats, 2 weeks of shRNA viral infection was not sufficient to impair ventilation.

Four weeks of PHOX2B knockdown impairs the CO_2-chemoreflex.

Since the shRNAs integrate into the genome of the infected cells and are continuously produced, it

is reasonable to expect that knockdown efficiency is time-dependent, and an extended infection

time would result in a higher knockdown effect. To test this hypothesis, a subgroup of rats (naïve,

NT-shRNA and PHOX2B-shRNA) were investigated four weeks post-viral injection (Fig. 3A,B).

The total number of RTN neurons in PHOX2B-shRNA rats (Nmb^+/PHOX2B$^+$ plus Nmb^+/PHOX2B$^-$; Fig.

3C) was significantly reduced compared to naïve rats (naïve: 347.8 ± 12 cells, n=4; PHOX2B-shRNA:

249.5 ± 33.9 cells; n=6; -28.3%; p=0.009) but no difference was observed compared to NT-shRNA

rats (294.9 ± 52.8 cells; n=10; -15.4%;). Furthermore, there was no difference in the total number

of Nmb^+ cells in RTN between 2 and 4 weeks post infection for both NT-shRNA (week 2: 286 ±

10.4; week 4: 294.9 ± 52.8, p=0.783) and PHOX2B-shRNA rats (week 2: 218.8 ± 7.8; week 4: 249.5

± 33,9, p=0.118), suggesting that progressive PHOX2B knockdown was not accompanied by

increased cell death.

The number of PHOX2B expressing Nmb^+ cells in the RTN was significantly reduced in PHOX2B-

shRNA rats by 67.0% and 60.7% compared to naïve and NT-shRNA rats, respectively (naïve: 284.5 ±

19.8; NT-shRNA: 238.7 ± 56; PHOX2B-shRNA: 93.8± 11.9 cells, p<0.001; Fig. 3D). Moreover, a

significant increase in the fraction of Nmb^+/PHOX2B$^-$ was observed with PHOX2B knockdown

(PHOX2B-shRNA: 155.7 ± 22.7; naïve: 63.2 ± 12cells; + 146.1% increase vs naïve; NT-shRNA: 56.2 ±

13.8cells; + 177% increase vs NT-shRNA, p<0.001; Fig. 3E).

The respiratory function prior to histological examination (4 weeks post-viral injection) showed

significant differences in PHOX2B-shRNA rats compared to both baseline, naïve and NT-shRNA

rats. Similar to what we reported at 2 weeks post-surgery, f_R at 4 weeks was significantly lower in

every treatment group compared to baseline recordings (Air: p=0.001; 5% CO_2: p< 0.001; 7.2% CO_2

p< 0.001; Fig. 4A).

During room air, V_T was modestly reduced in naïve rats compared to baseline (baseline: 5.9 ± 0.5

ml·kg^{-1}; week 4: 5.2 ± 0.6 ml·kg$^{-1;}$ -12%; p=0.001; Fig. 4B). Moreover, V_T in PHOX2B-shRNA rats was

reduced compared to NT-shRNA (NT-shRNA: 5.6 ± 0.4 ml·kg^{-1}; PHOX2B-shRNA: 5.3 ± 0.4 ml·kg^{-1}; -

10%; p=0.043) and to the baseline (6.1 ± 0.7 ml·kg^{-1}; -18%; p=0.002). No significant changes were

observed at 5% CO_2 although a decrease in V_T occurred at 7.2% CO_2 in PHOX2B-shRNA rats

compared to baseline pre-surgery (baseline: 7.7 ± 1.2 ml·kg^{-1}; PHOX2B-shRNA: 6.7 ± 0.6 ml·kg^{-1}; -

31%; p<0.001), naïve rats (7.9 ± 0.9 ml·kg^{-1}; -26%; p<0.001), and NT-shRNA rats (8.1 ± 0.7 ml·kg^{-1}; -

23%; p=0.002).

Decreased V_T observed for PHOX2B-shRNA led to a ventilation (V_E ALLO) impairment during

exposure to hypercapnia (7.2% CO_2, PHOX2B-shRNA vs. NT-shRNA: -17%; p=0.021; PHOX2B-shRNA

vs. naïve; -21% ; p=0.007; PHOX2B-shRNA vs. baseline, -18% p=0.002; Fig. 4C). Since O_2-

consumption (VO_2 ALLO) did not change (data not shown) a significant reduction in V_E/VO_2 ALLO

confirmed alveolar hypoventilation in PHOX2B-shRNA animals compared to NT-shRNA rats both at

5% (NT-shRNA: 5.8 ± 0.5; PHOX2B-shRNA: 4.8 ± 0.4; -16%; p=0.023) and 7.2% CO_2 (NT-shRNA: 7.9 ±

1.2; PHOX2B-shRNA: 6.1 ± 0.8; -24%; p=0.004; Fig. 4D). Moreover, the HCVR (normalized to room

air; Fig. 4E) at 7.2% CO_2 was lower in PHOX2B-shRNA (40.3 ± 9.4 ml·min^{-1}) rats compared to the

pre-surgery baseline (58.1 ± 8.9 ml·min^{-1}; -31%; p=0.015), naïve rats (63.6 ± 14.3 ml·min^{-1}; -37%;

p=0.008) and NT-shRNA rats (56.0 ± 10.4 ml·min^{-1}; -28%; p=0.016). Analysis of individual rats in

each treatment group over time showed that the HCVR was only consistently impaired in PHOX2B-

shRNA rats between weeks 2 and 4 (Fig. 1F; p=0.007).

Given that previous studies have proposed a role for the RTN in the hypoxic chemoreflex (Barna et

al., 2016; Basting et al., 2015; Gourine et al., 2005; Wickström et al., 2004), and we observed

impairment of the CO_2-chemoreflex at 4 weeks post-infection, we investigated whether PHOX2B

knockdown would also affect respiratory response to hypoxia (10% O_2). The 4 week V_E ALLO was

increased relative to baseline values in naïve (baseline: 62.3 ± 5.2 ml·min^{-1}; naïve: 71.0 ± 10.1

ml·min^{-1}; +14%), NT-shRNA (baseline: 71.8 ± 11.1 ml·min^{-1}; NT-shRNA: 75.8 ± 12.8 ml·min^{-1}; +6%)

and PHOX2B-shRNA rats (baseline: 65.9 ± 9.8 ml·min^{-1}; PHOX2B-shRNA: 79.3 ± 6.4 ml·min^{-1}; +29%;

p<0.001). However, analysis of the convective requirement ratio (V_E/VO_2 ALLO) indicates that

changes in hypoxic ventilation were due to metabolic adjustments that occurred in all treatment

groups (VE/VO_2 ALLO naive: 6.5 ± 1.3; NT-shRNA: 6.7 ± 1.7; PHOX2B-shRNA: 6.6 ± 0.6), suggesting

that the hypoxic ventilatory response was not affected by PHOX2B knockdown.

PHOX2B knockdown does not extend to C1 catecholaminergic neurons and does not affect

mRNA expression of *Nmb* in RTN neurons

Our data shows that PHOX2B knockdown specifically impairs breathing during hypercapnia. To

exclude that the observed impairment was due to unintended knockdown of PHOX2B in adjacent

areas of the brain, we quantified the total number of TH^+/$PHOX2B^+$ catecholaminergic C1 neurons

located caudal to the RTN in PHOX2B-shRNA rats compared to naïve and NT-shRNA (Fig. 5A). No

differences were observed in either the TH^+ cell numbers or in the PHOX2B expression between

the 3 treatment groups (naïve: 378.3 ± 23.81; NT-shRNA: 391.7 ± 33.08; PHOX2B-shRNA: 370.2 ±

33.04 cells).

NMB is currently considered the most selective anatomical marker for chemosensitive RTN

neurons (Shi et al., 2017) and was used in this study to identify RTN location, cell numbers and

relative PHOX2B expression. Since changes in the level of the transcription factor PHOX2B could

regulate the expression of its target genes, we investigated whether NMB is affected by shRNA

knockdown and assessed *Nmb* expression in PHOX2B-shRNA rats compared to both naïve and NT-

shRNA rats (Fig 5B). *Nmb* mRNA expression at the level of RTN was calculated as total fluorescence

intensity (CTCF) ratio relative to *Nmb* expression at the level of NTS, which served as an internal

control (Fig. 5B). We observed no changes in relative fluorescence of *Nmb* in RTN neurons across

the 3 groups (p=0.608), suggesting that NMB expression is not affected by PHOX2B knockdown.

To better analyse the association between PHOX2B knockdown in RTN neurons and the

attenuation of the HCVR, we evaluated the correlation between the number of Nmb^+/PHOX2B$^+$ or

Nmb$^+$/PHOX2B$^-$ cells in the RTN and the resulting HCVR using linear regression (Fig. 5C,D). Our

results indicate that there is a significant correlation between HCVR and number of

Nmb$^+$/PHOX2B$^+$ ($r^2 = 0.739$ p<0.001; Fig. 5C), and Nmb$^+$/PHOX2B$^-$ ($r^2 = 0.482$ p<0.001 Fig. 5D) cells

suggesting that the number of PHOX2B expressing cells in the RTN is a good predictor of the

chemoreflex response. Moreover, the nature of the correlation between PHOX2B$^-$ cells and HCVR

is the opposite of PHOX2B$^+$, another important indicator that PHOX2B protein contributes to the

CO_2 sensing and consequently, the reduction of PHOX2B protein impairs the CO_2-chemoreflex.

Gpr4 and Task2 mRNAs expression is affected by PHOX2B Knockdown

We next examined the mRNA expression of *Gpr4* and *Task2*, two important pH sensors for the

central respiratory chemoreflex response in RTN neurons (Gestreau et al., 2010; Guyenet et al.,

2016; Kumar et al., 2015). As previously reported, the two pH sensors are expressed in the NMB

RTN neurons in partially overlapping populations of chemosensory neurons (Gestreau et al., 2010;

Kumar et al., 2015). In naïve rats 91.6 ± 11.2 % of NMB cells were *Gpr4* positive, and 76.6 ± 14.8 %

of NMB cells were also *Task2* positive (n=4 rats). These values were not significantly different in

either NT-shRNA or PHOX2B shRNA rats (NT-shRNA: 90.0 ± 15.7% *Nmb$^+$/Gpr4$^+$* and 76.2 ± 21.8%

Nmb$^+$/Gpr4$^+$, n=10 rats; PHOX2B-shRNA: 89.4 ± 18.3% *Nmb$^+$/Gpr4$^+$* and 75.6 ± 27.8% *Nmb$^+$/Gpr4$^+$*,

n=6 rats).

To further investigate whether the expression of these two sensors was affected by the reduction

in PHOX2B protein we quantified the total fluorescence intensity (CTCF) for their mRNA in *Nmb$^+$*

cells. Total CFCT for *Gpr4* and *Task2* in PHOX2B-shRNA rats was significantly reduced compared to

naïve and NT-shRNA rats (*Gpr4*: -34.4 ± 4.79% vs naïve, -27.9 ± 5.71% vs NT-shRNA p<0.001; *Task2*:

-39.0 ± 3.02% vs naïve, -34 ± 2.62% vs NT-shRNA p<0.001; Fig. 6B,D). To better understand

whether this reduction was ascribed to the specific loss of PHOX2B expression, we compared the

mRNA expression levels between *Nmb$^+$*/PHOX2B$^-$ and *Nmb$^+$*/PHOX2B$^+$ neurons in PHOX2B-shRNA

rats. Both *Gpr4* and *Task2* mRNA expression was reduced in *Nmb$^+$*/PHOX2B$^-$ cells compared to

Nmb$^+$/PHOX2B$^+$ (*Gpr4*: -63.8 ± 3.9%; p= 0.022; *Task2*: -63.2 ± 2.77%; p= 0.029; Fig. 6C,E). Our data

suggest that PHOX2B knockdown not only causes a reduction in the hypercapnic respiratory

response but also a reduction in the expression of *Gpr4* and *Task2* mRNAs.

DISCUSSION

The role of the transcription factor PHOX2B in the adult brain is still not fully understood. In this study we reduced the expression of PHOX2B in the key chemosensory structure of the RTN with a selective viral shRNA approach to test whether a knockdown of PHOX2B expression in these neurons negatively impacts ventilation. Two weeks after viral infection we observed a modest reduction of PHOX2B/NMB expressing neurons in the RTN but no significant changes in basal V_E or in the CO_2 chemoreflex. Four weeks after shRNA infection we observed a further reduction in the expression of PHOX2B in RTN neurons and a significant reduction of the CO_2 chemoreflex, while ventilation in normoxia or hypoxia was unaffected. Furthermore, the level of expression of *Gpr4* and *Task2* mRNA, the two proton sensors responsible for the RTN neurons chemosensory properties were significantly reduced in RTN neurons that lacked PHOX2B protein expression, whereas *Nmb* mRNA expression level was unaffected by PHOX2B knockdown. These results suggest that PHOX2B reduction affects RTN neurons transcriptional activity and decreases the CO_2 response, possibly by reducing the expression of key proton sensors in the RTN.

The role of PHOX2B in development and in the adult brain

PHOX2B is a transcription factor with an important role in the development and differentiation of neuronal structures in both the central and peripheral nervous system. Genetic loss of PHOX2B expression in mice shows that PHOX2B is essential for the correct development of all autonomic ganglia in sympathetic, parasympathetic, and enteric nervous system and the distal ganglia of the facial (VII), glossopharyngeal (IX), and vagus (X) cranial nerves (Pattyn et al., 1999), in addition to carotid bodies and the solitary tract nucleus (Dauger, 2003). Furthermore, PHOX2B is involved in the formation of all branchial and visceral hindbrain motor neurons and its absence during embryonic period impairs the development of motoneurons of the facial, trigeminal, ambiguus

and dorsal motor nucleus of the vagal nerve. PHOX2B is also necessary for the specification and

differentiation of the noradrenergic phenotype in multiple centres in the brain (Brunet & Pattyn,

2002; Pattyn et al., 1999, 2000a).

Interestingly, the expression of PHOX2B persists into adulthood in selected neuronal populations,

such as the ones involved in peripheral and central chemoreception, i.e., the carotid bodies, the

solitary tract nucleus and the RTN, in addition to neurons in the dorsal motor nucleus of the vagus,

the area postrema, the nucleus ambiguus, C1 catecholaminergic neurons and other scattered

neurons throughout the brainstem (Stornetta et al., 2006). Such widespread expression within the

adult brain suggests an important transcriptional role in neuronal survival and/or function. With

the exception of its role in maintaining the noradrenergic phenotype, its function in the postnatal

brain is unknown, in part because a full investigation of its transcriptional activity in vivo has not

yet been performed.

Heterozygote polyalanine expansion mutations in the exon 3 of *PHOX2B* are responsible for the

majority of cases of CCHS, a rare genetic disease that affect the autonomic nervous system and

central CO_2 chemoreflexes (Amiel et al., 2003; Di Lascio et al., 2021; Weese-Mayer et al., 2017).

While studies in transgenic mice expressing the mutated PHOX2B protein suggest that the RTN

may not form or be severely impaired (Dubreuil et al., 2008, 2009; Ramanantsoa et al., 2011), the

function of the wild-type protein in these neurons has never been tested in adulthood. Thus, in

order to determine whether PHOX2B is necessary for the survival and/or chemosensory function

of adult PHOX2B$^+$ RTN neurons in vivo, we used a shRNA viral approach to progressively

knockdown the PHOX2B protein in these neurons.

Modest reduction of PHOX2B in RTN does not impair ventilation

We identified RTN neurons by the mRNA expression of the neuropeptide, NMB (Shi et al., 2017)

and observed a reduced number in both total *Nmb*$^+$ and *Nmb*$^+$/PHOX2B$^+$ neurons in shRNA rats

compared to naïve and NT-shRNA rats two weeks after viral infection. Interestingly, despite the

small reduction in total Nmb^+ RTN neurons in both NT-shRNA and PHOX2B-shRNA injected rats,

possibly due to some off-target effects due to activation of innate immunity or saturation of the

microRNA pathway (Mcbride et al., 2008; Van Gestel et al., 2014), no significant impairment in

ventilation was observed. Nonetheless, the fraction of Nmb^+ cells expressing PHOX2B was reduced

indicating that shRNA knockdown occurred in a fraction of RTN neurons. Even though PHOX2B

may be critical for chemosensory function in RTN neurons, the lack of significant ventilatory

impairment with a modest reduction of PHOX2B expressing neurons is not surprising. In our

previous study (and the ones of others), in which RTN chemosensory neurons were lesioned with

Substance P saporin toxin, CO_2-chemoreflex impairment became significant (~30% V_E Allo

reduction vs. pre-surgical baseline) only when the sum of Nmb^+/PHOX2B$^+$ and Nmb^+/PHOX2B$^-$

neurons was reduced by ~63% compared to naïve controls (medium lesion range: 50%-81% loss of

Nmb^+ neurons; Janes et al., 2024; Souza et al., 2023), therefore, even though loss of PHOX2B

severely altered the chemosensitive function of NMB cells, we would not expect significant

changes in ventilation with a <50% RTN cell loss.

Four weeks of PHOX2B knockdown impairs central CO_2 chemoreception

375
Four weeks post- shRNA viral injection, the fraction of Nmb^+ cells expressing PHOX2B was further

reduced by 67% compared to naïve and by 61% compared to NT-shRNA rats. PHOX2B knockdown

was also restricted to RTN neurons, as more caudal C1 TH$^+$ neurons and the adjacent FN

motoneurons did not show any change in number or PHOX2B expression. Although we did not

specifically quantify the relative mRNA expression change in intracellular $Phox2b$ levels within

infected neurons, we measured the success of PHOX2B knockdown by assessing the overall

change in the number of Nmb^+ cells that had no detectable levels of PHOX2B protein in the nuclei.

Thus, it is possible that this method led us to underestimate the success of the overall PHOX2B

shRNA knockdown in the RTN.

In PHOX2B-shRNA rats we did not observe any respiratory function changes in room air or in

hypoxia, similar to RTN lesioning studies (Janes et al., 2024; Souza et al., 2023). When tested in

hypercapnia though, PHOX2B-shRNA rats displayed a reduced HCVR compared to both baseline

(i.e., pre-surgery), naïve and NT-shRNA rats. The impairment of V_E Allo was primarily a result of

blunted V_T, and analysis of the convective exchange ratio (i.e. V_E/VO_2 ALLO) confirmed

hypoventilation in 5% and 7.2% CO_2 only for PHOX2B-shRNA rats, suggesting a specific impairment

of chemosensitive properties in RTN neurons with PHOX2B knockdown. These results are in line

with previous studies in which medium RTN lesions (i.e. loss of >60 % Nmb^+ cells) elicited by

Saporin toxin injection resulted in a significant CO_2-chemoreflex impairment (Janes et al., 2024;

Souza et al.,2018).

Changes in the CO_2 response following 4 weeks of PHOX2B-shRNA treatment could occur because

the reduced expression of the transcription factor PHOX2B causes changes in the transcriptional

machinery of RTN neurons and alters transcription of CO_2/pH sensing proteins or other key

elements for their chemosensitive function. These transcriptional changes could also affect

neuronal survival. In fact, a small but significant loss of NMB expressing RTN neurons may

contribute to the reduction in the CO_2 response, although multiple studies have demonstrated

that significant cell loss must occur in the RTN in order to show a respiratory function impairment

(Janes et al., 2024; Souza et al.,2018; Nattie & Li, 2002; Takakura et al., 2008, 2014).

In our experiments, we observed a 35% reduction in total Nmb^+ cells compared to naïve rats and

23% compared to NT-shRNA, possibly due to cell death within the first 14 days post-infection.

Interestingly the loss of Nmb^+ cells did not increase with time (although the number of PHOX2B$^+$

neurons decreased) and the central chemoreflex at 2 weeks post-surgery was not impaired,

suggesting that: i) the observed reduction in overall RTN cell number is not responsible for the CO_2

chemoreflex impairment; ii) PHOX2B expression is most likely not necessary for neuronal survival

of RTN neurons; iii) a large reduction in RTN PHOX2B$^+$ neurons is necessary to impair the HCVR.

The interpretation that PHOX2B expression contributes positively to the chemoreflex is further

supported by our linear regression analysis showing that *Nmb*$^+$/PHOX2B$^+$ cell number was the best

predictor of the HCVR (r^2 = 0.739). The initial cell loss, which we observed also in NT-shRNA, and

more prominently with larger injection volume of virus, may be attributed to some inherent cell

death associated with the surgical procedure or, more likely, with the potential off-target toxic

effects associated with shRNA procedures (Mcbride et al., 2008; Van Gestel et al., 2014).

416
PHOX2B knockdown reduces expression of proton sensors in the RTN

Current theories on central respiratory chemosensitivity postulate that changes in CO_2/pH in the

RTN are detected through two pH sensors, the proton-activated receptor GPR4 and the pH-

sensitive K$^+$ channel TASK2 that are expressed in partially overlapping populations of NMB

expressing RTN neurons (Kumar et al., 2015). Because it is not known whether PHOX2B has any

effect on the expression of TASK2, GPR4 or even the expression of the neuropeptide NMB used to

anatomically identify RTN neurons, we determined their mRNA levels within the RTN (calculated

as CFTC) and observed no changes in relative fluorescence of *Nmb*$^+$ mRNA in RTN neurons relative

to *Nmb* expression in unrelated cells in the NTS, suggesting that NMB is not affected by PHOX2B

shRNA knockdown and that *Nmb* is most likely not a target gene of PHOX2B in adult RTN neurons.

Furthermore, we determined the mRNA expression levels of *Task2* and *Gpr4* in both

Nmb$^+$/PHOX2B$^+$ and Nmb$^+$/PHOX2B$^-$ neurons in shRNA rats. Our results indicate that the total CTCF

for *Gpr4* and *Task2* in PHOX2B-shRNA rats was reduced compared to naïve and NT-shRNA group.

Furthermore, there was a significant reduction of both *Gpr4* and *Task2* in *Nmb*$^+$/PHOX2B$^-$ cells

compared to *Nmb*$^+$/PHOX2B$^+$ cells. Because no good antibodies are currently available to detect

protein levels of either NMB, GPR4 or TASK2, our results are only based on changes in mRNA

levels, thus we can only speculate that a reduction in *Gpr4* and *Task2* mRNA would translate in a

reduction in the protein levels and consequent reduction of RTN neurons chemosensitive

properties. Direct electrophysiological recordings in RTN neurons will be able to address the

changes in CO_2/pH sensitivity of RTN neurons at cellular level.

PHOX2B knockdown may impair CO_2-sensing through additional transcriptional targets

Since loss of PHOX2B reduces expression of TH and dopamine beta hydroxylase enzymes in

catecholaminergic neurons *in vivo* and *in vitro* (Fan et al., 2011), it is possible that in NMB[+] RTN

neurons, reduction of PHOX2B alters the transcriptional machinery of other proteins (in addition

to GPR4 and TASK2) and impairs their function in central chemoreception. For example, a change

in the expression of enzymes, neurotransmitters, receptors, and ion channels in RTN neurons

could affect the excitatory signal transmission to preBötzinger Complex and the respiratory

network specifically during CO_2 challenges (Guyenet et al., 2016). Further studies will be needed to

address and identify additional transcriptional changes induced by PHOX2B knockdown *in vivo.*

An interesting observation from our histological data was the reduction in the overall number of

Nmb[+] cells at two- and four-weeks following shRNA viral injections. Although some cell death may

be associated with either surgical procedures or off-target effects associated with the shRNA

approach (Mcbride et al., 2008; Van Gestel et al., 2014), it is intriguing to speculate that loss of

PHOX2B expression may have some effects on viability of RTN neurons *in vivo*. Even though this is

a possibility, especially with an even more severe knockdown, we did observe a large proportion

of *Nmb*[+] cells devoid of PHOX2B protein expression, suggesting that absence of PHOX2B is

compatible with neuronal survival, at least in our experimental time frame.

455
Relevance to CCHS and its pathogenic mechanisms

Our results contribute to further our understanding on potential pathogenic mechanisms in CCHS

(Di Lascio et al., 2018). Homozygous knockout mice for PHOX2B die during gestation (Pattyn et al.,

1999, 2000) demonstrating a key role of this transcription factor in cellular proliferation, migration

and differentiation during embryonic development, whereas heterozygotes for PHOX2B survive

birth and live into adulthood, but display apneas (Durand et al., 2005) and impairment of the

hypoxic and hypercapnic ventilatory responses in the neonatal period (Dauger, 2003). Here we

show that, independent of the PHOX2B role during development, PHOX2B is still required to

maintain proper CO_2 chemoreflex responses in the adult brain.

Heterologous expression of the mutant PHOX2B alters development of RTN in mice and impairs

their chemoreflex response (Dubreuil et al., 2008, 2009). The effects of the expression of the

mutant PHOX2B on wild-type PHOX2B and its transcriptional activity is not known *in vivo* yet,

although *in vitro* data indicates that the mutated PHOX2B protein alters wild-type PHOX2B

conformation and its ability to form homo- and hetero-dimers (Di Lascio et al., 2016; Trochet et

al., 2005, 2008), its affinity for DNA and coactivators, as well as its degradation rate (Nagashimada

et al., 2012; Wu et al., 2009). It has also been shown that mutant PHOX2B causes the formation of

cytoplasmic aggregates (Bachetti et al., 2005; Trochet et al., 2005, 2008) and fibrils *in vitro* (Pirone

et al., 2019), and dysregulates the transcriptional function of wild-type PHOX2B (Di Lascio et al.,

2013; Parodi et al., 2012), thus changing expression of important target genes such as DBH,

PHOX2A and TLX2 (Bachetti et al., 2005; Di Lascio et al., 2013; Trochet et al., 2005). Based on this

evidence, different CCHS pathogenic mechanisms have been proposed (e.g., PHOX2B loss-of-

function, dominant-negative or toxic function of the mutant protein) including gene and cell

specific transcriptional dysregulation (Di Lascio et al., 2018, 2021). Interestingly, only a handful of

PHOX2B target genes have been identified and the function of PHOX2B and its mutated forms in

respiratory control and CCHS pathogenesis is still under investigation.

Our data suggest that, in addition to potential effect of mutated PHOX2B on development and

cellular function in CCHS pathogenesis, the expression of wild type PHOX2B has an important role

in respiratory control that extends the developmental period and its reduction in CCHS may

contribute to the respiratory impairment in this disorder.

MATERIALS AND METHODS

Experiments were performed using male Sprague–Dawley rats (starting weight range 250-350g)

born in the University of Alberta animal facility to pregnant dams obtained from Charles River

(Senneville, QC). Rats were housed at the University of Alberta Health Sciences Animal Housing

Facility and maintained on a 12-hr dark/light cycle with food and water available *ad libitum*.

Handling and experimental procedures were approved by the Health Science Animal Policy and

Welfare Committee at the University of Alberta (AUP#461) and performed in accordance with

guidelines established by the Canadian Council on Animal Care.

ShRNA viral injection in the retrotrapezoid nucleus

In order to knockdown Phox2b expression at the level of RTN neurons, we used a mix of two

shRNA clones targeting two different sequences of the Phox2b mRNA carried by a non-replicating

lentivirus vector ($1x10^9$ VP/ml; TRCN000041283: GCCTTAGTGAAGAGCAGTATG and

TRCN0000096437: CCTCTGCCTACGAGTCCTGTA; Sigma Aldrich, St. Louis, MA, USA). The shRNAs

were designed against the Phox2b mouse sequence (Ref Seq NM_008888) which shares 100%

homology with the Phox2b rat sequence (Ref Seq XM_008770167).

Metacam analgesic was administered 1 hour prior to surgery (2mg/kg) to reduce post-surgical

pain. Rats were anesthetized with a ketamine/xylazine mixture (100mg/kg + 10mg/kg,

respectively; i.p.); the anaesthetic level was assessed by lack of paw pinch reflex and maintenance

of a regular breathing rate. Additional anaesthesia was administered as needed. The rat was

positioned on a stereotactic apparatus (Kopf Instruments, Tujunga, CA, USA) and a total of four

microinjections (200 or 100 nL per injection; two rostro-caudally aligned injections per side) were

made lateral to the midline and ventral to the facial motor nucleus under aseptic conditions.

Coordinates were as follows in mm from obex (medio-lateral/rostro-caudal/dorso-ventral): ±1.8,

+2.0,-3.5; ±1.8,+2.4,-3.6. Injections were made using a glass microelectrode (30 µm diameter tip,

Drummond Scientific, PA, USA). Twenty-five rats (8 rats 200nl/injection; 17 rats 100nl/injection)

underwent shRNA viral injection (PHOX2B-shRNA) and 23 rats (6 rats 200nl/injection; 17 rats

100nl/injection) received a non-target shRNA viral injection (NT-shRNA; SHC016:

MISSION® pLKO.1-puro non-Target shRNA Control Plasmid DNA; Sigma Aldrich, St. Louis, MA, USA)

to determine any effects of surgery alone on respiratory behaviour. Eight naïve rats, receiving no

surgery and maintained in the same housing conditions, were used as additional controls for both

respiratory function and anatomical experiments. Following each injection, the glass electrodes

were left in place for 3-5 minutes to minimize backflow of virus up the electrode track. At the end

of the surgery, the incision was sutured, and rats received local anaesthetic bupivacaine (0.1 mL,

s.c.) and were treated with metacam analgesic for 72 hours. Rats recovered for 7 days before the

next respiratory function testing.

Data acquisition and analysis of respiratory measurements

Rats were habituated to whole-body plethysmography chambers (Buxco, 5L) once, 3-4 days before

baseline recordings. On the day of the experiment, rats were placed in the chamber and testing

commenced once the animals were quietly awake (typically 20-30 mins). Gas mixtures were

delivered at 1.5 L/min using a GSM-3 (CWE Inc., Ardmore, PA, USA) and monitored using a Gas

Analyzer (AD Instruments, Sydney, Australia). Ventilatory parameters were measured during

exposure to different air compositions for 8-15 min each. Room air: $21\%O_2 + 0\%CO_2$ (balanced

with N_2); Normoxic Hypercapnia: $21\%O_2 + 5\%CO_2$; $21\%O_2 + 7.2\%CO_2$; Normocapnic Hypoxia:

$10.6\%O_2 + 0\% CO_2$.

As previously described (Cardani et al., 2022; Janes et al., 2024) respiratory data were analysed

from rats during quiet wakefulness in the last 5 min period of each gas composition using the

barometric method (open-flow system) (Cardani et al., 2022; Janes et al., 2024; Mortola &

Frappell, 1998; Seifert et al., 2000). Raw pressure signals were acquired with a Validyne

differential pressure transducer connected to a CD15 carrier demodulator (Validyne Engineering,

Northridge, CA, USA) and digitized using a Powerlab 8/35 (AD Instruments, Sydney, Australia).

Analysis was done using Labchart 8 (v8.1.19, AD Instruments, Colorado Springs, CO, USA) to

determine tidal volume (V_T) and breathing frequency (f_R). The amplitude of the pressure signal was

converted to V_T ($ml \cdot kg^{-1}$) using the equations of Drorbaug and Fenn 1955 (Drorbaugh & Fenn,

1955) and calibrated against 1 mL of dry air injected into the empty chamber using a rodent

ventilator (Harvard Rodent Ventilator Model 683, Holliston, MA, USA f = 50, 75, 100, 150

$beats \cdot min^{-1}$). Humidity and chamber temperature were monitored using a RH-300 water vapour

analyzer (Sable Systems, Las Vegas, NV, USA) and a HPR Plus Handheld PIT Tag reader (Biomark,

Boise, ID, USA) respectively, depending on the setup. Rectal temperature was read through the

temperature probes of a Homeothermic Monitor (507222F, Harvard Apparatus, Holliston, MA,

USA). Minute ventilation ($\dot{V}_E$) was calculated as f_R x V_T and expressed as $ml \cdot min^{-1} \cdot kg^{-1}$. Rats gained,

on average, 160 ± 4g over the experimental paradigm (naive: 158 ± 19g ; NT-shRNA: 202 ± 25g ;

PHOX2B-shRNA: 133 ± 7g); we therefore applied an allometric correction to $\dot{V}_E$ to account for non-

linear effects of weight gain (V_E ALLO) (Mortola et al., 1994).

Metabolic parameters VO_2 and V_E/VO_2 were calculated by pull-mode indirect calorimetry and

allometerically scaled to be consistent with ventilation (Mortola et al., 1994). The % composition

of dry gas flowing into, and out of the recording chamber was measured by using an AD

Instruments Gas Analyzer (Colorado Springs, CO) and applying the equations of Depocas and Hart

(Depocas F, Hart JS, 1957) and Lighton (Lighton, 2021).

Weight measurements were taken prior to any recording. Rats breathing function was measured

the week prior to surgery (baseline recording) and weekly following shRNA injections to determine

the time course of ventilatory impairment.

Respiratory data were compared for naive, NT-shRNA and PHOX2B-shRNA pre- and post-injection

using a mixed factor ANOVA (independent factor = treatment, repeated measures factor = pre-

and post-lesion; Fig.1,4). Post-hoc analysis used the Bonferroni correction factor, with $p < 0.05$

considered to be significant. Ventilatory data are reported as mean ± standard deviation (SD).

In Situ hybridization (RNAScope) and immunofluorescence

Either 2 or 4-weeks after viral injection, rats were transcardially perfused with saline 0.9% NaCl

followed by 4% paraformaldehyde (PFA) and the brains were post-fixed in 4% PFA overnight and

cryoprotected in 30% sucrose in 1X Phosphate Buffered Saline (PBS). Brains were then frozen in

O.C.T compound (Fisher Scientific) and sectioned on a cryostat (MODEL CM1950, Leica

Biosystems, Buffalo grove, IL, USA) at 30 μm and stored in cryoprotectant buffer at -20°C until

processing.   Sections were mounted on slides for combined RNAScope® in situ hybridization

(Advanced Cell Diagnostics-ACD Bio, Newark CA, USA) and immunofluorescence assay and

processed as previously detailed (Biancardi et al., 2021; Cardani et al., 2022). The mRNA

expression for *Neuromedin B* (*Nmb)* was used as marker of RTN CO_2-sensing neurons (Shi et al.,

2017). Slides were incubated with probes for *Nmb* (Rn-NMB-C2 #494791-C2, ACDBio, Newark, CA,

USA), *Phox2b* (Rn-Phox2b-O1-C1 #1064121-C1, ACDBio), G-protein-coupled receptor 4 (*Gpr4*)

(Mm-Gpr4-C1, #427941, ACDBio), and potassium channel, subfamily K, member 5 (*Kcnk5* or *Task-2*) (Mm-Kcnk5-C3, #427951-C3, ACDBio) for 2 hr at 40°C. Gpr4 and Kcnk5 probes are design against

the mouse mRNA sequence (Ref Seq NM_175668.4 and NM_021542.4, respectively). However,

the alignment of the mouse mRNA sequences with those of rat (GPR4: Ref Seq NM_001025680.1;

TASK2: Ref Seq NM_1039516.2) showed an identity of 94% and 96%, respectively, at the level of

the target region (Base Pairs 1030-150) (BLAST ® program, National Library of Medicine, National

Center for Biotechnology Information, NIH). In parallel, two sections/rat were treated with

positive (low copy housekeeping gene), and negative (non-specific bacterial gene) control probes

provided by ACDBio. Finally, slides were processed using the RNAScope Multiplex Fluorescent

Assay kit V2 (ACDBio) according to the manufacturer's instructions. The probes were visualized

using Opal 520 and Opal 570 reagent (1:500 and 1:1000, respectively; PerkinElmer, Woodbridge,

ON, CA). PHOX2B immunoreactivity was detected using mouse monoclonal PHOX2B antibody (B-11: sc-376997, 1:100, Santa Cruz Biotechnology) incubated overnight in 0.3% TritonX-100, 1%NDS

in PBS followed by donkey CY5-conjugated anti-mouse IgG (1:200; Jackson Immuno Research

Laboratories Inc) in PBS + 1% NDS for 2 hours. Slides were then washed in PBS and cover-slipped

with Fluorosave mounting media (EMD Millipore).

We also performed staining for tyrosine hydroxylase (TH) to identify and quantify C1 cells

(TH[+]/PHOX2B[+]) following shRNA injection. Briefly, perfusion and tissue fixation/freezing were done

as described above. Floating sections were stained with rabbit anti TH antibody (AB152,1:1000,

Millipore SIGMA, USA), and mouse anti PHOX2B (B-11: sc-376997, 1:100, Santa Cruz

Biotechnology). Antibodies were visualized using donkey CY3 (TH) and CY5 (PHOX2B) conjugated

IgG (1:200; Jackson Immuno Research Laboratories Inc). The sections were washed in PBS and

mounted on slides and cover-slipped with Fluorosave mounting media.

Cell counting, imaging and data analysis.

To quantify the number of neurons Nmb^+/PHOX2B$^-$ *and* Nmb^+/PHOX2B$^+$ within the RTN region, we

analysed one every 7 sections (210 μm interval; 8 sections in total; from -12.5 to -11.1 mm from

Bregma, according to the Paxinos and Watson atlas coordinates (Paxinos & Watson, 2023) and

compared total bilateral cell counts of PHOX2B-shRNA rats with non-target control (NT-shRNA)

and naïve rats. Cells that express *Nmb* and *Phox2b* mRNA but do not show co-localization with

PHOX2B protein were considered Nmb^+/PHOX2B$^-$.

The Correct Total Cell Fluorescence (CTCF) signal for *Nmb, Gpr4* and *Task2* mRNA was quantified

as previously described (Cardani et al., 2022; McCloy et al., 2014). A Leica TCS SP5 (B-120G) Laser

Scanning Confocal microscope was used to acquire images of the tissue. Exposure time and

acquisition parameters were set for the naïve group and kept unchanged for the entire dataset

acquisition. For each image, three background areas were used to normalize against

autofluorescence. We used 4 sections/rat to count *Nmb*, *Gpr4* and *Task2* mRNA CTCF in the RTN

area (-12.3 to -11.7 mm from Bregma). For each section two images were acquired with a 20×

objective, so that at least fifty cells per tissue sample were available for analysis. To evaluate

changes in *Nmb* mRNA expression following PHOX2B knockdown at the level of the RTN, we

compared, on the same tissue section, the fluorescence intensity of RTN Nmb^+ cells with the signal

of Nmb^+ cells in the NTS (*Nmb* CTCF ratio RTN/NTS) as the latter would not be affected by the

shRNA infection and knockdown.

To quantify *Gpr4* and *Task2* mRNA expression in the RTN, we first quantified CTCF for either *Gpr4*

or *Task2* mRNA in *Nmb*$^+$ RTN neurons in the 3 experimental groups. We then compared CTCF of

Gpr4 and *Task2* mRNA between *Nmb*$^+$/PHOX2B$^+$ and *Nmb*$^+$/PHOX2B$^-$ RTN neurons in PHOX2B-

shRNA rats to address changes in mRNA expression induced by PHOX2B knockdown.

To assess whether shRNA knockdown affected either the number or the expression of PHOX2B in

C1 neurons, we counted the number of TH$^+$/PHOX2B$^+$ cells along the ventrolateral medulla (-13.2

to -11.5 mm from bregma; 8 sections/rat; 240 μm interval) using an EVOS fluorescent microscope

(ThermoFisher Scientific).

Digital colour photomicrographs were acquired using a Leica TCS SP5 (B-120G) Laser Scanning

Confocal microscope. Image J (version 1.54f; National Institutes of Health, Bethesda, MD, USA)

and Excel (Microsoft Office 365 for Windows) were used for cell counting and the measurements

of fluorescence intensity. Statistical analysis of cell counts, and fluorescent signals was made using

one- way ANOVA (Fig. 2C, 3C, S1B) and repeated-measures ANOVA (Fig 5A, B and 6) with

GraphPad Prism 8 Software (GraphPad Software, Inc., San Diego, CA, USA). Linear regressions to

determine the effect of cell counts (*Nmb*$^+$/PHOX2B$^+$, *Nmb*$^+$/PHOX2B$^-$, total cells) on chemoreflex

magnitude were run using a combined dataset for naive, NT-shRNA and PHOX2B-shRNA rats (SPSS

version 13.0; Fig.5C-D). *p* values of < 0.05 were considered significant and data are reported as

mean ± standard deviation (SD).

 FIGURE LEGENDS

Figure 1. Respiratory data two weeks post viral shRNA injection. **(A)** breathing frequency (f_R,), **(B)**

tidal volume (V_T), **(C)** allometric minute ventilation (V_E ALLO), **(D)** oxygen consumption (VO_2 ALLO),

(E) convective requirement ratio (V_E/VO_2 ALLO), **(F)** hypercapnic ventilatory response (HCVR

absolute change in V_E ALLO vs. corresponding room air) of baseline (pre-surgery, grey filled box),

naïve (black n=8), non-target control shRNA (NT-shRNA, grey n=17) and PHOX2B shRNA (PHOX2B-

shRNA, red n=17) rats 2-weeks post injection during room air, hypercapnia 5% and 7.2% CO_2. f_R is

equally impaired in all experimental group compared to baseline but no treatment effect was

observed (A). V_T is significantly impaired following RTN injection in PHOX2B-shRNA group

compared to naïve animals (p=0.022) and baseline (p< 0.001) at 7.2% CO_2 (B). V_E ALLO is increased

in naïve rats compared to the other treatment groups (p=0.005) in room air (C). Boxplots: median,

$1^{st} - 3^{rd}$ quartiles and $10^{th} - 90^{th}$ percentiles, outliers = dots, "+" indicates arithmetic mean.

Bonferroni post-hoc as indicated. Black*, different from naïve; Grey*, different from NT-shRNA;

Red*, different from PHOX2B sh-RNA.

Figure 2. PHOX2B and *Nmb* expression and total cell count within the RTN area in naïve, NT-

shRNA and PHOX2B-shRNA injected rats two weeks post viral shRNA injection. (A) Schematic

and representative image of a transverse brainstem section at the level of the RTN (-11.5 mm

distance from Bregma) showing the area of investigation containing RTN neurons. **(B)** Expression

of *Phox2b* mRNA (red), PHOX2B protein (white) and *Nmb* mRNA (green) in RTN $Nmb^+/PHOX2B^+$

and $Nmb^+/PHOX2B^-$ neurons in naïve (left), NT-shRNA (middle), and PHOX2B-shRNA (right) rats

(magnified view inserts below). Arrowheads indicate absence of PHOX2B protein. Scale

bar=400μm (top figures), 150μm (inserts below). **(C)** The number of total cells (Nmb^+/PHOX2B$^+$ plus Nmb^+/PHOX2B$^-$) comprising the RTN are reduced in PHOX2B-shRNA rats (n=4) as compared to naïve (n=4) and NT-shRNA (n=4), and in NT-shRNA rats as compared to naïve rats (black[#], One-way ANOVA, p<0.001). The number of Nmb^+/PHOX2B$^+$ cells are reduced in PHOX2B-shRNA rats as compared to naïve and NT-shRNA (Blue*, One-way ANOVA, p<0 0.001). The number of Nmb^+/PHOX2B$^-$ cells was unchanged across groups. **(D,E)** Rostral-caudal distribution of Nmb^+/PHOX2B$^+$ (D) and Nmb^+/PHOX2B$^-$ (E) neurons along the RTN.

Figure 3. PHOX2B and *Nmb* expression and total cell count within the RTN area in naïve, NT-shRNA and PHOX2B-shRNA injected rats 4 weeks post viral shRNA injection. (A) Schematic and representative image of a transverse brainstem section at the level of the RTN (-11.5 mm distance from Bregma) showing the area of investigation containing RTN neurons. **(B)** Expression of *Phox2b* mRNA (red), PHOX2B protein (white) and *Nmb* mRNA (green) in naïve (left), NT-shRNA (middle), and PHOX2B-shRNA (right) rats (magnified view insert). Arrowheads indicate absence of PHOX2B protein. Scale bar = 400μm (top figures), 150μm (inserts below). **(C)** The number of total cells (Nmb^+/PHOX2B$^+$ + Nmb^+/PHOX2B$^-$) comprising the RTN are reduced in PHOX2B-shRNA rats (n=6) as compared to naïve (n=4) (Black[#], One-way ANOVA, p=0.0087) but not to NT-shRNA (n=10). The number of Nmb^+/PHOX2B$^+$ cells are reduced in PHOX2B-shRNA rats as compared to naïve and NT-shRNA (Blue*, one-way ANOVA, p<0.001). The number of Nmb^+/PHOX2B$^-$ cells are increased in PHOX2B-shRNA rats as compared to both naïve and NT-shRNA (Green*, one-way ANOVA p<0.001). **(D, E)** Rostral-caudal distribution of Nmb^+/PHOX2B$^+$ (D) and Nmb^+/PHOX2B$^-$ (E) neurons along the RTN.

Figure 4. Respiratory data following 4 weeks post viral shRNA injection. **(A)** breathing frequency

(f_R,), **(B)** tidal volume (V_T), **(C)** allometric minute ventilation (V_E ALLO), **(D)** convective requirement

ratio (V_E/VO_2 ALLO), **(E)** hypercapnic ventilatory response (HCVR, absolute change in V_E ALLO vs.

corresponding room air), **(F)** HCVR at baseline, week 2 and week 4 post-viral injections in naïve

(black n=8), non-target control shRNA (NT-shRNA, grey n=10) and PHOX2B-shRNA (PHOX2B-

shRNA, red n=6). f_R is equally impaired in all experimental group compared to baseline but no

treatment effect was observed (A). V_T is significantly impaired following RTN injection in PHOX2B-

shRNA group compared to baseline pre-surgery (p<0.001), naïve rats p<0.001), and NT-shRNA rats

(p=0.002) at 7.2% CO_2. (B). V_E ALLO is impaired in PHOX2B-shRNA rats during exposure to

hypercapnia (7.2% CO_2) compared to baseline (p=0.0025), naïve rats (p=0.007), and NT-shRNA rats

(p=0.002). (C). V_E/VO_2 ALLO is reduced in PHOX2B-shRNA animals compared to NT-shRNA rats

both at 5% (p=0.023) and 7.2% (p=0.004.) CO_2 (D). HCVR during 7.2% CO_2 is lower in PHOX2B-

shRNA rats compared to baseline (p=0.007), naïve rats (p=0.001), and NT-shRNA rats (p=0.016) (E).

Boxplots: median, 1^{st} – 3^{rd} quartiles and 10^{th} – 90^{th} percentiles, outliers = dots, "+" indicates

arithmetic mean. Bonferroni post-hoc as indicated. HCVR is significantly impaired only in PHOX2B-

shRNA rats 4-weeks post-surgery (One-way ANOVA p=0.007) (F). Black*, different from naïve;

Grey*, different from NT-shRNA; Red*, different from PHOX2B- shRNA.

Figure 5. PHOX2B knockdown in RTN neurons does not alter TH or *Nmb* expression but impairs

the hypercapnic ventilatory response. (A) No differences are observed in the rostral-caudal

distribution of TH^+/$PHOX2B^+$ catecholaminergic C1 neurons cells caudal to the RTN between naïve

(black, n=4), NT-shRNA (grey, n=10) and PHOX2B-shRNA (red, n=6) rats. **(B)** Quantification of *Nmb*

mRNA fluorescence staining intensity along the rostro-caudal extension of RTN in naïve (black,

n=4), NT-shRNA (grey, n=10) and PHOX2B-shRNA (red, n=6) rat calculated as ratio RTN/NTS shows

no difference between treatment groups. **(D-E)** X-Y plot of HCVR during 7.2% CO_2 exposure

relative to the number of Nmb^+/PHOX2B$^+$ (D, slope is different from "0" at p<0.001; r^2=0.739) and

Nmb^+/PHOX2B$^-$ (E, p<0.001 for difference between slopes; r^2=0.482) in the RTN.

Figure 6. Gpr4 and Task2 mRNA expression within the RTN area in naïve, and PHOX2B-shRNA

injected rats 4 weeks post viral shRNA injection. (A,D) *Nmb* (green), *Gpr4*, *Task2* mRNA (red) and

PHOX2B protein (grey) expression in RTN neurons in naïve (top) and PHOX2B-shRNA (bottom) rats.

Scale bar = 150 µm. Arrowheads indicate colocalization of *Gpr4* (A) and *Task2* (D) with

Nmb^+/PHOX2B$^+$ neurons. Asterisks indicate colocalization of *Gpr4* (A) and *Task2* (D) with

Nmb^+/PHOX2B$^-$ neurons. **(B,E)** Quantification of *Gpr4* and *Task2* mRNA fluorescence staining

intensity along the rostro-caudal extension of RTN in naïve (black), NT-shRNA (grey) and PHOX2B-

shRNA (red) rats. Black*, different from naïve, Grey*, different from NT-shRNA (**C and F**)

Quantification of *Gpr4* and *Task2* mRNA fluorescence staining intensity along the rostro-caudal

extension of RTN in Nmb$^+$/PHOX2B$^+$ (red filled dot) and Nmb$^+$/PHOX2B$^-$ (red empty dot) neurons in

PHOX2B-shRNA rats. Mean corrected total cell fluorescence (CTCF) value ± SEM combined (naïve,

n=4; NT-shRNA, n=10; PHOX2B-shRNA, n=6) (see methods for detail). One-way ANOVA repeated

measures.

Supplementary Fig. 1. (A) Allometric V_E is equally impaired following RTN injection of non-target

control (NT-shRNA n=6) and PHOX2B-shRNA (n=8). Asterisks, difference from baseline. **(B)** The

number of Nmb$^+$/PHOX2B$^+$ cells comprising the RTN are reduced in NT-shRNA (n=5), and shRNA

rats (n=5) as compared to naive controls (n=4; Asterisks, difference from baseline). The number of

Nmb$^+$/PHOX2B$^-$ cells was unchanged in both surgical treatments compared to baseline.

REFERENCES

Abbott, S. B. G., Stornetta, R. L., Coates, M. B., & Guyenet, P. G. (2011). *Phox2b-Expressing Neurons of the Parafacial Region Regulate Breathing Rate, Inspiration, and Expiration in Conscious Rats*. https://doi.org/10.1523/JNEUROSCI.3280-11.2011

Amiel, J., Laudier, B., Attié-Bitach, T., Trang, H., De Pontual, L., Gener, B., Trochet, D., Etchevers, H., Ray, P., Simonneau, M., Vekemans, M., Munnich, A., Gaultier, C., & Lyonnet, S. (2003). Polyalanine expansion and frameshift mutations of the paired-like homeobox gene PHOX2B in congenital central hypoventilation syndrome. *Nature Genetics*, *33*(4), 459–461. https://doi.org/10.1038/ng1130

Bachetti, T., Matera, I., Borghini, S., Duca, M. Di, Ravazzolo, R., & Ceccherini, I. (2005). Distinct pathogenetic mechanisms for PHOX2B associated polyalanine expansions and frameshift mutations in congenital central hypoventilation syndrome. *Human Molecular Genetics*, *14*(13), 1815–1824. https://doi.org/10.1093/hmg/ddi188

Barna, B. F., Takakura, A. C., Mulkey, D. K., & Moreira, T. S. (2016). Purinergic receptor blockade in the retrotrapezoid nucleus attenuates the respiratory chemoreflexes in awake rats. *Acta Physiologica (Oxford, England)*, *217*(1), 80–93. https://doi.org/10.1111/APHA.12637

Basting, T. M., Burke, P. G. R., Kanbar, R., Viar, K. E., Stornetta, D. S., Stornetta, R. L., & Guyenet, P. G. (2015). *Hypoxia Silences Retrotrapezoid Nucleus Respiratory Chemoreceptors via Alkalosis*. https://doi.org/10.1523/JNEUROSCI.2923-14.2015

Berry-Kravis, E. M., Zhou, L., Rand, C. M., & Weese-Mayer, D. E. (2006). Congenital central hypoventilation syndrome PHOX2B mutations and phenotype. *American Journal of Respiratory and Critical Care Medicine*, *174*(10), 1139–1144. https://doi.org/10.1164/rccm.200602-305OC

Biancardi, V., Saini, J., Pageni, A., Prashaad M., H., Funk, G. D., & Pagliardini, S. (2021). Mapping of the excitatory, inhibitory, and modulatory afferent projections to the anatomically defined active expiratory oscillator in adult male rats. *Journal of Comparative Neurology*, *529*(4), 853–884. https://doi.org/10.1002/CNE.24984

Bourdeaut, F., Trochet, D., Janoueix-Lerosey, I., Ribeiro, A., Deville, A., Coz, C., Michiels, J. F., Lyonnet, S., Amiel, J., & Delattre, O. (2005). Germline mutations of the paired-like homeobox 2B (PHOX2B) gene in neuroblastoma. In *Cancer Letters* (Vol. 228, Issues 1–2, pp. 51–58). Elsevier. https://doi.org/10.1016/j.canlet.2005.01.055

Brunet, J. F., & Pattyn, A. (2002). Phox2 genes - From patterning to connectivity. In *Current Opinion in Genetics and Development* (Vol. 12, Issue 4, pp. 435–440). https://doi.org/10.1016/S0959-437X(02)00322-2

Cardani, S., Janes, T. A., Saini, J. K., Di Lascio, S., Benfante, R., Fornasari, D., & Pagliardini, S. (2022). Etonogestrel Administration Reduces the Expression of PHOX2B and Its Target Genes in the Solitary Tract Nucleus. *International Journal of Molecular Sciences*, *23*(9). https://doi.org/10.3390/ijms23094816

Cutsforth-Gregory, J. K., & Benarroch, E. E. (2017). Nucleus of the solitary tract, medullary reflexes, and clinical implications. *Neurology*, *88*(12), 1187–1196. https://doi.org/10.1212/WNL.0000000000003751

Dauger, S. (2003). Phox2b controls the development of peripheral chemoreceptors and afferent visceral pathways. *Development*, *130*(26), 6635–6642. https://doi.org/10.1242/dev.00866

Depocas, F., & Sanford Hart, J. (n.d.). *Use of the Pdzlling Oxygen Analyzer for Memwement of Oxygen conszlm~tion of Ahmds iti Open-Circuit Systems ad in a Short-Lag, Closed-Cirwit Appma tml*.

Di Lascio, S., Bachetti, T., Saba, E., Ceccherini, I., Benfante, R., & Fornasari, D. (2013). Transcriptional dysregulation and impairment of PHOX2B auto-regulatory mechanism induced by polyalanine expansion mutations associated with congenital central hypoventilation syndrome. *Neurobiology of Disease*, *50*(1), 187–200. https://doi.org/10.1016/j.nbd.2012.10.019

Di Lascio, S., Belperio, D., Benfante, R., & Fornasari, D. (2016). Alanine expansions associated with congenital central hypoventilation syndrome impair PHOX2B homeodomain-mediated dimerization and nuclear import. *Journal of Biological Chemistry*, *291*(25), 13375–13393. https://doi.org/10.1074/jbc.M115.679027

Di Lascio, S., Benfante, R., Cardani, S., & Fornasari, D. (2018). Advances in the molecular biology and pathogenesis of congenital central hypoventilation syndrome—implications for new therapeutic targets. *Expert Opinion on Orphan Drugs*, *00*(00), 1–13. https://doi.org/10.1080/21678707.2018.1540978

Di Lascio, S., Benfante, R., Cardani, S., & Fornasari, D. (2021). Research Advances on Therapeutic Approaches to Congenital Central Hypoventilation Syndrome (CCHS). *Frontiers in Neuroscience*, *14*. https://doi.org/10.3389/FNINS.2020.615666

Di Lascio, S., Benfante, R., Zanni, E. Di, Cardani, S., Adamo, A., Fornasari, D., Ceccherini, I., & Bachetti, T. (2018). *Structural and functional differences in PHOX2B frameshift mutations underlie isolated or syndromic congenital central hypoventilation syndrome*. https://doi.org/10.1002/humu.23365

Drorbaugh, J. E., & Fenn, W. O. (1955). A barometric method for measuring ventilation in newborn infants. *Pediatrics*, *16*(1), 81–87. https://doi.org/10.1542/peds.16.1.81

Dubreuil, V., Ramanantsoa, N., Trochet, D., Vaubourg, V., Amiel, J., Gallego, J., Brunet, J.-F., & Goridis, C. (2008). A human mutation in Phox2b causes lack of CO2 chemosensitivity, fatal central apnea, and specific loss of parafacial neurons. *Proceedings of the National Academy of Sciences*, *105*(3), 1067–1072. https://doi.org/10.1073/pnas.0709115105

Dubreuil, V., Thoby-Brisson, M., Rallu, M., Persson, K., Pattyn, A., Birchmeier, C., Brunet, J.-F., Fortin, G., & Goridis, C. (2009). Defective Respiratory Rhythmogenesis and Loss of Central Chemosensitivity in Phox2b Mutants Targeting Retrotrapezoid Nucleus Neurons. *Journal of Neuroscience*, *29*(47), 14836–14846. https://doi.org/10.1523/JNEUROSCI.2623-09.2009

Durand, E., Dauger, S., Pattyn, A., Gaultier, C., Goridis, C., & Gallego, J. (2005). Sleep-disordered breathing in newborn mice heterozygous for the transcription factor phox2b. *American Journal*

of Respiratory and Critical Care Medicine, *172*(2), 238–243.
https://doi.org/10.1164/rccm.200411-1528OC

Fan, Y., Huang, J., Duffourc, M., Kao, R. L., Ordway, G. A., Huang, R., & Zhu, M. Y. (2011).
Transcription factor Phox2 upregulates expression of norepinephrine transporter and
dopamine β-hydroxylase in adult rat brains. *Neuroscience*, *192*, 37–53.
https://doi.org/10.1016/j.neuroscience.2011.07.005

Gestreau, C., Heitzmann, D., Thomas, J., Dubreuil, V., Bandulik, S., Reichold, M., Bendahhou, S.,
Pierson, P., Sterner, C., Peyronnet-Roux, J., Benfriha, C., Tegtmeier, I., Ehnes, H., Georgieff, M.,
Lesage, F., Brunet, J.-F., Goridis, C., Warth, R., & Barhanin, J. (2010). Task2 potassium channels
set central respiratory CO2 and O2 sensitivity. *Proceedings of the National Academy of Sciences*,
107(5), 2325–2330. https://doi.org/10.1073/pnas.0910059107

Goridis, C., Dubreuil, V., Thoby-brisson, M., & Fortin, G. (2010). *Seminars in Cell & Developmental*
Biology Phox2b , congenital central hypoventilation syndrome and the control of respiration. *21*,
814–822. https://doi.org/10.1016/j.semcdb.2010.07.006

Gourine, A. V, Llaudet, E., Dale, N., & Spyer, K. M. (2005). *Behavioral/Systems/Cognitive Release of*
ATP in the Ventral Medulla during Hypoxia in Rats: Role in Hypoxic Ventilatory Response.
https://doi.org/10.1523/JNEUROSCI.3763-04.2005

Guyenet, P. G., Bayliss, D. A., Stornetta, R. L., Ludwig, M. G., Kumar, N. N., Shi, Y., Burke, P. G. R.,
Kanbar, R., Basting, T. M., Holloway, B. B., & Wenker, I. C. (2016). Proton detection and
breathing regulation by the retrotrapezoid nucleus. *Journal of Physiology*, *594*(6), 1529–1551.
https://doi.org/10.1113/JP271480

Guyenet, P. G., Stornetta, R. L., Souza, G. M. P. R., Abbott, S. B. G., Shi, Y., & Bayliss, D. A. (2019).
The Retrotrapezoid Nucleus: Central Chemoreceptor and Regulator of Breathing Automaticity.
Trends in Neurosciences, *42*(11), 807–824. https://doi.org/10.1016/J.TINS.2019.09.002

Harper, R. M., Kumar, R., Macey, P. M., Harper, R. K., & Ogren, J. A. (2015). Impaired neural
structure and function contributing to autonomic symptoms in congenital central
hypoventilation syndrome. *Frontiers in Neuroscience*, *9*(OCT), 415.
https://doi.org/10.3389/FNINS.2015.00415

Janes, T. A., Cardani, S., Saini, J. K., & Pagliardini, S. (2024). *Title: Etonogestrel promotes respiratory*
recovery in an in vivo rat model of central chemoreflex impairment. Running title: Chemoreflex
recovery by etonogesterel.

Kang, B. J., Chang, D. A., Mackay, D. D., West, G. H., Moreira, T. S., Takakura, A. C., Gwilt, J. M.,
Guyenet, P. G., & Stornetta, R. L. (2007). Central nervous system distribution of the
transcription factor Phox2b in the adult rat. *Journal of Comparative Neurology*, *503*(5), 627–
641. https://doi.org/10.1002/cne.21409

Kumar, N. N., Velic, A., Soliz, J., Shi, Y., Li, K., Wang, S., Weaver, J. L., Sen, J., Abbott, S. B. G.,
Lazarenko, R. M., Ludwig, M. G., Perez-Reyes, E., Mohebbi, N., Bettoni, C., Gassmann, M., Suply,
T., Seuwen, K., Guyenet, P. G., Wagner, C. A., & Bayliss, D. A. (2015). Regulation of breathing by
CO2 requires the proton-activated receptor GPR4 in retrotrapezoid nucleus neurons. *Science*,
348(6240), 1255–1260. https://doi.org/10.1126/science.aaa0922

Lighton, J. R. B. (2021). *MEASURING METABOLIC RATES : a manual for scientists.*
https://global.oup.com/academic/product/measuring-metabolic-rates-9780198869320

Madani, A., Pitollat, G., Sizun, E., Cardoit, L., Ringot, M., Bourgeois, T., Ramanantsoa, N., Delclaux,
C., Dauger, S., Thoby-Brisson, M., Gallego, J., & Matrot, B. (2021). Obstructive Apneas in a
Mouse Model of Congenital Central Hypoventilation Syndrome. American journal of respiratory
and critical care medicine, 204(10), 1200–1210. https://doi.org/10.1164/rccm.202104-0887OC.

Marina, N., Abdala, A. P., Trapp, S., Li, A., Nattie, E. E., Hewinson, J., Smith, J. C., Paton, J. F. R., &
Gourine, A. V. (2010). Essential Role of Phox2b-Expressing Ventrolateral Brainstem Neurons in
the Chemosensory Control of Inspiration and Expiration. *Journal of Neuroscience*, *30*(37),
12466–12473. https://doi.org/10.1523/JNEUROSCI.3141-10.2010

Matera, I. (2004). PHOX2B mutations and polyalanine expansions correlate with the severity of the
respiratory phenotype and associated symptoms in both congenital and late onset Central
Hypoventilation syndrome. *Journal of Medical Genetics*, *41*(5), 373–380.
https://doi.org/10.1136/jmg.2003.015412

Mcbride, J. L., Boudreau, R. L., Harper, S. Q., Staber, P. D., Monteys, A. M., Martins, I. S., Gilmore,
B. L., Burstein, H., Peluso, R. W., Polisky, B., Carter, B. J., & Davidson, B. L. (2008). Artificial
miRNAs mitigate shRNA-mediated toxicity in the brain: implications for the therapeutic
development of RNAi. Proceedings of the National Academy of Sciences of the United States of
America, 105(15), 5868–5873. https://doi.org/10.1073/pnas.0801775105

McCloy, R. A., Rogers, S., Caldon, C. E., Lorca, T., Castro, A., & Burgess, A. (2014). Partial inhibition
of Cdk1 in G2 phase overrides the SAC and decouples mitotic events. *Cell Cycle*, *13*(9), 1400–
1412. https://doi.org/10.4161/CC.28401/SUPPL_FILE/KCCY_A_10928401_SM0001.ZIP

Mortola, J. P., & Frappell, P. B. (1998). On the barometric method for measurements of
ventilation, and its use in small animals.

Mortola, J. P., Matsuoka, T., Saiki, C., & Naso, L. (1994). RESPIRATION PHYSIOLOGY Metabolism
and ventilation in hypoxic rats: effect of body mass. *Respiration Physiology*, *97*, 225–234.

Nagashimada, M., Ohta, H., Li, C., Nakao, K., Uesaka, T., Brunet, J. F., Amiel, J., Trochet, D.,
Wakayama, T., & Enomoto, H. (2012). Autonomic neurocristopathy-associated mutations in
PHOX2B dysregulate Sox10 expression. *Journal of Clinical Investigation*, *122*(9), 3145–3158.
https://doi.org/10.1172/JCI63401

Nattie, E. E., & Li, A. (1994). Retrotrapezoid nucleus lesions decrease phrenic activity and CO2
sensitivity in rats. *Respiration Physiology*, *97*(1), 63–77. https://doi.org/10.1016/0034-
5687(94)90012-4

Nattie, E. E., & Li, A. (2002). Substance P-saporin lesion of neurons with NK1 receptors in one
chemoreceptor site in rats decreases ventilation and chemosensitivity. *The Journal of
Physiology*, *544*(2), 603–616. https://doi.org/10.1113/JPHYSIOL.2002.020032

Nobuta, H., Cilio, M. R., Danhaive, O., Tsai, H.-H., Tupal, S., Chang, S. M., Murnen, A., Kreitzer, F.,
Bravo, V., Czeisler, C., Gokozan, H. N., Gygli, P., Bush, S., Weese-Mayer, D. E., Conklin, B., Yee,
S.-P., Huang, E. J., Gray, P. A., Rowitch, D., & Otero, J. J. (2015). Dysregulation of locus coeruleus

development in congenital central hypoventilation syndrome. *Acta Neuropathologica*, *130*(2), 171–183. https://doi.org/10.1007/s00401-015-1441-0

Parodi, S., Di Zanni, E., Di Lascio, S., Bocca, P., Prigione, I., Fornasari, D., Pennuto, M., Bachetti, T., Ceccherini, I., Zanni, E. Di, Lascio, D., Bocca, P., Prigione, I., Fornasari, D., Pennuto, M., & Bachetti, T. (2012). The E3 ubiquitin ligase TRIM11 mediates the degradation of congenital central hypoventilation syndrome-associated polyalanine-expanded PHOX2B. *Journal of Molecular Medicine*, *90*(9), 1025–1035. https://doi.org/10.1007/s00109-012-0868-1

Pattyn, A., Hirsch, M. R., Goridis, C., & Brunet, J. F. (2000). Control of hindbrain motor neuron differentiation by the homeobox gene Phox2b. *Development (Cambridge, England)*, *127*(7), 1349–1358. https://doi.org/10.1242/DEV.127.7.1349

Pattyn, A., Morin, X., & Cremer, H. (1999). *The homeobox gene Phox2b is essential for the development of autonomic neural crest derivatives*. *399*(May).

Pattyn, A., Morin, X., Cremer, H., Goridis, C., & Brunet, J. F. (1997). Expression and interactions of the two closely related homeobox genes Phox2a and Phox2b during neurogenesis. *Development (Cambridge, England)*, *124*(20), 4065–4075. http://www.ncbi.nlm.nih.gov/pubmed/9374403

Paxinos, G., & Watson, C. *The rat brain in stereotaxic coordinates*.

Pirone, L., Caldinelli, L., Di Lascio, S., Di Girolamo, R., Di Gaetano, S., Fornasari, D., Pollegioni, L., Benfante, R., & Pedone, E. (2019). Molecular insights into the role of the polyalanine region in mediating PHOX2B aggregation. *FEBS Journal*, *286*(13), 2505–2521. https://doi.org/10.1111/FEBS.14841

Ramanantsoa, N., Hirsch, M.-R., Thoby-Brisson, M., Dubreuil, V., Bouvier, J., Ruffault, P.-L., Matrot, B., Fortin, G., Brunet, J.-F., Gallego, J., & Goridis, C. (2011). Breathing without CO2 Chemosensitivity in Conditional Phox2b Mutants. *Journal of Neuroscience*, *31*(36), 12880–12888. https://doi.org/10.1523/JNEUROSCI.1721-11.2011

Seifert, E. L., Knowles, J., & Mortola, J. P. (2000). Continuous circadian measurements of ventilation in behaving adult rats. *Respiration Physiology*, *120*, 179–183. www.elsevier.com/locate/resphysiol

Shi, Y., Stornetta, R. L., Stornetta, D. S., Suna Onengut-Gumuscu, X., Farber, E. A., Turner, S. D., Guyenet, P. G., & Bayliss, D. A. (2017). *Neuromedin B Expression Defines the Mouse Retrotrapezoid Nucleus*. https://doi.org/10.1523/JNEUROSCI.2055-17.2017

Shimokaze, T., Sasaki, A., Meguro, T., Hasegawa, H., Hiraku, Y., Yoshikawa, T., Kishikawa, Y., & Hayasaka, K. (2015). Genotype-phenotype relationship in Japanese patients with congenital central hypoventilation syndrome. *Journal of Human Genetics*, *60*(9), 473–477. https://doi.org/10.1038/jhg.2015.65

Souza, G. M. P. R., Kanbar, R., Stornetta, D. S., Abbott, S. B. G., Stornetta, R. L., & Guyenet, P. G. (2018). Breathing regulation and blood gas homeostasis after near complete lesions of the retrotrapezoid nucleus in adult rats. *Journal of Physiology*, *596*(13), 2521–2545. https://doi.org/10.1113/JP275866

Souza, G. M. P. R., Stornetta, D. S., Shi, Y., Lim, E., Berry, F. E., Bayliss, D. A., & Abbott, S. B. G. (2023). Neuromedin B-Expressing Neurons in the Retrotrapezoid Nucleus Regulate Respiratory Homeostasis and Promote Stable Breathing in Adult Mice. *Journal of Neuroscience*, *43*(30), 5501–5520. https://doi.org/10.1523/JNEUROSCI.0386-23.2023

Stanke, M., Junghans, D., Geissen, M., Goridis, C., Ernsberger, U., & Rohrer, H. (1999). The Phox2 homeodomain proteins are sufficient to promote the development of sympathetic neurons. *Development*, *126*(18), 4087–4094. https://doi.org/10.1242/dev.00165

Stornetta, R. L., Moreira, T. S., Takakura, A. C., Kang, B. J., Chang, D. A., West, G. H., Mulkey, D. K., Bayliss, D. A., & Guyenet, P. G. (2006). *Expression of Phox2b by Brainstem Neurons Involved in Chemosensory Integration in the Adult Rat*. *26*(40), 10305–10314. https://doi.org/10.1523/JNEUROSCI.2917-06.2006

Takakura, A. C., Barna, B. F., Cruz, J. C., Colombari, E., & Moreira, T. S. (2014). Phox2b-expressing retrotrapezoid neurons and the integration of central and peripheral chemosensory control of breathing in conscious rats. *Experimental Physiology*, *99*(3). https://doi.org/10.1113/expphysiol.2013.076752

Takakura, A. C., Moreira, T. S., Stornetta, R. L., West, G. H., Gwilt, J. M., & Guyenet, P. G. (2008). Selective lesion of retrotrapezoid Phox2b-expressing neurons raises the apnoeic threshold in rats. *The Journal of Physiology*, *586*(Pt 12), 2975. https://doi.org/10.1113/JPHYSIOL.2008.153163

Tiveron, M. C., Hirsch, M. R., & Brunet, J. F. (1996). The expression pattern of the transcription factor Phox2 delineates synaptic pathways of the autonomic nervous system. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *16*(23), 7649–7660. https://doi.org/10.1523/JNEUROSCI.16-23-07649.1996

Trang, H., Dehan, M., Beaufils, F., Zaccaria, I., Amiel, J., & Gaultier, C. (2005). The French Congenital Central Hypoventilation Syndrome Registry. *Chest*, *127*(1), 72–79. https://doi.org/10.1378/chest.127.1.72

Trochet, D., de Pontual, L., Estêvao, M. H., Mathieu, Y., Munnich, A., Feingold, J., Goridis, C., Lyonnet, S., & Amiel, J. (2008). Homozygous mutation of the PHOX2B gene in congenital central hypoventilation syndrome (Ondine's Curse). *Human Mutation*, *29*(5), 770–770. https://doi.org/10.1002/humu.20727

Trochet, D., Hong, S. J., Lim, J. K., Brunet, J. F., Munnich, A., Kim, K. S., Lyonnet, S., Goridis, C., & Amiel, J. (2005). Molecular consequences of PHOX2B missense, frameshift and alanine expansion mutations leading to autonomic dysfunction. *Human Molecular Genetics*, *14*(23), 3697–3708. https://doi.org/10.1093/hmg/ddi401

Van Gestel, M. A., Van Erp, S., Sanders, L. E., Brans, M. A. D., Luijendijk, M. C. M., Merkestein, M., Pasterkamp, R. J., & Adan, R. A. H. (2014). shRNA-induced saturation of the microRNA pathway in the rat brain. *Gene Therapy*, *21*(2), 205–211. https://doi.org/10.1038/GT.2013.76

Van Limpt, V., Schramm, A., Lakeman, A., Van Sluis, P., Chan, A., Van Noesel, M., Baas, F., Caron, H., Eggert, A., & Versteeg, R. (2004). The Phox2B homeobox gene is mutated in sporadic neuroblastomas. *Oncogene*, *23*(57), 9280–9288. https://doi.org/10.1038/sj.onc.1208157

Weese-Mayer, D. E., Berry-Kravis, E. M., Zhou, L., Maher, B. S., Silvestri, J. M., Curran, M. E., &
Marazita, M. L. (2003). Idiopathic congenital central hypoventilation syndrome: Analysis of
genes pertinent to early autonomic nervous system embryologic development and
identification of mutations in PHOX2b. *American Journal of Medical Genetics*, *123A*(3), 267–
278. https://doi.org/10.1002/ajmg.a.20527

Weese-Mayer, D. E., Rand, C. M., Zhou, A., Carroll, M. S., & Hunt, C. E. (2017). Congenital central
hypoventilation syndrome: A bedside-to-bench success story for advancing early diagnosis and
treatment and improved survival and quality of life. In *Pediatric Research* (Vol. 81, Issues 1–2).
https://doi.org/10.1038/pr.2016.196

Wickström, H. R., Berner, J., Holgert, H., Hökfelt, T., & Lagercrantz, H. (2004). Hypoxic response in
newborn rat is attenuated by neurokinin-1 receptor blockade. *Respiratory Physiology and
Neurobiology*, *140*(1), 19–31. https://doi.org/10.1016/j.resp.2004.01.008

Wu, H., Su, Y., Hung, C., Hsieh, W., & Wu, K. (2009). *Interaction Between PHOX2B and CREBBP
Mediates Synergistic Activation : Mechanistic Implications of PHOX2B Mutants Human
Mutation*. https://doi.org/10.1002/humu.20929

Zoccal, D. B., Furuya, W. I., Bassi, M., A Colombari, D. S., Colombari, E., Hildreth, C., & Putnam, R.
W. (2014). *The nucleus of the solitary tract and the coordination of respiratory and sympathetic
activities*. https://doi.org/10.3389/fphys.2014.00238

A
Retrotrapezoid Nucleus
(RTN)

Bregma -11.5 mm

B
Phox2b mRNA   PHOX2B protein   *Nmb* mRNA
naïve   NT-shRNA   PHOX2B-shRNA

C
Nmb⁺/PHOX2B⁺
Nmb⁺/PHOX2B⁻

D
naïve
NT-shRNA
PHOX2B-shRNA

E
naïve
NT-shRNA
PHOX2B-shRNA