Extreme in Every Way: Exceedingly Low Genetic Diversity in Snow Leopards Due to Persistently Small Population Size

Katherine A. Solari[1&], Simon Morgan[1], Andrey D. Poyarkov[2], Byron Weckworth[3], Gustaf Samelius[4,5], Koustubh Sharma[4,6], Stephane Ostrowski[7], Uma Ramakrishnan[8], Zairbek Kubanychbekov[9], Shannon Kachel[3], Örjan Johansson[4,10], Purevjav Lkhagvajav[11], Heather Hemmingmoore[10], Dmitry Y. Alexandrov[2], Munkhtsog Bayaraa[12], Alexey Grachev[13], Miroslav P. Korablev[2], Jose A. Hernandez-Blanco[2], Bariushaa Munkhtsog[14], Barry Rosenbaum[15], Viatcheslav V. Rozhnov[2], Ali Madad Rajabi[16], Hafizullah Noori[16], Kulbhushansingh Suryawanshi[3,17,18], Ellie E. Armstrong[1,19]*, Dmitri A. Petrov[1,20,21*&]

[1]Department of Biology, Stanford University, Stanford, CA, USA
[2]A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Science, Moscow, Russia
[3]Panthera, New York, NY, USA
[4]Snow Leopard Trust, Seattle, WA, USA
[5]Nordens Ark, Hunnebostrand, Sweden
[6]Snow Leopard Foundation in Kyrgyzstan, Bishkek, Kyrgyz Republic
[7]Wildlife Conservation Society, New York, NY, USA
[8]National Centre for Biological Sciences, TIFR, Bangalore, India
[9]Ilbirs Foundation, Bishkek, Kyrgyzstan
[10]Grimsö Wildlife Research Station, Swedish University of Agricultural Sciences, Riddarhyttan, Sweden
[11]Snow Leopard Conservation Foundation, Ulanbaatar, Mongolia
[12]Wildlife Institute, College of Nature Conservation of the Beijing Forestry University, Beijing, China
[13]Institute of Zoology of Republic of Kazakhstan, Almaty, Kazakhstan
[14]Institute of General and Experimental Biology, Academy of Sciences of Mongolia, Ulaanbaatar, Mongolia
[15]Altai Institute for Research and Conservation, Boulder, CO, USA
[16]Wildlife Conservation Society, Afghanistan Program, Kabul, Afghanistan
[17]Nature Conservation Foundation, Mysore, India
[18]Canadian Institute for Advanced Research, Toronto, Ontario, Canada
[19]Washington State University, Pullman, WA, USA
[20]Chan Zuckerberg BioHub, San Francisco, CA, USA
[21]Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, USA
*Contributed equally
&Corresponding authors emails: ksolari@stanford.edu, dpetrov@stanford.edu

Abstract:

Snow leopards (*Panthera uncia*) serve as an umbrella species whose conservation benefits their high-elevation Asian habitat. Their numbers are believed to be in decline due to numerous Anthropogenic threats; however, their conservation is hindered by numerous knowledge gaps. They are the least studied genetically of all big cat species with more to learn regarding their population structure, historical population size, and current levels of genetic diversity. Here, we use whole-genome sequencing data for 41 snow leopards (37 newly sequenced) to offer new insights into these unresolved questions. Among our samples, we find evidence of a primary genetic divide between the northern and southern part of the range around the Dzungarian Basin, as previously identified, and a secondary divide south of Kyrgyzstan around the Taklamakan Desert. Most noteworthy, we find that snow leopards have the lowest genetic diversity of any big cat species, due to a persistently small population size (relative to other big cat species) throughout their evolutionary history rather than recent inbreeding. Without a large population size or ample standing genetic variation to help buffer them from any forthcoming Anthropogenic challenges, snow leopard persistence may be more tenuous than currently appreciated.

Introduction:

Residing in some of the most extreme and remote areas of the world, snow leopards (*Panthera uncia*) are rarely encountered and are challenging to study, making them one of the most enigmatic of the large charismatic mammals. They are among the largest carnivores in the high-elevation habitat in which they reside and their persistence relies on healthy mountain ungulate populations (Jumabay-Uulu et al. 2014) sometimes supplemented by livestock (Shehzad et al. 2012; Wegge et al. 2012; Aryal et al. 2014; Johansson et al. 2015; Chetri et al. 2017; Suryawanshi et al. 2017; Lu et al. 2021; Thapa et al. 2021). Snow leopard habitat

consists of mountainous areas of Asia, spanning 12 countries (Fig. 1A), habitat that offers

immense ecosystem services–acting as an important source of carbon storage (Mu et al. 2020)

and providing water to almost 2 billion people. Snow leopards serve as an umbrella species

whose conservation benefits this globally crucial Asian mountain ecosystem. In spite of the

global benefits of snow leopard conservation, efforts are often impeded by the many knowledge

gaps regarding this elusive species (McCarthy and Mallon 2016).

Snow leopards were listed as Endangered by the International Union for Conservation of

Nature (IUCN) for 45 years but were downlisted to Vulnerable in 2017 due to not meeting

specific criteria of population size (fewer than 2,500 mature individuals) and percent population

decline (more than 20% over two generations) for Endangered status. This change of status has

been controversial (Aryal 2017; Ale and Mishra 2018) as snow leopard numbers are presumed

to be in decline due to habitat loss, decreased availability of primary prey (high-elevation,

mountain-dwelling ungulates), retaliatory killings for livestock predation (Jackson et al. 2010),

and poaching for their skins (Network 2014). As climate change in high mountain Asia is

occurring at an even more rapid rate than elsewhere in the Northern hemisphere, excluding the

Arctic (Lalande et al. 2021), it is also likely to become an increasing threat to snow leopards (Li

et al. 2016).

Currently, the global snow leopard population size is estimated to be anywhere from

4,700 to 7,500 individuals and little is known about their historical population size and range

(McCarthy et al. 2024) or their current population trends (McCarthy et al. 2017). While many

other big cat species experienced historical declines due to range contractions during the Last

Glacial Maximum (Cooper et al. 2016; Cooper et al. 2021) and are facing a second,

contemporary human-driven decline (Dobrynin et al. 2015; Pečnerová et al. 2021), it remains

unclear if the snow leopard was ever more abundant than is currently estimated. One study

using fecal microsatellite data suggests that snow leopards may have undergone a bottleneck

~8,000 years ago (Janečka et al. 2017). Estimates of genetic diversity and levels of inbreeding

are also limited due to a dearth of genomic data for the species. They are the least studied

genetically of all the big cat species (Weckworth 2021) with whole-genome sequencing (WGS)

data available for only two wild snow leopards (Cho et al. 2013) and two captive individuals

(Armstrong et al. 2022) prior to this study. Genetic diversity assessed using fecal microsatellite

data (Janečka et al. 2017; Aruge et al. 2019; Atzeni et al. 2021; Korablev et al. 2021; Hacker et

al. 2023) as well as genomic data from one of the previously sequenced wild snow leopards

suggests a lack of diversity (Cho et al. 2013), but genomic data from additional samples are

required to determine if this is a characteristic of the species across its range.

Additionally, there is still more to learn about snow leopard population structure and

connectivity across their range (McCarthy et al. 2016). Numerous studies have used

microsatellite markers from fecal samples to assess genetic structure and connectivity at

varying geographic scales. Such studies have found connectivity at the local scale to be location

dependent, with evidence of continuous habitat connectivity across at least 75 km in Pakistan

(Aruge et al. 2019) and >1000 km in Mongolia (Hacker et al. 2023), weak genetic differentiation

among snow leopards samples across 400 km of mountainous terrain in Northern China (Atzeni

et al. 2021), and signs of genetic structure between sampling areas in Russia about 500 km

apart (Korablev et al. 2021). At a larger geographic scale, fecal microsatellite studies (Janečka

et al. 2017; Korablev et al. 2021; Hacker et al. 2023) and other lines of evidence (Riordan et al.

2016; Li et al. 2020) suggest a geographic divide between Mongolia/Russia and the Southern

part of the range due to the Dzungarian Basin and Gobi Desert. However, snow leopards are

known to cross long distances between mountain ranges (McCarthy et al. 2005; Sharma et al.

2014). In addition to this prominent north-south divide, the largest scale fecal microsatellites

study also identified a second divide within the southern group separating the east and west of

the Himalayas-Tibetan Plateau complex (Janečka et al. 2017). Janecka et al. (Janečka et al.

2017) argue that each of these three distinct groups constitute unique subspecies; however, this

subspecies demarcation and the level of connectivity across the landscape remains

controversial (Janečka et al. 2018; Senn et al. 2018). Population structure and connectivity have

never been assessed in snow leopards using whole-genome data.

In addition to the wild snow leopard population, the international community has worked

for decades to establish a sustainable zoo population. As of 2008 there were 445 snow leopards

across 205 institutions globally, not including China, representing the genetic diversity of 56 wild

founders (Blomqvist 2008), most of which came from the wild in the 1960s-1990s, often from

unknown locations. As it is the goal of zoos to maintain a genetically diverse population (Tetzloff

and Schwartz 2024) and to act as reserves of genetic diversity for endangered species, it is

important to know what portion of the global genomic diversity of snow leopards this population

represents.

Here, we generate WGS data for 33 wild snow leopards from multiple locations across

their range in addition to four captive snow leopards. We combine this novel data with existing

data for four individuals to achieve three main objectives: 1) characterize snow leopard

population structure and connectivity to see how estimates from WGS data compare to fecal

microsatellite data; 2) assess the current level of genomic diversity in snow leopards, how this

compares to other big cat species, and how this relates to historical population size and

inbreeding levels; and 3) assess the ancestry of the current zoo snow leopard population.

Among our samples, which do not include most of the central/southern part of the range, we find

evidence of a genetic divide between the north and south, as well as a divide between the

Kyrgyz population and populations farther south. However, low levels of genetic differentiation

between groups suggests recent separation and/or weak levels of connectivity. We also find the

current zoo population to be dominated by Kyrgyzstan-region ancestry. Most notably, we find

snow leopards to be the least genetically diverse contemporary big cat species, owed to a

persistently small population throughout their evolutionary history rather than to recent

inbreeding. We believe these results have significant implications for snow leopard

conservation.

Results:

We generated WGS data for 37 snow leopards–33 wild born snow leopards representing

six countries (Mongolia, Russia, Kyrgyzstan, Tajikistan, Afghanistan, and Pakistan) and four

from the American zoo population. After removing samples based on sequence quality and

breadth we were left with 34 samples from this newly generated sample set with an average

depth of coverage of 7.3X (minimum of 3X and maximum of 16.8X). We also included all

published WGS data for snow leopards in our analysis, which consisted of data for two

additional wild samples, one from Mongolia (Cho et al. 2013) and one from India, and one

captive individual from the American zoo population (Armstrong et al. 2022) with an average

depth of coverage of 23X (minimum of 12X and maximum of 28.9X). Published sequencing data

for an additional captive individual was not included due to sequencing quality issues. After

removing samples based on sequence quality and breadth, we were left with 37 samples–34

from our newly generated data and three previously published samples. This final dataset

consisted of 32 wild born snow leopards representing all seven countries included in the pre-

filtered sample set (Mongolia, Russia, Kyrgyzstan, Tajikistan, Afghanistan, Pakistan, and India)

(Fig. 1B) and five captive samples (Supplementary Table 1).

We mapped these data to the snow leopard reference genome (NCBI accession

PRJNA602938) (Armstrong et al. 2022) and called single nucleotide polymorphisms (SNPs) as

described in the methods, resulting in a final SNP set of 1,591,978. Within this final dataset, we

identified one pair of first-degree relatives and three pairs of second-degree relatives (all within

the wild samples). We removed one representative from each of these four related pairs before

conducting analyses that could be impacted by the presence of related individuals as indicated

in the methods.

Population structure and dispersal barriers:

We assessed population structure among our samples using principal component analysis (PCA). PCA results indicated that the one Indian sample, and to a lesser extent, one of the Tajikistan samples (U13) were genetically distinct from all of the other samples (Fig. 2A, B). In order to more clearly visualize the groupings among the other samples, we also visualized the PCA with the Indian and U13 sample removed. In this PCA, three distinct groups are apparent–Mongolia and Russia; Kyrgyzstan and captive; and Tajikistan, Afghanistan and Pakistan (Fig. 2C).

We also investigated population structure using Admixture (Alexander et al. 2009) to test models with one to ten ancestral populations (K=1-10) over ten independent runs. Due to issues that can arise with having a small sample size for a genetically distinct group (Nazareno et al. 2017), we ran Admixture analyzes both with and without the India sample (Fig. 2D, Supplementary Fig. 1). Indeed, when including the India sample, Admixture had difficulty assigning it at K=2, with five runs grouping it with the southern samples and five runs showing ancestry split between the north and the south (Supplementary Fig. 1A). Once removing the India sample, Mongolia and Russia separated clearly from all other samples with the same ancestry assignments supported in all ten iterations at K=2. At K=3, also with the India sample removed, nine of the ten iterations supported a clear separation of the samples into three groups–Mongolia and Russia; Tajikistan, Afghanistan, and Pakistan; captives and Kyrgyzstan (Fig. 2D, Supplementary Fig. 1B)–recapitulating what was observed in the PCA. Admixture and PCA results were robust to the removal of captive samples from the analyses (Supplementary Fig. 2).

Maximum likelihood phylogeny construction also showed the Indian sample to be genetically distinct (Supplementary Fig. 3). Thus, PCA, Admixture and phylogenetic results all corroborate that the one Indian sample included in this study is genetically unique from all of the other samples.

Among the other samples (excluding India), Admixture and PCA results suggest three

genetically distinct groups. We see a primary genetic divide between the north (Mongolia and

Russia) and the south (all other samples) with a secondary divide within the southern group

between Kyrgyzstan and populations farther south (Afghanistan, Tajikistan, and Pakistan).

These results also show all of our captive samples, whose lineages encompass more than half

of all of the founders of the current captive population (33 of 56 founders are represented,

Supplementary Fig. 4), group most closely with the Kyrgyzstan samples.

Assessment of gene flow:

Based on Admixture and PCA results, we quantified population structure between the

two groups identified in Admixture at K=2 and among the three groups identified at K=3 (Fig.

2D) by assessing shared versus private SNPs, F_{ST}, and rare allele sharing. We excluded

captive samples and the Indian sample from these analyses. We also excluded one sample

from each related pair from F_{ST} and rare allele sharing analyses, but not shared versus private

SNP assessments.

At K=2, each group (North and South, down sampled to n=15) had more shared SNPs

(598,449 SNPs) than private SNPs (379,861 SNPs private to the north and 364,010 SNPs

private to the south) (Supplementary Fig. 5A). At K=3, we compared groups that we will refer to

as "North" (consisting of Russia and Mongolia), "Kyrgyzstan", and "Far South" (consisting of

Tajikistan, Afghanistan and Pakistan). We downsampled the North and Far South groups to

seven individuals such that all three groups had the same sample size. We found that the Far

South and North group had more than twice as many private SNPs (222,569 and 219,175,

respectively) as Kyrgyzstan (111,392) (Supplementary Fig. 5B) and that Kyrgyzstan shared a

similar number of SNPs with both the North and the Far South (394,020 SNPs shared among all

three groups, 65,181 shared only between Kyrgyzstan and North, 64,871 SNPs shared only

between Kyrgyzstan and Far South). It is worth noting that the Kyrgyz group represents the

smallest geographic area and this limitation in sampling could contribute to our observation of

fewer private SNPs in this group.

We calculated Weir and Cockerham's weighted pairwise F_{ST} after removing one

individual for each first and second degree related pair and downsampling groups to equal size.

At K=2, F_{ST} between the north and south was 0.091. At K=3, the pairwise F_{ST} between the North

and Far South was 0.123, between North and Kyrgyzstan was 0.115, and between Kyrgyzstan

and the Far South was 0.093. For context, these values fall well below F_{ST} observed between

different tiger subspecies, which range from 0.164 to 0.318, and are on par with the lower bound

F_{ST} observed between Bengal tiger sub-populations, which range from 0.094 to 0.3 (Armstrong

et al. 2021). Observed pairwise F_{ST} at K=3 was compared to a null distribution representing

panmixia created by randomly shuffling population assignments 10,000 times and recalculating

F_{ST} between populations. This analysis showed observed F_{ST} values to be highly significant

(p<0.0001) (Supplementary Fig. 6).

We identified doubletons (SNPs with an alternate allele count of two) where each

alternate allele occurs in separate individuals. We then assessed doubleton sharing (Anopheles

gambiae 1000 Genomes Consortium et al. 2017) among the three groups–North, Kyrgyzstan,

and Far South–after downsampling each group to six individuals. The amount of doubleton

sharing between populations can offer a sense of population connectivity as diverged

populations have very little rare variant sharing (Gravel et al. 2011). Observed doubleton

sharing was compared to a null distribution representing panmixia in the same way as pairwise

F_{ST} –sample population assignments were randomly shuffled and observed doubleton sharing

between populations was recalculated 10,000 times. Doubleton sharing corroborated the pre-

identified groups with individuals sharing a significantly higher fraction of doubletons (0.40-0.72)

with individuals within the same group (p-value North - 0.0006, Kyrgyzstan - 0.008, Far South -

0.01) (Fig. 2E). However, we also found that each individual shared 11-32% of their doubletons

with each of the other groups, with the Kyrgyzstan population showing the highest fraction of

doubleton sharing with outside groups - even sharing enough with the Far South to not be

significantly different from panmixia (p = 0.2). Both Kyrgyzstan and Far South also share

enough rare alleles with the North to only be marginally significantly different from panmixia

(p=0.04 and p=0.03, respectively). These results support the presence of genetic divergence

among these three groups, but also indicate either recent separation or some low level of gene

flow among the groups.

Figure 1. Snow leopard distribution and sample maps. A) The IUCN snow leopard

distribution (McCarthy et al. 2017) is shown and the largest snow leopard range countries are

labeled. B) Sample locations are indicated with different sized circles indicating the number of

samples from each location. The basemap (ESRI 2013) indicates elevation and landscape

features discussed in the text are labeled. Grey dashed ovals indicate the geographic

distribution of the three subspecies suggested by Janecka et al. 2017– a) *P. u. irbis*, b) *P. u.*

uncia, c) *P. u. uncioides*. In both maps, country boundaries available from the Snow Leopard

Trust (The Snow Leopard Trust) are shown in dark grey. Not all country boundaries are in

agreement, so a dotted line is used for India and a dashed line is used for China to make

overlapping boundaries visible. Maps were created using ArcGIS software by ESRI.

Figure 2. Principal components analysis (PCA), Admixture, and rare allele sharing. A)

PCA of genetic variation of 37 unrelated snow leopards using 1,448,657 SNPs. B) PCA after

removing one outlier sample from India. C) PCA after removing two outlier samples, India and

sample U13 from Tajikistan. PCA axis labels include the percent variation explained by PC1 and

PC2. D) Admixture results for ten independent runs for K=2 and K=3 distinct ancestry groups.

The ancestry assignments shown for K=2 were supported by all ten iterations and the ancestry

assignments shown for K=3 were supported by nine of the ten iterations (the alternate ancestry

assignments supported by one run is shown in Supplementary Fig. 1B). E) Doubleton sharing

between populations compared to null distributions under panmixia. We identified all doubletons

where each allele occurred in a different individual. For each group, identified above each

graph, we then calculated the fraction of doubletons occurring in an individual of that group that

were shared with individuals of each other group. Observed values are shown in color and null

distributions are shown in gray. We made null distributions by randomly shuffling population

assignment among the samples and recalculating doubleton sharing 10,000 times. P-values for

comparisons between observed data and the null distribution using the Wilcoxon Rank Sum

Test are shown. The lower and upper edges of the boxes correspond to the first and third

quartiles and the whiskers extend to the lowest/highest value that is no further than 1.5*IQR

(inter-quartile range) from the box. Points falling further than 1.5*IQR from the box are plotted

individually. In all analyses, we have removed one member of each related pair and in the case

of doubleton sharing, we have downsampled populations to n=6 for each group.

Heterozygosity and historical population size:

We used publicly available data to call SNPs in all big cat species using the Genome

Analysis Toolkit (GATK) (Van der Auwera and O'Connor 2020) and calculated heterozygosity

using VCFtools (Danecek et al. 2011). Here, we defined big cat species as species with an

average adult body weight of 40kg or more, which includes all Panthera species (lion (*P. leo*),

tiger (*P. tigris*), leopard (*P. pardus*), jaguar (*P. onca*), and snow leopard) as well as cheetah

(*Acinonyx jubatus*) and puma (*Puma concolor*) (Johansson et al. 2021). We found snow

leopards to have the lowest heterozygosity of any big cat species, with heterozygosity for every

snow leopard sample included in this study falling lower than that observed in any other big cat

(Fig. 3A). Notably, snow leopard heterozygosity was lower than that of cheetahs, which have

long been considered the archetype of low heterozygosity in big cats (Dobrynin et al. 2015;

Prost et al. 2022). The relative values of observed heterozygosity among all the other species

for which we calculated heterozygosity were consistent with previous work (Paijmans et al.

2021; Pečnerová et al. 2021; Armstrong et al. 2023; Wang et al. 2023).

We used pairwise sequentially Markovian coalescent (PSMC) (Li and Durbin 2011) to

reconstruct historical effective population size using the highest coverage individual from each

genetically distinct group (North-29X, Kyrgyzstan/captive-28X, South-9X, India-12X).

Reconstructions showed consistent results across all populations sampled and across all

bootstrap replicates (Fig. 3B). All reconstructions show a consistently small effective population

size over the last ~300,000 years with snow leopard effective population size never exceeding

14,000 individuals. Reconstructions suggested a snow leopard effective population size of

8,000-6,000 individuals from ~200,000-20,000 years ago and a decrease to an effective

population size of ~2,500-4,000 individuals ~20,000 years ago, coincident with the end of the

Last Glacial Maximum. Effective population size is generally smaller than census size

(Frankham 2007) and PSMC historical effective population size estimates can be impacted by

numerous species-specific parameters (e.g., population structure, inbreeding, mutation rate,

generation time) (Mazet et al. 2016; Mather et al. 2020) which have not been thoroughly

characterized in snow leopards, and thus exact effective population size estimates should be

interpreted cautiously.

Inbreeding and runs of homozygosity (ROH):

Using the same dataset that we used to calculate genome-wide heterozygosity across

big cat species, we also calculated the inbreeding coefficient (F) across big cats. These results

showed the average inbreeding coefficient of snow leopards to be lower than the puma,

cheetah, lion, Asian leopard and numerous tiger subspecies, indicating that the lower genetic

diversity observed in snow leopards is not explained by higher inbreeding (Fig. 4A).

327   We also used the software GARLIC to assess ROH in snow leopards. GARLIC identifies

ROH in three different size classes, each likely the results of different processes– Class A

(short) likely reflects homozygosity in ancient ancestral haplotypes, Class B (intermediate) is

highly correlated with Class A and indicates recent bottlenecks or background levels of

relatedness due to small population size (Pemberton et al. 2012), and Class C (long) reflects

recent inbreeding. GARLIC identified 130bp to be the ideal window size for our data set. Garlic

uses a sliding window to make an initial logarithm of the odds calculation. Class A was defined

as segments under 453,793bp, Class B was defined as segments between 453,793bp -

1,488,700bp, and Class C was defined as segments greater than 1,488,700bp. The proportion

of the genome in each class of ROH for each individual was calculated by dividing the total

sequence length in each size class by the total mappable length of the genome

(1,818,166,894). These proportions were compared to those observed in tigers using GARLIC

output provided in Armstrong et al. (Armstrong et al. 2023).

340   We found the general trends in the proportion of each ROH size class to be different

between snow leopards and tigers, which are known to have undergone recent inbreeding due

to small population sizes (Khan et al. 2021). On average, snow leopards had a similar

proportion of the genome in Class A and B ROH (~10%) and a little less in Class C ROH (~8%)

with some high outliers. Conversely, in numerous tiger subspecies (Bengal, Amur and Generic),

the average proportion of the genome in Class C ROH was much higher than the proportion of

the genome in Class A and B.

347   When comparing ROH trends among snow leopards, we found the proportion of the

genome in Class A ROH to be extremely consistent across samples. The proportion of the

genome in Class B ROH was also quite consistent across groups, with only a few individuals

from the North group showing higher than average values. We observed more variability in the

proportion of the genome in Class C ROH across snow leopard samples, with two outlier

samples, one from the North and one from the Far South, showing high Class C ROH values.

Figure 3. Genome-wide heterozygosity across all big cats and demographic history of

snow leopards. A) Comparison of genome-wide heterozygosity across all big cat species. We

used publicly available data to call SNPs for every big cat species using GATK and calculated

observed heterozygosity using VCFtools. In the case of leopard and tiger, we called SNPs

separately for genetically distinct groups. We calculated snow leopard heterozygosity from

SNPs called using the same in-house pipeline that was used for SNP calling in all the other big

cat species. Only snow leopard samples with a depth of 8X or higher are included (n=15). The

lower and upper edges of the boxes correspond to the first and third quartiles and the whiskers

extend to the lowest/highest value that is no further than 1.5*IQR (inter-quartile range) from the

box. Points falling further than 1.5*IQR from the box are plotted individually. B) Reconstruction

of effective population sizes using PSMC using a mutation rate of 0.7×10^{-8} per site per

generation and a generation time of five years (Cho et al. 2013). Thirty bootstraps are shown for

each sample in thinner, fainter lines of the same color. The average depth of coverage of each

sample used is indicated.

Figure 4. Inbreeding estimates across all big cats and ROH estimates for snow leopards and tigers. A) Comparison of inbreeding coefficient F measured by method of moments across all big cat species. We used publicly available data to call SNPs for every big cat species using GATK and calculated the inbreeding coefficient F using VCFtools. Only snow leopard samples with 8X coverage or greater are included (n=15). B) The proportion of the genome in Class A ROH (small) for snow leopards (colored) and tigers (gray). C) The proportion of the genome in

Class B ROH (intermediate) for snow leopards (colored) and tigers (gray). D) The proportion of

the genome in Class C ROH (long) for snow leopards (colored) and tigers (gray). All ROH

estimates were calculated using GARLIC. We pulled tiger ROH data from Armstrong et al.

(Armstrong et al. 2023). Snow leopard ROH is shown separated by geographic group (North,

Far South, India, Cpative) as well as all samples together (in gold). Tiger ROH is shown

separated by tiger subspecies. All ROH estimates, for snow leopard and tiger, only include

samples with greater than 8X coverage. In all plots, each point represents one individual. The

lower and upper edges of the boxes correspond to the first and third quartiles and the whiskers

extend to the lowest/highest value that is no further than 1.5*IRQ (inter-quartile range) from the

box. Points falling further than 1.5*IRQ from the box are plotted individually.

Discussion:

Population structure and dispersal barriers:

We used WGS from 37 snow leopards to investigate population structure of the species.

Our results corroborate many results of previous studies. Population structure analyses show

that India is genetically distinct from all other samples (Fig. 2A, Supplementary Fig. 3). The

uniqueness of this sample could suggest that this individual is our one representative of the third

phylogenetic group (*P. u. uncioides*, Fig. 1B) suggested by Janečka et al. (Janečka et al. 2017),

but more samples from this area will be necessary to resolve how genetically distinct Indian

snow leopards may be in future analyses.

Among the other samples (excluding India), we identify three genetically distinct groups.

Admixture and PCA results identify the most pronounced divide among our samples to occur

between the northern and southern part of the range around the Dzungarian Basin (Fig. 2B-D),

consistent with previous microsatellite analyses (Janečka et al. 2017; Korablev et al. 2021;

Hacker et al. 2023) and models (Riordan et al. 2016; Li et al. 2020). Admixture and PCA results

also identify a secondary divide occurring south of Kyrgyzstan (Fig. 2C,D), around the

Taklamakan Desert, consistent with previous microsatellite analyses (Korablev et al. 2021).

Both of these genetic divides are also supported by highly significant F_{ST} values between

groups (Supplementary Fig. 6).

These results indicate that the Dzungarian Basin and Taklamakan Desert present

barriers to dispersal for snow leopards. However, while F_{ST} values are significant, they are much

lower than those observed between tiger subspecies. This, in combination with low levels of

differentiation as measured through shared versus private alleles (Supplementary Fig. 5), and

shared rare alleles (Fig. 2E) suggest that these genetic divides may be very recent or there may

be some level of weak connectivity among these groups. This is consistent with previous fecal

microsatellite work which has also found evidence of weak connectivity between Mongolia and

China (Hacker et al. 2023), as well as between Kyrgyzstan and Mongolia/Russia and between

Kyrgyzstan and Tajikistan (Korablev et al. 2021). Additional samples will be necessary to

confidently estimate gene flow in future analyses.

As recommended in Hacker et al. (Hacker et al. 2023), we agree that maintaining this

trace level of connectivity should not be prioritized. Rather, connectivity within genetically

distinct groups should be targeted especially as human development continues and border

fencing, which can potentially act as an absolute barrier to snow leopards, proliferates in this

area (Linnell et al. 2016; Li et al. 2020; Johansson et al. 2024).

Heterozygosity, demographic history and ROH:

Our results show that snow leopards have the lowest genetic diversity of any big cat

species (Fig. 3A), lower than previously appreciated (Kim et al. 2016; Pečnerová et al. 2021).

Demographic history and ROH assessments indicate the exceptionally low heterozygosity in

snow leopards is likely due to a persistently small population size over the last 500,000 years

(Fig. 3B) rather than from recent inbreeding events. With a persistently small population size

and low genetic diversity throughout their evolutionary history, snow leopards serve as yet

another example that high genetic diversity is not a requirement for the long-term persistence of

a species (Xue et al. 2015; Westbury et al. 2018; Westbury et al. 2019; Morin et al. 2021). Our

demographic history assessment (Fig. 3B) did not pick up the more recent population bottleneck

~8,000 years ago suggested by Janecka et al. (Janečka et al. 2017) from microsatellite data

likely because PSMC analyzes loose power at more recent time scales.

Consistent with demographic history assessments, ROH analyses show snow leopards

have a greater proportion of their genome in smaller length ROH (Class A and B) (Fig. 4B,C),

which likely reflects homozygosity in ancient haplotypes and small historic population size

(Pemberton et al. 2012). Conversely, many tiger subspecies, which are known to have high

levels of recent inbreeding due to small isolated populations (Khan et al. 2021), show the

opposite trend, with the highest proportion of their genome in longer ROH (Class C) (Armstrong

et al. 2023) (Fig. 4D). However, we do observe examples of wild snow leopards in the North and

the Far South with large portions of their genome in Class C ROH suggesting that there may be

sporadic inbreeding events occurring in wild populations.

Origin of captive samples:

Studbook-based pedigrees for the five captive individuals sequenced in this study show

that the origin of their wild ancestors is mostly unknown. The few ancestors of known origin are

documented as originating from the USSR, Kyrgyzstan, Kazakhstan, and the western

Himalayas (Supplementary Fig. 4). Admixture and PCA results both show a strong signal of

genetic similarity between the five captive individuals and the current Kyrgyzstan population

(Fig. 2A-D). The current Kyrgyzstan population is likely genetically similar to their close

neighbors in Kazakhstan and northeastern China, for which we have no genomic data. Thus,

these results suggest that the ancestors of the captive individuals largely came from what is

currently Kyrgyzstan, or the surrounding area, and fewer ancestors have come from other

portions of the range.

Conservation implications:

Snow leopards live in arid, cold, low-productivity, high-elevation habitats where few

species can persist – an environment that has evidently only ever been able to support a limited

number of snow leopards (Fig. 3B). As a result, they have likely always had an effective

population size much lower than other big cats (as suggested in Pečnerová et al.(Pečnerová et

al. 2021) and Cho et al. (Cho et al. 2013)), and harbor less genetic diversity than even the

cheetah (Fig. 3A).

Thanks to their extreme environment, snow leopards have not yet been exposed to the

same level of acute Anthropogenic pressures as have big cats living in habitats more easily

accessible to humans. Yet, even having been spared the most intense human impacts, they

already have a small population size and extremely low genomic diversity compared to other big

cats. While these characteristics have not hindered their long-term persistence thus far, this

means that snow leopards do not have the ability to rely on a large population size or standing

genetic variation to help them survive any forthcoming Anthropogenic challenges, as other big

cats have done (Dinerstein et al. 2007; Bauer et al. 2015). Unfortunately, the most intense

Anthropogenic pressures may lie ahead for snow leopards. Anthropogenic climate change

threatens to shrink snow leopard range through habitat change (Li et al. 2016) and intensify

interspecific competition (e.g. (Pal et al. 2022, Lovari et al. 2024)); shifting grazing practices risk

facilitating spillover of novel pathogens from domestic animals into snow leopards and their prey

(Ostrowski and Gilbert 2024); and accelerating mining, energy, and infrastructure development

threaten to fragment and degrade previously remote snow leopard habitats (Heiner et al. 2024;

Zahler and Victurine 2024). Protection of snow leopards and their habitat, to the greatest degree

possible, will be pivotal as this low-density carnivore appears genetically and demographically

ill-equipped to bounce back from Anthropogenic perturbations (Markert et al. 2010). More

broadly, our findings underscore how the IUCN Red List assessment criteria could benefit from

the inclusion of genetic factors (Garner et al. 2020) as our results suggest that snow leopards

may be more imperiled than their current IUCN Red List status would indicate (Ale and Mishra

2018; Suryawanshi et al. 2019).

Methods:

Sample collection and sequencing:

Through an international collaborative effort, we collected a total of 37 snow leopard

blood or tissue samples, composed of 32 wild caught samples from five countries (Afghanistan,

Kyrgyzstan, Tajikistan, Mongolia, and Russia), one currently captive but wild-born individual

from Pakistan (included in the wild group throughout the manuscript), and four captive samples

from mixed/unknown ancestry. Sample details, including DNA extraction and library preparation

methods used, can be found in Supplementary Table 1. We sequenced all samples on an

Illumina sequencing platform using paired end 150bp reads with an aim of sequencing each

sample to ~5-8X coverage.

All samples from captive individuals were collected as part of routine animal care in the

respective AZA zoos and all wild samples were collected as part of ongoing monitoring of snow

leopards by local conservation groups with all necessary collection permits. Samples were

processed in labs in the United States, France, and Russia and were transported to respective

locations with appropriate transport permits.

Additionally, we collected all currently published WGS data for snow leopards, which

included data for two captive individuals (NCBI accessions SRR16227515 (Armstrong et al.

2022) and SRR12437590), one wild individual from Mongolia (NCBI accession SRR836372

(Cho et al. 2013)), and one wild individual from India (NCBI BioProject PRJNA1051290). In

total, we gathered WGS data for 41 snow leopards.

Note that there is WGS data for an additional sample identified as a snow leopard on

Genbank (biosample SAMN17432540, SRA accession SRR13500277, used in the following

publications: (Zhou, Wang, et al. 2021; Zhou, Li, et al. 2021)) that we did not use in this study

because we concluded that this data was from an Asian leopard (*P. pardus*), not a snow leopard

(details of the analyses done to identify this sample as an Asian leopard are provided in the

Supplementary Methods and Supplementary Fig. 7).

We were unable to include any samples from a large portion of the snow leopard range

in the southeast, but are hopeful that data from this area will be available to include in future

analyses.

Mapping and variant calling:

We used the snow leopard reference genome (NCBI accession PRJNA602938

(Armstrong et al. 2022)) and added the lion Y chromosome (from GCA_018350215.1) in order

to have the Y-chromosome represented in this female reference genome. After removing

duplicate reads, reads were mapped and variants called by Gencove using BWA-MEM (Li 2013)

and the GATK (McKenna et al. 2010), respectively. Gencove is a service-providing company

that has developed cloud based pipelines to offer quick and streamlined SNP calling. We

calculated the depth and breadth of coverage for each sample from bam files using SAMtools

(Li et al. 2009) and are provided in Supplementary Table 1.

Variant filtering:

We first filtered variants based on mappability across the genome. We indexed the

reference genome and generated mappability scores using GenMap v1.3.0 (Pockrandt et al.

2019) with flags '-K 30' and '-E 2'. GenMap mappability scores represent the uniqueness of k-

mers (k-mer size given by flag -K) for each position in the genome while allowing for a certain

number of mismatches (given by the flag -E) where a mappability score of one at a position

indicates that the k-mer at that position occurs only once with up to E mismatches. We sorted

the reference genome and mappability bed file using BEDTools (Quinlan and Hall 2010) and

removed SNPs falling in a region of the genome with a mappability score less than one using

BEDTools *intersect* with the flags '-v', '-header', and '-sorted' (leaving 63,199,070 sites). We

then used VCFtools (Danecek et al. 2011) to remove indels using the option '--remove-indels'

(leaving 51,199,725 SNPs). We removed individuals with less than 60% of the genome

represented (samples U02, U12, and U20) using the VCFtools option '--remove-ind'. Most SNPs

were removed at this step because sample U02 had over 1.3 million unique singletons

(Supplementary Table 1) and over 44 million unique doubletons, likely due to some unknown

contamination. We then used VCFtools to remove sites that no longer had any variation using

flag '--non-ref-ac-any 1' (leaving 2,379,069 SNPs), SNPs that did not fall on putative autosomes

(leaving 2,213,174 SNPs), and non-biallelic SNPs using the flags '--min-alleles 2 --max-alleles

2' (leaving 2,146,294 SNPs). We also split scaffold 22 at base pair 67,650,000 into

chromosomes E1 and F1, respectively, to remedy the missassembly described in Armstrong et

al.(Armstrong et al. 2022). Next, with bam files as input, we used GATK v4.1.4.1 to add the

following annotations to our filtered vcf file using *VariantAnnotator*: QD (QualByDepth - variant

confidence normalized by unfiltered depth of variant samples), FS (FisherStrand - strand bias

estimated using Fisher's exact test), MQ (RMSMappingQuality - root mean square of the

mapping quality of reads across all samples), MQRankSum (MappingQualityRankSumTest -

rank sum test for mapping qualities of reference versus alternate reads), and

ReadPosRankSum (ReadPosRankSumTest - rank sum test for relative positioning of reference

versus alternate alleles within reads). After adding these annotations, we used GATK

VariantFiltration to filter SNPs using the following flag: '--filter-expression "QD < 2.0 || FS > 60.0

|| MQ < 40.0 || MQRankSum < -12.5 || ReadPosRankSum < -8.0"'. We then used VCFtools to

remove SNPs that did not pass the GATK *VariantFiltration* step using option '--remove-filtered-

all' (leaving 2,065,453 SNPs). Lastly, we removed SNPs missing data in more than 10% of the

individuals using flag '--max-missing 0.9' (leaving a final SNP set of 1,591,978). We calculated

the number of singletons and private doubletons using the VCFtools flag '--singletons' and

removed one of the captive samples (10x_SDzoo) due to an excess of singletons

(Supplementary Table 1) which we suspected to be due to the 10X library prep method unique

to this sample.

Relatedness assessment:

We estimated relatedness among samples using SNPrelate (Zheng et al. 2012) in R (R

Core Team 2020). We divided the samples into three genetically distinct groups based on

preliminary PCA and Admixture analyses–Mongolia and Russia, Kyrgyzstan and captive, and all

samples South of Kyrgyzstan. We conducted preliminary PCA and Admixture analyses using

the same methods described below but in this case included samples that were later removed

due to relatedness. We ran each group through SNPrelate separately. We used the function

snpgdsBED2GDS to convert bed files to gds files. We used *snpgdsLDpruning* to prune the input

data using the following flags - 'method="corr", slide.max.n=50, ld.threshold=0.3, maf = 0.05,

autosome.only = F'. We then calculated kinship coefficients using the function *snpgdsIBDMLE*

which calculates IBD coefficients for individual pairs using Maximum Likelihood Estimation. A

kinship coefficient of 0.5 indicates a duplicate sample, 0.25 indicates a first-order related pair

(parent-offspring or full sibling), and 0.125 indicates a second-order related pair (half-siblings or

grandparent-grandchild). All sample pairs with non-zero kinship coefficients are listed in

Supplementary Table 2. We identified one sample pair with a kinship coefficient greater than

that expected from first-order relatives–U01 and U09 (kinship coefficient of 0.335), and we

identified three sample pairs with kinship coefficients consistent with second-order related

pairs–SL_KGZ_F1 and SL_KGZ_F4 (kinship coefficient of 0.148), AF_SL_07 and AF_SL_06

(kinship coefficient of 0.135), and U14 and U08 (kinship coefficient of 0.124). We removed the

sample with lower sequencing coverage from each of these pairs (U01, SL_KGZ_F4,

AF_SL_06, and U08) from indicated analyses.

Pedigrees for captive-bred samples:

Using snow leopard studbooks (Blomqvist 2008; Tupa 2011), we compiled information

on all of the ancestors of the five captive-born individuals included in this study. The wild origin

of their ancestors is mostly unknown; however, each sample has a few ancestors for which

there is a wild origin location listed (Supplementary Fig. 4), these origin locations include wild

ancestors listed as coming from the USSR (between 1964-1974), Kazakhstan (in 1972),

Kyrgyzstan (between 1974-1980), Przewalsk (between 1974-1975, which we believe to refer to

Karakol, Kyrgyzstan which was previously named Przewalsk), and Aksai (a contested region

between China and India in the Western Himalayas, in 1979).

Using only information from snow leopard studbooks (Blomqvist 2008; Tupa 2011), we

visualized pedigrees for each individual with the kinship2 package (Sinnwell et al. 2014) in R

(Supplementary Fig. 4).

Principal component analysis:

We conducted principal component analyses (PCAs) on the dataset after filtering for first

and second degree relatives (N=33, 1,448,657 SNPs) using PLINK2 (Purcell et al. 2007). First,

we generated bed files from vcf files using the flags '--make-bed' and '--allow-extra-chr' and

allele frequencies were computed using the flags '--freq' and '--allow-extra-chr'. Then, we ran

the PCA by inputting the resultant bed file with flag '--bfile' and frequency file with flag '--read-

freq' and using the flags '--allow-extra-chr' and '--pca 38'. We visualized PCA output using

ggplot2 in R.

Admixture analysis:

We conducted Admixture analyses on the dataset after filtering for first and second

degree relatives. Since these analyses can be sensitive to having a small sample size for a

genetically distinct group (Nazareno et al. 2017), we ran Admixture analyzes both with and

without the India sample (Puechmaille 2016; Toyama et al. 2020). We used VCFtools to convert

vcf files into PLINK format files. We then used PLINK1.9 to recode alleles numerically using the

flags '--allow-extra-chr --recode12'. We then ran Admixture (Alexander et al. 2009) for K=1-10

ten times using the flag '-s time' so that a different random seed was used for each run based

on the time. We visualized all runs using Clumpak (Kopelman et al. 2015).

We also conducted PCA and Admixture analyses on the dataset after removing captive

samples in order to assess the impact of captive samples on population structure results

(Supplementary Fig. 2) as we know these individuals have at least somewhat mixed ancestry

and have been separated from the wild population for many decades.

Phylogenetic tree construction:

We constructed a phylogenetic tree as an additional way to visualize genetic groupings

among the samples using the same data set used for PCA and Admixture analyses. We

included two tiger samples (SRR13647578 and SRR13500748) to serve as an outgroup. We

first mapped the tiger WGS data to the snow leopard reference genome using BWA-MEM. We

then used the function *vcf2bed* from bedops (Neph et al. 2012) with the flags '-snvs' and '-d' to

convert our filtered snow leopard vcf file to a bed file. We then used BCFtools *mpileup* with the

flags '-A -a AD,DP -R' followed by BCFtools *call* with the flag '-m' to generate a vcf file of SNP

calls for just the snow leopard SNP sites from the tiger data. We then merged the tiger calls to

our snow leopard vcf file using *vcf-merge* from vcftools. We used vcf2phylip (Creators Edgardo

M. Ortiz1 Show affiliations 1. Technical University of Munich) to convert the merged vcf to phylip

format and constructed a maximum likelihood phylogeny using IQtree (Nguyen et al. 2015) with

the flags '-st DNA -m GTR+ASC -T 10 -B 1000'. We visualized the tree using FigTree v1.4.4

(Anon).

Quantifying population structure:

We further characterized population divides identified in Admixture and PCA analyses by

calculating the number of shared versus private SNPs among groups, pairwise F_{ST}, and the rate

of rare variant sharing among groups. We excluded captive samples from these analyses as

well as the sample from India as the PCA indicates that this sample is genetically distinct from

all groups.

We did not filter for relatedness when calculating shared versus private SNPs among the

groups; however, we did downsample as necessary to have an equal number of samples in

each group for all comparisons (n=15 for the K=2 north vs. south comparison and n=7 for the

K=3 North-Kyrgyzstan-FarSouth comparison). We used BCFtools (Li 2011) to create and index

vcf files with only the individuals from each group using the *view* and *index* function,

respectively. We then used BCFtools *isec* to calculate how many SNPs were shared and private

among the groups.

Before calculating F_{ST}, we removed one individual for each first or second degree related

pair. We also downsampled to have an equal number of samples in each group (n=13 for the

K=2 north vs. south comparison and n=6 for the K=3 North-Kyrgyzstan-FarSouth comparison).

We then used vcftools to calculate the Weir and Cockerham weighted F_{ST} estimate (Weir and

Cockerham 1984) between each group using the flag '--weir-fst-pop'. In order to assess the

significance of F_{ST} estimates at K=3, we conducted 10,000 permutations by shuffling the

population assignments and recalculating F_{ST} to generate a null distribution.

To assess doubleton sharing among groups, we excluded one individual for each related

pair and downsampled all groups so that each had an equal number of individuals (N=6). We

then used VCFtools to retain only SNPs with a minor allele count of two (doubletons) using the

flags '--mac 2 --max-mac 2'. We removed any SNPs where the doubletons were present in the

same individual and identified allele sharing among samples using the flag '--genome full' in

PLINK. We created a null distribution under panmixia by calculating doubleton sharing between

populations after randomly shuffling population assignments among the samples 10,000 times.

We compared fractions of observed doubleton sharing to the null distribution using the Wilcoxon

Rank Sum Test in R.

PSMC:

We used PSMC (Li and Durbin 2011) to estimate snow leopard effective population size

back in time using the highest coverage sample for each population cluster. We generated

diploid consensus sequences for each individual as recommended by PSMC. Starting with bam

files filtered to include only putative autosomes, we used SAMtools *mpileup* and BCFtools *call* to

generate a vcf. We then used *vcfutils.pl vcf2fq* to generate diploid fastq files using the '-D' and '-

d' flags to reflect the average depth of coverage of each sample as recommended in PSMC

documentation. We then ran PSMC with the default settings. We also performed 30 rounds of

bootstrapping using random sampling with replacement for each sample. For plotting, we used

a mutation rate of 0.7×10^{-8} per site per generation as suggested for snow leopards by Cho et al.

(Cho et al. 2013) and a generation time of five years.

Heterozygosity in other big cats:

We calculated heterozygosity in all big cat species using publicly available data in order

to see how snow leopard heterozygosity levels compare to other big cat species. We included

all other species in the genus *Panthera* (leopard, lion, tiger, jaguar) as well as cheetah and

puma. Accession numbers of publicly available WGS data and reference genomes used in this

analysis are listed in Supplementary Table 3 and 4, respectively. We mapped all fastq data to

the corresponding reference genome using BWA-MEM (Li 2013). We sorted and indexed

resulting bam files using SAMtools (Li et al. 2009). We added read groups and marked

duplicates using picard (Anon). We calculated depth of coverage and breadth of coverage for

each sample using SAMtools. We then used GATK (Van der Auwera and O'Connor 2020)

HaplotypeCaller to generate a gvcf file for each sample and GATK *CombineGVCFs* to combine

the gvcf files for every species (or subgroup in the case of tigers and leopards). Lastly, we used

GATK *GenotypeGVCFs* to create a final vcf file for every group.

In order to limit the need for excessive computing power, we downsampled some

samples with over 30X coverage using SAMtools *view* as indicated in Supplementary Table 3.

The leopard samples required a larger amount of computing power than the other species to

call haplotypes, so we split these samples into one bam file per contig using BAMtools *split* and

ran each contig through GATK *HaplotypeCaller* and *CombineGVCFs* in parallel. We also found

it necessary to split the jaguar and snow leopard data into intervals to combine gvcf files using

GATK *GenomicsDBImport* followed by *CombineGVCFs*. In both cases, we concatenated

resultant vcf files, one per contig or interval, using BCFtools *concat*.

We filtered each reference genome for mappability using genmapv1.3.0 to first index

and then calculate mappability using flags '-K 30' and '-E 2'. We then filtered each vcf to only

include nucleotides with a mappability score of one. To do this, we sorted both vcf files and

mappability bed files using BEDTools and retained only SNPs falling in regions with a

mappability score of one in each vcf using BEDTools *intersect*. We then filtered each vcf using

GATK *VariantFiltration* with the following flag: '--filter-expression "QD < 2.0 || FS > 60.0 || MQ <

40.0 || MQRankSum < -12.5 || ReadPosRankSum < -8.0"'. Then, we filtered vcf files to remove

indels and non-bialellic SNPs using VCFtools flags '--remove-indels --min-alleles 2 --max-alleles

2'.

We calculated the mappable length of each genome using mappability bed files to

calculate the number of nucleotides with a mappability score of one. Using the filtered vcf for

each species or group, we calculated observed homozygosity for each sample using VCFtools

(Danecek et al. 2011) with the flag '--het'. In R, we calculated the number of heterozygous sites

by subtracting the number of observed homozygous sites column ((O)HOM) from the total

number of sites column (NSITES). We then calculated heterozygosity by dividing the number of

heterozygous sites by the length of the genome consisting of putative autosomes with a

mappability score of one. We used ggplot2 in R to create box plots of heterozygosity results.

Heterozygosity in snow leopards:

Although SNPs had already been called by Gencove (as described above), we re-called

SNPs from the snow leopard dataset using the same in-house pipeline used to call SNPs in all

of the other big cat species to ensure comparability of heterozygosity calculations among

species. We calculated observed heterozygosity in snow leopards using both SNP datasets

(Gencove's and ours) in the same way described above. We calculated heterozygosity for all

individuals, regardless of relatedness (N=37); however, because heterozygous SNPs are most

accurately called with higher coverage data (Maruki and Lynch 2017), we only used samples

with 8X coverage or more.

We compared snow leopard heterozygosity calculated from SNPs called using our in-

house pipeline to heterozygosity calculated from SNPs called by Gencove using Pearson

correlation coefficient calculated using the ggpubr (Kassambara 2020) package in R

(Supplementary Fig. 8). We found heterozygosity calculated from the two different SNP calling

pipelines to be extremely correlated (R = 0.97, p=5.1E-9); however, all heterozygosity estimates

calculated from our in-house SNP calls were slightly higher (average of ~0.000056) than that

estimated from Gencove SNP calls.

Runs of homozygosity:

We converted vcf files to transposed PLINK files using plink version 1.90b5.3 using the

flags '--allow-extra-chr –const-fid 0 --recode transpose'. We then inferred ROH in each

individual using the software GARLIC (Szpiech et al. 2017) with an error rate of 0.001 and the

centromere location of each chromosome set to 0, 0; since centromere location is unknown. In

addition, we used '--auto-win-size, --auto-overlap-frac, –winsize 100' to allow GARLIC to

determine the ideal window size to use. We excluded runs of homozygosity that were less than

50,000bp.

ROH were automatically divided into different size classes by GARLIC - A (short), B

(intermediate), and C (long). We calculated the proportion of the genome in each size class by

dividing the total sequence length in each size class by the total sequence length of putative

autosomes with a mappability score greater than one. The proportion of the genome in each

size class was visualized in box plots using ggplot2 in R.

We pulled ROH values for tigers, calculated using GARLIC, from Armstrong et al.

(Armstrong et al. 2023) and calculated the proportion of the genome in each size class of ROH

for each individual in the same way as described above, but using the total mappable length

across all autosomes in the tiger reference genome.

As with heterozygosity assessments in snow leopards, in order to limit any biases

caused by allelic drop out in lower coverage samples, we only used samples with 8X coverage

or higher in ROH assessments in both snow leopards and tigers.

Inbreeding across big cats:

We calculated the inbreeding coefficient F in all big cat species using the same SNP

datasets used to calculate heterozygosity. As with heterozygosity assessments, we only

calculated inbreeding for snow leopards samples with 8X coverage or higher. We calculated the

inbreeding coefficient F using the method of moments ((observed homozygous count - expected

count)/(total observations - expected count)) using vcftools with the flag '-het'.

Samples included in each analysis:

763   We provide a detailed list of which samples were included in each analysis in

764 Supplementary Table 5.

765 **Data Availability:**

766 Data associated with this study has been deposited into bioproject PRJNA1048427 and will be

767 released upon publication.

769 **Code Availability:**

770 The code used for analyses in this project is available on the project's github:

771 https://github.com/ksolari/SL_WGS

773 **Acknowledgements:**

774 This work was supported by funding from the Snow Leopard Trust. We would like to thank those

775 who helped us in acquiring the captive samples used in this study, including Karen Ingerman

776 and Susie Bartlett from the Bronx Zoo. We would like to thank Haiqing Xu for his help in

777 conducting simulations to produce the null distribution for the shared doubleton analysis.

779 **References:**

780 Ale SB, Mishra C. 2018. The snow leopard's questionable comeback. *Science* 359:1110.

781 Alexander DH, Novembre J, Lange K. 2009. Fast model-based estimation of ancestry in
782   unrelated individuals. *Genome Res.* 19:1655–1664.

783 Anon. FigTree. Available from: http://tree.bio.ed.ac.uk/software/figtree/

784 Anon. Picard. Available from: http://broadinstitute.github.io/picard/

785 Anopheles gambiae 1000 Genomes Consortium, Data analysis group, Partner working group,
786   Sample collections—Angola:, Burkina Faso:, Cameroon:, Gabon:, Guinea:, Guinea-Bissau:,
787   Kenya:, et al. 2017. Genetic diversity of the African malaria vector Anopheles gambiae.
788   *Nature* 552:96–100.

789 Armstrong EE, Campana MG, Solari KA, Morgan SR, Ryder OA, Naude VN, Samelius G,
790   Sharma K, Hadly EA, Petrov DA. 2022. Genome report: chromosome-level draft

assemblies of the snow leopard, African leopard, and tiger (Panthera uncia, Panthera
pardus pardus, and Panthera tigris). *G3* [Internet] 12. Available from:
http://dx.doi.org/10.1093/g3journal/jkac277

Armstrong EE, Khan A, Taylor RW, Gouy A, Greenbaum G, Thiéry A, Kang JT, Redondo SA,
Prost S, Barsh G, et al. 2021. Recent Evolutionary History of Tigers Highlights Contrasting
Roles of Genetic Drift and Selection. *Mol. Biol. Evol.* 38:2366–2379.

Armstrong EE, Mooney JA, Solari KA, Kim BY, Barsh GS, Grant V, Greenbaum G, Kaelin CB,
Panchenko K, Pickrell JK, et al. 2023. Unraveling the Genomic Diversity and Evolutionary
History of Captive Tigers in the United States. *bioRxiv* [Internet]:2023.06.19.545608.
Available from: https://www.biorxiv.org/content/10.1101/2023.06.19.545608v1.abstract

Aruge S, Batool H, Khan FM, Fakhar-i-Abbas, Janjua S. 2019. A pilot study—genetic diversity
and population structure of snow leopards of Gilgit-Baltistan, Pakistan, using molecular
techniques. *PeerJ* 7:e7672.

Aryal A. 2017. Poaching: Is snow leopard tally underestimated? *Nature* 550:457.

Aryal A, Brunton D, Ji W, Karmacharya D, McCarthy T, Bencini R, Raubenheimer D. 2014.
Multipronged strategy including genetic analysis for assessing conservation options for the
snow leopard in the central Himalaya. *J. Mammal.* 95:871–881.

Atzeni L, Cushman SA, Wang J, Riordan P, Shi K, Bauman D. 2021. Evidence of spatial genetic
structure in a snow leopard population from Gansu, China. *Heredity* 127:522–534.

Bauer H, Chapron G, Nowell K, Henschel P, Funston P, Hunter LTB, Macdonald DW, Packer C.
2015. Lion (Panthera leo) populations are declining rapidly across Africa, except in
intensively managed areas. *Proc. Natl. Acad. Sci. U. S. A.* 112:14894–14899.

Blomqvist L. 2008. International Pedigree Book for Snow leopards, Uncia uncia Volume 9.

Chetri M, Odden M, Wegge P. 2017. Snow Leopard and Himalayan Wolf: Food Habits and Prey
Selection in the Central Himalayas, Nepal. *PLoS One* 12:e0170549.

Cho YS, Hu L, Hou H, Lee H, Xu J, Kwon S, Oh S, Kim H-M, Jho S, Kim S, et al. 2013. The
tiger genome and comparative analysis with lion and snow leopard genomes. *Nat.*
Commun. 4:2433.

Cooper DM, Dugmore AJ, Gittings BM, Scharf AK, Wilting A, Kitchener AC. 2016. Predicted
Pleistocene–Holocene range shifts of the tiger (Panthera tigris). *Divers. Distrib.* 22:1199–
1211.

Cooper DM, Dugmore AJ, Kitchener AC, Metzger MJ, Trabucco A. 2021. A kingdom in decline:
Holocene range contraction of the lion () modelled with global environmental stratification.
PeerJ 9:e10504.

Creators Edgardo M. Ortiz1 Show affiliations 1. Technical University of Munich. vcf2phylip v2.0:
convert a VCF matrix into several matrix formats for phylogenetic analysis. Available from:
https://zenodo.org/record/2540861

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics*

27:2156–2158.

Dinerstein E, Loucks C, Wikramanayake E, Ginsberg J, Sanderson E, Seidensticker J, Forrest
J, Bryja G, Heydlauff A, Klenzendorf S, et al. 2007. The Fate of Wild Tigers. *Bioscience*
57:508–514.

Dobrynin P, Liu S, Tamazian G, Xiong Z, Yurchenko AA, Krasheninnikova K, Kliver S, Schmidt-
Küntzel A, Koepfli K-P, Johnson W, et al. 2015. Genomic legacy of the African cheetah,
Acinonyx jubatus. *Genome Biol.* 16:277.

ESRI. 2013. Terrain: Elevation Tinted Hillshade. Available from:
https://elevation.arcgis.com/arcgis/rest/services/WorldElevation/Terrain/ImageServer

Frankham R. 2007. Effective population size/adult population size ratios in wildlife: a review.
Genet. Res. 89:491–503.

Garner BA, Hoban S, Luikart G. 2020. IUCN Red List and the value of integrating genetics.
Conserv. Genet. 21:795–801.

Gravel S, Henn BM, Gutenkunst RN, Indap AR, Marth GT, Clark AG, Yu F, Gibbs RA, 1000
Genomes Project, Bustamante CD. 2011. Demographic history and rare allele sharing
among human populations. *Proc. Natl. Acad. Sci. U. S. A.* 108:11983–11988.

Hacker C, Atzeni L, Munkhtsog B, Munkhtsog B, Galsandorj N, Zhang Y, Liu Y, Buyanaa C,
Bayandonoi G, Ochirjav M, et al. 2023. Genetic diversity and spatial structures of snow
leopards (Panthera uncia) reveal proxies of connectivity across Mongolia and northwestern
China. *Landsc. Ecol.* 38:1013–1031.

Heiner M, Oakleaf J, Davaa G, Kiesecker J. 2024. Chapter 10 - Emerging threats to snow
leopards from energy and mineral development. In: Mallon D, McCarthy T, editors. Snow
Leopards (Second Edition). Academic Press. p. 113–121.

Jackson RM, Mishra C, McCarthy TM, Ale SB. 2010. Snow leopards: conflict and conservation.
The biology and conservation of wild felids 2:417–430.

Janečka JE, Janecka MJ, Helgen KM, Murphy WJ. 2018. The validity of three snow leopard
subspecies: response to Senn et al. *Heredity* 120:586–590.

Janečka JE, Zhang Y, Li D, Munkhtsog B, Bayaraa M, Galsandorj N, Wangchuk TR,
Karmacharya D, Li J, Lu Z, et al. 2017. Range-Wide Snow Leopard Phylogeography
Supports Three Subspecies. *J. Hered.* 108:597–607.

Johansson Ö, Alexander JS, Lkhagvajav P, Mishra C, Samelius G. 2024. Natal dispersal and
exploratory forays through atypical habitat in the mountain-bound snow leopard. *Ecology*
105:e4264.

Johansson Ö, Ausilio G, Low M, Lkhagvajav P, Weckworth B, Sharma K. 2021. The timing of
breeding and independence for snow leopard females and their cubs. *Mamm. Biol.*
101:173–180.

Johansson Ö, McCarthy T, Samelius G, Andrén H, Tumursukh L, Mishra C. 2015. Snow leopard
predation in a livestock dominated landscape in Mongolia. *Biol. Conserv.* 184:251–258.

Jumabay-Uulu K, Wegge P, Mishra C, Sharma K. 2014. Large carnivores and low diversity of
optimal prey: a comparison of the diets of snow leopards Panthera uncia and wolves Canis
lupus in Sarychat-Ertash Reserve in Kyrgyzstan. *Oryx* 48:529–535.

Kassambara A. 2020. ggpubr:"ggplot2" based publication ready plots.

Khan A, Patel K, Shukla H, Viswanathan A, van der Valk T, Borthakur U, Nigam P, Zachariah A,
Jhala YV, Kardos M, et al. 2021. Genomic evidence for inbreeding depression and purging
of deleterious genetic variation in Indian tigers. *Proc. Natl. Acad. Sci. U. S. A.* [Internet] 118.
Available from: http://dx.doi.org/10.1073/pnas.2023018118

Kim S, Cho YS, Kim H-M, Chung O, Kim H, Jho S, Seomun H, Kim J, Bang WY, Kim C, et al.
2016. Comparison of carnivore, omnivore, and herbivore mammalian genomes with a new
leopard assembly. *Genome Biol.* 17:211.

Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I. 2015. Clumpak: a program
for identifying clustering modes and packaging population structure inferences across K.
Mol. Ecol. Resour. 15:1179–1191.

Korablev MP, Poyarkov AD, Karnaukhov AS, Zvychaynaya EY, Kuksin AN, Malykh SV, Istomov
SV, Spitsyn SV, Aleksandrov DY, Hernandez-Blanco JA, et al. 2021. Large-scale and fine-
grain population structure and genetic diversity of snow leopards (Panthera uncia Schreber,
1776) from the northern and western parts of the range with an emphasis on the Russian
population. *Conserv. Genet.* 22:397–410.

Lalande M, Ménégoz M, Krinner G, Naegeli K, Wunderle S. 2021. Climate change in the High
Mountain Asia in CMIP6. *Earth Syst. Dyn.* 12:1061–1098.

Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. *Bioinformatics* 27:2987–
2993.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv [q-bio.GN] [Internet]. Available from: http://arxiv.org/abs/1303.3997

Li H, Durbin R. 2011. Inference of human population history from individual whole-genome
sequences. *Nature* 475:493–496.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map
format and SAMtools. *Bioinformatics* 25:2078–2079.

Li J, McCarthy TM, Wang H, Weckworth BV, Schaller GB, Mishra C, Lu Z, Beissinger SR. 2016.
Climate refugia of snow leopards in High Asia. *Biol. Conserv.* 203:188–196.

Li J, Weckworth BV, McCarthy TM, Liang X, Liu Y, Xing R, Li D, Zhang Y, Xue Y, Jackson R, et
al. 2020. Defining priorities for global snow leopard conservation landscapes. *Biol. Conserv.*
241:108387.

Linnell JDC, Trouwborst A, Boitani L, Kaczensky P, Huber D, Reljic S, Kusak J, Majic A,
Skrbinsek T, Potocnik H, et al. 2016. Border Security Fencing and Wildlife: The End of the
Transboundary Paradigm in Eurasia? *PLoS Biol.* 14:e1002483.

Lovari S, Kachel S, Xueyang L, Ferretti F. 2024. Chapter 13 - Snow leopard, common leopard,
and wolf: Are they good neighbors? In: Mallon D, McCarthy T, editors. Snow Leopards
(Second Edition). Academic Press. p. 137–147.

Lu Q, Xiao L, Cheng C, Lu Z, Zhao J, Yao M. 2021. Snow Leopard Dietary Preferences and
Livestock Predation Revealed by Fecal DNA Metabarcoding: No Evidence for Apparent
Competition Between Wild and Domestic Prey. *Frontiers in Ecology and Evolution* [Internet]
9. Available from: https://www.frontiersin.org/articles/10.3389/fevo.2021.783546

Markert JA, Champlin DM, Gutjahr-Gobell R, Grear JS, Kuhn A, McGreevy TJ Jr, Roth A,
Bagley MJ, Nacci DE. 2010. Population genetic diversity and fitness in multiple
environments. *BMC Evol. Biol.* 10:205.

Maruki T, Lynch M. 2017. Genotype Calling from Population-Genomic Sequencing Data. *G3*
7:1393–1404.

Mather N, Traves SM, Ho SYW. 2020. A practical introduction to sequentially Markovian
coalescent methods for estimating demographic history from genomic data. *Ecol. Evol.*
10:579–589.

Mazet O, Rodríguez W, Grusea S, Boitard S, Chikhi L. 2016. On the importance of being
structured: instantaneous coalescence rates and human evolution--lessons for ancestral
population size inference? *Heredity* 116:362–371.

McCarthy T, Mallon D eds. 2016. Snow Leopards: Biodiversity of the World: Conservation from
Genes to Landscapes. Academic Press

McCarthy T, Mallon D, Jackson R, Zahler P, McCarthy K. 2017. Panthera uncia. *The IUCN Red*
List of Threatened Species [Internet] e.T22732A50664030. . Available from:
https://dx.doi.org/10.2305/IUCN.UK.2017-2.RLTS.T22732A50664030.en

McCarthy T, Mallon D, Sanderson EW, Zahler P, Fisher K. 2016. What is a Snow Leopard?
Biogeography and Status Overview. In: McCarthy T, Mallon D, editors. Snow Leopards.
San Diego, CA: Elsevier. p. 23–42.

McCarthy T, Mallon D, Zahler P. 2024. Chapter 3 - What is a snow leopard? Biogeography and
status overview. In: Mallon D, McCarthy T, editors. Snow Leopards (Second Edition).
Academic Press. p. 31–41.

McCarthy TM, Fuller TK, Munkhtsog B. 2005. Movements and activities of snow leopards in
Southwestern Mongolia. *Biol. Conserv.* 124:527–537.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler
D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a MapReduce framework
for analyzing next-generation DNA sequencing data. *Genome Res.* 20:1297–1303.

Morin PA, Archer FI, Avila CD, Balacco JR, Bukhman YV, Chow W, Fedrigo O, Formenti G,
Fronczek JA, Fungtammasan A, et al. 2021. Reference genome and demographic history
of the most endangered marine mammal, the vaquita. *Mol. Ecol. Resour.* 21:1008–1020.

Mu C, Abbott BW, Norris AJ, Mu M, Fan C, Chen X, Jia L, Yang R, Zhang T, Wang K, et al.
2020. The status and stability of permafrost carbon on the Tibetan Plateau. *Earth-Sci. Rev.*

211:103433.

Nazareno AG, Bemmels JB, Dick CW, Lohmann LG. 2017. Minimum sample sizes for
population genomics: an empirical study from an Amazonian plant species. *Mol. Ecol.*
Resour. 17:1136–1147.

Neph S, Kuehn MS, Reynolds AP, Haugen E, Thurman RE, Johnson AK, Rynes E, Maurano
MT, Vierstra J, Thomas S, et al. 2012. BEDOPS: high-performance genomic feature
operations. *Bioinformatics* 28:1919–1920.

Network SL. 2014. Snow leopard survival strategy. *Seattle, Washington, USA*:1–145.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.*
32:268–274.

Ostrowski S, Gilbert M. 2024. Chapter 9 - Diseases of wild snow leopards and their wild
ungulate prey. In: Mallon D, McCarthy T, editors. Snow Leopards (Second Edition).
Academic Press. p. 95–111.

Paijmans JLA, Barlow A, Becker MS, Cahill JA, Fickel J, Förster DWG, Gries K, Hartmann S,
Havmøller RW, Henneberger K, et al. 2021. African and Asian leopards are highly
differentiated at the genomic level. *Curr. Biol.* 31:1872–1882.e5.

Pal R, Panwar A, Goyal SP, Sathyakumar S. 2022. Changes in ecological conditions may
influence intraguild competition: inferring interaction patterns of snow leopard with co-
predators. *PeerJ* 10:e14277.

Pečnerová P, Garcia-Erill G, Liu X, Nursyifa C, Waples RK, Santander CG, Quinn L, Frandsen
P, Meisner J, Stæger FF, et al. 2021. High genetic diversity and low differentiation reflect
the ecological versatility of the African leopard. *Curr. Biol.* 31:1862–1871.e5.

Pemberton TJ, Absher D, Feldman MW, Myers RM, Rosenberg NA, Li JZ. 2012. Genomic
patterns of homozygosity in worldwide human populations. *Am. J. Hum. Genet.* 91:275–
292.

Pockrandt C, Alzamel M, Iliopoulos CS, Reinert K. 2019. GenMap: Fast and Exact Computation
of Genome Mappability. *Cold Spring Harbor Laboratory* [Internet]:611160. Available from:
https://www.biorxiv.org/content/10.1101/611160v1.abstract

Prost S, Machado AP, Zumbroich J, Preier L, Mahtani-Williams S, Meissner R, Guschanski K,
Brealey JC, Fernandes CR, Vercammen P, et al. 2022. Genomic analyses show extremely
perilous conservation status of African and Asiatic cheetahs (Acinonyx jubatus). *Mol. Ecol.*
31:4208–4223.

Puechmaille SJ. 2016. The program structure does not reliably recover the correct population
structure when sampling is uneven: subsampling and new estimators alleviate the problem.
Mol. Ecol. Resour. 16:608–627.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de
Bakker PIW, Daly MJ, et al. 2007. PLINK: a tool set for whole-genome association and
population-based linkage analyses. *Am. J. Hum. Genet.* 81:559–575.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* 26:841–842.

R Core Team. 2020. R: A language and environment for statistical computing. Available from:
https://www.R-project.org/

Riordan P, Cushman SA, Mallon D, Shi K, Hughes J. 2016. Predicting global population
connectivity and targeting conservation action for snow leopard across its range.
Ecography 39:419–426.

Senn H, Murray-Dickson G, Kitchener AC, Riordan P, Mallon D. 2018. Response to Janečka et
al. 2017. *Heredity* 120:581–585.

Sharma K, Bayrakcismith R, Tumursukh L, Johansson O, Sevger P, McCarthy T, Mishra C.
2014. Vigorous dynamics underlie a stable population of the endangered snow leopard
Panthera uncia in Tost Mountains, South Gobi, Mongolia. *PLoS One* 9:e101319.

Shehzad W, McCarthy TM, Pompanon F, Purevjav L, Coissac E, Riaz T, Taberlet P. 2012. Prey
preference of snow leopard (Panthera uncia) in South Gobi, Mongolia. *PLoS One*
7:e32104.

Sinnwell JP, Therneau TM, Schaid DJ. 2014. The kinship2 R package for pedigree data. *Hum.*
Hered. 78:91–93.

Suryawanshi KR, Khanyari M, Sharma K, Lkhagvajav P, Mishra C. 2019. Sampling bias in snow
leopard population estimation studies. *Popul. Ecol.* 61:268–276.

Suryawanshi KR, Redpath SM, Bhatnagar YV, Ramakrishnan U, Chaturvedi V, Smout SC,
Mishra C. 2017. Impact of wild prey availability on livestock predation by snow leopards. *R*
Soc Open Sci 4:170026.

Szpiech ZA, Blant A, Pemberton TJ. 2017. GARLIC: Genomic Autozygosity Regions Likelihood-
based Inference and Classification. *Bioinformatics* 33:2059–2062.

Tetzloff J, Schwartz KR. 2024. Chapter 26 - Role of zoos in snow leopard conservation: The
Species Survival Plan® in North America. In: Mallon D, McCarthy T, editors. Snow
Leopards (Second Edition). Academic Press. p. 333–346.

Thapa K, Schmitt N, Pradhan NMB, Acharya HR, Rayamajhi S. 2021. No silver bullet? Snow
leopard prey selection in Mt. Kangchenjunga, Nepal. *Ecol. Evol.* 11:16413–16425.

The Snow Leopard Trust. Gallery for Snow Leopard Trust. *Arcgis Gallery for Snow Leopard*
Trust [Internet]. Available from: snowleopard.maps.arcgis.com

Toyama KS, Crochet P-A, Leblois R. 2020. Sampling schemes and drift can bias admixture
proportions inferred by structure. *Mol. Ecol. Resour.* 20:1769–1785.

Tupa L. 2011. North American Regional Snow Leopard Studbook.

Van der Auwera GA, O'Connor BD. 2020. Genomics in the Cloud: Using Docker, GATK, and
WDL in Terra. "O'Reilly Media, Inc."

Wang C, Wu D-D, Yuan Y-H, Yao M-C, Han J-L, Wu Y-J, Shan F, Li W-P, Zhai J-Q, Huang M,

et al. 2023. Population genomic analysis provides evidence of the past success and future
potential of South China tiger captive conservation. *BMC Biol.* 21:64.

Weckworth B. 2021. Snow Leopard (Panthera uncia) Genetics: The Knowledge Gaps, Needs,
and Implications for Conservation. *J. Indian Inst. Sci.* 101:279–290.

Wegge P, Shrestha R, Flagstad Ø. 2012. Snow leopard Panthera uncia predation on livestock
and wild prey in a mountain valley in northern Nepal: implications for conservation
management. *Wildlife Biol.* 18:131–141.

Weir BS, Cockerham CC. 1984. Estimating F-Statistics for the Analysis of Population Structure.
Evolution 38:1358–1370.

Westbury MV, Hartmann S, Barlow A, Wiesel I, Leo V, Welch R, Parker DM, Sicks F, Ludwig A,
Dalén L, et al. 2018. Extended and Continuous Decline in Effective Population Size Results
in Low Genomic Diversity in the World's Rarest Hyena Species, the Brown Hyena. *Mol.*
Biol. Evol. 35:1225–1237.

Westbury MV, Petersen B, Garde E, Heide-Jørgensen MP, Lorenzen ED. 2019. Narwhal
Genome Reveals Long-Term Low Genetic Diversity despite Current Large Abundance
Size. *iScience* 15:592–599.

Xue Y, Prado-Martinez J, Sudmant PH, Narasimhan V, Ayub Q, Szpak M, Frandsen P, Chen Y,
Yngvadottir B, Cooper DN, et al. 2015. Mountain gorilla genomes reveal the impact of long-
term population decline and inbreeding. *Science* 348:242–245.

Zahler P, Victurine R. 2024. Chapter 11 - Linear infrastructure and snow leopard conservation.
In: Mallon D, McCarthy T, editors. Snow Leopards (Second Edition). Academic Press. p.
123–128.

Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. 2012. A high-performance
computing toolset for relatedness and principal component analysis of SNP data.
Bioinformatics 28:3326–3328.

Zhou C, Li F, Wen Q, Price M, Yang N, Yue B. 2021. Characterization of microsatellites in the
endangered snow leopard based on the chromosome-level genome. *Mamm. Res.* 66:385–
398.

Zhou C, Wang L, Qiao L, Lan Y, Price M, Yang N, Yue B. 2021. Characterization of Olfactory
Receptor Repertoires in the Endangered Snow Leopard Based on the Chromosome-Level
Genome. *DNA Cell Biol.* 40:293–302.

1053