Extreme in Every Way: Exceedingly Low Genetic Diversity in Snow Leopards Due to Persistently Small Population Size

Katherine A. Solari[1&], Simon Morgan[1], Andrey D. Poyarkov[2], Byron Weckworth[3], Gustaf Samelius[4,5], Koustubh Sharma[4,6], Stephane Ostrowski[7], Uma Ramakrishnan[8], Zairbek Kubanychbekov[9], Shannon Kachel[3], Örjan Johansson[4,10], Purevjav Lkhagvajav[11], Heather Hemmingmoore[10], Dmitry Y. Alexandrov[2], Munkhtsog Bayaraa[12], Alexey Grachev[13], Miroslav P. Korablev[2], Jose A. Hernandez-Blanco[2], Bariushaa Munkhtsog[14], Barry Rosenbaum[15], Viatcheslav V. Rozhnov[2], Ali Madad Rajabi[16], Hafizullah Noori[16], Ellie E. Armstrong[1,17]*, Dmitri A. Petrov[1,18,19]*&

[1]Department of Biology, Stanford University, Stanford, CA, USA
[2]A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Science, Moscow, Russia
[3]Panthera, New York, NY, USA
[4]Snow Leopard Trust, Seattle, WA, USA
[5]Nordens Ark, Hunnebostrand, Sweden
[6]Snow Leopard Foundation in Kyrgyzstan, Bishkek, Kyrgyz Republic
[7]Wildlife Conservation Society, New York, NY, USA
[8]National Centre for Biological Sciences, TIFR, Bangalore, India
[9]Ilbirs Foundation, Bishkek, Kyrgyzstan
[10]Grimsö Wildlife Research Station, Swedish University of Agricultural Sciences, Riddarhyttan, Sweden
[11]Snow Leopard Conservation Foundation, Ulanbaatar, Mongolia
[12]Wildlife Institute, College of Nature Conservation of the Beijing Forestry University, Beijing, China
[13]Institute of Zoology of Republic of Kazakhstan, Almaty, Kazakhstan
[14]Institute of General and Experimental Biology, Academy of Sciences of Mongolia, Ulaanbaatar, Mongolia
[15]Altai Institute for Research and Conservation, Boulder, CO, USA
[16]Wildlife Conservation Society, Afghanistan Program, Kabul, Afghanistan
[17]Washington State University, Pullman, WA, USA
[18]Chan Zuckerberg BioHub, San Francisco, CA, USA
[19]Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, USA
*Contributed equally
&Corresponding authors emails: ksolari@stanford.edu, dpetrov@stanford.edu

Abstract:

Snow leopards (*Panthera uncia*) serve as an umbrella species whose conservation benefits their high-elevation Asian habitat. Their numbers are believed to be in decline due to numerous Anthropogenic threats; however, their conservation is hindered by numerous knowledge gaps. They are the least studied genetically of all big cat species and little is known about their historic population size and range, current population trends, or connectivity across their range. Here, we use whole genome sequencing data for 41 snow leopards (37 newly sequenced) to assess population connectivity, historic population size, and current levels of genetic diversity. Among our samples, we find evidence of a primary genetic divide between the northern and southern part of the range around the Dzungarian Basin and a secondary divide south of Kyrgyzstan around the Taklamakan Desert. However, we find evidence of gene flow, suggesting that barriers between these groups are permeable. Perhaps most noteworthy, we find that snow leopards have the lowest genetic diversity of any big cat species, likely due to a persistently small population size throughout their evolutionary history. Without a large population size or ample standing genetic variation to help buffer them from any forthcoming Anthropogenic challenges, snow leopard persistence may be more tenuous than currently appreciated.

Introduction:

Residing in some of the most extreme and remote areas of the world, snow leopards (*Panthera uncia*) are rarely encountered and challenging to study, making them one of the most enigmatic of the large charismatic mammals. Snow leopard habitat consists of mountainous areas of Asia, spanning 12 countries (Fig. 1A). They are among the largest carnivores in the high-elevation habitat in which they reside and their persistence relies on healthy mountain ungulate populations which in turn rely on healthy rangelands[1]. Healthy high-elevation rangeland also benefits pastoral communities and offers immense ecosystem services, acting as an important source of carbon storage[2] and Asia's water tower. In these regions, snow leopards serve as an umbrella species whose conservation benefits this globally crucial Asian mountain ecosystem. However, snow leopard conservation is impeded by the many knowledge gaps regarding this elusive species[1].

Snow leopards were listed as Endangered by the International Union for Conservation of Nature (IUCN) for 45 years but were downlisted to Vulnerable in 2017 due to not meeting specific criteria of population size and percent population decline for Endangered status. This change of status has been controversial[3,4] as snow leopard numbers are presumed to be in

decline due to habitat loss, decreased availability of primary prey (high-elevation, mountain-dwelling ungulates), retaliatory killings for livestock predation[5], and poaching for their skins[6]. As climate change in high mountain Asia is occurring at an even more rapid rate than elsewhere in the Northern hemisphere[7], it is also likely to become an increasing threat to snow leopards[8]. Currently, the global snow leopard population size is estimated to be anywhere from 4,700-7,500 individuals and little is known about their historic population size and range[9] or their current population trends[10]. While many other big cat species experienced historic declines and are facing a second, contemporary human-driven decline[11,12], it remains unclear if the highly-specialized snow leopard was ever more abundant than is currently estimated. Estimates of genetic diversity, which are likely to correlate positively with historic population size, are also limited due to a dearth of genomic data for the species. They are the least studied genetically of all the big cat species[13] with whole-genome sequencing data available for only two wild snow leopards[14] and two captive individuals[15] prior to this study. Genetic diversity assessed using genomic data from one of the previously sequenced wild snow leopards suggests a lack of diversity[14], but additional samples are required to determine if this is a characteristic of the species across its range.

Additionally, little is known about snow leopard population connectivity across their range[16]. A potentially significant gap between northern and southern populations has been defined in the Dzungarian Basin of northwestern China[17,18]; however, snow leopards are known to cross long distances between mountain ranges[19,20]. To date, most genetic work on snow leopards has been done using microsatellite markers to estimate population sizes, connectivity, and diversity at mostly small spatial scales[21–26]. Using range-wide samples, however, Janečka et al.[27] analyzed 33 microsatellite loci and 683 bp of mitochondrial DNA from fecal samples to suggest three snow leopard subspecies–one of them occurring north of the Dzungarian Basin and Gobi Desert and two subspecies south of this divide, separated between the east and west of the Himalayas-Tibetan Plateau complex. However, this subspecies demarcation and the level of connectivity across the landscape is controversial[28,29]. It remains unclear if any landscape features within the snow leopard range are impermeable or if all snow leopard populations are connected to some degree through stepping-stone and/or long distance dispersal events.

Here, we generate whole genome sequencing data for 33 wild snow leopards from multiple locations across their range in addition to four captive snow leopards. We use this dataset to shed light on two outstanding questions: 1) how connected are snow leopard populations across their range and 2) what was the historic population size of snow leopards and how is this reflected in their current levels of genetic diversity. We also compare the genetic

diversity of snow leopards to that of the other big cats. Among our samples, which do not include the southeast of the range, we find evidence of a genetic divide between the north and south, as well as a divide between the Kyrgyz population and populations farther south. Our data shows evidence of contemporary gene flow, suggesting that barriers between these groups are permeable. Most notably, we find snow leopards to be the least genetically diverse contemporary big cat species, likely owed to a persistently small population throughout their evolutionary history. We believe these results have significant implications for snow leopard conservation.

Results:

We generated whole genome sequencing data for 37 snow leopards–33 wild born snow leopards representing seven countries (Mongolia, Russia, Kyrgyzstan, Tajikistan, Afghanistan, and Pakistan) (Fig. 1A) and four from the American zoo population. We also included all published whole genome sequencing data for snow leopards in our analysis, which consisted of data for two additional wild samples, one from Mongolia[14] and one from India, and two captive individuals from the American zoo population[15]. After removing samples based on sequence quality and depth we were left with 37 samples–34 from our newly generated data with an average depth of coverage of 7.3X (minimum of 3X and maximum of 16.8X) and three previously published samples with an average depth of coverage of 23X (minimum of 12X and maximum of 28.9X) (Supplementary Table 1).

These data were mapped to the snow leopard reference genome (NCBI accession PRJNA602938)[15] and single nucleotide polymorphisms (SNPs) were called and filtered as described in the methods, resulting in a final SNP set of 1,591,978. Within this final dataset, we identified one pair of first-degree relatives and three pairs of second-degree relatives (all from the wild samples). One representative from each of these four related pairs was removed before conducting analyses that could be impacted by the presence of related individuals as indicated in the methods.

Population structure and dispersal barriers:

We assessed population structure among our samples using principal component analysis (PCA). PCA results indicated that the one Indian sample was genetically distinct from all of the other samples (Fig. 1C). Once removing two outlier samples, three distinct groups became evident–Mongolia and Russia; Kyrgyzstan and captive; and Tajikistan, Afghanistan and Pakistan (Fig. 1D).

We also investigated population structure using Admixture[30] to test models with one to ten ancestral populations (K=1-10) over ten iterations. Admixture analyses can be sensitive to differences in sample sizes among groups[31], so the India sample was not included in this analysis. At K=2, Mongolia and Russia separated clearly from all other samples, with the same ancestry assignments supported in all ten iterations. At K=3, nine of the ten iterations supported a clear separation of the samples into three groups–Mongolia and Russia; Tajikistan, Afghanistan, and Pakistan; captives and Kyrgyzstan (Fig. 1B), recapitulating what we observe in the PCA. These Admixture and PCA results are robust to the removal of captive samples from the analyses (Supplementary Fig. 3).

We used EEMS (Estimated Effective Migration Surfaces)[32] to assess migration patterns. EEMS results were consistent with PCA and Admixture analyses and also indicated a geographic barrier to migration from Kyrgyzstan to the northeast, around the Dzungarian Basin, and to the south, around the Taklamakan Desert (Fig. 1E).

Taken together, our results suggest three genetically distinct groups within our samples (excluding India). We see a primary genetic divide between the north (Mongolia and Russia) and the south (all other samples) with a secondary divide within the southern group between Kyrgyzstan and populations farther south (Afghanistan, Tajikistan, and Pakistan).

Assessment of gene flow:

Based on Admixture and PCA results, we quantified population structure between the two groups identified in Admixture at K=2 and among the three groups identified at K=3 (Fig. 1B) by assessing shared versus private SNPs, F_{ST}, and rare allele sharing. Captive samples and the Indian sample were excluded from these analyses. One sample from each related pair was excluded from F_{ST} and rare allele sharing analyses, but not shared versus private SNP assessments.

At K=2, each group had more shared SNPs (600,052 SNPs) than private SNPs (415,931 SNPs private to the north and 362,407 SNPs private to the south) (Supplementary Fig. 5A). At K=3, we compared groups that we will refer to as "North" (consisting of Russia and Mongolia), "Kyrgyzstan", and "Far South" (consisting of Tajikistan, Afghanistan and Pakistan). The North group was downsampled to eight individuals such that all three groups had a similar sample size (Far South N=8, Kyrgyzstan N=7). We found that the Far South and North group had more than twice as many private SNPs (234,825 and 224,417, respectively) than Kyrgyzstan (109,016) (Supplementary Fig. 5B) and that Kyrgyzstan shared a similar number of SNPs with both the North and the Far South (398,843 SNPs shared among all three groups, 64,319 shared

only between Kyrgyzstan and North, 63,286 SNPs shared only between Kyrgyzstan and Far South). It is worth noting that the Kyrgyz group represents the smallest geographic area and this limitation in sampling could contribute to our observation of fewer private SNPs in this group.

Pairwise F_{ST} analyses indicated low levels of differentiation among the groups. Weir and Cockerham's pairwise F_{ST} was calculated after removing one individual for each first or second degree related pair. At K=2, F_{ST} between the north and south was 0.091. At K=3, the pairwise F_{ST} between the North and Far South was 0.123, between North and Kyrgyzstan was 0.108, and between Kyrgyzstan and the Far South was 0.091.

Doubletons (SNPs with an alternate allele count of two), where each alternate allele occurs in separate individuals, were identified. Doubleton sharing was then assessed among the three groups–North, Kyrgyzstan, and Far South–after downsampling each group to six individuals as in[33]. The amount of doubleton sharing between populations can offer a sense of population connectivity as diverged populations have very little rare variant sharing[34]. Observed doubleton sharing was compared to a null distribution representing panmixia created by permuting the data 10,000 times. Doubleton sharing corroborated the pre-identified groups with individuals sharing a significantly higher fraction of doubletons (0.40-0.72) with individuals within the same group (p-value North - 0.0006, Kyrgyzstan - 0.008, Far South - 0.01) (Figure 1F). However, we also found that each individual shares 11-32% of their doubletons with each other group, with the Kyrgyzstan population showing the highest fraction of doubleton sharing with outside groups - even sharing enough with the Far South to not be significantly different from panmixia (p = 0.2). Both Kyrgyzstan and Far South also share enough rare alleles with the North to only be marginally significantly different from panmixia (p=0.04 and p=0.03, respectively). These results support the presence of genetic divergence among these three groups, but also indicate a lack of any absolute barriers as there is some level of gene flow among these groups.

Figure 1. Snow leopard distribution and sample map, Admixture, principal components analysis (PCA), estimated migration surfaces, and rare allele sharing. A) The IUCN snow leopard distribution is shown and sample locations are indicated with different sized circles indicating the number of samples from each location. The basemap indicates elevation and country outlines are shown in grey. B) Admixture results for ten independent runs for K=2 and K=3 distinct ancestry groups. The ancestry assignments shown for K=2 were supported by all ten iterations and the ancestry assignments shown for K=3 were supported by nine of the ten iterations. C) PCA of genetic variation of 37 snow leopards using 1,448,657 SNPs. D) PCA after removing two outlier samples, India and sample U13 from Tajikistan. PCA axis labels include the percent variation explained by PC1 and PC2. E) Estimate of migration surfaces based on 1,343,552 autosomal SNPs for 31 snow leopards. Log 10-transformed effective migration rates are shown where green indicates areas with relatively higher migration and brown indicates areas with relatively lower migration. Black circles represent sample locations as assigned to their closest deme. Country outlines are shown in solid gray and landscape features are shown

in dashed black lines and labeled (Dzungarian Basin (DB), Taklamakan Desert (TD), Gobi Desert (GD)). F) Doubleton sharing between populations compared to null distributions under panmixia. We identified all doubletons where each allele occurred in a different individual. For each group, identified above each graph, we then calculated the fraction of doubletons occurring in an individual of that group that were shared with individuals of each other group. Null distributions were made by permuting the data 10,000 times. P-values for comparisons between observed data and the null distribution using the Wilcoxon Rank Sum Test are shown. The lower and upper edges of the boxes correspond to the first and third quartiles and the whiskers extend to the lowest/highest value that is no further than 1.5*IQR (inter-quartile range) from the box. Points falling further than 1.5*IQR from the box are plotted individually.

Heterozygosity and historic population size:

We used publicly available data to call SNPs in all big cat species using the Genome Analysis Toolkit (GATK)[35] and calculated heterozygosity using VCFtools[36]. Here, we defined big cat species as species with an average adult body weight of 40kg or more, which includes all Panthera species (lion (*Panthera leo*), tiger (*Panthera tigris*), leopard (*Panthera pardus*), jaguar (*Panthera onca*), and snow leopard) as well as cheetah (*Acinonyx jubatus*) and puma (*Puma concolor*) [37]. We found snow leopards to have the lowest heterozygosity of any big cat species, with heterozygosity for every snow leopard sample lower than that observed in any other big cat (Fig. 2A). Notably, snow leopard heterozygosity was lower than that of cheetahs, which have long been considered the archetype of low heterozygosity in big cats[11,38]. The relative values of observed heterozygosity among all the other species for which we calculated heterozygosity were consistent with previous work[12,39–41]. Within snow leopards, we found observed heterozygosity to be consistently low across all sampled locations (Fig. 2B).

We used pairwise sequentially Markovian coalescent (PSMC)[42] to reconstruct historic effective population size using the highest coverage individual from each genetically distinct group (North-28X, Kyrgyzstan/captive-28X, South-9X, India-12X). Reconstructions showed consistent results across all populations sampled and across all bootstrap replicates (Fig. 2C). All reconstructions show a consistently small effective population size over the last ~300,000 years with snow leopard effective population size never exceeding 14,000 individuals. Reconstructions suggested a snow leopard effective population size of 8,000-6,000 individuals from ~200,000-20,000 years ago and a decrease to an effective population size of ~2,500-4,000 individuals ~20,000 years ago, potentially coincident with the end of the last glacial maximum. Effective population size is generally smaller than census size[43] and PSMC historic effective population size estimates can be impacted by numerous species-specific parameters (e.g., population structure, inbreeding, mutation rate, generation time)[44,45] which have not been

thoroughly characterized in snow leopards, and thus exact effective population size estimates should be interpreted cautiously.

Figure 2. Genome-wide heterozygosity and demographic history. A) Comparison of genome-wide heterozygosity across all big cat species. Publicly available data was used to call SNPs for every big cat species using GATK and observed heterozygosity was calculated using VCFtools. In the case of leopard and tiger, SNPs were called separately for genetically distinct groups. Snow leopard heterozygosity is calculated from SNPs called using the same in-house pipeline that was used for SNP calling in all the other big cat species B) Genome-wide heterozygosity of snow leopards separated by geographic location. Genome-wide heterozygosity shown is calculated from SNPs called by Gencove. A direct comparison of heterozygosity calculated from SNPs called in-house versus SNPs called by Gencove is shown in Supplementary Fig. 6. The lower and upper edges of the boxes correspond to the first and third quartiles and the whiskers extend to the lowest/highest value that is no further than 1.5*IQR (inter-quartile range) from the box. Points falling further than 1.5*IQR from the box are plotted individually. C) Reconstruction of effective population sizes using PSMC using a mutation rate of 0.7×10^{-8} per site per generation and a generation time of five years. Thirty bootstraps are shown for each sample in thinner, fainter lines of the same color.

Discussion:

Heterozygosity and demographic history:

We used whole genome sequencing from 37 snow leopards to investigate genetic diversity, demographic history, and population structure of the species. Our results show that snow leopards have the lowest genetic diversity of any big cat species (Fig. 2A), lower than previously appreciated[12,46]. Demographic history assessments indicated the exceptionally low heterozygosity in snow leopards is likely due to a persistently small population size over the last 500,000 years (Fig. 2C). With a persistently small population size and low genetic diversity throughout their evolutionary history, snow leopards serve as yet another example that high genetic diversity is not a requirement for the long-term persistence of a species[47–50]. In light of our results, very low population densities, as estimated in many contemporary landscapes (<0.5 animals/100 km2; e.g.,Suryawanshi et al.[51]), appears to be a familiar condition in snow leopard evolutionary history.

Population structure and dispersal barriers:

PCA analyses show that India is genetically distinct from all other samples (Fig. 1C). The uniqueness of this sample could suggest that this individual is our one representative of the third phylogenetic group suggested by Janečka et al.[27], but more samples from this area will be necessary to resolve the unique evolutionary position of Indian snow leopards in future analyses.

Among the other samples, we identify three genetically distinct groups. Structure and PCA results identify the most pronounced divide among our samples to occur between the

northern and southern part of the range around the Dzungarian Basin (Fig. 1B,D), consistent with previous microsatellite analyses[27] and resistant kernel models[18]. Our Structure and PCA results also identify a secondary divide occurring south of Kyrgyzstan (Fig. 1B,D), around the Taklamakan Desert, consistent with previous microsatellite analyses[52]. Both of these genetic divides were also apparent in effective migration surface maps (Fig. 1E).

These results indicate that the Dzungarian Basin and Taklamakan Desert are effective barriers to dispersal for snow leopards. However, low levels of differentiation – measured through F_{ST}, shared versus private alleles (Supplementary Fig. 5), and shared rare allele (Fig. 1F) analyses – indicate that there is likely still some level of functional connectivity among these three distinct groups despite intervening lowlands and large geographic distances between them. Additional samples will be necessary to confidently estimate gene flow in future analyses.

Origin of captive samples:

Studbook-based pedigrees for the five captive individuals sequenced in this study show that the origin of their wild ancestors is mostly unknown. The few ancestors of known origin are documented as originating from the USSR, Kyrgyzstan, Kazakhstan, and the western Himalayas (Supplementary Fig. 1). Admixture and PCA results both show a strong signal of genetic relatedness between the five captive individuals and the current Kyrgyzstan population (Fig. 1B,D). These results suggest that many of the wild ancestors listed as coming from the USSR or unknown wild locations likely came from the Kyrgyzstan region.

Conservation implications:

With these data, we have begun to visualize the geographic distribution of genomic diversity in snow leopards. Though this analysis does not contain samples from every part of the snow leopard range (i.e., China and eastern Himalayas), it provides substantial insight into the genetic divides across the portion of the range that we have sampled. We also provide evidence that even the most expansive low-elevation areas are likely circumventable or traversable by snow leopards, as our results suggest gene flow across all regions sampled. This indicates that as human development continues and border fencing proliferates in this area[17,53], it will be important to maintain the likely narrow and dispersed corridors between distant patches of snow leopard habitat to maintain existing levels of connectivity[17].

Snow leopards live in arid, cold, low-productivity, high-elevation habitats where few species can persist – an environment that has evidently only ever been able to support a limited number of snow leopards. As a result, they have likely always had an effective population size

much lower than other big cats (as suggested in Pečnerová et al.[12] and Cho et al.[14]), and harbor less genetic diversity than even the cheetah.

Thanks to their extreme environment, snow leopards have not yet been exposed to the same level of acute Anthropogenic pressures as have big cats living in habitats more easily accessible to humans. Yet, even having been spared the most intense human impacts, they already have extremely low population size and genomic diversity. While these characteristics have not hindered their long-term persistence thus far, this means that snow leopards do not have the ability to rely on a large population size or standing genetic variation to help them survive any forthcoming Anthropogenic challenges, as other big cats have done[54,55]. Unfortunately, it is likely that the most intense Anthropogenic pressures may lie ahead for snow leopards. In addition to their ongoing threats, Anthropogenic climate change threatens to shrink snow leopard range through habitat change[8] and intensify interspecific competition (e.g. Pal et al.[56], Lovari et al. [57]); shifting grazing practices risk facilitating spillover of novel pathogens from domestic animals into snow leopards and their prey[58]; and accelerating mining, energy, and infrastructure development threaten to fragment and degrade previously remote snow leopard habitats[59,60]. Our results reveal exceptionally low genetic diversity in the species, and make clear that the persistence of snow leopards is likely tenuous. Protection of snow leopards and their habitat, to the greatest degree possible, will be pivotal as this specialist, low-density carnivore appears genetically and demographically ill-equipped to bounce back from Anthropogenic perturbations[61]. More broadly, our findings underscore how the IUCN Red List assessment criteria could benefit from the inclusion of genetic factors[62] as our results suggest that snow leopards may be more imperiled than their current IUCN Red List status would indicate[3,63].

Methods:

Sample Collection:

Through an international collaborative effort, a total of 37 snow leopard blood or tissue samples were collected, composed of 33 wild caught samples from six countries (Afghanistan, Kyrgyzstan, Tajikistan, Mongolia, and Russia), one currently captive but wild-born individual from Pakistan, and four captive samples from mixed ancestry. Sample details can be found in Supplementary Table 1. We were unable to include any samples from a large portion of the snow leopard range in the southeast, but are hopeful that data from this area will be available to include in future analyses. All samples were sequenced on an Illumina sequencing platform using paired end 150 bp reads. Additionally, we collected all currently published whole genome

sequencing data for snow leopards, which included data for two captive individuals (NCBI accessions SRR16227515[15] and SRR12437590), one wild individual from Mongolia (NCBI accession SRR836372[14]), and one wild individual from India (NCBI BioProject PRJNA1051290). In total, we gathered whole genome sequencing data for 41 snow leopards.

Note that there is whole genome sequencing data for an additional sample identified as a snow leopard on Genbank (biosample SAMN17432540, SRA accession SRR13500277, used in the following publications: Zhou et al.[64,65]) that we did not use in this study because we concluded that this data was from a leopard (*Panthera pardus*), not a snow leopard (details of the analyses done to identify this sample as a leopard are provided in Supplementary File 1).

Mapping and variant calling:

We used the snow leopard reference genome (NCBI accession PRJNA602938[15]) and added the lion Y chromosome (from GCA_018350215.1) in order to have the Y-chromosome represented in this female reference genome. Reads were mapped and variants called by Gencove using BWA-MEM[66] and the Genome Analysis Toolkit (GATK)[67], respectively. The depth and breadth of coverage for each sample were calculated from bam files using SAMtools[68] and are provided in Supplementary Table 1.

Variant filtering:

Variants were first filtered based on mappability across the genome. The reference genome was indexed and mappability scores generated using genmapv1.3.0[69] with flags '-K 30' and '-E 2'. The reference genome and mappability bed file were sorted using BEDTools[70] and SNPs falling in a region of the genome with a mappability score less than one were removed using BEDTools *intersect* with the flags '-v', '-header', and '-sorted' (leaving 63,199,070 sites). We then used VCFtools[36] to remove indels using the option '--remove-indels' (leaving 51,199,725 SNPs). Individuals with less than 60% of the genome represented (samples U02, U12, and U20) were removed using the VCFtools option '--remove-ind'. We then used VCFtools to remove sites that no longer had any variation using flag '--non-ref-ac-any 1' (leaving 2,379,069 SNPs), SNPs that did not fall on putative autosomes (leaving 2,213,174 SNPs), and non-biallelic SNPs using th flags '--min-alleles 2 --max-alleles 2' (leaving 2,146,294 SNPs). We also split scaffold 22 at base pair 67,650,000 into chromosomes E1 and F1, respectively. Next, we used GATK v4.1.4.1 to add the following annotations to our filtered vcf file using *VariantAnnotator*: QD (QualByDepth - variant confidence normalized by unfiltered depth of variant samples), FS (FisherStrand - strand bias estimated using Fisher's exact test), MQ

(RMSMappingQuality - root mean square of the mapping quality of reads across all samples), MQRankSum (MappingQualityRankSumTest - rank sum test for mapping qualities of reference versus alternate reads), and ReadPosRankSum (ReadPosRankSumTest - rank sum test for relative positioning of reference versus alternate alleles within reads). After adding these annotations, we used GATK *VariantFiltration* to filter SNPs using the following flag: '--filter-expression "QD < 2.0 || FS > 60.0 || MQ < 40.0 || MQRankSum < -12.5 || ReadPosRankSum < -8.0"'. We then used VCFtools to remove SNPs that did not pass the GATK *VariantFiltration* step using option '--remove-filtered-all' (leaving 2,065,453 SNPs). Lastly, we removed SNPs missing data in more than 10% of the individuals using flag '--max-missing 0.9' (leaving a final SNP set of 1,591,978). Singletons and private doubletons were calculated using the VCFtools flag '--singletons' and one of the captive samples (10x_SDzoo) was removed due to an excess of singletons (Supplementary Table 1) which we suspected to be due to the 10X library prep method unique to this sample.

Relatedness assessment:

Relatedness among samples was estimated using SNPrelate[71] in R[72]. We divided the samples into three genetically distinct groups based on preliminary PCA and Admixture analyses–Mongolia and Russia, Kyrgyzstan and captive, and all samples South of Kyrgyzstan. Preliminary PCA and Admixture analyses were done using the same methods described below but in this case included samples that were later removed due to relatedness. Each group was run through SNPrelate separately. We used the function *snpgdsBED2GDS* to convert bed files to gds files. We used *snpgdsLDpruning* to prune the input data using the following flags - 'method="corr", slide.max.n=50, ld.threshold=0.3, maf = 0.05, autosome.only = F'. We then calculated kinship coefficients using the function *snpgdsIBDMLE* which calculates IBD coefficients for individual pairs using Maximum Likelihood Estimation. A kinship coefficient of 0.5 indicates a duplicate sample, 0.25 indicates a first-order related pair (parent-offspring or full sibling), and 0.125 indicates a second-order related pair (half-siblings or grandparent-grandchild). All sample pairs with non-zero kinship coefficients are listed in Supplementary Table 2. We identified one sample pair with a kinship coefficient greater than that expected from first-order relatives–U01 and U09 (kinship coefficient of 0.335), and we identified three sample pairs with kinship coefficients consistent with second-order related pairs–SL_KGZ_F1 and SL_KGZ_F4 (kinship coefficient of 0.148), AF_SL_07 and AF_SL_06 (kinship coefficient of 0.135), and U14 and U08 (kinship coefficient of 0.124). The sample with lower sequencing

coverage from each of these pairs (U01, SL_KGZ_F4, AF_SL_06, and U08) was removed from indicated analyses.

Pedigree-based measures of relatedness in captive-bred samples:

Using snow leopard studbooks[73,74], we compiled information on all of the ancestors of the five captive-born individuals included in this study. The wild origin of their ancestors is mostly unknown; however, each sample has a few ancestors for which there is a wild origin location listed (Supplementary Fig. 1), these origin locations include wild ancestors listed as coming from the USSR (between 1964-1974), Kazakhstan (in 1972), Kyrgyzstan (between 1974-1980), Prezewalsk (between 1974-1975, which we believe to refer to Karakol, Kyrgyzstan which was previously named Prezewalsk), and Aksai (a contested region between China and India in the Western Himalayas, in 1979).

Pedigrees for each individual were constructed (Supplementary Fig. 1) and kinship was estimated (Supplementary Table 3) among these individuals using the kinship2 package[75] in R. Kinship calculated from pedigrees was compared to kinship calculated from SNPrelate using Pearson correlation coefficient calculated using the ggpubr[76] package in R and plotted using ggplot2[77]. We found kinship calculated from pedigrees to be highly correlated with kinship calculated from SNPrelate (R = 0.91, p = 0.00023, Supplementary Fig. 2).

Principal component analysis:

Principal component analyses (PCAs) were conducted on the dataset after filtering for first and second degree relatives (N=33, 1,448,657 SNPs) using PLINK2[78]. First, bed files were generated from vcf files using the flags '--make-bed' and '--allow-extra-chr' and allele frequencies were computed using the flags '--freq' and '--allow-extra-chr'. Then, the principal component analysis was run by inputting the resultant bed file with flag '--bfile' and frequency file with flag '--read-freq' and using the flags '--allow-extra-chr' and '--pca 38'. PCA output was visualized using ggplot2 in R (Fig. 1C,D).

Admixture analysis:

Admixture analyses were also conducted on the dataset after filtering for first and second degree relatives. Since these analyses can be sensitive to uneven sample size, the Indian sample was excluded from this analysis (N=32)[31,79]. We used VCFtools to convert vcf files into PLINK format files. We then used PLINK1.9 to recode alleles numerically using the flags '--allow-extra-chr --recode12'. We then ran Admixture[30] for K1-10 ten times using the flag

'-s time' so that a different random seed was used for each run based on the time. All runs were then visualized using Clumpak[80] (Fig. 1B).

PCA and Admixture analyses were also conducted on the dataset after the removal of captive samples in order to assess the impact of captive samples on population structure results (Supplementary Fig. 3).

Estimate of Effective Migration Surfaces (EEMS):

We used a matrix of average pairwise genetic dissimilarities as input into EEMS[32] to estimate effective migration and diversity surfaces across the snow leopard range. To make this input file from the filtered vcf file, we used VCFtools to remove all captive-bred samples, the one Mongolia sample (SRR836372) lacking location data, and one individual per first or second degree related pair (U01, U08, AF_SL_06, SL_KGZ_F4) (N=27). We then removed sites for which there was no longer any variation present in the remaining samples and were left with 1,343,552 SNPs as input into EEMS. We then used PLINK1.9 to create a bed file from the vcf using the '--make-bed' flag. Lastly, we used *bed2diffs* in EEMS to generate the pairwise dissimilarity matrix from the bed file. The habitat outline used was selected to include the entire extant range of snow leopards as shown in Fig. 1A and was drawn manually using arcgis online. We ran *runeems_snps* with 8 million MCMC iterations, 1 million burn-in iterations, and 9999 iterations thinned between writing steps. We used grids of 200, 400 and 600 demes and averaged the results across three independent realizations at each deme size. Through numerous initial trial run iterations, we adjusted the proposed variance of parameters such that the accepted proportions of proposed variances all fell between 10-40%. Results of all nine independent runs were visualized together in R using the package rEEMSplots[32] (Fig. 1E).

We also ran EEMS on the microsatellite dataset from[27] which is composed of 32 variable microsatellite loci genotyped for 70 individuals. We analyzed this dataset using *runeems_sats* using the same parameters as above - 8 million MCMC iterations, 1 million burn-in iterations, 9999 iterations thinned between writing steps, and grid sizes of 200, 400, and 600 demes. Each deme size was run three independent times and results were averaged across all nine realizations (Supplementary Fig. 4).

Quantifying population structure:

We further characterized population divides identified in Admixture and PCA analyses by calculating the number of shared versus private SNPs among groups, pairwise F_{ST}, and the rate of rare variant sharing among groups. Captive samples were excluded from these analyses as

was the sample from India as the PCA indicates that this sample is genetically distinct from all groups.

We did not filter for relatedness when calculating shared versus private SNPs among the groups; however, we did downsample the Northern group at K=3 such that all groups had between 7-8 individuals. We used BCFtools[81] to create and index vcf files with only the individuals from each group using the *view* and *index* function, respectively. We then used BCFtools *isec* to calculate how many SNPs were shared and private among the groups.

Before calculating F_{ST}, we removed one individual for each first or second degree related pair. We first used PLINK1.9 to create transposed PLINK files using the flags '--allow-extra-chr --double-id --recode --transpose'. We then used PLINK2 to calculate F_{ST} among groups using the Weir and Cockerham estimator[82] using the flags '--allow-extra-chr --fst CATPHENO method=wc'.

To assess doubleton sharing among groups, we excluded one individual for each related pair and downsampled all groups so that each had an equal number of individuals (N=6). We then used VCFtools to retain only SNPs with a minor allele count of two (doubletons) using the flags '--mac 2 --max-mac 2'. We removed any SNPs where the doubletons were present in the same individual and identified allele sharing among samples using the flag '--genome full' in PLINK. A null distribution under panmixia was created by permuting the data 10,000 times. Fractions of observed doubleton sharing were compared to the null distribution using the Wilcoxon Rank Sum Test in R.

PSMC:

We used PSMC[42] to estimate snow leopard population size back in time using the highest coverage sample for each population cluster. We generated diploid consensus sequences for each individual as recommended by PSMC. Starting with bam files filtered to include only putative autosomes, we used SAMtools *mpileup* and BCFtools *call* to generate a vcf and then used *vcfutils.pl vcf2fq* to generate diploid fastq files. We then ran PSMC with the default settings. We also performed 30 rounds of bootstrapping using random sampling with replacement for each sample. For plotting, we used a mutation rate of 0.7×10^{-8} per site per generation as suggested for snow leopards by Cho et al.[14] and a generation time of five years.

Heterozygosity:

Observed homozygosity was calculated for every individual from the vcf file using VCFtools[36] with the flag '--het'. In R, we calculated the number of heterozygous sites by

subtracting the number of observed homozygous sites column ((O)HOM) from the total number of sites column (NSITES). We then calculated heterozygosity by dividing the number of heterozygous sites by the length of the genome consisting of putative autosomes with a mappability score of one (1,818,166,894). Heterozygosity was calculated for all individuals, regardless of relatedness (N=37). We used ggplot2 in R to create box plots of heterozygosity results. All snow leopard samples had a heterozygosity between 0.000039 - 0.000168 (Fig. 2B).

Heterozygosity across big cats:

Heterozygosity was calculated in all big cat species using publicly available data in order to see how snow leopard heterozygosity levels compare to other big cat species. We included all other species in the genus *Panthera* (leopard, lion, tiger, jaguar) as well as cheetah and mountain lion. Accession numbers of publicly available whole genome sequencing data and reference genomes used in this analysis are listed in Supplementary Table 4 and 5, respectively. We also recalculated heterozygosity in all snow leopard samples using the same in-house pipeline to ensure that heterozygosity calculated using SNPs called with the Gencove pipeline were comparable to heterozygosity calculated using SNPs called with our own pipeline.

All fastq data was mapped to the corresponding reference genome using BWA-MEM[66]. Resulting bam files were sorted and indexed using SAMtools[68]. Depth of coverage and breadth of coverage was calculated for each sample using SAMtools. We added read groups and marked duplicates using picard[83]. We then used GATK[35] *HaplotypeCaller* to generate a gvcf file for each sample and GATK *CombineGVCFs* to combine the gvcf files for every species (or subgroup in the case of tigers and leopards). Lastly, we used GATK *GenotypeGVCFs* to create a final vcf file for every group.

In order to limit the need for excessive computing power, some samples with over 30X coverage were downsampled using SAMtools *view* as indicated in Supplementary Table 4. The leopard samples required a larger amount of computing power than the other species to call haplotypes, so these samples were all split into one bam file per contig using BAMtools *split* and run through GATK *HaplotypeCaller* and *CombineGVCFs* in parallel. We also found it necessary to split the jaguar and snow leopard data into intervals to combine gvcf files using GATK *GenomicsDBImport* followed by *CombineGVCFs*. In both cases, resultant vcf files, one per contig or interval, were then concatenated using BCFtools *concat*.

Each reference genome was filtered for mappability using genmapv1.3.0 to first index and then calculate mappability using flags '-K 30' and '-E 2'. We then filtered each vcf to only include nucleotides with a mappability score of one. To do this, vcf files and mappability bed

files were both sorted using BEDTools and only SNPs falling in regions with a mappability score of one were retained in each vcf using BEDTools *intersect*. Each vcf was then filtered using GATK *VariantFiltration* with the following flag: '--filter-expression "QD < 2.0 || FS > 60.0 || MQ < 40.0 || MQRankSum < -12.5 || ReadPosRankSum < -8.0"'. Then, vcf files were filtered to remove indels and non-bialellic SNPs using VCFtools flags '--remove-indels --min-alleles 2 --max-alleles 2'.

The mappable length of each genome was calculated using mappability bed files to calculate the number of nucleotides with a mappability score of one. Using the filtered vcf for each species or group, observed heterozygosity was then calculated for each sample and plotted in the same way described above.

Snow leopard heterozygosity calculated from SNPs called using our in-house pipeline was compared to heterozygosity calculated from SNPs called by Gencove using Pearson correlation coefficient calculated using the ggpubr[76] package in R (Supplementary Fig. 6). We found heterozygosity calculated from the two different SNP calling pipelines to be extremely correlated (R = 0.98, p=2.2E-16); however, all heterozygosity estimates calculated from our in-house SNP calls were slightly higher (average of ~0.000041) than that estimated from Gencove SNP calls.

Data Availability:

Data associated with this study has been deposited into bioproject PRJNA1048427 and will be released upon publication.

Code Availability:

The code used for analyses in this project is available on the project's github: https://github.com/ksolari/SL_WGS

Acknowledgements:

We would like to thank those who helped us in acquiring the captive samples used in this study, including Karen Ingerman and Susie Bartlett from the Bronx Zoo. We would like to thank Haiqing Xu for his help in conducting simulations to produce the null distribution for the shared doubleton analysis.

References:

1. *Snow Leopards: Biodiversity of the World: Conservation from Genes to Landscapes*. (Academic Press, 2016).
2. Mu, C. *et al.* The status and stability of permafrost carbon on the Tibetan Plateau. *Earth-Sci. Rev.* **211**, 103433 (2020).
3. Ale, S. B. & Mishra, C. The snow leopard's questionable comeback. *Science* **359**, 1110 (2018).
4. Aryal, A. Poaching: Is snow leopard tally underestimated? *Nature* **550**, 457 (2017).
5. Jackson, R. M., Mishra, C., McCarthy, T. M. & Ale, S. B. Snow leopards: conflict and conservation. *The biology and conservation of wild felids* **2**, 417–430 (2010).
6. Network, S. L. Snow leopard survival strategy. *Seattle, Washington, USA* 1–145 (2014).
7. Lalande, M., Ménégoz, M., Krinner, G., Naegeli, K. & Wunderle, S. Climate change in the High Mountain Asia in CMIP6. *Earth Syst. Dyn.* **12**, 1061–1098 (2021).
8. Li, J. *et al.* Climate refugia of snow leopards in High Asia. *Biol. Conserv.* **203**, 188–196 (2016).
9. McCarthy, T., Mallon, D. & Zahler, P. Chapter 3 - What is a snow leopard? Biogeography and status overview. in *Snow Leopards (Second Edition)* (eds. Mallon, D. & McCarthy, T.) 31–41 (Academic Press, 2024).
10. McCarthy, T., Mallon, D., Jackson, R., Zahler, P. & McCarthy, K. *Panthera uncia*. *The IUCN Red List of Threatened Species* **e.T22732A50664030.** (2017).
11. Dobrynin, P. *et al.* Genomic legacy of the African cheetah, *Acinonyx jubatus*. *Genome Biol.* **16**, 277 (2015).
12. Pečnerová, P. *et al.* High genetic diversity and low differentiation reflect the ecological versatility of the African leopard. *Curr. Biol.* **31**, 1862–1871.e5 (2021).
13. Weckworth, B. Snow Leopard (*Panthera uncia*) Genetics: The Knowledge Gaps, Needs, and Implications for Conservation. *J. Indian Inst. Sci.* **101**, 279–290 (2021).
14. Cho, Y. S. *et al.* The tiger genome and comparative analysis with lion and snow leopard genomes. *Nat. Commun.* **4**, 2433 (2013).
15. Armstrong, E. E. *et al.* Genome report: chromosome-level draft assemblies of the snow leopard, African leopard, and tiger (*Panthera uncia*, *Panthera pardus pardus*, and *Panthera tigris*). *G3* **12**, (2022).
16. McCarthy, T., Mallon, D., Sanderson, E. W., Zahler, P. & Fisher, K. What is a Snow Leopard? Biogeography and Status Overview. in *Snow Leopards* (eds. McCarthy, T. & Mallon, D.) 23–42 (Elsevier, 2016).
17. Li, J. *et al.* Defining priorities for global snow leopard conservation landscapes. *Biol. Conserv.* **241**, 108387 (2020).
18. Riordan, P., Cushman, S. A., Mallon, D., Shi, K. & Hughes, J. Predicting global population connectivity and targeting conservation action for snow leopard across its range. *Ecography* **39**, 419–426 (2016).
19. Sharma, K. *et al.* Vigorous dynamics underlie a stable population of the endangered snow leopard *Panthera uncia* in Tost Mountains, South Gobi, Mongolia. *PLoS One* **9**, e101319 (2014).

20. McCarthy, T. M., Fuller, T. K. & Munkhtsog, B. Movements and activities of snow leopards in Southwestern Mongolia. *Biol. Conserv.* **124**, 527–537 (2005).
21. Janečka, J. E. *et al.* Population monitoring of snow leopards using noninvasive collection of scat samples: a pilot study. *Anim. Conserv.* **11**, 401–411 (2008).
22. Janečka, J. E. *et al.* Comparison of noninvasive genetic and camera-trapping techniques for surveying snow leopards. *J. Mammal.* **92**, 771–783 (2011).
23. Janečka, J. E., Jackson, R., Munkhtsog, B. & Murphy, W. J. Characterization of 9 microsatellites and primers in snow leopards and a species-specific PCR assay for identifying noninvasive samples. *Conserv. Genet. Resour.* **6**, 369–373 (2014).
24. Aruge, S., Batool, H., Khan, F. M., Fakhar-i-Abbas & Janjua, S. A pilot study—genetic diversity and population structure of snow leopards of Gilgit-Baltistan, Pakistan, using molecular techniques. *PeerJ* **7**, e7672 (2019).
25. Chetri, M., Odden, M., Sharma, K., Flagstad, Ø. & Wegge, P. Estimating snow leopard density using fecal DNA in a large landscape in north-central Nepal. *Global Ecology and Conservation* **17**, e00548 (2019).
26. Poyarkov, A. D. *et al.* Assurance of the existence of a trans-boundary population of the snow leopard (*Panthera uncia*) at Tsagaanshuvuut - Tsagan-Shibetu SPA at the Mongolia-Russia border. *Integr. Zool.* **15**, 224–231 (2020).
27. Janečka, J. E. *et al.* Range-Wide Snow Leopard Phylogeography Supports Three Subspecies. *J. Hered.* **108**, 597–607 (2017).
28. Senn, H., Murray-Dickson, G., Kitchener, A. C., Riordan, P. & Mallon, D. Response to Janečka et al. 2017. *Heredity* **120**, 581–585 (2018).
29. Janečka, J. E., Janecka, M. J., Helgen, K. M. & Murphy, W. J. The validity of three snow leopard subspecies: response to Senn et al. *Heredity* **120**, 586–590 (2018).
30. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).
31. Puechmaille, S. J. The program structure does not reliably recover the correct population structure when sampling is uneven: subsampling and new estimators alleviate the problem. *Mol. Ecol. Resour.* **16**, 608–627 (2016).
32. Petkova, D., Novembre, J. & Stephens, M. Visualizing spatial population structure with estimated effective migration surfaces. *Nat. Genet.* **48**, 94–100 (2016).
33. Anopheles gambiae 1000 Genomes Consortium *et al.* Genetic diversity of the African malaria vector *Anopheles gambiae*. *Nature* **552**, 96–100 (2017).
34. Gravel, S. *et al.* Demographic history and rare allele sharing among human populations. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 11983–11988 (2011).
35. Van der Auwera, G. A. & O'Connor, B. D. *Genomics in the Cloud: Using Docker, GATK, and WDL in Terra*. ('O'Reilly Media, Inc.', 2020).
36. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).
37. Johansson, Ö. *et al.* The timing of breeding and independence for snow leopard females and their cubs. *Mamm. Biol.* **101**, 173–180 (2021).
38. Prost, S. *et al.* Genomic analyses show extremely perilous conservation status of African and Asiatic cheetahs (*Acinonyx jubatus*). *Mol. Ecol.* **31**, 4208–4223 (2022).
39. Paijmans, J. L. A. *et al.* African and Asian leopards are highly differentiated at the genomic

level. *Curr. Biol.* **31**, 1872–1882.e5 (2021).

40. Armstrong, E. E. *et al.* Unraveling the Genomic Diversity and Evolutionary History of Captive Tigers in the United States. *bioRxiv* 2023.06.19.545608 (2023) doi:10.1101/2023.06.19.545608.

41. Wang, C. *et al.* Population genomic analysis provides evidence of the past success and future potential of South China tiger captive conservation. *BMC Biol.* **21**, 64 (2023).

42. Li, H. & Durbin, R. Inference of human population history from individual whole-genome sequences. *Nature* **475**, 493–496 (2011).

43. Frankham, R. Effective population size/adult population size ratios in wildlife: a review. *Genet. Res.* **89**, 491–503 (2007).

44. Mazet, O., Rodríguez, W., Grusea, S., Boitard, S. & Chikhi, L. On the importance of being structured: instantaneous coalescence rates and human evolution--lessons for ancestral population size inference? *Heredity* **116**, 362–371 (2016).

45. Mather, N., Traves, S. M. & Ho, S. Y. W. A practical introduction to sequentially Markovian coalescent methods for estimating demographic history from genomic data. *Ecol. Evol.* **10**, 579–589 (2020).

46. Kim, S. *et al.* Comparison of carnivore, omnivore, and herbivore mammalian genomes with a new leopard assembly. *Genome Biol.* **17**, 211 (2016).

47. Morin, P. A. *et al.* Reference genome and demographic history of the most endangered marine mammal, the vaquita. *Mol. Ecol. Resour.* **21**, 1008–1020 (2021).

48. Westbury, M. V. *et al.* Extended and Continuous Decline in Effective Population Size Results in Low Genomic Diversity in the World's Rarest Hyena Species, the Brown Hyena. *Mol. Biol. Evol.* **35**, 1225–1237 (2018).

49. Westbury, M. V., Petersen, B., Garde, E., Heide-Jørgensen, M. P. & Lorenzen, E. D. Narwhal Genome Reveals Long-Term Low Genetic Diversity despite Current Large Abundance Size. *iScience* **15**, 592–599 (2019).

50. Xue, Y. *et al.* Mountain gorilla genomes reveal the impact of long-term population decline and inbreeding. *Science* **348**, 242–245 (2015).

51. Suryawanshi, K. *et al.* Estimating snow leopard and prey populations at large spatial scales. *Ecol Sol and Evidence* **2**, (2021).

52. Korablev, M. P. *et al.* Large-scale and fine-grain population structure and genetic diversity of snow leopards (*Panthera uncia* Schreber, 1776) from the northern and western parts of the range with an emphasis on the Russian population. *Conserv. Genet.* **22**, 397–410 (2021).

53. Linnell, J. D. C. *et al.* Border Security Fencing and Wildlife: The End of the Transboundary Paradigm in Eurasia? *PLoS Biol.* **14**, e1002483 (2016).

54. Dinerstein, E. *et al.* The Fate of Wild Tigers. *Bioscience* **57**, 508–514 (2007).

55. Bauer, H. *et al.* Lion (*Panthera leo*) populations are declining rapidly across Africa, except in intensively managed areas. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 14894–14899 (2015).

56. Pal, R., Panwar, A., Goyal, S. P. & Sathyakumar, S. Changes in ecological conditions may influence intraguild competition: inferring interaction patterns of snow leopard with co-predators. *PeerJ* **10**, e14277 (2022).

57. Lovari, S., Kachel, S., Xueyang, L. & Ferretti, F. Chapter 13 - Snow leopard, common leopard, and wolf: Are they good neighbors? in *Snow Leopards (Second Edition)* (eds.

Mallon, D. & McCarthy, T.) 137–147 (Academic Press, 2024).

58. Ostrowski, S. & Gilbert, M. Chapter 9 - Diseases of wild snow leopards and their wild ungulate prey. in *Snow Leopards (Second Edition)* (eds. Mallon, D. & McCarthy, T.) 95–111 (Academic Press, 2024).

59. Heiner, M., Oakleaf, J., Davaa, G. & Kiesecker, J. Chapter 10 - Emerging threats to snow leopards from energy and mineral development. in *Snow Leopards (Second Edition)* (eds. Mallon, D. & McCarthy, T.) 113–121 (Academic Press, 2024).

60. Zahler, P. & Victurine, R. Chapter 11 - Linear infrastructure and snow leopard conservation. in *Snow Leopards (Second Edition)* (eds. Mallon, D. & McCarthy, T.) 123–128 (Academic Press, 2024).

61. Markert, J. A. *et al.* Population genetic diversity and fitness in multiple environments. *BMC Evol. Biol.* **10**, 205 (2010).

62. Garner, B. A., Hoban, S. & Luikart, G. IUCN Red List and the value of integrating genetics. *Conserv. Genet.* **21**, 795–801 (2020).

63. Suryawanshi, K. R., Khanyari, M., Sharma, K., Lkhagvajav, P. & Mishra, C. Sampling bias in snow leopard population estimation studies. *Popul. Ecol.* **61**, 268–276 (2019).

64. Zhou, C. *et al.* Characterization of Olfactory Receptor Repertoires in the Endangered Snow Leopard Based on the Chromosome-Level Genome. *DNA Cell Biol.* **40**, 293–302 (2021).

65. Zhou, C. *et al.* Characterization of microsatellites in the endangered snow leopard based on the chromosome-level genome. *Mamm. Res.* **66**, 385–398 (2021).

66. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv [q-bio.GN]* (2013).

67. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

68. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

69. Pockrandt, C., Alzamel, M., Iliopoulos, C. S. & Reinert, K. GenMap: Fast and Exact Computation of Genome Mappability. *Cold Spring Harbor Laboratory* 611160 (2019) doi:10.1101/611160.

70. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).

71. Zheng, X. *et al.* A high-performance computing toolset for relatedness and principal component analysis of SNP data. *Bioinformatics* **28**, 3326–3328 (2012).

72. R Core Team. *R: A language and environment for statistical computing*. (2020).

73. Tupa, L. North American Regional Snow Leopard Studbook. Preprint at (2011).

74. Blomqvist, L. International Pedigree Book for Snow leopards, Uncia uncia Volume 9. Preprint at (2008).

75. Sinnwell, J. P., Therneau, T. M. & Schaid, D. J. The kinship2 R package for pedigree data. *Hum. Hered.* **78**, 91–93 (2014).

76. Kassambara, A. ggpubr:'ggplot2' based publication ready plots. (2020).

77. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer International Publishing, 2016).

78. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

79. Toyama, K. S., Crochet, P.-A. & Leblois, R. Sampling schemes and drift can bias admixture proportions inferred by structure. *Mol. Ecol. Resour.* **20**, 1769–1785 (2020).

80. Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A. & Mayrose, I. Clumpak: a program for identifying clustering modes and packaging population structure inferences across K. *Mol. Ecol. Resour.* **15**, 1179–1191 (2015).

81. Li, H. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* **27**, 2987–2993 (2011).

82. Weir, B. S. & Cockerham, C. C. Estimating F-Statistics for the Analysis of Population Structure. *Evolution* **38**, 1358–1370 (1984).

83. Picard. http://broadinstitute.github.io/picard/.