1
2
3
4   Airway epithelial cells as a novel intracellular host reservoir for *Cryptococcus* spores
5
6

7   Sébastien C. Ortiz[1], Rachael Fortune-Grant[1], Joshua Davies[1], Robin C. May[2], Rebecca A.
8   Drummond[3] and Margherita Bertuzzi[1]
1. Manchester Fungal Infection Group, Faculty of Biology, Medicine and Health, The University of
Manchester, Manchester Academic Health Science Centre, Core Technology Facility, Grafton Street,
Manchester M13 9NT, UK
2. Institute of Microbiology & Infection and School of Biosciences, University of Birmingham,
Edgbaston, United Kingdom
3. Institute of Immunology & Immunotherapy, Institute of Microbiology & Infection, University of
Birmingham, Birmingham B15 2TT, UK

43   Running Title: *Cryptococcus* spores invade airway epithelial cells
44
45   Key Words: *Cryptococcus neoformans*, *C. deneoformans,* spores, airway epithelial cells,
46   phagocytosis, fungal pathogenesis, latency, dissemination
47

Abstract

Human fungal pathogens cause an estimated 1.5 million annual deaths and *Cryptococcus neoformans,* an inhaled facultative intracellular pathogen, is at the top of the WHO's priority list of fungal pathogens*. Cryptococcus* causes disease when it disseminates out of the lung and into the brain which can occur years after initial exposure. However, the mechanisms of both *Cryptococcus* dissemination and latency are still unclear. One potential reason that these mechanisms remain unknown is that the morphotype used for the vast majority of studies is the yeast form and not the dormant and stress resistant spore morphotype. The inhalation of *Cryptococcus* spores initiates infections and recent evidence *in vivo* has shown that spores display unique host-pathogen interactions and are able to escape the lung and disseminate to the brain more readily than yeast. Due to the difficulties associated working with these spores, little is known about how they interact with host cells. Airway Epithelial Cells (AECs), which cover the entire alveolar surface and comprise 24% of all cells in the human lung parenchyma, likely have instant and extensive contact with inhaled spores. In this study, we demonstrated that *Cryptococcus* spores can invade AECs both *in vitro* and *in vivo*. Once inside spores can germinate and subsequently replicate, persist and/or escape. This ability to enter AECs correlates with a preferential ability of spore to cross AEC barriers. Together our work indicates AECs may serve as a previously ignored intracellular host reservoir and challenges the current paradigms of both *Cryptococcus* dissemination and latency.

Introduction

Invasive fungal diseases cause over 1.5 million deaths annually, with mortality rates often over 50% due to lack of diagnostics and antifungal treatments [1,2]. No vaccines exist for fungal infections, and preventative treatments are rare [3]. By understanding how fungi establish infections, persist in hosts and disseminate, we can develop much needed diagnostics, prevention, and antifungal therapeutics to combat invasive fungal disease. *Cryptococcus neoformans,* an inhaled facultative intracellular pathogen, is at the top of the WHO's priority list of human fungal pathogens, largely due to the high mortality associated with cryptococcosis resulting in an estimated 181,000 annual deaths [4,5]. Key aspects of *Cryptococcus* disease progression are still not understood. Studies have focused on understanding the virulence of this pathogen and its ability to colonise the brain (the primary site of disease), but little is known about how *Cryptococcus* disseminates out of the lung (the initial site of infection). Additionally, via unknown mechanisms, *Cryptococcus* undergoes latency and is able to cause disease in hosts years or decades after initial infection [6,7]. Accordingly, after treatment for cryptococcosis, patients living with HIV/AIDS undergo a minimum of a year of secondary prophylaxis to prevent resurgence of the pathogen [8]. The mechanisms of *Cryptococcus* dissemination, latency and persistence remain unknown.

One of the potential reasons that the pathogenesis of *Cryptococcus* remains poorly understood is that the morphotype (morphologically distinct cell-type) used for the vast majority of studies is the yeast form; however, yeast are too large to reach the lower airway and thus unlikely to establish infection under normal circumstance. *Cryptococcus* produces spores (basidiospores), a dormant and stress resistant morphotype small enough to reach lower airways; consequently, spores are the presumed infectious morphotype in *Cryptococcus* disease [9,10]. Alas, difficulties associated with producing and working with *Cryptococcus* spores has resulted in a problematic knowledge gap regarding spore-mediated infections despite the fact that spores have distinct surface properties, interact with host immune cells differently than yeast, and, critically, disseminate out of the lung when yeast cannot [10-14]. The mechanisms of spore-specific dissemination out of the lung remain unknown, emphasising a need for the investigation of spore pathogenesis [14].

One of the first barriers inhaled pathogens encounter are non-professional phagocytic airway epithelial cells (AECs). Alveolar AECs make up 24% of the lung parenchyma, likely have extensive and prolonged contact with inhaled fungal spores and are key for the antimicrobial defence of the lung, which if mis-regulated may lead to an opportunity for pathogens to persist [15-18]. Studies looking at the interactions of *Cryptococcus* with AECs show that yeast are not readily internalised [19-22]. Given the unique characteristics of spores and the importance of AECs in the host lung defence, we hypothesised that spores interact differently with AECs when compared to yeast and found that *Cryptococcus* spores invade AECs both *in vitro* and *in vivo*. Once inside, spores can germinate, replicate, persist and even escape. Furthermore, spores show preferential crossing of epithelial barriers implicating spore-AEC interactions in dissemination out of the lung. Our results identify AECs as a novel host cell-type that *Cryptococcus* can invade in a spore-specific manner and potentially exploit. AECs also provide an intracellular environment which may facilitate latency/persistence of spores, a fundamentally dormant morphotype. These results challenge current paradigms of *Cryptococcus* dissemination, latency and persistence and emphasise the importance of evaluating different morphotypes in pathogenesis to open the door to novel opportunities for much needed diagnostic and therapeutic intervention.

Results

Cryptococcus spores, unlike yeast, are readily internalised by AECs and can germinate and replicate intracellularly

Previous studies investigating the ability of *Cryptococcus* yeast to enter AECs have shown that even at high infectious doses (Multiplicity of Infection (MOI) of 10) very few yeast are internalised (~0%) [21]. Internalisation increases slightly for acapsular mutants, but these are avirulent and thus unlikely to be a cause of human infection [19-22]. Spores are known to have distinct surface properties that differ from those of yeast cells, suggesting that spores may interact with AECs differently from yeast [10-12, 23].

To test this, we exposed AECs to spores and carried out differential staining followed by imaging flow cytometry (IFC) (**Fig 1**) [24]. A549 immortalised type-II alveolar epithelial cells were exposed to either mCherry tagged *Cryptococcus* spores (produced from prolific sporulator strains JEC20xJEC21), wildtype yeast of the same background, or yeast of the previously evaluated background (KN99) at a low MOI of 0.167. Internalisation was evaluated 6 hours post infection, and counter staining with calcofluor white (CFW) was used to differentiate between AECs associated with *Cryptococcus* extracellularly (AEC$_a$) versus AECs containing intracellular *Cryptococcus* (AEC$_i$) (**Fig 1A**). Consistent with previous studies, neither KN99 nor JEC20/21 yeast cells were taken up by AECs; however, JEC20/21 spores were readily internalised by AECs with 1.91% of AECs having 1 or more internalised spores (1.91% AEC$_i$), and 20% of which had more than one spore per AEC (average stoichiometry of 1.34) (**Fig 1B**) [21,22]. This rate of internalisation is equivalent to ~15.4% of infectious inoculum being internalised by AECs within 6 hours.

In order to cause disease, spores need to germinate into vegetatively growing yeast. *Cryptococcus* spores readily germinate in the lung environment with cryptococci recovered from the bronchial alveolar lavage of mice 18 hours after infection with spores primarily being in the yeast form [14]. In the absence of AECs, spore populations were found to germinate fully by 24 hours in tissue culture conditions (37°C + 5% CO_2) with a previously established bimodal phenotype (**Sup Fig 1**) [25]. To determine if spores germinate within AECs, differential staining and IFC was performed on cells after 24 hours of infection with either spores or yeast (MOI of 1) and intracellular germination was evaluated by quantifying the size (surface area) and circularity (aspect ratio) of internalised *Cryptococcus* using previously establish metrics [26]. As observed previously, with this higher MOI, spores were readily internalised by AECs (12.6% AEC$_i$) as opposed to yeast (0% AEC$_i$) (**Sup Fig 2Ai).** At 24 hours post infections, intracellular *Cryptococcus* was present as different morphotypes, i.e. 30.7% yeast, 46.0% intermediates, and a remaining 23.3% spores (**Fig 1C**), indicating that spores readily germinate but that germination is inhibited in the intracellular environment. This population spread demonstrates a previously characterised asynchronous phenotype, indicating that internalised spores either react non-uniformly within the intracellular environment or that AECs have a range of intracellular environments that spores must germinate in [25]. These possibilities are not mutually exclusive. In addition to germination state, we observed that germinated spores can actively replicate 24 hours post-infection, as evidenced by daughter cell formation. Enlarged intracellular *Cryptococcus* yeast were also observed, consistent with the large size variation that is classically observed in lung environments [27]. The intracellular stoichiometry did not dictate the level of germination given that comparable germination states were observed across AEC$_i$ having internalised different numbers of spores (**Sup Fig 3A**). On the other hand, when multiple spores were internalised by the same AEC$_i$, intracellular spores displayed similar germination kinetics with similar shapes observed 24 hours post infection (**Fig 1D**: Stoichiometry = 2, **Sup Fig 3B:** Stoichiometry ≥ 3), suggesting that

Figure 1. Spore-specific invasion of airway epithelial cellls (AECs) by *Cryptococcus*. A) Representative images from Imaging Flow Cytometry (IFC), 6 hours post infection (hpi). **mCherry** tagged *Cryptococcus* used to identify AECs interacting fungal cells and Calcofluor White was used to distinguish internalised from non-internalised *Cryptococcus*. **B)** Percent of AECs with internalization events (AECi) quantified for yeast (KN99 and JEC20/21), and spores (JEC20xJEC21), demonstrating that **spores** are readily internalised while **yeast** are not. A value of 1.91% AECi reflects 15.4% of spore infectious dose being internalised. Unpaired T-test: p < 0.0001 **C)** IFC was performed 24 hours post infection to evaluate intracellular germination kinetics. Morphotypes were quantified based on cell size (surface area - μm^2) and shape (aspect ratio), with spores being small and oval and yeast being large and round. Along with heterogeneous germination kinetics, both intracellular replication and cell enlargement were observed. **D)** Correlation of spore shape for *Cryptococcus* residing within the same AECs (Stoichiometry = 2) demonstrate a link between host AEC and germination kinetics p < 0.0001 two-way ANOVA.

172	germination is linked to each internalisation event. Together these results show that spores can
173	invade AECs, and once inside, can germinate heterogeneously, and subsequently replicate. This
174	implicates AECs as a novel spore-specific intracellular reservoir that *Cryptococcus* can invade
175	and proliferate within.
176	
177	**Once germinated, *Cryptococcus* can undergo rapid replication, non-lytic escape, or persist**
178	**inside AECs**
179	
To further examine the behaviour of internalised and germinated spores, fluorescence microscopy
was used to monitor intracellular populations hourly from 6 to 26 hours post infection. CFW
counterstaining was used at 6 hours post infection to demark internalisation events. We identified
three behaviours that have previously been identified for intracellular yeast in macrophages,
which are rapid replication (**Fig 2A**), large vacuole formation (**Fig 2B**) and non-lytic escape – also
termed vomocytosis (**Fig 2C**) [28-30]. Between 240 and 358 internalised *Cryptococcus* events
per biological replicate were evaluated for the occurrence of budding, rapid replication balloon
formation, vacuole formation, and non-lytic escape over 26 hours (**Fig 2D**). Intracellular budding
occurred in 10.6% of internalisation events (21.0% of germinated spores), while rapid replication
balloons occurred in 3.0% of events (6.3% of germinated spores) and was limited to when there
was more than one fungal cell present within a single AEC. Vacuole formation occurred in 30.1%
of germinated spores and only 2.0% of ungerminated spores (16.2% of all events). Finally, non-
lytic escape was observed for 12.3% of germinated intracellular spores, and 1.5% of non-
germinated spores (7.0% of all events). These results demonstrate a large heterogeneity of
intracellular behaviours, and that *Cryptococcus* can readily escape AECs once germinated.
195	
To investigate whether internalised *Cryptococcus* can persist within AECs for a prolonged period
of time, A549 cells were infected with spores or yeast for 6 hours, then re-seeded to a low density
to allow continue growth for an additional 5 days (120 hours). After 5 days, cells were evaluated
for the presence of internal *Cryptococcus* cells. Once again, yeast-infected AECs had no
intracellular *Cryptococcus* events; however, despite the daily duplication of AECs, we were still
able to detect intracellular *Cryptococcus* in spore-infected AECs after 126 hours of infection (**Sup**
Fig 2Aii). The germination state of these internalised *Cryptococcus* cells was evaluated and a full
range of germination morphotypes were identified with 59.6% yeast, 24.1% intermediates, and a
remaining 16.3% spores (**Fig 2E**). While this population was more germinated than at 24 hours
(up from 30.7% yeast), there were still spores and intermediates present after 126 hours. It is still
unclear how viable these *Cryptococcus* cells are, but the presence of the unquenched mCherry
reporter may suggest they are either alive or have not been dead long enough for their fluorescent
signal to be quenched [31]. AECs do not show significant signs of apoptosis or necrosis after 6,
24, and 126 hours of infection, likely due to the low virulence of JEC20/JEC21, making lysis and
re-entry much less likely than intracellular persistence (**Sup Fig 2B**) [32]. Together, these results
demonstrate that AECs provide an intracellular environment in which spores can germinate,
replicate, escape and even persist.
213	
214	***Cryptococcus* spores preferentially cross AEC barriers and are internalised *in vivo***
215	
Spores of avirulent yeast preferentially disseminate out of the lung resulting in disease in
intranasal murine models of infection [14]. Given the unique ability of spore to enter AECs, and
subsequently germinate, replicate and escape, we hypothesised that spore-specific invasion of
AECs would enable them to cross epithelial barriers better than yeast. To test this, we monitored
the crossing of *Cryptococcus* across trans-wells seeded with Calu-3 bronchial-like cell line, which
unlike A549 cells, form tight junctions and impermeable barriers. Spores are internalised by
bronchial Calu-3 cells but less efficiently than alveolar A549 cells (**Sup Fig 2Aiii**). Confluent Calu-

Figure 2. Internalised *Cryptococcus* spores display a variety of behaviors. Representative images of behaviours observed using timecourse microscopy of **A) Rapid Replication:** Internalised clusters of spores (6 hpi), efficiently germinating (16 hpi), and proceeded to rapidly replicate resulting in an AEC full of *Cryptoccus* yeast (24 hpi). **B) Vacuole formation** (25 hpi) **C) Non-lytic escape:** Internalised spore having germinated (24 hpi) undergoes non-lytic escape (25 hpi). **D) Quantification of intracellular behaviors:** Population precentages for All, Germinated (G) or Non-Germinated (NG) *Cryptococcus* that show the behaviours described: budding, rapid replication, vacuole formation, and non-lytic escape. **E) Intracellular Persistence:** IFC identified intracellular *Cryptococcus* in spore infected cells 126 hpi displaying high levels of germination, and the presence of ungerminated spores and intermediates.

223 3 cells were infected with either spores or yeast (MOI 5) and media from the lower compartment
224 was plated for colony forming units at 0, 24 and 48 hours (**Fig 3A**). After 48 hours, spores showed
225 preferential crossing as compared to yeast, with 88.9% spore infected wells showing crossing, as
226 opposed to 22.2% of yeast infected wells.
227
Given this ability of spores to invade, persist inside, and escape AECs, we evaluated the *in vivo*
internalisation of *Cryptococcus* by AECs using an intranasal murine model of cryptococcosis.
Mice were intranasally infected with $5x10^6$ spores or yeast (JEC20mCh/JEC21mCh) and the
infection was allowed to progress for 8 hours. This timepoint was chosen to ensure morphotype-
specific characterisation because 8 hours is not long enough for spores to have germinated into
yeast. Lungs were harvested, processed, pooled and differentially stained using our previously
established methodology to identify AECs via IFC (**Fig 3B,C**) [18]. We found that for spore-
infected mice, 0.96% of recovered AECs (EpCAM$^+$) and 0.97% of recovered type-II alveolar
epithelial cells (EpCAM$^+$, CD74$^+$) had internalised spores, with the majority of events having more
than one spore (58.6%) with an average stoichiometry of 2.28 spores/AEC$_i$ (**Sup Fig 3Ciii**). On
the other hand, for yeast-infected mice, only 0.09% of recovered AECs and 0.15% of recovered
type-II alveolar epithelial cells had internalised yeast, with 21.3% having 2 yeast with an average
stoichiometry of 1.14 yeast/AEC$_i$. These data indicate that spores are able to cross epithelial
barriers *in vitro* and invade AECs *in vivo* better than yeast. These results reveal AECs as a novel
intracellular host reservoir for *Cryptococcus* and offers potential mechanisms of spore-specific
dissemination and persistence within the host lung.
244
245
246

Figure 3. *Cryptococcus* **spores preferentially cross AEC barriers and are internalised by AECs in a murine model of infection. A)** Lower well colony forming units at 0, 24, and 48 hours in spore and yeast infected trans-wells indicating the crossing of AEC barrier. **B)** Percentage of i) AECs (EpCAM+) or ii) Type II alveolar epithelial cells (EpCAM+, CD74+) with intracellular *Cryptococcus* (mCherry+, CFW-), derived from mouse lungs 8 hours post intranasal infection with spores or yeast. **C)** Representative Imaging Flow Cytometry panels of *Cryptococcus* cells (both Spores and Yeast) inside type II airway epithelial cell, 8 hours post intranasal infection. **Ch1: Calcofluor White** staining differentiates between fungal cells inside and outside AECs. **Ch2: CD74** staining is used to identify Type II alveolar AECs. **Ch4:** Nuclear localized **mCherry** tag identifies fungal cells. **Ch5:** Brightfield image of cells. **Ch6: EpCAM** staining is used to identify AECs. All statistics derived from unpaired T-test across samples.

Discussion

Cryptococcus spores, along with 'desiccated' yeast, are the presumed infectious particles in cryptococcal disease; however, vegetatively growing yeast are not generally considered to be the primary infectious propagules due to their large size preventing them from reaching the lower airways. Contrary to this, and likely due to the difficulties associated with working with spores and the low viability of 'desiccated' yeast, the vast majority of studies on *Cryptococcus* host-pathogen interactions and disease kinetics have been performed with vegetatively growing yeast and not spores. While understanding yeast-host interactions is critical, ignoring other morphotypes prevents us from understanding the complete picture concerning *Cryptococcus* infections and subsequent disease. By neglecting to study these morphotypes, we miss out on opportunities to develop diagnostics, prevention and treatments for fungal disease. In this study, we demonstrate that spores, which have distinct surface properties, interactions with host cells, and disease kinetics to yeast, are internalised by AECs when yeast are not [10-14]. Once inside, these spores can germinate, replicate, escape and/or persist within AECs (**Fig 4**). These spore-specific host-pathogen interactions may have major implications for our understanding of both dissemination and latency.

Work by Walsh et al., demonstrated that spores of avirulent yeast (B3501/02) caused disease in an intranasal murine model 50 days after initial infection, while yeast did not [14]. Spore-specific virulence was directly linked to the ability of spores to disseminate out of the lungs when yeast could not, with dissemination to the kidneys within 6 days post infection, and eventually dissemination to the brain. Since spores readily germinate within 18 hours in the mouse lung, this work suggested that early host-pathogen interactions in the lung dictate spore-specific dissemination and disease. The authors postulated that spore-specific dissemination could result from a Trojan-Horse mechanism where spore hide inside macrophages to get shuttled out of the lung, a prominent hypothesis for yeast dissemination as well. The work demonstrated that spores were more efficient at reaching lung draining lymph nodes than yeast, and that trafficking to the lymph nodes was CD11c[+] dependent. The data we present here provides an alternative, but not mutually exclusive, hypothesis for the mechanism of dissemination out of the lung. Rather than a macrophage driven Trojan-Horse model, spores may be able to disseminate out of the lung through transcytosis, where spores are internalised by epithelial cells, germinate, and non-lytically escape on the other side of the epithelial barrier. This alternative mechanism is further supported by recent work that identified the infrequent occurrence of 'free floating' yeast in the vasculature (large blood vessels) of mouse lungs 7 days post intranasal yeast infection, consistent with the low levels of internalisation by AEC we observed in our murine yeast infections [33]. These two mechanisms may even work in tandem, as reported with *Mycobacterium tuberculosis (Mtb)*, with monocytes containing *Mtb* more efficiently translocating across an epithelial barrier when AECs were also infected with *Mtb* [34]. Subsequent studies investigating the precise role of AECs in *Cryptococcus* dissemination out of the lung will require the identification of both molecular mechanisms of internalisation and escape in order to modulate spore-epithelial interactions.

The first epidemiological evidence of *Cryptococcus* latency was obtained in 1999 [34]. Since this first study, a variety of evidence has supported the idea of latency/persistence in cryptococcal disease [6,7]. Relatively little is known about the mechanisms underlying latency/persistence; however, granulomas have largely been presumed to be the predominant source of latent *Cryptococcus* [36]. This idea originally stems from a comparison to *Mtb*, where similar granulomas have been the predominant paradigm for a mechanism of latency [6, 36, 37]. However, this has recently been challenged with evidence of *Mtb* DNA being present in non-professional phagocytic lung cells in the absence of tuberculous legions [37,38]. The frequency and importance of granulomas in Cryptococcal disease is unclear, and while granulomas are likely more feasible as

Figure 4. Schematics of our current understanding of *Cryptococcus*-AEC interactions. In this work we've demonstrated that spores, unlike yeast, are readily internalised by airway epithelial cells. This spore-specific interaction provides a unique intracellular niche where spores can germinate, and once germinated, can replicate, persist and non-lytically escape. The morphotype specific ability to enter AECs, survive, and escape provides a viable explanation for *Cryptococcus* spore-specific dissemination out of the lung. The ability of *Cryptococcus* to persist inside AECs, not only as fully germinated yeast, but as apparently dormant (ungerminated) spores provides an unexplored potential mechanism of latency.

a latency reservoir than free-floating *Cryptococcus* in the lung, our data provides an alternative mechanism where non-professional phagocytic AECs could provide an ideal intracellular environment, shielded from immune cells, in which *Cryptococcus* could lie dormant for extended periods of time.

Dormancy in *Cryptococcus* is poorly understood and the morphological state of *Cryptococcus* during latency is unknown. *In vivo* models have demonstrated the presence of low-metabolic activity yeast that match a dormant population [39]. Following this, *in vitro* conditions were developed to mimic this dormant population resulting in viable but non culturable cells (VBNC) [40]. Further research is required to properly understand the importance and role of this yeast subpopulation in *Cryptococcus* disease kinetics [7,39,40]. Importantly, spores are the classical example of a dormant particle, and for *Cryptococcus* these are exclusively produced in the environment, not inside the host. Spores shows distinct nutrient utilisation to yeast but the molecular mechanisms that control the escape from dormancy still require characterisation [9,12]. It is possible that spores, being a fundamentally dormant morphotype, are more readily able to remain dormant (or re-establish dormancy) in a host than yeast. The identification of AECs as a spore-specific intracellular environment, where spores can persist and germinate to varying degrees, provides a novel alternative for understanding latency and persistence in cryptococcal disease.

A growing body of evidence shows an important role for AECs in the antimicrobial potency of the lung, where the interactions of these non-professional phagocytes with pathogens can be either a benefit or detriment to the host [16,17]. This likely depends on both pathogen factors and host predispositions. The role of AECs in cryptococcal disease has largely been ignored which may be due to the lack of internalisation of the yeast morphotype. It will be interesting to understand if other understudied morphotypes present in the lung, such as VBNC yeast, seed cells, and Titanides, could interact similarly with AEC as spores do. This study expands on the already long list of differences between *Cryptococcus* spores and yeast and points towards AECs as a novel intracellular environment for the field to consider when studying *Cryptococcus* disease kinetics and potential therapeutics.

Methods

Fungal Strains and Strain Manipulation

Cryptococcus neoformans serotype D (*deneoformans*) strains JEC20-mCh (CHY4031), JEC21-mCh (CHY4028), and *Cryptococcus neoformans* serotype A strain KN99-mCh were handled using standard techniques and media as described previously [14,41,42]. *Cryptococcus* spores were isolated from cultures as described previously [10]. Briefly, yeast of both mating types (JEC20-mCh and JEC21-mCh) were grown at 30°C for 2 days on yeast-peptone-dextrose (YPD) agar, resuspended in 1X phosphate buffered saline (PBS) [Sigma-Aldrich: D8537], mixed in a ratio of 1:1, and intermittently spotted onto dried V8 pH 7.0 agar plates. Plates were incubated for 7 days at 25°C in the dark and crosses were resuspended in 75% Percoll® [Sigma-Aldrich: GE17-0891-01] in 1X PBS for separation via gradient centrifugation (2110xg, 25 min). Spores were recovered, counted using a hemocytometer and assessed for purity by visual inspection.

Epithelial Cell Culture

The cell lines used in this study were A549, human pulmonary carcinoma epithelial cells [ATCC: CCL185] and Calu-3, human pulmonary adenocarcinoma epithelial cells [ATCC: HTB-55]. A549 cells were maintained in DMEM [Sigma-Aldrich: D5796] supplemented with 10% Gibco^{TM} FBS [ThermoFisher Scientific: A5256801] and 1% penicillin/streptomycin cocktail [Sigma-Aldrich: P0781] (DMEMs). Calu-3 cells were maintained in DMEM/F12 [ThermoFisher Scientific: 11320033] supplemented with 10%FBS, 1% penicillin/streptomycin cocktail (DMEM/F12s). All cells were grown and maintained at 37°C, 5% CO_2, with all passaging and seeding performed with Trypsin-EDTA [Sigma-Aldrich: T3924] mediated detachment.

Differential Fluorescence Imaging Flow Cytometry

Differential fluorescence IFC was performed following published methodology [24]. Briefly, confluent monolayers of A549 cells across triplicate wells of 6-well plates [Greiner: 657160] were infected with either spores (JEC20-mChxJEC21-mCh) or yeast (JEC20-mCh+JEC21-mCh, KN99α-mCh) at varying MOIs (0.167, 1 or 5) for different lengths of times (6 hours, 24 hours). All experiments had uninfected controls processed identically as negative controls and for gating purposes. For intracellular persistence evaluation, after 6 hours of incubation A549 cells were reseeded to attain confluency 120 hours later (effectively 126 hpi). At the end of the infections, A549 cells were trypsin digested and collected via centrifugation at 500x g. Cells were incubated in 500 µL of Binding Buffer, 1X (5 µL) Annexin V-FITC [Abcam: ab14085], 100 nM TO-PRO^{TM}-3 [ThermoFisher Scientific: T3605] and 8 µg/mL Calcofluor White (CFW) [Sigma-Aldrich: F3543] for 5 minutes, centrifuged, and washed with 1X PBS. Cells were resuspended and evaluated on a Amnis ImageStream MkII. For each sample, 5000 single cells in focus were analysed across three independent acquisitions in biological triplicate. Analysis was performed for the identification of internalisation and host cell death using IDEAS® software according to previously defined analysis pipeline [24]. Uninfected A549 samples were used for background subtraction, and all samples were normalised based on infectious inoculum as determined by triplicate CFU counts. Visualisation of internalisation events was performed on both IDEAS® software and Fiji/ImageJ to quantify stoichiometry and germination. *Cryptococcus* germination quantification was performed by evaluation of area and aspect ratio of internalised *Cryptococcus* cells following previous established cut offs (Spores: Area <7.44 µm^2, Aspect Ratio <0.8 | Yeast: Area >7.84 µm^2, Aspect Ratio >0.8) as defined by Barkal et al. 2016 [26] across all biological replicates with technical replicates acquired for internalisation evaluation.

Microscopy

Intracellular Cryptococcus Behaviours

A549 cells were seeded into 24-well glass bottom plates [Greiner: 662892] for two days until reaching confluency, infected with JEC20-mChxJEC21-mCh spores at an MOI of 1. After 6 hours, media was removed and cells were stained with 8 µg/mL CFW for 5 minutes, washed with 500 µL 1X PBS and resuspended in Gibco™ FluoroBrite™ DMEM media [ThermoFisher Scientific: A1896701] supplemented with 10% FBS and 1% penicillin/streptomycin cocktail. All images were acquired on a Nikon Eclipse Ti Inverted Microscope, using a Nikon CFI Plan APO λ 40X objective and Hamamatsu ORCA-Flash4.0 LT C11440 camera, at 37°C and 5% CO_2 for 20 hours after staining. Images were acquired across 3 biological replicates, 3 wells per replicate (10 positions/well) in bright field, 400 nm (100 ms exposure) and 550 nm (300 ms exposure). For each biological replicate, 240-348 internalisation events (determined by CFW- signal at 6 hours post infection and localisation) were monitored hourly from 6 to 26 hours post infection. For each internalised *Cryptococcus* event, germination (determined by size/shape), replication (determined by the addition of intracellular *Cryptococcus* cells i.e. budding), rapid replication (determined by number of intracellular *Cryptococcus* cells at least tripling by 26 hours and visual ballooning), vacuole formation (determined by large ring surrounding intracellular *Cryptococcus* cells), and non-lytic escape (determined by intracellular *Cryptococcus* cells floating away from host cells) were evaluated and quantified.

Quantitative Germination Assay

Germination of spores was evaluated across 24 hours in tissue culture conditions (supplemented DMEM, 37C, 5%CO_2) using the previously published quantitative germination assay and custom algorithms [25,26] on images acquired on a Nikon Eclipse Ti Inverted Microscope, using a Nikon CFI Plan Fluor ELWD 20X objective and Hamamatsu ORCA-Flash4.0 LT C11440 camera.

Trans-well Crossing

Trans-well crossing experiments were performed by seeding Falcon® permeable support for 12-well plate with 8.0 µm transparent PET membrane [Corning: 353182] and associated companion plate [Corning: 353503] with Calu-3 cells at $1x10^6$ cells/well. Cells were allowed to reach confluency (8-12 days) as determined by a resistance measurement of >900 ohm using Milipore Millicell® ERS-2 Epithelial Volt-Ohm Meter and Milicell® ERS-22 Adjustable electrode set [MERSSTEX03]. Each upper well of the trans-well was infected with either spores or yeast (MOI of 5) and crossing was evaluated by plating lower well volumes onto YPD plates at 0, 24 and 48 hours post infection. YPD plates were incubated for 3 days, and total number of colonies/well/day were quantified. Impermeability of barrier was further confirmed by lack of fungal crossing following infection and resistance was monitored across all time points. Crossing was evaluated for 9 replicates of both spore and yeast infected wells.

Murine infection, lung processing, and staining for imaging flow cytometry

Murine infections were performed under UK Home Office Project License PBE275C33 at the Biomedical Service Unit, University of Birmingham. Mice were housed in individually ventilated cages under specific-pathogen free conditions, with access to standard chow and drinking water *ad libitum* under 12 hour light/dark cycle at 20-24ºC and 45-65% humidity. Spores (JEC20-mChxJEC21-mCh) and yeast (JEC20-mCh+JEC21-mCh 50:50 mix) were enumerated using hemocytometers and diluted in 1X PBS. Three groups of 8-10 week old female C57BL/6NCrl mice (Charles River), were left uninfected (6 mice), or intranasally infected with spores (4 mice), or

yeast (4 mice) at a final dose of 5×10^6 cell/mouse. Mice were humanely euthanised by cervical
dislocation at 8 hours post infection and lungs were perfused with 2 mL of 0.1 mg/ml Dispase
[Sigma-Aldrich: D4693] in HANKS buffer [Sigma-Aldrich: H9269] directly into trachea. Lungs were
processed as previously described [18]. Tissue digestion of whole lungs was carried out at 37°C
for 1 hour with gentle rotation (100 rpm). Digestion buffer (0.1 mg/ml Dispase, 0.4 U/ml Liberase
TL [Sigma-Aldrich: 5401020001] and 160 U/ml DNase [Sigma-Aldrich: D5025] was added and
lungs were incubated an additional 10 minutes. Samples were diluted with 1X PBS + 2% FBS +
2 mM EDTA, systematically filtered through 100 µm, 70 µm and 40 µm filters [Sigma-Aldrich:
431752, 431751, 431750] and centrifuged at 500x g for 6 minutes at 4°C. Supernatant was
removed, and pellet was resuspended in 10 mL of 1x RBC lysis buffer [BioLegend: 422401],
incubated at 25°C for 5 minutes. Samples were pooled together (2 mice per pool), diluted with 1X
PBS + 2% FBS + 2 mM EDTA, centrifuged for 6 minutes at 500x g and 4°C, supernatants were
discarded, and the pellets were resuspended in remaining volume (~200 µL). Samples containing
excess remaining red blood cells post lysis were not evaluated via IFC. Each sample was
incubated with 5 µg/mL anti-mouse CD16/32 [BioLegend, 101302] for 10 minutes at 4°C,
centrifuged at 500x g for 6 minutes at 4°C and pellets were resuspended in 100 µL of 1X PBS +
2% FBS + 2 mM EDTA. Samples were incubated with 1 µg/mL Ep-CAM (CD326) PE/Cy7
Antibody [BioLegend, 118215] and 1 µg/mL CD74 FITC Antibody [Santa Cruz Biotechnology, sc-
6262-FITC] for 25 minutes at 4°C, followed by 8 µg/mL CFW for 5 minutes. Samples were
centrifuged at 500x g, for 3 minutes and resuspended in 100 µL of 1X PBS + 2% FBS + 2 mM
EDTA prior to evaluation on an Amnis ImageStream MkII. For each pooled sample (three
uninfected, two yeast infected, and two spore infected), 3 acquisition of 3000-4000 EpCam$^+$ single
cells in focus were taken as technical replicates. Data analysis was performed using the IDEAS®
software and the pipeline previously described, using single stain controls for gating purposes.

Statistical analysis of data

GraphPad Prism was used to interpret data and p values were calculated through unpaired t tests
or, ordinary 2-way ANOVA as specified in the figure legends. Error bars show standard deviation.

Funding

This work was supported by grants to SCO and MB from the British Mycological Society (BMS
Research Grant 2022) and to MB from the Medical Research Council (MRC New Investigator
Research Grant MR/V031287/1) and to RAD from the Medical Research Council (MRC Career
Development Award MR/S024611).

Acknowledgments

We would like to thank Dr Christina Hull for the gift of the JEC20-mCh and JEC21-mCh strains,
Dr Gareth Howell and his team at the Flow Cytometry Core Facility, University of Manchester,
and Ferdus Sheik at the Flow Cytometry Facility, University of Birmingham for technical support.

Supplemental Figure 1. Germination of spores in tissue culture conditions. Two-dimentional histogram of spores (JEC20mChxJEC21mCh) germinated in supplemented DMEM, at 37°C, 5%CO_2 for 24 hours. Spores are small and oval (Area <7.44 µm², Aspect Ratio <0.8) and present in the lower left hand quandrant, while yeast are large and round (Area >7.84 µm², Aspect Ratio >0.8) and are present in the upper right han quandrant. Population displays a 'Bimodal' phenotype, with the population fully germinated by 24 hours.

Supplemental Figure 2. Relative *Cryptococcus* internalisation and host cell damage. A) Percent of AECs with intracellular *Cryptococcus* (%AECi) for spore and yeast infected i) A549s for 24 hours - MOI of 1 , ii) A549s for 126 hours - MOI of 5, and iii) Calu-3s for 6 hours - MOI of 1. **B)** Percentage of live AECs as determined by lack of necrosis (TO-PRO-3) or apoptosis (Annexin-FITC) for A549s infected for i) 6 hours with an MOI of 0.167, ii) 24 hours with an MOI of 1, or 126 hours with an MOI of 5.

Supplemental Figure 3. Intracellular *Cryptococcus* stoichiometry. A) Germination profiles of intracellular *Cryptococcus* 24 hours post infection seperated by number of fungal cells per AEC (stoichiometry) demonstrating that germination kinetics remain constatn across all intracellular stoichiometries. **B)** Correlation of spore shape for *Cryptococcus* residing within the same AECs (Stoichiometry = 3+) demonstrate a link between host AEC and germination kinetics p < 0.0001 two-way ANOVA. **C)** Stoichiometry, as percent of populaiton, for i) 6 hour post infection of A549 with spores (MOI 0.167), ii) 24 hours post infection with sporesA549 (MOI 1), and iii) AECs isolated from mice intranasally infected with either spores or yeast.

Citations

[1] Bongomin, F., Gago, S., Oladele, R. O., & Denning, D. W. (2017). Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. *Journal of Fungi (Basel, Switzerland)*, *3*(4), 57. https://doi.org/10.3390/jof3040057

[2] Brown, G. D., Denning, D. W., Gow, N. A., Levitz, S. M., Netea, M. G., & White, T. C. (2012). Hidden killers: human fungal infections. *Science Translational Medicine*, *4*(165), 165rv13. https://doi.org/10.1126/scitranslmed.3004404

[3] Oliveira, L. V. N., Wang, R., Specht, C. A., & Levitz, S. M. (2021). Vaccines for human fungal diseases: close but still a long way to go. *NPJ Vaccines*, *6*(1), 33. https://doi.org/10.1038/s41541-021-00294-8

[4] World Health Organization. WHO fungal priority pathogens list to guide research, development and public health action. *WHO* https://www.who.int/publications/i/item/9789240060241 (2022).

[5] Rajasingham, R., Smith, R. M., Park, B. J., Jarvis, J. N., Govender, N. P., Chiller, T. M., Denning, D. W., Loyse, A., & Boulware, D. R. (2017). Global burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. *The Lancet. Infectious Diseases*, *17*(8), 873–881. https://doi.org/10.1016/S1473-3099(17)30243-8

[6] Dromer, F., Casadevall, A., Perfect, J. and Sorrell, T. (2010). *Cryptococcus neoformans*: Latency and Disease. In Cryptococcus (eds J. Heitman, T.R. Kozel, K.J. Kwon-Chung, J.R. Perfect and A. Casadevall). https://doi.org/10.1128/9781555816858.ch31

[7] Alanio A. (2020). Dormancy in *Cryptococcus neoformans*: 60 years of accumulating evidence. *The Journal of Clinical Investigation*, *130*(7), 3353–3360. https://doi.org/10.1172/JCI136223

[8] World Health Organization. Guidelines for diagnosing, preventing and managing cryptococcal disease among adults, adolescents and children living with HIV. *WHO* https://www.who.int/publications/i/item/9789240052178 (2022).

[9] Ortiz, S. C. and Hull, C.M. 2023. Biogenesis, germination, and pathogenesis of *Cryptococcus* spores. *Microbiology and Molecular Biology Reviews*. Under Review 2023

[10] Botts, M. R., Giles, S. S., Gates, M. A., Kozel, T. R., & Hull, C. M. (2009). Isolation and characterization of *Cryptococcus neoformans* spores reveal a critical role for capsule biosynthesis genes in spore biogenesis. *Eukaryotic Cell*, *8*(4), 595–605.

[11] Giles, S. S., Dagenais, T. R., Botts, M. R., Keller, N. P., & Hull, C. M. (2009). Elucidating the pathogenesis of spores from the human fungal pathogen *Cryptococcus neoformans*. *Infection and Immunity*, *77*(8), 3491–3500.

[12] Ortiz, S.C., McKeon, M.C., Botts, M. R., Gage, H., Frerichs, A. B & Hull, C. M. (2003) Spores of the fungal pathogen *Cryptococcus* exhibit cell type-specific carbon source utilization during germination. *Preprint at bioRxiv*, 10.1101/2023.10.01.560341

[13] Davis, J. M., Huang, M., Botts, M. R., Hull, C. M., & Huttenlocher, A. (2016). A zebrafish model of cryptococcal infection reveals roles for macrophages, endothelial cells, and neutrophils

in the establishment and control of sustained fungemia. *Infection and Immunity*, *84*(10), 3047–3062.

[14] Walsh, N. M., Botts, M. R., McDermott, A. J., **Ortiz, S. C.,** Wüthrich, M., Klein, B., & Hull, C. M. (2019). Infectious particle identity determines dissemination and disease outcome for the inhaled human fungal pathogen *Cryptococcus*. *PLoS Pathogens*, *15*(6), e1007777.

[15] Crapo, J. D., Barry, B. E., Gehr, P., Bachofen, M., & Weibel, E. R. (1982). Cell number and cell characteristics of the normal human lung. *The American Review of Respiratory Disease*, *125*(6), 740–745. https://doi.org/10.1164/arrd.1982.125.6.740

[16] Bertuzzi, M., Hayes, G. E., Icheoku, U. J., van Rhijn, N., Denning, D. W., Osherov, N., & Bignell, E. M. (2018). Anti-*Aspergillus* activities of the respiratory epithelium in health and disease. *Journal of Fungi*, *4*(1), 8. https://doi.org/10.3390/jof4010008

[17] Bertuzzi, M., Hayes, G. E., & Bignell, E. M. (2019). Microbial uptake by the respiratory epithelium: outcomes for host and pathogen. *FEMS Microbiology Reviews*, *43*(2), 145–161. https://doi.org/10.1093/femsre/fuy045

[18] Bertuzzi, M., Howell, G.J., Thomson, D.D., Fortune-Grant, R., Möslinger, A., Dancer, P., Van Rhijn, N., Du, X., Codling, A., Sash, R., Demirbag, M., Bignell, E.M. (2022) Epithelial uptake of *Aspergillus fumigatus* drives efficient fungal clearance *in vivo* and is aberrant in Chronic Obstructive Pulmonary Disease (COPD). *Preprint at bioRxiv*, 10.1101/2022.02.01.478664

[19] Merkel, G. J., & Scofield, B. A. (1997). The in vitro interaction of *Cryptococcus neoformans* with human lung epithelial cells. *FEMS Immunology and Medical Microbiology*, *19*(3), 203–213. https://doi.org/10.1111/j.1574-695X.1997.tb01089.x

[20] Barbosa, F. M., Fonseca, F. L., Holandino, C., Alviano, C. S., Nimrichter, L., & Rodrigues, M. L. (2006). Glucuronoxylomannan-mediated interaction of *Cryptococcus neoformans* with human alveolar cells results in fungal internalization and host cell damage. *Microbes and Infection*, *8*(2), 493–502. https://doi.org/10.1016/j.micinf.2005.07.027

[21] Choo, K. K., Chong, P. P., Ho, A. S., & Yong, P. V. (2015). The role of host microfilaments and microtubules during opsonin-independent interactions of *Cryptococcus neoformans* with mammalian lung cells. *European Journal of Clinical Microbiology & Infectious Diseases*, *34*(12), 2421–2427. https://doi.org/10.1007/s10096-015-2497-4

[22] Taylor-Smith L. M. (2017). *Cryptococcus*-epithelial interactions. *Journal of Fungi (Basel, Switzerland)*, *3*(4), 53. https://doi.org/10.3390/jof3040053

[23] Walsh, N. M., Wuthrich, M., Wang, H., Klein, B., & Hull, C. M. (2017). Characterization of C-type lectins reveals an unexpectedly limited interaction between *Cryptococcus neoformans* spores and Dectin-1. *PloS One*, *12*(3), e0173866. https://doi.org/10.1371/journal.pone.0173866

[24] Bertuzzi, M., & Howell, G. J. (2021). Single-cell analysis of fungal uptake in cultured airway epithelial cells using differential fluorescent staining and imaging flow cytometry. *Methods in Molecular Biology (Clifton, N.J.)*, *2260*, 83–109. https://doi.org/10.1007/978-1-0716-1182-1_6

586 [25] Ortiz, S. C., Huang, M., & Hull, C. M. (2021). Discovery of fungus-specific targets and
587 inhibitors using chemical phenotyping of pathogenic spore germination. *mBio*, *12*(4), e0167221.
588 https://doi.org/10.1128/mBio.01672-21
589
590 [26] Barkal, L. J., Walsh, N. M., Botts, M. R., Beebe, D. J., & Hull, C. M. (2016). Leveraging a high
591 resolution microfluidic assay reveals insights into pathogenic fungal spore
592 germination. *Integrative Biology*, *8*(5), 603–615. https://doi.org/10.1039/c6ib00012f
593
594 [27] Okagaki, L. H., Strain, A. K., Nielsen, J. N., Charlier, C., Baltes, N. J., Chrétien, F., Heitman,
595 J., Dromer, F., & Nielsen, K. (2010). *Cryptococca*l cell morphology affects host cell interactions
596 and pathogenicity. *PLoS Pathogens*, *6*(6), e1000953.
597 https://doi.org/10.1371/journal.ppat.1000953
598
599 [28] Ma, H., Croudace, J. E., Lammas, D. A., & May, R. C. (2006). Expulsion of live pathogenic
600 yeast by macrophages. *Current Biology : CB*, *16*(21), 2156–2160.
601 https://doi.org/10.1016/j.cub.2006.09.032
602
603 [29] Alvarez, M., & Casadevall, A. (2006). Phagosome extrusion and host-cell survival after
604 *Cryptococcus neoformans* phagocytosis by macrophages. *Current Biology*, *16*(21), 2161–2165.
605 https://doi.org/10.1016/j.cub.2006.09.061
606
607 [30] Tucker, S. C., & Casadevall, A. (2002). Replication of *Cryptococcus neoformans* in
608 macrophages is accompanied by phagosomal permeabilization and accumulation of vesicles
609 containing polysaccharide in the cytoplasm. *Proceedings of the National Academy of Sciences of
610 the United States of America*, *99*(5), 3165–3170. https://doi.org/10.1073/pnas.052702799
611
612 [31] Shaner, N. C., Campbell, R. E., Steinbach, P. A., Giepmans, B. N., Palmer, A. E., & Tsien,
613 R. Y. (2004). Improved monomeric red, orange and yellow fluorescent proteins derived from
614 *Discosoma* sp. red fluorescent protein. *Nature Biotechnology*, *22*(12), 1567–1572.
615 https://doi.org/10.1038/nbt1037
616
617 [32] Lin, X., Nielsen, K., Patel, S., & Heitman, J. (2008). Impact of mating type, serotype, and
618 ploidy on the virulence of *Cryptococcus neoformans*. *Infection and Immunity*, *76*(7), 2923–2938.
619 https://doi.org/10.1128/IAI.00168-08
620
621 [33] Francis, V., Liddle, C., Camacho, E., Kulkarni, M., Junior S.R.S., Harvey, J.A., Ballou, E.R.,
622 Thomson, D.D., Hardwick J.M., Casadevall, A., Witton, J., Coelho, C. (2023). *Cryptococcus
623 neoformans* rapidly invades the murine brain by sequential breaching of airway and endothelial
624 tissues barriers, followed by engulfment by microglia. *Preprint at bioRxiv*,
625 10.1101/2023.11.13.564824
626
627 [34] Bermudez, L. E., Sangari, F. J., Kolonoski, P., Petrofsky, M., & Goodman, J. (2002). The
628 efficiency of the translocation of *Mycobacterium tuberculosis* across a bilayer of epithelial and
629 endothelial cells as a model of the alveolar wall is a consequence of transport within mononuclear
630 phagocytes and invasion of alveolar epithelial cells. *Infection and Immunity*, *70*(1), 140–146.
631 https://doi.org/10.1128/IAI.70.1.140-146.2002
632
633 [35] Garcia-Hermoso, D., Janbon, G., & Dromer, F. (1999). Epidemiological evidence for dormant
634 *Cryptococcus neoformans* infection. *Journal of Clinical Microbiology*, *37*(10), 3204–3209.
635 https://doi.org/10.1128/JCM.37.10.3204-3209.1999
636

637   [36] Ristow, L. C., & Davis, J. M. (2021). The granuloma in cryptococcal disease. *PLoS*
638   *Pathogens*, *17*(3), e1009342. https://doi.org/10.1371/journal.ppat.1009342
639
640   [37] Mayito, J., Andia, I., Belay, M., Jolliffe, D. A., Kateete, D. P., Reece, S. T., & Martineau, A. R.
641   (2019). Anatomic and cellular niches for *Mycobacterium tuberculosis* in latent tuberculosis
642   infection. *The Journal of Infectious Siseases*, *219*(5), 685–694.
643   https://doi.org/10.1093/infdis/jiy579
644
645   [38] Hernández-Pando, R., Jeyanathan, M., Mengistu, G., Aguilar, D., Orozco, H., Harboe, M.,
646   Rook, G. A., & Bjune, G. (2000). Persistence of DNA from *Mycobacterium tuberculosis* in
647   superficially normal lung tissue during latent infection. *Lancet (London, England)*, *356*(9248),
648   2133–2138. https://doi.org/10.1016/s0140-6736(00)03493-0
649
650   [39] Alanio, A., Vernel-Pauillac, F., Sturny-Leclère, A., & Dromer, F. (2015). *Cryptococcus*
651   *neoformans* host adaptation: toward biological evidence of dormancy. *mBio*, *6*(2), e02580-14.
652   https://doi.org/10.1128/mBio.02580-14
653
654   [40] Hommel, B., Sturny-Leclère, A., Volant, S., Veluppillai, N., Duchateau, M., Yu, C. H., Hourdel,
655   V., Varet, H., Matondo, M., Perfect, J. R., Casadevall, A., Dromer, F., & Alanio, A. (2019).
656   *Cryptococcus neoformans* resists to drastic conditions by switching to viable but non-culturable
657   cell phenotype. *PLoS pathogens*, *15*(7), e1007945. https://doi.org/10.1371/journal.ppat.1007945
658
659   [41] Upadhya, R., Lam, W. C., Maybruck, B. T., Donlin, M. J., Chang, A. L., Kayode, S., Ormerod,
660   K. L., Fraser, J. A., Doering, T. L., & Lodge, J. K. (2017). A fluorogenic *C. neoformans* reporter
661   strain with a robust expression of m-cherry expressed from a safe haven site in the
662   genome. *Fungal Genetics and Biology*, *108*, 13–25. https://doi.org/10.1016/j.fgb.2017.08.008
663
664   [42] Sherman F, Fink GR, Hicks JB. (1987). Laboratory course manual for methods in yeast
665   genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.