The functional type VI secretion system of *Sinorhizobium fredii* USDA257 is
required for a successful nodulation with *Glycine max* cv Pekin

Author list: Pedro José Reyes-Pérez[1], Ana Sánchez-Reina[1], Cristina Civantos[1], Natalia
Moreno-de Castro[1], Francisco Javier Ollero[1], Jacinto Gandullo[2], Irene Jiménez-
Guerrero[1*], Patricia Bernal[1]* and Francisco Pérez-Montaño[1]*.

Affiliations: 1. Departamento de Microbiología, Facultad de Biología, Universidad de
Sevilla, Avda. Reina Mercedes 6, Sevilla, Spain
2. Departamento de Biología Vegetal y Ecología, Facultad de Biología, Universidad de
Sevilla, Avda. Reina Mercedes 6, Sevilla, Spain
*Address correspondence to Irene Jiménez-Guerrero (ORCID ID: 0000-0001-8661-
5361), Patricia Bernal (ORCID ID: 0000-0002-6228-0496) and Francisco Pérez-
Montaño (ORCID ID: 0000-0002-6272-2063), ijimgue@us.es, pbernal@us.es and
fperezm@us.es

Abstract: For agriculture, the symbiosis carried out by rhizobia with legumes stands
out as crucial for both economic and environmental reasons. In this process, the bacteria
colonize the roots of the plants, inducing the formation of plant organs called nodules.
Within these structures, rhizobia fix the environmental nitrogen into ammonia reducing
the demand for this essential element required for plant growth. Various bacterial
secretion systems (TXSS, Type X Secretion System) are involved in the establishment
of this symbiosis, with the T3SS being the most extensively studied. The T6SS is a
nanoweapon present in 25% of gram-negative bacteria, commonly used against other
gram-negative bacteria, though some of them use it to manipulate eukaryotic cells.

Interestingly, although T6SS is widely distributed among rhizobia, whether it has a specific role in symbiosis with legumes remains elusive. *Sinorhizobium fredii* USDA257 is a fast-growing rhizobium with the capacity to nodulate a great variety of legume plants. This strain harbors a single T6SS cluster, containing the genes encoding all the structural components of the system and two genes encoding potential effectors that could target the cell wall of the plants and/or be acting as a toxin/antitoxin system. We have demonstrated that this system is active and can be induced in poor culture media. In addition, we have seen by fluorescence microscopy that the T6SS is active in nodules. Competition assays between USDA257 and different preys have shown that USDA257 cannot kill any of them using its T6SS under tested conditions. By constract, nodulation assays demonstrated that USDA257 utilizes this protein secretion system to enhance nodulation and competitiveness with its host *Glycine max* cv Pekin.

Keywords: rhizobium, legume, symbiosis, nodulation, competition, type VI secretion system, effector.

Funding: This work was funded with Projects PID2020-118279RA-I00 and PID2019-107634RB-I00 funded by MICIU/AEI/10.13039/501100011033.

Conflicts of Interest: The authors declare no conflict of interest.

1. Introduction

Rhizobia are α- and ß-Proteobacteria soil-borne microorganisms frequently found on

leguminous plant roots and rhizosphere. This environment is highly appropriate for

rhizobia development and, eventually, they form a symbiotic relationship with the

leguminous plant. The rhizosphere provides rhizobia with protection from desiccation,

extreme temperatures and light stress. At the same time, the legume supplies the

bacteria with nutrients exuded from the roots, including amino acids, organic acids,

sugars, aromatic compounds and secondary metabolites (Walker *et al.*, 2003). The

flavonoids are secondary metabolites exudated by the plants, that initiate the molecular

dialogue between rhizobia and legume. This dialogue culminates with the formation of

new organs in roots called nodules, where the bacterial reduction of atmospheric

nitrogen to the most-needed form -ammonia- takes place (Oldroyd & Downie, 2008).

Thus, when recognized by the appropriate rhizobia, the flavonoids induce the

production of rhizobial signal molecules, called nodulation factors (Nod factors). These

signal molecules can be specifically recognized by legume plants, initiating the

nodulation process that will culminate with nodule development, its occupation and the

transformation of the rhizobium into a nitrogen-fixing cell, the bacteroid (Oldroyd *et al*.,

2011). Besides the molecular recognition mediated by flavonoids and Nod factors, the

success of the nodulation process also depends on secretion systems effectors. Thus, in

the last fifteen years, the effector proteins secreted through the rhizobial type III

secretion system (T3SS) have been proven important, and in some cases essential, for

the symbiotic performance of several rhizobial genera, such as *Sinorhizhobium*,

Rhizobium, *Bradyrhizobium* and *Mesorhizobium* (López-Baena *et al*., 2016; Jiménez-

Guerrero *et al*., 2021, 2022). A novel secretion system has been described in the last

two decades, the type VI secretion system (T6SS). In fact, this machinery was first

identified in *Rhizobium leguminosarum* (Bladergroen *et al*., 2003), although the term

was not stablished until 2006, when the T6SS of *Vibrio cholerae* and *Pseudomonas*

aeruginosa were simultaneously characterized (Mougous, 2006; Pukatzki *et al*., 2009).

The T6SS is present in 25% of gram-negative bacteria, mainly in the Proteobacteria

phylum, where the α-, β- and γ-proteobacteria classes are included (Boyer *et al*., 2009).

Generally, this system secretes effectors/toxins into prokaryotic cells, playing a critical

role in inter-bacterial competition (Ho, *et al*., 2014). However, some type VI-secreted

effectors (T6Es) can target eukaryotic organism cells, having the ability to manipulate

the host during an infective process (Hachani *et al*., 2016).

The specific role of the T6Es in rhizobia symbiosis is mostly unknown, although recent

works suggest that these proteins could exert either neutral, positive, or negative effects,

depending on the symbiotic pair. Thus, it has been reported that the T6SS of

Paraburkholderia phymatum STM 815^{T} and *Azorhizobium caulinodans* do not appear

to be directly implicated in the symbiotic effectiveness with *Vigna unguiculata* and

Sesbania rostrata, respectively. However, it is described that both systems are involved

in symbiotic competitiveness against rhizobial competitors for nodulation (de Campos

et al., 2017; Lin *et al*., 2018). Conversely, in both *Rhizobium etli* Mim1 and

Bradyrhizobium sp. LmicA16, the T6SS is required for efficient nodulation with

Phaseolus vulgaris and *Lupinus* spp, respectively (Salinero-Lanzarote *et al*., 2019;

Tighilt *et al*., 2022). In case of *R. etli* Mim1, this system is synthesized at high cell

densities, in the presence of root exudates and within host-plant nodules (Salinero-

Lanzarote *et al*., 2019), being the T6E Re78 beneficial for competitiveness in bean

nodule occupancy (de Souza *et al*., 2023). Finally, *Rhizobium leguminosarum* RBL5787

is unable to form nitrogen-fixing nodules on pea (*Pisum sativum*) due to the presence of

a functional T6SS (Bladergroen *et al*., 2003).

Structurally, the T6SS is a multiprotein complex composed of thirteen main

constituents. The genes encoding these proteins are grouped in genetic clusters and

named *tss* (type six secretion) (Ho *et al.*, 2014). In some cases, there is an additional set

of genes, named *tag* genes (type six accessory genes), that encode for accessory proteins

with complementary functions . Those proteins have been involved in regulation (TagF,

TagQ, TagR, TagS, TagT) (Lin *et al.*, 2018; Hsu *et al.*, 2009) and fine-tuning the

system assembly (TagP, TagL, TagN, TagW, TagA, TagB, TagJ) (Aschtgen *et al.*,

2010; Santin *et al.*, 2018; Bernal *et al.*, 2021). The T6SS is structured into three main

compartments, and the TssA protein orchestrates the assembly among the different

parts. The membrane complex, formed by TssJ, TssL and TssM, anchors the system to

the cell envelope. TssA interacts with the membrane complex and recruits the baseplate,

composed by TssK, TssE, TssF and TssG. In this new position, TssA primes

polymerization of the tail, formed by an inner tube (Hcp), surrounded by a contractile

sheath (TssB and TssC) and ended in a needle-shaped tip (VrgG and PAAR). The T6Es

can be transported inside the tube, or connected to the tip, being released into the

intracellular environment of the target cell upon sheath contraction. The sheath, which

remains in the cytosol, is then disassembled by an AAA^{+} ATPase, named ClpV (or

TssH), which plays a crucial role in the recycling of the system (Cherrak *et al.*, 2018).

Genes encoding effector/toxin proteins can be found within the T6SS cluster or

scattered around the genome. These genes are usually located downstream genes

encoding VgrG, Hcp or/and PAAR proteins. In some cases, T6Es are encoded by the

same gene that encodes for Hcp, VrgG or PAAR proteins and, therefore, they are fused

to them at the C-terminal end. Consequently, structural proteins with a C-terminal

cytotoxic domain are termed evolved, named as Hcp_{e}, $VrgG_{e}$ or $PAAR_{e}$ (Durand *et al.*,

2014). The antibacterial T6Es can target the periplasm; these include peptidoglycan

hydrolases (Tae1-4 and Tge1-3), phospholipases (Tle1-4) and pore-forming effectors

(VasX). Other T6Es act in the bacterial cytoplasm, such as the nucleases effectors (Tde1

and Tke2), or those that degrade essential cofactors - $NAD(P)^+$, like the hydrolase

effectors (Tse6/Tne1 and Tne2) (Coulthurst, 2019). Bacteria with a functional T6SS

produce immunity proteins to protect from sister cells attack and self-intoxication.

Contrary, T6Es delivered into eukaryotic host cells are less widespread than

antimicrobial effectors, but the few identified so far are involved in different steps of

host manipulation to promote bacterial infection (Hachani *et al.*, 2016).

Sinorhizobium fredii USDA257, hereafter USDA257, is a fast-growing rhizobium that

was isolated from wild soybean (*Glycine soja*), but is also able to form nitrogen-fixing

nodules in a wide variety of legume species, such as *G. max*, *S. rostrata, V. radiata, P.*

vulgaris, *Cajanus cajan*, *Lotus japonicus*, and *L. burtii,* among others (Pueppke &

Broughton, 1999). USDA257 is one of the most versatile rhizobia, along with other

bacteria of the same species, *S. fredii* NGR234 and *S. fredii* HH103. Interestingly, only

USDA257 presents both T3SS and T6SS machinery. The T3SS of USDA257 has been

extensively studied,playing a prominent role in symbiosis, as well asin tdetermining of

its host-range of nodulation (Staehelin & Krishnan, 2015). Legume recognition of type

III-secreted effectors(T3Es) can exert both positive (induction of nodulation) or

negative (inhibition of nodulation) extreme effects, depending on the plant cultivar

(Jiménez-Guerrero *et al.*, 2022). However, ecological and physiological function of the

USDA257 T6SS is completely unknown in the *Sinorhizobium* genera.

In this work, we identified and characterized the USDA257 T6SS, which exhibits novel

characteristics of the structural proteins, implying an apparatus with a distinctive

assembly and a definite set of T6Es. We showed that this USDA257 secretion system is

especially active in non-rich media during stationary phase of growth and when

colonizes legume root nodules. In fact, nodulation assays demonstrated that USDA257

utilizes this protein secretion system to enhance nodulation and competitiveness with *G.*

max cv Pekin. According to *in silico* predictions, improvement in the symbiotic

performance could be directly mediated by essential T6Es and/or indirectly by secreting

antibacterial toxins during the nodulation process.

2. Materials and Methods

2.1. Bioinformatical analysis

Sequences of 161 TssB proteins from 153 strains, belonging to 12 genera were obtained

and compiled by using the BLASTp tool from NCBI website (Boratyn *et al*., 2013).

Sequences were aligned with the ClustalW software (Sievers *et al*., 2011) and the

construction of the phylogenetic tree was carried out using the MEGA7 software

(Kumar *et al*., 2016), applying the Maximum Likehood algorithm (Le *et al*., 2008) and a

JTT matrix model (bootstrap value=500). The phylogenetic tree was customized with

the iTOL tool (Letunic & Bork, 2016). Amino acid sequence searches were carried out

using SMART (Letunic *et al*., 2015) and Pfam (Finn *et al*., 2016). The Protein

Homology/analogy Recognition Engine (Phyre[2]) server was used to perform a template-

based approach to predict protein structural homology (Kelley *et al*., 2015). An

additional analysis to identify structural homologs was performed using the recent

algorithm Foldseek (Van Kempen *et al*., 2024), that is a structural alignment tool based

on a structural protein alphabet of tertiary interactions that operates using predicted

models from the Alphafold database (Abramson *et al*., 2024). Molecular weight and

isoelectric point for each protein were calculated with the ExPaSy software (Artimo *et*

al., 2012). PSORTb server was used to predict sub-cellular location of proteins (Yu *et*

al., 2010), TMHMM software to predict transmembrane domains (Krogh *et al*., 2001),

and SignalP and PSORTb to predict signal peptides (Almagro Armenteros *et al.*, 2019).

The figures for the cluster alignments were created using Easyfig (Sullivan *et al.*, 2011).

2.2. Bacterial strains and plasmids

Bacterial strains and plasmids used in this work are listed in **Table S1**. Rhizobial strains

used in this study were grown at 28 °C on tryptone yeast (TY) medium (Beringer,

1974), yeast extract mannitol (YM) medium (Vincent, 1970) or minimal (MM) medium

(Robertsen *et al.*, 1981) with two different mannitol concentrations (3 or 10 g mL^{-1},

YM3/MM3 or YM10/MM10, respectively). *Agrobacterium tumefaciens* and

Pectobacterium carotovorum strains were cultured on LB medium (Bertani, 1951) at 37

°C in the first case, and 28 °C in the last two ones. When required, the media were

supplemented with the appropriate antibiotics, as previously described (Lamrabet *et al.*,

1999). Genistein was dissolved in ethanol and used at 1 µg ml^{-1} to give a final

concentration of 3.7 µM.

Plasmids were transferred from *E. coli* to *Sinorhizobium* strains by triparental

conjugation, as described by Simon (1984), using plasmid pRK2013 as helper (Figurski

& Helinski, 1979). Recombinant DNA techniques were performed according to the

general protocols of Sambrook *et al.* (1989). PCR amplifications were performed as

previously described López-Baena *et al.*, (2009). Primer pairs used for the amplification

of the USDA257 genes are summarized in **Table S2**.

For the obtention of a USDA257 *tssA* mutant, an internal fragment of this gene was

amplified using specific primers, digested with *Eco*RI and *Bam*HI restriction enzymes

(both restriction sites were added in forward and reverse 5′ primer ends, respectively)

and cloned into pK18*mob* vector, which was previously digested with the same

enzymes, obtaining plasmid pMUS1480. This construction was employed for the

homogenotization of the mutated version of *tssA* in USDA257, generating the mutant

strain. Double recombination events was confirmed by Southern blot, PCR and

sequencing (data not shown). For hybridization, DNA was blotted to Hybond-N nylon

membranes (Amersham, UK), and the DigDNA method of Roche (Switzerland) was

employed according to the manufacturer's instructions (**Fig. S1**).

Promotor regions of the *ppkA* USDA257 T6SS gene was amplified with specific

primers (575 bp) and the resulting DNA fragment was digested with *Eco*RI and *Xba*I

(both restriction sites were added in forward and reverse 5´ primer ends, respectively)

and cloned into plasmid pMP220, upstream of the *lacZ* gene, being previously digested

with same enzymes, obtaining plasmid pMP220::P$_{ppkA}$.

Construction of the dual reporter strain of USDA257 was performed using the plasmid

developed by Shamal and Chatterjee (2021) and inserting the promoter region of the

ppkA gene (530 pb), that was previously amplified and digested with *Eco*RI (restriction

sites were added in forward and reverse 5´ primer ends). Briefly, transcriptional fusion

was constructed by fusing the promoter region upstream of the *ppkA* gene (P$_{ppkA}$) to the

red fluorescent protein gene (*rfp*) and cloned into a pBBR-MCS-4 plasmid, that is

constitutively expressing the *gfp* gene (P$_{kan}$::*gfp*). This plasmid was conjugated with the

USDA257 strain for obtaining the dual-reporter strain of USDA257 (pBBR4::P$_{kan}$::*gfp*-

P$_{ppkA}$::*rfp*). As a control, the plasmid pBBR4::P$_{kan}$::*gfp*-P$_{w/o}$::*rfp* (Shamal and Chatterjee,

2021), in which the *rfp* gene lacks the promoter region, was also transferred by

conjugation to USDA257. All plasmids were stably maintained during infection.

For USDA257 Hcp antibody production, the *hcp* ORF without the stop codon was

amplified by PCR with specific primers and cloned into the entry plasmid pDONR207

and then mobilized into pET42 plasmid following the Gateway cloning system

technology (Invitrogen, USA). The generated plasmid was transferred to *E. coli* BL21

(DE3) (Studier and Moffatt, 1986) by the heat shock method (Froger and Hall, 2007).

2.3. Determination of β-galactosidase activity

To determinate the β-galactosidase activity, the USDA257 wild-type strain was

conjugated with the plasmids pMP220::P_{ppkA} and pMP240, that contains transcriptional

fusions between the *ppkA* (USDA257) and *nodA* (*R. leguminosarum*) promoters,

respectively, both upstream of the *lacZ* gene. The empty plasmid pMP220 was also

transferred to USDA257 and used as negative control. Assays of β-galactosidase

activity were carried out as described by Zaat *et al* (Zaat *et al*., 1987). Units of β-

galactosidase activity were calculated according to Miller (Miller, 1972). The

experiments were repeated four times with three technical replicates each time.

2.4. RNA extraction and qRT-PCR experiments

Total RNA was isolated using a High Pure RNA Isolation Kit (Roche, Switzerland),

according to the manufacturer's instructions. Verification of the amount and quality of

total RNA samples was carried out using a Nanodrop 1000 spectrophotometer (Thermo

Scientific, USA) and a Qubit 2.0 Fluorometer (Invitrogen, USA). Four independent total

RNA extractions were obtained for each condition. This (DNA-free) RNA was reverse

transcribed into cDNA by using PrimeScript RT reagent Kit with gDNA Eraser (Takara,

Japan). Quantitative PCR was performed using a LightCycler 480 (Roche, Switzerland)

equipment with the following conditions: 95 °C, 10 min; 95 °C, 30 s; 50 °C, 30 s; 72

°C, 20 s; forty cycles, followed by the melting curve profile from 60 to 95 °C to verify

the specificity of the reaction. The USDA257 16S rRNA gene was used as an internal

control to normalize gene expression. The fold-changes of four biological samples with

three technical replicates of each condition were obtained using the ΔΔCt method

(Pfaffl, 2001). Selected genes and primers are listed in **Table S2**.

2.5. Expression and purification of recombinant Hcp for specific antibody generation

The *E. coli* BL21(DE3) strain carrying plasmid pET42::*hcp* was inoculated in 5 mL of

LB supplemented with carbenicillin and incubated overnight at 37 ℃ with shaking at

200 rpm. The culture was transferred to 200 mL in the same medium and incubated in

the same conditions. When the OD_{600} reached 0.6, protein expression was induced with

1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG). Then, cultures were grown for 4

hours at 37 ℃ with shaking at 200 rpm. Cells were harvested by centrifugation (5.000 g,

20 min, 4 °C) and the pellet was resuspended in 50 mM Tris-HCl (pH 7.5), 250 mM

NaCl (buffer NPI) containing 10 mM imidazole, 1 mg/mL of lysozyme and protease

inhibitors (Roche, Switzerland). The suspension was incubated for 30 min and sonicated

on ice 5 times for 30 sec, with 30-sec cooling intervals between sonication treatments.

Cell debris was eliminated by centrifugation (10.000 g, 30 min, 4 °C). Clarified lysate

was filtered with a 0.45 nm filter and incubated in a column containing 2.5 mL of

nickel-Sepharose resin (Protino Ni-NTA, Macherey-Nagel, Duren, Germany),

previosuly equilibrated with 10 volumes of buffer NPI allowing binding of His-tagged

Hcp to Protino Ni-NTA agarose. Columns were washed with 10 volumes of buffer NPI

containing 100 mM imidazole and, finally, the His-tagged Hcp protein was eluted off

the resin with 1 mL of buffer NPI containing 500 mM imidazole. The His-tagged Hcp

protein was subsequently washed with PBS and concentrated using Amicon Ultra

Centrifugal Filter Unit (Milipore Sigma, Burlington, MA, USA), following the

manufacturer's instructions. Expression and purification of His-tagged Hcp were

verified by SDS-PAGE and confirmed by Western blot using His-tag monoclonal

antibody. Polyclonal antibody production was carried out by the "Centro de

Experimentación Animal Óscar Pintado" from the University of Sevilla (Spain)

following the procedure described by Vidal et al. (1980). through their speedy 28-Day

program (1 rabbit; 4 protein injections and 2 bleedings).

2.6. Purification and analysis of extracellular proteins

Extracellular and intracellular proteins were recovered following the protocol described

by Hatachi et al. (2011) with some modification. Briefly, 20 ml of the different rhizobial

cultures grown on an orbital shaker (180 rpm) at 28 ℃ for 48 h (stationary phase, with

an adjusted OD_{600} of 1, which corresponds to approximately 10^9 bacteria ml^{-1}) were

centrifuged for 20 min at 10.000 g at 4 ℃. Bacterial pellets were normalized and added

directly to 200 μl of sample buffer (62.5 mM Tris-HCl [pH 6.8], 2% SDS [m/v], 10%

glycerol [v/v], 5% β-mercaptoethanol [m/v], and 0.001% bromophenol blue [m/v]). To

eliminate any remaining cell in the supernatant, 3 additional sequential centrifugations

(20 min, 10.000 g, 4 ℃) were performed. A volumen of 1.8 mL from each culture

supernatants were collected and precipitated with trichloroacetic (TCA) acid overnight

at -20 ℃. The mixtures were centrifuged for 30 min at 16.000 g at 4 ℃. Dried pellets,

previously washed with 90% acetone, were resuspended in sample buffer. For

immunostaining, proteins were separated on SDS 20-4% (m/v) polyacrylamide gels

(Bio-Rad, USA) and electroblotted to Immun-Blot polyvinylidene difluoride

membranes (Bio-Rad) using a Mini Trans-Blot electrophoretic transfer cell (Bio-Rad).

Membranes were blocked with TBS containing 2% (m/v) bovine serum albumin (BSA)

and then incubated with antibodies raised against the USDA257 Hcp protein diluted

1:1000 in the same solution. Anti-rabbit HRP-linked antibody was used as secondary

antibody and reaction results were visualized using CDP Start Detection Reagent kits

(GE Healthcare, USA) according to manufacturer's instructions. Bands were visualized

using an ImageQuant LAS 500 imaging workstation instrument (GE Healthcare, USA).

2.7. Interbacterial competition assays

In vitro competition assays were performed on the USDA257 T6SS-inducing YM3 and

MM3 agar (2% w/v) plates, as previously described (Bernal *et al.*, 2021). Briefly,

overnight bacterial cultures were washed and adjusted to an OD_{600} of 1.0 in sterile

PBS,and mixed at a 1:1 ratio (USDA257:prey). *A. tumefaciens*, *P. carotovorum* and *S.*

fredii HH103 were used as preys. 100 µl of mixtures were grown on YM3 (for *A.*

tumefaciens and *P. carotovorum*) or MM3 (for *S. fredii* HH103) agar plates at 30 °C for

24 or 48 hours and then collected using an inoculating loop and resuspended in sterile

PBS. The outcome of the competition was quantified by counting colony forming units

(CFUs) using antibiotic selection of the input (time = 0 hours) and output (time = 24-48

hours). *A. tumefaciens* and *P. carotovorum* prey strains harbored the plasmid pRL662,

which confers resistance to gentamicin and was used for antibiotic selection, whereas

USDA257 was naturally resistant to streptomycin. Three biologically independent

experiments were performed.

2.8. Plant tests

For the evaluation of the symbiotic phenotypes in nodulation assays, the wild-type and

its derivative mutant strains were grown in YM3 medium. Surface-sterilized seeds of *G.*

max cv Pekin were pre-germinated and placed in sterilized Leonard jars , containing

Fårhaeus N-free solution (Vincent, 1970). Germinated seeds were then inoculated with

1 mL of bacterial culture with an adjusted OD_{600} of 0.6. Growth conditions were 16 h at

26 ºC in the light and 8 h and 18 ºC in the dark, with 70% humidity. Nodulation

parameters were evaluated after 6 weeks. Shoots were dried at 70 ℃ for 48 h and

weighed. Nodulation experiments were performed three times with five replicates for

each treatment.

Competition for nodulation experiments (competitiveness) on *G. max* cv Pekin were

performed using USDA257 and the non-T6SS-carrying *S. fredii* HH103 rhizobium .

These bacteria were grown to 10^9 cells ml^{-1}, and four to five Leonard jar assemblies

containing two plant seedlings each were inoculated with 1 ml of a mixture of bacterial

competitors in 1:1, 1:10 and 10:1 ratios. Plants were grown for 6 weeks in a plant

growth chamber under the growth conditions described above. To identify bacteria

occupying the nodules,100 *G. max* cv Pekin nodules from each treatment were surface

sterilized by immersing them in 5% [wt/vol] sodium hypochlorite (for 5 min, followed

by five washing steps in sterilized distilled water. The effectiveness of the surface-

sterilizing treatment was checked by inoculating TY plates with 20-µl aliquots from the

last washing step. Individual surface sterilized nodules were crushed in 30 µl of

sterilized distilled water, and 20-µl aliquots were used to inoculate TY plates. Nodule

occupancy was determined by assessing the ratio of differential natural antibiotic

resistant of isolates (Rifampicin for USDA257 and Streptomycin for HH103). At least

10 colonies from each isolate were analyzed in order to check the possibility of nodules

containing both inoculants.

For nodule occupancy visualization, 30-day-old *L. burttii* nodules formed in plants

inoculated with the dual fluorescent reporters of USDA257 were embedded in 6%

agarose in water and sliced in thick layer sections (50 µm) using a Leica VT 1000S

vibratome (Wetzlar, Germany). Functional fluorophores of sections of nodules were

stained with 0.04% calcofluor and observed by using a Leica Stellaris 8 SPE Confocal

Microscope (Leica Microsystems)   (Jena, Germany) fluorescence microscope as

previously described (Kawarada *et al*., 2017). The image´s contrast and intensity were

adjusted using ImageJ software (Schneider *et al*., 2012).

2.9. Statistical analysis

The statistical tests performed in this work and indicated in the Figure legends were

done with Prism 8 (GraphPad, La Jolla, CA, USA).

3. Results

3.1. Genome-wide screening for T6SSs in N$_2$-fixing bacteria

We have performed a T6SS phylogenetic analyses of 160 N$_2$-fixing bacteria from 13

different genera all belonging to the Phylum Proteobacteria and including the main

genera containing root-nodule forming bacteria (**Table S3**). The selected species belong

to the order *Hyphomicrobiales* (better known as *Rhizobiales*) from the α-Proteobacteria

class and the order *Bulkholderiales* from the β-Proteobacteria class. We have included

well-described *Agrobacterium* and *Pseudomonas* T6SSs to identify and locate the

previously described T6SS phylogenetic groups (Wettstadt *et al*., 2020). The tree

displays in **Fig. 1** contains 160 TssB proteins**,** showing the phylogenetic distribution of

selected rhizobia T6SSs. Our analysis shows that rhizobial T6SSs are distributed among

the five main clades previously described (Boyer *et al*., 2009). Most of them, 140

(87.5%) belong to groups 3 (54, 33.75%) and 5 (86, 53.75%) (**Fig. 1**). Group 3 contains

a great variety of species from the *Ensinfer*, *Rhizobium*, *Bradyrhizobium* and

Mesorhizobium genera among others, as well as the first-discovered *P. aeruginosa* H1-

T6SS. Group 5 contains mostly *Rhizobium* and *Azorhizbium* species and includes the

well-studied *A. tumefaciens* T6SS. Although phylogenetically-distance *Cupriavidus* and

Parabulkholderia species can be found in both groups. Minority groups 2 and 4

principally contain *Paraburkholderia* species (2, 1.25% each) and group 1 (14, 9.75%)

contains predominantly species of the *Methylobacterium* genus.

Some bacteria groups, namely *Pseudomonas,* frequently contain 2 or more T6SS

clusters (Bernal *et al*, 2017). However, despite the number of T6SS clusters in a strain

ranging from 1 to 5, in a broader group such a phytobacteria, only an estimate 7% of

strains contain more than one cluster (Bernal *et al*., 2018). This number is even smaller

among rhizobial T6SSs where most strains contain only a single T6SS cluster.

3.2. The reference rhizobial strain S. fredii USDA257 possesses a complete T6SS

Inspection of USDA257 genome revealed 26 T6SS-related ORFs located in a

chromosomal cluster (**Fig. 2A, Table 1**). Thirteen genes (*tssABCDEFGHIJKLM*)

encode the structural proteins required for a functional T6SS, including the membrane

complex, the base plate, the tail and the ATPase that recycles the system. We further

identified genes encoding a previously described regulatory phosphorylation cascade

(Mougous *et al*., 2007), including the threonine kinase/phosphatase pair (PpkA-PppA)

and the phosphorylation substrate (Fha). TagF is a posttranslational repressor that

regulates T6SS via Fha interaction. TagF-linked T6SS repression have been studied in

several systems, including *P. aeruginosa* and *A. tumefaciens* T6SSs (Lin *et al*., 2018). A

putative TagF protein is encoded in USDA257 T6SS cluster, indicating that this system

might have a similar posttranscriptional regulation to both *P. aeruginosa* and *A.*

tumefaciens.

TssJ, TssL and TssM are the core components of the T6SS membrane complex that

docks the system to the cell envelope. TssL (DotU domain) and TssM (IcmF domains)

are anchored to the inner membrane through one and three transmembrane helices

respectively and form a channel through the cell envelope by interaction with each other

and the outer membrane lipoprotein TssJ. Additional membrane-complex-related

proteins have been described in some systems, including a set of proteins that anchor

the T6SS to the peptidoglycan (PG) through the PG-binding domain OmpA

(pfam00691) (Aschtgen *et al*., 2010). The set included the specialized TssL with a

canonical N-terminal DotU domain and a C-terminal OmpA domain that have been

identified in *P. aeruginosa* and *A. tumefaciens* among others. The authors also described

a ~250 amino acids protein with an N-terminal signal peptide and a C-terminal OmpA

domain identified in *Burkholderia pseudomallei* and *Ralstonia solanacearum* and

named TagN. The localization, topology, PG-binding, and interactions with the rest of

the membrane complex remains to be characterized for this T6SS protein.

An in-depth analysis of USDA257 T6SS components, we identified the three core

components of the membrane complex *i.e.,* TssJ, TssL and TssM, and two related

accessory proteins named, TagM and TagN (**Fig. 2B**). In addition, USDA257 TssJ and

TssM proteins contain the conventional T6SS domains, while USDA257 TssL is a

specialized TssL (TssL$_e$). We identified two additional accessory components, a TagN

homologue and TagM, an 821 amino acids protein that shares certain degree of

homology with both TssM and TssL (**Fig. S2**). This protein contains an N-terminal

DotU domain, a single transmembrane domain and two IcmF domains; its

transmembrane domain and the absence of a specific signal peptide indicates that this

protein might be anchored to the inner membrane. USDA257 membrane complex varies

from the canonical, presenting not only one but two membrane complex proteins with

PG-binding domains (TssL$_e$ and TagN) and a TssM-TssL hybrid protein (TagM) (**Fig.**

2B). This complexity might indicate a different assembling mode/capacity to anchor to

the cell envelope.

Lastly, we have identified a gene encoding an additional accessory protein we have

named *tagY* (**Fig. 2A**). TagY presents homology with an M15 peptidase, a D-alanyl D-

alanine carboxypeptidase that is involved in bacterial cell wall biosynthesis (Lessard *et*

al., 1999). Predictions indicate that TagY has a transmembrane domain at the N-

terminal end and depending on the software used, it might have a signal peptide,

suggesting that this protein could be anchored to the outer membrane with the C-

terminal end facing the periplasm or the cellular exterior.

3.3. *The S. fredii USDA257 T6SS cluster harbors genes encoding potential effectors*

Frequently, T6SS effectors are genetically linked to *vgrG* and *hcp* genes, which can be

found within a T6SS cluster or, in different numbers, scattered through the genome. For

instance, *P. aeruginosa* PAO1 genome contains 10 *vgrG* and 5 *hcp* genes (Hachani *et*

al., 2016). The *in silico* study of the USDA257 genome has revealed single copies of

vrgG and *hcp* genes in the T6SS cluster (**Fig. 2A**). *hcp* is found surrounded by the

structural genes *tssC* and *tssE* and there is no evidence of genes encoding putative

effectors in the *hcp* proximity. On the other hand, *vrgG* is located at the end of the

cluster and genetically linked to putative T6SS effectors and adaptors, displaying a

similar genetic architecture to *P. aeruginosa* PAO1 and *P. fluorescens* F113 *vrgG1b*

clusters (Hachani *et al*., 2011; Redondo-Nieto *et al*., 2013). These clusters encoded an

evolved VrgG, an oligonucleotide-binding(OB)-fold, a DUF2169 adaptor (an

immunoglobulin-like protein), a thiolase-like protein (PRK06147) proposed to be a

novel T6SS adaptor, a PAAR protein with a C-terminal cytotoxic domain, an immunity

protein and a heat repeat-containing protein. Both sets of genes only differ in the

sequence of the toxic domain and the immunity pair, a common characteristic of this

genetic island, previously described by Pissaridaou *et al*. (2018). In USDA257, genes

downstream *vrgG* are inverted, but aside from that, the genetic organization of the

region is conserved compared to previously described ones. Thus, USDA257 contains

the genes encoding an evolved VrgG with a similar C-terminal domain, the OB-fold

protein, the DU2169 adaptor named *tsar* (type six adapter rhizobium), followed by a

thiolase-like protein (T6SS adaptor), an evolved PAAR protein named Tsre1 (*tsre*, type

six rhizobial effector) with a C-terminal domain of unknown function and, lastly, a

protein named Tsre2 (**Fig. 2A, Tables 1 and 2**). According to Phyre[2], in which model is

built based on the experimental structure of a homologue that serves as a structural

template (Kelley *et al*., 2015), Tsre2 shows structural homology with a glycoside

hydrolase enzyme GH74 with a xyloglucan binding domain from *Caldicellulosiruptor*

lactoaceticus 6A (69% alignment, 99.3% confidence). In addittion, the predicted model

of the USDA257 Tsre2 protein was collected from the Alphafold database and queried

using the FoldSeek tool, that predict the structural homology based on the terciary

interactions of proteins in a sequence-independent manner (Van Kempen *et al.*, 2023).

Using this approach, best prediction for the structural homology is found with the

immune protein Tsi7 from *P. aeruginosa* PAO (probable structural homology between

amino acid residues 2 and 340 with an E-Value of 3.77e-14), whereas the structural

homology with the glycoside hydrolase enzyme ranges between residues 11 and 338

(1.10e-1 E-Value). Interestingly, the evolved PAAR protein named Tsre1, which does

not display any structural homology in its C-terminal domain according to Phyre[2], is

predicted to be homologous to the DNase toxin Tse7 of *P. aeruginosa* PAO1 (ranging

from 3 to 321 amino acid positions, 5.25e-24 E-Value).

Not only the *paar* genes of USDA257 but also *vgrG* gene encode for evolved proteins,

in which C-terminal ends contain putative effector domains (Pukatzki *et al*., 2009).

According to sequence similarity, the C-terminal domain of VgrG corresponds to a 225

amino acid fragment from the FhaB protein, a chaperone that prevents the premature

folding of hemagglutinin filaments, one of the virulence factors of *Bordetella pertussis*

(Coutte *et al*., 2001). However, the analysis performed with Phyre[2] software indicates

that this C-terminal domain also shows structural homology with the Cthe_2159 protein

from *Hungateiclostridium thermocellum* ATCC 27405 (85% alignment, 86.5%

confidence), a polysaccharide lyase-type protein that binds to cellulose and

polygalacturonic acid (**Fig. 2A, Table 2**). The predicted model of the evolved VrgG

from USDA257 using Foldseek determined that the *Nitrospirae bacterium* structural

homologous VgrG is the most similar protein among all proteins (between amino acid

residues 3 and 768, 1.40e-75 E-Value).

Interestingly, we identified one potential orphan effector, since Tsar adaptor and

evolved PAAR proteins, termed Tsre3, were found separately from the T6SS cluster

(**Fig. 2A**). Curiously, this orphan PAAR protein displays sequence dissimilarity with the

Tsre1 effector both in the N-terminal and C-terminal domains and does not exhibit

structural homology with any previously described proteins according to Phyre[2]. Using

FoldSeek, the structure of a DUF4150 domain-containing protein with unknown

function from *Methilobacterium* sp. yr596 is identified with probable structural

homology only with the N-terminal part (amino acid residues from 38 to 168 with a E-

Value 1.11e-16).

3.4. *The vrg cluster is well conserved among rhizobia except for the effector region*

To obtain a broader perspective of functioning and distribution of *tsre1* and *tsre2* genes

among rhizobia, we performed a comparison of the *vgrG* clusters from 18 rhizobial

strains. **Fig. 3** shows sequence conservation across the entire cluster. As expected, the

genes encoding the VgrG and the OB-fold proteins are conserved and the region

encoding the putative proteins Tsre1 and Tsre2 is the one varying the most. Curiously, the 3´end of the gene encoding the DUF2169 adaptor and 5´end the gene encoding the thiolase-like protein are less conserved and could indicate that they are effector/toxin-specific.

A neighbor-joining phylogeny of the Tsre1 and Tsre2 proteins from these rhizobial strains clearly group USDA257 with *Sinorhizobium americanum* strains CCGM7 and CFNEI 73 for both proteins, whereas most Tsre1 and Tsre2 *Rhizobium phaseoli* versions are distributed in two separated branches, one grouping versions from strains R620, R650, R611, N771, N671, N261, R723 and Brasil 5, and the other one clustering proteins from strains N841, R630, N831, N931 and R744 (**Fig. 4**). Alignment of the Tsre1 protein sequences indicated a high degree of similarity among the N-terminal but not in the C-terminal domains in all versions, that were grouped as in the phylogenetic tree (**Fig. S3-5**). In this case, the first half of the three versions of Tsre1 proteins showed a clear structural homology with PAAR domain according to Phyre[2] (**Table 2**). Despite alignment of Tsre2 versions showeing a lower degree of similarity among proteins from different branches of the phylogenetic tree (**Fig. S6-8**), the structural homology predicted using this approach indicates that the other two versions are structurally similar than the same GH74 glycoside hydrolase enzyme, differencing in their range of alignment. When using the Foldseek tool, the proteins most structurally similar to the N841 versions of both Tsre1 and Tsre2 are Tse7 and Tsi7 from *Pseudomonas aeruginosa* PAO1. In contrast, for the R620 version, the lack of an AlphaFold prediction made structural comparison of the protein impossible using this approach (**Table 2**).

3.5. The T6SS of S. fredii USDA257 is functional and induced in minimal medium and

in the nodules

To determine the conditions under which the T6SS of *S. fredii* USDA257 is expressed,

we cloned the promoter region of the first gene of the T6SS structural operon (*ppkA*)

upstream of the reporter gene *lacZ*. Thus, by ß-galactosidase assays, we aimed to

measure the expression levels of the USDA257 T6SS genes. We tested USDA257

cultures 30 hours post inoculation in selected rich, standard and minimal media for

growing rhizobia: TY, YM, and MM respectively (**Fig. 5A**). YM is a standard medium

possessing moderate quantities of yeast extract that supply the limiting organic nitrogen

and micronutrients, whereas MM is a minimal medium that only has glutamate as

nitrogen source. Both YM and MM media can be prepared with different concentrations

of mannitol as carbon source. Thus, the activation pattern in the ß-galactosidase assays

of the wild-type strain carrying the construct containing the USDA257 *ppkA* promoter

region fused to the *lacZ* reporter gene indicated that the expression of the T6SS genes is

highest in the minimal medium MM3 (3 g L^{-1} of mannitol), whereas the lowest values

were obtained when this rhizobium was grown in the rich medium TY (about 3-fold and

1,8-fold, respectively, in comparison to those values obtained by the strain carrying the

empty plasmid) (**Fig. 5A**). To confirm these findings, we followed a complementary

approach, measuring the transcriptional activation of the USDA257 *ppkA* gene. *q*RT-

PCR experiments shed the same transcriptional activation trend for the *ppkA* gene in

USDA257 cultures after 48 hours post inoculation. The highest transcriptional values

were found in cultures growing at MM3 medium (around 8-fold) in comparison to the

values obtained for the same genes in TY medium (**Fig. 5B**). To delve further into the

regulation of T6SS, we performed ß-galactosidase assays at different points of the

USDA257 growth curve, inoculating the bacterium in the MM3 inducing-medium.

Interestingly, the activation pattern of the wild-type strain carrying the P_{ppkA}::*lacZ*

fusion pointed out that the expression of the T6SS follows the archetypical growth

curve profile, with a saturation of the ß-galactosidase values around 54 hours after

inoculation (about 7-fold in comparison to those values obtained by the strain carrying

the empty plasmid) (**Fig. 5C**).

To gain insight about the potential role of the USDA257 T6SS in the early steps of the

symbiotic process, β-galactosidase assays were performed in the presence of genistein, a

nod-gene inducing flavonoid in several related *S. fredii* strains (Jiménez-Guerrero *et al*.,

2015, 2018; Pérez-Montaño *et al*., 2014). Activation of the *ppkA* promoter was

measured in MM3 medium at stationary phase in the presence or the absence of

genistein, using as control of *nod*-gene induction the wild-type strain carrying the

pMP240 plasmid, which harbors the conserved *nodA* promoter of *R. leguminosarum* bv.

viciae fused to the *lacZ* reporter gene. In the presence of the USDA257 *nod* gene-

inducing flavonoid genistein, induction values of strains carrying the *ppkA* promotor

fused to *lacZ* were similar among conditions (**Fig. S9**), indicating that the expression of

the T6SS is not controlled by the presence of this symbiotic-plant molecule.

The extracellular secretion of the Hcp protein into a culture medium is a hallmark to

assess functionality of the T6SS system. Therefore, we produced a specific polyclonal

antibody against USDA257 Hcp to assess USDA257 T6SS activity in the previously

described conditions, *i.e.* the different culture media after 48 hours of growth (**Fig. 6A**).

As anticipated, Western blot analysis corroborated the results obtained from both

transcriptomic assays, revealing that the presence of Hcp was lowest in TY medium and

highest in both MM media. We also used a USDA257 *tssA* mutant strain to disable the

system, since TssA is an essential component for T6SS activity. As expected, we

detected Hcp in the supernatant of the wild-type strain grown in MM3 but not in the

isogenic *tssA* mutant, while it was present in the cytosol of both strains (**Fig. 6B**).

Altogether, results stablishe that USDA257 T6SS is a functional secretion machinery,

and it is found most active in poor media at the stationary phase of growth.

Finally, to elucidate the USDA257 T6SS potential functioning specifically during

nodule colonization in the rhizobial symbiosis, we conducted a detailed investigation by

confocal microscopy into the localization of rhizobial cells within nodules, along with

the activation pattern of their T6SS. The expression of both constitutive GFP and T6SS-

responsive RFP allowed to account for all dual-bioreporter cells *in vivo* during the

infection of legume nodules. Interestingly, strong constitutive GFP and T6SS-

responsive RFP fluorescence expression were observed for all rhizobial cells that are

colonizing the symbiotic cells of *L. burttii* nodules (**Fig. 7C**), which reveales that the

T6SS of USDA257 is fully active when rhizobial cell invade the host nodule tissue.

3.6. The T6SS of S. fredii USDA257 is required for a successful nodulation and

competitivenes with G. max cv Pekin

Due to many bacterial T6SS are mainly involved in interbacterial competition, we first

performed interbacterial competition assays. These experiments we carried out between

the wild-type and a *ttsA* mutant (T6SS-defective) strains of USDA257 as predators and

other bacterial species (*A. tumefaciens*, *P. carotovorum* and *S. fredii* HH103) as preys.

The bacteria were mixed 1:1 (predator:prey) and co-cultured on USDA257 T6SS-

inducing medium for 24 and 48 hours. Spots containing the competing bacteria were

collected and the bacterial CFUs were determined on selective plates. The number of

predator and prey bacteria after competing is shown in **Table 3**. In addition to the fact

that no differences were observed between the wild-type and mutant strains in all

conditions, both wild-type and the *tssA* mutant strains displayed a reduced number of

bacteria when competing with any prey. These results may indicate that the USDA257

T6SS is not involved in interbacterial competition *in vitro* against another potential

rhizospheric competitor as the pathogenic *A. tumefaciens* and *P. carotovorum* or the

symbiont *S. fredii* HH103 (Bernal *et al*., 2017).

To gain insight into the role and functioning of the *S. fredii* USDA257 T6SS, the

symbiotic abilities of the wild-type or its derivate *tssA* mutant strains with *G. max* cv

Pekin, considered its natural host, were inspected (**Fig. 7A**). Interestingly, the number of

nodules in examined plants inoculated by the *tssA* mutant was statistically lower than

those obtained in plants infected by the wild-type strain. Similar but not statistically

trends were observed for both nodule fresh mass and plant-top dry mass parameters. On

the other hand,the impact of the T6SS on the bacterial competitive nodulation capacity

with *G. max* cv Pekin was assessed by co-inoculation experiments with USDA257 and

the *tssA* mutant. In all ratios (1:1, 1:10 and 10:1), the T6SS defected strain was clearly

outcompeted by the wild-type (**Fig. 7B**). Thus, results obtained from nodulation assays

and competitiveness for nodule occupancy indicate that the T6SS of USDA257 is

required for an optimal nodule development in soybean.

4. Discussion

Application of symbiotic rhizobia is a cheap and ecofriendly alternative source of

nitrogen for plants, which favors a moderate use of nitrogen fertilizers mitigating their

excessive utilization (Martinelli *et al*., 2020; Pérez-Montaño *et al*., 2014). The

efficiency of symbiosis depends on various factors, such as the soil, the environment,

and the symbiotic pair. From the rhizobia point of view, several molecules are important

in this intimate interaction, highlighting Nod factors, extracellular polysaccharides, and

protein secretion systems (López-Baena *et al*., 2016). In case of T3SS, numerous

examples have been described to exert neutral, positive or negative effects on symbiosis

depending on the plant host (Nelson & Sadowsky, 2015; Jiménez-Guerrero *et al*.,

2022). However, studies about the role of rhizobial T6SS on symbiosis are scarce,

despite the first report dealing with T6SS indicated that this protein secretion system

impairs nodulation between *R. leguminosarum* and pea (Bladergroen *et al*., 2003).

Conversely, positive effect of T6SS on symbiosis has been recently reported in the

symbiosystems *R. etli*-bean and *Bradyrhizobium-Lupinus* (Salinero-Lanzarote *et al*.,

2019, Tighilt *et al*., 2022). In two additional studies, it was demonstrated a role of T6SS

on interbacterial competition but not in symbiosis (de Campos *et al*., 2017; Lin *et al*.,

2018). In the case of *Sinorhizobium* genera, no studies have been performed so far.

The T6SSs distributed among *Proteobacteria* phylum are classified in five phylogenetic

groups (Boyer *et al*., 2009). Reference strains for each phylogenetic group belong to the

order *Pseudomonadales*, except for group 5, whose reference strain is *Agrobacterium*

tumefaciens. TssB is commonly used to group and classify the different T6SSs among

the five phylogenetic groups, since this highly conserved protein conforms the

contractile sheath of the T6SS in all microorganisms analyzed so far (Bernal *et al*.,

2017; Cascales & Cambillau, 2012). In case of the USDA257, according to its TssB

sequence, the T6SS belongs to the group 3, which is one of the most represented

phylogenetic groups among the *Rhizobiales* order. An in deep analysis indicates that the

genome of USDA257 contains the thirteen genes coding for all structural proteins

required for a functional T6SS in one chromosomal cluster. Besides to these T6SS core

proteins, *in silico* analysis points out that USDA257 synthetizes two accessory proteins

of the membrane complex, TagL and TagM. In addition, most T6SSs described so far

are controlled by regulatory proteins (Lin *et al*., 2018; Mougous *et al*., 2007). A well-

studied example is the kinase-phosphatase PpkA-PppA couple, which mediates the

system formation through a phosphorylation cascade that ends with the structural

components assembly. In the case of the USDA257 cluster, we found not only genes

coding for PpkA-PppA regulatory proteins but also those encoding the TagF and Fha

proteins, which also interact with the system and regulate its activity at post-

transcriptional level (Lin *et al*., 2018; Mougous *et al*., 2007). Genes coding effector

proteins are often found inside of the T6SS cluster, specifically in the *vgr* gene

neighborhood (Durand *et al*., 2014). In USDA257, both *vgrG* and *paar* (*tsre1*) genes

encode for evolved proteins. The C-terminal effector domain of VgrG is structurally

similar than a polysaccharide lyase-type enzime, which points out that this USDA257

effector could bind to cellulose and polygalacturonic acid, being involved in the

attachment to vegetal cell wall. Tsre2 shows structural homology -according to Phyre[2]-

with a glycoside hydrolase enzyme with a xyloglucan binding domain able to hydrolyze

the plant cell wall (Arnal *et al*., 2019), but also -using Foldseek- with the DNase toxin

Tse7 from *P. aeruginosa* PAO1 (Pissaridou *et al*., 2018). This tool also predicts that the

Tsre1 protein possesses structural similarity to the antitoxin protein Tse7, further

supporting their potential role as a toxin-antitoxin system. Comparisons of this effector

across several rhizobial strains indicate that, despite the sequence dissimilarity among

the three different versions of the effector, Tsre2 proteins consistently exhibit structural

homology with the same two enzymes. This suggests that, regardless of the specific

function of this protein, it is likely to be conserved among rhizobia. Moreover, it cannot

ruled out that Tsre2 could be a moonlighting effector, a class of multifunctional protein

capable of performing more than one distinct biological function, often depending on

factors like its location within the cell, interaction with different molecules, or changes

in environmental conditions (Jeffery, 1999). It has been recently proposed that

moonlighting proteins may be particularly important when the symbiotic interaction

becomes more intimate, that is, during the development of the symbiosome (Ma *et al.*,

2024).

Overall, the exhaustive *in silico* analysis performed in this work points to a functional

T6SS of *S. fredii* USDA257, which might be involved in the interaction with host plant

cells and/or with bacteria present in the plant rhizosphere. But is indeed this T6SS fully

functional and involved in the direct or indirect optimization of the nodulation process?

According to our results, the USDA257 T6SS genes are highly expressed within legume

nodules and in poor media at exponential phase, but not in the presence of *nod*-gene

inducing flavonoids. Findings described for *R. etli* T6SS are largely similar, since this

protein secretion system is fully functional in yeast mannitol medium at high optical

densities of growth and in the bean nodules (Salinero-Lanzarote *et al.*, 2019). In this

bacterium, the T6SS exerts a positive effect in the symbiotic process with its natural

legume, whereas information about its role on bacterial competition is missing. The

involvement on the symbiotic process of the USDA257 T6SS has been confirmed in

nodulation and competitiveness for nodule occupancy assays, where the absence of this

protein secretion system reduces the formation of nodules and their relative nodule

colonization in soybean. In summary, *in vivo* assays strenghthen, at least in some

extend, what emerged from *in silico* analysis, i.e. this protein secretion system is fully

functional playing a positive effect on the symbiosis stablished between *S. fredii*

USDA257 and *G. max* cv Pekin, its natural host.

So far, the involvement of rhizobial T6SS on interbacterial competition seems to be

incompatible for its role on symbiosis and *vicerversa*, since both abilities have never

been described at the same time in the same rhizobial strain (Bladergroen *et al.*, 2003;

de Campos *et al.*, 2017; Lin *et al.*, 2018; Salinero-Lanzarote *et al.*, 2019; Tighilt *et al.*,

2022). Competition assays performed against various rhizobacteria under *in vitro* T6SS-

inducing conditions indicate that this protein secretion system does not confer killing

ability to USDA257. However, this result disagrees with part of the *in silico* analysis of

Tsre2 effector, since this protein seems to share structurally homology with a DNase-

type toxin (Pissaridou *et al.*, 2018). Perhaps the ability to kill other competing

rhizobacteria occurs in later stages of symbiosis, during root colonization and infection.

This could explain why, under controlled laboratory conditions, we did not detect the

toxic effect mediated by Tsre2.

The ecological success of rhizobia is based on their great capacity to adapt, not only to

environmental changes as free-living soil bacteria, but also to the challenges they face

during root colonization, invasion through the infection tube, and their establishment as

mature nitrogen-fixing bacteroids in nodule cells. So far, studies have primarily focused

on the characterization of nodulation factors, surface polysaccharides, and the T3SS.

However, our results point to a new level of specificity mediated by T6SS effectors,

which optimize the symbiotic efficiency by enhancing rhizobial competitiveness during

nodulation. Despite possible functions have been suggested for some of these effector

proteins, future work is needed to elucidate their functioning and to understand the

specific role they play in rhizosphere and symbiosis.

Supplementary Materials: Figure S1: Southern blot hybridization with probes of *tssA*

genes. Figure S2: Multiple alignment of the TssM, TssL, TagM and TagL sequences.

Figure S3-5: Multiple alignment of the Tsre1 sequences. Figure S6-8: Multiple

alignment of the Tsre2 sequences. Figure S9: Induction of *ppkA* promoter in the

presence of *nod* gene inducing-flavonoids. Table S1. Bacterial strains and plasmid used

in this study. Table S2. DNA oligonucleotide primers used in this study. Table S3.

Nitrogen-fixing and related bacteria harboring a T6SS.

Acknowledgments and Funding: This work was funded with Projects PID2020-118279RA-I00 and PID2019-107634RB-I00 funded by MICIU/AEI/10.13039/501100011033.

Author Contributions: Conceptualization, F.P-M., P.B., and I.J-G.; Methodology, P.R-P., I.J-G., and P.B.; Validation, P.R-P., A.S-R., C.C., J.G., N.M-C., F.P-M, F.O., and I.J-G.; Formal Analysis, P.R-P., F.P-M., P.B. and I.J-G.; Investigation, P.R-P., A.S-R., C.C., N.M-C., F.O., F.P-M, and I.J-G.; Writing – Original Draft Preparation, F.P-M.; Writing – Review & Editing, P.B. and I.J-G.; Visualization, P.R-P., A.S-R., F.P-M., and P.B.; Project Administration, I. J-G, P. B., and F.P-M.; Funding Acquisition, P. B., and F.P-M.; Supervision: I. J-G., P. B., J.G., and F.P-M.

Conflicts of Interest Statement: The authors declare no conflict of interest.

Data Availability Statement: The data are available in the manuscript.

References

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., … & Jumper, J.M. (2024). Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature, 630*, 493–500. https://doi.org/10.1038/s41586-024-07487-w

Almagro Armenteros, J. J., Tsirigos, K. D., Sønderby, C. K., Petersen, T. N., Winther, O., Brunak, S., von Heijne, G., & Nielsen, H. (2019). SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nature Biotechnology*, *37*(4). https://doi.org/10.1038/s41587-019-0036-z

Arnal, G., Stogios, P. J., Asohan, J., Attia, M. A., Skarina, T., Viborg, A. H., ... & Brumer, H. (2019). Substrate specificity, regiospecificity, and processivity in

glycoside hydrolase family 74. *Journal of Biological Chemistry*, *294*(36).
https://doi.org/10.1074/jbc.RA119.009861

Artimo, P., Jonnalagedda, M., Arnold, K., Baratin, D., Csardi, G., de Castro, E.,
Duvaud, S., Flegel, V., Fortier, A., Gasteiger, E., Grosdidier, A., Hernandez, C.,
Ioannidis, V., Kuznetsov, D., Liechti, R., Moretti, S., Mostaguir, K., Redaschi, N.,
Rossier, G., … Stockinger, H. (2012). ExPASy: SIB bioinformatics resource
portal. *Nucleic Acids Research*, *40*(W1). https://doi.org/10.1093/nar/gks400

Aschtgen, M. S., Thomas, M. S., & Cascales, E. (2010). Anchoring the type VI
secretion system to the peptidoglycan: TssL, TagL, TagP... what else? *Virulence*,
1(6). https://doi.org/10.4161/viru.1.6.13732

Beringer J. E. (1974). R Factor Transfer in *Rhizobium leguminosarum. Microbiology*,
84(1). https://doi.org/10.1099/00221287-84-1-188

Bernal, P., Allsopp, L. P., Filloux, A., & Llamas, M. A. (2017). The *Pseudomonas*
putida T6SS is a plant warden against phytopathogens. *ISME Journal*, *11*(4), 972–
987. https://doi.org/10.1038/ismej.2016.169

Bernal, P., Llamas, M. A., & Filloux, A. (2018). Type VI secretion systems in
plant☐associated bacteria. *Environmental microbiology*, *20*(1).
https://doi.org/10.1111/1462-2920.13956

Bernal, P., Furniss, R. C. D., Fecht, S., Leung, R. C. Y., Spiga, L., Mavridou, D. A. I.,
& Filloux, A. (2021). A novel stabilization mechanism for the type VI secretion
system sheath. *Proceedings of the National Academy of Sciences*, *118*(7).
https://doi.org/10.1073/pnas.2008500118

Bertani, G. (1951). Studies on lysogenesis I. *Journal of Bacteriology*, *62*(3).
https://doi.org/10.1128/jb.62.3.293-300.1951

Bladergroen, M. R., Badelt, K., & Spaink, H. P. (2003). Infection-Blocking Genes of a
Symbiotic *Rhizobium leguminosarum* Strain That Are Involved in Temperature-
Dependent Protein Secretion. *Molecular Plant-Microbe Interactions®*, *16*(1).
https://doi.org/10.1094/MPMI.2003.16.1.53

Boratyn, G. M., Camacho, C., Cooper, P. S., Coulouris, G., Fong, A., Ma, N., Madden,
T. L., Matten, W. T., McGinnis, S. D., Merezhuk, Y., Raytselis, Y., Sayers, E. W.,
Tao, T., Ye, J., & Zaretskaya, I. (2013). BLAST: a more efficient report with
usability improvements. *Nucleic Acids Research*, *41*(W1).
https://doi.org/10.1093/nar/gkt282

Boyer, F., Fichant, G., Berthod, J., Vandenbrouck, Y., & Attree, I. (2009). Dissecting the bacterial type VI secretion system by a genome wide *in silico* analysis: what can be learned from available microbial genomic resources? *BMC Genomics*, *10*(1). https://doi.org/10.1186/1471-2164-10-104

Buendía-Clavería, A. M., Chamber, M. & Ruiz-Sainz, J. E. (1989). A comparative study of the physiological characteristics, plasmid content and symbiotic properties of different *Rhizobium fredii* in European soils. Systematic and Applied Bacteriology *17*, 155–160. https://doi.org/10.1016/S0723-2020(89)80015-3

Cascales, E., & Cambillau, C. (2012). Structural biology of type VI secretion systems. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1592). https://doi.org/10.1098/rstb.2011.0209

Cherrak, Y., Rapisarda, C., Pellarin, R., Bouvier, G., Bardiaux, B., Allain, F., Malosse, C., Rey, M., Chamot-Rooke, J., Cascales, E., Fronzes, R., & Durand, E. (2018). Biogenesis and structure of a type VI secretion baseplate. *Nature Microbiology*, *3*(12). https://doi.org/10.1038/s41564-018-0260-1

Coulthurst, S. (2019). The Type VI secretion system: a versatile bacterial weapon. *Microbiology*, *165*(5). https://doi.org/10.1099/mic.0.000789

Coutte, L., Antoine, R., Drobecq, H., Locht, C., & Jacob-Dubuisson, F. (2001). Subtilisin-like autotransporter serves as maturation protease in a bacterial secretion pathway. *The EMBO Journal*, *20*(18). https://doi.org/10.1093/emboj/20.18.5040

de Campos, S. B., Lardi, M., Gandolfi, A., Eberl, L., & Pessi, G. (2017). Mutations in Two *Paraburkholderia phymatum* Type VI Secretion Systems Cause Reduced Fitness in Interbacterial Competition. *Frontiers in Microbiology*, *8*. https://doi.org/10.3389/fmicb.2017.02473

de Maagd, R. A., Wijffelman, C. A., Pees, E., & Lugtenberg, B. J. (1988). Detection and subcellular localization of two Sym plasmid-dependent proteins of *Rhizobium leguminosarum* biovar viciae. *Journal of Bacteriology*, *170*(9). https://doi.org/10.1128/jb.170.9.4424-4427.1988

de Sousa, B.F.S., Domingo-Serrano, L., Salinero-Lanzarote, A., Palacios, J.M., & Rey, L. (2023). The T6SS-dependent effector Re78 of *Rhizobium etli* Mim1 benefits Bacterial Competition. *Biology*, 12(678). https://doi.org/10.3390/biology12050678

Durand, E., Cambillau, C., Cascales, E., & Journet, L. (2014). VgrG, Tae, Tle, and
beyond: the versatile arsenal of Type VI secretion effectors. *Trends in*
Microbiology, *22*(9). https://doi.org/10.1016/j.tim.2014.06.004

Figurski, D. H., & Helinski, D. R. (1979). Replication of an origin-containing derivative
of plasmid RK2 dependent on a plasmid function provided in trans. *Proceedings of*
the National Academy of Sciences, *76*(4). https://doi.org/10.1073/pnas.76.4.1648

Finn, R. D., Coggill, P., Eberhardt, R. Y., Eddy, S. R., Mistry, J., Mitchell, A. L., Potter,
S. C., Punta, M., Qureshi, M., Sangrador-Vegas, A., Salazar, G. A., Tate, J., &
Bateman, A. (2016). The Pfam protein families database: towards a more
sustainable future. *Nucleic Acids Research*, *44*(D1).
https://doi.org/10.1093/nar/gkv1344

Hachani, A., Wood, T. E., & Filloux, A. (2016). Type VI secretion and anti-host
effectors. *Current Opinion in Microbiology*, *29*.
https://doi.org/10.1016/j.mib.2015.11.006

Ho, B. T., Dong, T. G., & Mekalanos, J. J. (2014). A view to a kill: the bacterial type VI
secretion system. *Cell Host & Microbe*, *15*(1).
https://doi.org/10.1016/j.chom.2013.11.008

Hsu, F., Schwarz, S., & Mougous, J. D. (2009). TagR promotes PpkA☐catalysed type
VI secretion activation in *Pseudomonas aeruginosa*. *Molecular Microbiology*,
72(5). https://doi.org/10.1111/j.1365-2958.2009.06701.x

Jeffery, C. J. (1999). Moonlighting proteins. *Trends in Biochemical Sciences*, *24*(1), 8-
https://doi.org/11. 10.1016/S0968-0004(98)01335-8

Jiménez-Guerrero, I., Acosta-Jurado, S., del Cerro, P., Navarro-Gómez, P., López-
Baena, F. J., Ollero, F. J., Vinardell, J. M., & Pérez-Montaño, F. (2018).
Transcriptomic studies of the effect of nod gene-inducing molecules in rhizobia:
Different weapons, one purpose. *Genes 9*(1). https://doi.org/10.3390/genes9010001

Jiménez-Guerrero, I., Medina, C., Vinardell, J. M., Ollero, F. J., & López-Baena, F. J.
(2022). The Rhizobial Type 3 Secretion System: The Dr. Jekyll and Mr. Hyde in
the Rhizobium–Legume Symbiosis. *International Journal of Molecular Sciences*,
23(19), 11089. https://doi.org/10.3390/ijms231911089

Jiménez☐Guerrero, I., Moreno☐De Castro, N., & Pérez☐Montaño, F. (2021). One door
closes, another opens: when nodulation impairment with natural hosts extends

rhizobial host□range. *Environmental Microbiology*, *23*(4), 1837-1841. https://doi.org/10.1111/1462-2920.15353

Jiménez-Guerrero, I., Pérez-Montaño, F., Medina, C., Ollero, F. J., & López-Baena, F. J. (2015). NopC is a rhizobium-specific type 3 secretion system effector secreted by *Sinorhizobium* (*ensifer*) *fredii* HH103. *PLoS ONE*, *10*(11). https://doi.org/10.1371/journal.pone.0142866

Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N., & Sternberg, M. J. E. (2015). The Phyre2 web portal for protein modeling, prediction and analysis. *Nature Protocols*, *10*(6). https://doi.org/10.1038/nprot.2015.053

Kovach, M.E., Elzer, P.H., Hill, D.S., Robertson, G.T., Farris, M.A., Roop, R.M., Peterson, K.M. (1995). Four new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene*, *166*(1):175-6. https://doi.org/10.1016/0378-1119(95)00584-1.

Krogh, A., Larsson, B., von Heijne, G., & Sonnhammer, E. L. L. (2001). Predicting transmembrane protein topology with a hidden markov model: application to complete genomes11Edited by F. Cohen. *Journal of Molecular Biology*, *305*(3). https://doi.org/10.1006/jmbi.2000.4315

Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*, *33*(7). https://doi.org/10.1093/molbev/msw054

Lamrabet, Y., Bellogín, R. A., Cubo, T., Espuny, R., Gil, A., Krishnan, H. B., Megias, M., Ollero, F. J., Pueppke, S. G., Ruiz-Sainz, J. E., Spaink, H. P., Tejero-Mateo, P., Thomas-Oates, J., & Vinardell, J. M. (1999). Mutation in GDP-Fucose Synthesis Genes of *Sinorhizobium fredii* Alters Nod Factors and Significantly Decreases Competitiveness to Nodulate Soybeans. *Molecular Plant-Microbe Interactions®*, *12*(3). https://doi.org/10.1094/MPMI.1999.12.3.207

Le, S. Q., Lartillot, N., & Gascuel, O. (2008). Phylogenetic mixture models for proteins. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *363*(1512). https://doi.org/10.1098/rstb.2008.0180

Lessard, I. A., & Walsh, C. T. (1999). VanX, a bacterial D-alanyl-D-alanine dipeptidase: resistance, immunity, or survival function? *Proceedings of the National Academy of Sciences*, *96*(20). https://doi.org/10.1073/pnas.96.20.1102

Letunic, I., & Bork, P. (2016). Interactive tree of life (iTOL) v3: an online tool for the
display and annotation of phylogenetic and other trees. *Nucleic Acids Research*,
44(W1). https://doi.org/10.1093/nar/gkw290

Letunic, I., Doerks, T., & Bork, P. (2015). SMART: recent updates, new developments
and   status   in   2015.   *Nucleic   Acids   Research*,   *43*(D1).
https://doi.org/10.1093/nar/gku949

Lien, Y.-W., & Lai, E.-M. (2017). Type VI Secretion Effectors: Methodologies and
Biology.   *Frontiers   in   Cellular   and   Infection   Microbiology*,   *7*.
https://doi.org/10.3389/fcimb.2017.00254

Lin, H.-H., Huang, H.-M., Yu, M., Lai, E.-M., Chien, H.-L., & Liu, C.-T. (2018).
Functional Exploration of the Bacterial Type VI Secretion System in Mutualism:
Azorhizobium caulinodans ORS571– *Sesbania rostrata* as a Research Model.
Molecular Plant-Microbe Interactions®, *31*(8). https://doi.org/10.1094/MPMI-01-
18-0026-R

Lin, J.-S., Pissaridou, P., Wu, H.-H., Tsai, M.-D., Filloux, A., & Lai, E.-M. (2018).
TagF-mediated repression of bacterial type VI secretion systems involves a direct
interaction with the cytoplasmic protein Fha. *Journal of Biological Chemistry*,
293(23). https://doi.org/10.1074/jbc.RA117.001618

López-Baena, F. J., Monreal, J. A., Pérez-Montaño, F., Guasch-Vidal, B., Bellogín, R.
A., Vinardell, J. M., & Ollero, F. J. (2009). The Absence of Nops Secretion in
Sinorhizobium fredii HH103 Increases *GmPR1* Expression in Williams Soybean.
Molecular Plant-Microbe Interactions®, *22*(11). https://doi.org/10.1094/MPMI-
22-11-1445

López-Baena, F., Ruiz-Sainz, J., Rodríguez-Carvajal, M., & Vinardell, J. (2016).
Bacterial Molecular Signals in the *Sinorhizobium fredii*-Soybean Symbiosis.
International   Journal   of   Molecular   Sciences,   *17*(5).
https://doi.org/10.3390/ijms17050755

Ma, C., Zhang, X., Bao, X., & Zhu, X. (2024). In the symbiosome: moonlighting
functions   across   kingdoms.   *New   Crops*,   100015.
https://doi.org/10.1016/j.ncrops.2024.100015

Martinelli, F., Ollero, F. J., Giovino, A., Perrone, A., Bekki, A., Sikora, S., el Nabbout,
R., Bouhadida, M., Yucel, D., Bazzicalupo, M., Mengoni, A., & Pérez-Montaño, F.
(2020). Proposed research for innovative solutions for chickpeas and beans in a
climate change scenario: The Mediterranean basin. *Sustainability (Switzerland)*,
12(4). https://doi.org/10.3390/su12041315

Miller, J. H. (1972). *Experiment in molecular genetics*. Cold Spring Harbor Laboratory.

Mougous, J. D. (2006). A Virulence Locus of *Pseudomonas aeruginosa* Encodes a
Protein Secretion Apparatus. *Science*, *312*(5779).
https://doi.org/10.1126/science.1128393

Mougous, J. D., Gifford, C. A., Ramsdell, T. L., & Mekalanos, J. J. (2007). Threonine
phosphorylation post-translationally regulates protein secretion in *Pseudomonas*
aeruginosa. *Nature Cell Biology*, *9*(7). https://doi.org/10.1038/ncb1605

Nelson, M. S., & Sadowsky, M. J. (2015). Secretion systems and signal exchange
between nitrogen-fixing rhizobia and legumes. *Frontiers in Plant Science*, *6*.
https://doi.org/10.3389/fpls.2015.00491

Oldroyd, G. E. D., & Downie, J. A. (2008). Coordinating Nodule Morphogenesis with
Rhizobial Infection in Legumes. *Annual Review of Plant Biology*, *59*(1).
https://doi.org/10.1146/annurev.arplant.59.032607.092839

Oldroyd, G. E. D., Murray, J. D., Poole, P. S., & Downie, J. A. (2011). The Rules of
Engagement in the Legume-Rhizobial Symbiosis. *Annual Review of Genetics*,
45(1). https://doi.org/10.1146/annurev-genet-110410-132549

Pérez-Montaño, F., Alías-Villegas, C., Bellogín, R. A., del Cerro, P., Espuny, M. R.,
Jiménez-Guerrero, I., López-Baena, F. J., Ollero, F. J., & Cubo, T. (2014). Plant
growth promotion in cereal and leguminous agricultural important plants: From
microorganism capacities to crop production. In *Microbiological Research* (Vol.
169, Issues 5–6, pp. 325–336). Urban und Fischer Verlag Jena.
https://doi.org/10.1016/j.micres.2013.09.011

Pérez-Montaño, F., Jiménez-Guerrero, I., Acosta-Jurado, S., Navarro-Gómez, P.,
Ollero, F. J., Ruiz-Sainz, J. E., López-Baena, F. J., & Vinardell, J. M. (2016). A
transcriptomic analysis of the effect of genistein on *Sinorhizobium fredii* HH103
reveals novel rhizobial genes putatively involved in symbiosis. *Scientific Reports*,
6(1). https://doi.org/10.1038/srep31592

Pérez-Montaño, F., Jiménez-Guerrero, I., del Cerro, P., Baena-Ropero, I., López-Baena,
F. J., Ollero, F. J., Bellogín, R., Lloret, J., & Espuny, R. (2014). The symbiotic
biofilm of *Sinorhizobium fredii* SMH12, necessary for successful colonization and
symbiosis of *Glycine max* cv Osumi, is regulated by quorum sensing systems and
inducing flavonoids via NodD1. *PLoS ONE*, *9*(8).
https://doi.org/10.1371/journal.pone.0105901

Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time
RT-PCR. *Nucleic Acids Research*, *29*(9). https://doi.org/10.1093/nar/29.9.e45

Pissaridou, P., Allsopp, L. P., Wettstadt, S., Howard, S. A., Mavridou, D. A., & Filloux,
A. (2018). The *Pseudomonas aeruginosa* T6SS-VgrG1b spike is topped by a
PAAR protein eliciting DNA damage to bacterial competitors. *Proceedings of the*
National Academy of Sciences, *115*(49). https://doi.org/10.1073/pnas.1814181115

Pueppke, S. G., & Broughton, W. J. (1999). *Rhizobium* sp. Strain NGR234 and *R. fredii*
USDA257 Share Exceptionally Broad, Nested Host Ranges. *Molecular Plant-*
Microbe Interactions®, *12*(4). https://doi.org/10.1094/MPMI.1999.12.4.293

Pukatzki, S., McAuley, S. B., & Miyata, S. T. (2009). The type VI secretion system:
translocation of effectors and effector-domains. In *Current Opinion in*
Microbiology (Vol. 12, Issue 1, pp. 11–17). Elsevier Ltd.
https://doi.org/10.1016/j.mib.2008.11.010

Redondo-Nieto, M., Barret, M., Morrissey, J., Germaine, K., Martínez-Granero, F.,
Barahona, E., ... & Rivilla, R. (2013). Genome sequence reveals that *Pseudomonas*
fluorescens F113 possesses a large and diverse array of systems for rhizosphere
function and host interaction. *BMC Genomics*, *14*(1). https://doi.org/10.1186/1471-
2164-14-54

Robertsen, B.K.; Aiman, P.; Darwill, A.G.; Mcneil, M.; Albersheim, P.(1981). The
structure of acidic extracellular polysaccharides secreted *by Rhizobium*
leguminosarum and *Rhizobium trifolii*. *Plant Physiol*y, *67*, 389–400.

Salinero-Lanzarote, A., Pacheco-Moreno, A., Domingo-Serrano, L., Durán, D.,
Ormeño-Orrillo, E., Martínez-Romero, E., Albareda, M., Palacios, J. M., & Rey, L.
(2019). The Type VI secretion system of *Rhizobium etli* Mim1 has a positive effect
in symbiosis. *FEMS Microbiology Ecology*, *95*(5).
https://doi.org/10.1093/femsec/fiz054

Sambrook, J., Fritsch, E. F., & Maniatis, T. (1989). *Molecular Cloning. A Laboratory*
Manual. Cold Spring Harbor Laboratory Press.

Santin, Y. G., Doan, T., Lebrun, R., Espinosa, L., Journet, L., & Cascales, E. (2018). In
vivo TssA proximity labelling during type VI secretion biogenesis reveals TagA as
a protein that stops and holds the sheath. *Nature Microbiology*, *3*(11).
https://doi.org/10.1038/s41564-018-0234-3

Schäfer, A., Tauch, A., Jäger, W., Kalinowski, J., Thierbach, G., & Pühler, A. (1994).
Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli*
plasmids pK18 and pK19: selection of defined deletions in the chromosome of
Corynebacterium glutamicum. Gene, *145*(1). https://doi.org/10.1016/0378-
1119(94)90324-7

Samal B, Chatterjee S. (2021). Bacterial quorum sensing facilitates *Xanthomonas*
campesteris pv. campestris invasion of host tissue to maximize disease symptoms.
Journal   of   Experimental   Botany,   *72*(18):6524-6543.
https://doi.org/10.1093/jxb/erab211.

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R.,
McWilliam, H., Remmert, M., Söding, J., Thompson, J. D., & Higgins, D. G.
(2011). Fast, scalable generation of high□quality protein multiple sequence
alignments using Clustal Omega. *Molecular Systems Biology*, *7*(1).
https://doi.org/10.1038/msb.2011.75

Silverman, J. M., Brunet, Y. R., Cascales, E., & Mougous, J. D. (2012). Structure and
Regulation of the Type VI Secretion System. *Annual Review of Microbiology*,
66(1). https://doi.org/10.1146/annurev-micro-121809-151619

Simon, R. (1984). High frequency mobilization of gram-negative bacterial replicons by
the in vitro constructed Tn5-Mob transposon. *Molecular and General Genetics*
MGG, *196*(3). https://doi.org/10.1007/BF00436188

Spaink, H. P., Okker, R. J. H., Wijffelman, C. A., Pees, E., & Lugtenberg, B. J. J.
(1987). Promoters in the nodulation region of the *Rhizobium leguminosarum* Sym
plasmid   pRL1JI.   *Plant   Molecular   Biology*,   *9*(1).
https://doi.org/10.1007/BF00017984

Staehelin, C., & Krishnan, H. B. (2015). Nodulation outer proteins: double-edged
swords   of   symbiotic   rhizobia.   *Biochemical   Journal*,   *470*(3).
https://doi.org/10.1042/BJ20150518

Studier, F.W. and Moffatt, B.A. (1986). Use of bacteriophage T7 RNA polymerase to
direct selective high-level expression of cloned genes. *Journal of Molecular*
Biology, *189*: 113-130. https://doi.org/10.1016/0022-2836(86)90385-2

Sullivan, M. J., Petty, N. K., & Beatson, S. A. (2011). Easyfig: a genome comparison
visualizer. *Bioinformatics*, *27*(7). https://doi.org/10.1093/bioinformatics/btr039

Tighilt, L., Boulila, F., De Sousa, B. F. S., Giraud, E., Ruiz-Argüeso, T., Palacios, J. M., ... & Rey, L. (2022). The *Bradyrhizobium* Sp. LmicA16 Type VI Secretion System Is Required for Efficient Nodulation of *Lupinus* Spp. *Microbial Ecology, 84*(3). https://doi.org/10.1007/s00248-021-01892-8

Van Kempen, M., Kim, S. S., Tumescheit, C., Mirdita, M., Lee, J., Gilchrist, C. L., ... & Steinegger, M. (2024). Fast and accurate protein structure search with Foldseek. *Nature biotechnology*, *42*(2), 243-246. https://doi.org/10.1038/s41587-023-01773-0

Vincent, J. M. (1970). The modified Fåhraeus slide technique. In *A Manual for the Practical Study of Root Nodule Bacteria* (pp. 144–145). Blackwell Scientific Publications.

Walker, T. S., Bais, H. P., Grotewold, E., & Vivanco, J. M. (2003). Root Exudation and Rhizosphere Biology. *Plant Physiology*, *132*(1). https://doi.org/10.1104/pp.102.019661

Yu, N. Y., Wagner, J. R., Laird, M. R., Melli, G., Rey, S., Lo, R., Dao, P., Sahinalp, S. C., Ester, M., Foster, L. J., & Brinkman, F. S. L. (2010). PSORTb 3.0: improved protein subcellular localization prediction with refined localization subcategories and predictive capabilities for all prokaryotes. *Bioinformatics*, *26*(13). https://doi.org/10.1093/bioinformatics/btq249

Zaat, S. A., Wijffelman, C. A., Spaink, H. P., van Brussel, A. A., Okker, R. J., & Lugtenberg, B. J. (1987). Induction of the *nodA* promoter of *Rhizobium leguminosarum* Sym plasmid pRL1JI by plant flavanones and flavones. *Journal of Bacteriology*, *169*(1). https://doi.org/10.1128/jb.169.1.198-204.1987

Figures

Figure 1. Phylogenetic tree of T6SSs of nitrogen-fixing bacteria. Maximum likelihood tree with 500 bootstrap replicates were built with Mega 7 software for the core component protein TssB. T6SS cluster nomenclature is used to show the main phylogenetic clusters (Boyer *et al.*, 2009). Branches with black circles indicate a confidence level higher than 0.75. Five groups are distinguished: group 1 (red), group 2 (green), group 3 (blue), group 4 (purple) and group 5 (orange). Bacterium of the genus *Bacteroides* (brown) represents the tree root. Red star indicates the position of *S. fredii* USDA257 T6SS.

Figure 2. The T6SS of *S. fredii* USDA257. (A) Organization of the T6SS cluster of USDA257. Color codes: Genes that encode the components of membrane complex (*tssMLJ*) and accessory proteins (*tagLYM*) (veige). Genes coding for components of the base plate (*tssKGFE*) (purple). *tssA* (pink). Genes *tssB* and *tssC* (contractile sheath) (light green). The *hcp* gene, that encodes the protein forming the inner tube (dark blue). Genes that encode regulatory proteins (*tagF*, *fha*, *ppkA* and *pppA*) (light blue). *clpV*, that codes for ATPase (red). Genes coding for possible effectors of the system (C-term domain of *vgrG*, *tsre1*, *tsre2* and *tsre3*) (grey). Genes that encode a possible adapter (*tsar*) (dark green). Gene of unknown function (*usda_c28460*) (orange). The *vgrG* and *OB-fold* genes (brown). *paar* domain (brown). (B) Schematic representation of the structural protein domains of TssM, TssL, TssJ and the accessory proteins TagM and TagN. DotU cytosolic domain (dark green), transmembrane domain (light blue), IcmF structural domain (dark blue), OmpA peptidoglycan-binding domain (red), and signal peptide (light green). OM: outer membrane, IM: inner membrane, PG: peptidoglycan layer.

Sinorhizobium fredii USDA257
(100%)

Sinorhizobium americanum CCGM7
(89.52%)

Sinorhizobium americanum CFNEI 73
(89.28%)

Rhizobium gallicum bv. *gallicum* R602
(83.98%)

Rhizobium etli 8C-3
(82.93%)

Rhizobium phaseoli N841
(79.07%)

Rhizobium phaseoli N261
(79.04%)

Rhizobium phaseoli N671
(79.04%)

Rhizobium phaseoli N771
(79.04%)

Rhizobium phaseoli R611
(79.04%)

Rhizobium phaseoli R620
(79.04%)

Rhizobium phaseoli R630
(79.04%)

Rhizobium phaseoli R650
(79.04%)

Rhizobium phaseoli R723
(79.04%)

Rhizobium phaseoli N161
(79.00%)

Rhizobium phaseoli N831
(79.01%)

Rhizobium phaseoli N931
(79.01%)

Rhizobium phaseoli Brasil 5
(78.88%)

Rhizobium phaseoli R744
(78.94%)

vrgG   *tsre2*   *tsre1*

1 Kbp   100%

66%

Figure 3. Genome sequence alignment of the *vgrG* region demonstrating the divergence of *tsre1* and *tsre2* in eighteen rhizobial strains using Easyfig. Clusters are sorted by homology degree respect to the VgrG protein from *S. fredii* USDA257 (Blastp E value). In parenthesis is indicated the percentage of identity for each VrgG. Putative protein domains encoded for each gene of the *vgrG* region are boxed above alignments.

Figure 4. Neighbor-joining tree of rhizobia T6SS putative effectors. Bootstrap values ≥40 is indicated for each node. The cluster analysis to group the strains by Tsre1 (A) or Tsre2 (B) sequence similarities was done using the program CLUSTALW in the MEGA7 software package with the algorithm neighbor-joining method. Tree robustness was assessed by bootstrap resampling (1000 replicates each). Asterisks indicate representative Tsre1 and Tsre2 versions for each main branch.

Figure 5. Expression of the *S. fredii* USDA257 *ppkA* gene. (A) Fold-change values of the β-galactosidase activity of USDA257 carrying a pMP220 plasmid containing the *ppkA* promoter region fused to the *lacZ* gene respect to those values obtained by the strain carrying the empty pMP220 plasmid. Assayed conditions were TY, YM10, YM3, MM10 and MM3 media 30 hours after inoculation. Data followed by the same letter are not significantly different at the level of α = 5% (One-Way ANOVA with multiple comparisons, P<0.05). (B) Quantitative RT-PCR analysis of the expression of the *ppkA* gene of USDA257 grown in the TY, YM10, YM3, MM10 and MM3 media 48 hours after inoculation. Expression data shown are the mean (± standard deviation of the mean) for three biological replicates. Data followed by the same letter are not significantly different at the level of α = 5% (One-Way ANOVA with multiple comparisons, P<0.05). (C) Fold-change values of the β-galactosidase activity of USDA257 carrying a pMP220 plasmid containing the *ppkA* promoter region fused to the *lacZ* gene respect to those values obtained by the strain carrying the empty pMP220 plasmid. The assayed condition was MM3 medium at different hours after inoculation. Data followed by the same letter are not significantly different at the level of α = 5% (One-Way ANOVA with multiple comparisons, P<0.05).

Figure 6. Functionality of the *S. fredii* USDA257 T6SS. (A) Production and secretion of Hcp in the USDA257 wild-type strain grown for 48 h in rich, standard and minimal media: TY, YM, and MM, repectively. YM3 and MM3 represent derivatives of their corresponding media with reducedi carbon source content,. (B) Production and secretion of Hcp in the USDA257 wild-type and the *tssA* mutant strains grown in MM3. For both panels, the Hcp protein was detected by Western blot analysis using a specific anti-Hcp antibody. Detection of the β-subunit of the RNA polymerase (β-RNApol) was used as control. "Whole cells" represents the intracellular protein fraction, wh The position of the molecular size marker (in kDa) is indicated.

Figure 7. Plant responses to inoculation with *S. fredii* USDA257 strains. (A) Plant responses to inoculation of *G. max* cv Pekin with different *S. fredii* USDA257 strains. Plants were evaluated 6 weeks after inoculation. NI: non-inoculated plants. Data followed by the same letter are not significantly different at the level of α = 5% (One-Way ANOVA with multiple comparisons, P<0.05). (B) Competition for nodulation assays of different *S. fredii* USDA257 strains. Percentages of nodule occupancy were evaluated by co-inoculation at different ratios of both strains at 30 dpi. Light grey: USDA257 wild-type strain, dark grey: USDA257 *tssA* mutant strain. (C) Confocal microscopy of *L. burttii* 30 dai-nodules infected by the *S. fredii* USDA257 carrying a dual fluorescent reporter plasmid with (pDUAL with P_{ppkA}) or without (pDUAL w/o P_{ppkA}) the *ppkA* promoter region upstream of the *rfp* gene.

Tables

Table 1. Genes of the T6SS cluster of *S. fredii* USDA257.

Locus name	Protein name	Conserved domains[a]	Celullar localization prediction
USDA257_c28440	VgrG$_e$	COG3501 (VgrG) + COG3210 (FhaB)/ pfam06715 (Gp5_C)	C
USDA257_c28450	VgrG dom	COG3501 (VgrG)/ TIGR01646 (Vrg_GE)	D
USDA257_c28460		TIGR02270	C
USDA257_c28470	Tre2	Glycosyl hydrolase GH74	C
USDA257_c28480	Tre1	PAAR-pfam13665 (DUF4150)	C
USDA257_c28490	FabB	COG0304 (FabB)	MI
USDA257_c28500	Tar	DUF2169	C
USDA257_c28510	TagF	COG3913 (SciT)	D
USDA257_c28520	TssM	COG3523 (IcmF)	MI
USDA257_c28530	TssL	COG3455 (DotU) + Pfam00691 (OmpA)	ME
USDA257_c28540	TssK	COG3522	C
USDA257_c28550	TssJ	COG3521	MI
USDA257_c28560	Fha	COG3456	C
USDA257_c28570	TssG	COG3520	C
USDA257_c28580	TssF	COG3519	C
USDA257_c28590	TssE	COG3518	C
USDA257_c28600	Hcp	COG3157 (Hcp)	C
USDA257_c28610	TssC	COG3517	C
USDA257_c28620	TssB	COG3516	C
USDA257_c28630	TssA	COG3515 + TIGR03363	C
USDA257_c28640	PpkA	COG0515 (SPS1)	P + PC
USDA257_c28650	ClpV	COG0542 (ClpA) + pfam07724 (AAA_2)	C
USDA257_c28660	TagL	COG2885 (OmpA)	ME
USDA257_c28670	PppA	COG0631 (PTC) + TIGR02865 (spore_II_E)	C

USDA257_c28680	TagY	Pfam04102 (SlyX) + pfam13539 (peptidasa_M15_4)	D
USDA257_c28690	TagM	COG3523 (IcmF) + COG3455 (DotU)	D

OM: outer membrane, IM: inner membrane, C: cytoplasmic, P: periplasm, PC: cell wall,

D: unknown, aa: amino acid.

a: conserved domains according to COG / pfam / TIGR / Phyre[2].

Table 2. Best predictions for the structural homology from representative rhizobial versions coded by same loci than *tsre1* and *tsre2* genes of *S. fredii* USDA257.

Rhizobial strain	Tsre1-like	Tsre2-like
Sinorhizobium fredii USDA257	P: PAAR (25-125 a.a.) F: DNAse toxin Tse7 (3-321 a.a.) Size: 324 a.a.	P: Glycosyl hydrolase GH74 (1-340 a.a.) F: Immune protein Tsi7 (2-340 a.a.) Size: 351 a.a.
Rhizobium phaseoli N841	P: PAAR (26-126 a.a.). F: DNAse toxin Tse7 (1-327 a.a.) Size: 328 a.a.	P: Glycosyl hydrolase GH74 (106-301 a.a.) F: Immune protein Tsi7 (17-228 a.a.) Size: 309 a.a.
Rhizobium phaseoli R620	P: PAAR (26-126 a.a.). F: Not available. Size: 330 a.a.	P: Glycosyl hydrolase GH74 (98-302 a.a.) F: Not available. Size: 309 a.a.

Structural homologies were determined by Phyre[2] (P:) software and (F:) Foldseek tool. In parenthesis are indicated range of aligned amino acids.

Table 3. Average of CFUs obtained in competition assays after 0, 24 or 48 hours of interaction in T6SS-inducing media.

Strains	0 hours	24 hours	48 hours
P. carotovorum (vs WT)	1.8×10^7 $\pm 4.5 \times 10^6$	1.1×10^6 $\pm 1.0 \times 10^5$	9.5×10^4 $\pm 5.0 \times 10^3$
P. carotovorum (vs *tssA*)	1.6×10^7 $\pm 5.0 \times 10^6$	1.0×10^6 $\pm 3.0 \times 10^5$	1.1×10^5 $\pm 3.0 \times 10^4$
WT (vs *P. carotovorum*)	2.0×10^7 $\pm 7.0 \times 10^6$	7.5×10^5 $\pm 1.5 \times 10^5$	1.8×10^3 $\pm 9.0 \times 10^2$
tssA (vs *P. carotovorum*)	2.3×10^7 $\pm 1.7 \times 10^7$	1.4×10^6 $\pm 4.5 \times 10^5$	2.5×10^2 $\pm 5.0 \times 10^1$
A. tumefaciens (vs WT)	5.3×10^7 $\pm 5.0 \times 10^5$	1.3×10^7 $\pm 3.0 \times 10^6$	1.4×10^7 $\pm 1.5 \times 10^6$
A. tumefaciens (vs *tssA*)	5.6×10^7 $\pm 1.5 \times 10^6$	1.3×10^7 $\pm 2.0 \times 10^6$	1.7×10^7 $\pm 5.0 \times 10^6$
WT (vs *A. tumefaciens*)	1.8×10^7 $\pm 6.5 \times 10^6$	7.0×10^5 $\pm 1.0 \times 10^5$	9.5×10^5 $\pm 5.0 \times 10^4$
tssA (vs *A. tumefaciens*)	2.7×10^7 $\pm 1.9 \times 10^7$	1.0×10^6 $\pm 7.0 \times 10^5$	1.5×10^6 $\pm 8.0 \times 10^5$
S. fredii HH103 (vs WT)	1.8×10^7 $\pm 4.5 \times 10^6$	1.1×10^6 $\pm 1.0 \times 10^5$	9.5×10^4 $\pm 5.0 \times 10^3$
S. fredii HH103 (vs *tssA*)	1.6×10^7 $\pm 5.0 \times 10^6$	1.0×10^6 $\pm 3.0 \times 10^5$	1.1×10^5 $\pm 3.0 \times 10^4$
WT (vs *S. fredii* HH103)	2.0×10^7 $\pm 7.0 \times 10^6$	7.5×10^5 $\pm 1.5 \times 10^5$	1.8×10^3 $\pm 9.0 \times 10^2$
tssA (vs *S. fredii* HH103)	2.3×10^7 $\pm 1.7 \times 10^7$	1.4×10^6 $\pm 4.5 \times 10^5$	2.5×10^2 $\pm 5.0 \times 10^1$

*WT and *tssA* indicate *S. fredii* USDA257 wild-type and the *tssA* mutant strain used in these assays.