═ R E V I E W   C O M M O N S   M A N U S C R I P T ═

IMPORTANT:

- Manuscripts submitted to Review Commons are peer reviewed in a journal-agnostic way.
- Upon transfer of the peer reviewed preprint to a journal, the referee reports will be available in full to the handling editor.
- The identity of the referees will NOT be communicated to the authors unless the reviewers choose to sign their report.
- The identity of the referee will be confidentially disclosed to any affiliate journals to which the manuscript is transferred.

GUIDELINES:

- For reviewers: https://www.reviewcommons.org/reviewers
- For authors: https://www.reviewcommons.org/authors

CONTACT:

The Review Commons office can be contacted directly at: office@reviewcommons.org

Neuroepithelial depletion schedules cessation of neurogenesis in the *Drosophila* optic

lobes

Phuong-Khanh Nguyen[1,3], Louise Y Cheng[1,2,3,#]

1. Peter MacCallum Cancer Centre, Melbourne, VIC 3000, Australia

2. Sir Peter MacCallum Department of Oncology, The University of Melbourne, VIC 3010,

Australia

3. Department of Anatomy and Physiology, The University of Melbourne, VIC 3010, Australia

Corresponding author

Email: louise.cheng@petermac.org

Keywords: *Drosophila*, medulla neuroblast, gliogenesis, neuroepithelium

Running title: medulla neuroblast termination

Abstract

The brain is consisted of diverse neurons arising from a limited number of neural stem cells.

Drosophila neural stem cells called neuroblasts (NBs) produces specific neural lineages of

various lineage sizes depending on their location in the brain. In the *Drosophila* visual

processing centre - the optic lobes (OLs), medulla NBs derived from the neuroepithelium (NE)

give rise to neurons and glia cells of the medulla cortex. The timing and the mechanisms

responsible for the cessation of medulla NBs are so far not known. In this study, we show that

the termination of medulla NBs during pupal development is determined by the exhaustion of

the NE stem cell pool. Altering NE-NB transition during larval neurogenesis disrupts the timely

termination of medulla NBs. Medulla NBs terminate neurogenesis via a combination of cell

death, terminal symmetric division, and a switch to gliogenesis. We show that temporal

progression is not required for the termination of medulla NBs. The timing of NB cessation

can be altered through the acquisition of a glial cell fate via Glial cells missing, or through

conversion to type II NB cell fate via Tailless, or by inhibition of differentiation via Prospero

knockdown. As the *Drosophila* OL shares a similar mode of division with mammalian

neurogenesis, determining how and when these progenitors cease proliferation during

development can have important implications for mammalian brain size determination and

regulation of its overall function.

Introduction

The central nervous system (CNS) is the cognitive control centre of the body and is generated by neural stem cells (NSCs) during development. *Drosophila* NSCs, called neuroblasts (NBs), produce different neural and glial types and subtypes that contribute to the formation of the adult brain. The cellular diversity of the CNS is determined in a region-specific manner, via differential control of NB division mode and temporal identity (reviewed by Harding & White, 2018). The majority of NBs are Type I NBs located in the ventral nerve cord (VNC) and the central brain (CB). They divide asymmetrically to generate a NB and a smaller ganglion mother cell (GMC), that undergoes terminal differentiation to give rise to neurons or glial cells (Harding & White, 2018). In addition, there are eight Type II NBs located on the dorsal surface of the CB, where NBs asymmetrically divide to give rise to a NB and an intermediate neural progenitor (INP) (Bello et al., 2008; Boone & Doe, 2008; Bowman et al., 2008). INPs undergo maturation prior to additional rounds of asymmetric division, to give rise to neurons and glial cells.

The optic lobes (OLs) of the brain are the visual processing centre of the adult fly brain (Figure 1A). During neural development, neurogenesis in the OLs takes place in two proliferation centres: the inner and the outer proliferation centres, which produce different neural cell types that populate the various cortexes of the visual system (Sato *et al*, 2019; Apitz & Salecker, 2014). In the outer proliferation centre (OPC), medulla NBs are generated from a pseudostratified neuroepithelium through a differentiating "proneural wave" (Figure 1A') (Egger et al., 2007; Li et al., 2013; Yasugi et al., 2008). This proneural wave is characterised by the expression of proneural factors such as Lethal of scute (L'sc) which precedes NB generation from NE cells (Yasugi et al., 2008). During early larval stages, OPC NE cells undergo symmetric cell divisions to expand the progenitor population. By early-third instar, progression of the proneural wave is initiated in the NE cells at the medial edge, leading to their differentiation into medulla NBs. Medulla NBs sequentially express a series of temporal transcription factors (Konstantinides et al., 2022; Li et al., 2013; Zhu et al., 2022). During the early-mid temporal stages, NBs divide asymmetrically to produce two neurons, upon acquiring the terminal temporal identity, NBs switch to become glioblasts that produce glial cells (Zhu *et al*, 2022).

At the end of neurogenesis, NBs of different neural lineages are eliminated through distinct mechanisms. For instance, type I NBs in the VNC and the CB exit the cell cycle and

terminally differentiate at around 24 hr after pupal formation (APF), whereas type I

abdominal NBs are eliminated via apoptosis prior to pupation (Maurange *et al*, 2008; Bello *et*

al, 2003; Ito & Hotta, 1992). Mushroom body NBs are the last NBs to undergo termination,

and they do so through apoptosis or autophagy prior to adult eclosion (Siegrist *et al*, 2010;

Pahl *et al*, 2019). So far, the timing and mode by which medulla NBs terminate is not

known; furthermore, it is unclear whether the temporal series, which schedules NB cessation

in other NB lineages (Maurange et al., 2008; Syed et al., 2017) is also involved in the

termination of medulla NBs.

Here, we find that medulla NBs terminate between 12-18 hr APF. Unlike other NB

lineages, the timing of medulla NB cessation is mainly regulated by the depletion of the NE

pool. The exhaustion of the NE during pupal development results in the lack of the formation

of new NBs. Hence, altered NE-NB transition is sufficient to change the timing of NB

termination. Additionally, we demonstrate that medulla NBs terminate via a combination of

cell death, size-symmetric terminal differentiation and a switch to gliogenesis. While medulla

NBs undergo temporal transitions, the temporal series is not required for NB termination.

However, overexpression of Tll, Gcm and knockdown of Pros prevents timely NB cessation

during the pupal stages.

Results

Medulla NBs terminate during early pupal development.

To characterize the timing at which medulla NBs are eliminated during pupal development, we first assessed the number of NBs in the OPC using the pan-NB marker Deadpan (Dpn). We found that the number of NBs was significantly reduced between 12-18 hr APF and by 30 hr APF, all the NBs were eliminated (Figure 1B-C).

In most regions of the CNS, NB termination is associated with a reduction in cell growth and proliferation rate, scheduled by a peak of ecdysone between larval and pupal development (Homem *et al*, 2014). To test if ecdysone signalling plays a role in the termination of medulla NBs, we first looked at the expression of the ecdysone receptor (EcR) in the medulla NBs at 12 hr APF, prior to the reduction in the number of medulla NBs in the OL. Immunostainings showed that EcR is expressed at very low levels in the medulla NBs at 12 hr APF (Supplementary Figure 1A). Additionally, when EcR was knocked down via RNAi in the medulla NBs with *eyR16F10-GAL4,* which is expressed in a subset of medulla NBs and their progeny (Veen *et al*, 2023), we could not detect any persistent NBs at 24 hr APF (Supplementary Figure 1B, C, E). A similar phenotype was observed using a dominant negative form of EcR (EcR^{DN}) (Supplementary Figure 1D, E). Together, these data suggest that ecdysone signalling in the medulla NBs is not required for their cessation during pupal development.

During larval development, NBs regrow to their original size after each division empowered by aerobic glycolysis providing the NBs with energy and metabolites for rapid cell growth and proliferation (Homem *et al*, 2014; Maurange *et al*, 2008). To assess if a reduction in cell size also precedes NB elimination in the medulla, we monitored the nuclear size of NBs marked by Dpn. Between 12-18 hr APF when the majority of medulla NBs disappear, we found that there was a significant reduction in the nuclear sizes of the medulla NBs (Figure 1D-E), suggesting that decreased cell growth precedes the cessation of medulla NB. However, increasing cellular growth through the overexpression of a constitutive active form of the PI3K catalytic subunit Dp110 (dp110^{CAAX}) (Leevers et al., 1996), or the activation of the cellular growth regulator Myc (Rust *et al*, 2018) did not prolong NB persistence in the OL at 24 hr APF (Supplementary Figure 1F, G, J, data not shown).

In type I NBs of the CB and VNC, it was previously shown that NB termination is dependent on core components of the mediator complex, which promote key enzymes to increase oxidative phosphorylation (OxPhos) to slow down NB regrowth (Homem *et al*, 2014). Nevertheless, neither inhibition of OxPhos itself via the knockdown of Complex I (Ameele &

Brand, 2019), nor the knockdown of Med27 via RNAi (Homem *et al*, 2014), a component of

the Mediator complex, significantly affected medulla NB persistence at 24 hr APF

(Supplementary Figure 1H-J). As such, although cell size reduction coincides with NB

elimination during pupal development, a decrease in cell growth, OxPhos, and the ecdysone-

induced Mediator complex are not required for the termination of medulla NBs.

The depletion of the NE determines the timing of medulla NB cessation.

NBs of the medulla are continuously generated from NE cells from mid-late larval

development by the proneural wave (Egger et al., 2007; Yasugi et al., 2008). To assess if NE

depletion during pupal stages is linked to the cessation of medulla NBs, we first characterised

the volume of NE cells during pupal development. From 12 hr to 24 hr APF, the NE

gradually decreased in volume, concordant with the reduction in the number of medulla NBs

during this time window (Figure 2A-B, Figure 1B-C). This suggests that the depletion of the

progenitor NE pool may drive the cessation of medulla NBs.

To test if the depletion of the NE can change the timing of NB termination, we

manipulated Notch signalling, a negative regulator of the NE-NB transition (Egger et al.,

2010; Wang et al., 2011; Yasugi et al., 2010), where Notch signalling promotes NE self-

renewal and prevents their differentiation into NBs (Egger et al., 2010; Wang et al., 2011;

Yasugi et al., 2010). Control and *notch* RNAi clones were induced in the medulla NE at 24 hr

after larval hatching (ALH) and examined at 96 hr ALH. Control clones spanned the NE and

NB regions, demarcated by a sharp NE-NB boundary indicated by PatJ (NE) and Mira (NB)

(Figure 2D). In contrast, *notch* RNAi clones consisted mostly of only NBs, suggesting that

Notch inhibition has caused an accelerated NE-NB transition (Figure 2E) (Egger et al., 2010;

Wang et al., 2011; Yasugi et al., 2010). By 12 hr APF, we found that *notch* RNAi clones

were smaller in size and contained very few NBs compared to control (Figure 2F-H). Thus, it

is likely that an accelerated NE-NB transition caused by *notch* RNAi resulted in a premature

depletion of the NE which in turn led to the early termination of NBs. To test this further, we

overexpressed the proneural gene *lethal of scute* (*l'sc*), which has also been shown to

promote NE-NB transition during the larval stages (Yasugi et al., 2008) (Figure 2I). Similar

to *notch* RNAi clones, *l'sc* overexpression caused medulla NBs to undergo premature

termination at 12 hr APF (Figure 2J-K). Together, our data indicate that the premature

depletion of the NE results in earlier medulla NB termination.

Conversely, we found that the overexpression clones of Notch signalling via N^{ACT}

caused a delay in the NE-NB transition, resulting in clones consisting only of NE cells (and

no NBs) at 96 hr ALH (Figure 2L). We only started to observe NBs in N^{ACT} clones in the OLs

(Figure 2M, n = 8/8 lobes) at 72 hr APF, a time point where control medulla NBs have long

disappeared. These medulla NBs continued to persist into the adult OL (data not shown).

Together, our results suggest that early depletion of the NE results in the early cessation of

medulla NBs, whereas delayed formation of the NB from the NE results in late termination of

medulla NBs. Thus, the timing of the NE-NB transition determines when medulla NBs

terminate.

Medulla NBs cease divisions via cell death.

Next, we examined how medulla NBs undergo termination. First, we examined mechanisms

that are responsible for the termination of other NB lineages. Mushroom Body NBs are the

last NBs to undergo termination (Ito & Hotta, 1992). These NBs are eliminated via a

combination of apoptosis and autophagy (Siegrist *et al*, 2010). To test if medulla NBs

terminate via similar mechanisms, we first assessed cell death, where we assayed for the

expression of the apoptotic marker Caspase-1 (Dcp-1) in the medulla NBs marked by

Dpn::GFP. At 12 hr APF, we observed that some NBs expressed Dcp-1 (Figure 3A). To test

if apoptosis is required for the termination of medulla NBs, we inhibited cell death via the

overexpression of the baculovirus anti-apoptotic gene *p35* in clones as well as in medulla

NBs using *eyR16F10-GAL4*. Upon the inhibition of cell death, we observed a small number

of persistent NBs at 24 hr APF through both approaches (Figure 3B-D, Supplementary Figure

2A-B, E). However, no persistent NBs were found at 48 hr APF, suggesting that additional

factors may be involved to terminate medulla NBs (Supplementary Figure 2D, n = 5/5 lobes).

Next, we asked if autophagy is required to eliminate medulla NBs. To address this,

we inhibited autophagy via the expression of a RNAi against Atg1 in the medulla NBs with

eyR16F10-GAL4 (Xu *et al*, 2015). We found that the timing of medulla NB termination was

not significantly affected by the inhibition of autophagy (Supplementary Figure 2C, E).

Together, our data suggest that cell death, and not autophagy, makes a minor contribution to

medulla NB termination.

Medulla NBs undergo Pros-mediated termination.

In type I NBs of the VNC, the nuclear localisation of a differentiation factor called Prospero

(Pros) precedes NB termination via size-symmetric cell division (Choksi et al., 2006; Li et

al., 2013; Maurange et al., 2008). To assess if medulla NBs also undergo Pros-mediated

terminal differentiation, we first examined Pros expression together with the pan-NB reporter

Dpn::GFP at 12 hr APF. We found that many Dpn$^+$ NBs expressed nuclear Pros (Figure 4A).

Then, we examined the expressions of Dpn::GFP with Miranda (Mira) marks mature NBs

and asymmetric cell division at 12 hr APF. Interestingly, we occasionally observed some

small Dpn$^+$Mira$^-$ cells, that form doublets (Figure 4B), which we think are likely to be

progenies of NB symmetric divisions. Therefore, it is possible that medulla NBs also undergo

size symmetric divisions via Pros localisation like NBs of other lineages. To test if Pros is

required for the termination of medulla NBs, we induced *pros* RNAi clones (Shaw *et al*,

2018). This manipulation caused the formation of ectopic NBs in the deep layers of the

medulla where neurons reside during late larval stages (Figure 4C-B). In the *pros RNAi*

clones, many persistent NBs were found at 24 hr APF in contrast to the control (Figure 4E-

G). Together, our data suggest Pros is required to promote differentiation of the medulla NBs,

and consequently shut down medulla neurogenesis during pupal life (Figure 4H).

Tailless is involved in controlling the timing of medulla NB termination.

In the OL, medulla NBs sequentially express an array of temporal transcription factors as

they age to promote cellular diversity (Konstantinides et al., 2022; Li et al., 2013; Zhu et al.,

2022). The simplified temporal series includes Homothorax (Hth), Eyeless (Ey), Sloppy-

paired (Slp), Dichaete (D) and Tailless (Tll) (Figure 5A). It was previously shown that

nuclear Pros localisation occurred during the Tll$^+$ temporal window (Li et al., 2013); and that

the glial regulatory gene, Gcm, is expressed in the Tll$^+$ NBs prior a switch from neurogenesis

to gliogenesis (Konstantinides et al., 2022; Li et al., 2013; Zhu et al., 2022).

At 12 hr APF, we found some Tll$^+$ NBs were also *gcm*$^+$ (*gcm*-lacZ, Vincent *et al*,

1996) (Figure 5B), indicating that a neurogenic to gliogenic switch could also contribute

towards the disappearance of NBs at this time point. Consistent with this, at 12 hr APF, we

observed some superficially located Dpn$^+$ NBs that expressed both the differentiation marker

Pros and *gcm*-lacZ (Figure 5C, yellow arrowheads), suggesting that these NBs could be

undergoing a NB to glioblast cell fate transition. Deeper within the medulla, we found *gcm*$^+$

cells which express neither Dpn nor Pros (Figure 5C, white arrowhead), that are likely to be

the progenies of the glioblasts. Finally, we found a few medially located Dpn$^+$ NBs that

expressed Pros but not *gcm*-lacZ (Figure 5G, red arrowhead). This is consistent with what we

observed in Figure 4A-B, where some NBs undergo terminal differentiation through likely

via symmetric cell division.

To test if members of the temporal transcription factor series can alter NB persistence

and prevent the neurogenic to gliogenic switch, we first examined the role of Tll, which

defines the temporal window where terminal differentiation as well as the gliogenic switch

occur (Li et al., 2013; Zhu et al., 2022). First, we induced *tll* overexpression clones and found

that *tll* overexpression caused the formation of ectopic NBs in the larval CNS (Figure 5D, and

these NBs persisted at 24h APF (Figure 5E, n = 4/4 lobes). *tll* overexpression has been

previously shown to convert type I NBs to a type II NB cell fate that are capable of tumour

formation in type I NBs of the CB (Hakes & Brand, 2020). Indeed, we found that the *tll*

overexpressing NBs were mostly Dpn$^+$Ase$^-$ which is characteristic of type II NBs (Figure 5D-

E). Therefore, it is possible that *tll* overexpression caused type I to type II conversions that

accounts for why these clones failed to undergo termination on time.

Consistent with reports that Tll is not required for glial production in the larval

medulla NBs (Zhu et al., 2022), we found that inhibition of *tll* via a validated RNAi (Hakes &

Brand, 2020) was not able to alter the number of glial cells in the larval medulla

(Supplementary Figure 3A, B, E), nor the timing of medulla NB termination in pupal stages

(Supplementary Figure 3C, D, F). Moreover, the knockdown of *tll* did not alter the number of

NBs undergoing cell death assayed via Dcp-1 expression (Supplementary figure 3G-H) nor

the expression of Pros which promotes terminal size symmetric divisions (Supplementary

figure 3I-J). Together, while Tll is not required for medulla NB termination, ectopic

expression of Tll can induce persistent NBs. Furthermore, Tll acts independently of cell death

and symmetric NB termination mechanisms.

To assess whether early temporal transcription factors are required for the termination

of medulla NBs, we overexpressed Hth and inhibited Ey or Scarecrow (Scro) (Konstantinides

et al., 2022; Zhu et al., 2022) via RNAi in clones, which prevents the progression through the

temporal series. Nevertheless, none of these manipulations influenced the timing of NB

termination (data not shown). Taken together, altering temporal progression is not sufficient

to affect the timing of medulla NB termination in the pupal stages.

Gcm-induced gliogenic switch can affect the timing of NB termination.

We next assessed if manipulating the gliogenic switch can affect the timing of medulla NB

termination. Here, we induced *gcm* overexpression clones using a reagent previously

validated in Zhu et al., 2022. This manipulation was able to induce the formation of ectopic

glial cells during larval neurogenesis (Figure 5F-G). At 12 hr APF, we recovered

significantly fewer NBs in the OL upon *gcm* overexpression (Figure 5H-J), suggesting that

the induction of a gliogenic switch can cause precocious medulla NB termination. To assess if

the downregulation of *gcm* can cause prolonged NB persistence in the medulla, we knocked

down *gcm* using a previously validated RNAi (Zhu *et al*, 2022) in clones. This manipulation
was able to inhibit the expression of the glial cell marker Repo in clones during larval stages
(Supplementary figure 3K-L). However, no persistent medulla NBs was found at 24 hr APF
(Supplementary figure 3M-N, n = 5/5 lobes). This suggests that while the downregulation of
gcm is not sufficient to induce NB persistence, *gcm* overexpression can cause a neurogenic to
gliogenic switch and thereby, premature termination of NBs.
263
Discussion

Robust brain development requires the tight coordination between stem cell maintenance and
differentiation. In the *Drosophila* OL, there are two main stem cell pools, the NE cells that
divide symmetrically, and the NBs. Medulla NBs are derived from the OPC NE via the
proneural wave, which switch to asymmetric division where they self-renew and give rise to
GMCs that will produce post-mitotic neurons and glial cells of the medulla cortex (Egger et
al., 2007; Li et al., 2013; Yasugi et al., 2008). Our study shows that during early pupal
development, medulla NBs terminate via a combination of mechanisms including cell death,
Pros-mediated differentiation and a switch from neurogenesis to gliogenesis. While factors
such as ecdysone signalling and OxPhos of the NBs are involved in NB termination in other
lineages outside of the OL (Homem et al., 2014; Syed et al., 2017; van den Ameele & Brand,
2019; Yang et al., 2017), we found that these factors do not play a significant role in the
termination of medulla NBs. Instead, the main regulator of the timely NB cessation in the
pupal medulla is regulated by the timing of the conversion of NE-NB conversion during
larval stages. Intriguingly, while a specific cascade of temporal identity transcription factors
in the medulla NBs has been shown to generate the cellular diversity of medulla neurons and
glia (Konstantinides et al., 2022; Li et al., 2013; Zhu et al., 2022), temporal progression in the
medulla NBs appears dispensable for the timing of NB cessation in the pupal stages (Figure
6). Diseases such as microcephaly are mediated by the premature depletion of the NE pool, or
through a precocious commitment to neurogenic divisions (Götz & Huttner, 2005).
Consistent with this, our data present evidence that during development, the regulation of the
NE stem cell pool, and its timely conversion into NSCs is of critical importance to the overall
size and function of the brain.

NE-NB transition determines the timely termination of medulla NBs.
Our data support a model whereby once the NE pool is exhausted, no new NBs can be
generated. NBs transit through their temporal series to terminally differentiate to become

glial cells or neurons, which marks the end of medulla neurogenesis. Thus, unlike type I NB

lineages in which NB termination is scheduled by ecdysone signalling-mediated temporal

progression at the level of the NBs (Syed et al., 2017; Yang et al., 2017), in the OL, NB

termination is pre-determined in the NE. In the OL, besides medulla NBs, NE cells also

sequentially express an array of temporal factors, such as Chinmo and Broad downstream of

ecdysone signalling (Arain et al., 2022; Dillard et al., 2018; Lanet et al., 2013; Zhou et al.,

2019). Moreover, inhibiting the temporal Chinmo-to-Broad expression switch in the NE was

shown to delay the NE-NB transition, resulting in tumours with persistent NBs in the adult

OLs (Dillard et al., 2018; Zhou et al., 2019). The differences in the regulation of NB

termination in the OL medulla versus other lineages may be due to some fundamental

differences in the origins of these NBs. For example, in the OL, NBs are derived from the

NE, that undergoes rapid growth during larval development (Egger et al., 2007; Lanet et al.,

2013; Yasugi et al., 2008). Therefore, NB termination in the OL is scheduled at the level of

the NE, likely by the activation of ecdysone signalling during late larval development. In

contrast, NBs from other lineages are produced directly from the neuroectoderm in the

embryo without undergoing the transition from the NE (Harding & White, 2018). Hence,

their termination mechanism acts predominantly in the NBs during pupal development.

Cell death makes a small contribution to NB termination.

Consistent with previous reports, we found that cell death occurred in a subset of medulla

NBs at 12 hr APF (Hara *et al*, 2018). However, inhibition of cell death did not fully prevent

medulla NBs from termination in the pharate OLs (Supplementary Figure 2D), suggesting

that NBs can adopt alternative routes for timely termination before adulthood. From our data,

it is not yet clear whether a specific subset of medulla NBs undergo cell death. Although the

timing of cell death in the medulla NBs coincides with a small pulse of ecdysone around 10

hr APF (Riddiford, 1993), we showed that inhibition of ecdysone signalling in the medulla

NBs did not phenocopy the inhibition of apoptosis (Supplementary Figure 1B-E). Therefore,

unlike neurons in the OL which require ecdysone signalling for cell death during pupal

development (Hara et al., 2013), medulla NBs undergo cell death independently of ecdysone

signalling. Altogether, while some NBs terminate proliferation through cell death, it is likely

that cell death of the NBs makes a minor contribution to the termination of neurogenesis in

the medulla.

A neurogenic-gliogenic switch and symmetric divisions cause NB cessation.

Our data show that many medulla NBs displayed nuclear Pros expression during early pupal

stages (Figure 3A). Consistent with the essential role of Pros in promoting differentiation

(Choksi et al., 2006; Maurange et al., 2008), *pros* knockdown caused the persistence of the

NBs in the pupal OLs (Figure 3C-G), similar to other regions of the brain. In addition to

differentiation, another mechanism responsible for the cessation of neurogenesis, is the

commencement of gliogenesis. We found glioblasts in the early pupal medulla, which make

glial cells at the end of the temporal transcription factor series, express the NB marker Dpn,

the glial marker Gcm as well as the differentiation marker Pros (Figure 5B-C). As such, it is

possible that similar to the embryonic glioblasts, Pros is required for glial cell fate acquisition

via Gcm (Freeman & Doe, 2001; Ragone et al., 2001). As Gcm knockdown did not induce

any persistent medulla NBs during pupal stages (Supplementary Figure 3M-N), the presence

of Pros may likely induce symmetric division of the NBs, resulting in their termination.

Together, cell death, symmetric cell division and a gliogenic switch appear to be employed as

fail-safe mechanisms that help to eliminate the NB progenitor pool in the pupal OL. Hence, it

would be interesting to explore in the future the cross-regulatory relationship between these

termination programs.

Materials and Methods

Fly husbandry

Fly stocks were reared on standard media at 25°C. For larval dissection, brains were dissected at wandering L3 stages (120 hr ALH). For pupal dissection, white pupae were selected and allow to age until the desired stages (12 hr, 24 hr, 30 hr, 48 hr, and 72 hr APF).

Wildtype fly was w^{1118}. *eyR16F10-GAL4* (BL 48737) was used to induce expression of transgenes in a subset of medulla NBs and their progeny from larval development (Veen *et al*, 2023). Knockdown and overexpression clones were induced by flip-out system using *hsFLP,act>CD2>GAL4;UAS-Dcr2,UAS-GFP* (a gift from Lei Zhang). Flip-out clones were induced at 24 hr ALH for 10 min. *FRT42D;UAS-tll* clones were induced using the MARCM system: *UAS-mCD8::GFP,hsFLP;FRT42D, tub-GAL80; tub-GAL4* (a gift from Alex Gould). Clones were induced at 48 hr ALH for 15 min.

The fly strains used were as followed: *Dpn::GFP* (Morin *et al*, 2001), *FRT42D* (a gift from Alex Gould), *gcm-lacZ* (BL 5445), *Tll::EGFP* (BL 30874), *Pros::GFP* (VDRC 318463), *UAS-Atg1 RNAi* (BL 26731), *UAS-dp110^{CAAX}* (BL 25908), *UAS-EcR RNAi* (VDRC 37058), *UAS-EcR^{DN}* (BL 9451), *UAS-gcm* (BL 5446), *UAS-gcm RNAi* (VDRC 110539), *UAS-l'sc* (BL 51670), *UAS-lacZ* (BL 8529), *UAS-mCherry RNAi* (BL 35785), *UAS-med27 RNAi* (BL 34576), *UAS-N^{ACT}* (a gift from Helena Richardson), *UAS-ND75 TRiP* (BL 33911), *UAS-notch RNAi* (VDRC 100002), *UAS-p35* (BL 5072), *UAS-pros RNAi* (BL 26745), *UAS-tll* (a gift from Claude Desplan), *UAS-tll RNAi* (VDRC 330031), *UAS-D* (BL 8861).

Immunostaining

Larval and pupal brains were dissected out in phosphate buffered saline (PBS), fixed in 4% formaldehyde for 20 min and rinsed three time in 0.5% PBST (PBS + 0.5% Triton) at room temperature. For immunostaining, brains were incubated in primary antibodies overnight at 4°C, followed by two washes in 0.5% PBST and then an overnight secondary antibody incubation at 4°C or 3 hrs at room temperature. Brains were afterwards rinsed

two times in 0.5% PBST and incubated in 50% glycerol in PBS for 20 min. Samples were

mounted in 80% glycerol in PBS, on glass slides and sealed with coverslips (22x22 mm,

No.1.5, Knittel) for image acquisition. The primary antibodies used were: rat anti-Dpn

(1:200, Abcam 195172), chick anti-GFP (1:1000, Abcam 13970), mouse anti-EcR^{common}

(1:50, DSHB Ag10.2), rabbit anti-Dcp-1 (1:100, Cell Signalling 95785), rat anti-Mira

(1:100, Abcam 197788), rat anti-Pros (a gift from Fumio Matsuzaki), rabbit anti-β-

Galactosidase (1:200, a gift from Helena Richardson), rabbit anti-Ase (a gift from Lily Jan

and Yuh Nung Jan), mouse anti-Repo (1:50, DSHB 8D12), and rat anti-DE-Cad (1:200,

DSHB DCAD2), rabbit anti-PatJ (1:500, a gift from Helena Richardson). Secondary

donkey antibodies conjugated to Alexa 555 and Alexa 647, and goat antibodies conjugated

to Alexa 405, 488, 555, and 647 (Molecular Probes) were used at 1:500.

Imaging and image processing

Images were acquired using Olympus FV3000 confocal microscope with 40x (NA 0.95,

UPLSAPO) and 60x (NA 1.30, UPLSAPO) objectives. Δz = 1.5 μm. Images were processed

using Fiji (https://imagej.net/Fiji). Single section or maximum projections were shown as

representative images in the figures.

Quantification was performed in Fiji or via 3D reconstruction in Imaris (Bitplane).

For quantifications of NB numbers (Figures 1C, Supplementary Figure 1E, 2E), number of

Dpn$^+$ NBs in the medulla/OL were measured. For quantifications of nuclear sizes of

medulla NBs (Figure 1E), nuclear diameters identified by Dpn were measured. For

quantifications of the ratio of NBs in clones (Figures 2H, 2K, 3D, 4E, 5L, Supplementary

figure 1J, 3H), GFP$^+$ clones were first detected, and clonal volumes were measured. The

GFP$^+$ areas were then utilised to make a mask for GFP$^+$Dpn$^+$ or GFP$^+$Mira$^+$ NBs within the

clones, which was in turn used to measure the total Dpn volume. The NB ratio in clones

is calculated as the ratio of Dpn$^+$GFP$^+$ or GFP$^+$Mira$^+$ volume over GFP$^+$ volume. For

quantification of the numbers of glia in the medulla (Supplementary Figure 3G), the total

numbers of Repo$^+$ cells in the deep section of the medulla were measured in each lobe.

For quantification of the NE volumes (Figure 2B), DE-Cad$^+$ volumes were measured in

each lobe. Images were assembled in Adobe InDesign 2024 and Affinity Publisher 2. Scale

bars = 10 μm or 50 μm per indicated in figures.

Statistical analyses

At least two animals per genotype were used for all experiments. Statistical analyses were

performed in GraphPad Prism 9. In graphs, data (crossbars) = mean ± standard error of

the mean (SEM). n = lobes unless specified. Graphs were plotted using RStudio. For

comparisons between two conditions, P-values were calculated by non-parametric Mann-

Whitney tests when data show non-normally distribution. For comparisons between

more than two conditions, P-values were calculated by either ordinary one-way ANOVA

tests for normally distributed data, or Kruskal-Wallis tests for non-normally distributed

data. Dunnet or Dunn's tests were used to correct for multiple comparisons following

one-way ANOVA and Kruskal-Wallis tests, respectively. *(*) $p<0.05$ (**) $p< 0.01$, (***)*

$p< 0.001$, (****) $p < 0.0001$, (ns) $p>0.05$

Acknowledgements

We are grateful to Lei Zhang, Alex Gould, Claude Desplan, Fumio Matsuzaki, Helena

Richardson, Yuh Nung Jan for generous sharing of fly stocks and antibodies. We would also

like to thank Holger Apitz, Édel Alvarez-Ochoa, and Qian Dong for fruitful discussions and

critical reading of the manuscript. We also thank Bloomington Drosophila Stock Centre,

Vienna Drosophila Resource Centre, and Developmental Studies Hybridoma Bank for fly

stocks and antibodies. We would like to also thank OZDros for *Drosophila* quarantine, Peter

MacCallum Cancer Institute CAHM for microscopy assistance. LYC's laboratory is supported

by funding from the NHMRC Ideas Grant (APP2011289), and the Peter MacCallum Cancer

Foundation.

Disclosure and competing interests statement

The authors declare no competing interests.

References

Ameele J van den & Brand AH (2019) Neural stem cell temporal patterning and brain tumour growth rely on oxidative phosphorylation. *eLife* 8

van den Ameele J & Brand AH (2019) Neural stem cell temporal patterning and brain tumour growth rely on oxidative phosphorylation. *eLife* 8: e47887

Apitz H & Salecker I (2014) A challenge of numbers and diversity: neurogenesis in the Drosophila optic lobe. *J Neurogenet* 28: 233–249

Arain U, Islam IM, Valentino P & Erclik T (2022) Concurrent temporal patterning of neural stem cells in the fly visual system. 2022.10.13.512100 doi:10.1101/2022.10.13.512100 [PREPRINT]

Bello BC, Hirth F & Gould AP (2003) A Pulse of the Drosophila Hox Protein Abdominal-A Schedules the End of Neural Proliferation via Neuroblast Apoptosis. *Neuron* 37: 209–219

Bello BC, Izergina N, Caussinus E & Reichert H (2008) Amplification of neural stem cell proliferation by intermediate progenitor cells in Drosophila brain development. *Neural Development* 3: 5

Boone JQ & Doe CQ (2008) Identification of Drosophila type II neuroblast lineages containing transit amplifying ganglion mother cells. *Dev Neurobiol* 68: 1185–1195

Bowman SK, Rolland V, Betschinger J, Kinsey KA, Emery G & Knoblich JA (2008) The tumor suppressors Brat and Numb regulate transit-amplifying neuroblast lineages in Drosophila. *Dev Cell* 14: 535–546

Choksi SP, Southall TD, Bossing T, Edoff K, de Wit E, Fischer BE, van Steensel B, Micklem G & Brand AH (2006a) Prospero acts as a binary switch between self-renewal and differentiation in Drosophila neural stem cells. *Dev Cell* 11: 775–789

Choksi SP, Southall TD, Bossing T, Edoff K, de Wit E, Fischer BE, van Steensel B, Micklem G & Brand AH (2006b) Prospero acts as a binary switch between self-renewal and differentiation in Drosophila neural stem cells. *Dev Cell* 11: 775–789

Dillard C, Narbonne-Reveau K, Foppolo S, Lanet E & Maurange C (2018) Two distinct mechanisms silence chinmo in Drosophila neuroblasts and neuroepithelial cells to limit their self-renewal. *Development* 145

Egger B, Boone JQ, Stevens NR, Brand AH & Doe CQ (2007) Regulation of spindle orientation and neural stem cell fate in the Drosophila optic lobe. *Neural Dev* 2: 1

Egger B, Gold KS & Brand AH (2010) Notch regulates the switch from symmetric to asymmetric neural stem cell division in the Drosophila optic lobe. *Development* 137: 2981–2987

Freeman MR & Doe CQ (2001) Asymmetric Prospero localization is required to generate mixed neuronal/glial lineages in the Drosophila CNS. *Development* 128: 4103–4112

Götz M & Huttner WB (2005) The cell biology of neurogenesis. *Nat Rev Mol Cell Biol* 6: 777–788

Hakes AE & Brand AH (2020) Tailless/TLX reverts intermediate neural progenitors to stem cells driving tumourigenesis via repression of asense/ASCL1. *eLife* 9: e53377

Hara Y, Hirai K, Togane Y, Akagawa H, Iwabuchi K & Tsujimura H (2013) Ecdysone-dependent and ecdysone-independent programmed cell death in the developing optic lobe of Drosophila. *Developmental Biology* 374: 127–141

Hara Y, Sudo T, Togane Y, Akagawa H & Tsujimura H (2018) Cell death in neural precursor cells and neurons before neurite formation prevents the emergence of abnormal neural structures in the Drosophila optic lobe. *Developmental Biology* 436: 28–41

Harding K & White K (2018a) Drosophila as a Model for Developmental Biology: Stem Cell-Fate Decisions in the Developing Nervous System. *J Dev Biol* 6

Harding K & White K (2018b) Drosophila as a Model for Developmental Biology: Stem Cell-Fate Decisions in the Developing Nervous System. *J Dev Biol* 6

Homem CCF, Steinmann V, Burkard TR, Jais A, Esterbauer H & Knoblich JA (2014) Ecdysone and mediator change energy metabolism to terminate proliferation in Drosophila neural stem cells. *Cell* 158: 874–888

Ito K & Hotta Y (1992a) Proliferation pattern of postembryonic neuroblasts in the brain of Drosophila melanogaster. *Developmental Biology* 149: 134–148

Ito K & Hotta Y (1992b) Proliferation pattern of postembryonic neuroblasts in the brain of Drosophila melanogaster. *Developmental Biology* 149: 134–148

Konstantinides N, Holguera I, Rossi AM, Escobar A, Dudragne L, Chen Y-C, Tran TN, Martínez Jaimes AM, Özel MN, Simon F, *et al* (2022) A complete temporal transcription factor series in the fly visual system. *Nature* 604: 316–322

Lanet E, Gould AP & Maurange C (2013a) Protection of neuronal diversity at the expense of neuronal numbers during nutrient restriction in the Drosophila visual system. *Cell Rep* 3: 587–594

Lanet E, Gould AP & Maurange C (2013b) Protection of neuronal diversity at the expense of neuronal numbers during nutrient restriction in the Drosophila visual system. *Cell Rep* 3: 587–594

Leevers SJ, Weinkove D, MacDougall LK, Hafen E & Waterfield MD (1996) The Drosophila phosphoinositide 3-kinase Dp110 promotes cell growth. *EMBO J* 15: 6584–6594

Li X, Erclik T, Bertet C, Chen Z, Voutev R, Venkatesh S, Morante J, Celik A & Desplan C (2013a) Temporal patterning of Drosophila medulla neuroblasts controls neural fates. *Nature* 498: 456–462

Li X, Erclik T, Bertet C, Chen Z, Voutev R, Venkatesh S, Morante J, Celik A & Desplan C (2013b) Temporal patterning of Drosophila medulla neuroblasts controls neural fates. *Nature* 498: 456–462

Maurange C, Cheng L & Gould AP (2008) Temporal transcription factors and their targets schedule the end of neural proliferation in Drosophila. *Cell* 133: 891–902

Morin X, Daneman R, Zavortink M & Chia W (2001) A protein trap strategy to detect GFP-tagged proteins expressed from their endogenous loci in Drosophila. *Proceedings of the National Academy of Sciences* 98: 15050–15055

Pahl MC, Doyle SE & Siegrist SE (2019) E93 integrates neuroblast intrinsic state with developmental time to terminate neurogenesis via autophagy. *Current biology : CB* 29: 750

Ragone G, Bernardoni R & Giangrande A (2001) A Novel Mode of Asymmetric Division Identifies the Fly Neuroglioblast 6-4T. *Developmental Biology* 235: 74–85

Riddiford LM (1993) Hormones and Drosophila development. In *The development of Drosophila melanogaster* pp 899–939.

Rust K, Tiwari MD, Mishra VK, Grawe F & Wodarz A (2018) Myc and the Tip60 chromatin remodeling complex control neuroblast maintenance and polarity in Drosophila. *EMBO J* 37: e98659

Sato M, Yasugi T & Trush O (2019) Temporal patterning of neurogenesis and neural wiring in the fly visual system. *Neuroscience Research* 138: 49–58

Shaw RE, Kottler B, Ludlow ZN, Buhl E, Kim D, Morais da Silva S, Miedzik A, Coum A, Hodge JJ, Hirth F, *et al* (2018) In vivo expansion of functionally integrated GABAergic interneurons by targeted increase in neural progenitors. *EMBO J* 37: e98163

Siegrist SE, Haque NS, Chen C-H, Hay BA & Hariharan IK (2010) Inactivation of both foxo and reaper promotes long-term adult neurogenesis in Drosophila. *Current biology : CB* 20: 643

Syed MH, Mark B & Doe CQ (2017a) Steroid hormone induction of temporal gene expression in Drosophila brain neuroblasts generates neuronal and glial diversity. *eLife* 6: e26287

Syed MH, Mark B & Doe CQ (2017b) Steroid hormone induction of temporal gene expression in Drosophila brain neuroblasts generates neuronal and glial diversity. *eLife* 6: e26287

Veen K, Nguyen P-K, Froldi F, Dong Q, Alvarez-Ochoa E, Harvey KF, McMullen JP, Marshall O, Jusuf PR & Cheng LY (2023) Dedifferentiation-derived neural stem cells exhibit perturbed temporal progression. *EMBO reports* 24: e55837

Vincent S, Vonesch JL & Giangrande A (1996) Glide directs glial fate commitment and cell fate switch between neurones and glia. *Development* 122: 131–139

Wang W, Liu W, Wang Y, Zhou L, Tang X & Luo H (2011) Notch signaling regulates
neuroepithelial stem cell maintenance and neuroblast formation in Drosophila optic
lobe development. *Developmental Biology* 350: 414–428

Xu T, Nicolson S, Denton D & Kumar S (2015) Distinct requirements of Autophagy-related
genes in programmed cell death. *Cell Death Differ* 22: 1792–1802

Yang C-P, Samuels TJ, Huang Y, Yang L, Ish-Horowicz D, Davis I & Lee T (2017) Imp and
Syp RNA-binding proteins govern decommissioning of Drosophila neural stem cells.
Development 144: 3454–3464

Yasugi T, Sugie A, Umetsu D & Tabata T (2010) Coordinated sequential action of EGFR and
Notch signaling pathways regulates proneural wave progression in the Drosophila
optic lobe. *Development* 137: 3193–3203

Yasugi T, Umetsu D, Murakami S, Sato M & Tabata T (2008a) Drosophila optic lobe
neuroblasts triggered by a wave of proneural gene expression that is negatively
regulated by JAK/STAT. *Development* 135: 1471–1480

Yasugi T, Umetsu D, Murakami S, Sato M & Tabata T (2008b) Drosophila optic lobe
neuroblasts triggered by a wave of proneural gene expression that is negatively
regulated by JAK/STAT. *Development* 135: 1471–1480

Zhou Y, Yang Y, Huang Y, Wang H, Wang S & Luo H (2019) Broad Promotes
Neuroepithelial Stem Cell Differentiation in the Drosophila Optic Lobe. *Genetics*
213: 941–951

Zhu H, Zhao SD, Ray A, Zhang Y & Li X (2022) A comprehensive temporal patterning gene
network in Drosophila medulla neuroblasts revealed by single-cell RNA sequencing.
Nat Commun 13: 1247

Figure 1: Medulla NBs are terminated during early pupal development.

(A-A') Schematic depicting the larval CNS that is consisted of three main regions: the central brain (CB), the ventral nerve cord (VNC) and the optic lobe (OL). Box area indicates inset shown in A'. (A') The apical view of the outer proliferation centre (OPC) of the OL. Here, NE cells first symmetrically divide to self-renew. At mid-larval development, a proneural wave is initiated and travels from the medial to lateral of the NE. Upon the proneural wave, NE cells differentiate into NBs and switch to asymmetric cell division for neurogenesis.

(B-B''') In the wildtype (w^{1118}) OL, the number of medulla NBs marked by Dpn (dashed lines), gradually reduces from 12 hr to 24 hr APF. Medulla NBs completely disappear at 30 hr APF.

(C) Quantification of the numbers of medulla NBs at 12 hr, 18 hr, 24 hr, 30 hr APF in the wildtype OLs. Kruskal-Wallis test and Dunn's test to correct for multiple comparisons: *p = 0.0314. 12 hr: n = 6, m = 436.3 ± 25.21. 18 hr: n = 9, m = 45.22 ± 6.616. 24 hr: n = 5, m = 43.6 ± 9.255. 30 hr: n = 5, m = 0.000 ± 0.000.

(D-D') In the wildtype, the nuclear size of medulla NBs marked by Dpn decreases from 12 hr to 18 hr APF.

(E) Quantification of the nuclear diameters of medulla NBs at 12 hr and 18 hr APF. Mann-Whitney test: ****p < 0.0001. 12 hr: n = 109 cells, m = 5.182 ± 0.099 µm. 18 hr APF: n = 61 cells, m = 4.100 ± 0.082 µm.

Figure 2: The termination of medulla NBs is scheduled by the timing of the larval NE-NB transition.

(A-A'') The NE pool marked by DE-Cad (green, dashed lines) becomes gradually depleted from 12 hr to 24 hr APF. DAPI (blue)

(B) Quantification of DE-Cad volumes at 12 hr, 18 hr, and 24 hr APF. Kruskal-Wallis test and Dunn's test to correct for multiple comparisons: ***p = 0.0003. 12 hr: n = 5, m = 17922 ± 5129. 18 hr: n = 5, m = 6508 ± 747.0. 24 hr: n = 9, m = 1597 ± 280.9.

(C) Schematic depicting the larval OL at 120 hr ALH, where the NE-NB transition is boxed, and is the region of interest in (D-M).

(D-E') Representative images of *hsFLP* clones (dashed lines) at 120 hr ALH, induced with (D-D'') *UAS-mCherry RNAi* and (E-E'') *UAS-notch RNAi*. GFP (green), PatJ (grey), Mira (magenta).

(F-G') Representative maximum projections of the OL at 12 hr APF, in which *hsFLP* clones (dashed lines) are induced with (F-F') *UAS-mCherry RNAi* and (G-G') *UAS-notch RNAi*. DAPI (blue), GFP (green), Mira (magenta).

(H) Quantification of the NB ratio in clones in the OL at 12 hr APF in which *hsFLP* clones are induced with *UAS-mCherry RNAi* and *UAS-notch RNAi*. Mann-Whitney test: **p = 0.0041. *mCherry RNAi*: n = 7, m = 0.179 ± 0.054. *notch RNAi*: n = 11, m = 0.044 ± 0.013.

(I-I') Representative *hsFLP* clone (dashed line) at 120 hr ALH, induced with *UAS-l'sc*.

(J-J') Representative maximum projections of the OL at 12 hr APF, in which *hsFLP* clones (dashed line) are induced with *UAS-l'sc*. Actin (blue), GFP (green), Dpn (magenta)

(K) Quantification of the NB ratio in clones in the OL at 12 hr APF in which *hsFLP* clones are induced with *UAS-lacZ* and *UAS-l'sc*. Mann-Whitney test: **p = 0.0057. *lacZ*: n = 8, m = 0.009 ± 0.003. *l'sc*: n = 7, m = 0.002 ± 0.001.

(L-L') Representative *hsFLP* clone (dashed line) at 120 hr ALH, induced with $UAS\text{-}N^{ACT}$.

(M-M') Representative maximum projections of the OL at 72 hr APF, in which *hsFLP* clones (dashed lines) are induced $UAS\text{-}N^{ACT}$. Note that N^{ACT} clones do not generate NBs until late pupal development.

DAPI (blue), GFP (green), Mira (magenta)

Figure 3: A subset of medulla NBs terminate via cell death.

(A) At 12 hr APF, a subset of medulla NBs marked by Dpn::GFP (cyan) express the apoptotic marker Dcp-1 (grey) (arrowheads). A'-A''' is the magnified view of the boxed inset in A.

(B-C') Representative maximum projections of the OL (dashed line) at 24 hr APF with *hsFLP* clones induced with (B-B') *UAS-mCherry RNAi* and (C-C') *UAS-p35*. Ectopic NBs are indicated by arrowhead. DAPI (blue), GFP (green), Dpn (magenta). The same *hsFLP>mCherry RNAi* representative image is used in Supplementary figure 1F and Figure 4E.

(D) Quantification of the NB ratio in *hsFLP* induced with *UAS-mCherry RNAi* and *UAS-p35* at 24 hr APF in the OL. Mann-Whitney test: **p = 0.005. *mCherry RNAi*: n = 7, m = 0.001 ± 0.0004. *p35*: n = 5, m = 0.017 ± 0.006.

Figure 4: The termination of medulla NBs requires Pros-mediated size symmetric division.

(A-A''') Representative images of the OL at 12 hr APF. Some NBs marked by Dpn::GFP (cyan) in the medulla co-express Pros (red) (arrowheads). A'-A''' are magnified images of boxed inset in A.

(B-B''') Representative images of the OL at 12 hr APF. On the superficial layer where medulla NBs reside, we have occasionally observed a pair of similarly-sized cells at the most medial position (arrowheads), that express Dpn::GFP (cyan) but not Mira (magenta). B'-B''' are magnified images of boxed inset in B.

(C-D') Representative images of the deep layers of the medulla with *hsFLP* clones (dashed lines) driving (C-C) *UAS-mCherry RNAi* and (D-D') *UAS-pros RNAi* at 120 hr ALH. Arrowhead indicates ectopic NBs in the deep section of the medulla.

(E-F') Representative maximum projects of an OL (dashed line) with *hsFLP* clones driving (E-E') *UAS-mCherry RNAi* and (F-F') *UAS-pros RNAi* at 24 hr ALH. Arrowhead indicates

persistent NBs. The same *hsFLP>mCherry RNAi* representative image is used in Supplementary figure 1F and Figure 3B. DAPI (blue), GFP (green), Dpn (magenta).

(G) Quantification of the NB ratio in clones in the OL at 24 hr APF, in which *hsFLP* clones are induced with *UAS-mCherry RNAi* and *UAS-pros RNAi*. Mann-Whitney test: ***p = 0.0006. *mCherry RNAi*: n = 7, m = 0.001 ± 0.0004. *pros RNAi*: n = 7, m = 0.019 ± 0.005.

(H) Schematic depicting a medulla NB expresses nuclear Pros to promote symmetric division.

Figure 5: Gcm is sufficient to induce gliogenesis in the medulla at the expense of NBs.

(A) Schematic of the temporal progression in the OL medulla.

(B-B''') Representative images of the superficial layer of the medulla NBs at 12 hr APF. Some of the most medial NBs (arrowheads) express Dpn (gray), Tll::EGFP (green), and *gcm*-lacZ (red).

(C-C'''') Representative images of the OL at 12 hr APF. Here, some of the most medial NBs are $Dpn^+Pros^+gcm^+$ (yellow arrowheads) whereas some are $Dpn^+Pros^+gcm^-$ (red arrowheads) on the superficial layers. In the deep layers, some cells are $Dpn^-Pros^-gcm^+$ (white arrowheads). C'-C''' are magnified images of C.

(D-D''') Representative images of the OL at 120 hr ALH, in which MARCM clones overexpressed for Tll are induced. Box indicates inset shown in L'-L'''. In the clone (dashed line), medulla NBs expressed low levels of Ase compared to wildtype medulla NBs outside of the clone. D'-D''' are magnified images of D.

(E-E''') Representative maximum projections of the OL at 24 hr APF, in which MARCM overexpressed for Tll are induced. In the clone (dashed line), ectopic NBs do not express Ase. E'-E''' are magnified images of E. GFP (green), Dpn (magenta), Ase (cyan).

(F-G') Representative *hsFLP* clones (dashed lines) induced with (F-F') *UAS-lacZ*, and (G-G') *UAS-gcm* at 120 hr APF.

(H-I') Representative images of the OLs (dashed lines) at 12 hr APF, in which *hsFLP* clones are induced with (H-H') *UAS-lacZ*, and (I-I') *UAS-gcm*.

(J) Quantification of the NB ratio in clones in the OLs at 12 hr APF, in which *hsFLP* clones are induced with *UAS-lacZ* and *UAS-gcm*. Mann-Whitney test: *p = 0.0379. *lacZ*: n = 8, m = 0.009 ± 0.003. *gcm*: n= 8, m = 0.003 ± 0.0004.

(K) Schematic depicting a Dpn^+ medulla NB expresses nuclear Pros and the glial cell fate determinant Gcm to induce a switch from neurogenic-to-gliogenic program.

Figure 6: Working model of the termination of medulla NBs.

Timing of NB termination in the early pupal stages is scheduled by the timing of the NE-NB transition in the larval stages. Medulla NBs can be terminated via three mechanisms: (1) cell death, (2) Pros-mediated symmetric division and (3) Gcm-mediated gliogenic switch of the NBs.