Neurexins control the strength and precise timing of glycinergic inhibition in the

auditory brainstem

He-Hai Jiang[1,2,3], Ruoxuan Xu[1], Xiupeng Nie[1], Zhenghui Su[1], Xiaoshan Xu[1], Ruiqi Pang[4,5], Yi

Zhou[4], Fujun Luo[1,2,3]*

1. Guangzhou National Laboratory, 510005, Guangzhou, Guangdong, China

2. Bioland Laboratory, 510005, Guangzhou, Guangdong, China

3. School of Basic Medical Sciences, Guangzhou Medical University, 510005, Guangzhou,

Guangdong, China

4. Department of Neurobiology, School of Basic Medicine, Army Medical University, 400038,

Chongqing, China

5. Advanced Institute for Brain and Intelligence, School of Medicine, Guangxi University,

530004, Nanning, China

*Correspondence author. luo_fujun@gzlab.ac.cn (F.L.)

Abstract

Neurexins play diverse functions as presynaptic organizers in various glutamatergic and

GABAergic synapses. However, it remains unknown whether and how neurexins are involved

in shaping functional properties of the glycinergic synapses, which mediate prominent

inhibition in the brainstem and spinal cord. To address these issues, we examined the role of

neurexins in a model glycinergic synapse between the principal neuron in the medial nucleus

of the trapezoid body (MNTB) and the principal neuron in the lateral superior olive (LSO) in the

auditory brainstem. Combining RNAscope with stereotactic injection of AAV-Cre in the MNTB

of neurexin1/2/3 conditional triple knockout mice, we showed that MNTB neurons highly

express all isoforms of neurexins although their expression levels vary remarkably. Selective

ablation of all neurexins in MNTB neurons reduced not only the amplitude but also the kinetics

of the glycinergic synaptic transmission at LSO neurons. The synaptic dysfunctions were

primarily caused by impaired Ca^{2+} sensitivity of release and tightness of coupling between

voltage-gated Ca^{2+} channels and synaptic vesicles. Together, our current findings demonstrate

that neurexins are essential in controlling the strength and temporal precision of the glycinergic

synapse, which therefore corroborates the role of neurexins as key presynaptic organizers in

all major types of fast chemical synapses.

Introduction

Neurexins are evolutionarily conserved synaptic adhesion molecules and their genetic mutations are highly associated with autism and schizophrenia (Gomez et al., 2021; Sudhof, 2008; 2017). In the vertebrate, neurexins are encoded by three genes Nrxn1, Nrxn2, Nrxn3, each of which contains distinct promoters for expressing longer α- and shorter β-neurexins (Ullrich et al., 1995; Ushkaryov et al., 1992). Both α- and β-neurexins are critical and non-redundant in regulating the formation and function of synapses (Anderson et al., 2015; Missler et al., 2003; Sudhof, 2017). Triple knockout (KO) of α-neurexins causes a significant decrease in the density of GABAergic but not glutamatergic synapses while producing a strong reduction in both inhibitory and excitatory synaptic transmission(Missler *et al.*, 2003). Triple KO of β-neurexins reveals a significant phenotype only in glutamatergic synapses, but not in GABAergic synapses (Anderson *et al.*, 2015). Deletion of all neurexins, including Nrxn1/2/3 α- and β-neurexins, leads to profound but strikingly distinct phenotypes in different synapses, which further supports diverse actions of neurexins in specific neurons (Chen et al., 2017; Luo et al., 2020). Consistently, transcriptomic analysis has revealed that the expression profiles of various neurexins differ remarkably in excitatory neurons versus inhibitory neurons even among distinct subtypes of interneurons (Foldy et al., 2016; Fuccillo et al., 2015; Lukacsovich et al., 2019; Uchigashima et al., 2019). Despite the multifaceted roles of neurexins widely observed in various glutamatergic and GABAergic synapses (Boxer and Aoto, 2022; Chen *et al.*, 2017; Luo et al., 2021; Sudhof, 2017), however, it remains largely unknown how neurexins may regulate the functional properties of the glycinergic synapses, which mediate essential inhibition in the spinal cord, brainstem and retina (Assareh et al., 2023; Brill et al., 2020; Chang et al., 2023; Feng et al., 1998; Foster et al., 2015; Gomeza et al., 2003a; Gomeza et al., 2003b; Zeilhofer et al., 2012).

In the mammalian auditory brainstem, it is well-characterized that the principal neurons in the medial nucleus of the trapezoid body (MNTB) project glycinergic synaptic inputs to the lateral superior olive (LSO) (Fischer et al., 2019; Kotak et al., 1998; Nabekura et al., 2004; Sanes and Friauf, 2000). These synapses are robust, precise and extremely reliable even during high frequency stimulation(Krachan et al., 2017). Strong evidence has shown that the

MNTB-LSO glycinergic synapses are subject to extensive refinement during development,

including transmitter type switch, the elaboration of dendrites and axonal arbors followed by

synaptic pruning, to achieve exquisite control of fast inhibition(Brill *et al.*, 2020; Kim and

Kandler, 2003; Kotak *et al.*, 1998; Nabekura *et al.*, 2004; Sanes and Friauf, 2000). However,

the molecular mechanisms of shaping their functional characteristic remain incompletely

understood.

We utilized the triple Nrxn1/2/3 conditional KO mice (Nrxn123 cTKO) (Chen *et al.*, 2017)

to ablate all neurexins in MNTB principal neurons with stereotactic injection of associated

adeno-virus (AAV) expressing Cre-recombinase. Combined with RNAscope technique, we

verified that MNTB neurons highly express various isoforms of neurexins. Selective injection of

AAV-Cre, but not AAV-EGFP, successfully blocked expression of all neurexins in MNTB

neurons. Functionally, pan-neurexin deletion significantly impaired both the amplitude and the

kinetics of the glycinergic synaptic transmission. Surprisingly, deletion of all neurexins led to a

significantly higher number of glycinergic synapses innervating the LSO and higher frequency

of spontaneous neurotransmitter release. Application of low extracellular Ca^{2+} blocked the

glycinergic neurotransmission of both control and neurexin-deficient synapses, but with a

significant difference in its effectiveness. Moreover, treatment with EGTA, a high-affinity Ca^{2+}

chelator with slow binding kinetics, caused a much stronger inhibition of neurotransmission in

neurexin-deficient synapses, suggesting a significant disruption of the tight coupling between

voltage-gated Ca^{2+} channels and synaptic vesicles in the MNTB-LSO glycinergic synapses.

Results

Neurexins are highly expressed in MNTB neurons

To study the role and mechanism of neurexins in the glycinergic synapse between MNTB

neurons and LSO neurons, which are critical in computing sound localization in mammalian

auditory brainstem (Grothe et al., 2010), we first characterized the expression of Nrxn1-3 in the

principal neurons of the MNTB by fluorescent *in-situ* hybridization (FISH) using RNAscope

technique. As shown in Figure 1, MNTB neurons in wild-type (WT) mice expressed Nrxn1,

Nrxn2, and Nrxn3, although the expression level of each gene appeared to vary remarkably.

As an initial exploration, we took advantage of Nrxn123 cTKO mice that enable deletion of all

neurexins following the expression of Cre recombinase (Chen *et al.*, 2017), to bypass the

functional redundancy of different neurexins. Neuronal specific deletion of all neurexins was

achieved by stereotactic injection of AAV-Cre-EGFP into the MNTB at P0 (TKO). As control,

AAV-EGFP was injected (Ctrl). Successful virus injection and expression of Cre recombinase

in MNTB neurons were confirmed by strong EGFP fluorescence (Figure 1d-e). FISH data

further demonstrated that the recombinase dramatically blocked the expression of all

neurexins, as compared to the normal expression of neurexins when EGFP was expressed

(Figure 1f, compared to 1b). Altogether, these data show that the glycinergic neurons in the

MNTB highly express various neurexin genes, which can be specifically deleted by

stereotactic injection of AAV-Cre in Nrxn123 cTKO mice.

Neurexins are essential for dictating the strength and kinetics of the MNTB-LSO

glycinergic synapse

Since neurexins play an essential role in orchestrating functional organization of various

synapses including both glutamatergic and GABAergic synapses, we hypothesize that deletion

of all neurexins may have a significant impact on glycinergic synapses. To test the hypothesis,

we sought to examine how neurexins may regulate the function of the MNTB-LSO glycinergic

synapse in the auditory brainstem. Because LSO neurons receive excitatory glutamatergic

inputs and inhibitory GABAergic/glycinergic inputs, we used a pharmacologic approach to

isolate the glycinergic inputs. By blocking glutamatergic inputs with CNQX and AP5, we were

able to selectively record inhibitory postsynaptic currents (IPSCs) from LSO neurons, which

can be completely inhibited by strychnine, a potent glycine receptor inhibitor (Figure S1).

To specifically examine the role of neurexins in regulating the glycinergic synapse, we

used an optogenetic approach in which two AAVs, AAV-Cre-EGFP and

AAV-DIO-ChR2-mCherry, were co-injected in Nrxn123 cTKO mice. As control,

AAV-ChR2-EYFP was used for injection. Virus injection was performed at P0 and acute slices

derived from these mice were analyzed at P13-14. We verified the successful expression of

ChR2 by examining the fluorescence and further by recording optogenetic-evoked action

potentials from the soma of MNTB neurons. Reliable action potential (AP) firing can be

induced by single or repetitive high-frequency light stimulation (Figure S1c). In control mice,

optogenetic stimulation of MNTB neurons robustly triggered large IPSCs recorded from LSO

neurons (Figure 2a-b), which can be fully blocked by strychnine (Figure S1d) and are thus

purely glycinergic. In TKO mice with co-injection of Cre and ChR2, however, optogenetic

stimulation induced significantly smaller IPSCs (Figure 2a-b). On average, we observed ~60%

reduction in the peak amplitude (Figure 2b), similar to previous findings at other different

synapses (Chen *et al.*, 2017; Luo *et al.*, 2020). Furthermore, the kinetics of IPSCs were

dramatically reduced as indicated by significantly slower rise time and decay time in TKO mice

as compared to control (Figure 2c-d). Together, these data strongly suggest that neurexins are

essential for the intact function of the glycinergic synapse.

To further address the physiological role of neurexins in action potential evoked

neurotransmission at the glycinergic synapse, we applied electric stimulation to activate

presynaptic fibers of MNTB neurons by placing a stimulating electrode in the midway between

the MNTB and the LSO. Consistent with our optogenetic approach, afferent-evoked large

IPSCs were predominantly glycinergic as they can also be fully blocked by strychnine (Figure

S1). A modest but significant decrease in synaptic strength was observed in Nrxn123 TKO

mice as compared to control mice (Figure 2e-f), confirming the important role of neurexins in

the glycinergic synapses. The modest effect of the pan-neurexin deletion on synaptic strength

in the MNTB-LSO synapses is likely due to the incomplete AAV virus transfection in MNTN

neurons. Therefore, it is likely that a small fraction of MNTB neurons may still express

relatively normal expression of neurexins. It is worth pointing out that extensive efforts were

made to optimize our virus injection protocols by adjusting the injection parameters including

the needle angle, virus titers, and volume. On average, it was estimated that approximately

~80% neurons in the MNTB expressed GFP and thus infected by AAV-Cre. It is also possible

that other sources of glycinergic inputs other than the MNTB afferents are likely to present but

obscured in wild-type animals by the predominant glycinergic inputs from MNTB neurons

(Jalabi et al., 2013).

The mechanism of neurexins for regulating the MNTB-LSO glycinergic synapse

Since neurexins play multifaceted roles in regulating pre- and/or post-synaptic functions,

we explored the potential mechanisms by initially examining the paired-pulse ratio (PPR) of

IPSCs, which have been routinely used for identifying a presynaptic origin of regulation. As

shown in Figure 2h, IPSCs evoked by two consecutive stimuli at various intervals showed a

strong depression in both control and TKO mice. However, the PPRs were significantly

increased by deletion of neurexins (Figure 2i), suggesting that neurexins are required for

maintaining a high release probability of the glycinergic nerve terminal.

We then recorded and analyzed spontaneous IPSCs from LSO neurons. Compared to

control, there was no significant difference in both the amplitude and kinetics of sIPSCs

(Figure 3a-d) in TKO mice, indicating the normal content of synaptic vesicle as well as the

normal function of postsynaptic glycine receptors. Surprisingly, however, we found that

deletion of all neurexins caused a small but significant increase in the frequency of sIPSCs

(Figure 3a-b), which is opposite to previous reports of either no change or decrease in the

miniature frequency at glutamatergic or GABAergic synapse (Chen *et al.*, 2017; Luo *et al.*,

2020). To test whether the increase in sIPSC frequency may be caused by an increase in the

number of glycinergic synapses, we performed immunostaining of brainstem sections

containing the LSO with antibodies specific for GlyT2 or VGAT, synaptic markers for the

glycinergic neurons. Consistent with the increased frequency of sIPSCs, we found a significant

increase in the intensity of both GlyT2 (Figure 3e-f) and VGAT (Figure 3h-i) staining,

suggesting more abundant glycinergic terminals at the LSO of TKO mice. These data together

show that the pan-neurexin deletion in MNTB neurons has surprisingly increased the baseline

glycinergic neurotransmission likely by promoting the synapse formation/maintenance, which

is distinct from the phenotypes observed in glutamatergic and GABAergic neurons (Chen *et al.*,

2017; Luo *et al.*, 2021).

Deletion of neurexins impairs the tight coupling between voltage-gated Ca^{2+} channels

and synaptic vesicles at the glycinergic synapse

Because neurexins are shown to exert diverse impacts including specifying the function of

voltage-gated Ca^{2+} channels and their spatial clustering within the presynaptic active zone

(Chen *et al.*, 2017; Luo *et al.*, 2020; Missler *et al.*, 2003), we addressed these issues by first

exploring the effect of pan-neurexin deletion on the calcium sensitivity of neurotransmitter

release at the MNTB-LSO synapse. We perfused the brain slice with low Ca^{2+} (0.2 mM) ACSF

while recorded IPSCs continuously. In both control and Nrxn123 TKO mice, the perfusion of

low Ca^{2+} ACSF gradually decreased the amplitude of IPSCs (Figure 4a, c). However, the

reduction rate was much faster in TKO mice such that IPSC amplitudes dropped more than 50%

in TKO vs ~10% in control when extracellular $[Ca^{2+}]$ reached to approximately 1.5 mM (Figure

4b, d). The extracellular $[Ca^{2+}]$ blocking 50% of IPSCs were estimated as 1.1 mM and 1.5 mM

respectively for control and TKO mice, which are significantly different (Figure 4e, f),

suggesting that neurexins indeed are crucial for regulating calcium sensitivity of transmitter

release at the glycinergic synapse.

One of the critical factors affecting the calcium sensitivity of release is the spatial distance

between Ca^{2+} channels and synaptic vesicles (Eggermann et al., 2011; Luo *et al.*, 2020).

Additionally, neurexins have been shown to promote tight coupling of Ca^{2+} channels with

synaptic vesicle at the glutamatergic calyx of Held synapse (Luo *et al.*, 2020). Because the

MNTB-LSO glycinergic synapse is similar on multiple aspects of the functional characteristics,

being fast, robust, and precise, to the calyx of Held synapse, we tested whether neurexins are

also required for the tight coupling of Ca^{2+} channels with synaptic vesicle. We incubated the

brain slice with high-concentration EGTA-AM and continuously recorded IPSCs from individual

LSO neurons. As illustrated in Figure 5, EGTA-AM treatment slightly blocked IPSCs at control

synapses, suggesting the relative ineffectiveness of EGTA in interfering neurotransmitter

release (Figure 5a, b). By contrast, at the neurexin-deficient synapses, EGTA-AM treatment

remarkably impaired IPSCs (Figure 5c, d). On average, the blocking effects of EGTA at

neurexin-deficient synapses (53.2 ± 5.4%) was much stronger than control synapses (15.6 ±

4.3%). Therefore, consistent with previous studies at the calyx of Held synapse (Luo *et al.*,

2020), our data provide further support that neurexins are key synaptic organizers for

clustering voltage-gated Ca^{2+} channels tightly with synaptic vesicles at the glycinergic

synapse.

Discussion

Using the MNTB-LSO synapse as a model, here we provide first assessment of the role of

neurexins at glycinergic synapses that mediate prominent synaptic inhibition in the spinal cord,

brainstem and retina. Our findings in general strongly corroborate the idea that neurexins are

central presynaptic organizers in all major types of fast chemical synapses including

glutamatergic, GABAergic and glycinergic.

Specifically, we have shown that various neurexin genes express concurrently in these

glycinergic neurons but the expression level of each appears to vary significantly (Figure 1).

This finding is consistent with many studies in glutamatergic and GABAergic neurons, which

converge to show that multiple genes, isoforms, and splicing sites of neurexins normally

co-express in individual neurons and the expression profiles are highly cell-type specific and

brain region specific (Fuccillo *et al.*, 2015; Schreiner et al., 2014; Treutlein et al., 2014;

Uchigashima *et al.*, 2019). Distinct neurexins may interact separately and/or redundantly with

diverse signaling molecules to form specific molecular codes for shaping neuronal connectivity

and functional properties (Aoto et al., 2015; Aoto et al., 2013; Dai et al., 2019; Lin et al., 2023;

Lloyd et al., 2023; Sudhof, 2017).

We further show that neurexins are critical in controlling the strength and precise timing of

the glycinergic synaptic inhibition at LSO neurons by clustering Ca^{2+} channels tightly with

synaptic vesicles in presynaptic release sites (Figures 2, 4 and 5). These data strongly

corroborate two perspectives on the function of neurexins as central presynaptic organizers.

First, deletion of all neurexins not only weakens the glycinergic synaptic connectivity but also

impairs the temporal precision of synaptic inhibition, exactly as it does at the glutamatergic

calyx of Held synapse (Luo *et al.*, 2020). Since both the glutamatergic calyx of Held synapse

and the glycinergic MNTB-LSO synapse in the auditory brainstem are extremely strong and

fast (Beiderbeck et al., 2018; Grothe, 2003), the orchestrating role of neurexins in controlling

the timing and magnitude of their synaptic transmission is therefore desirable for the auditory

information processing. Secondly, neurexins execute their role at the glycinergic synapse

mainly by tightly organizing Ca^{2+} channels within the release sites, again exactly as it does at

the calyx of Held synapse (Luo *et al.*, 2020). Nanodomain coupling between Ca^{2+} channels

and synaptic vesicles has been found in many fast central synapses and is functionally crucial

for increasing the efficacy and speed of neurotransmission (Eggermann *et al.*, 2011). This

similar role of neurexins in clustering Ca^{2+} channels tightly at presynaptic active zones of not

only the glutamatergic synapse but also the glycinergic synapse prompts a shared mechanism

and further research on the potential molecular candidates.

On the other hand, our data surprisingly reveal that deletion of neurexins may promote

the formation/maintenance of the glycinergic synapse, as evidenced by the increase in the

frequency of spontaneous synaptic events measured at LSO neurons as well as the increase

in the glycinergic synaptic density in Nrxn123 TKO mice (Figure 3). Early studies of TKOs of

α-neurexins or β-neurexins lead to general conclusion that neurexins are not essential for the

formation/maintenance of excitatory and inhibitory synapses (Anderson *et al.*, 2015; Missler *et*

al., 2003). Moreover, deletion of all α-neurexins and β-neurexins has no impact on the

formation/maintenance of glutamatergic synapse (Luo *et al.*, 2020) and GABAergic synapse

made by somatostatin-positive interneurons (Chen *et al.*, 2017), but produces partial loss

(~30%) of GABAergic synapse made by parvalbumin-positive interneurons (Chen *et al.*, 2017).

More recent data even show that Nrxn2 unexpectedly restricts excitatory synapse formation in

the hippocampus but has no effect on the inhibitory synapse (Lin *et al.*, 2023). Our present

findings on the glycinergic synapse suggest a mix of pro- and anti-synaptogenic effects of

neurexins and may raise three possibilities. First, Nrxn2 may dominate its function at this

synapse during development and play an inhibitory role in the formation of glycinergic synapse,

similar as the previous report of Nrxn2 impact on the hippocampal excitatory synapse (Lin *et*

al., 2023). Second, deletion of neurexins impairs the strength of the MNTB-LSO synapse,

which may indirectly interfere with synaptic elimination process and therefore lead to an

increase in synaptic density. Third, the more abundant glycinergic synapses may arise from

the developmental compensation of synapse formation by other sources of glycinergic

neurons (Jalabi *et al.*, 2013). Strong reduction in the strength of the MNTB-LSO synapse

caused by selective deletion of neurexins may shift the winner of synaptic competition toward

other parallel but much weaker glycinergic inputs in normal animals.

Methods

Mouse breeding and virus injection.

All animal experiments were approved by the Institutional Animal Care and Use Committee at

Bioland Laboratory, Guangzhou National Laboratory, Guangzhou, China. Transgenic Nrxn123

cTKO mice of either sex were used for all experiments. Mice were housed at room

temperature on a 12 h light–dark cycle (7:00 to 19:00, light) with food and water freely

available.

To delete neurexins in MNTB neurons, we injected P0 mice with

AAV2/9-hSyn-EGFP-P2A-Cre-WPRE-hGH-Poly (Cat:PT-0156, BrainVTA Co., China). Mice

were anesthetized on ice and immediately transferred to a stereotaxic apparatus (RWD Life

Technology, China). Injection was achieved by connecting a glass pipette with tip diameter

20-30 μm to Nanoject III programmable nanoliter injector (Drummond Scientific Company,

USA). Bilateral injections were performed at the following coordinates: AP from the most

rostral point: 5.61 mm, ML: ±0.27 mm, DV: 3.58 mm. For optogenetics experiments, we

injected 200 nl (AAV2/9-hSyn-EGFP-P2A-Cre-WPRE-hGH-PolyA and

AAV2/9-EF1α-DIO-hChR2(H134R)-mCherry-WPRE-hGH-Poly, 1:1, Cat: PT-0002, BrainVTA

Co., China) or 100 nl of AAV2/9-hSyn-hChR2(H123R)-EYFP-WPRE-hGH-PolyA (Cat:

PT-1317, BrainVTA Co., China) into each-side MNTB. P13-14 mice of either sex were used for

most experiments including electrophysiology, FISH and immunohistochemistry.

Acute slice preparation for electrophysiology

Coronal brain slices containing both the MNTB and LSO were prepared similarly as described

previous. In brief, mice of P13-14 were decapitated; brains were rapidly isolated and fixed on

the cutting platform a vibratome (VT1200s; Leica, USA), which was immersed in oxygenated

cold ACSF containing (in mM): 119 NaCl, 26 NaHCO$_3$, 10 glucose, 1.25 NaH$_2$PO$_4$, 2.5 KCl,

0.05 CaCl$_2$, 3 MgCl$_2$, 2 Na-pyruvate, and 0.5 ascorbic acid, pH 7.4. Transverse slices of 250

μm were sectioned and transferred into a beaker with bubbled ACSF containing (in mM): 119

NaCl, 26 NaHCO3, 10 glucose, 1.25 NaH$_2$PO$_4$, 2.5 KCl, 2 CaCl$_2$, 1 MgCl$_2$, 2 Na-pyruvate, and

0.5 ascorbic acid, pH 7.4. Slices were recovered at 35°C for 30, and stored at room

temperature (~21-23°C) for experiments.

Whole-cell voltage-clamped recordings

Whole-cell voltage-clamped recordings were made from visually identified LSO neurons in

acute brain slices using an Axopatch 700B amplifier (Molecular Devices, USA) and pClamp 11

software (Molecular Devices, USA). Patch pipettes were pulled from borosilicate glass

capillaries using a two-stage vertical puller (PC-100, Narashige, Japan). Pipette resistance

was 4-6 MΩ when filled with an internal solution containing (in mM): 145 K-gluconate, 6 KCl,

10 HEPES, 3 Na_2-phosphocreatine, 4 Mg-ATP, 0.3 Na_2-GTP, 0.2 EGTA, 2 Qx-314; pH 7.2

with KOH. With the extracellular $[Cl^-]$ of 133 mM and a temperature of 22°, the equilibrium

potential of the Cl^- was approximately −78.8 mV. In voltage-clamp configuration, IPSCs were

recorded by holding cells at 10 mV in ACSF containing CNQX (20 μM), D-AP5 (50 μM), to

block AMPA, NMDA receptors, respectively. For some experiments, 2 μM strychnine, a potent

glycine receptor antagonist, was added to ACSF to verify that IPSCs recorded at LSO neurons

are glycinergic. Membrane potentials were not corrected for liquid junction potential. Electrical

fiber stimulation was performed using a concentric bipolar electrode (CBAEB75, FHC Inc, USA)

positioned medially to the LSO and MNTB. Short pulses (0.1 ms) were delivered via an

isolated stimulator (Model 2100; A-M Systems, USA). For optogenetics, a light simulation of 5

ms was delivered by LED (pE-300 white, CoolLED, UK).

Whole-cell current-clamp recordings

317 Whole-cell current-clamped recordings were performed similarly as voltage-clamped

318 recordings, except that the internal solution did not contain Qx-314. Action potentials at MNTB

319 neurons were recorded by holding the membrane potential at approximately −70 mV. For

320 optogenetics, a light simulation of 5 ms was delivered by LED (pE-300 white, CoolLED, UK).

RNAscope fluorescent in-situ hybridization (FISH) and immunohistochemistry

Mice were anesthetized and perfused at speed of 1ml/min with 1×PBS for 10 min followed by 4%

paraformaldehyde (PFA) for 10 min. Brains were removed and stored in 4% PFA overnight at

4C. After going through a 10%, 20%, and 30% sucrose gradient for cryo-protection, the tissue

was embedded in OCT (Cat : 14020108926, Leica) and rapidly frozen on dry ice. Transverse

sections at 16 μm were cut at -20 °C using a cryostat (CM3050-S, Leica) and mounted directly

on Superfrost Plus slides (Cat: PRO-04, Matsunami Platinum Pro). FISH and

immunohistochemistry were performed using the RNAscope multiplex platform (Multiplex

Reagent Kit, Cat: 323100; Co-detection ancillary Kit, Cat: 323180, Advanced Cell Diagnostics)

following the manufacturer's instructions. RNA probes for Nrxn1, Nrxn2, and Nrxn3 (Cat:

461511-C3, 533531-C2, 505431 respectively, Advanced Cell Diagnostics) and primary

antibody against VGluT1 (guinea pig, polyclonal, 1:500, Millipore) were used. Secondary

antibodies were Alexa Fluor conjugates (1:400; Thermo Fisher). Samples were mounted with

a Vectashield hard-set antifade mounting medium (Cat: H-1500, Vector Laboratories). Images

were acquired using Zeiss LSM800 confocal microscopy with a 63x oil-immersion objective

(1.4 numerical aperture) or OLYMPUS IXplore SpinSR confocal microscopy with a 60x

oil-immersion objective (1.5 numerical aperture). Image analysis were performed using Zeiss

2.6, cellSens Dimension 3.2 and Image J.

Pharmacological manipulations

To estimate the spatial relationship between voltage-gated Ca^{2+} channels and synaptic

vesicles, we perfused the slice with ACSF containing 20 μM EGTA-AM for 25~30 mins while

continuously recording eIPSC. The blocking effect of EGTA on eIPSCs were evaluated by

comparing eIPSCs before and after EGTA treatment.

Statistical analysis

Data were analyzed in the IgorPro (Version 8) and GraphPad Prism 9 software. Summary of

data was displayed as mean ± SEM. Statistical analysis was performed using Student's *t* test,

two-way ANOVA, and Mann-Whitney U test. Significance of difference was accepted at P <

0.05.

References

Anderson, G.R., Aoto, J., Tabuchi, K., Foldy, C., Covy, J., Yee, A.X., Wu, D., Lee, S.J., Chen, L.,
Malenka, R.C., and Sudhof, T.C. (2015). beta-Neurexins Control Neural Circuits by Regulating
Synaptic Endocannabinoid Signaling. Cell *162*, 593-606. 10.1016/j.cell.2015.06.056.
Aoto, J., Foldy, C., Ilcus, S.M., Tabuchi, K., and Sudhof, T.C. (2015). Distinct circuit-dependent
functions of presynaptic neurexin-3 at GABAergic and glutamatergic synapses. Nat Neurosci
18, 997-1007. 10.1038/nn.4037.
Aoto, J., Martinelli, D.C., Malenka, R.C., Tabuchi, K., and Sudhof, T.C. (2013). Presynaptic
neurexin-3 alternative splicing trans-synaptically controls postsynaptic AMPA receptor
trafficking. Cell *154*, 75-88. 10.1016/j.cell.2013.05.060.
Assareh, N., Fenech, C., Power, R., Uddin, M.N., Otsu, Y., and Aubrey, K.R. (2023).
Bidirectional Modulation of Nociception by GlyT2(+) Neurons in the Ventrolateral
Periaqueductal Gray. eNeuro *10*. 10.1523/ENEURO.0069-23.2023.
Beiderbeck, B., Myoga, M.H., Muller, N.I.C., Callan, A.R., Friauf, E., Grothe, B., and Pecka, M.
(2018). Precisely timed inhibition facilitates action potential firing for spatial coding in the
auditory brainstem. Nat Commun *9*, 1771. 10.1038/s41467-018-04210-y.
Boxer, E.E., and Aoto, J. (2022). Neurexins and their ligands at inhibitory synapses. Front
Synaptic Neurosci *14*, 1087238. 10.3389/fnsyn.2022.1087238.
Brill, S.E., Maraslioglu, A., Kurz, C., Kramer, F., Fuhr, M.F., Singh, A., and Friauf, E. (2020).
Glycinergic Transmission in the Presence and Absence of Functional GlyT2: Lessons From
the Auditory Brainstem. Front Synaptic Neurosci *12*, 560008. 10.3389/fnsyn.2020.560008.
Chang, Z., Skach, J., and Kam, K. (2023). Inhibitory subpopulations in preBotzinger Complex
play distinct roles in modulating inspiratory rhythm and pattern. bioRxiv.
10.1101/2023.08.07.552303.
Chen, L.Y., Jiang, M., Zhang, B., Gokce, O., and Sudhof, T.C. (2017). Conditional Deletion of
All Neurexins Defines Diversity of Essential Synaptic Organizer Functions for Neurexins.
Neuron *94*, 611-625 e614. 10.1016/j.neuron.2017.04.011.
Dai, J., Aoto, J., and Sudhof, T.C. (2019). Alternative Splicing of Presynaptic Neurexins
Differentially Controls Postsynaptic NMDA and AMPA Receptor Responses. Neuron *102*,
993-1008 e1005. 10.1016/j.neuron.2019.03.032.
Eggermann, E., Bucurenciu, I., Goswami, S.P., and Jonas, P. (2011). Nanodomain coupling
between Ca(2)(+) channels and sensors of exocytosis at fast mammalian synapses. Nat Rev
Neurosci *13*, 7-21. 10.1038/nrn3125.
Feng, G., Tintrup, H., Kirsch, J., Nichol, M.C., Kuhse, J., Betz, H., and Sanes, J.R. (1998).
Dual requirement for gephyrin in glycine receptor clustering and molybdoenzyme activity.
Science *282*, 1321-1324. 10.1126/science.282.5392.1321.
Fischer, A.U., Muller, N.I.C., Deller, T., Del Turco, D., Fisch, J.O., Griesemer, D., Kattler, K.,
Maraslioglu, A., Roemer, V., Xu-Friedman, M.A., et al. (2019). GABA is a modulator, rather
than a classical transmitter, in the medial nucleus of the trapezoid body-lateral superior olive
sound localization circuit. J Physiol *597*, 2269-2295. 10.1113/JP277566.
Foldy, C., Darmanis, S., Aoto, J., Malenka, R.C., Quake, S.R., and Sudhof, T.C. (2016).
Single-cell RNAseq reveals cell adhesion molecule profiles in electrophysiologically defined
neurons. Proc Natl Acad Sci U S A *113*, E5222-5231. 10.1073/pnas.1610155113.
Foster, E., Wildner, H., Tudeau, L., Haueter, S., Ralvenius, W.T., Jegen, M., Johannssen, H.,
Hosli, L., Haenraets, K., Ghanem, A., et al. (2015). Targeted ablation, silencing, and activation

establish glycinergic dorsal horn neurons as key components of a spinal gate for pain and itch.
Neuron *85*, 1289-1304. 10.1016/j.neuron.2015.02.028.

Fuccillo, M.V., Foldy, C., Gokce, O., Rothwell, P.E., Sun, G.L., Malenka, R.C., and Sudhof, T.C.
(2015). Single-Cell mRNA Profiling Reveals Cell-Type-Specific Expression of Neurexin
Isoforms. Neuron *87*, 326-340. 10.1016/j.neuron.2015.06.028.

Gomez, A.M., Traunmuller, L., and Scheiffele, P. (2021). Neurexins: molecular codes for
shaping neuronal synapses. Nat Rev Neurosci *22*, 137-151. 10.1038/s41583-020-00415-7.

Gomeza, J., Hulsmann, S., Ohno, K., Eulenburg, V., Szoke, K., Richter, D., and Betz, H.
(2003a). Inactivation of the glycine transporter 1 gene discloses vital role of glial glycine uptake
in glycinergic inhibition. Neuron *40*, 785-796. 10.1016/s0896-6273(03)00672-x.

Gomeza, J., Ohno, K., Hulsmann, S., Armsen, W., Eulenburg, V., Richter, D.W., Laube, B., and
Betz, H. (2003b). Deletion of the mouse glycine transporter 2 results in a hyperekplexia
phenotype and postnatal lethality. Neuron *40*, 797-806. 10.1016/s0896-6273(03)00673-1.

Grothe, B. (2003). New roles for synaptic inhibition in sound localization. Nat Rev Neurosci *4*,
540-550. 10.1038/nrn1136.

Grothe, B., Pecka, M., and McAlpine, D. (2010). Mechanisms of sound localization in
mammals. Physiol Rev *90*, 983-1012. 10.1152/physrev.00026.2009.

Jalabi, W., Kopp-Scheinpflug, C., Allen, P.D., Schiavon, E., DiGiacomo, R.R., Forsythe, I.D.,
and Maricich, S.M. (2013). Sound localization ability and glycinergic innervation of the superior
olivary complex persist after genetic deletion of the medial nucleus of the trapezoid body. J
Neurosci *33*, 15044-15049. 10.1523/JNEUROSCI.2604-13.2013.

Kim, G., and Kandler, K. (2003). Elimination and strengthening of glycinergic/GABAergic
connections during tonotopic map formation. Nat Neurosci *6*, 282-290. 10.1038/nn1015.

Kotak, V.C., Korada, S., Schwartz, I.R., and Sanes, D.H. (1998). A developmental shift from
GABAergic to glycinergic transmission in the central auditory system. J Neurosci *18*,
4646-4655. 10.1523/JNEUROSCI.18-12-04646.1998.

Krachan, E.G., Fischer, A.U., Franke, J., and Friauf, E. (2017). Synaptic reliability and
temporal precision are achieved via high quantal content and effective replenishment: auditory
brainstem versus hippocampus. J Physiol *595*, 839-864. 10.1113/JP272799.

Lin, P.Y., Chen, L.Y., Jiang, M., Trotter, J.H., Seigneur, E., and Sudhof, T.C. (2023). Neurexin-2:
An inhibitory neurexin that restricts excitatory synapse formation in the hippocampus. Sci Adv
9, eadd8856. 10.1126/sciadv.add8856.

Lloyd, B.A., Han, Y., Roth, R., Zhang, B., and Aoto, J. (2023). Neurexin-3 subsynaptic
densities are spatially distinct from Neurexin-1 and essential for excitatory synapse nanoscale
organization in the hippocampus. Nat Commun *14*, 4706. 10.1038/s41467-023-40419-2.

Lukacsovich, D., Winterer, J., Que, L., Luo, W., Lukacsovich, T., and Foldy, C. (2019).
Single-Cell RNA-Seq Reveals Developmental Origins and Ontogenetic Stability of Neurexin
Alternative Splicing Profiles. Cell Rep *27*, 3752-3759 e3754. 10.1016/j.celrep.2019.05.090.

Luo, F., Sclip, A., Jiang, M., and Sudhof, T.C. (2020). Neurexins cluster Ca(2+) channels within
the presynaptic active zone. EMBO J *39*, e103208. 10.15252/embj.2019103208.

Luo, F., Sclip, A., Merrill, S., and Sudhof, T.C. (2021). Neurexins regulate presynaptic
GABAB-receptors at central synapses. Nat Commun *12*, 2380. 10.1038/s41467-021-22753-5.

Missler, M., Zhang, W., Rohlmann, A., Kattenstroth, G., Hammer, R.E., Gottmann, K., and
Sudhof, T.C. (2003). Alpha-neurexins couple Ca2+ channels to synaptic vesicle exocytosis.

Nature *423*, 939-948. 10.1038/nature01755.

Nabekura, J., Katsurabayashi, S., Kakazu, Y., Shibata, S., Matsubara, A., Jinno, S., Mizoguchi,
Y., Sasaki, A., and Ishibashi, H. (2004). Developmental switch from GABA to glycine release in
single central synaptic terminals. Nat Neurosci *7*, 17-23. 10.1038/nn1170.

Sanes, D.H., and Friauf, E. (2000). Development and influence of inhibition in the lateral
superior olivary nucleus. Hear Res *147*, 46-58. 10.1016/s0378-5955(00)00119-2.

Schreiner, D., Nguyen, T.M., Russo, G., Heber, S., Patrignani, A., Ahrne, E., and Scheiffele, P.
(2014). Targeted combinatorial alternative splicing generates brain region-specific repertoires
of neurexins. Neuron *84*, 386-398. 10.1016/j.neuron.2014.09.011.

Sudhof, T.C. (2008). Neuroligins and neurexins link synaptic function to cognitive disease.
Nature *455*, 903-911. 10.1038/nature07456.

Sudhof, T.C. (2017). Synaptic Neurexin Complexes: A Molecular Code for the Logic of Neural
Circuits. Cell *171*, 745-769. 10.1016/j.cell.2017.10.024.

Treutlein, B., Gokce, O., Quake, S.R., and Sudhof, T.C. (2014). Cartography of neurexin
alternative splicing mapped by single-molecule long-read mRNA sequencing. Proc Natl Acad
Sci U S A *111*, E1291-1299. 10.1073/pnas.1403244111.

Uchigashima, M., Cheung, A., Suh, J., Watanabe, M., and Futai, K. (2019). Differential
expression of neurexin genes in the mouse brain. J Comp Neurol *527*, 1940-1965.
10.1002/cne.24664.

Ullrich, B., Ushkaryov, Y.A., and Sudhof, T.C. (1995). Cartography of neurexins: more than
1000 isoforms generated by alternative splicing and expressed in distinct subsets of neurons.
Neuron *14*, 497-507. 10.1016/0896-6273(95)90306-2.

Ushkaryov, Y.A., Petrenko, A.G., Geppert, M., and Sudhof, T.C. (1992). Neurexins: synaptic
cell surface proteins related to the alpha-latrotoxin receptor and laminin. Science *257*, 50-56.
10.1126/science.1621094.

Zeilhofer, H.U., Wildner, H., and Yevenes, G.E. (2012). Fast synaptic inhibition in spinal
sensory processing and pain control. Physiol Rev *92*, 193-235. 10.1152/physrev.00043.2010.

ACKNOWLEDGEMENTS

We thank Dr. B. Zhang (Shenzhen Bay Laboratory) for helpful comments on the manuscript.

This study was supported by grants from the National Natural Science Foundation of China

(31970914 to F.L.; 32171001, 32371050 to Y.Z.).

AUTHOR CONTRIBUTIONS

Conceptualization, F.L.; Experiments, H.J., R.X., X.N., Z.S., X.X., R. P.; Data analysis, H.J.,

R.X., X.N., Z.S., X.X., Y.Z, F.L.; Writing – Original Draft, F.L.; Writing – Review & Editing, H.J,

Y.Z, F.L.

DECLARATION OF INTEREST

The authors declare no competing financial interests.

Figure legends

Fig.1 Neurexins are highly expressed in MNTB neurons. (**a**) Immunohistochemistry and

RNAscope FISH showing expression of various isoforms of neurexins in the MNTB, which

make glycinergic synapses with LSO neurons (**b**) Summary of expression of various isoforms

of neurexins in the MNTB of WT mice. (**c**) Diagram of auditory brainstem and virus injection in

Nrxn123 cTKO mice. (**d**) Representative images of Nrxn3 FISH in the MNTB of mice injected

with AAV-EGFP or AAV-Cre-EGFP. Scale bar: 50 μm. (**e**) Selected ROI in high magnification

showing Nrxn3 FISH in MNTB neurons expressing EGFP or Cre-EGFP. Scale bar: 10 μm. (**f**)

Summary of expression of various neurexins in the MNTB of Nrxn123 cTKO mice injected with

AAV-EGFP and AAV-Cre-EGFP.

Data are means ± SEM. Number of sections analyzed are indicated in the bars (**b**, **f)**;

Statistical differences were assessed by Student's t test. (**P < 0.01, ***P < 0.001). Source

data are provided as a Source Data file.

Figure 2 Deletion of neurexins reduces the strength and kinetics of glycinergic synapse.

(**a**) Representative IPSC traces of LSO neurons evoked by optogenetic stimulation of ChR2

expressed in MNTB neurons. (**b-d**) Summary of IPSC amplitude, rise time, and decay time

constant. P = 0.0119, P = 0.0004, P = 0.002. unpaired two-sided t-test. (**e-h**) Same as a-d,

except that IPSCs were evoked by afferent fiber stimulation. (**i**) Representative IPSCs traces

of LSO neurons evoked by a pair of fiber stimulation at 50 ms interval. (**j**) Summary of

paired-pulse ratio (PPR) of IPSCs in relationship to different paired pulse intervals. P < 0.0001,

two-way ANOVA.

Data are means ± SEM. Number of neuron are indicated in the bars (**b-d**, **f-h**) or graph (**j**).

Statistical differences were assessed by Student's t test or two-way ANOVA test. (*P < 0.05,

**P < 0.01, ***P < 0.001).

Figure 3 Deletion of neurexins increases sIPSC frequency and glycinergic synapse

density of LSO neurons. (**a**) Representative sIPSC traces of LSO neurons. (**b**) Summary of

sIPSC amplitude and frequency. P = 0.057, P = 0.015, unpaired two-sided t-test. (**c**)

Representative sIPSC waveforms of LSO neurons. (**d**) Summary of sIPSC rise time and decay

time constant P = 0.376, P = 0.82, unpaired two-sided t-test. (**e**) Representative confocal

microscopy images of MNTB-LSO-containing brainstem slice with specific labeling of GlyT2

(red) and EGFP (green) from both control and Nrxn123 TKO mice at P14. Scale bar, 10 μm. (**f**)

Summary of GlyT2 immunostaining intensity (normalized to control). P = 2.49E-07, unpaired

two-sided t-test. (**g-h**), Same as (**e-f**) except for specific labeling of VGAT (red) and EGFP

(green). P = 0.002, unpaired two-sided t-test.

Data are means ± SEM. Number of neuron or sections/animals for immunostaining are

indicated in the bars (**b**, **d**, **f** and **h**). Statistical differences were assessed by Student's t test.

(*P < 0.05, **P < 0.01, ***P < 0.001). Source data are provided as a Source Data file.

Figure 4 Deletion of neurexins increases the Ca^{2+} sensitivity of release at the

glycinergic synapse. (**a**) Representative time courses of IPSC amplitudes for individual LSO

neurons of control mice during the perfusion of low Ca^{2+} ACSF. (**b**) Representative IPSC

traces of control mice before and during low Ca^{2+} perfusion at the indicated points in **a**. (**c** & **d**)

Same as **a** & **b**, except of TKO mice. (**e**) Summary of normalized IPSC amplitudes plotted

against the calculated extracellular Ca^{2+} concentration for control and TKO mice. P = 0.0311

for 0.92 mM Ca^{2+}, P = 0.0104 for 1.10 mM Ca^{2+}, P = 0.0208 for 1.28 mM Ca^{2+}, P = 0.0067 for

1.46 mM Ca^{2+}, P = 0.0007 for 1.64 mm Ca^{2+}, unpaired two-sided t-test. (**f**) Summary of IC_{50} of

Ca^{2+} for half-blocking of IPSCs and the blocking effect of IPSCs by 0.2 mM Ca^{2+}. P = 0.0139, P

= 0.3985, unpaired two-sided t-test.

Data are means ± SEM. Number of neuron are indicated in the graph (**e**) or bars (**f**). Statistical

differences were assessed by Student's t test. (*P < 0.05, **P < 0.01, ***P < 0.001). Source

data are provided as a Source Data file.

Figure 5 Deletion of neurexins increases the blocking effect of EGTA on glycinergic

IPSCs. (**a**) Representative time courses of IPSC amplitudes for individual LSO neurons of

control mice during the treatment of high concentration of EGTA-AM. (**b**) Representative IPSC

traces of control mice before and after EGTA. (**c, d**) Same as a, b, except of TKO mice. (**e**)

Summary of IPSC amplitude before and after EGTA treatment of control mice. P = 0.0201,

paired two-sided t-test. (**f**) Summary of IPSC amplitude before and after EGTA treatment of

TKO mice. P = 0.0123, paired two-sided t-test. (**g**) Summary of the blocking percentage of

IPSCs by EGTA. P = 0.0001, unpaired two-sided t-test.

Data are means ± SEM. Number of neuron are indicated in the bars (**e** to **g**). Statistical

differences were assessed by Student's t test. (*P < 0.05, ***P < 0.001). Source data are

provided as a Source Data file.

550
551

552 **Figure 1**

556 **Figure 2**

561 **Figure 3**

564 **Figure 4**

568   **Figure 5**

Figure S1

a Representative sIPSC traces of LSO neurons before and after application of potent glycine receptor blocker strychine (stry) (left) and summary of sIPSC amplitude without and with stry (right).
b Representative eIPSC traces of LSO neurons before and after stry application (left) and summary of eIPSC amplitude without and with stry (right).
c Representative action potential (AP) firing of MNTB neurons infected with ChR2 during high frequency repetitive pulse stimulation of 473 nm.
d Representative oIPSC traces of LSO neurons before and after stry application (left) and summary of oIPSC amplitude without and with stry (right).

575
576

Figure s2

a Summary of eIPSC amplitude of control during 0.2 mM Ca^{2+} perfusion. Mean ± SEM.
b Summery of eIPSC amplitude of TKO during 0.2 mM Ca^{2+} perfusion. Mean ± SEM.
c Average of eIPSC amplitude before and after 0.2 mM Ca^{2+} perfusion. Mean ± SEM.
(**P < 0.01, ***P < 0.001), paired t-test.

577
578

a Summary of eIPSC amplitude of control during 10 μM EGTA-AM treatment. Mean ± SEM.
b Summary of eIPSC amplitude of TKO during 10 μM EGTA-AM treatment. Mean ± SEM