Chromosome-level genome assembly for Sichuan taimen (*hucho bleekeri*) reveals the extraordinary tandem repeat proportions and its persistent population shrinkage

Xinmiao Zhang[1], Dongmei Xiong[1], Shenglong Jian[2], Yu Jiang[3*], Lixin Wang[1*]

[1]College of Animal Science and Technology, Northwest A&F University, Yangling, 712100, China

[2]Qinghai Provincial Fishery Technology Extension Center, Xining, China

[3]Center for Ruminant Genetics and Evolution, College of Animal Science and Technology, Northwest A&F University, Yangling 712100, China

*Correspondence authors. E-mail: fisherwanglx@nwsuaf.edu.cn (Lixin Wnag) and yu.jiang@nwafu.edu.cn (Yu Jiang).

ABSTRACT:

Salmonid fishes are globally renowned and valuable, with most members of the Salmonidae family living in seawater and exhibiting migratory behavior. In contrast, huchonid fishes (*Hucho* spp.and *Brachymystax* spp.), an ancient evolutionary branch within Salmoninae, are entirely landlocked. The Sichuan taimen (*Hucho bleekeri* Kimura) is a critically endangered fish that has attracted widespread concern and is one of eleven national first-level protected fishes in China. However, genome resources for all *Hucho* spp., including *H. bleekeri*, are scarce, and the genomic characteristics of this ancient evolutionary lineage remain unclear, hindering conservation biology efforts. Here, we present the first chromosome-level genome for the Sichuan taimen, with a final genome size of approximately 3.45 Gb across 44 chromosomes. The Sichuan taimen genome contains 44.15% tandem repetitive sequences, exceeding those of all sequenced salmonid fishes. We also identified 44 Ss4R homeologous block pairs in the assembled genome. Genome synteny analysis suggested a ~5 fold tandem repeat expansion in the Sichuan taimen compared to the Salmonidae ancestor Northern pike. Phylogenetic analysis estimated the divergence time between huchonid and other Salmoninae fishes at approximately 33.29 million years ago (Mya). The divergence time between Siberia taimen and Sichuan taimen was estimated at around 2.29 Mya, with their effective population size declining from around 1 Mya. The genomic resource provided in this article will promote the protection of the Sichuan taimen and evolutionary genetics studies of salmonids.

Keywords: *Hucho bleekeri*; Genome assembly; Tandem repeats; Salmonid

INTRODUCTION

The Sichuan taimen (*Hucho bleekeri* Kimura, 1934) belonging to the subfamily Salmoninae, is a glacial remnant species and endemic to China (Wang et al., 2016). It has the lowest distribution latitude among all taimen species, and is primarily found in the upper tributaries of the Yangtze River and Wei River (Hu et al., 2008). Sichuan taimen, characterized by their large body size and predatory behavior, occupy the top of the aquatic food chain (Wang et al., 2016). However, their late sexual maturation makes them particularly vulnerable to human activities and ecological changes (Allendorf & Hard, 2009; Sadovy et al., 2013). Individuals of Sichuan taimen are found sporadically and scattered in their distribution, and have been listed in China red data book of endangered animals since 1998. Despite this, they continues to be illegally captured by poachers for their high nutritional value and taste (Sung et al., 1998). Due to the continuous decline of the population size, the Sichuan taimen was listed as critically endangered (CR) level in the International Union for Conservation of Nature (IUCN) Red List of Threatened Species in 2012 (Song, 2012). As of 2021, the Sichuan taimen is under the highest level of protection from the Chinese government, and is the only salmonid fish with national protected animal in "class I". Genomic data provide significant opportunities for advancements in the conservation and evolutionary research of threatened animals (Allendorf et al., 2010; Kohn et al., 2006). As sequencing costs have decreased, the genomes of many endangered fish species have been published, including *Trachidermus fasciatus* Heckel (Xue et al., 2022), *Myxocyprinus asiaticus* (Liu et al., 2022), and *Brachymystax tsinlingensis* (Zhu

et al., 2022), etc. Based on genome data, scientific and rational conservation policies have been formulated, enhancing the protection benefits of these endangered animals (Brandies et al., 2019; Waples et al., 2020). However, the lack of genome data for *Hucho bleekeri* creates a gap in nuclear-based phylogenetic research of Salmonidae and affects genomic studies, such as taxonomic studies based on SNPs (Chen et al., 2020; Zhang et al., 2020) and researches on transcriptome (Chen et al., 2021).

Due to the whole genome duplication and lineage-specific rediploidization, the genome of salmonids have became complex and variable, increasing the difficulty of assembly (Dysin et al., 2022; Robertson et al., 2017). The circular consensus sequencing (CCS) mode in single-molecule real-time (SMRT) sequencing by Pacific Biosciences can produce long and high fidelity (HiFi) reads, which facilitates the genome assembly process and improves quality (Wenger et al., 2019). In this research, we utilized HiFi and Hi-C sequencing technology to construct a high-quality genome for Sichuan taimen. From the genome we identified a high proportion of tandem repetitive sequences. Additionally, we analyzed the phylogenetic relationships of salmonid fishes and the divergence history of *H. bleekeri* and *H. taimen*, another species found in Heilongjiang. Our work provides insights into the genome evolution of salmonid fishes and the conservation of the endangered Sichuan taimen.

MATERIALS AND METHODS

Sample collection and sequencing

Two wild individuals of *H. bleekeri* were collected from Taibai river (Baoji, Shaanxi province), one for genome assembly the other for resequencing. Additionally, a

cultured individual of *H. taimen* was collected from Heilongjiang. All the experiments

in this study were performed in accordance with the guidelines of China Council on

Animal Care, and approved by the Experimental Animal Ethics Committee of

Northwest A&F University (NWAFU-314020038).

Genomic DNA was extracted from kidney tissue with cetyltrimethylammonium

bromide (CATB) method (Stefanova et al., 2013). The reads for genome size

estimation were generated on Illumina NovaSeq-6000 platform after the constructed

of 150 bp library. For genome assembly, a 15 kb library preparation was performed by

SMRTbell Express Template Prep Kit 2.0 (Pacific Biosciences, CA, USA), and

sequenced on the PacBio Sequel II platform (Pacific Biosciences, CA, USA) with two

cells of single molecule real time sequencing (SMRT). The CCS software v.6.0.0

(https://github.com/PacificBiosciences/ccs) was performed to integrate HiFi reads

with the parameters "--min-passes 3 --min-length 10 --min-rq 0.99". The Hi-C library

was constructed and sequenced on Illumina NovaSeq-6000 platform. The RNA

samples were extracted from fin，brain, gill, gonad, and eye, then sequenced on

Illumina NovaSeq-6000 platform. The two wild samples for demographic history

analysis were sequenced on DNB-seq platform (BGI).

Genome size estimation

The raw Illumina paired-end reads of Sichuan taimen were filtered using Fastp

version 0.20.0 (Chen et al., 2018) with default parameters. The genome size

estimation of Sichuan taimen was performed by GCE software version 1.0.2 (Liu et

al., 2013). A total of 58.89 Gb clean reads were used to calculate the distribution of

k-mer frequency using kmerfreq with k=17. Then the gce software was ran in both

homozygous and heterzygous mode to evaluate the genome size and heterozygosity.

The genome heterozygous was calculated using the formula "a[1/2]/k".

Chromosome level genome assembly

The PacBio HiFi reads were assembled using Hifiasm version 0.13-r308 (Cheng et al.,

2021) to obtain *de novo* contigs. After quality control by Fastp (Chen et al., 2018), the

clean Hi-C reads were used to integrate the contigs. The Hi-C interaction matrix was

generated using Juicer (version 1.6)(Durand et al., 2016b)scripts with -s MboI

parameter, then, the 3D-DNA (version 201008)(Dudchenko et al., 2017)with 0 rounds

was used to link the contigs into chromosome level based on the Hi-C interaction

matrix. The initial assembly was manually corrected according to the Hi-C heatmap

using Juicebox version 1.11.08 (Durand et al., 2016a) and run the 3D-DNA again to

generate the final assembly. The BUSCO (Benchmarking universal single-copy

orthologues) version 3.0.2 (Simão et al., 2015) was used to assess the accuracy and

integrity of the genome with the gene database of ray-finned fishes

(actinopterygii_odb9).

Repeat sequence detection and gene annotation

To recognize repeat sequences of the Sichuan taimen genome, we build a *de novo*

library of transposable elements (TEs). First, the RepeatModeler version 2.1 (Flynn et

al., 2020) with default parameters was ran to constructed a repeat library; then an

additional long terminal repeat (LTR) library of the Sichuan taimen was built by

LTR_FINDER version 1.07 (Xu & Wang, 2007) and LTR_RETRIEVER version 2.9.6

(Ou & Jiang, 2018); third the final repeat library was integrated by replacing the LTR elements of the first repeat library with the LTR library. To further classify the "Unknown" sequence, the BLAST version 2.14.0 (Ye et al., 2006) was performed between the final repeat library and public repeat library (http://www.hrt.msu.edu/uploads/535/78637/Tpases020812.gz).

The TEs of the genome were described and masked by RepeatMasker version 4.1.2 (Tarailo-Graovac & Chen, 2009) with parameters "-xsmall -s -cutoff 255 -frag 20000 –gff -engine ncbi". The tandem repeats (TRs) were detected through Tandem Repeats Finder (TRF) version 4.09 (Benson, 1999) with parameters "2 7 7 80 10 50 500 -f -d -m". The TRs were further divided into short tandem repeats (STRs) and variable number tandem repeats (VNTRs) according to the length of repeat unit (STR: unit number at 1 to 6; VNTR: unit number more than 6).

The gene annotation of the Sichuan taimen genome was carried out using BRAKER2 pipeline version 2.1.5 (Hoff et al., 2016; Hoff et al., 2019). For transcriptome based prediction, the RNAs were mapping to the assembled genome using STAR version 2.7.1a (Dobin et al., 2013) with default parameters, then the RNA sequence data were used to predict genes through auto BRAKER2 pipeline. For homology based prediction, the protein sequence of six species including *Lepisosteus oculatus* (GCF_000242695.1)(Braasch et al., 2016), *Esox Lucius* (GCF_011004845.1)(Ishiguro et al., 2003), *Oncorhynchus mykiss* (GCF_013265735.2)(Gao et al., 2021), *Oncorhynchus nerka* (GCF_006149115.2) , *Salmo salar* (GCF_000233375.1)(Lien et al., 2016), and *Brachymystax tsinlingensis* (GCA_031305455.1) which represent the

sister clade or the ancestor species of taimen were download, and predicted through

BRAKER2 pipeline. The results from RNA and protein-based annotation were

combined using TSEBRA version 1.0.3 (Gabriel et al., 2021) with default parameters

to generate the final gene prediction file. The gene expression level was measured by

transcript per million (TPM) which were calculated using Kallisto (Bray et al., 2016)

with default parameters. To remove the pseudogenes caused by tandem repetitive

sequence, the gene set was filtered based on the following criteria: 1) contain only one

exon; 2) TPM<1; 3) no matched BLAST results with *B. tsinlingensis*.

Chromosome synteny identification

Homology blocks were reconstructed using whole-genome duplicated genes.

Diamond (v0.8.24.86) with default parameter was used to align against the predicted

genes. Then selected paired genes with highest identity and identity $\geq$ 75% from the

Diamond results. Finally, one-to-one paralogs were extracted and conducted manual

linkage. The *Esox lucius* genome was download from NCBI (GCF_011004845.1).

Phylogenetic analysis

Protein sequences of 10 species *Esox lucius*, *Coregonus* sp. *'balchen'*

(GCA_902810595.1)(Christensen et al., 2020), *Thymallus thymallus*

(GCA_023634145.1), *Oncorhynchus mykiss*, *Oncorhynchus nerka*, *Salvelinus*

namaycush (GCF_016432855.1)(Smith et al., 2022), *Salmo salar*, *Salmo trutta*

(GCF_901001165.1)(Hansen et al., 2021), *Hucho bleekeri*, and *Brachymystax*

tsinlingensis were used to construct phylogenetic tree. The single copy orthologous

proteins among the 10 species were recognized by Orthfinder version 2.3.12 (Emms

& Kelly, 2019) with "msa" method. After alignment using MUSCLE version 3.8.31

(Edgar, 2004), the species tree was calculated through RAxML version 8.2.12

(Stamatakis, 2014) with 1000 bootstraps. The expansions and contractions of gene

family was estimated using CAFE5 (Mendes et al., 2020).

The divergence time among all these species were calculated using MCMCtree

program of PAML software version 4.9d (Yang, 2007). Four fossil records were

downloaded from TimeTree (Kumar et al., 2022) and calibrated the results, the

Markov chain Monte Carlo (MCMC) was sampled with every 1,000 step and conduct

10,000 samples, the pre-burnin number was set at 20,000.

175 Divergence time estimation of two taimen species

The raw resequencing data of *H. taimen* was mapping to the assembled genome using

BWA-MEM version 0.7.17 after quality control. The whole genome sequence for *H.*

taimen was generated using angsd version 0.918. To obtain the gene annotation

information of *H. taimen*, the liftoff (v1.6.3) was operated with default parameters,

and further removed genes with a proportion of unknown nucleotides above 1%. Then,

four species were selected to generated 1:1 single copy genes using blastn version

2.2.31+, including *H. taimen*, *H. bleekeri*, *B. tsinlingensis*, and the outgroup *Salmo*

salar. After the construction of phylogenetic tree using the methods described above,

we estimated the divergence time using BEAST version 2.6.60, the nucleotide

substitution model F81 was predicted by jmodeltest version 2.1.10, the chain length

was set to 50,000,000 and pre-burnin 10%.

Demographic history estimation

The pairwise sequentially Markovian coalescent (PSMC) method version 0.6.5-r67 (Li & Durbin, 2011) was utilized to predict the effective population size. Besides the two wild taimen sample, we used two lenok samples with highest sequencing depths from Heilongjiang River (*B.tumernsis*) and Qinling Mountains (*B. tsinlingensis*). Due to the different sequencing depth, we used different strategies to generate the diploid consensus sequences for taimen and lenok, -D 60 for taimen and -D 20 for lenok at vcfutils.pl packages of bcftools version 1.12 (Danecek et al., 2021). The psmc was run with parameters "-N25 -t30 -r4 -p 8+4+25*2+4+6" to control the divergence at left of the results.

RESULTS

Genome size estimation and assembly

A total of 426,685,475 paired-end reads were obtained from the Illumina NovaSeq-6000 platform, yielding a total length of 64.00 Gb. The genome size estimated by GCE (k=17) was ~3.21 Gb with a low heterozygous ratio of 0.44% (Supplementary Figure S1; Supplementary Table S1). Based on the estimated genome size, the PacBio HiFi and Illumina Hi-C sequencing produced 76.48 Gb (23.8×) and 350.63 Gb (109.23×) of raw reads for genome assembly, respectively. Besides, 155.88 Gb of resequencing data from two additional wild samples (~22 ×) were generated.

After *de novo* genome assembly using the HiFi reads, a 3.48 Gb primary genome was generated, which included 4,373 high quality contigs (>2 Kb) with a contig N50 of 2.4 Mb (Table 1). Following Hi-C assisted genome assembly and scaffold

deduplication, the contigs were anchored into 44 chromosome-level scaffolds,

covering 95.26% of the total scaffolds length, with a scaffold N50 of 73.71 Mb

(Figure 1A; Supplementary Figure S2; Table 1). The final Sichuan taimen genome,

closely matching the estimated size, had a total size of 3.45 Gb with a GC content of

41.99% (Figure 1A, B; Table 1). The BUSCO results showed that the assembled

genome included 95.3% complete genes from the actinopterygii_odb9 database,

demonstrating high completeness (Supplementary Table S2).

Landscape of repeat sequences

The non-redundancy length of the total repeat sequences recognized by three

annotation methods was 2.60 Gb, covering 74.21% of the assembled genome, which

represents the highest proportion of repetitive sequences among salmonids (Figure 1C;

Figure 2; Supplementary Table S3, S4 and S5). The interspersed repeats reported by

the Repeatmasker and LTR_FINDER composed 36.18% of the genome, the lowest

proportion among salmonids; however, the total length of interspersed repeats in the

Sichuan taimen genome was similar to that of other salmonids (Supplementary Table

S3 and S4). The proportion of retroelements in the Sichuan taimen genome was

13.11%, with 7.03% LINEs and 5.98% LTR elements (Supplementary Table S3).

DNA transposons composed 9.93% of the genome, with the Tc1-IS630-Pogo family

being the most abundant TE, similar to other salmonids, covering 9.38% of the

Sichuan taimen genome (Supplementary Table S3).

Tandem repetitive sequences covered 44.15% of the Sichuan taimen genome,

representing highest proportion among salmonids (Figure 2A; Supplementary Table

S5). The lengths of STRs and VNTRs were 45,561,499 bp and 1,553,872,291,

accounting for 1.30% and 44.40% of the genome, respectively. Additionally, 59.12%

of STRs were less than 1 Kb away from the surrounding VNTRs (Figure 2A). The

distribution of TR was not uniform. We found that 29.44% of the regions in the

Sichuan taimen genome had low TR ratios (One-tenth of the genome-wide scale;

Figure 2A), and some chromosomes had an enrichment of VNTR at their ends (Figure

2A). The distribution of STRs in the Sichuan taimen genome was relatively scattered,

with only the top 200 STRs by length exceeding the average length (16,620 bp) of the

HiFi reads (Figure 2B). Dinucleotide repeats accounted for 81.53% of STRs

(Supplementary Table S6), which is higher than the average level for vertebrates

(Adams et al., 2016). We identified 3,337 VNTRs with lengths over 100 Kb, totaling

0.97 Gb and covering 28.20% of the genome (Figure 2C). In addition, the presence of

TRs affected GC content in local regions of the genome; however, the GC content of

the TR regions was 41.16%, similar to the genome-wide range (Figure 2A).

Gene annotation and genomic synteny

A total of 40.00 Gb of RNA reads were generated from five tissues for RNA-based

gene annotation. After filtering pseudogenes, the BRAKER2 annotation pipeline

predicted 49,052 genes, with an average length of 14,577 bp, containing 338,448

exons in total. The average number of exons per gene was ~7 (Figure 1D). After

mapping to the NR databases, only 79.02% of the predicted genes were functionally

annotated. The TR ratio in intron regions was 22.74%, significantly lower than the

genome-wide level (Z-test; $P<0.01$). Based on kernel density estimation, 161 gene

hotspots were identified (Supplementary Figure S3). NR database annotation results showed that the top three regions with the highest gene density were located in histone gene family loci on chromosome 8 and 22. The BUSCO analysis for the predicted genes indicated that 88.1% were complete BUSCOs (C: 88.1%[S: 43.3%, D: 44.8%], F: 4.3%, M: 7.6%, n: 4584).

After the protein mapping process, we identified 15,787 paired homeologous genes, accounting for 64.37% of the total predicted genes. Eventually, 44 homeologous blocks were located based on the genomic synteny of the genes (Figure 1E; Supplementary Supplementary Table S7). The circos plot of homeologous blocks showed that the longer chromosomes were more likely to be composed of multiple homologous blocks, especially for the first five chromosomes larger than 120 Mb (Figure 1E). Notably, more than half of the chromosomes (27/44) were composed of a single homologous block, and the repeat distribution pattern was similar between blocks (Figure 1E; Figure 2A).

Analysis of phylogenetic and evolution of gene family

We identified 35,589 orthologous gene families between Sichuan taimen and nine other species, including 481 gene families specific to the Sichuan taimen and 342 single-copy orthologues. The results of the phylogenetic study showed that, as an ancient branch of Salmoninae, the divergence time between huchonids (*Hucho* spp. and *Brachymystax* spp.) and other Salmoninae fishes was estimated at approximately 33.29 Mya (Million years ago)(Figure 3; Supplementary Figure S4), which is consistent with previous studies (Horreo, 2017; Lecaudey et al., 2018). The

divergence time estimated by PAML was 15.81 Mya between the taimen and its sister

group, lenok (*Brachymystax* spp.)(Figure 3).

We identified 235 significant expanded genes in the Sichuan taimen genome ($P <$

0.01). Gene Ontology analysis showed the highest enrichment in nucleosome

(GO:0000786) and chromosome (GO:0005694) categories (Supplementary Table S8).

Similarly, a large number of histone genes were found in the expanded gene set,

which we speculated might be influenced by the expansion of repetitive sequences.

Besides, the GO term fin development (GO:0033333) was annotated, implying that

the Sichuan taimen might have evolved enhanced swimming or predation abilities

(Supplementary Table S8). KEGG pathway analysis showed enrichment in the

Hedgehog signaling pathway (dre04340) and Insulin signaling pathway (dre04910),

suggesting changes in embryonic development, energy metabolism and growth, which

might be related to its special body size and habits (Supplementary Table S9).

Ancestral chromosome status

To evaluate the chromosome evolution of Sichuan taimen, genomic synteny blocks

were recognized against the Northern pike (*Esox lucius*), which serves as an

appropriate outgroup preceding the Ss4R. A total of 12,077 orthologs (1:2 or 1:1 for *E.*

lucius to *H. bleekeri*) were identified, resulting in the linkage of 80 homologous

blocks covering 90.89% of the length of the 44 Sichuan taimen chromosomes

(Supplementary Table S10). We recovered Sichuan taimen chromosome

equivalents for all Northern pike chromosomes, with each Northern pike chromosome

corresponding to two Sichuan taimen homologues (Figure 4A). The results showed

that half of the Sichuan taimen chromosomes (26/44) mapped completely to individual Northern pike chromosomes (Figure 4A).

The total chromosome lengths were 3.32 Gb and 0.92 Gb for Sichuan taimen and Northern pike, respectively. However, the length of the Sichuan taimen genome far exceeded what would be expected from whole-genome duplication alone. The expansion of TR in the Sichuan taimen genome was also evident in the circos plot (Figure 4A). We further evaluated the TR expansion in the Sichuan taimen by comparing the ratio of total length to TR length in the 80 homologous blocks. The results revealed that all blocks showed TR expansion in the Sichuan taimen genome, with the TR length ratio increasing proportionally with total length growth (Figure 4B). Linear regression analysis indicated that the TR ratio in the Sichuan taimen genome was approximately fivefold higher than that of the Northern pike (Figure 4B).

Demographic history and divergence for two taimen species

The geographical distribution pattern of the other landlocked huchonid fish, lenok, is similar to that of taimen. Both are found in the Qinling Mountain regions and the Heilongjiang River, and the competitive relationship is existed between them (Galland et al., 2021; Kaus et al., 2019). To reveal the divergence time between the two taimen species, one Siberia taimen (*H. taimen*) sample were collected from the Heilongjiang River (Figure 5A). A total of 44,120 genes were captured from the Siberia taimen genome, and we identified 4,860 1:1 orthologous genes among the four species. The result of BEAST showed that two taimen species separated at 2.29 Mya (Figure 5B), which was similar to the previous inference (Kucinski & Fopp-Bayat, 2022).

To illustrate the trends in historical effective population sizes (N_e) changes of taimen

and lenok in Shaanxi and Heilongjiang, four resequencing samples from these regions

were used for joint analysis. The results showed that the histories of the four species

trace back to different times: ~9 Mya for the two Heilongjiang samples (*H. taimen*

and *B.tumernsis*), and ~1 Mya for the two Shaanxi samples (*H. bleekeri* and *B.*

tsinlingensis)(Figure 5C). *H. taimen* had a higher N_e level than *H. bleekeri* throughout

their entire history (Figure 5C). The N_e curve suggested that a large-scale population

expansion event occurred around 2 Mya in both *H. taimen* and *B.tumernsis*, followed

by a continuous decline in population size over millions of years (Figure 5C). In

summary, the Heilongjiang populations had higher N_e level than the Shaanxi

populations since ~200 Kya. Although the biomass of Sichuan taimen is currently

smaller than that of Qinling lenok, there might have been times in history when the

opposite was true (Figure 5C), which might reflect the competitive dynamics between

them.

DISCUSSION

Complete genome assembly under high tandem repeat ratio

The combination of HiFi and Hi-C reads allowed us to achieve the chromosome-level

assembly of the largest salmonid genome to date. The final size of the Sichuan taimen

genome was approximately 3.45 Gb, which exceeds the size of the Atlantic salmon

genome (~2.97 Gb)(Lien et al., 2016). The chromosome count of the Sichuan taimen

genome was 44, which approached to other salmonids such as *Salvelinus namaycush*

(42), *B. tsinlingensis* (41), and *Salmo trutta* (40)(Supplementary Table S11). The

assembled genome recovered 95.20% complete BUSCO genes, with a relatively high

percentage of duplicated BUSCOs (46.20%), similar to its sister group *B. tsinlingensis*

(Smith et al., 2022). Additionally, the chromosomal-level gene synteny implied the

integrity of the genome assembly and annotation.

The genome assembly process and outcome were affected by the ratio of repetitive

sequences (Tørresen et al., 2017; Treangen & Salzberg, 2011). The highly accurate

reads produced by CCS technology improved both the process and quality of the

assembly, especially inrepetitive sequences regions (Logsdon et al., 2020; Wenger et

al., 2019). We found that the assembled Sichuan taimen genome contains a high

proportion of repetitive sequences, with nearly half of the genome covered by tandem

repeat sequences. This may contribute to the increased genome size and pose

challenges for genome assembly. Fortunately, most of the TR regions were covered

and assembled by the HiFi reads, demonstrating the advantage of long reads.

However, some long TRs accounted for a relatively large proportion of the total

genome length, causing gaps around them. Generally, the number of gaps in the

assembled Sichuan taimen genome was similar to or even smaller than other salmonid

genomes, except for GCF_013265735.2, indicating that our assembly was not

severely disturbed by TRs (Supplementary Table S11).

Long repetitive regions can affect gene annotation, leading to the appearance of false

and redundant genes (Lang et al., 2020). In our preliminary gene annotation, we found

many tandem pseudogenes in long TR regions, which interfered with subsequent

analysis. Despite the presence of TRs in specific protein domains (Simon & Hancock,

363 2009), each of these pseudogenes usually had only one exon according to our manual

364 inspection. Therefore, we removed these single-exon genes located within repetitive

365 sequence regions based on expression levels (TPM<1). In parallel, we blasted these

366 pseudogenes to confirm their absence in other species.

Phylogenetic relationships of Salmoninae based on single-copy orthologous genes

Salmonidae contains three subfamilies: Coregoninae, Thymallinae, and Salmoninae.

However, their evolutionary relationships remains uncertain (Dysin et al., 2022).

Lineage-specific rediploidization, incomplete lineage sorting, hybridization, and

introgression have resulted in a complex and disordered phylogenetic relationship

within Salmonidae (Campbell et al., 2020; Lecaudey et al., 2018; Robertson et al.,

2017). Some studies suggest that Thymallinae is the sister group of Salmoninae

(Crête-Lafrenière et al., 2012; Mckay et al., 2004; Wang et al., 2011), while other

studies place Thymallinae at the root of the Salmonidae phylogenetic tree (Horreo,

2017; Wang et al., 2022). Based on the phylogenetic tree constructed using the

orthologous genes, we suggest that Coregoninae and Thymallinae have a sister

relationship, supported by 97% bootstraps (Supplementary Figure S4). This is

consistent with other trees based on orthologues (Macqueen & Johnston, 2014;

Robertson et al., 2017), mitochondrial genome (Campbell et al., 2013; Shedko et al.,

2013), and ultra-conserved elements (Campbell et al., 2020). The inferred topology

within Salmoninae aligns with other studies, with the genera *Hucho* and

Brachymystax forming the ancient branch of the subfamily, constituting a sister group

to the genera *Salmo, Salvelinus*, *and Oncorhynchus* (Crespi & Fulton, 2004; Kucinski

& Fopp-Bayat, 2022). The divergence time between taimen and lenok was estimated

at 9.41 to 23.51 Mya, which aligns with previous results based on mitochondria

(Kucinski & Fopp-Bayat, 2022), and coincides with the Miocene Climate Optimum

(MCO) at 15-18 Mya (Böhme, 2003; Sun & Zhang, 2008).

Effective population size changes and the urgency of conservation

All species of the genera *Hucho* and *Brachymystax* are endangered and protected by

Chinese government law (Sung et al., 1998). The results of PSMC revealed that the

effective populations size of four *Hucho* and *Brachymystax* species have declined

over the past million years, following the separation of *H. bleekeri* and *H. taimen*.

Similar patterns of N_e curves were observed in species with the same geographical

distribution. The curves of *H. bleekeri* and *B. tsinlingensis* in Shaanxi showed a close

coalescent endpoint, we speculated that these two salmonids shared the same

immigration route during the glacial period and were retained in low-latitude areas

during the interglacial period. The Sichuan taimen had a lower N_e level than Siberia

taimen in our calculation result, and current resource surveys indicate that Sichuan

taimen are very rare in the Qinling Mountains, having disappeared from most of their

original distribution areas (Shen et al., 2006; Wang et al., 2016). Consequently, more

effective conservation measures should be implemented to protect the Sichuan

taimen.

CONCLUSIONS

This study provides a comprehensive genome assembly of the Sichuan taimen, uncovering a novel tandem repeat sequences expansion phenomenon and offering valuable insights into the evolutionary history of this species. The detailed analysis of phylogenetic relationships and effective population size changes highlights the historical population dynamics of the Sichuan taimen and its relatives. The assembled genome serves as an essential resource for future research on the evolutionary biology and conservation genetics of salmonids. As the Sichuan taimen remains critically endangered, this genomic resource provides a foundation for developing effective conservation measures to ensure the survival of this ancient and unique species.

COMPETING INTERESTS

The authors declare that they have no competing interests.

AUTHOR CONTRIBUTIONS

Xin-miao Zhang, Li-xin Wang and Yu Jiang conceived and supervised the project. Dong-mei Xiong, and Sheng-long Jian the samples. Xin-miao Zhang, and Dong-mei Xiong carried out the bioinformatic analyses. Xin-miao Zhang wrote the manuscript. Sheng-long Jian, Yu Jiang and Li-xin Wang revised the manuscript. All authors have read and approved the final manuscript.

ACKNOWLEDGEMENTS

This study was financially supported by the National Natural Science Foundation of

China (32273138) and the latitudinal project foundation of China Power Construction

Group Beijing Survey, Design and Research Institute Co., Ltd. (K4050422184) . We

thank the High-Performance Computing Platform of Northwest A&F University and

the National Supercomputing Center in Xi'an and Hefei for providing computing

resources.

REFERENCE

Adams RH, Blackmon H, Reyes-Velasco J, et al. 2016. Microsatellite landscape evolutionary dynamics
across 450 million years of vertebrate genome evolution. *Genome*, **59**(5): 295-310.

Allendorf FW & Hard JJ. 2009. Human-induced evolution caused by unnatural selection through
harvest of wild animals. *Proceedings of the National Academy of Sciences of the United States of*
America, **106**(1): 9987-9994.

Allendorf FW, Hohenlohe PA, Luikart G. 2010. Genomics and the future of conservation genetics.
Nature Reviews Genetics, **11**(10): 697-709.

Benson G. 1999. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids*
Researchearch, **27**(2): 573-580.

Böhme M. 2003. The Miocene climatic optimum: evidence from ectothermic vertebrates of Central
Europe. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **195**(3-4): 389-401.

Braasch I, Gehrke AR, Smith JJ, et al. 2016. The spotted gar genome illuminates vertebrate evolution
and facilitates human-teleost comparisons. *Nature Genetics*, **48**(4): 427-437.

Brandies P, Peel E, Hogg CJ, et al. 2019. The value of reference genomes in the conservation of
threatened species. *Genes (Basel)*, **10**(11).

Bray NL, Pimentel H, Melsted P, et al. 2016. Near-optimal probabilistic RNA-seq quantification.
Nature Biotechnology, **34**(5): 525-527.

Campbell MA, Buser TJ, Alfaro ME, et al. 2020. Addressing incomplete lineage sorting and paralogy
in the inference of uncertain salmonid phylogenetic relationships. *PeerJ*, **8**: e9389.

Campbell MA, López JA, Sado T, et al. 2013. Pike and salmon as sister taxa: detailed intraclade
resolution and divergence time estimation of Esociformes + Salmoniformes based on whole
mitochondrial genome sequences. *Gene*, **530**(1): 57-65.

Chen S, Zhou Y, Chen Y, et al. 2018. Fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics, **34**(17): i884-i890.

Chen Y, Yang H, Chen Y, et al. 2021. Full-length transcriptome sequencing and identification of
immune-related genes in the critically endangered *Hucho bleekeri*. *Developmental & Comparative*
Immunology, **116**: 103934.

Chen Y, Yang H, Gong Q, et al. 2020. Isolation and characterization of 34 SNP markers in *Hucho*
bleekeri. *Conservation Genetics Resources*, **12**: 157-160.

Cheng H, Concepcion GT, Feng X, et al. 2021. Haplotype-resolved *de novo* assembly using phased
assembly graphs with hifiasm. *Nature Methods*, **18**(2): 170-175.

Christensen KA, Rondeau EB, Minkley DR, et al. 2020. The sockeye salmon genome, transcriptome, and analyses identifying population defining regions of the genome. *PLoS One*, **15**(10): e0240935.

Crespi BJ & Fulton MJ. 2004. Molecular systematics of Salmonidae: combined nuclear data yields a robust phylogeny. *Molecular Phylogenetics and Evolution*, **31**(2): 658-679.

Crête-Lafrenière A, Weir LK, Bernatchez L. 2012. Framing the Salmonidae family phylogenetic portrait: a more complete picture from increased taxon sampling. *PLoS One*, **7**(10): e46662.

Danecek P, Bonfield JK, Liddle J, et al. 2021. Twelve years of SAMtools and BCFtools. *Gigascience*, **10**(2):giab008.

Dobin A, Davis CA, Schlesinger F, et al. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, **29**(1): 15-21.

Dudchenko O, Batra SS, Omer AD, et al. 2017. De novo assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. *Science*, **356**(6333): 92-95.

Durand NC, Robinson JT, Shamim MS, et al. 2016a. Juicebox provides a visualization system for Hi-C contact maps with unlimited zoom. *Cell Systems*, **3**(1): 99-101.

Durand NC, Shamim MS, Machol I, et al. 2016b. Juicer provides a one-click system for analyzing loop-resolution Hi-C experiments. *Cell Systems*, **3**(1): 95-98.

Dysin AP, Shcherbakov YS, Nikolaeva OA, et al. 2022. Salmonidae genome: features, evolutionary and phylogenetic characteristics. *Genes (Basel)*, **13**(12):2221.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, **32**(5): 1792-1797.

Emms DM & Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biology*, **20**(1): 1-14.

Flynn JM, Hubley R, Goubert C, et al. 2020. RepeatModeler2 for automated genomic discovery of transposable element families. *Proceedings of the National Academy of Sciences of the United States of America*, **117**(17): 9451-9457.

Gabriel L, Hoff KJ, Brůna T, et al. 2021. TSEBRA: transcript selector for BRAKER. *BMC Bioinformatics*, **22**(1): 566.

Galland LM, Simmons JB, Jahner JP, et al. 2021. Hierarchical genetic structure and implications for conservation of the world's largest salmonid, *Hucho taimen. Sci Rep*, **11**(1): 20508.

Gao G, Magadan S, Waldbieser GC, et al. 2021. A long reads-based de-novo assembly of the genome of the Arlee homozygous line reveals chromosomal rearrangements in rainbow trout. *G3 (Bethesda)*, **11**(4):jkab052.

Hansen T, Fjelldal PG, Lien S, et al. 2021. The genome sequence of the brown trout, *Salmo trutta* Linnaeus 1758. *Wellcome Open Research*, **6**: 108.

Hoff KJ, Lange S, Lomsadze A, et al. 2016. BRAKER1: Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS. *Bioinformatics*, **32**(5): 767-769.

Hoff KJ, Lomsadze A, Borodovsky M, et al. 2019. Whole-Genome Annotation with BRAKER. *Methods in Molecular Biology*, **1962**: 65-95.

Horreo JL. 2017. Revisiting the mitogenomic phylogeny of Salmoninae: new insights thanks to recent sequencing advances. *PeerJ*, **5**: e3828.

Hu M, Wang Y, Cao L, et al. 2008. Threatened fishes of the world: *Hucho bleekeri* Kimura, 1934 (Salmonidae). *Environmental Biology of Fishes*, **82**: 385-386.

Ishiguro NB, Miya M, Nishida M. 2003. Basal euteleostean relationships: a mitogenomic perspective on the phylogenetic reality of the "Protacanthopterygii". *Molecular Phylogenetics and Evolution*, **27**(3):

476-488.

Kaus A, Michalski S, Hänfling B, et al. 2019. Fish conservation in the land of steppe and sky: Evolutionarily significant units of threatened salmonid species in Mongolia mirror major river basins. *Ecology and Evolution*, **9**(6): 3416-3433.

Kimura S. 1934. Description of the fishes collected from the Yangtze-Kiang China by late Dr. K. Kishinouye and his party in 1927–1929. Shanghai: Journal of the Shanghai Science Institute.

Kohn MH, Murphy WJ, Ostrander EA, et al. 2006. Genomics and conservation genetics. *Trends in Ecology Evolution*, **21**(11): 629-637.

Kucinski M & Fopp-Bayat D. 2022. Phylogenetic analysis of *Brachymystax* and *Hucho* genera-Summary on evolutionary status within the Salmoninae subfamily. *Journal of Applied Ichthyology*, **38**(4): 403-411.

Kumar S, Suleski M, Craig JM, et al. 2022. TimeTree 5: An Expanded Resource for Species Divergence Times. *Molecular Biology and Evolution*, **39**(8).

Lang D, Zhang S, Ren P, et al. 2020. Comparison of the two up-to-date sequencing technologies for genome assembly: HiFi reads of Pacific Biosciences Sequel II system and ultralong reads of Oxford Nanopore. *Gigascience*, **9**(12):giaa123.

Lecaudey LA, Schliewen UK, Osinov AG, et al. 2018. Inferring phylogenetic structure, hybridization and divergence times within Salmoninae (Teleostei: Salmonidae) using RAD-sequencing. *Molecular Phylogenetics and Evolution*, **124**: 82-99.

Li H & Durbin R. 2011. Inference of human population history from individual whole-genome sequences. *Nature*, **475**(7357): 493-496.

Lien S, Koop BF, Sandve SR, et al. 2016. The Atlantic salmon genome provides insights into rediploidization. *Nature*, **533**(7602): 200-205.

Liu B, Shi Y, Yuan J, et al. 2013. Estimation of genomic characteristics by analyzing k-mer frequency in *de novo* genome projects. *arXiv.org arXiv*: 1308.2012.

Liu X, Zeng H, Wang C, et al. 2022. Improved genome assembly of Chinese sucker (*Myxocyprinus asiaticus*) provides insights into the identification and characterization of pharyngeal teeth related maker genes in Cyprinoidei. *Water Biology and Security*, **1**(3): 100049.

Logsdon GA, Vollger MR, Eichler EE. 2020. Long-read human genome sequencing and its applications. *Nature Reviews Genetics*, **21**(10): 597-614.

Macqueen DJ & Johnston IA. 2014. A well-constrained estimate for the timing of the salmonid whole genome duplication reveals major decoupling from species diversification. *Proceedings of the Royal Society B-Biological Sciences*, **281**(1778): 20132881.

Mckay SJ, Trautner J, Smith MJ, et al. 2004. Evolution of duplicated growth hormone genes in autotetraploid salmonid fishes. *Genome*, **47**(4): 714-723.

Mendes FK, Vanderpool D, Fulton B, et al. 2020. CAFE 5 models variation in evolutionary rates among gene families. *Bioinformatics*, **36**(22-23): 5516-5518.

Ou S & Jiang N. 2018. LTR_retriever: a highly accurate and sensitive program for identification of long terminal repeat retrotransposons. *Plant Physiology*, **176**(2): 1410-1422.

Robertson FM, Gundappa MK, Grammes F, et al. 2017. Lineage-specific rediploidization is a mechanism to explain time-lags between genome duplication and evolutionary diversification. *Genome Biology*, **18**(1): 111.

Sadovy De Mitcheson Y, Craig MT, Bertoncini AA, et al. 2013. Fishing groupers towards extinction: a global assessment of threats and extinction risks in a billion dollar fishery. *Fish and Fisheries*, **14**(2):

119-136.

Shedko SV, Miroshnichenko IL, Nemkova GA. 2013. Phylogeny of salmonids (Salmoniformes,

Salmonidae) and molecular dating: analysis of mtDNA data. *Genetika*, **49**(6): 718-734.

Shen Z, Tang W, Li K. 2006. The analysis of population dynamics of *Hucho bleekeri* in Markehe River,

Qinghai Province. *Reservoir Fisheries*, **26**: 71-73 (In Chinese).

Simão FA, Waterhouse RM, Ioannidis P, et al. 2015. BUSCO: assessing genome assembly and

annotation completeness with single-copy orthologs. *Bioinformatics*, **31**(19): 3210-3212.

Simon M & Hancock JM. 2009. Tandem and cryptic amino acid repeats accumulate in disordered

regions of proteins. *Genome Biology*, **10**(6): R59.

Smith SR, Normandeau E, Djambazian H, et al. 2022. A chromosome-anchored genome assembly for

Lake Trout (*Salvelinus namaycush*). *Molecular Ecology Resources*, **22**(2): 679-694.

Song Z. 2012. *Hucho bleekeri*. The IUCN Red List of Threatened Species. Version 2014.3.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics*, **30**(9): 1312-1313.

Stefanova P, Taseva M, Georgieva T, et al. 2013. A modified CTAB method for DNA extraction from

soybean and meat products. *Biotechnology & Biotechnological Equipment*, **27**(3): 3803-3810.

Sun J & Zhang Z. 2008. Palynological evidence for the Mid-Miocene Climatic Optimum recorded in

Cenozoic sediments of the Tian Shan Range, northwestern China. *Global and Planetary Change*, **64**(1):

53-68.

Sung W, Peiqi Y, Yiyu C, et al. 1998. China Red data book of endangered animals: Pisces. Beijing:

Science press.

Tarailo-Graovac M & Chen N. 2009. Using RepeatMasker to identify repetitive elements in genomic

sequences. *Current Protocols in Bioinformatics*, **4**: 1-4.10.

Tørresen OK, Star B, Jentoft S, et al. 2017. An improved genome assembly uncovers prolific tandem

repeats in Atlantic cod. *BMC Genomics*, **18**(1): 95.

Treangen TJ & Salzberg SL. 2011. Repetitive DNA and next-generation sequencing: computational

challenges and solutions. *Nature Reviews Genetics*, **13**(1): 36-46.

Wang K, Zhang SH, Wang DQ, et al. 2016. Conservation genetics assessment and phylogenetic

relationships of critically endangered *Hucho bleekeri* in China. *Journal of Applied Ichthyology*, **32**(2):

343-349.

Wang Y, Guo R, Li H, et al. 2011. The complete mitochondrial genome of the Sichuan taimen (*Hucho*

bleekeri): repetitive sequences in the control region and phylogenetic implications for Salmonidae.

Marine Genomics, **4**(3): 221-228.

Wang Y, Xiong F, Song Z. 2022. Molecular phylogeny and adaptive mitochondrial DNA evolution of

salmonids (Pisces: Salmonidae). *Frontiers in Genetics*, **13**: 903240.

Waples RS, Naish KA, Primmer CR. 2020. Conservation and management of salmon in the age of

genomics. *Annual Review of Animal Biosciences*, **8**: 117-143.

Wenger AM, Peluso P, Rowell WJ, et al. 2019. Accurate circular consensus long-read sequencing

improves variant detection and assembly of a human genome. *Nature Biotechnology*, **37**(10):

1155-1162.

Xu Z & Wang H. 2007. LTR_FINDER: an efficient tool for the prediction of full-length LTR

retrotransposons. *Nucleic Acids Researchearch*, **35**(2): W265-W268.

Xue DX, Xing TF, Liu JX. 2022. A high-quality chromosome-level genome of the endangered

roughskin sculpin provides insights into its evolution and adaptation. *Molecular Ecology Resources*,

22(5): 1892-1905.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution*, **24**(8): 1586-1591.

Ye J, Mcginnis S, Madden TL. 2006. BLAST: improvements for better sequence analysis. *Nucleic Acids Research*, **34**(Web Server issue): W6-9.

Zhang Y, Luan P, Ren G, et al. 2020. Estimating the inbreeding level and genetic relatedness in an isolated population of critically endangered Sichuan taimen (*Hucho Bleekeri*) using genome-wide SNP markers. *Ecology and Evolution*, **10**(3): 1390-1400.

Zhu W, Wang Z, Li H, et al. 2022. A chromosome-level genome of *Brachymystax tsinlingensis* provides resources and insights into salmonids evolution. *G3 (Bethesda)*, **12**(8):jkac162.

603 **Figure 1 Circos plot of the genome assembly statistics of Sichuan taimen**

604 A: Chromosome names and length scales. Chromosomes consisting of multiple homologous
605 regions were shown in gray (The same below). B: GC ratio (scale: 0-100%). C: Proportion of total
606 repeats (TR and TE)(scale: 0-100%). D: Gene density after log transformation (scale: 0-7). E:
607 Homeologous regions in the Sichuan taimen genome recognized by 1:1 duplicated proteins.

Figure 2 Distribution and statistics of repetitive sequence in the Sichuan taimen genome.

A: Visualization of genomic GC content, STR, and VNTR distribution based on 100 Kb windows; the bar graph reflects the TR frequency distribution in genome-wide 100 Kb windows. B and C were statistics of tandem repeat unit length and total tandem repeat sequence length of STRs and VNTRs respectively; The average length of the HiFi reads (16620 bp) was labeled as dashed line.

Figure 3 Maximum likelihood phylogenetic tree constructed based on single-copy orthologous genes.

Fossil calibration points used by MCMC are marked in red. Black numbers next to each nodes are the divergence times (Mya) and the 95% confidential intervals estimated by PAML. The green positive and red negative integers indicate the number of genes at expanded or contracted gene families at that node, respectively ($P < 0.05$). Ss4R: salmonid-specific fourth vertebrate whole-genome duplication.

Figure 4 Chromosome evolution of *H. bleekeri*

A: Circos plot of homologous blocks and TR ratio in *E. lucius* and *H. bleekeri*. From the outside to the inside were chromosome length, TR ratio, and homologous block connections. B: TR expansion in taimen genome. The abscissa and ordinate were the ratio of the total length and TR length in homologous blocks between Sichuan taimen and Northern pike, respectively. The size of the circles was expressed as the ratio of the proportion of TR (RPT) in homologous blocks between them.

Figure 5 Species divergence history between *H. taimen* and *H. bleekeri*

A: Map of the two taimen species distribution. B: Phylogeny and divergence time of huchonid. 95% confidence interval was labeled as blue frame. Fossil calibration points were marked in red. C: Demographic history of two Heilongjiang sample (*H. taimen* and *B.tumernsis*) and two Qinling sample (*H. bleekeri* and *B.tsinlingensis*). The generation (g) time was set at 4 years and the neutral mutation rate per generation (μ) was set at 6.54 × 10-9.

Table 1. Summary statistics of the Sichuan taimen genome assembly

Category	Length (bp)		Number	
	Contig	Scaffold	Contig	Scaffold
Max	25,857,850	143,121,144	-	-
Min	3,656	18,648	-	-
N50	2,400,826	73,705,547	354	19
N90	418,871	59,067,955	1680	39
Total	3,479,724,502	3,451,243,059	4373	505

SUPPLEMENTARY FIGURE AND TABLE LEGEND

Supplementary Figure S1. Frequency distribution map in 17-mer.

Supplementary Figure S2. Hi-C interaction heatmap of Hucho bleekeri genome between 44 chromosomes and unmapped scaffolds.

Supplementary Figure S3. Gene hotspots of the hucho genome.

Supplementary Figure S4. Phylogenetic trees under 1,000 bootstraps.

Supplementary Table S1. Genome size estimation table at k=17.

Supplementary Table S2. Summarized benchmarking in BUSCO notation.

Supplementary Table S3. Summary of TE in 10 salmonids (%).

Supplementary Table S4. Summary of TE in 10 salmonids (length).

Supplementary Table S5. Statistics of TR for 10 salmonids.

Supplementary Table S6. Summary of STRs' unit and total length.

Supplementary Table S7. Genomic synteny of 44 homeologous blocks.

Supplementary Table S8. The Gene Ontology annotation of the 235 positive selected genes.

Supplementary Table S9. The KEGG pathway annotation of the 235 positive selected genes.

Supplementary Table S10. Genomic synteny of 80 homeologous blocks between Northern pike (luc) and Sichuan taimen.

Supplementary Table S11. Summary of genome assembly results of 10 salmonids.