Chromosome-level genome assembly for *hucho bleekeri* reveals the extraordinary tandem repeat proportions and its persistent population shrinkage

Xinmiao Zhang[1], Dongmei Xiong[1], Shenglong Jian[2], Yu Jiang[3], Lixin wang[1*]

[1]College of Animal Science and Technology, Northwest A&F University, Yangling, 712100, China

[2]Qinghai Provincial Fishery Technology Extension Center, Xining, China

[3]Center for Ruminant Genetics and Evolution, College of Animal Science and Technology, Northwest A&F University, Yangling 712100, China

[*]Corresponding author. E-mail: fisherwanglx@nwsuaf.edu.cn

Abstract

The Chuanshan hucho (*Hucho bleekeri*) is one of the largest freshwater salmonids, and located at the base of the Salmoninae phylogeny. Facing a serious risk of extinction, the Chuanshan hucho has attracted widespread concern and has been listed as national first-level protected animal in China. Currently, biological research and conservation policies of the Chuanshan hucho are in full progress, but the complete genome sequence is still unknown which hindered the advance of its conservation genomics. In this study, we provided the first chromosome-level genome for Chuanshan hucho, the final genome size was ~3.45 Gb with 44 chromosomes. The hucho genome contained 44.15% tandem repetitive sequence, which exceeded all sequenced salmonid fishes. According to the nuclear sequence, the divergence time of *Hucho* and *Brachymystax* was estimated as 12.29 Mya, and the effective population size of Chuanshan hucho declined from 1 Mya. The genomic resource in this article will promote the protection of Chuanshan hucho and evolutionary genetics studies of salmonids.

1 Introduction

The Chuanshan hucho (*Hucho bleekeri*) belonging to Salmoninae, is a glacial remnant species and endemic to China (Wang, et al. 2016). It has the lowest distribution latitude among all the fishes in *Hucho* genus, and mainly exists in the upper tributaries of the Yangtze River (Hu, et al. 2008). Hucho have a large body size and a ferocious habit, which occupied the top of the aquatic food chain (Wang, et al. 2016), and the late sexual maturation makes it more vulnerable to human activities and

ecological changes (Allendorf and Hard 2009; Sadovy de Mitcheson, et al. 2013).

Individuals of *H. bleekeri* were found sporadically and scattered in their distribution,

and had been protected since 1998, however, they continues to be illegally captured

by poachers for its high nutritional value and taste (Sung, et al. 1998). As the

continuous decline, *H. bleekeri* is listed as critically endangered (CR) level in the

International Union for Conservation of Nature (IUCN) Red List of Threatened

Species in 2012 (Song 2012). At present, *H. bleekeri* is under the highest level of

protection from the Chinese government (since 2021), and it is the only national

protected animal with "class I" among all distributed salmonid fishes.

Genomic data provide great opportunities for the progress on conservation and

evolution research of threatened animals (Kohn, et al. 2006; Allendorf, et al. 2010).

As sequencing prices have decreased, the genomes of many endangered fishes have

been published, such as *Trachidermus fasciatus* Heckel (Xue, et al. 2022),

Myxocyprinus asiaticus (Liu, et al. 2022), and *Brachymystax tsinlingensis*

(GCA_031305455.1), etc. Based on the genome data, the scientific and rational

conservation policies were formulated and further promoted the protection benefits of

the endangered animals (Brandies, et al. 2019; Waples, et al. 2020). However, lacking

of genome data of *H. bleekeri*, a blank was existence in the nuclear-based

phylogenetic research of Salmonidae, and also influenced the genomic studies such as

taxonomic studies based on SNPs (Chen, et al. 2020; Zhang, et al. 2020) and

researches on transcriptome (Chen, et al. 2021).

Due to the whole genome duplication and lineage-specific rediploidization in

salmonids, the genome of salmonids became complex and variable, increased the

difficulty of assembly (Robertson, et al. 2017; Dysin, et al. 2022). The circular

consensus sequencing (CCS) mode in single molecule real-time (SMRT) sequencing

of Pacific Biosciences can produce long and high fidelity (HiFi) reads, which

facilitates the genome assembly process and improves the quality (Wenger, et al.

2019). In this research, we utilized HiFi and Hi-C sequencing technology to construct

a high-quality genome for Chuanshan hucho, from the genome we found the existence

of high-proportioned tandem repetitive sequence. Additionally, we analyzed the

phylogenetic relationships of Chuanshan hucho and the demographic history of *H.*

bleekeri and *H. taimen*. Our work provides insights into the genome evolution of

salmonid fishes and the conservation of the endangered Chuanshan hucho.

2 Methods

2.1 Sample collection and sequencing

Two wild individuals of *H. bleekeri* were collected from Taibai river (Baoji, Shannxi

province), one for genome assembly the other for resequencing, and an additional

wild individual of *H. taimen* was collected from Heilongjiang. All the experiments in

this study were performed in accordance with the guidelines of China Council on

Animal Care.

Genome DNA was extracted from kidney tissue with cetyltrimethylammonium

bromide (CATB) method (Stefanova, et al. 2013). The reads for genome size

estimation were generated on Illumina NovaSeq-6000 platform after the constructed

of 150 bp library. For genome assembly, a 15 kb library preparation was performed by

SMRTbell Express Template Prep Kit 2.0 (Pacific Biosciences, CA, USA), and

sequenced on the PacBio Sequel II platform (Pacific Biosciences, CA, USA) with two

cells of single molecule real time sequencing (SMRT). The CCS software v.6.0.0

(https://github.com/PacificBiosciences/ccs) was performed to integrate HiFi reads

with the parameters "--min-passes 3 --min-length 10 --min-rq 0.99". The Hi-C library

was constructed and sequenced on Illumina NovaSeq-6000 platform. The RNA

samples were extracted from fin，brain, gill, gonad, and eye, then sequenced on

Illumina NovaSeq-6000 platform. The two wild samples for demographic history

analysis were sequenced on DNBseq platform (BGI).

2.2 Genome size estimation

The raw Illumina paired-end reads of hucho were filtered using fastp version 0.20.0

(Chen, et al. 2018) with default parameters. The genome size estimation of hucho was

performed by GCE software version 1.0.2 (Liu, et al. 2013). A total of 58.89 Gb clean

reads were used to calculate the distribution of k-mer frequency using kmerfreq with

k=17. Then the gce was ran in both homozygous and heterzygous mode to evaluate

the genome size and heterozygosity. The genome heterozygous was calculated using

the formula "a[1/2]/k".

2.3 Genome assembly

The PacBio HiFi reads were assembled using hifiasm version 0.13-r308 (Cheng, et al.

2021) to obtain *de novo* contigs. After quality control by fastp (Chen, et al. 2018), the

clean Hi-C reads were used to integrate the contigs using Juicer version 1.6 (Durand,

Shamim, et al. 2016) and 3D-DNA version 201008 (Dudchenko, et al. 2017). First,

the Hi-C interaction matrix was generated using Juicer scripts with -s MboI parameter.

And then, the 3D-DNA with 0 rounds was used to link the contigs into chromosome

level based on the Hi-C interaction matrix. The initial assembly was manually

corrected according to the Hi-C heatmap using Juicebox version 1.11.08 (Durand,

Robinson, et al. 2016) and run the 3D-DNA again to generate the final assembly. The

BUSCO (Benchmarking universal single-copy orthologues) version 3.0.2 (Simão, et

al. 2015) was used to assess the accuracy and integrity of the genome with the gene

database of ray-finned fishes (actinopterygii_odb9).

2.4 Repeat detection and gene annotation

To recognize repeat sequences of the hucho genome, we build a *de novo* library of

transposable elements (TEs). First, we ran the RepeatModeler version 2.1 (Flynn, et al.

2020) with default parameters to constructed a repeat library; then an additional long

terminal repeat (LTR) library of the hucho was built by LTR_FINDER version 1.07

(Xu and Wang 2007) and LTR_RETRIEVER version 2.9.6 (Ou and Jiang 2018); third

the final repeat library was integrated by replacing the LTR elements of the first

repeat library with the LTR library. To further classify the "Unknown" sequence, the

BLAST version 2.14.0 (Ye, et al. 2006) was performed between the final repeat

library   and   public   repeat   library

(http://www.hrt.msu.edu/uploads/535/78637/Tpases020812.gz).

The TEs of the genome were described and masked by RepeatMasker version

4.1.2 (Tarailo-Graovac and Chen 2009) with parameters "-xsmall -s -cutoff 255 -frag

20000 –gff -engine ncbi". The tandem repeats (TRs) were detected through Tandem

Repeats Finder (TRF) version 4.09 (Benson 1999) with parameters "2 7 7 80 10 50

500 -f -d -m". The TRs were further divided into short tandem repeats (STRs) and

variable number tandem repeats (VNTRs) according to the length of repeat unit (STR:

unit number at 1 to 6; VNTR: unit number more than 6).

The gene annotation of the hucho genome was carried out using BRAKER2

pipeline version 2.1.5 (Hoff, et al. 2016; Hoff, et al. 2019). For transcriptome based

prediction, the RNAs were mapping to the assembled genome using STAR version

2.7.1a (Dobin, et al. 2013) with default parameters, then the RNA sequence data were

used to predict genes through auto BRAKER2 pipeline. For homology based

prediction, the protein sequence of six species including *Lepisosteus oculatus*

(GCF_000242695.1) (Braasch, et al. 2016), *Esox Lucius* (GCF_011004845.1)

(Ishiguro, et al. 2003), *Oncorhynchus mykiss* (GCF_013265735.2) (Gao, et al. 2021),

Oncorhynchus nerka (GCF_006149115.2) , *Salmo salar* (GCF_000233375.1) (Lien,

et al. 2016), and *Brachymystax tsinlingensis* (GCA_031305455.1) which represent the

sister clade or the ancestor species of hucho were download, and predicted through

BRAKER2 pipeline. The results from RNA and protein-based annotation were

combined using TSEBRA version 1.0.3 (Gabriel, et al. 2021) with default parameters

to generate the final gene prediction file. The gene expression level was measured by

transcript per million (TPM) which were calculated using Kallisto (Bray, et al. 2016)

with default parameters. To remove pseudogenes caused by tandem repetitive

sequence, we filtered the gene set based on the following criteria: 1) contain only one

exon; 2) TPM<1; 3) no matched BLAST results with *Brachymystax tsinlingensis*.

2.5 Phylogenetic analysis

Protein sequences of 10 species *Esox lucius*, *Coregonus* sp. *'balchen'*

(GCA_902810595.1) (Christensen, et al. 2020), *Thymallus thymallus*

(GCA_023634145.1), *Oncorhynchus mykiss*, *Oncorhynchus nerka*, *Salvelinus*

namaycushs (GCF_016432855.1) (Smith, et al. 2022), *Salmo salar*, *Salmo trutta*

(GCF_901001165.1) (Hansen, et al. 2021), *H. bleekeri*, and *Brachymystax*

tsinlingensis were used to construct phylogenetic tree. The single copy orthologous

proteins among the 10 species were recognized by Orthfinder version 2.3.12 (Emms

and Kelly 2019) with "msa" method. After alignment using MUSCLE version 3.8.31

(Edgar 2004), the species tree was calculated through RAxML version 8.2.12

(Stamatakis 2014) with 1000 bootstraps. The expansions and contractions of gene

family was estimated using CAFE5 (Mendes, et al. 2020).

The divergence time among all these species were calculated using MCMCtree

program of PAML software version 4.9d (Yang 2007). Four fossil records were

downloaded from TimeTree (Kumar, et al. 2022) and calibrated the results, the

Markov chain Monte Carlo (MCMC) was sampled with every 1000 step and conduct

10000 samples, the pre- burnin number was set at 20000.

2.6 Demographic history estimation

The pairwise sequentially Markovian coalescent (PSMC) method version 0.6.5-r67

(Li and Durbin 2011) was utilized to predict the effective population size. Besides the

two wild hucho sample, we used two lenok samples with highest sequencing depths

form Heilongjiang (*B.tumernsis*) and Shaanxi (*B.tsinlingensis*). Due to the different

sequencing depth, we used different strategies to generate the diploid consensus

sequences for huhco and lenok, -D 60 for hucho and –D 20 for lenok at vcfutils.pl

packages of bcftools version 1.12 (Danecek, et al. 2021). The psmc was run with

parameters "-N25 -t30 -r4 -p 8+4+25*2+4+6" to control the divergence at left of the

results.

3 Results

3.1 Genome size estimation and assembly

A total of 426685475 pair-end reads were obtained from the Illumina NovaSeq-6000

platform, with a total length of 64.00 Gb. The genome size estimated by GCE (k=17)

was ~3.21 Gb with low heterozygous ratio 0.44% (Figure S1, Table S1). Based on the

estimated genome size, the PacBio HiFi and Illumina Hi-C sequencing produced

76.48 Gb (23.8×) and 350.63 Gb (109.23×) raw reads separately for genome assembly,

besides, 155.88 Gb resequencing data of the additional two wild sample (~22×,

average) were generated.

After *de novo* genome assembly using the HiFi reads, a 3.48 Gb primary genome

were generated which had 4373 high quality contigs (>2 Kb), and the contig N50 was

2.4 Mb (Table 1). After the Hi-C assisted genome assembly and scaffold

deduplication, the contigs were anchored into 44 chromosome-level scaffolds,

covered 95.26% total length of all scaffolds, and the scaffold N50 was 73.71 Mb

(Figure 1; Table 1). Closed to the estimated size, the final hucho genome had a total

size of 3.45 Gb with 41.99% GC ratio (Table 1). The BUSCO results showed that the

assembled genome included 95.3% complete genes of actinopterygii_odb9 database,

indicated the high completeness (Table S2).

3.2 Landscape of repeat sequences

The non-redundancy length of the total repeat sequence recognized by three

annotation methods was 2.60 Gb, covered 74.21% of the hucho genome, which had

the highest proportion of repetitive sequences among salmonids (Tables S3, S4 and

S5). The interspersed repeats reported by the Repeatmasker and LTR_FINDER

composed 36.18% of the genome, which had the lowest proportion among salmonids,

but the total length in hucho genome was similar to others (Tables S3 and S4). The

proportion of retroelements in hucho genome was 13.11%, with 7.03% LINEs and

5.98% LTR elements (Tables S3). The DNA transposons composed 9.93% of the

genome, and the Tc1-IS630-Pogo family was the most abundant TE like other

salmonids and covered 9.38% of the hucho genome (Tables S3).

The tandem repetitive sequences covered 44.15% of the hucho genome, had the

highest proportion among salmonids (Figure 2; Tables S5). The lengths of STRs and

VNTRs were 45561499 bp and 1553872291, which accounted for 1.30% and 44.40%

of the genome respectively, and 59.12% of STRs had less than 1 Kb distance away

from the surrounding VNTRs (Figure 2). The distribution of TR was not uniformly,

we found 29.44% regions in hucho genome with low TR ratios (One-tenth of the

genome-wide scale; Figure 2), and some chromosomes had an enrichment of VNTR

at their ends (Figure 2). The distribution of STRs in hucho genome was relatively

scattered, only the top 200 STRs arranged by length exceeded the average length

(16620 bp) of the HiFi reads, had a total length of 15.64 Mb (Figure S2a).

Dinucleotide repeats accounted for 81.53% of STRs (Tables S6), which was higher

than the average level for vertebrates (Adams, et al. 2016).We identified 3337 VNTRs

with lengths over 100 Kb, which had a total length of 0.97 Gb and covered 28.20% of

the genome (Figure S2b). In addition, the presence of TRs affected GC content in

local regions of the genome, but the GC content of the TR region was 41.16%, similar

to the whole genome range (Figure 2).

3.3 Gene annotation

A total of 40.00 Gb RNA reads were generated from five tissues for RNA-based gene annotation. After filtered out the pseudogenes, the BRAKER2 annotation pipeline predicted 49052 genes, with an average length of 14576.6 bp. We generated 338448 exons, the average number of exon in gene was 6.90. After mapping with the NR databases, only 79.02% of the predict genes were functionally annotated. The TR ratio in intron regions was 22.74% which significantly lower than whole genome level (Z-test; $P<0.01$). Based on the kernel density estimation, 161 gene hotspots were identified (Figure S3). The results of NR database annotation showed that the top three regions with the highest gene density were both histone gene family loci (chr8 and chr22). The result of BUSCO for the predicted genes showed that 88.1% were complete BUSCOs (C:88.1%[S:43.3%,D:44.8%],F:4.3%,M:7.6%,n:4584).

3.4 Analysis of phylogenetic and evolution of gene family

We identified 36292 orthologous gene families between hucho and the other 9 species, including 479 hucho specific gene families and 267 single copy orthologues. The result of phylogenetic tree based on single copy orthologues revealed that the hucho was a sister group to lenok, and the divergent time estimated by PAML was 12.29 Mya between this two species (Figure 3), which was closed to previous results based on mitochondria (Kucinski and Fopp‐Bayat 2022). We generated 916 significant positive expanded gene families in hucho genome, and the results of KEGG pathway showed the enrichments the enrichment of motor and metabolic pathways such as tight junction, adrenergic signaling in cardiomyocytes, glyoxylate and dicarboxylate metabolism and regulation of actin cytoskeleton (Figure S4; Table S7), which was fits

239 with its ferocious hunting habits and large body size.

3.5 Analysis of demographic history

The geographical distribution pattern of *Brachymystax* is similar to that of *Hucho*, and the competitive relationship is existed between them (Kaus, et al. 2019; Galland, et al. 2021). Due to the relatively larger body size and fiercer temperament, the hucho dominated the niche and often preyed on lenok. In order to illustrate the trends of historical effective population sizes (N_e) change of hucho and lenok in Shaanxi and Heilongjiang, four samples of them were used for joint analysis. The result showed that the history of four species traced back to different times respectively, ~9 Mya for the two Heilongjiang sample *H. taimen* and *B.tumernsis*, and ~1 Mya for the two Shaanxi sample *H. bleekeri* and *B.tsinlingensis* (Figure 4). The *H. taimen* had a higher N_e level than *H. bleekeri* throughout the history (Figure 4). The curve suggested that a large scale population expansion event occurred at ~2 Mya in both *H. taimen* and *B.tumernsis*, and before ~250 Kya the N_e of *H. taimen* was greater than the *B.tumernsis* (Figure 4). On the contrary, the *B.tsinlingensis* had a larger N_e than *H. bleekeri* before ~150 Kya (Figure 4). In summary, the Heilongjiang populations had higher N_e level than Shannxi populations since ~200 Kya, although the biomass of hucho was currently smaller than that of lenok, there had been times in history with the opposite case, which reflected the competition results between them.

4.Discussion

The combination of HiFi and Hi-C reads allowed us to achieve the chromosome-level assembly of the largest salmonid genomes to date. The final size of hucho genome was ~3.45 Gb, which exceeds that of the largest Atlantic salmon genome (~2.97 Gb)

(Lien, et al. 2016). The chromosome count of the hucho genome was 44, and

approached to some salmonids, such as lake trout (42), *B. tsinlingensis* (41), and

brown trout (40) (Table S8). The assembled genome recovered 95.20% complete

BUSCO genes, of which the percentage of duplicated BUSCOs (46.20%) was

relatively high among salmonids (Smith, et al. 2022), but similar to its sister group *B.*

tsinlingensis.

The genome assembly process and the outcome were affected by the ratio of

repetitive sequences (Treangen and Salzberg 2011; Tørresen, et al. 2017). The highly

accurate reads produced by circular consensus sequencing technology resulted in a

significant improvement in both the process and the quality of the assembly,

especially in the repetitive sequences region (Wenger, et al. 2019; Logsdon, et al.

2020). We found that the assembled hucho genome contains a high proportion of

repetitive sequences, and nearly half region of the genome were covered by tandem

repeat sequence, which may contribute to the increased genome size. We discovered

that most of the TR regions were covered and assembled by the HiFi reads, which

revealed the merit of long reads. However, there were some long TRs account for a

relatively large of the total genome length, and causing many gaps around them.

Generally, the number of gaps in the assembled hucho genome was similar to or even

smaller than other salmonid genomes, except for GCF_013265735.2, which indicated

that our assembly was not severely disturbed by TRs (Table S8).

The existence of long repetitive regions can affect the results of gene annotation,

leading to the appearance of false and redundant genes (Lang, et al. 2020). In our

preliminary gene annotation results, we found a mass of end-to-end pseudogenes in

long TR regions, which seriously interfered with our subsequent analysis. Despite the

presence of TRs in specific domains of some proteins (Simon and Hancock 2009), we

found that each of these pseudogenes had only one exon according to our manual

inspection, so we filtered out these single-exon genes located within the repetitive

sequence region based on expression (TPM<1). In parallel, we also blasted these

pseudogenes to demonstrate their absence in other species.

Salmonidae contains three subfamilies Coregoninae, Thymallinae, and

Salmoninae, but the evolutionary relationships remains uncertain (Dysin, et al. 2022).

Lineage-specific rediploidization, incomplete lineage sorting, hybridization, and

introgression resulted in the disordered messy phylogenetic relationship of

Salmonidae (Robertson, et al. 2017; Lecaudey, et al. 2018; Campbell, et al. 2020).

Some studies suggested that Thymallinae is the sister group of Salmoninae (McKay,

et al. 2004; Wang, et al. 2011; Crête-Lafrenière, et al. 2012), while some studies put

Thymallinae in the root of the Salmonidae phylogenetic tree (Horreo 2017; Wang, et

al. 2022). Based on the phylogenetic tree constructed by the orthologous genes, we

suggested that Coregoninae and Thymallinae had sister relationship which supported

by 100% bootstraps, and was consistent with other trees based on orthologues

(Macqueen and Johnston 2014; Robertson, et al. 2017), mitochondrial genome

(Campbell, et al. 2013; Shedko, et al. 2013), and ultra-conserved element (Campbell,

et al. 2020). The topology inner Salmoninae we inferred was consistent with other

studies, the *Hucho* and *Brachymystax* formed the ancient branch of the subfamily

salmoninae, and constituted the sister group to the genera *Salmo, Salvelinus and Oncorhynchus* (Crespi and Fulton 2004; Kucinski and Fopp‑Bayat 2022). The estimated divergence time based on nuclear genome was similar to the previous researches based on mitochondrial genome (Crête-Lafrenière, et al. 2012; Horreo 2017; Kucinski and Fopp‑Bayat 2022). The divergence time between the *Hucho* and *Brachymystax* was estimated at ~9.28 to ~17.96 Mya, which coincides with the Miocene Climate Optimum (MCO) at 15-18 Mya (Böhme 2003; Sun and Zhang 2008).

All species of genera *Hucho* and *Brachymystax* were endangered and were protected by Chinese government (Sung, et al. 1998), the results of PSMC revealed that the effective populations size of four *Hucho* and *Brachymystax* species had declined over 1 million years. The similar patterns of N_e curves were observed from the species in same geographical distribution. The curves of two species *H. bleekeri* and *B.tsinlingensis* in Shannxi had closed coalescent ending point, we speculated that this two salmonids shared the same immigration route during the glacial period and were retained in the low-latitude area at the interglacial period. The *H. bleekeri* had lower N_e level than *H. taimen* in our calculation, and it was very rare in the current resource survey in the Qinling Mountains, which had disappeared from most of the original distribution areas (Shen, et al. 2006; Wang, et al. 2016). Consequently, more effective measures should be taken to promote the conservation of *Hucho bleekeri*.

Author contributions

Xinmiao Zhang, and Lixin Wang conceived and supervised the project. Dongmei

328   Xiong, and Shenglong Jian the samples. Xinmiao Zhang, and Dongmei Xiong carried

329   out the bioinformatic analyses. Xinmiao Zhang wrote the manuscript. Shenglong Jian,

330   Yu Jiang and Lixin Wang revised the manuscript. All authors have read and approved

331   the final manuscript.

Competing interests

333   The authors declare no competing interests.

Acknowledgements

This study was financially supported by the National Natural Science Foundation of

China (32273138) and the latitudinal project foundation of China Power Construction

Group Beijing Survey, Design and Research Institute Co., Ltd. (K4050422184) . We

thank the High-Performance Computing Platform of Northwest A&F University and

the National Supercomputing Center in Xi'an and Hefei for providing computing

resources.

Reference

Adams RH, Blackmon H, Reyes-Velasco J, Schield DR, Card DC, Andrew AL, Waynewood N, Castoe TA. 2016. Microsatellite landscape evolutionary dynamics across 450 million years of vertebrate genome evolution. Genome 59:295-310.

Allendorf FW, Hard JJ. 2009. Human-induced evolution caused by unnatural selection through harvest of wild animals. Proc Natl Acad Sci U S A 106 Suppl 1:9987-9994.

Allendorf FW, Hohenlohe PA, Luikart G. 2010. Genomics and the future of conservation genetics. Nat Rev Genet 11:697-709.

Benson G. 1999. Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res 27:573-580.

Böhme M. 2003. The Miocene climatic optimum: evidence from ectothermic vertebrates of Central Europe. Palaeogeogr, Palaeoclimatol, Palaeoecol 195:389-401.

Braasch I, Gehrke AR, Smith JJ, Kawasaki K, Manousaki T, Pasquier J, Amores A, Desvignes T, Batzel P, Catchen J, et al. 2016. The spotted gar genome illuminates vertebrate evolution and facilitates human-teleost comparisons. Nat Genet 48:427-437.

Brandies P, Peel E, Hogg CJ, Belov K. 2019. The value of reference genomes in the conservation of threatened species. Genes 10(11): 846.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol 34:525-527.

Campbell MA, Buser TJ, Alfaro ME, López JA. 2020. Addressing incomplete lineage sorting and paralogy in the inference of uncertain salmonid phylogenetic relationships. PeerJ 8:e9389.

Campbell MA, López JA, Sado T, Miya M. 2013. Pike and salmon as sister taxa: detailed intraclade resolution and divergence time estimation of Esociformes + Salmoniformes based on whole mitochondrial genome sequences. Gene 530:57-65.

Chen S, Zhou Y, Chen Y, Gu J. 2018. Fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics 34:i884-i890.

Chen Y, Yang H, Chen Y, Song M, Liu B, Song J, Liu X, Li H. 2021. Full-length transcriptome sequencing and identification of immune-related genes in the critically endangered *Hucho bleekeri*. Dev Comp Immunol 116:103934.

Chen Y, Yang H, Gong Q, Chen Y, Tu Q, Li H. 2020. Isolation and characterization of 34 SNP markers in *Hucho bleekeri*. Conserv Genet Resour 12:157-160.

Cheng H, Concepcion GT, Feng X, Zhang H, Li H. 2021. Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm. Nat Methods 18:170-175.

Christensen KA, Rondeau EB, Minkley DR, Sakhrani D, Biagi CA, Flores AM, Withler RE, Pavey SA, Beacham TD, Godin T, et al. 2020. The sockeye salmon genome, transcriptome, and analyses identifying population defining regions of the genome. PLoS One 15:e0240935.

Crespi BJ, Fulton MJ. 2004. Molecular systematics of Salmonidae: combined nuclear

data yields a robust phylogeny. Mol Phylogenet Evol 31:658-679.

Crête-Lafrenière A, Weir LK, Bernatchez L. 2012. Framing the Salmonidae family
phylogenetic portrait: a more complete picture from increased taxon sampling.
PLoS One 7:e46662.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,
Keane T, McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and
BCFtools. Gigascience 10(2): giab008.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics
29:15-21.

Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS,
Machol I, Lander ES, Aiden AP, et al. 2017. De novo assembly of the *Aedes*
aegypti genome using Hi-C yields chromosome-length scaffolds. Science
356:92-95.

Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, Aiden EL.
2016. Juicebox provides a visualization system for Hi-C contact maps with
unlimited zoom. Cell Syst 3:99-101.

Durand NC, Shamim MS, Machol I, Rao SS, Huntley MH, Lander ES, Aiden EL.
2016. Juicer provides a one-click system for analyzing loop-resolution Hi-C
experiments. Cell Syst 3:95-98.

Dysin AP, Shcherbakov YS, Nikolaeva OA, Terletskii VP, Tyshchenko VI,
Dementieva NV. 2022. Salmonidae genome: features, evolutionary and
phylogenetic characteristics. Genes 13(12): 2221.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and
high throughput. Nucleic Acids Res 32:1792-1797.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for
comparative genomics. Genome Biol 20:1-14.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. 2020.
RepeatModeler2 for automated genomic discovery of transposable element
families. Proc Natl Acad Sci U S A 117:9451-9457.

Gabriel L, Hoff KJ, Brůna T, Borodovsky M, Stanke M. 2021. TSEBRA: transcript
selector for BRAKER. BMC Bioinf 22(1): 1-12.

Galland LM, Simmons JB, Jahner JP, Luzuriaga-Neira AR, Sloat MR, Chandra S,
Hogan Z, Jensen OP, Parchman TL. 2021. Hierarchical genetic structure and
implications for conservation of the world's largest salmonid, Hucho taimen. Sci
Rep 11:20508.

Gao G, Magadan S, Waldbieser GC, Youngblood RC, Wheeler PA, Scheffler BE,
Thorgaard GH, Palti Y. 2021. A long reads-based *de-novo* assembly of the
genome of the Arlee homozygous line reveals chromosomal rearrangements in
rainbow trout. G3 11(4): jkab052.

Hansen T, Fjelldal PG, Lien S, Smith M, Corton C, Oliver K, Skelton J, Betteridge E,
Doulcan J, Fedrigo O, et al. 2021. The genome sequence of the brown trout,
Salmo trutta Linnaeus 1758. Wellcome Open Res 6:108.

Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2016. BRAKER1:

unsupervised RNA-seq-based genome annotation with GeneMark-ET and
AUGUSTUS. Bioinformatics 32:767-769.

Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. 2019. Whole-genome annotation
with BRAKER. Methods Mol Biol 1962:65-95.

Horreo JL. 2017. Revisiting the mitogenomic phylogeny of Salmoninae: new insights
thanks to recent sequencing advances. PeerJ 5:e3828.

Hu M, Wang Y, Cao L, Xiong BJEbof. 2008. Threatened fishes of the world: *Hucho*
bleekeri Kimura, 1934 (Salmonidae). Environ Biol Fishes 82:385-386.

Ishiguro NB, Miya M, Nishida M. 2003. Basal euteleostean relationships: a
mitogenomic perspective on the phylogenetic reality of the "Protacanthopterygii".
Mol Phylogenet Evol 27:476-488.

Kaus A, Michalski S, Hänfling B, Karthe D, Borchardt D, Durka W. 2019. Fish
conservation in the land of steppe and sky: Evolutionarily significant units of
threatened salmonid species in Mongolia mirror major river basins. Ecol Evol
9:3416-3433.

Kohn MH, Murphy WJ, Ostrander EA, Wayne RK. 2006. Genomics and conservation
genetics. Trends Ecol Evol 21:629-637.

Kucinski M, Fopp-Bayat DJJoAI. 2022. Phylogenetic analysis of *Brachymystax* and
Hucho genera-Summary on evolutionary status within the Salmoninae subfamily.
J Appl Ichthyol 38:403-411.

Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, Stecher G,
Hedges SB. 2022. TimeTree 5: an expanded resource for species divergence
times. Mol Biol Evol 39(8): msac174.

Lang D, Zhang S, Ren P, Liang F, Sun Z, Meng G, Tan Y, Li X, Lai Q, Han L, et al.
2020. Comparison of the two up-to-date sequencing technologies for genome
assembly: HiFi reads of Pacific Biosciences Sequel II system and ultralong reads
of Oxford Nanopore. Gigascience 9(12): giaa123.

Lecaudey LA, Schliewen UK, Osinov AG, Taylor EB, Bernatchez L, Weiss SJ. 2018.
Inferring phylogenetic structure, hybridization and divergence times within
Salmoninae (Teleostei: Salmonidae) using RAD-sequencing. Mol Phylogenet
Evol 124:82-99.

Li H, Durbin R. 2011. Inference of human population history from individual
whole-genome sequences. Nature 475:493-496.

Lien S, Koop BF, Sandve SR, Miller JR, Kent MP, Nome T, Hvidsten TR, Leong JS,
Minkley DR, Zimin A, et al. 2016. The Atlantic salmon genome provides insights
into rediploidization. Nature 533:200-205.

Liu B, Shi Y, Yuan J, Hu X, Zhang H, Li N, Li Z, Chen Y, Mu D, Fan W. 2013.
Estimation of genomic characteristics by analyzing k-mer frequency in *de novo*
genome projects. arXiv.org arXiv:1308.2012.

Liu X, Zeng H, Wang C, Bo J, Gan X, Fang C, He S. 2022. Improved genome
assembly of Chinese sucker (*Myxocyprinus asiaticus*) provides insights into the
identification and characterization of pharyngeal teeth related maker genes in
Cyprinoidei. Water Biology and Security 1(3): 100049.

Logsdon GA, Vollger MR, Eichler EE. 2020. Long-read human genome sequencing

and its applications. Nat Rev Genet 21:597-614.

Macqueen DJ, Johnston IA. 2014. A well-constrained estimate for the timing of the salmonid whole genome duplication reveals major decoupling from species diversification. Proc Biol Sci 281:20132881.

McKay SJ, Trautner J, Smith MJ, Koop BF, Devlin RH. 2004. Evolution of duplicated growth hormone genes in autotetraploid salmonid fishes. Genome 47:714-723.

Mendes FK, Vanderpool D, Fulton B, Hahn MW. 2020. CAFE 5 models variation in evolutionary rates among gene families. Bioinformatics 36:5516-5518.

Ou S, Jiang N. 2018. LTR_retriever: a highly accurate and sensitive program for identification of long terminal repeat retrotransposons. Plant Physiol 176:1410-1422.

Robertson FM, Gundappa MK, Grammes F, Hvidsten TR, Redmond AK, Lien S, Martin SAM, Holland PWH, Sandve SR, Macqueen DJ. 2017. Lineage-specific rediploidization is a mechanism to explain time-lags between genome duplication and evolutionary diversification. Genome Biol 18: 1-14.

Sadovy de Mitcheson Y, Craig MT, Bertoncini AA, Carpenter KE, Cheung WW, Choat JH, Cornish AS, Fennessy ST, Ferreira BP, Heemstra PCJF. 2013. Fishing groupers towards extinction: a global assessment of threats and extinction risks in a billion dollar fishery. Fish and fisheries 14:119-136.

Shedko SV, Miroshnichenko IL, Nemkova GA. 2013. Phylogeny of salmonids (Salmoniformes, Salmonidae) and molecular dating: analysis of mtDNA data. Genetika 49:718-734.

Shen Z, Tang W, Li K. 2006. The analysis of population dynamics of *Hucho bleekeri* in Markehe River, Qinghai Province. Reserv Fisheries 26:71-73 (In Chinese).

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics 31:3210-3212.

Simon M, Hancock JM. 2009. Tandem and cryptic amino acid repeats accumulate in disordered regions of proteins. Genome Biol 10(6): 1-16.

Smith SR, Normandeau E, Djambazian H, Nawarathna PM, Berube P, Muir AM, Ragoussis J, Penney CM, Scribner KT, Luikart G, et al. 2022. A chromosome-anchored genome assembly for Lake Trout (*Salvelinus namaycush*). Mol Ecol Resour 22:679-694.

Song Z. 2012. *Hucho bleekeri*. The IUCN Red List of Threatened Species. Version 2014.3.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30:1312-1313.

Stefanova P, Taseva M, Georgieva T, Gotcheva V, Angelov A. 2013. A modified CTAB method for DNA extraction from soybean and meat products. Biotechnol Biotechnol Equip 27:3803-3810.

Sun J, Zhang Z. 2008. Palynological evidence for the Mid-Miocene Climatic Optimum recorded in Cenozoic sediments of the Tian Shan Range, northwestern China. Global and Planetary Change 64:53-68.

Sung W, Peiqi Y, Yiyu C, Commission ESS. 1998. China Red data book of

endangered animals: Pisces. Science Press, Beijing, Hong Kong, New York, pp 32–34 (In Chinese).

Tarailo-Graovac M, Chen N. 2009. Using RepeatMasker to identify repetitive elements in genomic sequences. Curr Protoc Bioinformatics 4:1-4.10.

Tørresen OK, Star B, Jentoft S, Reinar WB, Grove H, Miller JR, Walenz BP, Knight J, Ekholm JM, Peluso P, et al. 2017. An improved genome assembly uncovers prolific tandem repeats in Atlantic cod. BMC Genomics 18:95.

Treangen TJ, Salzberg SL. 2011. Repetitive DNA and next-generation sequencing: computational challenges and solutions. Nat Rev Genet 13:36-46.

Wang K, Zhang SH, Wang DQ, Wu JM, Wang CY, Wei QW. 2016. Conservation genetics assessment and phylogenetic relationships of critically endangered *Hucho bleekeri* in China. J Appl Ichthyol 32:343-349.

Wang Y, Guo R, Li H, Zhang X, Du J, Song Z. 2011. The complete mitochondrial genome of the Sichuan taimen (*Hucho bleekeri*): repetitive sequences in the control region and phylogenetic implications for Salmonidae. Mar Genomics 4:221-228.

Wang Y, Xiong F, Song Z. 2022. Molecular Phylogeny and adaptive mitochondrial DNA evolution of Salmonids (Pisces: Salmonidae). Front Genet 13:903240.

Waples RS, Naish KA, Primmer CR. 2020. Conservation and management of salmon in the age of genomics. Annu Rev Anim Biosci 8:117-143.

Wenger AM, Peluso P, Rowell WJ, Chang PC, Hall RJ, Concepcion GT, Ebler J, Fungtammasan A, Kolesnikov A, Olson ND, et al. 2019. Accurate circular consensus long-read sequencing improves variant detection and assembly of a human genome. Nat Biotechnol 37:1155-1162.

Xu Z, Wang H. 2007. LTR_FINDER: an efficient tool for the prediction of full-length LTR retrotransposons. Nucleic Acids Res 35:W265-W268.

Xue DX, Xing TF, Liu JX. 2022. A high-quality chromosome-level genome of the endangered roughskin sculpin provides insights into its evolution and adaptation. Mol Ecol Resour 22:1892-1905.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol Biol Evol 24:1586-1591.

Ye J, McGinnis S, Madden TL. 2006. BLAST: improvements for better sequence analysis. Nucleic Acids Res 34:W6-9.

Zhang Y, Luan P, Ren G, Hu G, Yin J. 2020. Estimating the inbreeding level and genetic relatedness in an isolated population of critically endangered Sichuan taimen (*Hucho Bleekeri*) using genome-wide SNP markers. Ecol Evol 10:1390-1400.

Table

TABLE 1 Summary statistics of the Chuanshan hucho (*Hucho bleekeri*) genome assembly.

Category	Length (bp)		Number	
	Contig	Scaffold	Contig	Scaffold
Max	25,857,850	143,121,144	-	-
Min	3,656	18,648	-	-
N50	2,400,826	73,705,547	354	19
N90	418,871	59,067,955	1680	39
Total	3,479,724,502	3,451,243,059	4373	505

Figures

Figure 1 Hi-C interaction heatmap of *Hucho bleekeri* genome between 44 chromosomes and unmapped scaffolds. The intensity of red indicates the interaction strength. The 44 chromosome-level scaffolds are represented by blue outlines, and the preliminarily assembled contigs are shown by green outlines.

Figure 2 Visualization of genomic GC content, STR, and VNTR distribution based on 100 Kb windows. The bar graph reflects the TR frequency distribution in genome-wide 100 Kb windows.

Figure 3 Maximum likelihood tree constructed based on single-copy orthologous genes from 9 salmonids and outgroup (*Esox lucius*). Fossil calibration points used by MCMC are marked in red. Black numbers next to each nodes are the divergence times (million years ago, MYA) and the 95% confidential intervals estimated by PAML. The green positive and red negative integers indicate the number of genes at expanded or contracted gene families at that node, respectively.

Figure 4 Demographic history of two Heilongjiang sample (*H. taimen* and *B.tumernsis*) and two Qinling sample (*H. bleekeri* and *B.tsinlingensis*). The generation (g) time was set at 4 years and the neutral mutation rate per generation (μ) was set at 6.54 × 10-9.

Supplementary figure and table legend

Supplementary Figure S1. Frequency distribution map in 17-mer.

Supplementary Figure S2. Statistics of tandem repeat unit length and total tandem repeat sequence length of STRs (a) and VNTRs (b).

Supplementary Figure S3. Gene hotspots of the hucho genome.

Supplementary Figure S4. Visualization of the KEGG pathway annotation of the 1472 positive selected genes.

Supplementary Table S1. Genome size estimation table at k=17.

Supplementary Table S2. Summarized benchmarking in BUSCO notation.

Supplementary Table S3. Summary of TE in 10 salmonids (%).

Supplementary Table S4. Summary of TE in 10 salmonids (length).

Supplementary Table S5. Statistics of TR for 10 salmonids.

Supplementary Table S6. Summary of STRs' unit and total length.

Supplementary Table S7. The KEGG pathway annotation of the 1472 positive selected genes.

Supplementary Table S8. Summary of genome assembly results of 10 salmonids.