The use of threshold methods to detect beta bursts may inadequately characterize the beta band

Zachary D Langford[1*], Charles R E Wilson[1]

[1]Univ Lyon, Université Claude Bernard Lyon 1, Inserm, Stem Cell and Brain Research Institute U1208, France

*corresponding author

ABSTRACT

In neurophysiological research, the traditional view of beta band activity as sustained oscillations is being reinterpreted as transient bursts. Bursts are characterized by a distinct wavelet shape, high amplitude, and, most importantly, brief temporal occurrence. The primary method for their detection relies on a threshold-based analysis of spectral power, and this presents two fundamental issues. First, the threshold selection is effectively arbitrary, being influenced by both local and global factors in the signal. Second, the method necessarily detects temporal events, as such it is susceptible to misidentifying sustained signals as transient bursts. To address these issues, this study systematically explores burst detection through simulations, shedding light on the method's robustness across various scenarios. Although the method is effective in detecting transients in numerous cases, it can be overly sensitive, leading to spurious detections. Moreover, when applied to simulations featuring exclusively sustained events, the method frequently yields events exhibiting characteristics consistent with a transient burst profile. This capacity to produce misleading outcomes challenges the reinterpretation of sustained beta oscillations as transient bursts and prompts a critical reassessment of the existing literature.

INTRODUCTION

Beta bursts are transient spectral events, occurring between 15 and 29 Hz, in neurophysiological recordings. Bursts are usually characterized as being of relatively high amplitude, occurring briefly (<300 ms), and having the distinct appearance of a wavelet shape (Sherman *et al.*, 2016; Szul et al. 2022; Zich et al. 2023). Beta bursts have been linked to sensorimotor cortex during motor actions (Wessel 2020; Bonaiuto *et al.*, 2021; Zich et al. 2023; Rayson *et al.*, 2023), prefrontal cortex during working memory tasks (Lundqvist et al. 2018; Liljefors *et al.*, 2023), and frontal cortex in cognitive control tasks (Langford *et al.*, 2023). However, despite their functional relevance, the precise nature and underlying mechanisms of beta bursts remain incompletely understood, and this may largely be determined by the methodologies employed to characterize them.

The most common method used to detect bursts is based on identifying a threshold (PT-method), or cutoff point, in power - at which any value occurring above this threshold is considered a putative burst (e.g., Shin et al. 2017) if it fulfills a number of other criteria. The threshold is often chosen as the point that maximizes the correlation between the single-trial area above the threshold and averaged single-trial power. This threshold is contended to be appropriate because it most fully accounts for the variance in single trial power (Shin et al. 2017), yet in of itself this contention makes the strong assumption that the transient bursts are the only factor contributing to single trial power. It has further become commonplace to employ a predetermined threshold, commonly set at six factors-of-medians (FOM), across a gamut of experimental designs, recording modalities, cortical locations, and species

(e.g., Wessel et al. 2020; Morris *et al.*, 2023; Errington *et al.*, 2020; Law *et al.*, 2022; and additionally in the gamma band: McKeon *et al.*, 2023).

In this study, our aim was to conduct a comprehensive investigation into the performance of the PT-method using simulated data. This strategy is particularly useful because it enables the modeling of numerous plausible burst scenarios, allowing us insight into the ground truth of each, which is not attainable through neural data alone. We narrow our focus on addressing what we view as two unresolved issues. First, the establishment of an optimal cutoff using correlation maximization appears to harbor an element of arbitrariness. This stems from the fact that the identification of individual bursts hinges on the calculation of medians across experimental data spans, regardless of the content of those spans, and whether they are readily comparable between recordings, individuals, or conditions. Specifically the process is susceptible to the signal-to-noise ratio (SNR), to the presence of sustained signals, and to the presence of both temporally proximate and distant bursts, and their associated amplitude distributions. To address this, we conducted an examination by simulating a medium enriched with bursts, elucidating the PT-method's proficiency in recovering intended bursts and the inadvertent retrieval of unintended ones. Second, any plausible neurophysiological data will necessarily produce a median and the requisite single-trial maximum correlation, consequently detecting events with a set of characteristics to be classified as either transient or sustained. Hence, the potential of the methodology to erroneously identify transient events within signals of a sustained nature merits consideration. To confront this issue, we conducted a second simulation, presuming a substrate characterized by oscillatory and sustained behavior. This analysis focused on the classification of event characteristics, with the aim of affirming or negating our assumption of a sustained medium's presence, based solely on the recovered features.

We establish that the PT-method exhibits robust performance in detecting transient bursts when a relatively high SNR prevails. But its efficacy wanes significantly under conditions of relatively low SNR, leading to the inadvertent detection of numerous undesired events. Furthermore, our examination reveals that when applied to simulations that contain **only** sustained events, the method frequently registers beta events that are clearly within the duration of a burst event, even in cases of a high SNR. That is, if the brain does produce either sustained or transient beta events, analyzing them with the PT method will necessarily favor a transient interpretation. Lastly, we searched across all simulations for datasets that produce a single-trial correlation maximum at six FOM and detailed the extent to which we can differentiate the varied data generation processes based solely on the attributes of the recovered bursts. We conclude that the method has a clearly limited discriminative capacity, in that it consistently yields distributions of burst characteristics that do not easily permit us to draw a distinction between event generation processes. As such determining whether a set of bursts originates from a ground truth process of a burst-filled medium, a largely noise-filled medium, or a medium filled with sustained events is thus a highly challenging task. Of course, this lack of discriminative capacity is particularly problematic in the analysis of real neural data, where we have no access to any form of ground truth.

METHODS

Simulations

To create the simulations we embedded either short duration wavelets or long duration sine waves, in continuous coloured noise with a slope of 2 using NeuroDSP (Cole *et al.*, 2019). The simulations and analyses are all freely available online at https://github.com/langfordzd/burst_simulations. There were 100 trials per simulation, which included 1 second of burst-activity with noise, surrounded by a 1/2

second of pure noise. The sine-waves were asymmetric with rise-decay symmetry uniformly drawn at random between 0.25 and 0.75. The wavelets were created using the morlet2 function in the python sciPy module (Virtanen *et al.*, 2020) with the whole wavelet being scaled by the amplitude, i.e., the center point of the wavelet was the peak amplitude. In either the wavelet or sine-wave instantiation we randomly flipped the sign of the signal so it would either start with an ascent or descent. The burst frequencies were always drawn from a truncated normal distribution with a mean of 21, a standard deviation of 1, and being truncated between a low of 15 and a high of 29 Hz. The SNR was calculated as the ratio of mean spectral power for the 0.5 seconds in the middle of the trials (the burst-activity period) and 0.5 seconds before the trial (the initial noise period), within the defined beta band.

In these simulations the main parameters of interest were the burst probabilities, the burst durations, and the burst amplitude distributions. The burst probability was manipulated by drawing randomly from a weighted binary distribution at each time-step to determine if that time-step would start a burst or not – with there being at the least 1 cycle between bursts. The probability of the next time step was specifically weighted to 0.9 in the high, 0.15 in the middle, and 0.03 in the low probability manipulations of the transient event simulations. The duration of the sustained oscillations were uniformly drawn between 0.5 and 0.8 seconds for both the wavelet and sine wave variants. The probability of the next time step was specifically weighted to 0.9 in the high, 0.1 in the low probability manipulations of the sustained simulations. For the transient event simulations we used a lognormal distribution with shape of 0.5, location of 0.04, and scale of 0.12, which we then truncated at 0.4 seconds to model the durations. This gave us a distribution similar to empirical research (Shin *et al.*, 2017) that starts around 0.04 seconds, peaks at 0.16 seconds, and tapers off with the highest values somewhere around 0.4 seconds. We simply used a uniform distribution between 0.5 and 0.8 seconds for the durations of the sustained simulations. The burst amplitude distributions were modeled again with a lognormal distribution, but we varied the parameters so as to produce significant variation in these amplitude properties and test the effects of the distribution on the outcomes. The impact on amplitude of variations in scale and shape parameters are illustrated in Fig 1B. The location parameter provides a scalar shift along the x-axis of the distribution, in our case therefore a shift in the amplitude. Location was equally spaced in 3 steps between 0.001 and 0.05. The shape parameter is also the standard deviation of the log of the distribution, and affects the general shape of the distribution as seen in Fig 1B. Here it was equally spaced in 9 steps from 0.1 to 0.9. Increasing the scale parameter will spread the distribution along the amplitude axis, and in fact the scale is the median of the distribution. Here the scale parameter was logarithmically spaced in 12 steps between 0.01 and 0.25.

Power-threshold detection method

In our instantiation of the PT-method we first translated the simulated signal into a time-frequency representation (TFR) where the signals are convolved with a complex Morlet wavelet at 7 cycles using MNE-Python (Gramfort et al. 2013). The TFR at any time-frequency point was then normalized by the simulation median power value at each frequency; i,e, it was put into factors of medians (FOM). The tested power thresholds ranged from 0.1 to 20 in 30 equally sized steps. For each simulation the best threshold was determined by calculating the maximum correlation between mean single trial power and the area, represented as a proportion of total TFR area per trial, above the cutoff (Shin et al. 2017). Following this, we found local maxima that were above the previously calculated cutoff. These were then considered burst peaks and were used as the dataset to determine the relevant characteristics for each simulation.

We were specifically interested in the power, duration, frequency, and frequency span of all of the recovered bursts. For power we simply took the power value of the detected local maxima, and this was

done similarly for frequency. Frequency span was calculated as the full-width half maximum (FWHM) surrounding the local peak in the frequency domain. In a similar vein, duration was calculated as the FWHM surrounding the local peak in the time domain. Following this we determined if each burst was either intended or not, and if more than one detected burst was being assigned to an intended burst we considered it a repeat. A detected burst was considered intended if it had *any* temporal overlap with a simulated burst, but not twice the temporal length. Repeat detections of intended bursts were collapsed with unintended bursts after an initial evaluation to see if it played a large role in the outcome, which it did not. We then calculated the total number of unintended bursts, the proportion of intended bursts recovered, and importantly for the sustained simulations, the total number of events detected under 0.3 seconds - a duration that we took as a reasonable value to be the transition from transient to sustained events. We then, for each recovered burst, found the maximum absolute value in the time domain signal, and sign flipped it if it was negative. These were then all aligned at the temporal location of the maximum and averaged, which gave us burst related potentials (BURPS; Langford *et al.*, 2023) of each simulation's collection of detected events.

RESULTS

We set out to evaluate the PT-method of burst detection using simulated data over a range of plausible parameters of event characteristics. We consider two specific types of simulation. In the first, we've assumed that the underlying beta-band process is filled with transient events (0.04-0.3 second durations) and tried to faithfully represent a variety of possibilities close to the burst characteristics that are found in the literature (Shin *et al.*, 2017; Morris *et al.*, 2023). In the second type of simulation we evaluated the results of the t-threshold method if it were applied to data with a beta-band process that was filled with sustained events (0.5-0.8 second durations), and not transient events. Here, we were interested in what would happen if we incorrectly apply a method to detect transients when they don't exist. To create the simulations we embedded either short duration wavelets, or long duration sine waves, in continuous fractional noise. In both types of simulations, importantly, we parameterize the amplitudes, the spread of the amplitude distributions, and the probability of oscillatory event occurrence. Taken together, these manipulations inform the signal-to-noise ratio of any given simulation.

Burst detection assuming that transient events populate the beta-band.

Figure 1 (A) Simulation results when transient events populate the beta band. *In each plot, the measure indexed at the top of the column is plotted across variations in shape and scale of the lognormal distribution that has been used to generate burst amplitude. Top row is low event probability, middle is middle event probability, and the bottom is high event probability; all of which are marked on the right of the graphic. Leftmost column is average recovery rate, middle-left is the average unintended count of the 100 trial simulation, middle-right is the average maximum correlation threshold, and the rightmost column is the SNR. Gray boxes indicate points in which the average of the maximum-correlation threshold over the simulations is approximately 6 FOM and indicate where you would most likely find an amplitude distribution that produces this threshold.* ***(B) Effects of shape and scale on the amplitude distributions.*** *Representation of the amplitude distributions for each of the shape and scale parameters held constant at the most extreme values of the other (which is indicated in the title of each graph). The location parameter of the distribution was conveniently set to zero, but the effects on the distribution can be thought as a simple shift along the amplitude axis.*

The purpose of this simulation was to model the characteristics of bursts found in the literature, assume that they are true, and then see the PT-method's reaction to different manipulations. Most notably the amplitude distribution and the burst rate are of interest, because in calculating the median cutoffs these should, a priori, have a large effect on the result. We modulated the burst-rate at three levels by weighting the probability of a random draw at each time step (the exact burst count distributions can be

seen in column 6 of Figure 3). The burst frequency was assumed to be a truncated normal distribution centralized at 21 Hz with a standard deviation of 1, and truncated between 15 and 29 Hz. The amplitude distribution was a lognormal distribution and we manipulated the location, shape, and scale; the effects of which can be seen in Figure 1B. The SNR then varies, most notably, over the scale parameter of the lognormal distribution, as well as the three steps on the location parameter. The overall duration of the bursts was also set as a lognormal distribution with shape set at 0.5, scale at 0.12, and location set to 0.04. This gave us a distribution similar to empirical research that starts around 0.04 seconds, peaks at 0.16 seconds, and tapers off with the highest values somewhere around 0.4 seconds (Shin *et al.*, 2017). The bursts themselves were modeled as wavelets as this has been the proposed shape in past work. In total, we simulated 20 datasets, each of which had the same noise structure across the different parameters varied in the amplitude distribution.

In this analysis presented in Fig 1 we see two general trends; the first is that when the amplitude distribution shape is low, across all scales, the method recovers the highest proportion of intended bursts for a given probability distribution. Conversely, as the shape increases the amount of bursts recovered decreases. The second is that as the scale of the distribution increases (increasing the central tendency of the amplitudes) the amount of unintended bursts decreases, for any given shape parameter. Unsurprisingly maybe, this means that as the burst amplitude (SNR) goes up, the chances of an unintended burst goes down, but this relationship is modulated by the spread of the distribution. Going along with this, the maximum correlation threshold both increases as the shape increases, and increases as the scale increases. In Figure 1 the average maximum-correlation threshold at approximately 6 FOM manifests as a diagonal line with a downward-shift in the x-axis, which is the scale parameter of the amplitude distribution. Figure 1 further indicates that the SNR is relatively low when 6 FOM is the correct maximum. In most instances where 6 FOM is applied, the method consistently achieves a high recovery rate, generally exceeding 50%. Higher recovery rates are observed in simulations involving lower probability event detection. A notable challenge lies in the method's pronounced ability to recover unintended bursts. This phenomenon is evident across all probability levels but becomes more problematic in scenarios with low burst probability. This is attributed to the significantly higher ratio of intended to unintended bursts in such cases.

Burst detection assuming that sustained events populate the beta-band.

Figure 2. Simulation results when sustained events populate the beta band. *Top column is low event probability and the bottom is high event probability. Leftmost column is the count of recovered events that could be considered bursts, the middle is the average maximum correlation threshold, and the rightmost is the SNR. Gray boxes indicate points in which the average threshold is approximately 6 FOM.*

In the second type of simulation, we assessed the performance of the PT method when applied to data featuring a beta-band process characterized by sustained events, and not containing the transient events considered above. The key distinction from the transient event simulation lies in the fact that the bursts in this scenario consistently lasted between 0.5 and 0.8 seconds—always sustained. To elaborate, these sustained bursts ranged from a minimum of 7.5 cycles (0.5 seconds at 15 Hz) to approximately 23 cycles (0.8 seconds at 29 Hz), with the duration modeled as a uniform distribution.

For simplicity and to avoid redundancy, we employed only two probability levels, 0.1 or 0.9, recognizing that modeling an even lower probability event would yield results very similar to the lowest probability event in the transient event simulations (i.e., in both there would be very little real beta activity). The frequency distribution mirrored that of the transient simulations. The sustained nature of the signals inherently led to fewer oscillatory events compared to the transient simulation, given temporal constraints. To address the potential limitations of wavelets for sustained oscillations, we modeled the bursts as sine waves.

As discussed above, real neural signals are reported to contain transient beta bursts, but there is little consideration of whether such transients are the ***only*** instantiation of beta in the signal, or indeed whether more sustained beta processes might be interpreted as transient bursts under certain analysis conditions. As such we wanted to see how the threshold detection method will perform when presented ***only*** with sustained beta events. To test how the method interprets these sustained beta processes, our primary analysis focus shifted from recovering simulated and unsimulated bursts to assessing the characteristic of any recovered bursts that lasted for less than 0.3 seconds—our defined cutoff for a detected event to be considered transient. Correctly identifying sustained events was crucial for

evaluating the method's performance in these circumstances. While recovery of intended sustained events was less relevant to transient burst detection, we didn't exclude detected sustained events for analysis in the subsequent section.

Figure 2's leftmost column vividly illustrates that when applying the PT-method to sustained oscillations, it detects <0.3-second transients in the sine wave instantiations, even though these were not included in the simulated data. At the low probability level, the detection pattern on the shape-scale surface closely resembles that of the transient event simulations, reflecting the same minimal oscillatory activity. However, at the high probability level, a distinct profile emerges suggesting that even with a very high signal-to-noise ratio (SNR), the method detects numerous transients.

Evaluating the characteristics of the recovered bursts.

Figure 3. Comparison of the burst characteristics for each simulation type show remarkable similarity. *Row 1-3 are transient event simulations at Low (1), Mid (2), Hi (3) burst probability levels. Row 4-5 are sustained event simulations at Low (4) and Hi (5) burst probability level. Columns 1-7. (1) is the correlation at each given threshold showing a peak roughly around 6 FOM. 2, 3, and 4 are the power, frequency span, and duration distributions of the recovered bursts. Grey dashed line marks the median. (5) is the trial count of the number of detected bursts for each of the datasets, while (6) is the simulated or intended distribution of the same. (7) is the maximum centered burst related potentials (BURPS) for each of the data sets that fell around a 6 FOM maximum. Sustained events lasted between 0.5 seconds and 0.8 seconds, whereas transient bursts were modeled to be brief 'burst' events.*

We have therefore demonstrated a capacity of threshold methods to capture unintended bursts in transient simulations and transient events within sustained simulations. As a result, there is an

important need for a manner to validate the plausibility of all of the captured bursts. To do so we now consider the specific characteristics of these bursts captured from the simulated data where we have an established ground truth. Given the prevailing use of a threshold maximum correlation set at 6 FOM in past observations, we focused on simulations where the peak correlation fell within the range of 5.5 to 6.5 FOM.

Figure 3 compiles data from simulations within this peak correlation range, facilitating a qualitative comparison of burst characteristics across various simulated substrates. Our analysis delved into four key characteristics—burst power, frequency span, duration, and burst counts—commonly explored in the literature. Surprisingly, our findings revealed that qualitatively distinguishing between the five types of oscillatory event simulations was not straightforward. Notably, if there is abundant oscillatory signal in the simulation, the median amplitudes of the bursts tend to be higher, even with the same 6 FOM cutoff. This is likely due to the ample signal elevating the median calculation during the FOM procedure and thus possibly contributing to the lower recovery rates in high-probability transient simulations.

In the frequency domain, we observed minimal differences among simulation types, except in high-probability sustained event simulations, which exhibited a narrower frequency range. This could be explained by fewer bursts overall, albeit with a higher SNR. Interestingly, there was little disparity in the durations of bursts extracted from the different simulations, although sustained simulations appeared to have broader tails in their distributions. Further scrutiny of the threshold correlation profiles and burst count histograms failed to reveal distinct features associating a specific profile with a particular substrate. While some differences were discernible across substrates, a definitive signal of a burst's origin—from a sustained signal, noise-dominated signal, or one rich in oscillatory signals—remained elusive.

To deepen our analysis, we aligned each detected burst at the largest deflection, incorporating sign-flipping where necessary, and generated averaged burst-related potentials (BURPS) visible in column 7 of Figure 3. BURPS provide a unique perspective on the average rhythmicity of the detected bursts. In both the high-probability burst-filled substrate and the high-probability sustained substrate, the averaged plots displayed a somewhat more rhythmic pattern, characterized by visually more zero-crossings. However, while this rhythmicity provided insight into the generative substrate for each simulation type, again each was subject to be interpreted as a plausible 'wavelet-like' waveform of bursts within the broader literature; from very brief ricker-like wavelets (e.g., Sherman *et al.*, 2016) to more temporally extended waveforms (e.g., Karvat *et al.*, 2020).

DISCUSSION

Here we have assessed the effectiveness of the widely used thresholding method in detecting beta bursts using simulated neural signals. We manipulated the SNR by varying burst-rate probability and burst amplitudes in simulated data. We conducted two distinct types of simulation, each based on different assumptions. In the first type, where a "bursty" substrate was assumed, we examined the PT-method's ability to recover intended bursts while also evaluating its capacity to identify unintended bursts. The method's performance hinged on the shape and scale of amplitude distributions, along with the prevalence of transient events. The method recovers intended events at a reasonable rate, yet is prone to detecting an over-abundance of unintended events. In the second type of simulation, assuming a substrate without bursts but featuring sustained oscillations, our results were more concerning. Even in cases of clearly prolonged events and a high SNR, the method demonstrated a tendency to recover numerous short-duration events that could be misinterpreted as bursts. Subsequent analysis of each simulation type emphasized that while the PT-method may perform well

under specific conditions, differentiating between bursty signals, predominantly colored noise, and sustained signals based on the recovered characteristics presents a considerable challenge. This challenges the reliability of research utilizing the PT-method and urges a reevaluation of its application.

Given the challenges and limitations associated with the PT-method in accurately identifying and characterizing beta bursts, it becomes crucial to complement the analysis with analysis of behavioral correlates, when available. In instances where the method detects true bursts that coincide with behavioral events, investigating the precise temporal dynamics and examining the relationship between burst occurrence, burst properties, and behavior can provide meaningful insights into cognitive processes. However, it is equally important to remain cautious and consider alternative explanations. It is possible that the broader beta power dynamics or, possibly, even the underlying aperiodic noise could exhibit temporal patterns or fluctuations that may influence burst detection and apparent associations with behavior, e.g., it is known the beta power can fluctuate over entire recording sessions (Stoll *et al.*, 2016). One might want to devise methods to determine, for example, the stability of beta power or even of the slope of aperiodic activity over an entire neural recording. Such temporal dynamics could introduce complexities in the interpretation of burst-behavior relationships, highlighting the importance of carefully considering the broader context and potential confounding factors.

Crucially, even when the PT-method identifies bursts strongly correlated with behavior, asserting a high degree of confidence in characterizing the phenomenon as transient, rather than sustained, is unwarranted. Simply pinpointing these bursts doesn't guarantee their transient nature. Realistic neural signals inherently exhibit a median and corresponding single-trial maximum correlation, resulting in events that might superficially appear transient – a tendency observed in many cases. This issue stems from imperfections in the PT method. For sustained signals, setting a threshold too high may capture only the most powerful segment, potentially overlooking the signal's sustained nature. Essentially, the most powerful segments, often of shorter duration, excel at explaining variance in single-trial power compared to lower power and longer segments in a neural signal; and as such the threshold is set to high. In this context, we posit that the PT-method coerces numerous realistic scenarios into being incorrectly interpreted as being populated by transient events - a point overlooked in the broader literature. Therefore, before placing confidence in its outcomes, it becomes crucial to establish the equally difficult assumption that the data indeed comprises transient events rather than sustained ones.

Acknowledging this limitation underscores a twofold research imperative. Firstly, a critical re-evaluation of existing literature on beta bursts is essential, involving a meticulous examination to determine the transient nature of the phenomenon. This reassessment serves as a foundational step toward refining our knowledge of the beta band and its temporal dynamics. Thus the second research imperative, which we argue will provide a means to achieve the first , is the necessity for a comprehensive investigation using multiple methods. The utilization of diverse and complementary techniques, such as elaborating on past research using hidden Markov models (Heideman *et al.*, 2020) or adapting the cycle-by-cycle method (Cole and Voytek, 2019; Langford *et al.*, 2023), *might* facilitate the cross-validation of burst detection results and offer a more robust insight into the underlying neural dynamics.

Bibliography

Bonaiuto JJ, Little S, Neymotin SA, Jones SR, Barnes GR, Bestmann S. Laminar dynamics of high amplitude beta bursts in human motor cortex. Neuroimage 2021; 242: 118479.

Cole S, Donoghue T, Gao R, Voytek B. NeuroDSP: A package for neural digital signal processing. JOSS 2019; 4: 1272.

Cole S, Voytek B. Cycle-by-cycle analysis of neural oscillations. J Neurophysiol 2019; 122: 849–61.

Errington SP, Woodman GF, Schall JD. Dissociation of Medial Frontal β-Bursts and Executive Control. J Neurosci 2020; 40: 9272–82.

Heideman SG, Quinn AJ, Woolrich MW, van Ede F, Nobre AC. Dissecting beta-state changes during timed movement preparation in Parkinson's disease. Prog Neurobiol 2020; 184: 101731.

Karvat G, Schneider A, Alyahyay M, Steenbergen F, Tangermann M, Diester I. Real-time detection of neural oscillation bursts allows behaviourally relevant neurofeedback. Commun Biol 2020; 3: 72.

Langford ZD, Procyk E, Wilson CR. Frontal oscillatory beta bursts have rhythmically distinct regimes with differing functional relevance. BioRxiv 2023

Law RG, Pugliese S, Shin H, Sliva DD, Lee S, Neymotin S, et al. Thalamocortical Mechanisms Regulating the Relationship between Transient Beta Events and Human Tactile Perception. Cereb Cortex 2022; 32: 668–88.

Liljefors J, Almeida R, Rane G, Lundstrom JN, Herman PA, Lundqvist M. Distinct functions for beta and alpha bursts in gating of human working memory. BioRxiv 2023

McKeon SD, Calabro F, Thorpe RV, de la Fuente A, Foran W, Parr AC, et al. Age-related differences in transient gamma band activity during working memory maintenance through adolescence. Neuroimage 2023; 274: 120112.

Morris AT, Temereanca S, Zandvakili A, Thorpe R, Sliva DD, Greenberg BD, et al. Fronto-central resting-state 15-29 Hz transient beta events change with therapeutic transcranial magnetic stimulation for posttraumatic stress disorder and major depressive disorder. Sci Rep 2023; 13: 6366.

Rayson H, Szul MJ, El-Khoueiry P, Debnath R, Gautier-Martins M, Ferrari PF, et al. Bursting with Potential: How Sensorimotor Beta Bursts Develop from Infancy to Adulthood. J Neurosci 2023; 43: 8487–503.

Sherman MA, Lee S, Law R, Haegens S, Thorn CA, Hämäläinen MS, et al. Neural mechanisms of transient neocortical beta rhythms: Converging evidence from humans, computational modeling, monkeys, and mice. Proc Natl Acad Sci USA 2016; 113: E4885-94.

Shin H, Law R, Tsutsui S, Moore CI, Jones SR. The rate of transient beta frequency events predicts impaired function across tasks and species. BioRxiv 2017

Stoll FM, Wilson CRE, Faraut MCM, Vezoli J, Knoblauch K, Procyk E. The effects of cognitive control and time on frontal beta oscillations. Cereb Cortex 2016; 26: 1715–32.

Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods 2020; 17: 261–72.