Task dependent coarticulation of movement sequences

Hari Teja Kalidindi[1, 2], Frederic Crevecoeur[1, 2]

[1]Institute for Information and Communication Technologies, Electronics and Applied Mathematics (ICTEAM), Université Catholique de Louvain, Louvain-la-Neuve 1348, Belgium

[2]Institute of Neuroscience (IoNS), Université Catholique de Louvain, Brussels 1200, Belgium

Corresponding author:

Frederic Crevecoeur

4 avenue Georges Lemaitre

1348 Louvain-la-Neuve, Belgium

email : frederic.crevecoeur@uclouvain.be

Abstract

Classical theories propose that multiple goals of a sequence are specified simultaneously in the motor system, enabling each elements' execution to be influenced by others in the sequence, a process known as *coarticulation*. However, recent neural recordings suggest independent execution within circuits generating motor commands, with each individual goal separately specified to the controller. Consequently, the manner in which movement sequences are produced remains unclear. In a two-reaches sequence task, we tested the idea that both coarticulation and separation are related to flexible feedback control. More precisely, simulations revealed that an optimal feedback controller planning for multiple reaches at once can produce both separated and coarticulated sequences based on constraints imposed at intermediate goals. Human experiments confirmed that the coarticulation of sequential reaches was flexibly modulated by task instructions, matching simulation results closely. Moreover, the location of the second goal impacted long-latency muscle stretch responses to mechanical perturbations applied prior to the first goal, indicating that sequence combination was expressed in the sensorimotor network mediating fast feedback control. Overall, our results uncover the role of long-latency feedback pathway as a flexible control policy, supporting efficient sequence production without the need to individually specify each target one-by-one to the controller.

Significance

From preparing coffee to executing a tennis serve, our nervous system combines multiple elementary actions into longer sequences. Traditional research has explored whether these elements are combined in low-level control circuits for rapid execution or independently recalled from higher-level circuits. Surprisingly, the role of feedback control in the production of movement sequences has remained unexplored. Here, using simulations of optimal feedback control alongside human experiments, we shed light on the crucial role of task-dependent feedback control in shaping how sequences manifest within the motor system. Our findings reveal that the combination of sequence elements can result from flexible processing of multiple goals within the motor system involved in fast feedback control.

Introduction

Everyday behavior involves performing sequences of actions to achieve a given goal. Sequential motor skills depend on the ability to combine individual actions with short-term goals into a longer sequence consisting of multiple goals (1). However, the computations that underlie the combination of sequence elements and the neural circuits where this combination occurs remain a topic of debate (2–4).

Behavioral data from various sensorimotor repertoires indicate that the trajectory of sub-movements in a sequence is affected by multiple future goals (5–10). For example, in human speech and bird song, the pronunciation of individual elements is influenced by subsequent elements (11, 12), a phenomenon known as coarticulation. Ramkumar and colleagues (13) employed a computational model, revealing that concatenating multiple sub-movements into larger units, referred to as chunks (14, 15), induces coarticulation within the velocity patterns of sub-movements. This coarticulation serves as a natural solution to enhance the efficiency of chunked movements. Collectively, behavioral findings highlight a pattern where sub-movements depend on the whole sequence, or a chunk.

It remains unclear how future goals are integrated in the neural circuits to produce coarticulation. For rapid execution of a sequence, one possible solution is to represent multiple goals within low-level control circuits (3, 16), so that several elements can be executed together as a chunk. However, recent neural recordings in the primary motor cortex (M1), a region involved in the production of motor commands, showed no specific influence of future goals on the population responses governing the control of ongoing action (17, 18). Specifically, Zimnik and Churchland (18) observed in a two-reach sequence task that, there was no coarticulation in sub-movement kinematics although the execution got faster with practice. Notably, M1 displayed separate phases of execution related activity for each sub-movement. Using a neural network model, they interpreted that sequence goals could be separated and serially specified to the controller from regions upstream of M1 (Figure 1A). These findings diverge from

earlier studies demonstrating coarticulation of sub-movements and whole sequence representations in

M1 (19–21). Hence, coarticulation and separation of sub-movements have been formulated as distinct

hypothesis, with possibly dedicated neural processes, where the former presumably involves the

representation and production of a sequence as a whole, while the latter involves independent control of

each sub-movement (4, 18).

Strikingly, most studies have put much emphasis on potential differences in the state of the

sensorimotor system during preparatory activity (18–22). As a consequence, the role of feedback control

in this debate has been unexplored, although it is known that human motor control involves continuous

processing of sensory feedback (23, 24). The theory of optimal feedback control (OFC) has been

particularly useful in predicting the influence of numerous task parameters on the controller (25–32), thus

reproducing goal-directed motor commands during both unperturbed movements and feedback

responses to disturbances (28). In this framework, all targets in a sequence can be presented as

simultaneous inputs to the controller, and the overlap between control gains applied to each target

determines when they influence the sequence (Figure 1B). The control gains themselves depend on task

requirements (33–35). Thus, OFC predicts two experimentally testable features of sequence production:

first, even if the sequence is planned as a whole, control gains can show different amounts of overlap

based on task requirements, leading to either coarticulation or separation of the sequence elements;

second, in sequences with strong co-articulation, feedback responses to perturbations are expected to

depend on future targets due to the overlap of control gains.

Here we studied these predictions of OFC in a two-reach sequence task. Simulations revealed that

the instruction to slow-down at the first target influenced the coarticulation of the two reaches.

Importantly, the penalty for velocity at the first target produced an apparent separation of two reaches,

even if the controller was always computed jointly for both reaches. We conducted two experiments with

human participants and observed that their behaviour was fully consistent with the simulations. We

observed both flexible coarticulation dependent on task instruction about the intermediate target, as well

as a modulation of long-latency feedback responses to the mechanical loads applied during the first reach

dependent on location of the second target when coarticulation was present. These results demonstrate

that both coarticulation and separation of sub-movements may arise as the consequence of efficient

feedback control. Recordings of muscle activity suggest that this operation is carried out by the

sensorimotor network that mediates fast state-feedback control.

Results

The results section is organized as follows: we first introduce the computational model and

describe the results of the simulations showing that separate or coarticulated sequence production can

result from flexible feedback control without implying a different structure in the controller (Figure 1A-B).

Second, we present the two behavioural experiments in which participants were instructed to perform a

sequence of two reaching movements with an intermediate goal (Figure 1C-F). Our key manipulation

guided by the model simulations was to give them different instructions regarding the velocity at the first

target, which produced the modulation of sequence production predicted in theory.

Figure 1 Schematic illustration of hypotheses and experiment method. (A) independent control hypothesis
where each target is specified one at a time onto the controller (B) flexible feedback control hypothesis
where multiple targets are simultaneously considered (C) Experimental setup of two-reach sequences with

Simulations of sequence execution

The model considers a task dependent state-feedback controller derived in the framework of stochastic

optimal control (28). The controller was optimized to produce translation of a point mass on a horizontal

plane. A given estimate of a state vector $\hat{x}$ (position, velocity, target location and external load) can

influence the motor output u through multiplicative functions called 'feedback control gains', represented

by $k(t)$. Typically, the motor output at any time (t) is a result of a state feedback control law:

$$u(t) = k(t)\hat{x}(t).$$

Note that the control law is dependent on the state estimate $(\hat{x})$, which is computed by integrating

delayed and noisy sensory feedback (x) with internal prediction of the state. Importantly, feedback gains

are tuned to task demands (33, 36), and vary within a movement (34). This task dependency gives rise to

instances where all state variables $(\hat{x})$ are simultaneously available to the feedback controller, but some

of them may not influence the motor output if deviations of the state vector in the corresponding

dimension do not impact the performance.

We applied this control formalism to a task composed of a two-target sequence of reaching

movements, such that two different sequences shared the first target, after which they diverged towards

the second target (T1 and T2, Figure 1C). In the model, the estimated positions of the targets $(\widehat{T1}, \widehat{T2})$ are

in the state estimate $(\hat{x})$, hence the control law can be decomposed and expressed in the following

equation (also see Figure 2A):

$$u(t) = k_1(t)\widehat{T1} + k_2(t)\widehat{T2} + k_x(t)\hat{x}(t),$$

where (k_1, k_2) are the feedback gains that determine how much and when each target $(\widehat{T1}, \widehat{T2})$ influences

the motor output (Figure 2A). This formulation highlights that, even when both targets are simultaneously

available to the feedback controller, the overlap between control gains (k_1 and k_2) will determine if and

when the second target influences the first reach (Figure 2B). In the simulations, the control gains and the

overlap between the two components of the sequence were not hardcoded but were found as an optimal

control solution that improves the efficiency of the sequence as a whole (i.e., as a solution to holistic

sequence planning in the sense that all targets are specified to the controller at once). For holistic

sequence planning, we formulated the efficiency in terms of a cost function on the terminal movement

errors and speed of reaching at both targets (T1 and T2), and total motor output across movement

duration (see model with terminal costs in Methods).

Figure 2 Simulations of the OFC model in two-reaches sequence task. (A) Schematic illustration of the
feedback control policy. Targets influence motor output through control gains, which are themselves
derived by minimizing movement costs (B) Illustration of how overlap of control gains can explain
coarticulated reaches (top panel), and late influence of second target on first reach (middle). (C) Hand
speed (left), control gains corresponding to the two targets (middle), and hand path (middle) when dwell
velocity at target 1 was not constrained. (D) same as (C) except the dwell velocity was penalized.

The model simulations revealed that, even though the sequence was planned as a whole, the

necessity to slow-down at the first target, which was induced by an intermediate cost on speed at the

corresponding time, influenced when the first reach became sensitive to the second target (Figure 2C, D),

i.e., when the two reaches in the sequence coarticulate. When there was no constraint on speed at the

first target, the simulated hand reached the first target with a high velocity (Figure 2C, left panel), and

subsequently continued towards the second target at a high initial speed of > 0.2 m/s. In this rapid transfer

scenario, the control gains showed a cross-over effect where the control gain associated with the second

target (k2) was non-zero, albeit small, during the first reach, showing an early overlap with the feedback

gain (k1) (Figure 2C, middle panel). As a result, the location of the second target influenced the lateral

hand deviation of the first reach (Figure 2C, right panel). The first reach was thus different depending on

the location of the second target. Notably, the first reach was similarly influenced by the second target

even when temporal buildup costs were considered instead of terminal costs (see temporal buildup cost

model in Methods), to penalize the hand deviation from the targets across the duration of the sequence

(Supplementary Figure 1A).

Because the control gains also determine the response to external perturbations, the path

corrections following external loads applied to the mass were also dependent on the second target

(Supplementary Figure S2A). Interestingly, lateral deviation along the perturbed direction was relatively

smaller when the second target was in the direction of perturbation. Although this may appear intriguing,

the model shows that it is an efficient solution, because a higher acceleration linked to reversal at T1, when

the perturbation is in the direction of T2, incurs a higher motor cost. Hence, a smaller hand deviation is

necessary to avoid high acceleration at T1. This result qualitatively persisted across broad variations in

parameters related to the costs, delays and point-mass dynamics (not shown). A similar result was

observed when we used temporal buildup of kinematic error in the cost function (Supplementary Figure

2B), instead of using the simple terminal costs.

To the contrary, the requirement to slow down at the first target, captured by an increased cost

on speed, reduced the velocity at the end of first reach and altered the shape of the trajectories (Figure

2D, left panel) compared to the rapid transfer scenario (Figure 2C, left panel). The control gain

corresponding to the second target became non-zero at a later time, at the end of the first reach (Figure

2D, middle panel). Due to the later increase in this gain value, the influence of the second target on the

motor output during the first reach was much reduced, hence generating similar hand path to reach the

first target across the two sequences (Figure 2D, right panel). Similarly, feedback responses to lateral

perturbations that were applied early-on in the first reach were not dependent on the location of the

second target (Supplementary Figure S3A). The amount of lateral deviation when an unexpected

perturbation was applied did not vary as a function of the direction of the second target. Qualitatively

similar influence of the need to slow down at the first target on the trajectory of first reach was observed

when temporal buildup costs were included instead of terminal errors (Supplementary Figure S1B and

S3B).

Overall, the model predicted that even if all sequential movements were planned holistically over

a long time-horizon, the requirements associated with intermediate goals determines how early in the

sequence the second (future) target can influence the feedback controller (compare Figure 2C and Figure

2D; Supplementary Figure S1A and S1B; Supplementary Figure S2 and S3).

Correspondence between Human Behaviour and Model Predictions

The experiments were designed to verify whether task requirements at the intermediate goals

influenced hand trajectories and feedback control in a way that was expected from the model simulations.

The first experiments tested the scenario in which there was no penalty on velocity during the stop-over

at the first target, referred to as the GO task. We instructed 15 human participants to perform a two-target

sequence of reaching movements (Figure 1C-F). Both targets in a given sequence were displayed from the

beginning of the trial, allowing for preparation of both reaches. After receiving a GO cue (500-1000 ms

after target display), participants were required to reach the first target within a prescribed time (500 -

650 ms), and then to proceed to the second target. There was no instruction to stop at the first target. To

encourage fast transfer between the two reaches, the participants were allowed to remain in the first

target for a maximum of 150 ms after a successful reach, before initiating the second reach, otherwise the

second target disappeared. To determine if the feedback controller was tuned to the location of the

second target, mechanical loads (spanning 200 ms width) in randomized direction (rightward or leftward)

were applied randomly in 20% of the trials soon after the hand left the starting location (4.5 cm in forward

direction from the starting location (this occurred typically at ~200ms after the movement onset)

Figure 3 Hand kinematics in the GO task (experiment 1). (A) hand speed and (B) hand path in unperturbed
baseline trials. (C, E) same as (A) but for perturbation trials. (D, F) same as (B) but for perturbation trials.
(G) Baseline EMG response of PEC and PD muscles (H) Perturbation response (baseline subtracted) of PEC
and PD muscle during rightward and (I) leftward perturbations. Light curves in the hand path panels are
average paths of individual participants. Shaded area around the average curves represent the standard
error (±sem) across participants around the mean. Gray colored box indicates the average duration of the
first reach. Red and blue colors correspond to reaches made towards left and right second targets
respectively. t = 0 ms represents the time of perturbation onset. *p < 0.05, **p < 0.005, ***p < 0.001.

The model prediction about the influence of the second target during the first reach was clearly

borne out of the experimental results. Figure 3A illustrates the average hand speed across baseline trials

in the rapid transfer scenario. It is clear that the hand did not come to rest at the end of first reach, but

continued towards the second target at a relatively high transfer speed of ~0.2 m/sec. The hand speed at

the first target was slightly different between the two second target conditions (Figure 3A, inset; paired t-

test: t(14) = -3.8, p = 0.002). In line with model predictions, in the rapid transfer scenario, the location of

the second target influenced the hand path during the first reach (Figure 3B). Statistically, across trials and

participants, lateral hand deviation during the first reach was significantly different between the two

sequences (Figure 3B, inset; paired t-test: t(14) = -5.9, p < 10^{-4}).

Note that the experiments involve a block design where a sequence direction was maintained for

10 trials (including 2 randomly interleaved perturbation trials), while the blocks themselves were randomly

interleaved (Figure 1E). To test whether the influence of second target on hand deviation was due to some

trials occurring late within a block, we computed the hand path in the first and last trials and averaged

across all blocks for each participant. Group data in Supplementary Figure 4 illustrates that the hand

deviation was significantly different between the leftward and rightward sequence conditions from the

very first trial within a block (paired t-test: t(14) = 5.7, p < 10^{-3}). Similarly, the hand deviation was

significantly different between leftward and rightward sequences in the last trial within a block (paired t-

test: t(14) = 5.9, p < 10^{-3}). Additionally, a temporal effect was observed in the leftward sequence condition,

where the last trial incurred a higher hand deviation compared to the first trial within blocks (paired t-test:

t(14) = 3.9, p = 0.003). Note that the above results include correcting for multiple comparisons using Holm-

Bonferroni method. Overall, the block design of the experiment may lead to small differences between

first and last trials within a block, but the effect of second target on the lateral hand deviation was present

from very first trial within a block, suggesting that similar results can be expected in experimental designs

without block design.

During perturbation trials, hand trajectories during the first reach were influenced by the location

of the second target. In rightward perturbation trials, the dwell velocity and hand deviation (Figure 3C, D)

were significantly different across the two second target conditions (paired t-test: t(14) = -7.1, p < 10^{-5} for

dwell velocity; t(14) = -7.9, p < 10^{-5} for hand deviation). Similar differences across the two sequences were

observed in the leftward perturbation trials (Figure 3E, F) (paired t-test: t(14) = 6.9, p < 10^{-5} for dwell

velocity; t(14) = -8.2, p < 10^{-6} for hand deviation). The influence of second target on the lateral hand

deviation was qualitatively similar to that observed in model simulations, and counterintuitive to what we

might expect without the help of the model simulations. As observed in the model simulations (see also

Supplementary Figure 2), lateral hand deviation was smaller when the perturbation was in the direction

of the second target (T2) and vice-versa. This was consistent for both rightward and leftward perturbation

conditions. Both the model and humans expressed this strategy that can be seen as an emergent feature

of efficient feedback control during production of movement sequences.

We identified a pair of muscles in the upper-arm shoulder joint (Pectoralis Major, PEC, and

Posterior Deltoid, PD) that were strongly recruited by the lateral perturbation (see earlier studies with a

similar experimental setup (29, 37)). These muscles also steer the hand towards the first and second

targets in unperturbed condition. PEC muscle displayed agonist burst coinciding with acceleration of hand

towards the first target, while the PD muscle provided the antagonist burst to decelerate the hand near

the first target. Notably, both muscles displayed a clear difference in baseline EMG activity during the first

reach, depending on whether the following reach involved right or left second targets (Figure 3G).

Following the perturbation (left or right), the stretch responses were visibly larger in both PEC and PD

muscles compared to the baseline response (Figure 3H, I). The perturbation response in the stretched

muscle – PEC for rightward load (Figure 3H, left panel) and PD muscle for leftward load (Figure 3I, right

panel) – was visibly larger if the second reach involved movement reversal compared to the first reach

where the perturbation was in the direction of the second target. Interestingly, even the unloading

response (below baseline) in the unstretched (shortened) muscle – PD for rightward load (Figure 3H, right

panel) and PEC for leftward load (Figure 3I, left panel) – was larger if the second reach involved direction

reversal. Overall, the perturbation responses – both stretch and shortening muscle responses – indicated

vigorous feedback corrections against the load when the second target induced movement reversal at the

first target, explaining the smaller lateral deviation of the hand when the perturbation was in the direction

of the second target.

Sequence-dependent long-latency feedback responses

We assessed when the muscle stretch and shortening responses show dependency on the second

target. If the two targets were simultaneously used to form an efficient feedback controller for the whole

sequence, one would expect that the sensitivity of the response to the second target to become visible in

the long-latency epoch, considering responses in this epoch as a proxy of flexible, state-feedback controller

in the primate nervous system (28, 38). Overall, the EMG responses indicated a significant influence of the

second target on the long latency responses in both stretched and shortened muscles (Figure 4). When

perturbed in the rightward direction, PEC muscle (agonist) response increased in both sequence

conditions, which counters the perturbation by increasing the leftward pull. Given the lower hand

deviation in the rightward sequence (Figure 3D), we asked if the reduced deviation was due to a larger

leftward pull from the PEC muscle. Indeed the PEC stretch response was significantly larger in the

rightward sequence than the leftward sequence in the long-latency epoch, in both R2 (one tail paired t-

test: $t(14) = 3.5$, $p < 0.002$) and R3 (one tail paired t-test: $t(14) = 2.25$, $p = 0.02$). On the other hand, the PD

muscle shortening response was significantly larger (below baseline) in the rightward sequence than the

leftward sequence in both R2 (one tailed paired t-test: $t(14) = -3.2$, $p = 0.003$) and R3 epochs (one tailed

paired t-test: $t(14) = -3.5$, $p = 0.002$).

Figure 4 EMG perturbation responses in the GO task (Experiment 1). (A) PEC muscle response to a rightward
perturbation (B) PD muscle response to a rightward perturbation (C) PEC muscle response to a leftward
perturbation (D) PD muscle response to a leftward perturbation. Red and blue colors represent sequences
to leftward and rightward second targets respectively. Shaded area represents the standard error (±1 sem)
across participants around mean. All traces were aligned to perturbation onset. t = 0 ms indicates the time
of perturbation onset. *p < 0.05, **p < 0.005, ***p < 0.001 and n.s indicates non-significant effects of one-
tailed t-test with p > 0.05.

When the perturbation load was to the left, the PEC shortening response (below baseline) was significantly larger in the leftward sequence compared to the rightward sequence in both R2 (one tail t-test: $t(14) = 7$, $p < 10^{-5}$) and R3 epochs ($t(14) = 8.1$, $p < 10^{-6}$). Such a large unloading response in the PEC muscle could facilitate a larger correction in the leftward sequence by reducing the leftward pull in the direction of the applied load. Note that the PEC shortening response showed a weak significance in the R0 condition, when tested for absolute difference between the rightward and leftward sequence conditions, instead of testing if one was larger than the other (i.e., a two-tailed t-test in the R0 epoch: $t(14) = -2.3$, $p = 0.035$). Given this effect was transient and limited to a very short time-period, it could be a false positive. While the shortening response in PEC was larger, the stretch response in PD muscle was not significantly larger in the leftward sequence throughout the long-latency time period (one-tail paired t-test; R2 epoch:

t(14) = 2.8, p = 0.99; R3 epoch: t(14) = -0.13, p = 0.44). On the contrary, the PD stretch response was

significantly smaller in the leftward sequence in the R2 epoch (one-tailed t-test: t(14) = 2.8, p = 0.006). All

muscles exhibited differences in the voluntary time window (120 – 180 ms) during the first reach

depending on the location of the second target (two-tailed paired t-test: t (14) absolute value between

(3.5 – 7.4), and p-value between ($1e^{-6}$ – 0.012) across muscles and perturbation types).

Next, we verified that the differences reported above did not follow from mere differences in

baseline muscle activities at perturbation onset. Muscle responses exhibit a property called 'automatic

gain scaling' where the stretch response increases proportionally to the pre-perturbation muscle activity

(39). If gain-scaling was significant, then we expected to observe a difference in the R1 period response

between the two sequence conditions. We did not observe any sequence-related modulation of the

stretch responses in the R1 epoch, suggesting that any potential differences at baseline did not have a

strong impact. It is also important to realize that the sequence direction that involved a larger stretch

response in the corresponding muscle was always associated with lower levels of activity in the baseline

condition, thus potential biases induced by baseline differences were in the direction opposite to the effect

reported above.

Overall, the results indicate, when required to perform movement sequences rapidly, the

sensorimotor circuits that mediate long latency feedback responses showed modulation that was

dependent on the secondary target. The specificity of controller to the second target was evident when

perturbations were applied early-on during the first reach. These observations agree with the predictions

of the feedback control model that, with low or absent penalty on velocity at the first target, the second

target influences the hand path and feedback responses to perturbations prior to the first target.

Separation of Sequence Elements

Another clear prediction of the model is that when there is a cost on velocity at the first target,

the first reach should be less dependent on the second target. To test this prediction experimentally, we

performed a STOP task where 14 subjects were asked to perform the same two-target reaching

movements with a different speed constraint (Figure 1F). They were required to bring the hand to a near-

complete halt (speed < 0.5 cm/s) before starting the second reach.

Figure 5 illustrates that the hand speed at the first target dropped to near zero levels in all

unperturbed, leftward and rightward perturb conditions. Clearly, there was no influence of the second

target on the velocity of reaching the first target (two-tailed paired t-test: $t(13) = -1.9$, $p = 0.08$ in the

baseline condition; $t(13) = -0.01$, $p = 0.99$ for rightward perturbations; $t(13) = 2.1$, $p = 0.055$ for leftward

perturbations). The hand deviation during the first reach was similar between two sequences (two-tailed

paired t-test: $t(13) = -1.8$, $p = 0.08$ in the baseline condition; $t(13) = -1.2$, $p = 0.26$ for rightward

perturbations; $t(13) = -1.2$, $p = 0.25$ for leftward perturbations). Moreover, the unperturbed EMG

responses of the PEC and PD muscles were similar throughout the first reach, and diverged only after the

end of first reach, while moving towards second targets (T2). In rightward and leftward perturbation

conditions, the perturbation responses (*ΔEMG)* of the PEC and PD muscles (Figure 5D, E and Figure 6) were

not statistically different in pre-perturbation, R1, R2, or R3 epochs. Only the PEC perturbation response,

to a leftward perturbation, showed a weak effect of the second target location (R2 epoch: $t(13) = 2.1$, $p =$

0.055; R3 epoch: $t(13) = 2.44$, $p = 0.03$). To conclude, these results suggested that the second target did

not influence the first reach when the hand was required to slow-down at the first target.

Figure 5 Hand kinematics in the STOP task (experiment 2). (A) hand speed and (B) hand path in unperturbed baseline trials. (C, E) same as (A) but for perturbation trials. (D, F) same as (B) but for perturbed trials. (G) Baseline EMG response of PEC and PD muscles (H) Perturbation response (baseline subtracted) of PEC and PD muscle during rightward and (I) leftward perturbations. Light curves in hand path panels are average paths of individual participants. Shaded area around the average curves represent the standard error (±sem) across participants around the mean. Gray colored box indicates the average duration of the first reach. Red and blue colors correspond to reaches made towards left and right second targets respectively. t = 0 ms represents the time of perturbation onset. *p < 0.05, **p < 0.005, ***p < 0.001.

A straightforward interpretation could be that the stopping at the first target invoked a different strategy in which the control of the two reaches was performed independently, effectively separating the two movements, whereas executing them rapidly could produce the merging of the two sub-movements into a coarticulated sequence. While this is conceptually valid, it is not necessary and the model provides a more nuanced view: the apparent separation or coarticulation of the two motor patterns did not require any explicit formulation of whether they had to be combined into a sequence or not, rather it resulted

from the optimization of the controller, which in all cases considered the two targets as a whole and

derived an optimal control policy shaped by the instructions given about the intermediate goal.

Figure 6 EMG perturbation responses in the STOP task (Experiment 2). (A) PEC muscle response to a
rightward perturbation (B) PD muscle response to a rightward perturbation (C) PEC muscle response to a
leftward perturbation (D) PD muscle response to a leftward perturbation Red and blue colors represent
sequences to leftward and rightward second targets respectively. Shaded area represents the standard
error (±1 sem) across participants around mean. All traces were aligned to perturbation onset. t = 0 ms
indicates the time of perturbation onset. n.s indicates non-significant effects of one-tailed t-test with p >
0.05.

Discussion

The results presented here are consistent with recent behavioral studies showing that the

information about future targets in a sequential task is integrated at the stage of motor planning (6–8, 40,

41). Our key contribution is two-fold: we showed from computational and behavioral perspectives that

the coarticulation and separation of sequence components depended on task instructions, and likely

reflected efficient solutions of the same continuous control problem with distinct penalties at the

intermediate goals; second, the results highlighted that long-latency feedback control was modulated by

future targets in a coarticulated sequence, consistent with theories suggesting that movement planning is

synonymous with derivation of a control policy (42, 43).

A key question the model and experiments addressed was whether the integration of multiple

goals required processing in upstream voluntary circuits, indicating timed, voluntary recall of future goals.

Instead, our results support the idea of flexible and simultaneous processing of sequence elements, where

low-level controller plays a crucial role in combining sequence elements. The model based on optimal

feedback control showed that even if two reaches are planned as a whole, the second target does not

necessarily influence the controller. The constraints placed on swiftness of transition around the first

target modulated when the second target started to influence the sequence. This was clearly apparent in

the hand deviation during the first reach, and further revealed by perturbing participants' hand with

external loads, evoking sequence-dependent long-latency responses in Experiment 1. Hence, fast-

transcortical feedback control circuits were modulated early by the second target information in the

coarticulated sequence.

On the neural computations underlying coarticulation and chunking

Our study proposes a novel and precise definition of the neural difference to expect in order to

identify coarticulated versus independent activities from a computational point of view. Because each

condition (i.e., overlapping versus non-overlapping controllers as in Figure 2) was associated with different

cost-functions and time-varying control gains, it is the process of deriving these control gains, using the

internal representation of the task structure, that may differ across coarticulated and separated sequence

conditions. To our knowledge, how and where this operation is performed is unknown. Corollary

properties of this definition are that the preparatory activity (18, 44) may not discern independently

planned or coarticulated sequences because these situations imply different control policies (and cost

functions), as opposed to different initial states. Moreover, the nature of the sequence representation is

potentially not dissociable from its execution for the same reason.

Many studies highlight a phenomenon called chunking, where a subset of individual elements of

a sequence are concatenated and represented as a unified whole called chunks (13, 14, 22, 45). In the

pursuit of finding motor chunks, recent neural recordings explored for holistic (or) unified representations

in the circuits that produce motor commands, and found no such representations (4, 17, 18, 46). Hence it

was suggested that chunking resulted from a cognitive or high-level recall of intermediate goals sent to

the controller for independent execution (4) (see Figure 1A)(but, see (47)). On the contrary, our

developments also provide a computational definition of chunks. From our model, chunking (or a unified

representation) does not necessarily imply a single preparatory input to the controller representing

multiple spatial goals. Chunking may represent the formation of a composite cost function, hence a

composite control policy, that can jointly optimize multiple sub-movements (see Figure 1B). In this view,

chunking results from an acquired motor ability to overlap control policies for contiguous sequence

elements (Figure 1B and Figure 2). Importantly this property depends on the instruction at intermediate

goals, thus even a chunked sequence may falsely appear to be independently controlled (Figure 1A),

masking the signature of sequence representation in neural data even if the controller was formed at once.

Limitations

There are differences between the model and human movements that requires further study, such

as the influence of sequence direction on the dwell velocity, which was not observed in the model

simulations (compare velocity plots in Figure 2C and Figure 3A). A limitation is that nonlinear dynamics

were ignored in our model. However, the potential impact of non-linear mechanics on limb trajectory

could be completely suppressed by participants in Experiment 2, since we observed straight reach path,

thus making the curvature observed in Experiment 1 more likely related to the task constraints rather than

the biomechanics which did not differ between experiments. Yet, the quantification of how biomechanical

factors influence control may deserve a dedicated work, as it has been shown that biomechanics can also

influence coarticulation in a speech production task (48). Despite the limitations, the simple linearized OFC

models were crucial in determining the relationship between three properties of the task demands, its

impact on coarticulation, and feedback control. Our approach is consistent with how ideal actor/estimator

models have been predominantly employed as normative models in formalizing an initial hypothesis, and

simulating these to generate testable predictions (29, 33, 35–37, 49–55). Such normative models should

be distinguished from detailed biomechanical models that can trade-off interpretability and predictability

for achieving better quantitative fits. Furthermore, the main feature of the model – to generate a task-

dependent motor output as a combination of state feedback – is not limited to linear controllers and linear

body dynamics, but applicable even when the control policy is non-linear (see (56, 57)).

The modeling and empirical results presented here describe the computations underlying a two-

reaches sequence. Although two is admittedly the lowest possible length of a non-trivial sequence, it

enables comparison with similar tasks that have been used in recent neurophysiological studies (18–20)

to understand the neural computations underlying the control of sequences. We expect that longer

sequences be formed in a similar way as there is no theoretical constraint on the number of intermediate

goals. Observe that a recent study by Kashefi and colleagues (8), where longer sequences were used,

showed that at least two targets influence the hand path in a way that is similar to the coarticulation

condition presented in this study. Thus, we expect the combination of two elements can be the basis of

longer sequence formation constructed on the same principles.

Implications for sequential skill learning

Besides the implication of our computational model of co-articulation, it has been often

documented that swift execution of sequences was achieved by habituation, where a set of sequences are

practiced for several days to months (13, 17, 18, 21, 46, 47, 58, 59). Such process potentially involve a slow

time-scale linked to the acquisition of a motor skill (59, 60), which differs from the results presented here,

as the participants were able to execute accurate sequences after a pre-training of around 100 repetitions.

Here, the participants received explicit information about the dwell constraints at the intermediate goal,

resulting in quick coarticulation (also see (8, 10, 61–63)). This is in line with task-dependent modulation of

existing reaching controllers, which can be rapidly switched from one trial to another (29, 64). This suggests

that there exist several mechanisms operating at different time-scales and responsible for combination of

sequence elements. One mechanism is a slow habituation process in which the subject learns de novo to

build a novel controller by coarticulating different elements in the sequence (47, 65), whereas the other

mechanism is a fast modulation of this consolidated controller by task demands. In principle, the two

mechanisms are dissociable with perturbations, which may allow future studies to investigate how they

interact and to identify which brain regions mediate these components of skilled motor behaviour.

Acknowledgements

We thank our lab members and Kevin cross for their feedback on different drafts. HTK was supported by the Fonds de la Recherche Scientifique (F.R.S-FNRS) Chargé de recherche Grant CR 252 (FC 043127). F.C. was supported by the F.R.S.-FNRS Grant 1.C.033.18 (FC 036239). This work was additionally supported by a Concerted Research Action of Université Catholique de Louvain (ARC; "coAction").

Author contributions

HK and FC developed the models, designed the experiments and wrote the paper; HK conducted the experiments, and analyzed the data.

Declaration of Interests

The authors declare no competing interests.

Ethics Statement

Studies were approved by the ethics committee of UCLouvain.

Data and code availability

The data from simulations and experiments will be made available in the public repository 'figshare'.

Materials and Methods

Participants

A total of 29 volunteers (11 men, 18 women, age between 20 and 35 years) participated in 2 experiments (n = 15 with 9 women in Experiment 1; and n = 14 with 9 women in Experiment 2). All participants were neurologically healthy, right-handed and gave their informed consent according to a protocol approved by the ethics board at UCLouvain. Experiments did not exceed 2 h, and participants were compensated for their time.

Apparatus and Experiment Design

The experiments used a robotic device (endpoint KINARM, BKIN Technologies), permitting hand movement in the horizontal plane. In addition to recording the hand kinematics and forces on the robot handle, the KINARM can perturb the arm by applying mechanical loads at the handle. Projected targets (circles of chosen diameters) and hand feedback (white dot, 0.5 cm diameter) were presented to the participant from a heads-up display consisting of an overhead projector and a semitransparent mirror. During the experiment, direct vision of the arm and hand was occluded by a metal partition.

Participants were instructed to grasp a robotic handle and move the hand-aligned cursor to the home target (radius 0.6 cm). The home target was initially filled with red color but turned green when the participant's cursor entered the home target. Two targets (open circles with white edges) appeared simultaneously once the participants entered the home location. The first target was situated 18 cm in forward direction on the midline, while the second target was either situated to the left or right from the midline, displaced laterally from the first target by 8 cm (Figure 1C). Hence, at each trial, the participant could simultaneously view the first and second targets, T1 and T2 respectively. After holding the hand within the home location for 500 – 1000 ms ('preparatory period' uniformly distributed), the two targets

were filled with red color from open circles, acting as a GO cue, indicating the participant to start executing

the sequence. As the experiment involved many movements, in order to reduce the possibility that

participants missed a GO cue, a 50 ms beep sound was presented simultaneously as an additional, auditory

cue.

Experiment 1

Participants (n = 15) were instructed to initiate the sequence as soon as they receive the GO cues. For the

first reach towards T1, participants were asked to move as straight and as quickly as possible. The first

reach was deemed successful if T1 was reached between 500 – 650 ms after receiving the GO cue. Upon

successful first reach, the circle corresponding to the first target turned green. As soon as the first target

was successfully reached, a counter of 150 ms was initiated (unknown to the participant), leading to

disappearance of second target if the hand did not leave the first target quickly. The second target

remained in red color, if the hand left the first target within 150 ms, and subsequently turned green once

the hand reached T2. If participants successfully completed the sequence by acquiring T1 and T2, and

remained in T2 for at least 1 second, then they received 2 points. If they successfully acquired T1 but failed

to quickly leave T1 (< 150 ms after reaching) or to reach T2, then they received 1 point. The score was

projected on the screen and was updated only after the end of the trial and not during any reaching or

preparatory period. The short latency of dwelling (at most 150 ms) in the first target encouraged the

participants to swiftly move towards T2 soon after T1 was acquired. If the hand reached T1 in < 500 ms

after the GO signals, then both T1 and T2 disappeared indicating that movement was too fast, leading to

failure of this trial. If the hand reached T1 in > 650 ms after the GO signals, then the T1 circle never turned

red, indicating that the trial was too long. In both cases where T1 is not acquired properly, the participants

did not receive any points even if they reach T2. These cases of failed trials due to movements that were

too fast or too slow were included to encourage low variation in movement durations, but all trials were

used for analysis. Participants were asked to try to complete the sequence by reaching T2, even if they feel

that T1 was not reached within the prescribed time.

In a subset of random trials (20% of total number of trials), a mechanical load, of ±7 N in magnitude

with a 10 ms linear rise time and a 200 ms width, was applied randomly in the lateral direction (x axis in

Figure 1), once the hand reached 1/4th of the forward distance between the home location and the first

target (T1). Notably, in perturbation trials the successful reaching time to T1 was set to be between 500 –

800 ms, in contrast to 500 – 650 ms used in unperturbed trials. This additional time, not informed to the

participant, was necessary to account for the increased path length induced by the perturbation. This

ensured a good proportion of perturbation trials with complete sequence execution, instead of the

participants' not attempting the reach towards T2 due to failure of reaching T1 properly.

Experiment 2

Participants (n = 14) performed two-target reaching sequences similar to Experiment 1 except that they

were required to slow-down while reaching the first target (T1). Similar to Experiment 1, two targets were

simultaneously displayed (as open circles) once the hand enters home location. At the GO signals, only the

first target turned red, while T2 remained as an open circle. Participants were asked to reach T1 as straight

and quickly as possible, between 500 to 650 ms, but importantly to slow-down at T1 before continuing

towards T2. In this experiment, the condition for the target T2 to turn red, indicating the participant to

move towards it, was that the hand velocity had to fall below a threshold of 0.5 cm/s. The scores were

incremented in the same way as in Experiment 1.

Block design (Randomized within a set)

The experiments consisted of 9 sets, where each set composed of 10 blocks equally divided between

rightward and leftward sequences (i.e., 5 blocks with rightward and 5 blocks with leftward second target

location relative to T1) (see Figure 1). Notably, the rightward and leftward blocks within any given set

appeared in a pseudo-random order. Each block composed of 10 two-reach sequence trials. Hence, a total

of 900 trials were performed across 9 sets (9 sets × 10 blocks/set × 10 trials/block = 900 trials). After each

set, participants were given a 30-second to 1-minute break depending on their willingness to continue to

the next set. Across the 10 trials within a block, the location of the second target (T2) remained constant

(either to the right or to the left of T1). Within each block of 10 trials, 2 random trials involved

perturbations (1 rightward and 1 leftward perturbation), while the remaining 8 trials were unperturbed.

Hence, each perturbed direction included equal number of rightward and leftward sequences, allowing us

to examine the influence of location of the second target on the perturbation responses in each direction

of perturbation.

Pre-experiment training

The participants were first trained to execute single reaches to reach and stop at T1, without presenting

T2, for 50 practice trials. Then they performed a maximum of 1 set of two-reach sequences (10 blocks X

10 trials = 100 trials), without any perturbation trials. This allowed them to learn the task without the

complexities associated with reacting to external perturbations. Finally, they performed 1 more set,

including 10 blocks where each block is composed of 8 unperturbed trials and 2 perturbation (rightward

and leftward) trials. We added this pre-training before the main experiments to familiarize the participants

with the sequence task. We observed that the scores of most participants improved by the end of this

small pre-experiment training sets in both experiments.

Muscle recordings

We recorded the activity of mono-articular shoulder muscles using Delsys Bagnoli^{TM} surface EMG setup:

pectorialis major (PEC) and posterior deltoid (PD), involved in the production of lateral forces and known

to be largely recruited following similar perturbations with the same setup (66–68). The electrodes were

attached to the skin above the muscle belly after slight abrasion with alcohol. The signals were amplified

(gain: 10^4), digitally bandpass filtered with a dual-pass, fourth-order Butterworth filter (20 – 400 Hz

bandpass). EMG data were then normalized to the average activity across 1 s recorded when the

participants maintained postural control at the home target against a background load of 12 N applied

three times in each direction. This calibration was performed at the beginning of the 1st and 5th blocks.

Data Analysis

Behavior and Kinematics

We extracted the hand deviation, and dwell velocity at the first target, as defined below.

Lateral hand deviation: The average lateral hand deviation (in meters) from an imaginary straight-line

connecting the home location (H) to the first target location (T1), during the first reach. This is a signed

quantity where positive/negative sign indicates hand deviation to the right/left laterally.

Dwell velocity: The minimum root-mean-square velocity (of x and y dimension) in the duration in which

the hand dwells near the first target – below 0.5 cm distance from the edge of T1.

Kinematic variables of each participant were aligned to the time of perturbation onset (t = 0), then

averaged across trials for group data analysis. We performed paired t-tests (one-sampled and two-

sampled), unless explicitly stated otherwise, to determine significant differences in the extracted

kinematic variables.

Perturbation related muscle activity (ΔEMG)

Our primary interest was to compare different epochs of muscle activity to gain further insights into the

sensitivity of feedback responses to the location of the second target, when the hand was perturbed early-

on in the reaching towards the first target. The epochs of perturbation related muscle activity of most

interest were based on previous reports (32, 39, 69) and were categorized temporally: baseline = -100 to

0 ms, R0 = 0-20 ms, R1 = 20-45 ms, R2 = 45-75 ms, R3 = 75-105 ms, and voluntary (Vol) = 120-180 ms. Note

that all EMG signals for each participant were aligned to the perturbation onset.

We were most interested in comparing how the feedback responses are influenced by the location

of the second target, even though the perturbations were applied in the beginning of the first reach.

Accordingly, we calculated the average EMG response of each participant in the perturbed trials and

subtracted this with the average EMG responses across unperturbed trials for each block within a set. We

refer to this quantity as perturbation-related muscle activity (ΔEMG). Paired t-tests were performed using

group average to determine if the location of second target had any statistical effect on the perturbation-

related muscle activity. It should be noted that corrections for multiple comparisons do not apply here for

two reasons: first, the samples at each epoch are involved in only one comparison; second, consecutive

samples are not statistically independent. If there is a significant difference at a given epoch, it is very likely

that there will be a significant difference in the next epoch because signals do not vary instantaneously.

Hence, the risk of false-positive must not be controlled.

Model

The model describes the translation of a point-mass similar to the mass of an arm (m = 2.5 kg) in the

horizontal plane. The coordinates corresponded to the experiments, such that the first target was directly

in the forward reaching direction (along the y-dimension), and the second targets were at the same y-

coordinate but displaced in the x-dimension by 8 cm to the right or to the left of the first target.

<u>Dynamics of the effector</u>

The state-space representation of the point-mass (from (37, 50)), with mass 'm', was as follows:

596
$$m\ddot{x} = -g\dot{x} + f_x + f_{LOAD}$$

597 (1)

598
$$m\ddot{y} = -g\dot{y} + f_y$$

599 (2)

600
$$\tau\dot{f}_x = u_x - f_x$$

601 (3)

602
$$\tau\dot{f}_y = u_y - f_y$$

(4)

Dot(s) correspond to time derivatives. The variables f_x and f_y are the forces applied by the

controlled actuator to the mass, and these forces are a first order response to the actual control vector

denoted by variables u_x and u_y. The parameter g is a dissipative constant (0.1 Nsm^{-1}), Equations 3, 4

capture the first order muscle dynamics with time constant set to 0.1 s, and f_{LOAD} represents the external

load applied to perturb the point-mass (if any).

Defining the state vector as $\mathbf{z} = [x, y, \dot{x}, \dot{y}, f_x, f_y]$, and the control vector $\mathbf{u} = [u_x, u_y]$, we have $\dot{\mathbf{z}} = A\mathbf{z} +$

$B\mathbf{u}$, with:

$$A = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -g/m & 0 & 1/m & 0 \\ 0 & 0 & 0 & -g/m & 0 & 1/m \\ 0 & 0 & 0 & 0 & -1/\tau & 0 \\ 0 & 0 & 0 & 0 & 0 & -1/\tau \end{bmatrix}$$

612 (5)

613 and

614
$$B = \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 1/\tau & 0 \\ 0 & 1/\tau \end{bmatrix}$$

(6)

The system was discretized by using a first order Taylor series expansion over one time-step of δt : $A_d =$

$I + \delta t A$, and $B_d = \delta t B$ (I is the identity matrix). We used a discretization step of $\delta t = 0.01\ s$.

Importantly, the state space representation of body dynamics was augmented with the x and y-

coordinates of the two targets T1 and T2 (denoted by $x^*_{T1}, y^*_{T1}, x^*_{T2}, y^*_{T2}$), and the system was re-written in

discrete-time equations as follows:

621
$$z_{(t+1)} = A_d z_{(t)} + B_d u_{(t)} + \varepsilon_t$$

(7)

The subscript t is the time-step, and ε_t is a Gaussian disturbance with zero-mean and covariance $\sum_\varepsilon =$

$\sigma B B^T$, where $\sigma = 0.2$ is the scaling coefficient.

Control policy

The control policy (Π) produces motor commands (u_t) as a function of the estimated state of the point-

mass ($\hat{z}_t$). The state-estimate is computed using Kalman filtering as a weighted sum of internal model

prediction, and the sensory feedback representing the delayed state information ($z_{(t-\Delta)}$). The

sensorimotor delays were set at $\Delta = 50\ ms$ in the simulations. The details of the Kalman filtering can be

found in the previous paper (70). Overall, motor commands are computed by a linear control policy:

631
$$u_t = \Pi(\hat{z}_t) = K_t \hat{z}_t$$

(8)

The control gains (K_t) determine how much each state variable can influence the motor command,

and are computed to minimize a quadratic cost on the behavior (J), using standard dynamic programming

procedures (Riccatti equations) described in previous literature (50, 70). So, the control policy can be

formalized as finding the trajectory of the control gains 'K_t' that minimize the expected costs on the entire

sequence (i.e., a holistic plan):

$$\Pi^* = \underset{K_{t\epsilon[0,N]}}{\mathrm{argmin}}\, J\left(\hat{z}, u\right)$$

(9)

Importantly, the control policy from Equation 9, receives information about both targets within a sequence

as state feedback information, in addition to the continuous but delayed position, velocity and forces on

the end-effector.

<u>Cost function</u>

As described above, the controller design minimizes a quadratic cost function that captures the intended

behavior (in this case a two-reach sequence task). The cost function (J) was defined as a function of the

state-estimate ($\hat{z}$), and the motor commands (u) used to control the point-mass. In a linear quadratic

regulator (LQG), the control problem is to find a control sequence $\boldsymbol{u}^*$ which minimizes the expected value

of the sum of J($\hat{z}$, u).

Model with terminal costs: Typically, in the case of single reaches, the cost function penalizes the distance

between the target and the actual position of the end-effector (terminal or trajectory errors), speed, and

motor costs. But an efficient reaching movement in the context of single reach does not ensure the

efficiency of the sequence as a whole. Hence, we consider a composite cost function, that penalizes

terminal errors between the end-effector and both targets, to optimize the sequence elements as a whole.

Additionally, we can encourage swift or slower transfer between sequence elements by applying a cost on

the velocity at the moment when the system approaches the first target (i.e., the intermediate goal). Let

the duration of the sequence (N) be divided into two elements, where the first target (T1) should be

reached at time-step 'N1', and the second target (T2) at time-step 'N2'. The holistic cost can be formulated

as follows:

659
$$J(z, u) = J_{T1_{error}} + J_{T2_{error}} + J_{T1_{velocity}} + J_{T2_{velocity}} + J_{motorcost}$$

660   *(10)*

661
$$J_{T1_{error}} = w_1 \left[\left(x_{(N1)} - x_{T1}^* \right)^2 + \left(y_{(N1)} - y_{T1}^* \right)^2 \right]$$

662   *(11)*

663
$$J_{T2_{error}} = w_1 \left[\left(x_{(N2)} - x_{T2}^* \right)^2 + \left(y_{(N2)} - y_{T2}^* \right)^2 \right]$$

664   *(12)*

665
$$J_{T1_{velocity}} = w_2 \left[\dot{x}_{(N1)}^2 + \dot{y}_{(N1)}^2 \right]$$

666   *(13)*

667
$$J_{T2_{velocity}} = w_3 \left[\dot{x}_{(N2)}^2 + \dot{y}_{(N2)}^2 \right]$$

668   *(14)*

669
$$J_{motorcost} = w_4 \sum_{t=1}^{N2} \|u_t\|^2$$

(15)

Equations 11 and 12 penalize the terminal cost on the positional error relative to T1 and T2 at

time-steps N1 and N2 respectively. An equal weight ($w_1 = 500$) was used for both terminal error terms.

Equations 13 and 14 penalize the squared velocity at each target, and the last Equation 15 penalizes high

motor output. A rapid transfer between two reaches in a sequence can be simulated by removing the

penalty on the velocity at the first target. Hence, for this case, $w_2 = 0.1$, whereas a slower sequence

involves slowing down significantly at the first target, which correspond to a parameter set to $w_2 = 1$ in

the simulations. In all cases, we set $w_4 = 10^{-4}$ as the coefficient on the motor costs term ($\|u_t\|^2$).

Qualitatively similar results can be obtained with a broad range of parameter values (w_1, w_2, w_3, w_4).

Model with temporal buildup costs: The terminal costs, described above, are sufficient to produce accurate

movements with low endpoint errors, while neglecting the trajectory errors before the endpoint. Another

possibility is to penalize deviations in the trajectory from the desired endpoint targets throughout the

movement duration. Notably, by using temporal buildup of costs, the endpoint errors incur higher cost,

while incurring lower but non-zero cost in the movement duration before the endpoints. Such formulation

has been recently applied to emulate human reaching control under uncertain environmental dynamics,

in the context of elementary reaching movements (see 44, 62). To implement the temporal buildup costs

model, the position and velocity terms in Equations $11 - 14$ were modified as follows:

$$J_{T1_{error}} = \sum_{t=1}^{N1} w_1 (t/N1)^{b1} \left[\left(x_{(t)} - x_{T1}^* \right)^2 + \left(y_{(t)} - y_{T1}^* \right)^2 \right]$$

(16)

$$J_{T2_{error}} = \sum_{t=N1}^{N2} w_1 (\frac{t - N1}{N2 - N1})^{b2} \left[\left(x_{(t)} - x_{T2}^* \right)^2 + \left(y_{(t)} - y_{T2}^* \right)^2 \right]$$

(17)

$$J_{T1_{velocity}} = \sum_{t=1}^{N1} w_2 (t/N1)^{b1} \left[\dot{x}_{(t)}^2 + \dot{y}_{(t)}^2 \right]$$

(18)

$$J_{T2_{velocity}} = \sum_{t=N1}^{N2} w_2 (\frac{t - N1}{N2 - N1})^{b2} \left[\dot{x}_{(t)}^2 + \dot{y}_{(t)}^2 \right]$$

(19)

696 Where the terms $(t/N1)^{b1}$ and $(\frac{t-N1}{N2-N1})^{b2}$ assign higher cost for the errors that occur when t = N1

697 and t = N2 respectively, otherwise allowing for non-zero but time-increasing costs. The parameters $b1 =$

698 20 and $b2 = 4$ determine the steepness of increase in the cost for a given state from starting time to the

699 terminal times $N1$ and $N2$ respectively. Intuitively, these temporal buildup costs can encourage relatively

700 straight point-to-point reaching as the deviations from the targets (T1 and T2) are compensated

701 throughout the movement duration.

References

703 1. M. Botvinick, D. C. Plaut, Doing Without Schema Hierarchies: A Recurrent Connectionist Approach
704  to Normal and Impaired Routine Sequential Action. *Psychological Review* **111**, 395–429 (2004).

705 2. J. Diedrichsen, K. Kornysheva, Motor skill learning between selection and execution. *Trends in*
706  *Cognitive Sciences* **19**, 227–233 (2015).

707 3. O. Hikosaka, *et al.*, Parallel neural networks for learning sequential procedures. *Trends in*
708  *Neurosciences* **22**, 464–471 (1999).

709 4. A. L. Wong, J. W. Krakauer, Why Are Sequence Representations in Primary Motor Cortex So
710  Elusive? *Neuron* **103**, 956–958 (2019).

711 5. G. Ariani, J. Diedrichsen, Sequence learning is driven by improvements in motor planning. *Journal*
712  *of Neurophysiology* **121**, 2088–2100 (2019).

713 6. J. S. Diamond, D. M. Wolpert, J. R. Flanagan, Rapid target foraging with reach or gaze: The hand
714  looks further ahead than the eye. *PLOS Computational Biology* **13**, e1005504 (2017).

715 7. J. M. Gottwald, *et al.*, Infants prospectively control reaching based on the difficulty of future
716  actions – To what extent can infants' multiple step actions be explained by Fitts' law? (2018)
717  https:/doi.org/10.31234/osf.io/7wek4 (April 28, 2023).

718 8. M. Kashefi, *et al.*, Interaction of multiple future movement plans in sequential reaching.
719  2023.05.24.542099 (2023).

720 9. H. R. Sheahan, D. W. Franklin, D. M. Wolpert, Motor Planning, Not Execution, Separates Motor
721  Memories. *Neuron* **92**, 773–779 (2016).

722 10. R. Sosnik, B. Hauptmann, A. Karni, T. Flash, When practice leads to co-articulation: the evolution of
723  geometrically defined movement primitives. *Exp Brain Res* **156**, 422–438 (2004).

724 11. C. A. Fowler, E. Saltzman, Coordination and coarticulation in speech production. *Lang Speech* **36 (**
725  **Pt 2–3)**, 171–195 (1993).

12. M. J. Wohlgemuth, S. J. Sober, M. S. Brainard, Linked Control of Syllable Sequence and Phonology in
Birdsong. *J. Neurosci.* **30**, 12936–12949 (2010).

13. P. Ramkumar, *et al.*, Chunking as the result of an efficiency computation trade-off. *Nat Commun* **7**,
12176 (2016).

14. W. B. Verwey, Concatenating familiar movement sequences: the versatile cognitive processor. *Acta
Psychologica* **106**, 69–95 (2001).

15. W. B. Verwey, E. L. Abrahamse, Distinct modes of executing movement sequences: Reacting,
associating, and chunking. *Acta Psychologica* **140**, 274–282 (2012).

16. N. Kadmon Harpaz, K. Hardcastle, B. P. Ölveczky, Learning-induced changes in the neural circuits
underlying motor sequence execution. *Current Opinion in Neurobiology* **76**, 102624 (2022).

17. A. Yokoi, J. Diedrichsen, Neural Organization of Hierarchical Motor Sequence Representations in
the Human Neocortex. *Neuron* **103**, 1178-1190.e7 (2019).

18. A. J. Zimnik, M. M. Churchland, Independent generation of sequence elements by motor cortex.
Nat Neurosci **24**, 412–424 (2021).

19. Y. Ben-Shaul, *et al.*, Neuronal Activity in Motor Cortical Areas Reflects the Sequential Context of
Movement. *Journal of Neurophysiology* **91**, 1748–1762 (2004).

20. N. G. Hatsopoulos, L. Paninski, J. P. Donoghue, Sequential movement representations based on
correlated neuronal activity. *Exp Brain Res* **149**, 478–486 (2003).

21. X. Lu, J. Ashe, Anticipatory Activity in Primary Motor Cortex Codes Memorized Movement
Sequences. *Neuron* **45**, 967–973 (2005).

22. J. Diedrichsen, K. Kornysheva, Motor skill learning between selection and execution. *Trends in
Cognitive Sciences* **19**, 227–233 (2015).

23. F. Crevecoeur, I. Kurtzer, Long-latency reflexes for inter-effector coordination reflect a continuous
state feedback controller. *Journal of Neurophysiology* **120**, 2466–2483 (2018).

24. B. Parrell, J. Houde, Modeling the Role of Sensory Feedback in Speech Motor Control and Learning.
J Speech Lang Hear Res **62**, 2963–2985 (2019).

25. J. Diedrichsen, R. Shadmehr, R. B. Ivry, The coordination of movement: optimal feedback control
and beyond. *Trends in Cognitive Sciences* **14**, 31–39 (2010).

26. T. Cluff, S. H. Scott, Rapid Feedback Responses Correlate with Reach Adaptation and Properties of
Novel Upper Limb Loads. *Journal of Neuroscience* **33**, 15903–15914 (2013).

27. T. Cluff, S. H. Scott, Apparent and Actual Trajectory Control Depend on the Behavioral Context in
Upper Limb Motor Tasks. *Journal of Neuroscience* **35**, 12465–12476 (2015).

28. H. T. Kalidindi, F. Crevecoeur, Human reaching control in dynamic environments. *Current Opinion in
Neurobiology* **83**, 102810 (2023).

29. A. D. Comite, F. Crevecoeur, P. Lefèvre, Continuous Tracking of Task Parameters Tunes Reaching Control Online. *eNeuro* **9** (2022).

30. E. Todorov, M. I. Jordan, Optimal Feedback Control as a Theory of Motor Coordination. *Nature Neuroscience* **5**, 1226–1235 (2002).

31. S. H. Scott, Optimal feedback control and the neural basis of volitional motor control. *Nature Reviews Neuroscience* **5**, 532–546 (2004).

32. I. L. Kurtzer, J. A. Pruszynski, S. H. Scott, Long-Latency Reflexes of the Human Arm Reflect an Internal Model of Limb Dynamics. *Current Biology* **18**, 449–453 (2008).

33. D. Liu, E. Todorov, Evidence for the Flexible Sensorimotor Strategies Predicted by Optimal Feedback Control. *J. Neurosci.* **27**, 9354–9368 (2007).

34. M. Dimitriou, D. M. Wolpert, D. W. Franklin, The Temporal Evolution of Feedback Gains Rapidly Update to Task Demands. *Journal of Neuroscience* **33**, 10898–10909 (2013).

35. J. Česonis, D. W. Franklin, Mixed-horizon optimal feedback control as a model of human movement (2021) https:/doi.org/10.48550/arXiv.2104.06275 (November 8, 2022).

36. J. Y. Nashed, F. Crevecoeur, S. H. Scott, Influence of the behavioral goal and environmental obstacles on rapid feedback responses. *Journal of Neurophysiology* **108**, 999–1009 (2012).

37. F. Crevecoeur, S. H. Scott, T. Cluff, Robust Control in Human Reaching Movements: A Model-Free Strategy to Compensate for Unpredictable Disturbances. *J. Neurosci.* **39**, 8135–8148 (2019).

38. S. H. Scott, A Functional Taxonomy of Bottom-Up Sensory Feedback Processing for Motor Actions. *Trends in Neurosciences* **39**, 512–526 (2016).

39. J. A. Pruszynski, I. Kurtzer, T. P. Lillicrap, S. H. Scott, Temporal Evolution of "Automatic Gain-Scaling." *Journal of Neurophysiology* **102**, 992–1003 (2009).

40. G. Ariani, N. Kordjazi, J. A. Pruszynski, J. Diedrichsen, The planning horizon for movement sequences. *eNeuro* (2021) https:/doi.org/10.1523/ENEURO.0085-21.2021 (March 29, 2021).

41. K. C. Engel, M. Flanders, J. F. Soechting, Anticipatory and sequential motor control in piano playing. *Exp Brain Res* **113**, 189–199 (1997).

42. F. Crevecoeur, T. Cluff, S. H. Scott, Computational Approaches for Goal-Directed Movement Planning and Execution (2014) https:/doi.org/10.7551/mitpress/9504.003.0051 (September 8, 2023).

43. A. L. Wong, A. M. Haith, J. W. Krakauer, Motor Planning. *Neuroscientist* **21**, 385–398 (2015).

44. K. C. Ames, S. I. Ryu, K. V. Shenoy, Simultaneous motor preparation and execution in a last-moment reach correction task. *Nat Commun* **10**, 2718 (2019).

45. K. Bera, A. Shukla, R. S. Bapi, Motor Chunking in Internally Guided Sequencing. *Brain Sci* **11**, 292 (2021).

46. E. Berlot, N. J. Popp, J. Diedrichsen, A critical re-evaluation of fMRI signatures of motor sequence
learning. *eLife* **9**, e55241 (2020).

47. D. M. Huberdeau, A. M. Haith, J. W. Krakauer, Continuous motor skills as flexible control policies: a
video game study. 2023.09.21.558913 (2023).

48. D. J. Ostry, P. L. Gribble, V. L. Gracco, Coarticulation of jaw movements in speech production: is
context sensitivity in speech kinematics centrally planned? *J. Neurosci.* **16**, 1570–1579 (1996).

49. J. Diedrichsen, Optimal Task-Dependent Changes of Bimanual Feedback Control and Adaptation.
Current Biology **17**, 1675–1679 (2007).

50. F. Crevecoeur, J.-L. Thonnard, P. Lefèvre, A Very Fast Time Scale of Human Motor Adaptation:
Within Movement Adjustments of Internal Representations during Reaching. *eNeuro* **7**,
ENEURO.0149-19.2019 (2020).

51. J. Izawa, T. Rane, O. Donchin, R. Shadmehr, Motor Adaptation as a Process of Reoptimization. *J
Neurosci* **28**, 2883–2891 (2008).

52. T. Ikegami, *et al.*, Hierarchical motor adaptations negotiate failures during force field learning. *PLOS
Computational Biology* **17**, e1008481 (2021).

53. A. D. Comite, P. Lefèvre, F. Crevecoeur, Continuous evaluation of cost-to-go for flexible reaching
control and online decisions. *PLOS Computational Biology* **19**, e1011493 (2023).

54. M. Mistry, E. Theodorou, S. Schaal, M. Kawato, Optimal control of reaching includes kinematic
constraints. *Journal of Neurophysiology* **110**, 1–11 (2013).

55. T. Takei, S. G. Lomber, D. J. Cook, S. H. Scott, Transient deactivation of dorsal premotor cortex or
parietal area 5 impairs feedback control of the limb in macaques. *Current Biology* **31**, 1476-1487.e5
(2021).

56. H. T. Kalidindi, *et al.*, Rotational dynamics in motor cortex are consistent with a feedback controller.
eLife **10**, e67256 (2021).

57. T. P. Lillicrap, S. H. Scott, Preference Distributions of Primary Motor Cortex Neurons Reflect Control
Solutions Optimized for Limb Biomechanics. *Neuron* **77**, 168–179 (2013).

58. G. Ariani, J. Diedrichsen, Sequence learning is driven by improvements in motor planning. *J
Neurophysiol* **121**, 2088–2100 (2019).

59. P. Maceira-Elvira, *et al.*, Dissecting motor skill acquisition: Spatial coordinates take precedence. *Sci.
Adv.* **8**, eabo3505 (2022).

60. P. Beukema, T. Verstynen, Predicting and binding: interacting algorithms supporting the
consolidation of sequential motor skills. *Current Opinion in Behavioral Sciences* **20**, 98–103 (2018).

61. J. Friedman, M. Korman, Observation of an expert model induces a skilled movement coordination
pattern in a single session of intermittent practice. *Sci Rep* **9**, 4609 (2019).

62. R. Sosnik, T. Flash, B. Hauptmann, A. Karni, The acquisition and implementation of the smoothness
maximization motion strategy is dependent on spatial accuracy demands. *Exp Brain Res* **176**, 311–
331 (2007).

63. S. Sporn, X. Chen, J. M. Galea, Reward-based invigoration of sequential reaching.
2020.06.15.152876 (2020).

64. J. Česonis, D. W. Franklin, Contextual cues are not unique for motor learning: Task-dependant
switching of feedback controllers. *PLOS Computational Biology* **18**, e1010192 (2022).

65. C. S. Yang, N. J. Cowan, A. M. Haith, De novo learning versus adaptation of continuous control in a
manual tracking task. *eLife* **10**, e62578 (2021).

66. F. Crevecoeur, J. Mathew, M. Bastin, P. Lefèvre, Feedback Adaptation to Unpredictable Force Fields
in 250 ms. *eNeuro* **7**, ENEURO.0400-19.2020 (2020).

67. J. Mathew, P. Lefevre, F. Crevecoeur, Rapid Changes in Movement Representations during Human
Reaching Could Be Preserved in Memory for at Least 850 ms. *eNeuro* **7**, ENEURO.0266-20.2020
(2020).

68. A. De Comite, F. Crevecoeur, P. Lefèvre, Continuous Tracking of Task Parameters Tunes Reaching
Control Online. *eNeuro* **9**, ENEURO.0055-22.2022 (2022).

69. J. A. Pruszynski, *et al.*, Primary motor cortex underlies multi-joint integration for fast feedback
control. *Nature* **478**, 387–390 (2011).

70. F. Crevecoeur, R. J. Sepulchre, J.-L. Thonnard, P. Lefèvre, Improving the state estimation for optimal
control of stochastic processes subject to multiplicative noise. *Automatica* **47**, 591–596 (2011).

Supplementary Figures

Supplementary Figure S1. Simulations of the OFC model in two-reaches sequence task using temporal buildup costs. (A) Hand speed (left), control gains corresponding to the two targets (middle), and hand path (middle) when dwell velocity at target 1 was not constrained. (B) same as (A) except the dwell velocity was penalized.

A   holsitic planning with **unconstrained** dwell velocity : terminal costs model

B   holsitic planning with **unconstrained** dwell velocity : temporal buildup costs model

Supplementary Figure S2. Simulated perturbation trials when the whole sequence was planned for with no constraints on dwell velocity at the first target (A) control gains corresponding to the position errors relative to T1 (left column), control gains corresponding to T2 (middle column), and the perturbed hand path when a sudden load (+4 N) was applied in the rightward direction (right column) simulated using terminal costs. (B) Same as (A) where simulations used buildup costs, leading to straighter hand path in the beginning of the movement. Each curve in the hand path represents a single trial.

A   holistic (joint) planning with **constrained** dwell velocity : terminal costs model

B   holistic (joint) planning with **constrained** dwell velocity : temporal buildup costs model

Supplementary Figure S3. Simulated perturbation trials when the whole sequence was planned for but with a constraint on dwell velocity at the first target (A) control gains corresponding to the position errors relative to T1 (left column), control gains corresponding to T2 (middle column), and the perturbed hand path when a sudden load (+4 N) was applied in the rightward direction (right column) simulated using terminal costs. (B) Same as (A) where simulations used buildup costs, leading to straighter hand path in the beginning of the movement. Each curve in the hand path represents a single trial.

Supplementary Figure S4. Comparison between first and last trials averaged across blocks of each participant in the GO task (Experiment 1). Red color represents leftward sequence condition and blue color represents rightward sequence condition. Error bars represent $\pm 1\ sem.$ *** p < 0.0005, ** p < 0.005, n.s for p > 0.05 after correcting for multiple comparisons using Holm-Bonferroni method.