1 **An RXLR effector targets ER-Golgi interface to induce ER stress and necrotic cell**
2 **death**
3

4 Jihyun Kim[1,2], Jesse Kaleku[3], Jongchan Woo[3], Hongshi Jin[3], Hui Jeong Kang[1,2], Minji
5 Kang[1,2], Haeun Kim[1,2], Seungmee Jung[3], Cecile Segonzac[1,2], Eunsook Park[3*], Doil Choi[1,2*]
6

[1] Horticultural Biotechnology, Department of Agriculture, Forestry and Bioresources, Plant
Genomics and Breeding Institute, College of Agriculture and Life Sciences, Seoul National
University, Seoul, 08826, Republic of Korea
[2] Plant Immunity Research Center, Seoul National University, Seoul, 08826, Republic of
Korea
[3]Department of Molecular Biology, College of Agricultural, Life Sciences and Natural
Resources, University of Wyoming, WY, 82071, USA

14
15
16
17 **Short title**: RXLR effector in ER stress-triggered necrotic cell death
18

Abstract

To achieve successful colonization, the pathogen secretes hundreds of effectors into host cells to manipulate the host's immune response. Despite numerous studies, the molecular mechanisms underlying effector-induced necrotic cell death remain elusive. In this study, we identified a novel virulent RXLR effector named Pc12 from *P. capsici.* Pc12 induces necrosis by triggering a distinct ER stress response through its interaction with Rab13-2. Unlike conventional hypersensitive response cell death associated with effector-triggered immunity, Pc12-induced cell death does not coincide with defense gene expression. Instead, it induces the aggregation of ER-resident proteins and confines secretory proteins within the ER. Pc12 interacts with Rab13-2, exhibiting a specific affinity for the active form of Rab13-2. Thus, the complex of Pc12 and Rab13-2 mimics the conformation of the inactive state of Rab13-2, subsequently recruiting the Rab-escort protein (REP). This process results in disruptions in vesicle formation within the ER-Golgi trafficking pathway. Furthermore, the substitution of a single amino acid of Rab13-2 structurally predicted to be crucial for the Pc12 interaction decreased the interaction with Pc12 while maintaining the interaction with REP1. These findings offer valuable insights into the ER stress-mediated cell death as well as a potential strategy for enhancing resistance against pathogens.

Keywords:

Phytophthora capsici, RXLR effector, Necrotic cell death, Rab Small GTPase, ER stress, Vesicle trafficking

Introduction

P. capsici is an oomycete pathogen causing blight disease in economically important crops, including Solanaceae, Cucurbitaceae, Fabaceae, and Malvaceae (Kamoun et al. 2015; Parada-Rojas et al. 2021; Quesada-Ocampo et al. 2023). *P. capsici* can inflict up to 100% crop damage in fields upon successful plant infection. This is attributed to its rapid proliferation under various environmental conditions, robust persistence in the soil over five years, and high genetic variability, leading to its classification as the fifth most destructive oomycete worldwide (Kamoun et al. 2015; Quesada-Ocampo et al. 2023). *Phytophthora* species secrete an arsenal of RXLR effector proteins to infect host plants. RXLR effectors feature a highly conserved Arg-X-Leu-Arg (RXLR) motif, which is essential for their transport into host cells (Whisson et al. 2007; Wawra et al. 2017). They play a significant role in facilitating the successful establishment of the pathogen by disrupting plant physiology and defense responses (Whisson et al. 2007; Fan et al. 2018; Boevink et al. 2020). Therefore, a comprehensive understanding of the role of RXLR effectors is important for devising strategies to manage plant diseases effectively.

The endoplasmic reticulum (ER) functions as a cellular factory responsible for protein synthesis, folding, and export to subcellular organelles. These sequential processes must be balanced to maintain cellular homeostasis under environmental conditions (Cao and Kaufman 2012; Angelos et al. 2017; Adams et al. 2019). However, when an imbalance occurs leading to the accumulation of misfolded or unfolded proteins, known as ER stress, eukaryotic cells initiate the unfolded protein response (UPR). The UPR involves the translocation of key transcription factors like bZIP28 and bZIP60 into the nucleus via maturation processes, such as proteolytic cleavage or appropriate splicing. These transcription factors regulate the expression of UPR target genes crucial for protein folding and ER-associated degradation (ERAD), enhancing protein folding capacity in cells (Cao and Kaufman 2012; Angelos et al. 2017; Adams et al. 2019). However, if plants are subjected to severe or chronic stress, it can further lead to unresolved ER stress-induced programmed cell death (PCD) as a survival mechanism (Yang et al. 2014; Liu and Howell, 2016; Yang et al. 2021; Simoni et al. 2022).

Rab is a superfamily of small GTPases that plays a crucial role in intracellular vesicle trafficking, an essential process for the transfer of proteins, lipids, and other cellular materials between different membrane compartments within eukaryotic cells (Nielsen et al. 2008;

Stenmark et al. 2009; Pylypenko et al. 2018; Gray et al. 2020). Within a species, Rab proteins

exist in multiple isoforms, each defining membrane identity and mediating vesicle trafficking

to distinct subcellular compartments. Rab proteins function in the formation and guide of

vesicles by acting as molecular switches, cycling between an active GTP-bound form and an

inactive GDP-bound form (Nielsen et al. 2008; Stenmark et al. 2009; Pylypenko et al. 2018;

Gray et al. 2020). Dysfunctional Rab proteins in vesicle trafficking can result in ER stress

due to improper protein compartmentation (Kiral et al., 2018). Several effectors of various

pathogens are known to suppress antimicrobial protein secretion (PR1 and PDF1.2) or

subvert vesicle movement toward the pathogen's focal area by targeting Rab proteins

(Tomczynska et al. 2018; Pandey et al. 2021; Li et al. 2022). However, it remains elusive how

the RXLR effector leads to ER stress-induced cell death by disrupting Rab proteins in vesicle

trafficking.

In this study, we report a virulent RXLR effector Pc12 from *P. capsici*, triggering necrotic

cell death in plants of the Solanaceae. Transient overexpression and host-induced gene

silencing (HIGS) of *Pc12* in plants significantly affected the pathogenicity of *P. capsici*. Pc12

induced ER stress, resulting in distinct expression patterns of UPR genes and altering the

location of ER-resident proteins and the secretory pathway. Notably, in the presence of Pc12,

an active Rab13-2 attracted the Rab-escort protein 1 (REP1), enhancing its interaction. We

have also identified a crucial residue in Rab13-2 to avoid being targeted by Pc12 while

maintaining its original function to interact with REP1 and PRA1. Consequently, this study

provides valuable insights suggesting that modifying Rab13-2 to evade effector targeting

could offer a promising strategy for combating the pathogen infection while preserving its

essential functions.

Results

Pc12 induces cell death independently of the plant defense mechanism in Solanaceae

plants.

Among the previously selected RXLR effector candidates (Seo et al. 2023), Pc12 exhibited a

strong induction of cell death in *N. benthamiana*. Interestingly, Pc12 consistently induces cell

death in other Solanaceae species when transiently expressed in *N. tabacum, C. annuum,* and

S. lycopersicum (Figure 1A), suggesting that Pc12-induced cell death is more pronounced

within the Solanaceae family. To determine whether the observed cell death induced by Pc12

resulted from intracellular receptors of nucleotide-binding site leucine-rich repeats (NLRs) -

mediated hypersensitive response (HR), we investigated Pc12-mediated cell death in *N.*

benthamiana plants with individual silencing of NLR chaperon (*SGT1*) or NLR-downstream

signaling factors, including *EDS1, ADR1/NRG1,* and *NRC234* (Botër et al. 2007; Dongus and

Parker, 2021; Wu et al. 2017). Interestingly, Pc12-induced cell death remained unaffected by

the silencing of NLR chaperon or NLR-downstream signaling components (Figure 1B-D),

suggesting that Pc12 induces cell death through mechanisms that are distinct from those

typically associated with NLR-mediated effector-triggered immunity (ETI). To further

validate whether the cell death induced by Pc12 is distinct from HR cell death, we compared

the expression of defense-related genes during Pc12 expression in contrast to the expression

of a bacterial effector derived from *Xanthomonas spp.,* XopQ. XopQ is known to interact

with the resistance protein Roq1 in *N. benthamiana*, resulting in a mild chlorotic phenotype

(Schultink et al. 2017). Transcription of defense-related genes, including *PR1, RbohB,* and

WRKY8, exhibited significant upregulation upon transient overexpression of XopQ. In

contrast, Pc12 expression did not significantly induce the expression of defense-related genes

(Figure 1E), supporting the idea that Pc12-induced cell death may activate a different

molecular pathway compared to the ETI-induced HR cell death.

Pc12 functions as a virulence effector, enhancing the pathogenicity of *P. capsici*.

Pc12 expression is elevated during the biotrophic phase of *P. capsici* (Supplemental Figure 1).

Although Pc12 displays the canonical structure of an RXLR effector, it remains uncertain

whether Pc12 is a virulent effector for *P. capsici* colonization. To assess the impact of Pc12

on the virulence of *P. capsici*, the pathogen was inoculated in plants expressing Pc12 under

the control of an ethanol-inducible promoter without ethanol treatment (Figure 2A). Leaky

expression of Pc12 driven by the ethanol-inducible promoter (Lee et al. 2018) allows us to

generate weak expression of Pc12 avoiding rapid cell death. The *Agrobacterium* carrying

construct of RFP or RFP-Pc12 was infiltrated into *N. benthamiana*, followed by drop

inoculation of *P. capsici* onto the leaves 1 day post infiltration. We observed a substantial

increase in lesion size (Figure 2A) and *P. capsici* biomass (Figure 2B). Host-induced gene

silencing (HIGS) is a method that initiates targeted gene suppression in the pathogen through

the host's RNA interference (RNAi) machinery (Cheng et al. 2022). In all cases, the silencing of Pc12 by HIGS using TRV resulted in reduction in the pathogenicity of *P. capsici* compared to the control (Figure 2C-D). This suggests that Pc12 plays a role in enhancing the pathogenicity of *P. capsici* acting as a virulence effector.

Pc12 lineage expanded specifically in *P. capsici* and the C-terminal residue is crucial for inducing cell death.

To elucidate whether Pc12 functions as a single gene or as part of a gene family, we examined the copy number variation and sequence polymorphism of Pc12 homologs in *Phytophthora spp.* The sequence polymorphism of Pc12 was assessed through a BLAST search, encompassing genome sequences from *P. capsici* (ten strains; LT1534, KPC-7, MY-1, JHAI1-7, and CPV-219/262/267/270/277/302) and four other *Phytophthora spp.* (*P. ramorum* Pr102, *P. sojae* P6497, *P. cinnamoni* CBS144.22, and *P. infestans* T30-4) retrieved from the public database. A comparative genomic analysis revealed that the *P. capsici* strains exhibited a significantly higher number of Pc12 homologs in their genome compared to other *Phytophthora* pathogens (Supplemental Table 1, Reyes-Tena et al. 2019; NCBI reference sequence database (RefSeq)). In order to ascertain whether the Pc12 homologs of these *Phytophthora spp.* could induce cell death, three homologs with high similarity to Pc12, except for 90% identity with Pc12, from *P. capsici* LT1534 and one homolog each from *P. ramorum, P. sojae,* and *P. cinnamomi* were synthesized and transiently expressed in *N. benthamiana*. Interestingly, the cell death phenotype was only observed with *P. capsici* homologs, with the exception of PHYCA22034 (Supplemental Figure 2). Although Pc12 and PHYCA22034 share 85% identity, substantial differences are found at the C-terminal end, which may be responsible for their distinct abilities to induce cell death (green line in Figure 2E). To determine the importance of the C-terminus of Pc12 in triggering cell death, we generated a series of chimeric mutants with modified amino acids at the C-terminus of PHYCA22034 and a deletion mutant lacking five amino acids of Pc12 (Pc12ΔC5) (Figure 2F). The substitution of amino acids from the 91st to 95th position of Pc12 in PHYCA22034 resulted in mild cell death, whereas the substitution of amino acids from the 89th to the 95th position fully restored cell death in *N. benthamiana* (Figure 2G). Consistently, Pc12ΔC5 reduced the induction of cell death (Figure 2G). These results suggest that the functional

Pc12 lineage has expanded specifically within *P. capsici*. Furthermore, the C-terminal

sequence plays a pivotal role in triggering cell death in plants.

Pc12 interaction with Rab13-2 small GTPase via C-terminal five amino acids is essential

for cell death.

To identify Pc12 interacting proteins that would be responsible to induce necrotic cell death,

we performed immunoprecipitation and mass spectrometry (IP-MS) using Pc12 and

Pc12ΔC5. The list of proteins exclusively interacting with Pc12 but not with Pc12ΔC5

includes a small GTPase Rab13-2 protein in the highest peptide-spectrum match (PSM) score

(Supplemental Table 2). We confirmed *in planta* interaction between Pc12 and Rab13-2 by a

co-immunoprecipitation (co-IP) assay (Figure 3A). In addition, homologs of Pc12 that trigger

cell death also presented interaction with Rab13-2 (Supplemental Figure 2 and Figure 3A).

This suggests that the interaction between Pc12 homologs, which share a similar C-terminus

with Rab13-2, is crucial for the induction of cell death. Rab small GTPase is a superfamily

protein functionally conserved in eukaryotic cells (Nielsen et al. 2008, Pylypenko et al. 2018,

Nielsen et al. 2020). Typically, the family comprises more than 60 isoforms that reside in

specific membrane compartments generated during membrane trafficking in eukaryotic cells

(Stenmark 2009, Pylypenko et al. 2018, Nielsen et al. 2020). In the *N. benthamiana* genome,

the Rab protein family consists of 149 genes (Supplemental Figure 3). To explore the

specificity of the interaction between Pc12 and Rab13-2, co-immunoprecipitation

experiments were performed with Rab13-2 homologs. Pc12 was found to bind selectively to

Rab13-3 and Rab13-4, which share 98% and 91% amino acid sequence similarity with

Rab13-2, respectively (Figure 3B). This suggests that Pc12 preferentially interacts with

specific Rab13-2 homologs. Interestingly, one of the Pc12 homologs demonstrated to induce

cell death through its transient expression in *N. benthamiana* previously (Li et al. 2019) also

interacts with one of the Rab 13 proteins (Rab 13-4), which localizes with the ER and in the

nucleus (Li et al. 2022). GFP-Rab13-2 localizes close to the ER (Figure 3C) and colocalizes

with the Golgi apparatus (Figure 3D), suggesting a role for Rab13-2 at the ER-Golgi interface.

Rab proteins contain two loops in their globular region that undergo allosteric conformational

changes depending on whether the Rab protein is bound to GTP or GDP. In the GTP-bound

state, these loops move closer together, resulting in a compact structure. In contrast, Rab

proteins bound to GDP have a more relaxed structure (Gray et al. 2020). To determine the specific state of Rab13-2 targeted by Pc12, we generated mutants representing the constitutively active (Rab 13-2^{Q74L}) or inactive (Rab13-2^{S29N}) form of Rab13-2 based on the consensus sequences (Langemeyer et al., 2014; Bourne et al., 1991). In vitro GTPase assays confirmed the predicted enzyme activities of these mutants (Figure 4A). Co-IP assays showed that Pc12 interacts exclusively with the active form of Rab13-2 (Figure 4B). The direct interaction of Pc12 and Rab13-2 was further confirmed by a yeast two-hybrid (Y2H) assay (Figure 4C). In support, no interaction is observed between Rab 13-2 and Pc12ΔC5, highlighting the importance of the C-terminus of Pc12 in facilitating the interaction between Pc12 and Rab13-2.

The complex of Pc12 and Rab13-2 mimics the inactive conformation of Rab13-2.

Rab proteins are involved in vesicle budding, trafficking, and fusion processes by interacting with or recruiting other proteins (Stenmark 2009; Pylypenko et al. 2018; Nielsen et al. 2020). In the case of Rab13-2, potential interactors were identified using the STRING database (Supplemental Table 3). Among the interactors, the Rab escort protein 1 (REP1) and the prenylated Rab acceptor 1 (PRA1) were found to interact with Rab13-2 by co-IP screening (Supplemental Figure 4). REP1 has a dual role: it acts as a chaperone to facilitate the prenylation process of Rab and as a carrier to deliver Rab to its target membrane (Guo et al. 2008). PRA1 acts as a receptor for prenylated Rab proteins, promoting their binding to membranes (Figueroa et al. 2001). We hypothesized that Pc12 might affect the interaction between Rab13-2 and its interactors, REP1 or PRA1. To confirm the effect of Pc12 on these interactions, we performed an immunoprecipitation experiment. Rab13-2 and its interactors were continuously expressed in a plant, while Pc12 was induced by ethanol treatment at 1 dpi (day post-infiltration) to ensure that the action of Pc12 did not affect the accumulation of newly synthesized proteins. The results showed that Pc12 did not form a complex with Rab13-2 in the presence of PRA1 (Figure 4D), suggesting that Pc12 does not interact with Rab13-2 associated with PRA1. On the other hand, Pc12 interacts with Rab13-2 associated with REP1 (Figure 4D). Furthermore, the interaction between Rab13-2 and REP1 was enhanced by the expression of Pc12 (blue asterisk, α-FLAG/α-GFP = 1.3, Figure 4D). This suggests that Pc12 plays a regulatory role in the association between Rab13-2 and REP,

potentially affecting their cellular functions and signaling pathways. Pc12 displays a

preference for binding to the active form of Rab13-2 (Rab 13-2^{Q74L}, Figure 4B, C). However,

REP1 has been shown to interact predominantly with the inactive form (Stenmark 2009,

Supplemental Figure 5). Therefore, we hypothesized that binding of Pc12 to the active form

of Rab13-2 might alter the structural conformation similar to the inactive state of Rab13-2,

thereby facilitating the recruitment of REP1 to active Rab13-2. To validate this hypothesis,

we performed a co-IP with Pc12, Rab13-2^{Q74L}, Rab13-2^{S29N}, and REP1. Remarkably, REP1

showed a significant association with Rab13-2^{Q74L} in the presence of Pc12, compared to the

control RFP (Figure 4E). This experiment was repeated eight times, all with consistent results

(Figure 4F, Supplemental Figure 6). This finding indicates that Pc12 binds to the active state

of Rab13-2, thereby facilitating the recruitment of REP1. This recruitment may alter the

dynamics of Rab13-2 GTPase activity, ultimately leading to the disruption of vesicle

trafficking.

Pc12 interaction with Rab13-2 alters ER-Golgi trafficking

Rab proteins are known for their distinct localization within the plant endomembrane network

(Stenmark 2009; Nielsen et al. 2020). The hypervariable C-terminal region of Rab proteins

contributes to the specificity of membrane delivery for certain Rab proteins (Figueroa et al.

2001; Pylypenko et al. 2018). Previously, a Pc12 isoform was identified that interacts with a

Rab 13 subfamily isoform localizes to the ER (Li et al. 2022). To gain insight into the

localization of Pc12 within host cells, we transiently expressed EGFP-Pc12 and EGFP-

Pc12ΔC5 (without the signal peptide and RXLR-EER motifs) under the control of an ethanol-

inducible promoter. Due to the rapid induction of cell death by Pc12, cytosolic Pc12

expression was observed in the early time point following 1% ethanol spray treatment to

determine its subcellular localization (Figure 5A). Interestingly, in contrast to the previously

studied isoform (Li et al. 2022), the Pc12 proteins are defused in the cytoplasm with

preferential accumulation closer to the ER and Golgi (Figure 5A). We attempted to confirm

the Pc12 localization in the vicinity of the ER and Golgi apparatus by co-expressing ER

marker (Chakrabarty et al. 2007) and Golgi marker (Nelson et al. 2007, Park et al. 2017).

Surprisingly, we observed that cells showed a peculiar punctate localization of RFP-HDEL in

the cell (Figure 5B), whereas the reticular morphology of the ER visualized by RFP-HDEL is

unchanged under the expression of EGFP or EGFP-Pc12ΔC5 (Figure 5B). In addition, the cis-Golgi marker, soybean α-1,2 mannosidase I fused to mCherry (Nelson et al. 2007), was trapped in the ER under Pc12 expression (Figure 5C). The mislocalization of these two markers under the Pc12 expression suggests that perturbation of the Rab13-2 dynamics by Pc12 interaction strongly affects vesicle trafficking from the ER to the Golgi.

To better understand Rab13-2 function and Pc12 interference with vesicle trafficking at the ER-Golgi interface, we generated transgenic *N. benthamiana* plants expressing Rab13-2 together with the ER marker. As we observed with transient expression of GFP-Rab13-2 (Figure 3C, D), Rab13-2 proteins colocalized in the vicinity of ER and Golgi puncta structures (Figure 5D, top panel). Interestingly, under Pc12 expression, puncta structures shown by ER markers were located at ER tubules and often accumulated near Golgi (Figure 5D, yellow arrows). ER marker consist of mCherry, a red fluorescent protein fused to a topic signal peptide at the N-terminus and an ER retention signal peptide (HDEL) at the C-terminus (Park et al. 2017). RFP-HDEL typically displays a reticular ER morphology when observed under a microscope. This is because newly synthesized proteins containing HDEL are first synthesized within the ER lumen and then transported to the Golgi apparatus. They then return to the ER via signal peptide recognition (Alvim et al. 2023). The punctate accumulation of RFP-HDEL at the ER tubules and the proximity to the Golgi induced by Pc12 suggest that it may be caused by interference with the mCherry proteins secreted from the ER to the Golgi.

Rab small GTPases continuously switch enzyme properties between their GTP-bound and GDP-bound forms. Conventionally, the active state of Rab proteins was thought to be localized to membranes, while the inactive state was predominantly found in the cytoplasm, as dictated by their respective functions (Stenmark, 2009; Nielsen, 2020). To validate the localization of Rab13-2 mutants, we generated transgenic *N. benthamiana* expressing the constitutive active form Rab13-2^{Q74L} or the constitutive inactive form Rab13-2^{S29N} together with an ER marker (Figure 5E and F, respectively, supplemental Figure 7). Notably, the constitutive inactive form Rab13-2^{S29N} was frequently observed at the Golgi apparatus (Figure 5F, supplemental Figure 7C), but the constitutive active form Rab13-2^{Q74L} did not show the Golgi localization (Figure 5E, supplemental Figure 7B), contradicting the general expectation of Rab small GTPase dynamics of membrane binding upon their activation by

GTP recruitment (Stenmark, 2009; Nielsen, 2020). The localization pattern of Rab13-2^{Q74L} is

more similar to that of RabE1d^{Q74L} and Rab8^{Q74L} (Speth et al., 2009; Pandy et al., 2021).

Since Rab proteins are initially synthesized in a structurally stable active state, the host

interactors responsible for Rab prenylation and membrane anchoring are unable to bind to the

mutant due to its abnormal conformation.

With this in mind, we investigated whether the localization of Rab13-2 might be altered upon

coexpression of Pc12. We observed that larger punctate localizations of GFP-Rab 13-2^{Q74L}

were frequently observed upon expression of Pc12 (white arrows, Figure 5E), whereas they

were not observed with RFP or Pc12ΔC5. Furthermore, a more diffuse localization of GFP-

Rab13-2^{S29N} in the vicinity of the ER was observed under the Pc12 expression (yellow

arrowheads, Figure 5F). The results suggest that Pc12 alters the properties of Rab13-2^{Q74L} to

resemble those of Rab13-2^{S29N} and effectively recruiting REP1 and consequently localizing it

to the Golgi apparatus.

308 Pc12 interference with ER-Golgi vesicle trafficking primarily affects the secretory

309 pathway and induces ER stress leading to necrotic cell death.

The altered localization of Rab 13-2^{Q74L} with Pc12 indicates that the rapid dissociation of

active Rab proteins and their return to the ER periphery may be disrupted under Pc12

expression. Displacement of ER marker and Golgi marker under Pc12 expression (Figure 5B

and C) also suggested that the alteration of Rab13-2 dynamics might disrupt trafficking from

the ER to the Golgi. Therefore, we observed the localization of two subcellular compartment

markers (Figure 6A and B). Notably, an apoplast marker of moxVenus fluorescent proteins,

fused to an apoplastic signal peptide at the N-terminus to secrete the fluorescent protein

outside the plasma membrane (ApoSP-moxVenus, Balleza et al. 2017), is trapped in the ER

in the presence of RFP-Pc12 (Figure 6A). This observation suggests that Pc12 disrupts

vesicle trafficking at the ER-Golgi interface, leading to changes in the localization of the

markers.

Interestingly, the localization of a tonoplast marker, moxVenus fused to γ-TIP (Nelson et al.

2007), was not significantly affected approximately 18 h after induction of Pc12 expression

(Figure 6B, top panel), while the apoplast marker trafficking was completely blocked in the

ER (Figure 6A, Supplemental Figure 8A). At a later time point, 28 hours after Pc12 induction,

dead cells are often observed under the microscope, and the tonoplast markers are observed

in the ER (Figure 6B, lower panel). Later, around 36 h after Pc12 induction, most of cells are

already completely dead, but tonoplast markers rarely maintain their integrity (Supplemental

Figure 8B). An alternative pathway of vesicle trafficking from the ER to the vacuole,

bypassing the Golgi, has been investigated as an important route for ER-vacuole trafficking

(Rojas-Pierce, 2013). This observation suggests that the primary function of Rab13-2 in the

secretory pathway is through the ER-Golgi pathway. Thus, perturbation of this pathway by

Pc12 alternation of Rab13-2 dynamics could induce a significant traffic jam of secretion at

the ER-Golgi interface, ultimately inducing ER stress-mediated necrotic cell death.

Interestingly, the mutualistic fungus *Piriformospora indica* has been shown to induce ER

stress-induced cell death when it colonizes *Arabidopsis* roots (Qiang et al. 2012).

Furthermore, the altered ER morphology induced by *P. indica* infection is similar to that

induced by Pc12 expression (Qiang et al. 2012, Figure 5B). Therefore, we compared the

responses induced by Pc12 and tunicamycin, a known inducer of ER stress. These responses

include the accumulation of ER chaperone-binding immunoglobulin protein (BiP) and the

transcription of unfolded protein response (UPR) genes (Cao and Kaufman 2012; Angelos et

al. 2017; Adams et al. 2019; Yang et al. 2014). Following application of 5% ethanol to the

leaves, ethanol-induced expression of RFP-Pc12 and RFP resulted in the accumulation of

both proteins over time. As hypothesized, RFP-Pc12 induced a gradual accumulation of BiP

over time, similar to the effects of tunicamycin treatment (Figure 6C). To assess the

expression of UPR-related genes, samples were taken at 0 and 12 hours post-treatment.

Tunicamycin treatment resulted in the upregulation of transcripts for both ER chaperones

(BiP-like protein *BLP4* and calreticulin *CRT1*) and ER stress-related transcription factors

(*bZIP28* and *bZIP60*). However, Pc12 expression did not show a similar pattern of

upregulation of ER chaperone transcripts. Instead, Pc12 induced an increase in the expression

of transcription factors (Figure 6D). These results suggest that Pc12 induces a distinct form of

ER stress that is different from the effects of tunicamycin.

A specific residue of Rab13-2 is crucial for its interaction with Pc12 and its mutation

evading targeting by Pc12 affect to the virulence of *P. capsici*.

The failure of effectors to suppress plant immunity by manipulating the function of host

targets is thought to contribute factor to non-host resistance (NHR) (McLellan et al. 2022).

Thus, breaking down the interaction between Pc12 and Rab13-2 could reduce the

susceptibility of plants to pathogens. To obtain a better estimate of the biochemical nature of

the interaction between Pc12 and Rab13-2, we used the AlphaFold2 program (Jumper et al.

2021) to identify a pivotal residue on Rab13-2 that interacts with the C-terminal AHQMG

residues of Pc12. Rab13-2 was predicted to have a globular structure, whereas Pc12

displayed a stack of three parallel α-helices connected by a short helical linker (Figure 7A). A

docking model proposed that the three parallel α-helices of Pc12 interact with the globular

region of Rab13-2 (Figure 7A). In particular, L90, A91, and M94 at the C-terminus of Pc12

were predicted to engage with threonine at the 47th position of Rab13-2 (Figure 7B).

Remarkably, replacement of threonine with alanine (Rab13-2^{T47A}) abolished the Rab13-2

interaction with Pc12 (Figure 7C) without affecting its ability to bind REP1 and PRA1

(Figure 7D-E).

These results are important in considering the potential of Rab proteins as a target for

engineering plants resistant to *P. capsici*. To evaluate the applicability of the Rab13-2^{T47A}

mutation for resistance to *P. capsici*, *N. benthamiana* plants expressing Rab13-2^{T47A} were

inoculated with *P. capsici*. At 72 hpi, as the disease progressed, lesion size was significantly

reduced in Rab13-2^{T47A}-expressing leaf compared to Rab13-2-expressing leaf (Figure 7F).

Considering that the effect of endogenous Rab13-2 might diminish the effect of the

expression of Rab13-2^{T47A} expression, this result is significant to suggest that Pc12 is a key

player in determining the necrotrophic phase and Rab13-2^{T47A} may serve as a contributing

factor to delay the onset of the necrotrophic disease progression.

Discussion

Pc12 may serve as a transition effector, orchestrating the transition from the biotrophic

to the necrotrophic life cycle of *P. capsici*.

P. capsici is classified as a hemibiotrophic pathogen, characterized by a biphasic life cycle. It

begins with a biotrophic phase in which it strategically establishes itself within the host plant.

It then transitions to a necrotrophic phase aimed at extracting nutrients from dead cells and

releasing spores for subsequent dissemination. (Kamoun et al. 2015; Parada-Rojas et al. 2021;

Quesada-Ocampo et al. 2023). Although researchers have uncovered the role of RXLR

effectors in increasing susceptibility by interfering with host cell physiology or triggering defense responses through interactions with R proteins during the biotrophic phase, our understanding of how these effectors induce necrosis in host cells remains quite limited. Recently, a case of necrosis was reported in which a *P. infestans* RXLR effector induced nucleolar inflammation and interfered with pre-rRNA 25S processing, leading to necrosis through disruption of protein translation (Lee et al. 2023). This study identified the RXLR effector Pc12 as a critical factor in necrosis, which causes severe ER stress by preventing Rab13-2 recycling in vesicle trafficking. Furthermore, bypassing Pc12 targeting by Rab13-2^{T47A} led to a reduction in necrotrophic lesion size, highlighting that Pc12-induced necrosis contributes to the necrotrophic life cycle of the pathogen. Interestingly, Pc12 expression was highly upregulated at 6 hpi, while the transcripts of NPP1, a necrotrophic marker, gradually increased at 12 hpi (Supplemental Figure 1). This raises the question of why Pc12 transcription increased early in infection despite its role in inducing necrosis. In the experiment using the ethanol-inducible promoter, the accumulation of the Pc12 protein was detectable by immunoblots from 3 h after 5 % ethanol treatment (Figure 6C), and cell death was observed by microscopy before 24 h after 1 % ethanol treatment by microscopy (Supplemental Figure 8). Given the time delay between gene transcription and necrosis induction, it could be inferred that Pc12-induced necrosis serves as an initiator for entry into the necrotrophic phase.

The emergence of Pc12 may benefit *P. capsici* by broadening its host range and promoting survival.

Within the genus *Phytophthora*, only *P. capsici* possesses multiple copies of at least 8 homologs of Pc12 with the potential to induce necrosis. In contrast, *P. ramorum*, *P. cinnamoni*, and *P. sojae* lack this specificity (Supplemental Table 1, Supplemental Figure 2). *P. infestans* lacks a Pc12 homolog, despite its close evolutionary relationship with *P. capsici* (Lamour et al., 2012; Sharma et al., 2015). This suggests that the Pc12 family likely evolved after the emergence of *P. capsici* and underwent lineage-specific expansion within the species. Given the distinctive characteristics of *P. capsici*, the emergence of Pc12 has led to two positive outcomes. First, Pc12 disrupts vesicle trafficking in the basal plant physiology, allowing *P. capsici* to infect a wider range of hosts compared to *P. infestans* and *P. sojae*.

Second, Pc12 rapidly induces necrosis, causing *P. capsici* to shorten its biotrophic phase to

approximately 30 hpi, in contrast to *P. infestans* and *P. sojae* (2 days). This acceleration could

help *P. capsici* to colonize well before the onset of the host defense response, allowing the

pathogen to evade early immune responses. With these advantages conferred by Pc12, *P.*

capsici may have the potential to cause extensive damage to a wide range of hosts, often

resulting in crop losses of up to 100%.

Pc12 disrupts the Rab13-2 cycle by increasing the affinity of REP1 to the active

structure of Rab13-2.

Numerous studies have reported that pathogenic effectors target vesicle-associated proteins to

disrupt the secretion of defense-related proteins or to subvert autophagy to obtain nutrients

for pathogen nutrition (Gu et al. 2017; Pandey et al. 2021; Petre et al. 2021; Yuen et al. 2023).

In our research, we have shown that Pc12 targets Rab13-2 and disrupts vesicle trafficking

between the ER and the Golgi. This disruption leads to the accumulation of apoplastic

proteins in the ER lumen (Figure 6A, Supplemental Figure 8A). Similarly, the viral replicase

p27 of *Red clover necrotic mosaic virus* (RCNMV) inhibits vesicle trafficking between the

ER and Golgi by translocating Arf1 for COPI and Sar1 for COPII to the ER, resulting in the

accumulation of secretory proteins in the ER (Hyodo et al. 2013; Hyodo et al. 2014). We have

also discovered a novel strategy for the RXLR effector Pc12. It mimics the inactive

conformation of Rab13-2 by binding to the active structure of Rab13-2, thereby increasing

the affinity of REP1 for the active form of Rab13-2 (Figure 4, 8). This strategy is similar to

that of brefeldin A, a fungal toxin that increases the affinity of GEFs to bind to GDP-bound

Arf1, ultimately disrupting the cycling of Arf1 between GDP and GTP (Niu et al. 2005).

Given the shared goals of a viral protein and a fungal toxin, Pc12 could contribute

significatnly to the pathogenicity of the pathogen.

Necrosis caused by Pc12 results from a specific form of severe ER stress characterized

by hyper-accumulation of proteins.

Both animals and plants have evolutionarily conserved ER stress responses to a variety of

stimuli (Oakes et al. 2015; Angelos et al. 2017). When ER dysfunction is excessive or

prolonged, cells initiate death signaling, similar to a survival strategy in multicellular

organisms, aimed at eliminating dysfunctional cells (Oakes et al. 2015; Simoni et al. 2022).

ER stress-induced cell death has been extensively studied in animals, particularly in the

context of diseases such as Alzheimer's and Parkinson's (Oakes et al. 2015). However, our

understanding of ER stress-induced cell death in plants is still evolving and some aspects

have yet to be fully elucidated. Pc12-induced inhibition of Rab13-2 recycling in vesicle

trafficking leads to severe ER stress, ultimately culminating in necrosis. According to Yang et

al. 2014, the transcription factor NAC089, transcribed by bZIP28/60, plays a key role in

controlling ERstress-induced programmed cell death in plants by triggering the expression of

PCD-related genes (NAC094, MC5, and BAG6) (Yang et al. 2014). Notably, Pc12 induces a

substantial upregulation of bZIP28/60, but does not lead to a corresponding increase in the

expression of bZIP28/60-downsteam pathway genes (UPR-related genes, Figure 6C) and

PCD-related genes (not data shown). Coupled with the Pc12-induced disruption of vesicle

trafficking between the ER and the Golgi, this implies that Pc12 prevents bZIP28

translocation to the Golgi and prevents bZIP60 splicing in the absence of signaling of

unresolved ER stress. This results in the repression of genes regulated by bZIP28/60 (Figure

6C). Furthermore, Ko et al. (2023) recently reported that ABI5 activates the transcription of

bZIP60 under ER stress, suggesting the interpretation that Pc12 induces the upregulation of

bZIP28/60 independent of bZIP28/60 translocation to the nucleus (Ko et al. 2023). However,

the movement of ABI5 under ER stress remains unknown. Taken together, the failure to

alleviate ER stress by Pc12 may lead to an excessive accumulation of proteins in the ER,

effectively acting as a cellular "time bomb" (Figure 8).

The Rab13-2 point mutation has the potential to manipulate plant resistance to *P.*

***capsici*.**

The underlying molecular mechanism of non-host resistance is thought to arise from 1) pre-

existing barriers that prevent pathogen establishment, 2) the recognition of pathogen effector

proteins that induce robust immunity, and 3) the inability of pathogen effectors to suppress

immunity induced by molecular patterns (McLellan et al. 2022). The inability of effectors to

manipulate host targets for pathogen colonization results in a hostile environment, leading to

the failure of the pathogen to adapt to the host. Importantly, this implies that the evasion

strategy against effectors targets, acting like an orchestra, results in robust resistance,

allowing conversion from host to non-host plants. We identified the specific residue in
Rab13-2 that avoids Pc12 targeting by AF2. Mutation of this residue reduced the
pathogenicity of *P. capsici*. This study provides insight into the idea that host plants can be
engineered to resist *P. capsici* through the synergistic effects of different mutations that
bypass other effectors that collectively contribute to susceptibility.

Materials and methods

Plant materials and growth conditions

N. benthamiana, N. tabacum cv. Samsun*,* and *Solanum lycopersicum* cv. Heinz seeds were
directly sown into damp horticultural bed soil (Biogreen, Seoul, Korea) and cultivated within
a walk-in chamber at 22-24°C under a 16h/8h (day/night) cycle. *Capsicum annuum* cv. ECW
seeds were sterilized for 1 min in a 0.1% sodium hypochlorite solution and germinated in
darkness at 30°C for 7 days. Following this, the pepper seedlings were transplanted into soil
and grown within the same chamber. For virus-induced gene-silencing assays, 3-week-old *N.*
benthamiana plants were utilized. Transgenic *N. benthamiana* plants expressing RFP fused to
HDEL (ER marker, Chakrabarty et al. 2007) and GFP-Rab13-2, GFP-Rab13-2^{Q74L}, and GFP-
Rab13-2^{S29N} transgenic plants were purchased from the Nicotiana Genetic Stock Center (NGSC).
Plants were grown at 22-24°C for 4-5 weeks prior to the corresponding experiments.

P. capsici **culture condition and inoculation assays**

P. capsici strain 40476 was cultured on V8 agar medium for 7 days at 23°C in darkness. The
mycelia were scraped and then incubated under light for 12 hours. To induce the release of
zoospores, sporangia were harvested and placed in distilled water, incubating at 4°C for 30
minutes, followed by 23°C for 30 minutes. A total of 500 zoospores were inoculated on the
detached *N. benthamiana* leaves through droplets. The leaves were placed at 25°C for 2 days
under deem light. The lesion areas were measured at 2 dpi.

Constructs and markers preparation

RXLR effector and their mutants were amplified, incorporating an N-terminal 3xHA epitope
excluding signal peptide and RXLR-EER motifs. Rab proteins and REP1 or PRA were
amplified with N-terminal GFP and 3xFLAG, respectively, employing the overlap PCR

method (Kim et al. 2017). GOIs were inserted into pCambia2300-LIC vector using the

ligation-independent cloning method (Oh et al. 2010). RFP and RFP-Pc12 were amplified

with the attB site to insert into ethanol-inducible expression vector using Gateway cloning

(Invitrogen, USA). The primer sequence used for construction are presented in

Supplementary Table 3. mCherry targeted to Golgi generated previously was used for Golgi

marker (Addgene ID 97401, Park et al. 2017). The membrane bound apoplastic marker

constructs were generated by inserting a signal peptide of *At*Chitinase in front of moxVenus

yellow fluorescence protein sequences (Baleza et al. 2017, Berthold et al. 2019). A vacuolar

membrane maker, TONO-moxVenus was generated by *r*TIP, vacuolar membrane protein,

fusion at the N-terminal of moxVenus (Nelson et al. 2007). Plant expression constructs of

Rab variants, apoplastic side membrane marker, and Tonoplast marker were deposited to the

Addgene (Rab13-2, ID:213473; Rab 13-2^{Q74L}, ID:213474; Rab13-2^{S29N}, ID:213475; APO-

moxVenus, ID: 220490, TONO-moxVenus, ID: 220491) constructs are deposited in Addgene.

Agroinfiltration and quantification of cell death assays

Agrobacterium tumefaciens GV3101 strain or GV2260 containing the various constructs

were cultivated overnight at 28°C in LB medium supplemented with appropriate antibiotics.

The cells were harvested through centrifugation and then resuspended in an infiltration buffer

(10 mM MgCl$_2$, 10 mM MES (pH 5.6), and 200 μM Acetosyringone). Resuspensions were

adjusted to an optical density (OD600) of 0.1-0.3 and leaves of 4-week-plants were

agroinfiltrated for transient expression. To quantify the degree of cell death, *N. benthamiana*

leaves were detached and measured using chlorophyll fluorescence, employing the default

Fv/Fm protocol from a closed FluorCam (Photon Systems Instruments, Czech Republic). The

quantification analysis was conducted using the FluorCam 7.0 software.

Virus-induced gene silencing in *N. benthamiana* and host-induced gene silencing in *P.*

capsici

Virus-induced gene-silencing (VIGS) procedure was conducted described to the protocol

described by Liu et al. (2002). *A. tumefaciens* suspensions containing pTRV1 and pTRV2

with *SGT1, EDS1, ADR1/NRG1,* and *NRC2/3/4* were mixed at a 1:1 ratio in infiltration buffer

(10 mM MES, 10 mM MgCl$_2$, and 200 μM Acetosyringone, pH 5.6) to a final optical density

(O.D.) at 600 nm of 1.5. This mixture was infiltrated into two leaves of 2-week-old *N.*

benthamiana plants. All plants were grown within a walk-in chamber at 24℃ under a 16h/8h

(day/night) cycle. After 3 weeks, the upper leaves were utilized for further experiments

assessing the efficiency of silencing. Host-induced gene silencing (HIGS) was conducted

following the VIGS procedure. After 3 weeks, *P. capsici* strain 40476 was inoculated onto the

detached upper leaves. After 6 hours, four leaf discs containing zoospores were collected to

evaluate the silencing efficiency.

Extraction of DNA and RNA and gene expression analysis by qPCR

To assess the biomass of *P. capsici,* total genomic DNA was extracted from four leaf discs

adjacent to the *P. capsici*-inoculated area using cetyltrimethylammonium bromide (CTAB)

method. For gene expression analysis, total RNA was extracted from four leaf discs using

TRIzol reagent (TR118, MRC, USA), and cDNA synthesis was conducted using Superscript

II (18064014, Invitrogen, USA) following the manufacturer's instructions. Quantitative PCR

(qPCR) and quantitative reverse-transcription PCR (qRT-PCR) was performed utilizing

ExcelTaq™ 2X Q-PCR Master Mix (SYBR, ROX; TQ1110, SMOBIO, Taiwan) with a

CFX96 Touch Real-Time PCR Detection System (Bio-Rad, USA). The expression of the

PcActin gene was normalized to elongation factor-1a of *N. benthamiana* (NbEF-1a), and the

transcript levels were normalized using the internal standard (NbEF-1a). The primer

sequences used in this study are provided in Supplemental Table 4.

Yeast two hybrid assay

Pc12 and truncated Pc12 were cloned into pGBKT7 vector (Takara Bio, Japan), while Rab13-

2 and Rab13-2 mutants were inserted into pGADT7 vector. The recombinant plasmids,

pGBKT7 and pGADT7, were introduced into yeast strain Y2HGold and Y187, respectively.

For yeast two hybrid assay, yeast mating was performed in yeast peptone dextrose adenine

(YPDA) medium at 30℃ overnight. The resulting colonies were recovered on a Synthetic

Defined medium lacking tryptophan and leucine (SD/-Lue-Trp), and the interactions were

validated on SD medium lacking histidine, leucine, and tryptophan (SD/-Lue-Trp-His). The

medium plates were incubated at 28℃ and typically photographed in 5 days. For positive and

negative controls, commercial yeast constructs were used: positive control (pGBKT7-

p53/pGADT7-T) and negative control (pGBKT7-p53/pGADT7-Lam), both provided by

Matchmaker™ Gold Yeast Two-Hybrid System (630489, Clontech, USA).

Co-immunoprecipitation, immunoblot assays, and IP-MS analysis

N. benthamiana leaves infiltrated with *Agrobacterium* were sampled at 24–30 hpi for co-IP or

western blotting. Total protein was extracted using an extraction buffer (10% [v/v] glycerol,

25 mM Tris–HCl [pH 7.5], 1 mM EDTA, 150 mM NaCl, 1% [w/v] polyvinylpolypyrrolidone,

and 1× protease inhibitor cocktail). The extracted proteins were immunoprecipitated with 10

ul of anti-HA magnetic beads (M180-10, MBL, Japan), anti-GFP agarose beads (D153-8,

MBL, Japan), or anti-FLAG agarose beads (651502, BioLegend, USA) and incubated for 4

hours or overnight at 4°C. The beads were washed ten times with immunoprecipitation wash

buffer (GTEN extraction buffer with 0.015% [v/v] Triton X-100) and resuspended in 10 μl

SDS loading dye. Proteins were eluted from the beads by heating at 95°C for 5 min. For

western blotting, the immunoprecipitated and input proteins were separated on SDS–PAGE

gels and transferred onto PVDF membranes via a Trans-Blot Turbo Transfer System (Bio-

Rad, USA). After blocking the membranes with a solution of 5% skim milk prepared in

PBST with 0.1% Tween 20, they were incubated with HRP anti-RFP (1:12000, M204-7,

MBL, Japan), HRP anti-GFP (1:12000, AB6663, Abcam, UK), HRP anti-HA (1:12000,

AB173826, Abcam, UK), or HRP anti-FLAG (1:12000, A8592, Sigma, USA) antibodies at

room temperature for 1 h. The membrane was washed twice with PBST for 10 min each

before ECL (1705061, Bio-Rad, USA) detection was performed according to the

manufacturer's instructions. For IP-Mass analysis, immunoprecipitated proteins were

collected on SDS–PAGE gels and the samples were analyzed using High Resolution

LC/MSMS spectrometer (Q Exactive, Thermo Scientific, USA) in NICEM (National

Instrumentation Center for Environmental Management, College of Agriculture and Life

Sciences, Seoul National University Seoul 151-742, Korea)

GTPase activity assay

For the GTPase activity assay of Rab13-2, Rab 13-2Q74L, and Rab13-2S29N, E. coli codon-

optimized genes were synthesized. Each gene was constructed in a pGEX6-T vector and

transformed in BL21 *E. coli* cells to express GST fusion proteins. GPTase activities were

measured by GTPase-Glo^{TM} Assay (Promega, V7681). GST-tagged protein samples (1 μg)

purified with Pierce® Glutathione Agarose (Thermo, 16100) were used for analysis, and

negative controls were constructed by purifying GST. Both samples were incubated with

GTPase/GAP buffer containing 10 μM GTP and 1 mM DTT followed by the addition of

GTPase-Glo™ reagent. Detection Reagent detects the luminescence of ATP produced from

the remaining GTP after hydrolyzing GTP in the reaction. Luminescence was measured using

a luminometer (Tecan Spark 10M Multimode Plate Reader), Calculate the GTPase activity as

relative luminescence units (RLU). Reaction time was 120 minutes.

Confocal microscopy

For a confocal microscopy, 5 mm^2 *N. benthamiana* leaf discs of the infiltrated region were

observed 36 hours post *Agrobacteria* infiltration. Images were acquired with an Olympus

IX83spinning disk confocal microscope (yokogawa CSW-W1 SoRA) equipped with an

ORCA-Fusion Digital CMOS camera under 60 X oil immersion objective (N.A., 1.3) by

sequential detection of average of 76 Z stacks. Images were acquired by CellSens Dimension

32 software (Olympus). The 488 nm, and 561 nm laser lines (60% power) were used for

EGFP and RFP, respectively. Images were processed using Fiji ImageJ (National Institutes of

Health, Bethesda, Maryland, USA) from the maximum Z intensity projections of the confocal

images. RFP and moxVenus/GFP were pseudo-colored magenta and green respectively.

Protein structure prediction

625 The structures of Pc12 and Rab13-2 were expected using AlphaFold2 in the Google Colab

626 (https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb

627 #scrollTo=svaADwocVdwl). The predicted proteins were visualized using ChimeraX

628 (Pettersen et al. 2021).

Phylogenetic analysis

The 149 Rab family protein sequences were initially acquired from the *N. benthamiana*

genome (v.1.0.1) based on the Ras domain (PF00071) and manually trimmed while

considering the Rab description. Alignments were produced using the Multiple Sequence

Comparison by Log-Expectation (MUSCLE) software in MEGAX. A phylogenetic tree was

constructed using the Maximum Likelihood (ML) method with default parameters and 100

bootstrap replications in MEGAX (Tamura and Nei, 1993; Tamura et al., 2011).

Statistical analysis

Statistical analyses were performed as described in the figure legends. P values were

calculated by Student's t-test using GraphPad Prism software.

Data availability

Sequence information of protein-coding genes used in this study was obtained from the

FungiDB (https://fungidb.org/), NCBI (https://www.ncbi.nlm.nih.gov/), and Sol Genomics

Network (https://solgenomics.net/). The accession numbers for the sequences are as follows:

Niben101Scf09596g00001.1 (Rab13-2), Niben101Scf00684g00002.1 (Rab13-3),

Niben101Scf03277g02014.1 (Rab13-4), Niben101Scf05709g00001.1 (Rab13-10),

Niben101Scf05032g00003.1 (Rab1), Niben101Scf16705g00001.1 (Rab8B),

Niben261Chr06g1220001.1 (REP), and Niben261Chr08g0045015.1 (PRA1). All effector

sequence information is included in Supplemental Table 1. Rab13-2 (ID:213473) and

constitutively active (ID:213474) or inactive mutant (ID:213475), an apoplastic marker (ID:

220490), and a tonoplast marker (ID: 220491) constructs are deposited in Addgene.

Acknowledgements

This work was supported by grants from National Research Foundation of Korea (NRF)

grants funded by the Korean government (MSIT) (NRF-2018R1A5A1023599 [SRC], NRF-

2021R1A2B5B03001613, and NRF-2019R1C1C1008698) to D.C. and US National Science

Foundation awards (NSF EPS-1655726 for E.P.'s start-up and NSF IOS-2126256 to E.P. and

J.C.) and the intramural research program of the U.S. Department of Agriculture, National

Institute of Food and Agriculture Hatch Capacity (# 7000762) to E.P. We thank the UWYO

INBRE program supported by IDeA from NIGMS, NIH, 2P20GM103432 for providing the

spinning disc confocal microscope.

AUTHOR CONTRIBUTIONS

E.P. and D.C. conceptualized the project. J.Kim, J.Kaleku, J.W., H.J., and. S.J. performed

experiments. H.K., M.K., H.J.K., and C.S. provided initial data and materials. J.Kim,

J.Kaleku, J.W. and E.P. analyzed the results. J.Kim, E.P., J.W., and D.C. wrote the manuscript.

E.P., J.W., and D.C. provided funding to conduct the project.

Declaration of interest

The authors declare no competing interests.

Reference

Adams, C.J., Kopp, M.C., Larburu, N., Nowak, P.R., and Ali, M.M.U. (2019). Structure

and molecular mechanism of ER stress signaling by the unfolded protein response signal

activator IRE1. Frontiers in Molecular Biosciences **6,** 11.

Alvim, J.C., Bolt, R.M., An, J., Kamisugi, Y., Cuming, A., Silva-Alvim, F.A.L., Concha,

J.O., daSilva, L.L.P., Hu, M., Hirsz, D., and Denecke, J. (2023). The K/HDEL receptor

does not recycle but instead acts as a Golgi-gatekeeper. Nature Communications **14,** 1612.

Angelos, E., Ruberti, C., Kim, S.-J., and Brandizzi, F. (2017). Maintaining the factory: the

roles of the unfolded protein response in cellular homeostasis in plants. The Plant Journal

90, 671-682.

Balleza, E., Kim, J.M., and Cluzel, P. (2018). Systematic characterization of maturation

time of fluorescent proteins in living cells. Nature Methods **15,** 47-51.

Berthold, F., Roujol, D., Hemmer, C., Jamet, E., Ritzenthaler, C., Hoffmann, L., and

Schmitt-Keichinger, C. (2019). Inside or outside? A new collection of gateway vectors

allowing plant protein subcellular localization or over-expression. Plasmid **105,** 102436.

Boevink, P.C., Birch, P.R.J., Turnbull, D., and Whisson, S.C. (2020). Devastating

intimacy: the cell biology of plant–*Phytophthora* interactions. New Phytologist **228,** 445-

458.

Botër, M., Amigues, B.a., Peart, J., Breuer, C., Kadota, Y., Casais, C., Moore, G.,

Kleanthous, C., Ochsenbein, F., Shirasu, K., and Guerois, R.l. (2007). Structural and

functional analysis of SGT1 reveals that its interaction with HSP90 is required for the

accumulation of Rx, an R protein involved in plant immunity. The Plant Cell **19,** 3791-

3804.

Bourne, H.R., Sanders, D.A., and McCormick, F. (1991). The GTPase superfamily:
conserved structure and molecular mechanism. Nature **349**, 117-127.

Cao, S.S., and Kaufman, R.J. (2012). Unfolded protein response. Current Biology **22,**
R622-R626.

Chakrabarty, R., Banerjee, R., Chung, S.-M., Farman, M., Citovsky, V., Hogenhout,
S.A., Tzfira, T., and Coodin, M. (2007). pSITE vectors for stable integration or transient
expression of autofluorescent protein fusions in plants: probing *Nicotiana benthamiana-*
virus interactions. MPMI **20**, 740-750.

Chaparro-Garcia, A., Wilkinson, R.C., Gimenez-Ibanez, S., Findlay, K., Coffey, M.D.,
Zipfel, C., Rathjen, J.P., Kamoun, S., and Schornack, S. (2011). The receptor-like
kinase SERK3/BAK1 is required for basal resistance against the late blight pathogen
Phytophthora infestans in *Nicotiana benthamiana*. Plos One **6,** e16608.

Cheng, W., Song, X.-S., Li, H.-P., Cao, L.-H., Sun, K., Qiu, X.-L., Xu, Y.-B., Yang, P.,
Huang, T., Zhang, J.-B., Qu, B., and Liao, Y.-C. (2015). Host-induced gene silencing of
an essential chitin synthase gene confers durable resistance to *Fusarium* head blight and
seedling blight in wheat. Plant Biotechnology Journal **13,** 1335-1345.

Dongus, J.A., and Parker, J.E. (2021). EDS1 signalling: at the nexus of intracellular and
surface receptor immunity. Current Opinion in Plant Biology **62,** 102039.

Fan, G., Yang, Y., Li, T., Lu, W., Du, Y., Qiang, X., Wen, Q., and Shan, W. (2018). A
Phytophthora capsici RXLR effector targets and inhibits a plant PPIase to suppress
endoplasmic reticulum-mediated immunity. Molecular Plant **11,** 1067-1083.

Figueroa, C., Taylor, J., and Vojtek, A.B. (2001). Prenylated Rab acceptor protein is a
receptor for prenylated small GTPases. Journal of Biological Chemistry **276,** 28219-28225.

Gray, J.L., von Delft, F., and Brennan, P.E. (2020). Targeting the small GTPase
superfamily through their regulatory proteins. Angewandte Chemie International Edition
59, 6342-6366.

Gu, Y., Zavaliev, R., and Dong, X. (2017). Membrane trafficking in plant immunity.

Molecular Plant **10,** 1026-1034.

Guo, Z., Wu, Y.-W., Das, D., Delon, C., Cramer, J., Yu, S., Thuns, S., Lupilova, N.,
Waldmann, H., Brunsveld, L., Goody, R.S., Alexandrov, K., and Blankenfeldt, W.
(2008). Structures of RabGGTase–substrate/product complexes provide insights into the
evolution of protein prenylation. The EMBO Journal **27,** 2444-2456.

Hyodo, K., Kaido, M., and Okuno, T. (2014). Traffic jam on the cellular secretory pathway
generated by a replication protein from a plant RNA virus. Plant Signaling & Behavior **9,**
e28644.

Hyodo, K., Mine, A., Taniguchi, T., Kaido, M., Mise, K., Taniguchi, H., and Okuno, T.
(2013). ADP ribosylation factor 1 plays an essential role in the replication of a plant RNA
Virus. Journal of Virology **87,** 163-176.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C.,
Kohl, S.A.A., Ballard, A.J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R.,
Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M.,
Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A.W.,
Kavukcuoglu, K., Kohli, P., and Hassabis, D. (2021). Highly accurate protein structure
prediction with AlphaFold. Nature **596,** 583-589.

Kamoun, S., Furzer, O., Jones, J.D.G., Judelson, H.S., Ali, G.S., Dalio, R.J.D., Roy, S.G.,
Schena, L., Zambounis, A., Panabières, F., Cahill, D., Ruocco, M., Figueiredo, A.,
Chen, X.-R., Hulvey, J., Stam, R., Lamour, K., Gijzen, M., Tyler, B.M., Grünwald,
N.J., Mukhtar, M.S., Tomé, D.F.A., Tör, M., Van Den Ackerveken, G., McDowell, J.,
Daayf, F., Fry, W.E., Lindqvist-Kreuze, H., Meijer, H.J.G., Petre, B., Ristaino, J.,
Yoshida, K., Birch, P.R.J., and Govers, F. (2015). The top 10 oomycete pathogens in
molecular plant pathology. Molecular Plant Pathology **16,** 413-434.

Kim, J., Park, M., Jeong, E.S., Lee, J.M., and Choi, D. (2017). Harnessing anthocyanin-
rich fruit: a visible reporter for tracing virus-induced gene silencing in pepper fruit. Plant
Methods **13,** 3.

Kiral, F.R., Kohrs, F.E., Jin, E.J., and Hiesinger, P.R. (2018). Rab GTPases and

Membrane Trafficking in Neurodegeneration. Current Biology **28**, R471-R486.

Ko, D.K., Kim, J.Y., Thibault, E.A., and Brandizzi, F. (2023). An IRE1-proteasome

system signalling cohort controls cell fate determination in unresolved proteotoxic stress

of the plant endoplasmic reticulum. Nature Plants **9,** 1333-1346.

Lamour, K.H., Stam, R., Jupe, J., and Huitema, E. (2012). The oomycete broad-host-

range pathogen *Phytophthora capsici*. Molecular Plant Pathology **13,** 329-337.

Langemeyer, L., Nunes Bastos, R., Cai, Y., Itzen, A., Reinisch, K.M., and Barr, F.A.

(2014). Diversity and plasticity in Rab GTPase nucleotide release mechanism has

consequences for Rab activation and inactivation. eLife **3**, e01623.

Lee, S., Kim, J., Kim, M.-S., Min, C.W., Kim, S.T., Choi, S.-B., Lee, J.H., and Choi, D.

(2023). The *Phytophthora* nucleolar effector Pi23226 targets host ribosome biogenesis to

induce necrotrophic cell death. Plant Communications **4,** 100606.

Lee, S., Lee, Y.J., Choi, S., Park, S.-B., Tran, Q.-G., Heo, J., and Kim, H.-S. (2018).

Development of an alcohol-inducible gene expression system for recombinant protein

expression in *Chlamydomonas reinhardtii*. Journal of Applied Phycology **30,** 2297-2304.

Li, Q., Ai, G., Shen, D., Zou, F., Wang, J., Bai, T., Chen, Y., Li, S., Zhang, M., Jing, M.,

and Dou, D. (2019). A *Phytophthora capsici* effector targets ACD11 binding partners that

regulate ROS-mediated defense response in *Arabidopsis*. Molecular Plant **12,** 565-581.

Li, T., Ai, G., Fu, X., Liu, J., Zhu, H., Zhai, Y., Pan, W., Shen, D., Jing, M., Xia, A., and

Dou, D. (2022). A *Phytophthora capsici* RXLR effector manipulates plant immunity by

targeting RAB proteins and disturbing the protein trafficking pathway. Molecular Plant

Pathology **23,** 1721-1736.

Liu, J.-X., and Howell, S.H. (2016). Managing the protein folding demands in the

endoplasmic reticulum of plants. New Phytologist **211,** 418-428.

McLellan, H., Harvey, S.E., Steinbrenner, J., Armstrong, M.R., He, Q., Clewes, R.,

Pritchard, L., Wang, W., Wang, S., Nussbaumer, T., Dohai, B., Luo, Q., Kumari, P.,

Duan, H., Roberts, A., Boevink, P.C., Neumann, C., Champouret, N., Hein, I, Falter-

Braun, P., Beynon, J., Denby, K., and Birch, P.R.J. (2022). Exploiting breakdown in

nonhost effector–target interactions to boost host disease resistance. Proceedings of the
National Academy of Sciences USA **119,** e2114064119.

Nelson, B.K., Cai, X., and Nebenführ, A. (2007). A multicolored set of in vivo organelle
markers for co-localization studies in *Arabidopsis* and other plants. The Plant Journal **51,**
1126-1136.

Nielsen, E. (2020). The small GTPase superfamily in plants: A conserved regulatory module
with novel functions. Annual Review of Plant Biology **71,** 247-272.

Nielsen, E., Cheung, A.Y., and Ueda, T. (2008). The regulatory RAB and ARF GTPases for
vesicular trafficking. Plant Physiology **147,** 1516-1526.

Niu, T.-K., Pfeifer, A.C., Lippincott-Schwartz, J., and Jackson, C.L. (2005). Dynamics of
GBF1, a brefeldin A-sensitive Arf1 exchange factor at the Golgi. Molecular Biology of the
Cell **16,** 1213-1222.

Oakes, S.A., and Papa, F.R. (2015). The role of endoplasmic reticulum stress in human
pathology. Annual Review of Pathology: Mechanisms of Disease **10,** 173-194.

Pandey, P., Leary, A.Y., Tumtas, Y., Savage, Z., Dagvadorj, B., Duggan, C., Yuen,
E.L.H., Sanguankiattichai, N., Tan, E., Khandare, V., Connerton, A.J., Yunusov, T.,
Madalinski, M., Mirkin, F.G., Schornack, S., Dagdas, Y., Kamoun, S., and Bozkurt,
T.O. (2021). An oomycete effector subverts host vesicle trafficking to channel starvation-
induced autophagy to the pathogen interface. eLife **10,** e65285.

Parada-Rojas, C.H., Granke, L.L., Naegele, R.P., Hansen, Z., Hausbeck, M.K., Kousik,
C.S., McGrath, M.T., D. Smart, C., and Quesada-Ocampo, L.M. (2021). A diagnostic
guide for *Phytophthora capsici* infecting vegetable crops. Plant Health Progress **22,** 404-
414.

Park, E., Lee, H.-Y., Woo, J., Choi, D., and Dinesh-Kumar, S.P. (2017). Spatiotemporal
monitoring of *Pseudomonas syringae* effectors via type III secretion using split fluorescent
protein fragments. The Plant Cell **29,** 1571-1584.

Petre, B., Contreras, M.P., Bozkurt, T.O., Schattat, M.H., Sklenar, J., Schornack, S.,
Abd-El-Haliem, A., Castells-Graells, R., Lozano-Durán, R., Dagdas, Y.F., Menke,

F.L.H., Jones, A.M.E., Vossen, J.H., Robatzek, S., Kamoun, S., and Win, J. (2021).
Host-interactor screens of *Phytophthora infestans* RXLR proteins reveal vesicle
trafficking as a major effector-targeted process. The Plant Cell **33,** 1447-1471.

Pylypenko, O., Hammich, H., Yu, I.M., and Houdusse, A. (2018). Rab GTPases and their
interacting protein partners: structural insights into Rab functional diversity. Small
GTPases **9,** 22-48.

Qiang, X., Zechmann, B., Reitz, M.U., Kogel, K.-H., and Schäfer, P. (2012). The
mutualistic fungus *Piriformospora indica* colonizes *Arabidopsis* roots by inducing an
endoplasmic reticulum stress–triggered caspase-dependent cell death The Plant Cell **24,**
794-809.

Quesada-Ocampo, L.M., Parada-Rojas, C.H., Hansen, Z., Vogel, G., Smart, C.,
Hausbeck, M.K., Carmo, R.M., Huitema, E., Naegele, R.P., Kousik, C.S., Tandy, P.,
and Lamour, K. (2023). *Phytophthora capsici*: recent progress on fundamental biology
and disease management 100 years after its description. Annual Review of
Phytopathology **61,** 185-208.

Reyes-Tena, A., Huguet-Tapia, J.C., Lamour, K.H., Goss, E.M., Rodríguez-Alvarado,
G., Vázquez-Marrufo, G., Santillán-Mendoza, R., and Fernández-Pavía, S.P. (2019).
Genome sequence data of six isolates of *Phytophthora capsici* from Mexico. Molecular
Plant-Microbe Interactions **32,** 1267-1269.

Rojas-Pierce, M. (2013) Targeting of tonoplast proteins to the vacuole. Plant Science **211**,
132-136

Schultink, A., Qi, T., Lee, A., Steinbrenner, A.D., and Staskawicz, B. (2017). Roq1
mediates recognition of the *Xanthomonas* and *Pseudomonas* effector proteins XopQ and
HopQ1. The Plant Journal **92,** 787-795.

Seo, Y.-E., Lee, H.-Y., Kim, H., Yan, X., Park, S.A., Kim, M.-S., Segonzac, C., Choi, D.,
and Mang, H. (2022). The *Phytophthora capsici* RxLR effector CRISIS2 triggers cell
death via suppressing plasma membrane H$^+$-ATPase in the host plant. Journal of
Experimental Botany **74,** 1675-1689.

Sharma, R., Xia, X., Cano, L.M., Evangelisti, E., Kemen, E., Judelson, H., Oome, S.,
Sambles, C., van den Hoogen, D.J., Kitner, M., Klein, J., Meijer, H.J.G., Spring, O.,
Win, J., Zipper, R., Bode, H.B., Govers, F., Kamoun, S., Schornack, S., Studholme,
D.J., Van den Ackerveken, G., and Thines, M. (2015). Genome analyses of the
sunflower pathogen Plasmopara halstedii provide insights into effector evolution in downy
mildews and Phytophthora. BMC Genomics **16**, 741.

Simoni, E.B., Oliveira, C.C., Fraga, O.T., Reis, P.A.B., and Fontes, E.P.B. (2022). Cell
death signaling from endoplasmic reticulum stress: plant-specific and conserved features.
Frontiers in Plant Science **13,** 835738.

Stenmark, H. (2009). Rab GTPases as coordinators of vesicle traffic. Nature Reviews
Molecular Cell Biology **10,** 513-525.

Tomczynska, I., Stumpe, M., and Mauch, F. (2018). A conserved RxLR effector interacts
with host RABA-type GTPases to inhibit vesicle-mediated secretion of antimicrobial
proteins. The Plant Journal **95,** 187-203.

Wawra, S., Trusch, F., Matena, A., Apostolakis, K., Linne, U., Zhukov, I., Stanek, J.,
Koźmiński, W., Davidson, I., Secombes, C.J., Bayer, P., and van West, P. (2017). The
RxLR motif of the host targeting effector AVR3a of *Phytophthora infestans* is cleaved
before secretion. The Plant Cell **29,** 1184-1195.

Whisson, S.C., Boevink, P.C., Moleleki, L., Avrova, A.O., Morales, J.G., Gilroy, E.M.,
Armstrong, M.R., Grouffaud, S., van West, P., Chapman, S., Hein, I., Toth, I.K.,
Pritchard, L., and Birch, P.R.J. (2007). A translocation signal for delivery of oomycete
effector proteins into host plant cells. Nature **450,** 115-118.

Wu, C.-H., Abd-El-Haliem, A., Bozkurt, T.O., Belhaj, K., Terauchi, R., Vossen, J.H.,
and Kamoun, S. (2017). NLR network mediates immunity to diverse plant pathogens.
Proceedings of the National Academy of Sciences USA **114,** 8113-8118.

Yang, Y., Liu, X., Zhang, W., Qian, Q., Zhou, L., Liu, S., Li, Y., and Hou, X. (2021).
Stress response proteins NRP1 and NRP2 are pro-survival factors that inhibit cell death
during ER stress. Plant Physiology **187,** 1414-1427.

Yang, Z.-T., Wang, M.-J., Sun, L., Lu, S.-J., Bi, D.-L., Sun, L., Song, Z.-T., Zhang, S.-S.,
Zhou, S.-F., and Liu, J.-X. (2014). The membrane-associated transcription factor
NAC089 controls ER-stress-induced programmed cell death in plants. Plos Genetics **10,**
e1004243.

Yuen, E.L.H., Shepherd, S., and Bozkurt, T.O. (2023). Traffic control: subversion of plant
membrane trafficking by pathogens. Annual Review of Phytopathology **61,** 325-350.

 Figure and table legends

Figure 1. Pc12 causes cell death independent of defense response in Solanaceae.

(A) Pc12-triggered cell death in *Nicotiana benthamiana, Nicotiana tobacum, Capsicum*

annuum, and *Solanum lycopersicum*. Bax and EGFP were used as a positive control and a

negative control, respectively. The leaves of 4-week-old of plants were agroinfiltrated. The

images were photographed under white and UV light at 2 days after agroinfiltration.

(B) Cell death induced by Pc12 in *N. benthamiana* with silenced NLR-downstream signaling

genes *EDS1, ADR1/NRG1, NRC2/3/4*, and *SGT*. Positive controls consisted of the

combinations Rpiblb2+Avrblb2, R3a+Avr3a, R8+Avr8, and N+p50, while GFP served as the

negative control. Agrobacterium carrying each construct was infiltrated into the leaves of 5-

week-old plants with the respective gene components silenced, and images were taken at 2

dpi.

(C) Cell death observed in images (B) was quantified using the quantum yield (*Fv/Fm*) using

a closed FluorCam system. The data represents the mean ± standard deviation (SD, n = 7-10).

a indicates statistically significant differences by Student t-test (****, $P < 0.0001$).

(D) qRT-PCR analysis of the transcripts level in *N. benthamiana*-silencing *SGT1, EDS1,*

ADR1/NRG1, and *NRC2/3/4*. Leaf disks were sampled at 5-week-old in each components-

silenced *N. benthamiana*. Asterisks denote statistically significant differences by Student t-

test (****, $P < 0.0001$). Data are mean ± SD.

(E) Expression patterns of *PR1, RbohB*, and *WKRY8* in *N. benthamiana* expressing Pc12,

XopQ, and empty vector (EV). Total RNA was extracted in 30 hour-post-infiltration.

Asterisks denote statistically significant differences by Student t-test (*, $P < 0.1$; **, $P < 0.01$;

***, $P < 0.001$; ****, $P < 0.0001$). Data are mean ± SD. (3 experimental repeats).

Figure 2. Pc12 enhances the growth of *P. capsici*, triggering necrotic cell death through

its C-terminus.

(A) Increased growth of *P. capsici* under Pc12 expression compared to a control. RFP and

RFP-Pc12 were expressed by ethanol-inducible promoter in *N. benthamiana*. *P. capsici* was

inoculated on the leaves at 1 day-post-agroinfiltration without ethanol treatment. Images were

taken under UV light at 2 day-post-inoculation. The lesion sizes were measured using ImageJ.

Asterisks denote statistically significant differences using a Student t-test (****, P < 0.0001). Data are mean ± SD.

(B) Relative biomass of *P. capsici* in the leaves shown in Figure 2A. Leaf disks from around the inoculated area were sampled at 2 dpi. Total genomic DNA was extracted and subjected to qPCR analysis. The *P. capsici* biomass was quantified by the *PcActin* normalized to the *NbEF1α*. Asterisks denote statistically significant differences by Student t-test (****, P < 0.0001). Data are mean ± SD.

(C) Suppression of *P. capsici* symptoms through Pc12 silencing via HIGS. Plants were agroinfiltrated with TRV carrying GFP and Pc12. At 14 dpi, *P. capsici* was inoculated on the upper leaves. Images were taken at 2 dpi, and the lesion size was measured by ImageJ.

(D) Relative expression of Pc12 in Pc12-silenced and GFP leaves. The inoculated leaves were sampled at 6 hpi and utilized in qRT-PCR analysis. The transcripts level of Pc12 was normalized to the *PcTubulin*. Asterisks denote statistically significant differences by Student t-test (**, P < 0.01; ****, P < 0.0001). Data are mean ± SD.

(E) Pairwise sequence alignment comparisons of Pc12 homologs of *P. capsici* genome (LT1534). Alignments were obtained using the MUSCLE algorithm and were visualized via ESPript 3.0 (Robert and Gouet 2014). Strictly or highly conserved residues are highlighted in red boxes or blue empty boxes, respectively.

(F) Schematic representation of C-terminal chimeras between Pc12 and Pc22034.

(G) Cell death induced by the Pc12, Pc22034, Pc22034 mutants, and Pc12ΔC5 described in (F)

Figure 3. The Pc12 family interacts with the small GTPase Rab13-2 homologs, directly binding to its active form.

(A) Rab13-2 interacts with Pc12 homologs, inducing cell death. Rab13-2, with a N-terminal GFP, and 3xHA-Pc12 were transiently expressed in plants. Leaf samples were sampled at 30 hpi. Total protein extracts were subjected to co-IP using anti GFP agarose beads. (3 experimental repeats)

(B) Pc12 interacts with the subclade including Rab13-2, -3, and -4. The plants were transiently expressing 3xHA-Pc12 and GFP-Rab13 homologs and sampled at 30 hpi. (3 experimental repeats)

(C) GFP-Rab13-2 proteins partially co-localize at the proximity of ER tubules. GFP-Rab13-2 and RFP-HDEL were transiently expressed in the epidermal cells of *N. benthamiana* leaves. GFP was pseudo-colored to green, while RFP displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

(D) Puncta localization of GFP-Rab13-2 were colocalized with Golgi marker when transiently expressed them in the epidermal cells of *N. benthamiana* leaves. GFP was pseudo-colored to green, while mCherry displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

Figure 4. Pc12 directly binds to the active form of Rab13-2, enhancing the interaction between the active form of Rab13-2 and REP1.

(A) Intrinsic GTPase activity for Rab13-2 and its mutants Rab13-2^{Q74L} and Rab13-2^{S29N} was measured by using recombinant proteins expressed in *E. coli*. As expected, Rab13-2^{Q74L} showed stronger activity statistically significant, while Rab13-2^{S29N} lost substantial level of activity. Experiments were repeated more than five times with the similar results of the same conclusion. One representative graph is shown with 20 technical replicates. Dots on the graph indicate individual technical replicate.

(B) Interaction of Pc12 with Rab13-2 and Rab13-2^{Q74L}, not Rab13-2^{S29N} *in planta*. Pc12 and Pc12ΔC5 were transiently co-expressed with Rab13-2, Rab13-2^{Q74L}, and Rab13-2^{S29N} in *N. benthamiana*.

(C) Physical interaction of Pc12 with Rab13-2 and Rab13-2^{Q74L}, not Rab13-2^{S29N} *in vivo*. Yeast cells transformed with GAL4BD-Pc12 or Pc12ΔC5 and GAL4AD-Rab13-2 or mutants were grown on synthetic media lacking LTH or LT for 7 days. The combination of GAL4BD-p53 and GAL4AD-T was used as a positive control.

(D) Co-IP assays showing that Pc12 increased binding affinity between Rab13-2 and REP. The plants transiently expressing GFP-Rab13-2, REP- or PRA1-3xFLAG, and RFP or RFP-Pc12 were sampled at 12 hours after a 5% ethanol treatment at 1 day-post-agroinfiltration. Total protein extracts were subjected to co-IP using anti GFP agarose beads. The precipitated proteins were immunoblotted. The level of binding was calculated as the ratio between anti-FLAG and anti-FLAG of IP by imageJ. Red asterisks indicate expected band sizes.

(E) Pc12 strengthen the weak interaction between REP and Rab13-2^{Q74L}. The plants were

infiltrated with an agrobacterium carrying REP-3xFLAG and an agrobacterium containing

two plasmids, one plasmid with RFP or RFP-Pc12 and the other with GFP-Rab13-2^{Q74L} or

Rab13-2^{S29N}, in a 1:1 ratio in *N. benthamiana*. Samples treated with 5% ethanol at 1 dpi were

collected at 12 hours post treatment. The level of binding was calculated as the ratio between

anti-FLAG and anti-FLAG of IP by imageJ. Red asterisks indicate expected RFP and RFP-

Pc12 band sizes.

(F) Statistical graph depicting the results of 8 repeated experiments for (E).

Figure 5. Pc12 interrupts trafficking of RFP-HDEL and *Cis*-Golgi marker and relocates

Rab13-2, mimicking the inactive state of Rab13-2.

(A) Pc12 and Pc12ΔC5 are localized in the cytoplasm. GFP-Pc12 as well as GFP- Pc12ΔC5,

a control GFP were infiltrated in the leaves of a transgenic *N. benthamiana* expressing RFP-

HDEL as an ER marker (Magenta). Transient expressions of those proteins were imaged by a

spinning disc confocal microscope. More than 60 z-images in 0.3 *μ*m step size were

superimposed by a maximum z-projection function in ImageJ. The brightness of the original

micrograms was enhanced to improve the visibility of subcellular compartments. This

processing does not change the conclusion drawn from the images. GFP was pseudo-colored

to green, while RFP displayed in magenta. More than 12 images were acquired from 3

independent experiments. Scale bars, 20 *μ*m.

(B) Discontinuity and puncta localization of RFP-HDEL under the expression of Pc12. GFP,

GFP-Pc12 and GFP-Pc12ΔC5 were transiently expressed in transgenic *N. benthamiana*

expressing RFP-HDEL. More than 60 z-images in 0.3 *μ*m step size were acquired by a

spinning disc confocal microscope. Images were superimposed by a maximum z-projection

function in ImageJ. The brightness of the original micrograms was enhanced to improve the

visibility of subcellular compartments. This processing does not change the conclusion drawn

from the images. RFP displayed in magenta. More than 12 images were acquired from 3

independent experiments. Scale bars, 20 *μ*m.

(C) Rapid trafficking of *cis*-Golgi marker, Soybean α-1,2 mannosidase I fused to mCherry,

was interrupted by Pc12 expression. GFP, GFP-Pc12 and GFP-Pc12ΔC5 were transiently

expressed with Golgi marker *N. benthamiana*. Image acquisition and processing were

identical to those for (B). The brightness of the original micrograms was enhanced to

improve the visibility of subcellular compartments. This processing does not change the conclusion drawn from the images. mCherry displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

(D-F) For more accurate localization and dynamics of Rab13-2 and its mutants, transgenic *N. benthamiana* plants expressing both RFP-HDEL and GFP-Rab13-2 (D), GFP- Rab13-2^{Q74L} (E), or GFP-Rab13-2^{S29N} (F) were generated. Stable expressions of those proteins in the leaf epidermal cells were imaged by a spinning disc confocal microscope. More than 60 z-images in 0.3 μm step size were superimposed by a maximum z-projection function in ImageJ. The brightness of the original micrograms was enhanced to improve the visibility of subcellular compartments. This processing does not change the conclusion drawn from the images. GFP was pseudo-colored to green, while RFP displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

Figure 6. Pc12 inhibits the secretion from ER to Golgi, triggering ER stress-mediated cell death.

(A) Apoplastic membrane targeted moxVenus (apoSP-moxVenus-TM) remained in ER in the existence of Pc12. RFP-Pc12, RFP- Pc12ΔC5, and RFP driven by the ethanol promoter were induced by 1% ethanol treatment. Pc12 expression inhibited the secretion of the marker proteins resulted in their accumulation in the ER. Z-images for thicker than 3 μm sections in 0.75 μm step size were acquired by a laser scanning confocal microscope. Images of the corresponding sections were processed to improve the brightness for the clarity. This processing does not change the conclusion drawn from the images. moxVenus was pseudo-colored to green. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

(B) Tonoplast marker (*r*TIP, vacuolar membrane protein, fusion at the N-terminal of moxVenus) was transiently expressed in leaf epidermis of *N. benthamiana*. Trafficking of tonoplast marker takes longer time than other proteins. Thus, tonoplast marker was infiltrated 2 days ahead than RFP, RFP-Pc12, or RFP- Pc12ΔC5. Images were acquired 18 hours (top panel), 28 hours (bottom) panels, and 36 hours (supplemental figures 8B) after spraying 1% ethanol. Images were process with the same procedure described in (D). moxVenus was pseudo-colored to green. More than 12 images were acquired from 3 independent

experiments. Scale bars, 20 μm.

(C) Accumulation of the ER stress marker protein (Bip) in response to Pc12 expression. Plants were treated with mock and Tm (10 ug/ml) and sampled over time. Plants expressing RFP and RFP-Pc12 after 5% ethanol treatment were sampled over time.

(D) Transcripts level of UPR-related transcription factor and ER chaperon genes upon expression of Pc12. In (C), total RNA was extracted at 0 and 12 hours after tunicamycin or ethanol treatment. UPR-related genes, ER chaperon genes (*BLP4* and *CRT1*) and transcription factor genes (*bZIP28* and *bZIP60*), were normalized with *NbEF1a* in qRT-PCR.

Figure 7. A mutation in a key Rab13-2 residue weakens Pc12 binding without affecting REP1 and PRA1 binding, compromising *P. capsici* virulence.

(A-B) The predicted structures of Pc12 and Rab13-2 interaction by AlphaFold2. (B) zooms in on the black box in (A). The green lines depict the intracellular bonds between Pc12 and Rab13-2.

(C - E) *In planta*, a co-immunoprecipitation assay with Rab13-2, Rab13-2^{T47A}, and Pc12 (REP1 or PRA1). Total protein extracts were precipitated with anti HA magnetic beads or anti FLAG agarose. The precipitated protein and total protein were detected by western blotting.

(F) *P. capsici* inoculation on the leaves expressing GFP-Rab13-2 and GFP-Rab13-2^{T47A}. Agrobacterium carrying GFP-Rab13-2 and GFP-Rab13-2^{T47A} was infiltrated in *N. benthamiana,* followed by *P. capsici* inoculation at 1 dpi. Images were taken at 72 hpi, and the lesion size was measured by ImageJ (N = 31 - 32, 3 repeats).

Figure 8. Summary model of the inhibition of Pc12 in Rab13-2-mediated vesicle trafficking.

Prenylated Rab13-2 in its GDP-bound state is inserted to a membrane by Rab escort protein 1 (REP1). The guanine nucleotide exchange factor (GEF) exchanges GDP for GTP on Rab13-2 to initiate vesicle formation for vesicle trafficking. Pc12, secreted from *P. capsici*, binds to GTP-bound Rab13-2, subsequently recruiting REP1 to facilitate the extraction of the complex from the membrane or its retention on the membrane, consequently impeding vesicle formation mediated by Rab13-2. Figure was created using BioRender.com.

 Supplemental data

 Supplemental Table 1.

 Pc12 family in *P. capsici* isolates genome

 Supplemental Table 2.

 Candidates targeted by Pc12 in Mass spectrometry analysis

 Supplemental Table 3.

 Rab13-2 interactors from STRING

 Supplemental Table 4.

 Primer sequences used in this study

 Supplemental Figure 1. The expression pattern of Pc12 during the infection phases of *P.*

 capsici. *PcHmp1* and *PcNpp1* are utilized as markers for the biotrophic and necrotrophic

 phases, respectively. Leaf disks were sampled at regular time intervals. Each gene is

 normalized to *PcTubulin*.

 Supplemental Figure 2. Pc12 homologs from *P. capsici* induce cell death, except for

 PHYCA22034.

 (A) Cell death analysis of Pc12 homologs from *P. capsici, P. ramorum, P. cinnamoni,* and *P.*

 sojae. Pc12 homologs were expressed in 4-week-old *N. benthamiana* and images were

 photographed at 2 days after agroinfiltration.

 (B) The cell death in images (A) was quantified by quantum yield (*Fv/Fm*) using a closed

 FluorCam system. Data are mean ± SD (n = 9-12). a indicates statistically significant

 differences by Student t-test (a, $P < 0.0001$).

 Supplemental Figure 3. Phylogeny tree of Rab family in *N. benthamiana.*

 The red triangle represents the Pc12 target detected in MS, while black triangles represent

 Rab proteins used in the Co-IP shown in Figure 3B.

 Supplemental Figure 4. Screening of Rab13-2 interactor candidates from STRING

 database

1087 Leaves co-expressing Rab13-2 with interactor candidates in Supplemental Table 3 were

1088 sampled at 2 dpi, and total protein extracts were subjected to co-IP using anti-GFP agarose

1089 beads. (3 experimental repeats)

1090

1091 **Supplemental Figure 5. REP1 and PRA1 prefer to interact the inactive form of Rab13-2**

1092 **(Rab13-2^{S29N}).**

1093 Leaves expressing REP1 or PRA1 with Rab13-2 mutants were sampled at 2 dpi, and total

1094 protein extracts were subjected to co-IP using anti-FLAG agarose beads. (3 experimental

1095 repeats)

1096

1097 **Supplemental Figure 6. Repetitions of Co-IP assays with REP, Pc12, and Rab13-2**

1098 **mutants in Figure 4(E).**

1099 Leaves expressing REP, Pc12 with Rab13-2 mutants were sampled at 12 hpt after 5% ethanol

1100 at agrobacteria infiltration, and total protein extracts were subjected to co-IP using anti-GFP

1101 agarose beads.

1102

1103 **Supplemental Figure 7. Colocalization of Golgi and ER with Rab13-2 and its mutants in**

1104 ***N. benthamiana.***

1105 *Cis*-Golgi marker, GO-mCherry, were infiltrated in the epidermal cells of double transgenic

1106 plants of RFP-HDEL and GFP-Rab13-2 (A), GFP- Rab13-2^{Q74L} (B), or GFP-Rab13-2^{S29N} (C)

1107 *N. benthamiana* plants.

1108

Supplemental Figure 8. Different impact of Pc12 on secretory pathway and vacuolar

targeting pathway.

(A) Apoplastic marker displacement upon Pc12 expression occurs on 18 hours after the

induction (hpi) of Pc12. Interestingly RFP-Pc12 is undetectable under microscope at 18 hpi.

(B) 36 hpi of Pc12 expression, most of observed cells are dead. However, intact vacuolar

membranes labeled by tonoplast markers were seldom detected.

1115

1116

1117

1118

Figure 1. Pc12 causes cell death independent of defense response in Solanaceae.

(A) Pc12-triggered cell death in *Nicotiana benthamiana, Nicotiana tobacum, Capsicum annuum*, and *Solanaceae lycopersicum*. Bax and EGFP were used as a positive control and a negative control, respectively. The leaves of 4-week-old of plants were agroinfiltrated. The images were photographed under white and UV light at 2 days after agroinfiltration.

(B) Cell death induced by Pc12 in *N. benthamiana* with silenced NLR-downstream signaling genes *EDS1*, *ADR1/NRG1*, *NRC2/3/4*, and *SGT.* Positive controls consisted of the combinations Rpiblb2+Avrblb2, R3a+Avr3a, R8+Avr8, and N+p50, while GFP served as the negative control. Agrobacterium carrying each construct was infiltrated into the leaves of 5-week-old plants with the respective gene components silenced, and images were taken at 2 dpi.

(C) Cell death observed in images (B) was quantified using the quantum yield (*Fv/Fm*) using a closed FluorCam system. The data represents the mean ± standard deviation (SD, n = 7-10). a indicates statistically significant differences by Student t-test (****, P < 0.0001).

(D) qRT-PCR analysis of the transcripts level in *N. benthamiana*-silencing *SGT1, EDS1, ADR1/NRG1,* and *NRC2/3/4.* Leaf disks were sampled at 5-week-old in each components-silenced *N. benthamiana.* Asterisks denote statistically significant differences by Student t-test (****, P < 0.0001). Data are mean ± SD.

(E) Expression patterns of *PR1, RbohB*, and *WKRY8* in *N. benthamiana* expressing Pc12, XopQ, and empty vector (EV). Total RNA was extracted in 30 hour-post-infiltration. Asterisks denote statistically significant differences by Student t-test (*, P < 0.1; **, P < 0.01; ***, P < 0.001; ****, P < 0.0001). Data are mean ± SD. (3 experimental repeats).

Figure 2. Pc12 enhances the growth of *P. capsici*, triggering necrotic cell death through its C-terminus.
(A) Increased growth of *P. capsici* under Pc12 expression compared to a control. RFP and RFP-Pc12 were expressed by ethanol-inducible promoter in *N. benthamiana*. *P. capsici* was inoculated on the leaves at 1 day-post-agroinfiltration without ethanol treatment. Images were taken under UV light at 2 days post-inoculation. The lesion sizes were measured using ImageJ. Asterisks denote statistically significant differences using a Student t-test (****, P < 0.0001). Data are mean ± SD.
(B) Relative biomass of *P. capsici* in the leaves shown in Figure 2A. Leaf disks from around the inoculated area were sampled at 2 dpi. Total genomic DNA was extracted and subjected to qPCR analysis. The *P. capsici* biomass was quantified by the *PcActin* normalized to the *NbEF1α*. Asterisks denote statistically significant differences by Student t-test (****, P < 0.0001). Data are mean ± SD.
(C) Suppression of *P. capsici* symptoms through Pc12 silencing via HIGS. Plants were agroinfiltrated with TRV carrying GFP and Pc12. At 14 dpi, *P. capsici* was inoculated on the upper leaves. Images were taken at 2 dpi, and the lesion size was measured by ImageJ.
(D) Relative expression of Pc12 in Pc12-silenced and GFP leaves. The inoculated leaves were sampled at 6 hpi and utilized in qRT-PCR analysis. The transcripts level of Pc12 was normalized to the *PcTubulin*. Asterisks denote statistically significant differences by Student t-test (**, P < 0.01; ****, P < 0.0001). Data are mean ± SD.
(E) Pairwise sequence alignment comparisons of Pc12 homologs of *P. capsici* genome (LT1534). Alignments were obtained using the MUSCLE algorithm and were visualized via ESPript 3.0 (Robert and Gouet 2014). Strictly or highly conserved residues are highlighted in red boxes or blue empty boxes, respectively.
(F) Schematic representation of C-terminal chimeras between Pc12 and Pc22034.
(G) Cell death induced by the Pc12, Pc22034, Pc22034 mutants, and Pc12ΔC5 described in (F)

Figure 3.

Figure 3. The Pc12 family interacts with the small GTPase Rab13-2 homologs, directly binding to its active form.

(A) Rab13-2 interacts with Pc12 homologs, inducing cell death. Rab13-2, with a N-terminal GFP, and 3xHA-Pc12 were transiently expressed in plants. Leaf samples were sampled at 30 hpi. Total protein extracts were subjected to co-IP using anti GFP agarose beads. (3 experimental repeats)

(B) Pc12 interacts with the subclade including Rab13-2, -3, and -4. The plants were transiently expressing 3xHA-Pc12 and GFP-Rab13 homologs and sampled at 30 hpi. (3 experimental repeats)

(C) GFP-Rab13-2 proteins partially co-localize at the proximity of ER tubules. GFP-Rab13-2 and RFP-HDEL were transiently expressed in the epidermal cells of *N. benthamiana* leaves. GFP was pseudo-colored to green, while RFP displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

(D) Puncta localization of GFP-Rab13-2 were colocalized with Golgi marker when transiently expressed them in the epidermal cells of *N. benthamiana* leaves. GFP was pseudo-colored to green, while mCherry displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

Figure 4.

Figure 4. Pc12 directly binds to the active form of Rab13-2, enhancing the interaction between the active form of Rab13-2 and REP1.

(A) Intrinsic GTPase activity for Rab13-2 and its mutants Rab13-2^{Q74L} and Rab13-2^{S29N} was measured by using recombinant proteins expressed in *E. coli.* As expected, Rab13-2^{Q74L} showed stronger activity statistically significant, while Rab13-2^{S29N} lost substantial level of activity. Experiments were repeated more than five times with the similar results of the same conclusion. One representative graph is shown with 20 technical replicates. Dots on the graph indicate individual technical replicate.

(B) Interaction of Pc12 with Rab13-2 and Rab13-2^{Q74L}, not Rab13-2^{S29N} *in planta*. Pc12 and Pc12ΔC5 were transiently co-expressed with Rab13-2, Rab13-2^{Q74L}, and Rab13-2^{S29N} in *N. benthamiana*.

(C) Physical interaction of Pc12 with Rab13-2 and Rab13-2^{Q74L}, not Rab13-2^{S29N} *in vivo*. Yeast cells transformed with GAL4BD-Pc12 or Pc12ΔC5 and GAL4AD-Rab13-2 or mutants were grown on synthetic media lacking LTH or LT for 7 days. The combination of GAL4BD-p53 and GAL4AD-T was used as a positive control.

(D) Co-IP assays showing that Pc12 increased binding affinity between Rab13-2 and REP. The plants transiently expressing GFP-Rab13-2, REP- or PRA1-3xFLAG, and RFP or RFP-Pc12 were sampled at 12 hours after a 5% ethanol treatment at 1 day-post-agroinfiltration. Total protein extracts were subjected to co-IP using anti GFP agarose beads. The precipitated proteins were immunoblotted. The level of binding was calculated as the ratio between anti-FLAG and anti-FLAG of IP by imageJ. Red asterisks indicate expected band sizes.

(E) Pc12 strengthen the weak interaction between REP and Rab13-2^{Q74L}. The plants were infiltrated with an agrobacterium carrying REP-3xFLAG and an agrobacterium containing two plasmids, one plasmid with RFP or RFP-Pc12 and the other with GFP-Rab13-2^{Q74L} or Rab13-2^{S29N}, in a 1:1 ratio in *N. benthamiana*. Samples treated with 5% ethanol at 1 dpi were collected at 12 hours post treatment. The level of binding was calculated as the ratio between anti-FLAG and anti-FLAG of IP by imageJ. Red asterisks indicate expected RFP and RFP-Pc12 band sizes.

(F) Statistical graph depicting the results of 8 repeated experiments for (E).

Figure 5.

Figure 5. Pc12 interrupts trafficking of RFP-HDEL and *Cis*-Golgi marker and relocates Rab13-2, mimicking the inactive state of Rab13-2.

(A) Pc12 and Pc12ΔC5 are localized in the cytoplasm. GFP-Pc12 as well as GFP- Pc12ΔC5, a control GFP were infiltrated in the leaves of a transgenic *N. benthamiana* expressing RFP-HDEL as an ER marker (Magenta). Transient expressions of those proteins were imaged by a spinning disc confocal microscope. More than 60 z-images in 0.3 μm step size were superimposed by a maximum z-projection function in ImageJ. The brightness of the original micrograms was enhanced to improve the visibility of subcellular compartments. This processing does not change the conclusion drawn from the images. GFP was pseudo-colored to green, while RFP displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

(B) Discontinuity and puncta localization of RFP-HDEL under the expression of Pc12. GFP, GFP-Pc12 and GFP-Pc12ΔC5 were transiently expressed in transgenic *N. benthamiana* expressing RFP-HDEL. More than 60 z-images in 0.3 μm step size were acquired by a spinning disc confocal microscope. Images were superimposed by a maximum z-projection function in ImageJ. The brightness of the original micrograms was enhanced to improve the visibility of subcellular compartments. This processing does not change the conclusion drawn from the images. RFP displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

(C) Rapid trafficking of *cis*-Golgi marker, Soybean α-1,2 mannosidase I fused to mCherry, was interrupted by Pc12 expression. GFP, GFP-Pc12 and GFP-Pc12ΔC5 were transiently expressed with Golgi marker *N. benthamiana*. Image acquisition and processing were identical to those for (B). The brightness of the original micrograms was enhanced to improve the visibility of subcellular compartments. This processing does not change the conclusion drawn from the images. mCherry displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

(D-F) For more accurate localization and dynamics of Rab13-2 and its mutants, transgenic *N. benthamiana* plants expressing both RFP-HDEL and GFP-Rab13-2 (D), GFP- Rab13-2^{Q74L} (E), or GFP-Rab13-2^{S29N} (F) were generated. Stable expressions of those proteins in the leaf epidermal cells were imaged by a spinning disc confocal microscope. More than 60 z-images in 0.3 μm step size were superimposed by a maximum z-projection function in ImageJ. The brightness of the original micrograms was enhanced to improve the visibility of subcellular compartments. This processing does not change the conclusion drawn from the images. GFP was pseudo-colored to green, while RFP displayed in magenta. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

Figure 6.

Figure 6. Pc12 inhibits the secretion from ER to Golgi, triggering ER stress-mediated cell death.

(A) Apoplastic membrane targeted moxVenus (apoSP-moxVenus-TM) remained in ER in the existence of Pc12. RFP-Pc12, RFP- Pc12ΔC5, and RFP driven by the ethanol promoter were induced by 1% ethanol treatment. Pc 12 expression inhibited the secretion of the marker proteins resulted in their accumulation in the ER. Z-images fo r thicker than 3 μm sections in 0.75 μm step size were acquired by a laser scanning confocal microscope. Image s of the corresponding sections were processed to improve the brightness for the clarity. This processing does n ot change the conclusion drawn from the images. moxVenus was pseudo-colored to green. More than 12 image s were acquired from 3 independent experiments. Scale bars, 20 μm.

(B) Tonoplast marker (*r*TIP, vacuolar membrane protein, fusion at the N-terminal of moxVenus) was transiently e xpressed in leaf epidermis of *N. benthamiana*. Trafficking of tonoplast marker takes longer time than other protei ns. Thus, tonoplast marker was infiltrated 2 days ahead than RFP, RFP-Pc12, or RFP- Pc12ΔC5. Images were a cquired 18 hours (top panel), 28 hours (bottom) panels, and 36 hours (supplemental figures 8B) after spraying 1 % ethanol. Images were process with the same procedure described in (D). moxVenus was pseudo-colored to g reen. More than 12 images were acquired from 3 independent experiments. Scale bars, 20 μm.

(C) Accumulation of the ER stress marker protein (Bip) in response to Pc12 expression. Plants were treated with mock and Tm (10 ug/ml) and sampled over time. Plants expressing RFP and RFP-Pc12 after 5% ethanol treatm ent were sampled over time.

(D) Transcripts level of UPR-related transcription factor and ER chaperon genes upon expression of Pc12. In (C), total RNA was extracted at 0 and 12 hours after tunicamycin or ethanol treatment. UPR-related genes, ER chaperon genes (*BLP4* and *CRT1*) and transcription factor genes (*bZIP28* and *bZIP60*), were normalized with *NbEF1a* in qRT-PCR.

Figure 7

Figure 7 A mutation in a key Rab13-2 residue weakens Pc12 binding without affecting REP1 and PRA1 binding, compromising *P. capsici* virulence.

(A-B) The predicted structures of Pc12 and Rab13-2 interaction by AlphaFold2. (B) zooms in on the black box in (A). The green lines depict the intracellular bonds between Pc12 and Rab13-2.

(C - E) *In planta*, a co-immunoprecipitation assay with Rab13-2, Rab13-2^{T47A}, and Pc12 (REP1 or PRA1). Total protein extracts were precipitated with anti HA magnetic beads or anti FLAG agarose. The precipitated protein and total protein were detected by western blotting.

(F) *P. capsici* inoculation on the leaves expressing GFP-Rab13-2 and GFP-Rab13-2^{T47A}. Agrobacterium carrying GFP-Rab13-2 and GFP-Rab13-2^{T47A} was infiltrated in *N. benthamiana,* followed by *P. capsici* inoculation at 1 dpi. Images were taken at 72 hpi, and the lesion size was measured by ImageJ. The graph displays the mean and standard deviation of all the data (N = 26-35, 4 repeats).

Figure 8. Summary model of the inhibition of Pc12 in Rab13-2-mediated vesicle trafficking.

Prenylated Rab13-2 in its GDP-bound state is inserted to a membrane by Rab escort protein 1 (REP1). The guanine nucleotide exchange factor (GEF) exchanges GDP for GTP on Rab13-2 to initiate vesicle formation for vesicle trafficking. Pc12, secreted from *P. capsici*, binds to GTP-bound Rab13-2, subsequently recruiting REP1 to facilitate the extraction of the complex from the membrane, consequently impeding vesicle formation mediated by Rab13-2. Figure was created using BioRender.com.

Parsed Citations

Adams, C.J., Kopp, M.C., Larburu, N., Nowak, P.R., and Ali, M.M.U. (2019). Structure and molecular mechanism of ER stress signaling by the unfolded protein response signal activator IRE1. Frontiers in Molecular Biosciences 6, 11.
  Google Scholar: Author Only Title Only Author and Title

Alvim, J.C., Bolt, R.M., An, J., Kamisugi, Y., Cuming, A, Silva-Alvim, F.A.L., Concha, J.O., daSilva, L.L.P., Hu, M., Hirsz, D., and Denecke, J. (2023). The K/HDEL receptor does not recycle but instead acts as a Golgi-gatekeeper. Nature Communications 14, 1612.
  Google Scholar: Author Only Title Only Author and Title

Angelos, E., Ruberti, C., Kim, S.-J., and Brandizzi, F. (2017). Maintaining the factory: the roles of the unfolded protein response in cellular homeostasis in plants. The Plant Journal 90, 671-682.
  Google Scholar: Author Only Title Only Author and Title

Balleza, E., Kim, J.M., and Cluzel, P. (2018). Systematic characterization of maturation time of fluorescent proteins in living cells. Nature Methods 15, 47-51.
  Google Scholar: Author Only Title Only Author and Title

Berthold, F., Roujol, D., Hemmer, C., Jamet, E., Ritzenthaler, C., Hoffmann, L., and Schmitt-Keichinger, C. (2019). Inside or outside? A new collection of gateway vectors allowing plant protein subcellular localization or over-expression. Plasmid 105, 102436.
  Google Scholar: Author Only Title Only Author and Title

Boevink, P.C., Birch, P.R.J., Turnbull, D., and Whisson, S.C. (2020). Devastating intimacy: the cell biology of plant–Phytophthora interactions. New Phytologist 228, 445-458.
  Google Scholar: Author Only Title Only Author and Title

Botër, M., Amigues, B.a., Peart, J., Breuer, C., Kadota, Y., Casais, C., Moore, G., Kleanthous, C., Ochsenbein, F., Shirasu, K., and Guerois, R.I. (2007). Structural and functional analysis of SGT1 reveals that its interaction with HSP90 is required for the accumulation of Rx, an R protein involved in plant immunity. The Plant Cell 19, 3791-3804.
  Google Scholar: Author Only Title Only Author and Title

Bourne, H.R., Sanders, D.A., and McCormick, F. (1991). The GTPase superfamily: conserved structure and molecular mechanism. Nature 349, 117-127.
  Google Scholar: Author Only Title Only Author and Title

Cao, S.S., and Kaufman, R.J. (2012). Unfolded protein response. Current Biology 22, R622-R626.
  Google Scholar: Author Only Title Only Author and Title

Chakrabarty, R., Banerjee, R., Chung, S.-M., Farman, M., Citovsky, V., Hogenhout, S.A, Tzfira, T., and Coodin, M. (2007). pSITE vectors for stable integration or transient expression of autofluorescent protein fusions in plants: probing Nicotiana benthamiana-virus interactions. MPMI 20, 740-750.
  Google Scholar: Author Only Title Only Author and Title

Chaparro-Garcia, A, Wilkinson, R.C., Gimenez-Ibanez, S., Findlay, K., Coffey, M.D., Zipfel, C., Rathjen, J.P., Kamoun, S., and Schornack, S. (2011). The receptor-like kinase SERK3/BAK1 is required for basal resistance against the late blight pathogen Phytophthora infestans in Nicotiana benthamiana. Plos One 6, e16608.
  Google Scholar: Author Only Title Only Author and Title

Cheng, W., Song, X.-S., Li, H.-P., Cao, L.-H., Sun, K., Qiu, X.-L., Xu, Y.-B., Yang, P., Huang, T., Zhang, J.-B., Qu, B., and Liao, Y.-C. (2015). Host-induced gene silencing of an essential chitin synthase gene confers durable resistance to Fusarium head blight and seedling blight in wheat. Plant Biotechnology Journal 13, 1335-1345.
  Google Scholar: Author Only Title Only Author and Title

Dongus, J.A, and Parker, J.E. (2021). EDS1 signalling: at the nexus of intracellular and surface receptor immunity. Current Opinion in Plant Biology 62, 102039.
  Google Scholar: Author Only Title Only Author and Title

Fan, G., Yang, Y., Li, T., Lu, W., Du, Y., Qiang, X., Wen, Q., and Shan, W. (2018). A Phytophthora capsici RXLR effector targets and inhibits a plant PPIase to suppress endoplasmic reticulum-mediated immunity. Molecular Plant 11, 1067-1083.
  Google Scholar: Author Only Title Only Author and Title

Figueroa, C., Taylor, J., and Vojtek, A.B. (2001). Prenylated Rab acceptor protein is a receptor for prenylated small GTPases. Journal of Biological Chemistry 276, 28219-28225.
  Google Scholar: Author Only Title Only Author and Title

Gray, J.L., von Delft, F., and Brennan, P.E. (2020). Targeting the small GTPase superfamily through their regulatory proteins. Angewandte Chemie International Edition 59, 6342-6366.
  Google Scholar: Author Only Title Only Author and Title

Gu, Y., Zavaliev, R., and Dong, X. (2017). Membrane trafficking in plant immunity. Molecular Plant 10, 1026-1034.
Google Scholar: Author Only Title Only Author and Title

Guo, Z., Wu, Y.-W., Das, D., Delon, C., Cramer, J., Yu, S., Thuns, S., Lupilova, N., Waldmann, H., Brunsveld, L., Goody, R.S., Alexandrov, K., and Blankenfeldt, W. (2008). Structures of RabGGTase–substrate/product complexes provide insights into the evolution of protein prenylation. The EMBO Journal 27, 2444-2456.
Google Scholar: Author Only Title Only Author and Title

Hyodo, K., Kaido, M., and Okuno, T. (2014). Traffic jam on the cellular secretory pathway generated by a replication protein from a plant RNA virus. Plant Signaling & Behavior 9, e28644.
Google Scholar: Author Only Title Only Author and Title

Hyodo, K., Mine, A., Taniguchi, T., Kaido, M., Mise, K., Taniguchi, H., and Okuno, T. (2013). ADP ribosylation factor 1 plays an essential role in the replication of a plant RNA Virus. Journal of Virology 87, 163-176.
Google Scholar: Author Only Title Only Author and Title

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard, A.J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A.W., Kavukcuoglu, K., Kohli, P., and Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. Nature 596, 583-589.
Google Scholar: Author Only Title Only Author and Title

Kamoun, S., Furzer, O., Jones, J.D.G., Judelson, H.S., Ali, G.S., Dalio, R.J.D., Roy, S.G., Schena, L., Zambounis, A., Panabières, F., Cahill, D., Ruocco, M., Figueiredo, A., Chen, X.-R., Hulvey, J., Stam, R., Lamour, K., Gijzen, M., Tyler, B.M., Grünwald, N.J., Mukhtar, M.S., Tomé, D.F.A., Tör, M., Van Den Ackerveken, G., McDowell, J., Daayf, F., Fry, W.E., Lindqvist-Kreuze, H., Meijer, H.J.G., Petre, B., Ristaino, J., Yoshida, K., Birch, P.R.J., and Govers, F. (2015). The top 10 oomycete pathogens in molecular plant pathology. Molecular Plant Pathology 16, 413-434.
Google Scholar: Author Only Title Only Author and Title

Kim, J., Park, M., Jeong, E.S., Lee, J.M., and Choi, D. (2017). Harnessing anthocyanin-rich fruit: a visible reporter for tracing virus-induced gene silencing in pepper fruit. Plant Methods 13, 3.
Google Scholar: Author Only Title Only Author and Title

Kiral, F.R., Kohrs, F.E., Jin, E.J., and Hiesinger, P.R. (2018). Rab GTPases and Membrane Trafficking in Neurodegeneration. Current Biology 28, R471-R486.
Google Scholar: Author Only Title Only Author and Title

Ko, D.K., Kim, J.Y., Thibault, E.A., and Brandizzi, F. (2023). An IRE1-proteasome system signalling cohort controls cell fate determination in unresolved proteotoxic stress of the plant endoplasmic reticulum. Nature Plants 9, 1333-1346.
Google Scholar: Author Only Title Only Author and Title

Lamour, K.H., Stam, R., Jupe, J., and Huitema, E. (2012). The oomycete broad-host-range pathogen Phytophthora capsici. Molecular Plant Pathology 13, 329-337.
Google Scholar: Author Only Title Only Author and Title

Langemeyer, L., Nunes Bastos, R., Cai, Y., Itzen, A., Reinisch, K.M., and Barr, F.A. (2014). Diversity and plasticity in Rab GTPase nucleotide release mechanism has consequences for Rab activation and inactivation. eLife 3, e01623.
Google Scholar: Author Only Title Only Author and Title

Lee, S., Kim, J., Kim, M.-S., Min, C.W., Kim, S.T., Choi, S.-B., Lee, J.H., and Choi, D. (2023). The Phytophthora nucleolar effector Pi23226 targets host ribosome biogenesis to induce necrotrophic cell death. Plant Communications 4, 100606.
Google Scholar: Author Only Title Only Author and Title

Lee, S., Lee, Y.J., Choi, S., Park, S.-B., Tran, Q.-G., Heo, J., and Kim, H.-S. (2018). Development of an alcohol-inducible gene expression system for recombinant protein expression in Chlamydomonas reinhardtii. Journal of Applied Phycology 30, 2297-2304.
Google Scholar: Author Only Title Only Author and Title

Li, Q., Ai, G., Shen, D., Zou, F., Wang, J., Bai, T., Chen, Y., Li, S., Zhang, M., Jing, M., and Dou, D. (2019). A Phytophthora capsici effector targets ACD11 binding partners that regulate ROS-mediated defense response in Arabidopsis. Molecular Plant 12, 565-581.
Google Scholar: Author Only Title Only Author and Title

Li, T., Ai, G., Fu, X., Liu, J., Zhu, H., Zhai, Y., Pan, W., Shen, D., Jing, M., Xia, A., and Dou, D. (2022). A Phytophthora capsici RXLR effector manipulates plant immunity by targeting RAB proteins and disturbing the protein trafficking pathway. Molecular Plant Pathology 23, 1721-1736.
Google Scholar: Author Only Title Only Author and Title

Liu, J.-X., and Howell, S.H. (2016). Managing the protein folding demands in the endoplasmic reticulum of plants. New Phytologist

211, 418-428.
  Google Scholar: Author Only Title Only Author and Title

McLellan, H., Harvey, S.E., Steinbrenner, J., Armstrong, M.R., He, Q., Clewes, R., Pritchard, L., Wang, W., Wang, S., Nussbaumer, T., Dohai, B., Luo, Q., Kumari, P., Duan, H., Roberts, A, Boevink, P.C., Neumann, C., Champouret, N., Hein, I., Falter-Braun, P., Beynon, J., Denby, K., and Birch, P.R.J. (2022). Exploiting breakdown in nonhost effector–target interactions to boost host disease resistance. Proceedings of the National Academy of Sciences USA 119, e2114064119.
  Google Scholar: Author Only Title Only Author and Title

Nelson, B.K., Cai, X., and Nebenführ, A. (2007). A multicolored set of in vivo organelle markers for co-localization studies in Arabidopsis and other plants. The Plant Journal 51, 1126-1136.
  Google Scholar: Author Only Title Only Author and Title

Nielsen, E. (2020). The small GTPase superfamily in plants: A conserved regulatory module with novel functions. Annual Review of Plant Biology 71, 247-272.
  Google Scholar: Author Only Title Only Author and Title

Nielsen, E., Cheung, A.Y., and Ueda, T. (2008). The regulatory RAB and ARF GTPases for vesicular trafficking. Plant Physiology 147, 1516-1526.
  Google Scholar: Author Only Title Only Author and Title

Niu, T.-K., Pfeifer, A.C., Lippincott-Schwartz, J., and Jackson, C.L. (2005). Dynamics of GBF1, a brefeldin A-sensitive Arf1 exchange factor at the Golgi. Molecular Biology of the Cell 16, 1213-1222.
  Google Scholar: Author Only Title Only Author and Title

Oakes, S.A., and Papa, F.R. (2015). The role of endoplasmic reticulum stress in human pathology. Annual Review of Pathology: Mechanisms of Disease 10, 173-194.
  Google Scholar: Author Only Title Only Author and Title

Pandey, P., Leary, A.Y., Tumtas, Y., Savage, Z, Dagvadorj, B., Duggan, C., Yuen, E.L.H., Sanguankiattichai, N., Tan, E., Khandare, V., Connerton, A.J., Yunusov, T., Madalinski, M., Mirkin, F.G., Schornack, S., Dagdas, Y., Kamoun, S., and Bozkurt, T.O. (2021). An oomycete effector subverts host vesicle trafficking to channel starvation-induced autophagy to the pathogen interface. eLife 10, e65285.
  Google Scholar: Author Only Title Only Author and Title

Parada-Rojas, C.H., Granke, L.L., Naegele, R.P., Hansen, Z, Hausbeck, M.K., Kousik, C.S., McGrath, M.T., D. Smart, C., and Quesada-Ocampo, L.M. (2021). A diagnostic guide for Phytophthora capsici infecting vegetable crops. Plant Health Progress 22, 404-414.
  Google Scholar: Author Only Title Only Author and Title

Park, E., Lee, H.-Y., Woo, J., Choi, D., and Dinesh-Kumar, S.P. (2017). Spatiotemporal monitoring of Pseudomonas syringae effectors via type III secretion using split fluorescent protein fragments. The Plant Cell 29, 1571-1584.
  Google Scholar: Author Only Title Only Author and Title

Petre, B., Contreras, M.P., Bozkurt, T.O., Schattat, M.H., Sklenar, J., Schornack, S., Abd-El-Haliem, A, Castells-Graells, R., Lozano-Durán, R., Dagdas, Y.F., Menke, F.L.H., Jones, A.M.E., Vossen, J.H., Robatzek, S., Kamoun, S., and Win, J. (2021). Host-interactor screens of Phytophthora infestans RXLR proteins reveal vesicle trafficking as a major effector-targeted process. The Plant Cell 33, 1447-1471.
  Google Scholar: Author Only Title Only Author and Title

Pylypenko, O., Hammich, H., Yu, I.M., and Houdusse, A. (2018). Rab GTPases and their interacting protein partners: structural insights into Rab functional diversity. Small GTPases 9, 22-48.
  Google Scholar: Author Only Title Only Author and Title

Qiang, X., Zechmann, B., Reitz, M.U., Kogel, K.-H., and Schäfer, P. (2012). The mutualistic fungus Piriformospora indica colonizes Arabidopsis roots by inducing an endoplasmic reticulum stress–triggered caspase-dependent cell death The Plant Cell 24, 794-809.
  Google Scholar: Author Only Title Only Author and Title

Quesada-Ocampo, L.M., Parada-Rojas, C.H., Hansen, Z, Vogel, G., Smart, C., Hausbeck, M.K., Carmo, R.M., Huitema, E., Naegele, R.P., Kousik, C.S., Tandy, P., and Lamour, K. (2023). Phytophthora capsici: recent progress on fundamental biology and disease management 100 years after its description. Annual Review of Phytopathology 61, 185-208.
  Google Scholar: Author Only Title Only Author and Title

Reyes-Tena, A, Huguet-Tapia, J.C., Lamour, K.H., Goss, E.M., Rodríguez-Alvarado, G., Vázquez-Marrufo, G., Santillán-Mendoza, R., and Fernández-Pavía, S.P. (2019). Genome sequence data of six isolates of Phytophthora capsici from Mexico. Molecular Plant-Microbe Interactions 32, 1267-1269.
  Google Scholar: Author Only Title Only Author and Title

Rojas-Pierce, M. (2013) Targeting of tonoplast proteins to the vacuole. Plant Science 211, 132-136

Google Scholar: Author Only Title Only Author and Title

Schultink, A, Qi, T., Lee, A, Steinbrenner, A.D., and Staskawicz, B. (2017). Roq1 mediates recognition of the Xanthomonas and Pseudomonas effector proteins XopQ and HopQ1. The Plant Journal 92, 787-795.
Google Scholar: Author Only Title Only Author and Title

Seo, Y.-E., Lee, H.-Y., Kim, H., Yan, X., Park, S.A., Kim, M.-S., Segonzac, C., Choi, D., and Mang, H. (2022). The Phytophthora capsici RxLR effector CRISIS2 triggers cell death via suppressing plasma membrane H+-ATPase in the host plant. Journal of Experimental Botany 74, 1675-1689.
Google Scholar: Author Only Title Only Author and Title

Sharma, R., Xia, X., Cano, L.M., Evangelisti, E., Kemen, E., Judelson, H., Oome, S., Sambles, C., van den Hoogen, D.J., Kitner, M., Klein, J., Meijer, H.J.G., Spring, O., Win, J., Zipper, R., Bode, H.B., Govers, F., Kamoun, S., Schornack, S., Studholme, D.J., Van den Ackerveken, G., and Thines, M. (2015). Genome analyses of the sunflower pathogen Plasmopara halstedii provide insights into effector evolution in downy mildews and Phytophthora. BMC Genomics 16, 741.
Google Scholar: Author Only Title Only Author and Title

Simoni, E.B., Oliveira, C.C., Fraga, O.T., Reis, P.A.B., and Fontes, E.P.B. (2022). Cell death signaling from endoplasmic reticulum stress: plant-specific and conserved features. Frontiers in Plant Science 13, 835738.
Google Scholar: Author Only Title Only Author and Title

Stenmark, H. (2009). Rab GTPases as coordinators of vesicle traffic. Nature Reviews Molecular Cell Biology 10, 513-525.
Google Scholar: Author Only Title Only Author and Title

Tomczynska, I., Stumpe, M., and Mauch, F. (2018). A conserved RxLR effector interacts with host RABA-type GTPases to inhibit vesicle-mediated secretion of antimicrobial proteins. The Plant Journal 95, 187-203.
Google Scholar: Author Only Title Only Author and Title

Wawra, S., Trusch, F., Matena, A, Apostolakis, K., Linne, U., Zhukov, I., Stanek, J., Koźmiński, W., Davidson, I., Secombes, C.J., Bayer, P., and van West, P. (2017). The RxLR motif of the host targeting effector AVR3a of Phytophthora infestans is cleaved before secretion. The Plant Cell 29, 1184-1195.
Google Scholar: Author Only Title Only Author and Title

Whisson, S.C., Boevink, P.C., Moleleki, L., Avrova, A.O., Morales, J.G., Gilroy, E.M., Armstrong, M.R., Grouffaud, S., van West, P., Chapman, S., Hein, I., Toth, I.K., Pritchard, L., and Birch, P.R.J. (2007). A translocation signal for delivery of oomycete effector proteins into host plant cells. Nature 450, 115-118.
Google Scholar: Author Only Title Only Author and Title

Wu, C.-H., Abd-El-Haliem, A, Bozkurt, T.O., Belhaj, K., Terauchi, R., Vossen, J.H., and Kamoun, S. (2017). NLR network mediates immunity to diverse plant pathogens. Proceedings of the National Academy of Sciences USA 114, 8113-8118.
Google Scholar: Author Only Title Only Author and Title

Yang, Y., Liu, X., Zhang, W., Qian, Q., Zhou, L., Liu, S., Li, Y., and Hou, X. (2021). Stress response proteins NRP1 and NRP2 are pro-survival factors that inhibit cell death during ER stress. Plant Physiology 187, 1414-1427.
Google Scholar: Author Only Title Only Author and Title

Yang, Z.-T., Wang, M.-J., Sun, L., Lu, S.-J., Bi, D.-L., Sun, L., Song, Z.-T., Zhang, S.-S., Zhou, S.-F., and Liu, J.-X. (2014). The membrane-associated transcription factor NAC089 controls ER-stress-induced programmed cell death in plants. Plos Genetics 10, e1004243.
Google Scholar: Author Only Title Only Author and Title

Yuen, E.L.H., Shepherd, S., and Bozkurt, T.O. (2023). Traffic control: subversion of plant membrane trafficking by pathogens. Annual Review of Phytopathology 61, 325-350.
Google Scholar: Author Only Title Only Author and Title