Development of primary osteoarthritis during aging in genetically diverse UM-HET3 mice

Sher Bahadur Poudel[1], Ryan R Ruff[2], Gozde Yildirim[1], Richard A Miller[3], David E Harrison[4], Randy Strong[5,6], Thorsten Kirsch[7,8], and Shoshana Yakar[1].

[1]David B. Kriser Dental Center, Department of Molecular Pathobiology, New York University College of Dentistry, New York, NY 10010-4086, USA
[2]David B. Kriser Dental Center, Department of Epidemiology and Health Promotion, New York University College of Dentistry New York, NY 10010-4086
[3]Pathology and Geriatrics Center, University of Michigan, Ann Arbor, MI 48105, USA.
[4]The Jackson Laboratory, Bar Harbor, ME 04609, USA.
[5]Geriatric Research, Education and Clinical Center and Research Service, South Texas Veterans Health Care System, San Antonio, TX 78229, USA.
[6]Barshop Institute for Longevity and Aging Studies and Department of Pharmacology, The University of Texas Health Science Center, San Antonio, TX 78229, USA.
[7]Department of Orthopaedic Surgery, NYU Grossman School of Medicine, New York, NY 10100, USA
[8]Department of Biomedical Engineering, NYU Tandon School of Engineering, NY 10010, USA

Running title: Primary OA in UM-HET3 mice

Key words: osteoarthritis, UM-HET3, sub-chondral bone, cartilage, methylene blue, mitoquinone, antioxidants

Correspondence:
Shoshana Yakar, Ph.D.
Professor
New York University College of Dentistry
David B. Kriser Dental Center
Department of Molecular Pathobiology
345 East 24th Street New York, New York 10010-4086
Tel : 201-316-4979
sy1007@nyu.edu
ORCiD https://orcid.org/0000-0002-2352-1330

Statements:

- All authors concur with the submission. The material submitted for publication has not been previously reported and is not under consideration for publication elsewhere.

- **Data availability statement**: The datasets generated and analyzed during the current study are available upon request. Our studies do not include the use of custom code or mathematical algorithms. We have included citations for available data in the references section.

- **Funding statement**: Financial support received from the National Institutes of Health Grant R01AG056397, and B01 (2024) Department of Molecular Pathobiology Accelerator Award to SY. SBP is supported by New York University Provost's Postdoctoral Fellowship Program. This work was also supported by the National Institutes of Health grant to The Jackson Laboratory Nathan Shock Center of Excellence in the Basic Biology of Aging AG038070, U01-AG022303 to RAM, UO1-AG022308 to DEH, U01-AG013319 to RS, RS is supported by a Senior Research Career Scientist Award from the Department of Veterans Affairs Office of Research and Development, and S10 OD010751-01A1 for micro-computed tomography.

Conflict of interest disclosure: The authors declare no conflict of interest. All authors have discussed the results and approved the final version of the manuscript. SY is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

- **Ethics approval statement**: Animal protocol was reviewed and approved by the Institutional Animal Care and Use Committees of the ITP centers

Abstract:

This study investigated the prevalence and progression of primary osteoarthritis (OA) in aged UM-HET3 mice. Using the Osteoarthritis Research Society International (OARSI) scoring system, we assessed articular cartilage (AC) integrity in 182 knee joints of 22-25 months old mice.

We found a high prevalence of primary OA in both sexes of mice, with 85% of male and 90% of female mice showing varying degrees of OA during aging. Significant positive correlations were found between AC scores and synovitis, osteophyte formation, and ectopic chondrogenesis in control mice of both sexes. Significant direct correlations were found between AC scores and inflammatory markers in chondrocytes, including matrix metalloproteinase-13 (MMP-13), inducible nitric oxide synthase (iNOS), and the NLR family pyrin domain containing-3 (NLRP3) inflammasome in both sexes, indicating a link between OA severity and inflammation. Additionally, markers of cell cycle arrest, such as p16 and β-galactosidase, also correlated with AC scores. Using micro-CT, we examined the correlations between subchondral bone (SCB) morphology traits and AC scores. In male mice, no significant correlations were found between SCB volume and AC scores, while in female control mice, significant correlations were observed between AC scores and SCB volume in the femur.

Finally, we explored the effects of methylene blue (MB) and mitoquinone (MitoQ), two agents that affect mitochondrial function, on the prevalence and progression of OA during aging. Notably, MB and MitoQ treatments influenced the disease's progression in a sex-specific manner. In males, MB treatment exacerbated OA, while MitoQ showed mixed results. In contrast, both treatments in females increased the proportion of mice without OA, suggesting a potential slowing of OA progression.

In conclusion, our study provides comprehensive insights into the prevalence and progression of primary OA in aged UM-HET3 mice, highlighting the sex-specific effects of MB and MitoQ treatments. The correlations between AC scores and various pathological factors underscore the multifaceted nature of OA and its association with inflammation and subchondral bone changes.

Introduction:

Primary osteoarthritis (OA) refers to the development of OA without any known underlying factors, conditions, or injuries. Aging is a significant risk factor for OA, and during the aging process, there are progressive changes in joint tissues, including articular cartilage, subchondral bone, synovium, and ligaments. These changes involve increased matrix degradation, altered cell metabolism, impaired tissue repair mechanisms [1], and chronic low-grade inflammation [2-5]. In knee OA, which is commonly associated with aging, the articular cartilage gradually deteriorates, becoming thinner and exhibiting structural abnormalities. This compromises its ability to absorb shock and distribute load, resulting in joint pain and dysfunction. The subchondral bone also undergoes remodeling, including thickening and sclerosis, contributing to joint stiffness and further impacting cartilage health. Inflammation in the synovial membrane increases, leading to the production of inflammatory mediators that contribute to cartilage degradation and joint inflammation [6].

Our understanding of primary OA is derived from both clinical and preclinical research. In vivo preclinical animal models that focus mainly on genetic- or injury-induced OA have been particularly useful for two key purposes: investigating the underlying mechanisms of OA and evaluating the effectiveness of various treatment options. However, clinical studies of primary OA have shown variability in the actual onset of the disease, in the rate of OA progression, and in the severity of OA in different individuals. Thus, while inbred mouse models have provided valuable insights into OA mechanisms, they lack the genetic complexity of the human population [7, 8], and provide only limited insights into the natural development of the disease. Studying OA using a single genome compromises generalization and translation of findings [9]. The UM-HET3 mouse model, generated through a specific crossbreeding strategy (**Figure 1A**), offers a genetically heterogeneous population that better represents genetic diversity. The *UM-HET3 mouse model was selected by the Interventional Testing Program (ITP) at the National Institute of Aging (NIA) for studies of anti-aging drugs and was used in several studies to map genes controlling skeletal morphology* [10-13] (https://phenome.jax.org/projects/ITP1).

Since OA occurrence intensifies with age, evaluating the presence and severity of OA becomes a crucial factor in assessing the effectiveness of compounds designed to prolong life by influencing pathways that might also play a role in OA development. Therefore, this study aimed to assess the severity of OA associated with aging in genetically diverse male and female mice and to investigate whether the severity of this age-related OA was altered in mice treated with specific agents from the ITP study. Specifically, since common cellular processes are shared between OA and aging, such as mitochondrial dysfunction, oxidative stress, increased inflammation, and accumulation of senescent cells [14], we focused on two compounds that enhance mitochondrial function. Mitochondrial dysfunction and imbalanced production of reactive oxygen species (ROS) are considered contributing factors to OA development [15-17]. Impaired mitochondrial function in chondrocytes is linked to various cellular mechanisms involved in energy regulation, inflammation, and pro-catabolic responses.

Two compounds that affect mitochondrial function were tested by the ITP centers, methylene blue (MB) and mitoquinone (MitoQ). MB, an FDA-approved drug used to treat methemoglobinemia, has been shown to enhance mitochondrial function and has demonstrated beneficial effects in animal models of OA [18]. Intra-articular injection of MB in rat and rabbit OA models improved joint symptoms, protected cartilage, and reduced inflammation [19] [20]. MitoQ, a mitochondria-targeted antioxidant [21-23], also suppressed ROS production and ATP synthesis in an osteochondral explant model [24] .Here we report

the prevalence of primary OA and the effects of lifelong mitochondrial function enhancement on knee joint tissues during aging in the UM-HET3 mice [25].

Methods:

Animals:

UM-HET3 male and female mice were produced by a cross between (BALB/cByJ × C57BL/6J)F1 mothers (JAX stock #100009) and (C3H/HeJ × DBA/2J)F1 fathers (JAX stock #100004). Detailed housing conditions were specified elsewhere [26]. All experiments were approved by the Institutional Animal Care and Use Committee (IACUC) of each site.

Interventions: MB (28 ppm, starting at 4 months of age) or MitoQ (100 ppm, starting at 7 months of age) were administered in the diet. MB and MitoQ were formulated into irradiated Purina TestDiet 5LG6 diet, and supplied to all three sites such that all sites used the same batches of food.

Micro computed tomography (micro-CT):

Micro-CT of the knee joints was done in accordance with the American Society for Bone and Mineral Research (ASBMR) guidelines [27]. Intact femur and tibia including the knee joint were scanned using a high-resolution SkyScan micro-CT system (SkyScan 1172, Kontich, Belgium) containing 10-M digital detector set at a 10W energy level (100kV and 100 µA), with a 0.5 mm aluminum filter with a 9.7µm image voxel size. Subchondral bone (SCB) parameters were taken in the distal femur below the AC avoiding cortical bone and included bone volume fraction (bone volume/tissue volume, (BV/TV %), trabecular thickness (Tb.Th, mm), trabecular number (Tb.N, 1/mm), and bone mineral density (BMD, g/cc). Subchondral bone plate (SCBP) thickness and BMD were measured at the proximal tibia. Data reconstruction was done using NRecon software, and data analysis using CTAn software. 3D images and videos were obtained using CT Vox software (Version 3.3.1 64-bit).

Histology:

Following micro-CT scanning, knee joints were decalcified in 10% EDTA for 3-4 weeks, dehydrated using graded alcohol series and xylene, and processed for paraffin embedding and sectioning. Safranin-O-red staining was used for scoring the AC of the knee joint, including the presence and maturation stage of osteophytes as well as the ectopic chodrogenesis and ossification. H&E staining was used to score inflammation of synovial membrane, and overall morphology.

Osteoarthritis score: Cartilage damage at medial and lateral tibio-femoral joints was evaluated by two blinded observers using the Osteoarthritis Research Society International (OARSI) scoring system. Loss of cartilage proteoglycan scored by safranin-O-red; normal staining of non-calcified cartilage, scored 0; Loss of safranin-O-red staining without structural changes was scored 0.5; small fibrillations without loss of cartilage was scored 1; Vertical clefts down to the layer immediately below the superficial layer and some loss of surface lamina was scored 2; Vertical clefts/erosion to the calcified cartilage extending to <25% of the articular surface was scored 3; Vertical clefts/erosion to the calcified cartilage extending to 25–50% of the articular surface was scored 4; Vertical clefts/erosion to the calcified cartilage extending to 50–75% of the articular surface was scored 5; Vertical clefts/erosion to the calcified cartilage extending >75% of the articular surface was scored 6. Osteophyte maturity was scored 0 where no osteophytes were detected; 1 when osteophytes were composed of pre-cartilaginous lesion; 2 when osteophytes were composed predominantly of cartilage; 3 when osteophytes were composed of mixed cartilage and bone; and 4 when osteophytes were composed predominantly of bone. Synovitis was scored from H&E-stained sections. Briefly, the thickness of the synovial cell lining layer and the cell density within this layer was scored from 0 to 3, with 0 being the thinnest synovial cell lining layer and low cell density (1-2 cell layers), 1 with 3-5 cell layers, 2 with 6-8 cell layers, and 3 being the thickest synovial cell lining layer (>8

layers) and high cell density. <u>Ectopic chondrogenesis</u> was scored 0 when it was undetectable, 1 when it was found in the synovium and or the capsule, and 2 when it extended into the surrounding ligament and or musculature.

Immunohistochemistry:

Following deparaffinization and rehydration, the tissue sections were treated with citrate-based antigen unmasking solution (H-3300, Vector Laboratories, Inc, California, USA). Then endogenous peroxidase activity was inactivated by BLOXALL endogenous blocking solution (SP-6000, Vector Laboratories, Inc, California, USA). After blocking using the protein block solution (ab64226, Abcam, Massachusetts, USA), the sections were incubated overnight with the primary antibodies at 4^oC over-night, stained by rabbit specific HRP/DAB(ABC) detection kit (PK-4001/SK-4100, Vector Laboratories, Inc, California, USA) and counterstained by hematoxylin (Sigma, Missouri, USA)). The images were acquired by DMRXE universal microscope with objective imaging gigapixel montaging workstation (Leica Biosystems, IL, USA) and analyzed by Fiji Image J (Version 1.5r; NIH, Maryland, USA). For IHC, primary antibodies were against iNOS (1:300, PA3030A, Invitrogen, Massachusetts, USA), MMP-13 (1:50, 18165-1-AP, Proteintech, Illinois, USA), NLRP3 (1:300, PA5-88709, Invitrogen, Massachusetts, USA), β-galactosidase (1:400; #ab196838, Abcam), and p16 (1:100; #PA30670, Invitrogen).

Statistical analysis:

Individual descriptive statistics (e.g., means, medians, interquartile rangers) were calculated overall and by group for each outcome. Spearman's rank correlation coefficients were computed for each pairwise combination of categorical variables overall and by treatment and sex, with adjustment for the family wise error rate using the method of Benjamini and Hochberg [28]. Principal component analyses were then conducted for continuous multivariate outcomes by treatment and sex; any instance of missing data was imputed using the regularized iterative PCA algorithm [29]. Data analysis was conducted in R v4.1.3

Results:

Prevalence of primary OA in UM-HET3 mice: Knee joints of 22-25 months old mice were dissected and processed for safranin-O-red and H&E staining. We used The Osteoarthritis Research Society International (OARSI) scoring system [30-32] to assess the integrity of the AC at the distal femur and proximal tibia (**Figure 1B**) and the state of the synovial membrane at the medial and lateral sides of the joint (**Figure 1C**). The scores of control male and female mice varied both between and within sexes. The relative frequency of the cumulative AC scores at the *medial and lateral sides* of *both the tibia and the femur* (that can reach a maximal score of 24) revealed that 90% of control male mice and 98% of control female mice developed primary OA to varying degrees (**Figure 2A,B**). Interventions with MB or MitoQ increased the number of males with OA resulting in 100% of the male mice with various degrees of OA (**Figure 2A**). In females, however, treatment with MB or MitoQ increased the number of OA-free females to 4% and those with cumulative scores of $1 \leq X \leq 6$ to 87% as compared to 75% of control females with the same score range (**Figure 2B**).

The relative frequency of the cumulative scores of the AC specifically at the *medial side* of the tibia and the femur (that can reach a maximal score of 12) revealed that 85% of control male mice and 90% of control female mice developed primary OA to varying degrees (**Figure 2C,D**). Interventions with MB or MitoQ altered the natural history of the disease in a sex-specific manner. In male mice, treatment with MB worsened the outcomes, as all MB-treated males developed OA. Among them, 60% showed cumulative scores of $1 \leq X \leq 3$, 10% showed cumulative scores of $4 \leq X \leq 6$, and 34% showed cumulative scores of $7 \leq X \leq 12$ AC scores at the *medial side* of the tibia and the femur (**Figure 2C**). On the other hand, 9% of MitoQ-treated males showed no OA with age, 46% showed cumulative scores of $1 \leq X \leq 3$, while 41% showed a cumulative score of $7 \leq X \leq 12$ (**Figure 2D**). In female mice, 11% of control mice showed no OA with age. Treatment with MB or MitoQ increased that ratio to 25% and 23%, respectively. However, treatment with MB or MitoQ did not significantly affect the proportion of mice scored $1 \leq X \leq 3$ in female mice. Approximately 64% of control females, 67% of MB-treated females, and 69% of MitoQ-treated females exhibited an accumulative score of $4 \leq X \leq 6$ OA.

The relative frequency of the cumulative scores of the AC specifically at the *lateral side* of the tibia and the femur (that can reach a maximal score of 12) revealed that 80% of control male mice and 96% of control female mice developed primary OA to varying degrees (**Figure 2E,F**). Approximately 66% of control males, 86% of MB-treated males, and 82% of MitoQ-treated males exhibited an accumulative score of $1 \leq X \leq 3$ OA in the lateral side. In females, treatment with MitoQ associated with 14% of cumulative AC score of 0 and 87% of cumulative AC score of $1 \leq X \leq 3$. Overall, treatment with MB or MitoQ did not affect or even worsened the natural history of primary OA during aging, while in females, treatment with MB or MitoQ associated with increased fraction of mice with cumulative AC score of 0, suggesting that the progression of OA slowed down with these interventions.

AC scores correlate with osteophytosis, synovitis and ectopic chondrogenesis: We investigated the relationships between AC scores at the medial part of the knee joint and two other factors: medial osteophytosis and synovitis (**Figure 3**). In both control male and female mice, we observed significant positive correlations between AC scores and synovitis (**Figure 3A,B**). Treatment with MB or MitoQ did not alter these correlations in male mice. However, in females treated with MB, but not MitoQ, we did not find any correlations between the severity of AC scores and synovitis.

We used a semiquantitative method to score the degree of osteophytes present in the medial and lateral sides of the knee joint. In male mice, osteophytosis did not correlate with AC scores. However, treatment with MB or MitoQ resulted in increased osteophytosis with higher AC scores (**Figure 3C**). In untreated and MitoQ-treated females, osteophytosis showed an increase with higher AC scores (**Figure 3D**). Lastly, ectopic chondrogenesis exhibited significant correlations with AC scores in both untreated and MB-treated male and female mice. However, there were no correlations between ectopic chondrogenesis and AC scores in either sex for mice treated with MitoQ (**Figure 3E,F**).

AC scores correlate with markers of inflammation in chondrocyte: We used immunohistochemistry (IHC) approach to determine the levels of inflammatory markers in chondrocytes of the AC. Three inflammatory markers were selected including MMP-13, iNOS, and the NLRP3 inflammasome. We found significant direct correlations between the AC scores and the protein levels of MMP-13, iNOS, and NLRP3 in chondrocytes at the medial part of the knee joint in both male and female mice (**Figure 4A, supplement figure 1**). Similar correlations were found in the lateral part of the knee joint, although, not all were significant (**Figure 4B**). As expected, synovitis score correlated with the protein levels of iNOS, and NLRP3 at the synovial membrane at the medial part of the joint (**Figure 4C, supplement figure 2**). Finally, we found significant direct correlations between AC scores and the protein levels of markers of cell cycle arrest including p16 and β-galactosidase in chondrocytes (**supplement figure 3**).

SCB morphology in aged UM-HET3 mice was not modified by MB or MitoQ treatment: We examined the morphology of the subchondral bone (SCB) and subchondral plate (SCBP) using micro-CT analysis (**Table 1, Figure 5A-I, supplement figure 4**). In male mice, we did not observe any significant correlations between SCB volume (bone volume/total volume, BV/TV) and AC scores in the femur or tibia, regardless of whether they were untreated mice or mice treated with MB or MitoQ. However, in untreated female mice, we found significant correlations between AC scores and SCB BV/TV in the femur but not in the tibia (**Table 2**). Furthermore, we did not find any significant correlations between SCBP thickness and mineral density with AC scores. Results from principal component analyses (PCA) suggest that much of the variation in micro-CT measures across samples occurs in two dimensions (70%) (**Figure 5J,K**). In both males and females, the strongest influencing measurements included femur SCB BV/TV, femur SCB BMD, and tibia SCB BV/TV for the first component, while femur SCB Tb.Th had substantial influence on the second component. There was a strong positive correlation between femur and tibial measurements with the exception of strong negative correlations with both tibia and femur SCB Tb.Sp. We found that morphological traits of the SCB in the femur correlated significantly to those of the tibia, indicating that both SCB compartments are regulated in a similar manner (**supplement figure 5**). Surprisingly, PCA indicate that measurements for the tibia SCBP were largely orthogonal. In males, contributions from tibia SCB Tb.Th and tibia SCBP BMD were not as strong, nor were tibia SCB Tb.Th and tibia SCB Tb.Sp in females. Additionally, we did not find significant interactions between SCBP morphology and the cumulative AC scores at the lateral or medial of the tibia (**supplement figure 6**).

Discussion:

OA, a widespread and chronic joint disorder, is marked by ongoing deterioration of cartilage, alterations in subchondral bone, bone marrow lesions, damage to the meniscus, and synovitis. In the current study we utilized 182 stifles of genetically diverse UM-HET3 mouse to model the prevalence and severity of primary OA during aging in male and female mice. We found that at the medial side of the knee joint 90% of female and 85% of male mice developed primary OA to varying degrees, accompanied by synovitis, osteophytosis, and calcified menisci. In females, 64% showed cumulative OARSI scores of $1 \leq X \leq 3$, while males showed 43% of scores at that range at the *medial side* of the tibia and the femur.

Both male and female mice showed significant positive correlations between AC scores and synovitis. In female mice we found significant correlation between osteophytosis and AC scores. Notably, osteophytes contribute to both the functional properties of affected joints and clinically relevant symptoms. They are closely associated with cartilage damage, although they can also develop without explicit cartilage damage [33]. The precursor cells for osteophyte formation are debatable and include mesenchymal stem cells in the periosteum [34] or synovium-derived cells [35]. The TGFβ superfamily of growth factors and macrophages play important roles in osteophyte induction [36]. In males, osteophytosis did not correlate with AC scores, indicating that the progression of these two processes is independent in males.

The progression of osteoarthritis, triggers inflammatory reaction, characterized by increased expression of the inflammatory markers in AC chondrocytes and cells of the synovial membrane. Specifically, we report significant correlations between protein levels of MMP13, iNOS, and NLRP3 in AC chondrocytes and synovial membrane at the medial knee joint in both male and female mice. Release of inflammatory factors interfere with the anabolic and catabolic activities of chondrocytes and may lead to cell death [37, 38].

Previous studies have reported that mitochondrial dysfunction and oxidative stress contribute to OA development [39]. In chondrocytes, the production of mitochondrial ROS has been linked to a marked increase in cyclooxygenase, a powerful catabolic inflammatory agent [40] and MMP-13 [41], highlighting their potential role in degradation of the AC. Earlier research has established the significant role of oxidants in modulating metabolic processes in cartilage [42-44]. Together, it is expected that inhibiting excessive ROS accumulation can protect against chondrocyte damages and slow down OA progression. Further, mitochondria in chondrocytes are mechanically connected to the cell membrane through f-actin, facilitating their movement and distribution in the cytosol [45]. This connection between mitochondria and the cytoskeleton plays a crucial role in managing energy and ROS production [43], as well as maintaining calcium balance [46] and controlling mitochondrial apoptotic factors [47]. In light of these evidence, we studied the effects of lifelong treatment with MB or MitoQ on the natural history of primary OA in UM-HET3 mice.

MB and MitoQ affect oxidative stress in tissues of the knee joint via different mechanisms. MB ameliorates oxidative stress via stimulation of Nrf2 and its translocation to the nucleus in both in chondrocytes and synoviocytes and exerts anti-inflammatory effects in the synovium and neurons that innervate the knee joint [19]. Evidence from the current study indicate that lifelong treatment with MB does not alter the natural history of primary OA in UM-HET3 male mice. In females, however, MB increased the relative fraction of mice with cumulative AC score of 0. MitoQ, on the other hand, functions as a mitochondrial antioxidant [48]. Its lipophilic moiety allows MitoQ to concentrate within mitochondria

up to a thousand times more than other general antioxidants [49]. MitoQ is known for its protective effects against diseases related to oxidative damage. In a study using a model of bovine cartilage explants subjected to mechanical stress, MitoQ significantly reduced the production of ROS [24]. Furthermore, MitoQ treatment was found to mitigate the breakdown of the extracellular matrix (ECM), reduce oxidative stress, and lessen the inflammatory response in chondrocytes. This was observed in a mouse model of OA induced by destabilizing the medial meniscus (DMM) [50] and was likely due to its activation of the NRF2 pathway and the promotion of mitophagy in chondrocytes. As seen with MB, MitoQ altered OA progression only in UM-HET3 female but not male mice.

In this study, we report on the prevalence and severity of primary OA in the genetically diverse UM-HET3 mice. We also examined how treating these mice with the antioxidants MB and MitoQ from early adulthood into old age impacts the development of primary OA. Our research has certain limitations, including the fact that we only examined mice at one age point (22-25 months). We established the validity of our model by showing correlations between OA features and indicators of inflammation, synovitis, and osteophyte formation. However, our study did not explore the direct causes of these conditions. We also recognize that our research does not delve into the cellular or molecular mechanisms that might trigger or advance OA. Nevertheless, our findings demonstrate that primary OA in UM-HET3 mice reflects the prevalence and severity seen in human primary OA. This suggests that these mice are a suitable model for testing systemic treatments that could influence aging processes, particularly in the context of developing primary OA.

Author contribution:
Conceptualization: SY
Funding acquisition: SY
Formal analyses: RRR, SY
Investigation: SBP, GY
Statistical analyses: RRR
Resources: RAM, DEH, RS
Writing review and editing: SY, TK

Acknowledgements:
Financial support received from the National Institutes of Health Grant R01AG056397 and B01 (2024) Department of Molecular Pathobiology Accelerator Award to SY. SBP is supported by New York University Provost's Postdoctoral Fellowship Program. This work was also supported by the National Institutes of Health grant to The Jackson Laboratory Nathan Shock Center of Excellence in the Basic Biology of Aging AG038070, U01-AG022303 to RAM, UO1-AG022308 to DEH, U01-AG013319 to RLS, RLS is supported by a Senior Research Career Scientist Award from the Department of Veterans Affairs Office of Research and Development, and S10 OD010751-01A1 for micro-computed tomography.

Co-authors details:
- Sher Bahadur Poudel, Email: sbp4@nyu.edu
Affiliation: David B. Kriser Dental Center, Department of Molecular Pathobiology, New York University College of Dentistry, New York, NY
- Ryan R Raff, Email: rrr8@nyu.edu
[2]David B. Kriser Dental Center, Department of Epidemiology and Health Promotion, New York University College of Dentistry New York, NY 10010-4086
- Gozde Yildirim, Email: yildirimgina@gmail.com
Affiliation: David B. Kriser Dental Center, Department of Molecular Pathobiology, New York University College of Dentistry, New York, NY
- Richard A Miller, Email: millerr@umich.edu
Pathology and Geriatrics Center, University of Michigan, Ann Arbor, MI, USA.
- David E Harrison, Email: david.harrison@jax.org
The Jackson Laboratory, Bar Harbor, ME, USA.
- Randy Strong, Email: STRONG@uthscsa.edu
Geriatric Research, Education and Clinical Center and Research Service, South Texas Veterans Health Care System, San Antonio, TX, USA.
Barshop Institute for Longevity and Aging Studies and Department of Pharmacology, The University of Texas Health Science Center, San Antonio, TX, USA.
- Thorsten Kirsch, Email: Thorsten.Kirsch@nyulangone.org
Affiliation: Department of Orthopaedic Surgery, NYU Grossman School of Medicine, and Department of Biomedical Engineering, NYU Tandon School of Engineering, New York, NY
- Shoshana Yakar, Email: sy1007@nyu.edu
Affiliation: David B. Kriser Dental Center, Department of Molecular Pathobiology, New York University College of Dentistry, New York, NY

Figure Legends:

Figure 1: **OARSI scoring of stifles from UMHET3 mouse model**. (A) UM-HET3 is a genetically diverse mouse model, generated via 4way cross of (BALB/cByJ × C57BL/6J) F1 mothers and (C3H/HeJ × DBA/2J) F1 fathers. Each of the offspring is genetically unique but shares 50% of its genetic material with every other UM-HET3 mouse. (B) Shown are representative knee joint samples that were processed for safranin-O-red, or (C) H&E staining of the synovial membrane that were categorized based on the OARSI scoring system [30-32].

Figure 2: **Prevalence and severity of OA in UM-HET3 mice during aging**. The distribution of cumulative AC scores at the medial (A,B) and lateral (C,D) sides of the knee joints from control (CTL males n=47, CTL females n=45), MB- (males n=22, females n=24), and MitoQ-treated (males n=22, females n=22) mice.

Figure 3: **Correlations between cumulative AC scores and osteophytosis, synovitis or ectopic chondrogenesis**. (A,B) Cumulative AC scores at the medial side of the joint show positive correlation with synovitis in both control male and female mice. Treatment with MB alter these association in female mice. (C,D) Cumulative AC scores at the medial side of the joint do not show significant correlation with osteophyte formation in control male. However, treatment with MB or MitoQ result in positive correlation between AC scores and osteophyte formation. In female mice, cumulative AC scores at the medial side of the joint show positive correlation with osteophyte formation in control and MitoQ-treated mice. (E,F) Cumulative AC scores at the medial side of the joint show positive correlation with ectopic chondrogenesis in both control and MB-treated male and female mice, but not with MitoQ-treated mice. CTL males n=47, CTL females n=45, MB males n=22, MB females n=24, MitoQ males n=22, MitoQ females n=22.

Figure 4: **Correlations between cumulative AC scores and markers of inflammation**. Correlations between cumulative AC scores at the medial (A) and lateral (B) side of the joint with the levels of MMP13 (males n=21, females n=15), iNOS (males n=19, females n=19), and NLRP3 (males n=17, females n=15) in AC chondrocytes. (C) Sections of the medial synovial membrane in control mice immunostained with iNOS (males n=19, females n=19), and NLRP3 (males n=17, females n=15).

Figure 5: **Subchondral bone (SCB) and subchondral plate (SCP) morphology analyzed by micro-CT**. (A) 2D image of a micro-CT scan indicating the knee joint regions of interest that were analyzed including the SCB of the distal femur, SCB of the proximal tibia, and the SCB plate of the proximal tibia. Parameters assessed by micro-CT included femur (B) bone volume/total volume (BV/TV), (C) trabecular thickness (Tb.Th), and (D) SCB mineral density (BMD). Likewise, tibia BV/TV (E), Tb.Th of the tibia SCB (F), and tibia SCB mineral density (G). SCB plate (SCBp) thickness (H), and SCBp bone mineral density (I), were determined at the proximal tibia. Data presented as mean ± SEM, p values are indicated. CTL males n=44, CTL females n=44, MB males n=22, MB females n=24, MitoQ males n=22, MitoQ females n=22. SCB phenotypic measurements from PCA of both male (J) and female (K) mice. Positively correlated variables point to the same side of the plot; negatively correlated variables point to opposite sides of the plot. Length and color of arrows indicate contribution of the variable to the overall variance of the data. Arrows that are close together are strongly positively correlated, arrows that are about 90 degrees (orthogonal) to each other are uncorrelated, and those far away (such as 180 degrees) are strongly negatively correlated.

Supplement Figure Legends:

Supplement Figure 1: Shown are representative knee joint sections stained with MMP-13, iNOS, and NLRP3 antibodies in both male and female mice. Quantification is available in figure 4.

Supplement Figure 2: Shown are representative medial synovial membrane sections stained with iNOS and NLRP3 in both male and female mice. Quantification is available in figure 4.

Supplement Figure 3: Shown are representative knee joint sections stained with p16 (A) and quantification of positive chondrocytes in the (B) medial and (C) lateral side of the tibia. (D) Representative knee joint sections stained with β-Gal and quantification of positive chondrocytes in the (E) medial and (F) lateral side of the tibia in both male and female mice. Males n=19 and females n=19

Supplement Figure 4: (A) 3D images of a micro-CT scan of knee joints from male and female mice with different OA severities (cumulative scores at the medial side of the joint). A link to a short movie presenting 3D reconstruction of knee joints with no histological evidence of OA or with high cumulative AC score obtained by histology (https://data.mendeley.com/datasets/6nddwstfw3/1).

Supplement Figure 5: Femur and tibia SCB traits obtained by micro-CT are significantly directly correlated. Spearman's rank correlations between SCB BV/TV (A,B), SCB BMD (C,D), SCB Th. (E,F), and SCB Tb.N (G,H) of femur and tibia in male and female mice. Males n=47, Females n=45. P values are indicated for each correlation.

Supplement Figure 6: Cumulative AC scores of the medial or lateral tibia show no correlation to the morphology of the SCBP by micro-CT. Spearman's rank correlations between cumulative AC scores at the medial tibia with SCBP Th (A,B) or SCBP BMD (C,D) in male and female mice. Similar correlations were done between cumulative AC scores at the lateral tibia with SCBP Th (E,F) or SCBP BMD (G,H) in male and female mice. Males n=47, Females n=45. P values are indicated for each correlation.

Tables:

Table 1: SCB morphology of male and female mice determined by micro-CT. Bone volume/total volume (BV/TV), trabecular thickness (Tb.Th), and bone mineral density (BMD).

	M CTL (n=47)	M MB (n=22)	M MitoQ (n=22)	F CTL (n=45)	F MB (n=24)	F MitoQ (n=22)
Distal Femur Subchondral Bone						
BV/TV, %	21.7618±0.7489	20.9130±1.0306	21.8811±1.5200	17.6319±0.6837	17.5472±0.7915	17.7906±0.9468
Tb.Th, mm	0.0688±0.0012	0.0678±0.0020	0.0687±0.0020	0.0749±0.0016	0.0759±0.0022	0.0789±0.0023
Tb.Sp, mm	0.2532±0.0057	0.2539±0.0065	0.2512±0.0109	0.3296±0.0065	0.3243±0.0061	0.3336±0.0096
Tb.N, 1/mm	3.1403±0.0790	3.0609±0.0990	3.1812±0.1763	2.3486±0.0742	2.3010±0.0653	2.2506±0.1004
BMD, g/cc	0.2480±0.0072	0.2344±0.0090	0.2530±0.0137	0.2208±0.0072	0.2114±0.0087	0.2307±0.0097
Proximal Tibia Subchondral Bone						
BV/TV, %	20.3482±0.6811	20.6231±1.2275	18.6409±1.0443	17.0072±0.7781	16.7714±1.0041	15.9309±0.7885
Tb.Th, mm	0.0651±0.0010	0.0671±0.0015	0.0642±0.0013	0.0724±0.0014	0.0705±0.0018	0.0709±0.0018
Tb.Sp, mm	0.2135±0.0030	0.2180±0.0063	0.2314±0.0057	0.2601±0.0038	0.2558±0.0063	0.2674±0.0056
Tb.N, 1/mm	3.1047±0.0739	3.0392±0.1355	2.8963±0.1369	2.3249±0.0827	2.3618±0.1231	2.2370±0.0891
BMD, g/cc	0.2389±0.0070	0.2365±0.0105	0.2302±0.0103	0.2175±0.0075	0.2080±0.0105	0.2085±0.0082
Tibia Subchondral Plate						
SCB plate Th, mm	0.1080±0.0036	0.0918±0.0055	0.1027±0.0044	0.1009±0.0032	0.0890±0.0061	0.1059±0.0044
SCB plate BMD, g/cc	0.8638±0.0147	0.8529±0.0168	0.9381±0.0058	0.8914±0.0167	0.8690±0.0190	0.9620±0.0085

Table 2: P values of correlations between cumulative scores of the AC, synovitis, osteophytosis, ectopic chondrogenesis (Ec.Chon), and parameters of subchondral (SC) bone obtained by micro-CT. Bone volume/total volume (BV/TV), trabecular thickness (Tb.Th), and bone mineral density (BMD).

Comparisons			CTL F	MB F	MitoQ F	CTL M	MB M	MitoQ M
AC cumulative score (femur and tibia, medial and lateral)	vs	Osteophyte cumulative score	0.0001	>0.1	0.0119	0.0808	0.0162	0.0221
AC cumulative score (femur and tibia, medial and lateral)	vs	Synovitis cumulative score	0.0036	0.0975	>0.1	0.0004	0.0045	0.0004
AC cumulative score (femur and tibia, medial and lateral)	vs	Ec.Chond cumulative score	0.0031	0.0113	>0.1	0.0077	0.0031	0.0976
AC cumulative score (femur and tibia, medial and lateral)	vs	Femur SCB BV/TV	0.0224	0.0656	>0.1	>0.1	>0.1	>0.1
AC cumulative score (femur and tibia, medial and lateral)	vs	Femur SCB BMD	0.0088	0.0221	>0.1	>0.1	>0.1	>0.1
AC cumulative score (femur and tibia, medial and lateral)	vs	Tibia SCB BV/TV	>0.1	0.0153	>0.1	>0.1	>0.1	>0.1
AC cumulative score (femur and tibia, medial and lateral)	vs	Tibia SCB BMD	>0.1	0.0082	>0.1	>0.1	>0.1	>0.1
AC medial cumulative score (medial femur and tibia)	vs	Osteophyte medial cumulative scores	0.0004	>0.1	0.0031	>0.1	0.0077	0.0385
AC medial cumulative scores (medial femur and tibia)	vs	synovitis Medial	0.0077	>0.1	0.0167	0.0001	0.0153	0.0010
AC medial cumulative scores (medial femur and tibia)	vs	synovitis Lateral	0.06318	>0.1	>0.1	0.08928	>0.1	>0.1
AC cumulative score (femur and tibia, medial and lateral)	vs	Tibia SC plate Th mm	>0.1	>0.1	>0.1	>0.1	>0.1	>0.1
AC cumulative score (femur and tibia, medial and lateral)	vs	Tibia SC plate BMD	0.0487	>0.1	>0.1	>0.1	>0.1	>0.1

References:

1. Lotz, M. and R.F. Loeser, *Effects of aging on articular cartilage homeostasis.* Bone, 2012. **51**(2): p. 241-8.
2. Mobasheri, A., et al., *Glucose transport and metabolism in chondrocytes: a key to understanding chondrogenesis, skeletal development and cartilage degradation in osteoarthritis.* Histol Histopathol, 2002. **17**(4): p. 1239-67.
3. Salvioli, S., et al., *Inflamm-aging, cytokines and aging: state of the art, new hypotheses on the role of mitochondria and new perspectives from systems biology.* Curr Pharm Des, 2006. **12**(24): p. 3161-71.
4. Maggio, M., et al., *Interleukin-6 in aging and chronic disease: a magnificent pathway.* J Gerontol A Biol Sci Med Sci, 2006. **61**(6): p. 575-84.
5. De Martinis, M., et al., *Inflamm-ageing and lifelong antigenic load as major determinants of ageing rate and longevity.* FEBS Lett, 2005. **579**(10): p. 2035-9.
6. Greene, M.A. and R.F. Loeser, *Aging-related inflammation in osteoarthritis.* Osteoarthritis Cartilage, 2015. **23**(11): p. 1966-71.
7. Ashbrook, D.G., et al., *A platform for experimental precision medicine: The extended BXD mouse family.* Cell Syst, 2021. **12**(3): p. 235-247 e9.
8. Saul, M.C., et al., *High-Diversity Mouse Populations for Complex Traits.* Trends Genet, 2019. **35**(7): p. 501-514.
9. Williams, R.W., *Herding cats: the sociology of data integration.* Front Neurosci, 2009. **3**(2): p. 154-6.
10. Miller, R.A., et al., *Preservation of femoral bone thickness in middle age predicts survival in genetically heterogeneous mice.* Aging Cell, 2011. **10**(3): p. 383-91.
11. Reeves, G.M., et al., *Quantitative trait loci modulate vertebral morphology and mechanical properties in a population of 18-month-old genetically heterogeneous mice.* Bone, 2007. **40**(2): p. 433-43.
12. Volkman, S.K., et al., *Quantitative trait loci that modulate femoral mechanical properties in a genetically heterogeneous mouse population.* J Bone Miner Res, 2004. **19**(9): p. 1497-505.
13. Volkman, S.K., et al., *Quantitative trait loci for femoral size and shape in a genetically heterogeneous mouse population.* J Bone Miner Res, 2003. **18**(8): p. 1497-505.
14. Price, J.S., et al., *The role of chondrocyte senescence in osteoarthritis.* Aging Cell, 2002. **1**(1): p. 57-65.
15. Hui, W., et al., *Oxidative changes and signalling pathways are pivotal in initiating age-related changes in articular cartilage.* Ann Rheum Dis, 2016. **75**(2): p. 449-58.
16. Loeser, R.F., *Aging and osteoarthritis.* Curr Opin Rheumatol, 2011. **23**(5): p. 492-6.
17. Blanco, F.J., I. Rego, and C. Ruiz-Romero, *The role of mitochondria in osteoarthritis.* Nat Rev Rheumatol, 2011. **7**(3): p. 161-9.
18. Zheng, J. and Q. Li, *Methylene blue regulates inflammatory response in osteoarthritis by noncoding long chain RNA CILinc02.* J Cell Biochem, 2019. **120**(3): p. 3331-3338.
19. Li, J.W., et al., *Methylene blue prevents osteoarthritis progression and relieves pain in rats via upregulation of Nrf2/PRDX1.* Acta Pharmacol Sin, 2022. **43**(2): p. 417-428.
20. Li, X., et al., *Methylene blue relieves the development of osteoarthritis by upregulating lncRNA MEG3.* Exp Ther Med, 2018. **15**(4): p. 3856-3864.
21. Tauskela, J.S., *MitoQ--a mitochondria-targeted antioxidant.* IDrugs, 2007. **10**(6): p. 399-412.
22. Sanjuan-Pla, A., et al., *A targeted antioxidant reveals the importance of mitochondrial reactive oxygen species in the hypoxic signaling of HIF-1alpha.* FEBS Lett, 2005. **579**(12): p. 2669-74.
23. Kelso, G.F., et al., *Selective targeting of a redox-active ubiquinone to mitochondria within cells: antioxidant and antiapoptotic properties.* J Biol Chem, 2001. **276**(7): p. 4588-96.

24.   Wolff, K.J., et al., *Mechanical stress and ATP synthesis are coupled by mitochondrial oxidants in articular cartilage.* J Orthop Res, 2013. **31**(2): p. 191-6.

25.   Miller, R.A., et al., *An Aging Interventions Testing Program: study design and interim report.* Aging Cell, 2007. **6**(4): p. 565-75.

26.   Strong, R., et al., *Nordihydroguaiaretic acid and aspirin increase lifespan of genetically heterogeneous male mice.* Aging Cell, 2008. **7**(5): p. 641-50.

27.   Bouxsein, M.L., et al., *Guidelines for assessment of bone microstructure in rodents using micro-computed tomography.* J Bone Miner Res, 2010. **25**(7): p. 1468-86.

28.   Yoav Benjamini, Y.H., *Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing.* Journal of the Royal Statistical Society: Series B (Methodological), 1995. **57**(1): p. 289-300.

29.   Josse, J. and F. Husson, *missMDA: A Package for Handling Missing Values in Multivariate Data Analysis.* Journal of Statistical Software, 2016. **70**(1).

30.   Glasson, S.S., et al., *The OARSI histopathology initiative - recommendations for histological assessments of osteoarthritis in the mouse.* Osteoarthritis Cartilage, 2010. **18 Suppl 3**: p. S17-23.

31.   Chambers, M.G., M.T. Bayliss, and R.M. Mason, *Chondrocyte cytokine and growth factor expression in murine osteoarthritis.* Osteoarthritis Cartilage, 1997. **5**(5): p. 301-8.

32.   Kaneko, H., et al., *Synovial perlecan is required for osteophyte formation in knee osteoarthritis.* Matrix Biol, 2013. **32**(3-4): p. 178-87.

33.   van der Kraan, P.M. and W.B. van den Berg, *Osteophytes: relevance and biology.* Osteoarthritis Cartilage, 2007. **15**(3): p. 237-44.

34.   Shirasawa, S., et al., *In vitro chondrogenesis of human synovium-derived mesenchymal stem cells: optimal condition and comparison with bone marrow-derived cells.* J Cell Biochem, 2006. **97**(1): p. 84-97.

35.   Sakaguchi, Y., et al., *Comparison of human stem cells derived from various mesenchymal tissues: superiority of synovium as a cell source.* Arthritis Rheum, 2005. **52**(8): p. 2521-9.

36.   van Lent, P.L., et al., *Crucial role of synovial lining macrophages in the promotion of transforming growth factor beta-mediated osteophyte formation.* Arthritis Rheum, 2004. **50**(1): p. 103-11.

37.   Wang, Q., et al., *Curcumin attenuates collagen-induced rat arthritis via anti-inflammatory and apoptotic effects.* Int Immunopharmacol, 2019. **72**: p. 292-300.

38.   Song, J., et al., *Dysregulation of the NUDT7-PGAM1 axis is responsible for chondrocyte death during osteoarthritis pathogenesis.* Nat Commun, 2018. **9**(1): p. 3427.

39.   Coryell, P.R., B.O. Diekman, and R.F. Loeser, *Mechanisms and therapeutic implications of cellular senescence in osteoarthritis.* Nat Rev Rheumatol, 2021. **17**(1): p. 47-57.

40.   Cillero-Pastor, B., et al., *Mitochondrial dysfunction activates cyclooxygenase 2 expression in cultured normal human chondrocytes.* Arthritis Rheum, 2008. **58**(8): p. 2409-19.

41.   Ahmad, R., et al., *Involvement of H-Ras and reactive oxygen species in proinflammatory cytokine-induced matrix metalloproteinase-13 expression in human articular chondrocytes.* Arch Biochem Biophys, 2011. **507**(2): p. 350-5.

42.   Martin, J.A., et al., *Mitochondrial electron transport and glycolysis are coupled in articular cartilage.* Osteoarthritis Cartilage, 2012. **20**(4): p. 323-9.

43.   Sauter, E., et al., *Cytoskeletal dissolution blocks oxidant release and cell death in injured cartilage.* J Orthop Res, 2012. **30**(4): p. 593-8.

44.   Lee, R.B. and J.P. Urban, *Functional replacement of oxygen by other oxidants in articular cartilage.* Arthritis Rheum, 2002. **46**(12): p. 3190-200.

45.   Boldogh, I.R., et al., *A protein complex containing Mdm10p, Mdm12p, and Mmm1p links mitochondrial membranes and DNA to the cytoskeleton-based segregation machinery.* Mol Biol Cell, 2003. **14**(11): p. 4618-27.

46.   Yi, M., D. Weaver, and G. Hajnoczky, *Control of mitochondrial motility and distribution by the calcium signal: a homeostatic circuit.* J Cell Biol, 2004. **167**(4): p. 661-72.

47.   Gourlay, C.W. and K.R. Ayscough, *The actin cytoskeleton: a key regulator of apoptosis and ageing?* Nat Rev Mol Cell Biol, 2005. **6**(7): p. 583-9.

48.   James, A.M., et al., *Interaction of the mitochondria-targeted antioxidant MitoQ with phospholipid bilayers and ubiquinone oxidoreductases.* J Biol Chem, 2007. **282**(20): p. 14708-18.

49.   Ng, L.F., et al., *The mitochondria-targeted antioxidant MitoQ extends lifespan and improves healthspan of a transgenic Caenorhabditis elegans model of Alzheimer disease.* Free Radic Biol Med, 2014. **71**: p. 390-401.

50.   Hou, L., et al., *Mitoquinone alleviates osteoarthritis progress by activating the NRF2-Parkin axis.* iScience, 2023. **26**(9): p. 107647.

Figure 1

Figure 2

Overall cumulative AC score (medial+lateral, femur+tibia)

Medial cumulative AC score (femur+tibia)

Lateral cumulative AC score (femur+tibia)

Figure 3

Figure 4

Figure 5