**Activating SRC/MAPK signaling via 5-HT1A receptor contributes to**

**the effect of vilazodone on improving thrombocytopenia**

**Ling Zhou** [1, †]**, Chengyang Ni** [1, 2†]**, Ruixue Liao** [1, 3†]**, Xiaoqin Tang** [1]**, Taian Yi** [4]**,**

**Mei Ran** [1,2]**, Miao Huang** [4]**, Rui Liao**[1]**, Xiaogang Zhou** [1]**, Dalian Qin** [1]**, Long**

**Wang** [1]**, Feihong Huang** [1]**, Xiang Xie** [5]**, Ying Wan** [2]**, Jiesi Luo** [2*]**, Yiwei Wang** [2*]**,**

**Jianming Wu** [2,1,3*]

[1] Sichuan Key Medical Laboratory of New Drug Discovery and Druggability, Luzhou Key

Laboratory of Activity Screening and Druggability Evaluation for Chinese Materia Medica,

School of Pharmacy, Southwest Medical University, Luzhou 646000, China

[2] School of Basic Medical Sciences, Southwest Medical University, Luzhou 646000, China

[3] Education Ministry Key Laboratory of Medical Electrophysiology, Southwest Medical

University, Luzhou 646000, China

[4] School of Pharmacy, Chengdu University of Traditional Chinese Medicine, Chengdu 611137,

China

[5] School of Basic Medical Sciences, Public Center of Experimental Technology, Model Animal

and Human Disease Research of Luzhou Key Laboratory, Southwest Medical University,

Luzhou, 646000, China

† These authors contributed equally to this work.

**\* Corresponding author:**

jianmingwu@swmu.edu.cn (Jianming Wu)

ljs@swmu.edu.cn (Jiesi Luo)

wangyiwei0102@swmu.edu.cn (Yiwei Wang)

**Abstract**

Thrombocytopenia caused by long-term radiotherapy and chemotherapy exists in

cancer treatment. Previous research demonstrates that 5-Hydroxtrayptamine (5-HT)

and its receptors induces the formation of megakaryocytes (MKs) and platelets.

However, the relationships between 5-HT1A receptor (5-HTR1A) and MKs is unclear

so far. We screened and investigated the mechanism of vilazodone as a 5-HTR1A partial agonist in promoting MK differentiation and evaluated its therapeutic effect in thrombocytopenia. We employed a drug screening model based on machine learning (ML) to screen the megakaryocytopoiesis activity of VLZ. The effects of Vilazodone (VLZ) on megakaryocytopoiesis were verified in HEL and Meg-01 cells. Tg (itga2b: eGFP) zebrafish was performed to analyze the alterations in thrombopoiesis. Moreover, we established a thrombocytopenia mice model to investigate VLZ administration accelerates platelet recovery and function. We carried out network pharmacology, Western blotting and immunofluorescence to demonstrate the potential targets and pathway of VLZ. VLZ has been predicted to have a potential biological action. Meanwhile, VLZ administration promotes MK differentiation and thrombopoiesis in cells and zebrafish models. Progressive experiments showed that VLZ has a potential therapeutic effect on radiation-induced thrombocytopenia *in vivo*. The network pharmacology and associated mechanism study indicated that SRC and MAPK signaling are both involved in the processes of megakaryopoiesis facilitated by VLZ. Furthermore, the expression of 5-HTR1A during megakaryocyte differentiation is closely related to the activation of SRC and MAPK. Our findings demonstrated that the expression of 5-HTR1A on MK, VLZ could bind to the 5-HTR1A receptor and further regulate the SRC/MAPK signaling pathway to facilitate megakaryocyte differentiation and platelet production, which provides new insights into the alternative therapeutic options for thrombocytopenia.

**Keywords:** *thrombocytopenia; vilazodone; 5-HT1A receptor; megakaryocyte*

**1 Introduction**

5-HT has been identified as a vasoactive neurotransmitter mostly located in the periphery of the body, and less than 1% of total body serotonin circulates in its free form in the blood (Erspamer and Asero, 1952). Peripheral synthesis by enteric chromaffin cells and release into the blood are taken up by circulating platelets via serotonin transporters (SERTs) and stored in dense particles (Spohn and Mawe, 2017). In the blood, approximately 98% of 5-HT resides in platelets, which are released into

circulation when activated. The biological effects of 5-HT are exerted through

activation of any of 15 distinct receptors in 7 different classes 5-HT1 receptor to

5-HT7 receptor (5-HT1R to 5-HT7R) (Barnes et al., 2021). It has been assumed for a

long time that 5-HT plays an important role in the hematopoietic process. 5-HT is a

powerful mitogen that has been shown to have mitogenic action during embryonic

development in mice (Matsuda et al., 2004). In particular, 5-HT is essential for the

survival of hematopoietic stem cells and progenitor cells (HSPCs) *in vivo* throughout

embryogenesis and increases the generation of HSPCs in vitro (Lv et al., 2017).

Consistently, MKs were the first hematopoietic cells examined in relation to the

serotonergic system and were megakaryocytes due to the well-known link between

5-HT and platelets. Studies have shown that the 5-HT2A, 5-HT2B, and 5-HT2C

subtypes exist on the surface of human primary megakaryocyte progenitor cells,

megakaryocytes, and some major megakaryocyte lines (Yang et al., 2007). In the

process of megakaryocyte hematopoietic regulation, 5-HT increases the number of

megakaryocyte progenitor cells, promotes megakaryocyte proliferation by activating

the 5-HT2 receptor (5-HT2R) and promotes the formation and maturation of mouse

CFU-MK and platelet generation in a dose-dependent manner (Ye et al., 2014).

Additionally, 5-HT might contribute to erythropoiesis (Amireault et al., 2011). Overall,

the serotonergic system is a therapeutic target that shows promise due to its extensive

capacity for regeneration and regulation of hematopoietic lineages. In addition to

5-HT2R, the 5-HT1 receptor (5-HT1R) is also important for cell proliferation and

differentiation, among which the 5-HT1A receptor (5-HTR1A) is one of the main

subtypes involved. The presence of 5-HT1F receptor (5-HTR1F) on CD34[+] cells

suggested that 5-HT1R signaling may be involved in neuronal regulation of immature

hematopoietic progenitors (Skurikhin et al., 2001). Moreover, 5-HT dose-dependently

induces the migration of neural crest cells in culture through 5-HTR1A in mice to

interfere with development (Moiseiwitsch and Lauder, 1995). Further evidence

revealed that 5-HTR1A could exert effects on the phosphorylation of ERK1/2 (Della

Rocca et al., 1999).

Vilazodone (VLZ) is a new dual-mechanism antidepressant approved by the United

States Food and Drug Administration (FDA) for treating major depressive disorder (MDD). VLZ displays the property of an SSRI and also has partial activation of 5-HTR1A (McCormack, 2015). It selectively inhibits serotonin reuptake with a 50 % inhibitory concentration (IC50) of 1.6 nmol/L and binds selectively to human 5-HT1A receptors with high affinity (IC50 of 2.1 nmol/L) (Wang et al., 2015). The antidepressant effect of VLZ is through inhibiting the serotonin transporter's reuptake of 5-HT, while part of the stimulant effect of 5-HTR1A is directed at the auto receptor mediated negative neuronal feedback mechanism. Previous studies have shown that the 5-HTR1A partial activation of VLZ has a faster and larger effect on presynaptic 5-HTR1A (Wang et al., 2015). However, it is still unknown whether VLZ can regulate the fate of hematopoietic cells through 5-HTR1A.

Platelets are small nonnucleated cellular fragments that are essential for functions such as hemostasis, wound healing, angiogenesis, inflammation, and innate immunity (Machlus and Italiano, 2013). They are created in the bone marrow by megakaryocytes, which are enormous polyploid cells that develop from hematopoietic stem cells by megakaryopoiesis. The polyploid cells produced by this intricate process, which are remarkably similar in mice and humans, have a single polylobate nucleus and undergo repeated cycles of endomitosis after numerous rounds of unsuccessful cytokinesis (Marini et al., 2022). A low platelet count is a frequently encountered hematological abnormality. Thrombocytopenia is a medical condition associated with a reduced peripheral blood platelet count, which is characterized as a platelet count below $150 \times 10^9$/L of whole blood (Provan and Semple, 2022). Platelet production can be affected by bone marrow suppression, which is frequently observed in diseases including leukemia, chemotherapy, and sepsis (Lien et al., 2021). Among them, radiotherapy and chemotherapy for tumor treatment are the key factors leading to thrombocytopenia complications. Radiation-induced thrombocytopenia (RIT) is a significant cause of morbidity and mortality and is particularly important in the hematopoietic form of acute radiation syndrome (Tkaczynski et al., 2018). According to several studies, platelet count correlates more strongly with survival following total body irradiation than any other hematologic parameter (Mouthon et al., 1999; Kiang

et al., 2014; Lebois et al., 2016). Bleeding and thrombocytopenia account for a large

amount of morbidity and death in the context of radiation injury. Therefore, the

creation of novel therapeutic approaches to manage thrombocytopenia is essential.

The continual development of platelet synthesis is known as megakaryocytopoiesis.

The most fully researched regulator of MKs in this process is thrombopoietin (TPO),

which is one of a complex network of hemopoietic growth factors. Generally, platelet

production is regulated by TPO. The amount of TPO that is accessible to populations

of hematopoietic stem and progenitor cells is regulated by thrombopoietin receptor

(c-Mpl) on megakaryocytes and platelets (Ninos et al., 2006). In addition to TPO,

other regulators also have significant impacts on megakaryocytopoiesis. These

include the interleukin cytokines family, vascular endothelial growth factors (VEGFs),

SDF-1 (stromal cell-derived factor-1)/FGF-4 (fibroblast growth factor-4), insulin-like

growth factor-1 (IGF-1) and other cytokines (Zheng et al., 2008). The majority of

their effects are independent of TPO signaling to regulate MK differentiation and

thrombopoiesis. Understanding the TPO-independent regulation system is crucial

because it may present new options for understanding the process of development, the

mechanisms behind diseases with aberrant MK and platelet formation, and the

advancement of therapeutic research.

In the present study, we carried out virtual screening to predict the therapeutic activity

of VLZ in the early stage and the predicted results were verified through *in vitro* and

*in vivo* investigations. The molecular mechanism of the effect of VLZ on

thrombocytopenia was probed through network pharmacological methods. Then

experimental verification confirmed that the targeted activation of 5-HTR1A

promoted the maturation and differentiation of megakaryocytes and was related to the

regulation of the SRC-RAS-MEK-ERK1/2 signaling pathways. In conclusion, this

study reports a combination of computational analysis and experimental validation to

explore the thrombogenic effects and potential mechanisms by which 5-HTR1A

drives megakaryocytopoiesis and platelet production and enriches the research on the

regulatory mechanism of serotonergic signaling on hematopoietic lineages.

## 2 Materials and methods

### 2.1 Construction of a virtual screening model based on random forest

The virtual screening flow chart is depicted in **Fig. 1a**. First, we established a database of 39 active small molecule compounds that could promote MK differentiation or platelet generation and 691 inactive compounds. RDKit software (*http://www.rdkit.org*) was used to calculate the molecular descriptors of the 730 compounds to obtain their corresponding values. Molecular Descriptors contain multiple molecular properties, including MolWt, LabuteASA, Chi2n and TPSA. The values of molecular descriptors such as fr_thiocyan, fr_amidine, NumRadicalElectrons and SMR_VSA8 are meaningless. All of them need to be deleted to avoid affecting the results. A total of 160 Molecular Descriptors and their corresponding values are residual. The training sets consisted of data obtained from the remaining 713 small molecule compounds, while the validation sets comprised a random selection of 2 active and 15 inactive small molecule compounds. The equilibrium between the positive and negative samples was achieved using the synthetic minority oversampling technique (SMOTE) (Mathew et al., 2018). According to the value of the Gini index, the data of the top 50%, 60%, 70%, 80%, 90% and 100% of the fraction were obtained, and 6 training sets were generated. Then, 6 training sets were entered into the random forest. The area under the ROC curve (AUC) value was used to choose the greatest training set for the random forest, and the validation set was utilized to test the model. Eventually, the activities of compounds were predicted.

### 2.2 Reagents and antibodies

Vilazodone hydrochloride (V823988, purity of 98.0%) was obtained from Macklin (Shanghai, China). The substance was dissolved in DMSO and then stored at a concentration of 10 mM/L. Antibodies against p-SRC (6943S), SRC (2109S), p-MEK (2338S), MEK1/2 (4694S), p-ERK1/2 (4370S), ERK1/2 (4696S) were obtained from Cell Signaling Technology (Boston, MA, USA), against NFE2 (11089-1-AP), RUNX1 (25315-1-AP), FOS (66590-1-IG), GAPDH (60004-1-IG) were obtained

from Proteintech (Wuhan, China), against RAS (T56672S), EGR1 (T57177) was

obtained from Abmart (Shanghai, China), against HTR1A (BA1391) was obtained

from Boster Bio-Engineering Co. (Wuhan, China). Second antibody against rabbit

(4414) and against mouse (4410) were obtained from Cell Signaling Technology

(Boston, MA, USA).

**2.3 Cell culture**

HEL and Meg-01 cells were acquired from the American Type Culture Collection

(Manassas, VA, USA) and used throughout the experiments. Cells were kept at 37 °C

in a humid incubator with 5% $CO_2$ and grown in RPMI Medium 1640 Basic (CAT:

SP032020500, Sperikon Life Science & Biotechnology Co. Ltd., Chengdu, China)

with 10% fetal bovine serum (FBS, Gibco, Thermo Fisher Scientific, Waltham, MA,

USA), 100 U/mL penicillin and 100 μg/mL streptomycin (Beyotime, Sichuan, China).

**2.4 Cell Proliferation Assay**

HEL and Meg-01 cells were treated with VLZ in 96-well flat-bottomed plates for 1 to

5 days. The amount of lactic acid dehydrogenase (LDH) released from the cells was

measured to evaluate cytotoxicity. We used an LDH cytotoxicity assay kit (Beyotime,

Jiangsu, China). The maximum LDH control was used to calculate the total amount of

LDH present in the cells. The absorbance was measured at 490 nm. A Cell Counting

Kit-8 (CCK-8) assay (Dojindo, Kumamoto, Japan) was used to assess the

proliferation of HEL and Meg-01 cells. Briefly, $5 \times 10^3$ cells were seeded in 96-well

plates and then exposed to different concentrations of VLZ (2.5, 5, and 10 μM) at

37 °C and 5% $CO_2$ for 5 days. Each well received CCK-8 solution after the treatment,

which was then incubated for two hours at 37 °C. The absorbance (OD) was measured

at 450 nm.

**2.5 Analysis of cell morphology**

HEL and Meg-01 cells ($4\times10^4$ cells/well) were seeded in 6-well plates. Microscopy

(Olympus CKX53, Japan) at 10× resolution was used to detect the change in cell

shape after treatment with VLZ (2.5, 5 and 10 μM) or the positive control PMA (0.8

nM) for 5 days. Giemsa staining was used to analyze the formation of giant

multinucleated cells. Cells were spread out on slides using a cyto centrifuge and

centrifuged at 200 rpm while being treated with fixing solution (methanol: glacial

acetic acid = 3:1 (v/v)). Giemsa solution (Solarbio, Beijing, China) was added to each

sample for 15 min. Finally, the stained cells were photographed under an electron

microscope. Changes in F-actin were analyzed by phalloidin staining. Cells were

immobilized with 4% paraformaldehyde (Biosharp, Anhui, China) for 10 min,

permeabilized with 0.05% Triton X-100 solution for 10 min at room temperature and

washed with PBS. The sample was incubated for 30 minutes at room temperature in

the dark with 200 μL of TRITC-labeled phalloidin working solution (Solarbio, Beijing,

China). The sample was then counterstained with 200 μL DAPI (Solarbio, Beijing,

China) for 30 s. Under the guidance of a laser scanning confocal microscope (Leica,

Germany), cell samples were observed and photographed.

**2.6 Analysis of megakaryocyte differentiation and ploidy**

Cells incubated with VLZ (2.5, 5 and 10 μM) for 5 days were collected and stained

with FITC-CD41 and PE-CD42b (BioLegend, San Diego, CA, USA) antibodies for

15 min on ice in the dark. The percentages of $CD41^+CD42b^+$ cells were evaluated by

a flow cytometer (BD Biosciences, San Jose, CA, USA). Cells were fixed with 70%

ice-cold ethanol for an overnight period. The samples were washed twice with PBS.

The sample was then treated with 20 g/mL propidium iodide (PI) and incubated on ice

in the dark for 30 min. Finally, the cells were examined by flow cytometry. Cell

apoptosis was measured by flow cytometry using the Annexin V-FITC/PI Apoptosis

Detection Kit.

**2.7 Zebrafish model**

The Chinese National Zebrafish Resource Center (Wuhan, China) provided transgenic

zebrafish (itga2b: eGFP). All zebrafish procedures were conducted according to the

guidelines of Southwest Medical University's laboratory animal ethics committee

(License NO.20230716-002). The larvae of transgenic zebrafish expressing Tg

(itga2b: eGFP) were collected at 3 days post fertilization (dpf) and distributed evenly

in 12-well plates, with 20 larvae per plate. These larvae were subsequently exposed to

varying concentrations (10-40 μM) of VLZ. At 5 dpf, the zebrafish were fixed in 4%

paraformaldehyde overnight at 4 °C. Following fixation, the zebrafish larvae were

embedded in agarose on a petri dish for imaging purposes. The cells of 5 dpf zebrafish

treated with VLZ were extracted from the lysate and detected by flow cytometry.

**2.8 Radiation-induced thrombocytopenia mouse model**

We purchased specific pathogen-free (SPF) Kunming (KM) mice from Da-shuo

Biotechnology Limited in Chengdu, Sichuan, China. The mice were approximately 9

weeks old, weighed approximately 20 g and were kept under regular circumstances

(22±2 °C, 55±5% humidity, and a 12 h light and dark cycle). The experimental animal

ethics committee of Southwest Medical University was followed during all

procedures on mice (License NO.20230716-002). Except for the control group, the

other mice were given a total body X-ray dose of 4 Gy after a week of acclimatization

to induce a mouse model of thrombocytopenia. Each group of mice randomly

consisted of six males and six females: normal group; model group; TPO group (3000

U/kg); and VLZ treated group (2.5, 5 and 10 mg/kg). TPO and VLZ were injected

intraperitoneally daily post irradiation for 12 days. Normal saline was

intraperitoneally administered to the control and model groups.

**2.9 Hematologic parameters examination**

251  On the indicated days after injection, blood (40 μL) was collected from the orbit and

252  gently mixed with diluent (160 μL) to prevent clotting. The samples were measured

253  by a Sysmex XT-2000iV automatic hematology analyzer (Kobe, Japan), including

254  platelet counts (PLT), platelet volume (MPV), red blood cell counts (RBC), and white

255  blood cell counts (WBC).

**2.10 Flow cytometry analysis of cells in vivo**

Flow cytometry analysis of cells in vivo were performed as described previously

(Zhang et al., 2023). Briefly, PB, BM, and spleen cells were examined by flow

cytometry using the BD FACSCanto II flow cytometer (BD Biosciences, San Jose,

CA, USA). The following antibodies were used: anti-CD41 (BioLegend, San Diego,

CA, USA), anti-CD117 (c-Kit, BioLegend, San Diego, CA, USA), anti-61 (BD

Biosciences, San Jose, CA, USA), PI/RNase Staining Buffer (BD Bio- sciences, San

Jose, CA, USA), and anti-CD62P (BioLegend, San Diego, CA, USA). The antibodies

were incubated in the dark for 30 min on ice.

**2.11 Carotid artery thrombosis test**

The mice were administered normal saline, TPO, or VLZ daily post irradiation for 12 days and then subjected to the carotid artery thrombosis test following published protocol (Xu et al., 2017). In a nutshell, filter paper (1 mm × 3 mm) saturated with 10% $FeCl_3$ was placed on the left carotid artery for 3 minutes to induce thrombosis in the artery. The carotid artery was cleaned with PBS after the filter paper was removed. From the time of damage to stable occlusion (defined as no flow for 120 min), blood flow was constantly monitored with a vascular flow probe and Transonic TS420 flowmeter (Transonic Systems, Ithaca, NY, USA).

**2.12 Platelet aggregation**

Blood was collected from the inferior vena cava of the mice after they had been given general anesthesia and placed in a syringe containing anticoagulant as previously described (Lin et al., 2021). After centrifugation (100 × g) for 10 min to obtained platelet-rich plasma (PRP) and adjusted the concentration to $3 \times 10^9$ platelets/mL by saline solution. ADP (Helena Laboratories, TX, USA) was used to stimulate the platelets, and platelet aggregation was measured using a turbidimetric aggregation-monitoring device (Helena Laboratories, Beaumont, TX, USA).

**2.13 Platelet adhesion**

Each confocal dish was first filled with 1 mL of 5 g/mL collagen (Helena Laboratories, TX, USA) and left at 4 °C overnight. Each well was cleaned with PBS before being blocked for 1 h at room temperature with 1% BSA. Then, PRP ($2\times10^9$ cells/mL) was added, and the cells were incubated for 1 h at 37 °C. After unbound platelets were removed with PBS, the rest of the steps were similar to the phalloidin staining, and TRITC-conjugated phalloidin was used to stain the adhering platelets. Representative images were acquired under a fluorescence microscope. imageJ software was used to calculate the average cover of adhering platelets for quantification and statistical analysis (Zhou et al., 2023).

**2.14 Whole blood clot and washed platelet retraction**

Whole blood was diluted with saline at a 1:3 ratio in a siliconized flat-bottomed glass test tube. Thrombin (Solarbio, Beijing, China) and $CaCl_2$ were used for the initial

clotting and incubation at 37 °C for two hours was performed. After that, the

supernatant was eliminated, and clots were weighed. Washed platelets were obtained

as previously described. Washed platelets were diluted to $2 \times 10^8$ platelets/mL with

saline in glass test tubes. Then, retraction was initiated with thrombin and $CaCl_2$. The

test tubes were incubated at 37 °C for 2 hours. The percentage size of retracted

platelets over time was determined from images using ImageJ (Gauer et al., 2023).

**2.15 Network pharmacology and molecular docking simulation**

Network pharmacology and molecular docking simulation were performed according

to a previously described protocol (Ruan et al., 2023). The overlapping targets of VLZ

action and thrombocytopenia were obtained from the *SwissTargetPrediction,*

*DisGeNET* and *GeneCards* databases. For visibility and topology analysis, the eligible

data were entered into *Cytoscape_v3.7.1* software using the STRING_v11.5 database.

GO and KEGG analysis visualization using the *DAVID_v6.8* database, programs

*Cytoscape/ClueGO_v3.9.0* and *GraphPad Prism v9.1.1.233*. The crystal structures of

the core target proteins, including MAPK1 (PDB:2Y9Q) and SRC (PDB:1O4A), were

obtained from the *UniProt* website. These proteins were structurally altered using the

*Sybyl-X 2.0* program. The total Surflex-Dock scores represent the binding affinities.

Finally, the docking result was visualized utilizing *PyMOL_v2.5.2* and *Ligplus_v2.2*

*software* (Rosignoli and Paiardini, 2022; Liu et al., 2022).

**2.16 Western blot analysis**

Total cellular proteins were harvested by $1\times$ RIPA lysis buffer (Cell Signaling

Technology, MA, USA) containing phosphatase inhibitor and protease inhibitor. The

proteins were loaded and separated by SDS–PAGE, followed by transfer to a

polyvinylidene difluoride (PVDF) membrane. The PVDF membrane was probed with

the primary antibodies overnight at 4 °C. The blots were then incubated with

secondary antibodies after rinsing three times with PBST. The blots were detected

using ECL detection reagent (4A Biotech Co., Ltd., Beijing, China) and visualized

with a scanner. Then quantified the bands were performed with ImageJ software.

Positive signals were detected when the relative intensity of target proteins or

phosphorylated proteins was compared with that of GAPDH or total proteins.

**2.17 Immunofluorescence staining**

We collected cells treated with or without VLZ for 5 days and transferred them to microscope slides. These cells were fixed with 4% paraformaldehyde for 10 min, permeabilized with 0.1% Triton X-100, incubated with 1% BSA for 1 h at room temperature, and incubated with primary antibodies for an additional 6 h at 4 °C followed by the corresponding secondary antibody for 1 h. DAPI was used to counterstain the nuclei for 10 min at room temperature in the dark. Images were taken under a fluorescence microscope.

**2.18 Drug Affinity Responsive Target Stability Assay**

Meg-01 cells were harvested and lysed with RIPA lysis buffer for 15 min on ice. The samples were centrifuged at 12000 rpm for 15 min to obtain the supernatant. The protein concentration was measured with Bradford reagent and diluted to 5 mg/mL, followed by treatment with VLZ or DMSO for 1 h at room temperature. Next, various ratios of pronase (1:500, 1:1000 and 1:1500) were added and incubated for 10 min at 40 °C. Then, $5 \times$ loading buffer was added to the sample and boiled at 95 °C for 10 min. The sample proteins were analyzed by Western blotting.

**2.19 Statistical analysis**

The mean and standard deviation (SD) of at least three independent experiments are used to express all data. One-way analysis of variance (ANOVA) and Tukey's post hoc test were used to evaluate the statistical comparisons among several groups. Two-tailed Student's *t* tests were used to compare the differences between the two groups. Statistical significance was defined as a *P* value <0.05.

# 3 Results

**3.1 Prediction of potential megakaryocytopoiesis inducers via the virtual screening model**

We focused on building a drug screening model consisting of random forest in this work. The optimal training set for predicting drug activity was selected after acquiring 6 training sets (**Fig. 1a**) and utilizing random forest in Orange software (https://orangedatamining.com/) to perform operations on the 6 training sets. The Gini

index was used to rank the importance of 20 molecular descriptors to improve

predictive accuracy (**Fig. 1b**). To assess the prediction capability of these training sets,

we first determined the parameters of the model. When 30 trees and 2 attributes were

assessed at each split, we found that the training set with a score of 50% provided the

model with the best prediction performance, and the area under the AUC value was

0.954 (**Fig. 1c**). To further validate the performance of the model, the validation set

containing 2 active compounds and 15 inactive compounds was entered into the

model. The findings showed that the prediction accuracy of the model was 83.33%.

Subsequently, 100 chemical compounds from the Discovery Probe FDA-approved

Drug Library were then predicted using the model. VLZ was projected to have a

predicted score of 0.52, indicating that it possessed biological activity that promoted

MK differentiation or platelet formation.

**Fig.1 Drug screening model construction. a** Flowchart for the screening model construction. **b**

The top 20 molecular descriptors with high scores. **c** ROC curves of importance ratios.

## 3.2 VLZ dose-dependently promotes megakaryocyte maturation and differentiation

To verify the compound activity predicted by the model, we evaluated the effect on HEL and Meg-01 cells after VLZ intervention for the specified period to search for the ideal VLZ concentration for MK differentiation. For the in vitro study, we used drug dosages of 2.5, 5 and 10 μM. LDH assay was used to determine the cytotoxicity of VLZ on HEL and Meg-01 cells. The outcomes demonstrated that 2.5, 5 and 10 μM VLZ were within a safe concentration range for MK differentiation since LDH release at days 1, 3, and 5 in the VLZ group was not different from that in the control group. (**Supplementary Fig. 1a, b**). Furthermore, cells started to exhibit a decrease in proliferation rates after treatment on the 3rd day, and at the 5th day, there was a concentration-dependent inhibition of proliferation (**Supplementary Fig. 1c, d**). This may be the result of cell differentiation. Microscopic images showed that obvious large megakaryocyte-like cells were increased in the VLZ (2.5, 5 and 10 μM) and PMA (a positive control that is known to induce MK differentiation) groups on the 5th day (**Fig. 2a, b**). The data suggested that VLZ is a safe inducer to inhibit MK proliferation and induce differentiation.

Given that the established role of VLZ promotes megakaryocyte generation, we continued to investigate whether VLZ exerted excellent activity in inducing MK-typical differentiation and polyploidization. Wright-Giemsa staining showed that VLZ significantly increased the nuclear-to-cytoplasm ratio, large-sized cells, and deeply stained multilobulated nuclei compared with those in the control (**Fig. 2c, d**). Similarly, we evaluated the morphological alteration of VLZ-induced polyploidization, as determined by phalloidin staining. On the 5th day, significant numbers of multinucleated cells, cytoplasm enlargement and F-actin aggregation were observed in the VLZ and PMA treatment groups (**Fig. 2e, f**). Megakaryocyte differentiation and

maturate are frequently assessed using changes in the expression of distinctive molecular markers. MKs have specific antigens called CD41 and CD42b (Jimenez et al., 2015). After 5 days of treatment, flow cytometry analysis showed that VLZ obviously increased CD41$^+$CD42b$^+$ cells in a dose-dependent manner in HEL and Meg-01 cells (**Fig. 2g, h**). The typical feature of mature megakaryocytes is polyploidization of the nucleus (van Dijk et al., 2018). Flow cytometry analysis also showed that VLZ significantly increased the population of ≥4 N DNA content in HEL and Meg-01 cells in a dose-dependent manner after 5 days of VLZ intervention (**Fig. 2i, j**). Therefore, we further determined the action of VLZ on DNA content. We know that proplatelet formation requires local apoptosis. Annexin V/PI staining flow cytometry showed that Vilazodone induced excessive apoptosis of megakaryocytes on day 5 in a concentration-dependent manner (**Fig. 2k, l**). These data suggest a dose-dependent promotion of MK maturation and differentiation by VLZ.

**Fig.2 VLZ promotes megakaryocyte differentiation and enhances the DNA ploidy of HEL and Meg-01 cells. a, b** Representative images of HEL and Meg-01 cells treated with different concentrations of VLZ (2.5, 5 and 10 μM) for 5 days. Bars represent 100 μm. The positive control was PMA. **c, d** Giemsa staining of HEL and Meg-01 cells treated with VLZ (2.5, 5 and 10 μM) or PMA for 5 days. Bars represent 25 μm. **e, f** Phalloidin staining of HEL and Meg-01 cells treated with VLZ (2.5, 5 and 10 μM) or PMA for 5 days. DAPI staining nuclei (blue) and Phalloidin staining of F-actin (red). Bars represent 100 μm. **g, h** Flow cytometry analysis of the percentage of $CD41^+CD42b^+$ complexes surface expression on HEL and Meg-01 cells by VLZ (2.5, 5 and 10 μM) or PMA for 5 days. The histogram shows the percentage of $CD41^+CD42b^+$cells for each group. **i, j** Flow cytometry analysis of the DNA ploidy of HEL and Meg-01 cells treated with VLZ (2.5, 5 and 10 μM) or PMA for 5 days. The histogram shows the percentages of DNA ploidy. **k, l** Flow cytometry analysis of the cell apoptosis of HEL and Meg-01 cells treated with VLZ (2.5, 5 and 10 μM) or PMA for 5 days. Data are shown as the mean ± SD from three independent experiments. $*P \leq 0.05$, $**P \leq 0.01$, and $***P \leq 0.001$, vs the control group.

### 3.3 VLZ increases thrombocytes in zebrafish embryos

Compared with cell models, zebrafish are an excellent model system for drug delivery and biology (Kong et al., 2020). We further appraised the overall effect of VLZ on promoting thrombocyte production in zebrafish (**Fig. 3a**). The maximum dosage for subsequent investigations was set at 40 μM since toxicity analyses indicated that VLZ incubation had no discernible impact on the survival or general development of 3 dpf zebrafish embryos within a range of 2.5 to 40 μM (**Fig. 3b**). The embryos of platelet labeled Tg (itga2b: eGFP) transgenic zebrafish at 3 dpf were treated with VLZ (10-40 μM), and the itga2b: eGFP thrombocytes in the tail area were carefully examined at 5 dpf. Notably, VLZ increased the quantity of $GFP^+$ cells in the tail region, which represents platelets (**Fig 3c, d**). Next, we extracted total cells of zebrafish administered VLZ for quantification analysis of its effects on endogenous blood circulation through flow cytometry. As anticipated, there was a dose-dependent increase in the number of itga2b: eGFP cells among the total cells in the VLZ (10-40 μM) treatment groups compared to the control group, suggesting that circulating

platelets were increased after VLZ stimulation (**Fig. 3e**). These results demonstrate

that VLZ substantially increases thrombocytes in zebrafish embryos.

**Fig.3 VLZ administration enhances thrombogenesis in Tg (itga2b: eGFP) transgenic**

**zebrafish. a** Timeline of the zebrafish intervention. **b** Toxicity effects of different doses of VLZ on

fish survival and general development. **c** itga2b: eGFP thrombocytes in whole tail regions of

control and VLZ (2.5, 5 and 10 μM) treated zebrafish. Bars represent 200 μm. **d** Quantification of

itga2b: eGFP cells in each group (n=6 per group). **e** Flow cytometry analysis of GFP-labeled

platelets of Tg (itga2b: eGFP) zebrafish embryos treated with different doses of VLZ. The data are

shown as the mean ± SD. *$P \leq 0.05$, **$P \leq 0.01$, and ***$P \leq 0.001$, vs the control group.

**3.4 VLZ restores the quantity and function of peripheral platelets in**

**thrombocytopenic mice**

To confirm the in vivo therapeutic influence of VLZ on thrombocytopenia, a thrombocytopenia mouse model was constructed by 4 Gy X-ray total body irradiation, and KM mice were administered VLZ (2.5 mg/kg, 5 mg/kg and 10 mg/kg) and rhTPO (recombinant human thrombopoietin, 3000 U/kg) for 12 days after radiation (**Fig. 4a**). Owing to X-ray exposure, the platelet count in all irradiation groups declined to a minimum on the 7th day, whereas it remained the same in control mice. Nonetheless, VLZ was efficacious in promoting platelet production and resumption in a dose-dependent manner from the 7th day to the 12th day, indicating that VLZ administration boosted platelet recovery when the mice encountered irradiation (**Fig. 4b**). Complementary to the results obtained for zebrafish, the cell count and flow cytometry of peripheral blood showed that platelets increased significantly on the 12th day of 10 mg/kg VLZ administration in normal mice (**Supplementary Fig. 2a, b**). The mean platelet volume (MPV) was likewise discovered and revealed no difference between each group, indicating that VLZ had no impact on MPV (**Fig. 4c**). Additionally, we found that VLZ (5 mg/kg and 10 mg/kg) treatment improved RBC counts on the 10th day and 12th day (**Fig. 4d**). However, there was no significant difference in WBCs between the TPO, VLZ and model groups at any of the tested time points (**Supplementary Fig. 3**). Moreover, the expression of CD41 and CD61 in peripheral blood was determined by flow cytometry analysis. The results showed that VLZ and TPO significantly promoted the expression of CD41 and CD61, consistent with peripheral platelet counts (**Fig. 4e**). Additionally, H&E staining showed no significant differences between each group in major organs, suggesting that VLZ was not systemically toxic. The visceral indexes of the liver, spleen, kidney, and thymus were examined and showed that the index of the liver was high in the model group, which recovered after VLZ or TPO administration (**Supplementary Fig. 4a-e**).

To ascertain whether the platelets stimulated by VLZ therapy were functional, we subsequently studied platelet function in vivo and in vitro. P-selectin is sequestered in Weibel-Palade granules of endothelial cells and α-granules of platelets and translocated to the surface in response to various inflammatory and thrombogenic mediators (DuRoss et al., 2021). However, surface P-selectin (CD62P) increases after

radiation damage (Johansson et al., 2012). Peripheral blood cells were used in a

CD41/CD62P flow cytometry assay, and animals treated with VLZ and TPO had

significantly reduced levels of $CD41^+/CD62P^+$ expression compared to the model

group in the platelet resting state (**Fig. 4f**). In addition, animals treated with VLZ and

TPO showed dramatically increased platelet reactivity in the CD62P assay after ADP

stimulation (**Fig. 4g**). Moreover, platelet adhesion on solidified collagen improved

after VLZ or TPO treatment (**Fig. 4h**). Furthermore, we evaluated platelet aggregation

induced by ADP. Compared to the model group, platelet aggregation caused by ADP

was significantly strengthened in the VLZ-treated and TPO groups (**Fig. 4i**). In

addition, we assessed the impact of VLZ on thrombus development following

vascular damage in a carotid artery thrombosis model. The results suggested that VLZ

promoted thrombus development in mice with thrombopenia, since it took

considerably less time for the VLZ-treated group than the model group to form a

totally occluded thrombus after the commencement of arterial injury (**Fig. 4j**). The tail

bleeding time was used to quantitatively evaluate the hemostatic capacity of platelets.

The findings demonstrated that mice in the VLZ and TPO treatment groups had

considerably reduced tail bleeding time compared with mice in the model group (**Fig.**

**4k**). We next examined whole blood clot and washed platelet clot development and

retraction over time (120 min) following clotting triggered by thrombin and $CaCl_2$.

Differences in the proportion of whole blood clot retraction were observed for VLZ

and TPO and increased clot weight at the endpoint when compared with the model

group (**Fig. 4l**). Consistent with the results indicated for whole blood clot retraction,

VLZ and TPO completely improved washed platelet retraction compared to the model

group (**Fig. 4m**). Collectively, these data suggest that VLZ stimulation is conducive to

platelet recovery in vivo after encountering radiation. In the meantime, it also

contributed to RBC recovery. The platelet function of thrombopenia mice treated with

VLZ was complete, including hemostasis and coagulation, and the response

characteristics to agonists were almost normal.

**Fig.4 VLZ administration counteracts radiation-induced thrombocytopenia in *vivo*. a**
Schematic diagram of TPO (3000 U/kg) and VLZ (2.5, 5 and 10 mg/kg) administration in KM
mice with thrombocytopenia induced by IR. **b-d** Peripheral blood counts showing (b) platelet
counts, (c) MPV and (d) RBC on days 0, 4, 7, 10, and 12 post-IR. In B to E, (n=12 per group) the
data are expressed as the mean ± SD, and two-way ANOVA with Tukey's multiple comparisons
test was used unless otherwise specified. *$P \leq 0.05$, **$P \leq 0.01$, and ***$P \leq 0.001$, vs the model
group. **e** Flow cytometry analysis indicates the expression of CD41 and CD61 in peripheral blood
cells after receiving therapy for 12 days. The histogram represents the percentage of CD41⁺CD61⁺
cells in each group. **f** Representative image of CD62p in washed platelets in each group. **g**
Representative images of CD62p in ADP (10 mM) washed platelets in each group. **h** Micrographs

of collagen-coated slides with the same number of platelets perfused. Red represents platelets. The

histogram shows the average of red fluorescence on the whole surface by ImageJ software. **i**

Platelet aggregation was measured by a turbidimetric aggregation-monitoring device under ADP

stimulation. The histogram represents the aggregation results expressed as the maximum

aggregation amplitude of platelets in each group. **j** The rate tracings of carotid blood flow. The

histogram shows the mean carotid artery occlusion times of each group. **k** Tail bleeding time was

measured in each group. **l** Representative images of whole blood clots at time 0, 60 and 120 min

(end point) in each group. The histogram represents the percentage of final clot weight. **m**

Representative images of washed platelet clots at time 0 and 120 min (end point) in each group.

The histogram shows clot retraction at 120 min as a percentage clot area from 0 min. The data

represent the mean ± SD of three independent experiments. $*P \leq 0.05$, $**P \leq 0.01$, and $***P \leq$

$0.001$, vs the model group.

### 3.5 VLZ rescues bone marrow and splenic hematopoiesis after radiation injury

Bone marrow is the most important hematopoietic organ in the human body, and

extramedullary hematopoiesis in the spleen may occur in response to physiological

stress or illness and may serve as a substitute tissue site for bone marrow

hematopoiesis (Lucas, 2021; Short et al., 2019). We thus investigated whether VLZ

could promote the differentiation of hematopoietic progenitors into megakaryocytic

progenitors and megakaryocytes in the BM and spleen considering the increased

numbers of megakaryocytes in the VLZ-treated group. Following VLZ and TPO

treatment for 12 days, H&E staining results showed a clearly increased number of

megakaryocytes in the BM and spleen (**Fig. 5a, b**). According to flow cytometry

analysis, the percentages of c-Kit$^+$CD41$^-$ (hematopoietic progenitors), c-Kit$^+$CD41$^+$

(megakaryocytic progenitors), and c-Kit$^-$CD41$^+$ (megakaryocytes) cells were

considerably increased in the TPO- and VLZ-treated groups compared to the model

group (**Fig. 5c**). The findings imply that VLZ can stimulate megakaryocyte formation

at various megakaryopoiesis stages. CD41 and CD61 were detected by flow

cytometry as megakaryocyte-specific markers. The findings revealed that in the TPO-

and VLZ-treated groups, the fraction of CD41$^+$CD61$^+$ cells in the BM and spleen was

noticeably higher than that in the model group, which indicated that VLZ promoted

BM megakaryocyte differentiation (**Fig. 5d, e**). Megakaryocytes from the BM and

spleen were also examined for DNA content using flow cytometry. As anticipated, the

TPO- and VLZ-treated groups greatly outnumbered the model group in increasing

terms of the ploidy of BM and spleen megakaryocytes (**Fig. 5f, g**). All these data

indicate that VLZ promotes platelet formation in the BM and spleen of irradiated

mice by rescuing megakaryopoiesis. Collectively, these results demonstrate that VLZ

has excellent therapeutic effects on mice suffering from thrombocytopenia.

**Fig.5 VLZ rescues bone marrow and splenic hematopoiesis after radiation injury. a, b** H&E

staining shows the megakaryocytes in BM and spleen after mice were treated with normal saline,

TPO (3000 U/kg) and VLZ (2.5, 5 and 10 mg/kg) for 12 days. Ten microscopy fields per sample

were counted. The circles mark the megakaryocytes. The histogram shows the number of

megakaryocytes in the BM and spleen in each group. **c** The examination of the expression of c-Kit

and CD41 in each group by flow cytometry after receiving therapy for 12 days. The histogram

represents the percentage of c-Kit$^+$CD41$^-$, c-Kit$^+$CD41$^+$, and c-Kit$^-$CD41$^+$ cells in each group. **d, e**

Flow cytometry analysis indicates the expression of CD41 and CD61 in BM and spleen cells after

receiving therapy for 12 days. The histogram represents the percentage of $CD41^+CD61^+$ cells in

each group. **f, g** Flow cytometry analysis indicates the cell ploidy of BM and spleen cells of each

group. The histogram represents the cell ploidy in BM and spleen cells of each group. All data

represent the mean ± SD of three independent experiments. *$P \leq 0.05$, **$P \leq 0.01$, and ***$P \leq$

0.001, vs the model group.

**3.6 Potential therapeutic targets and signaling pathway prediction based on**

**network pharmacology and molecular docking**

We used the "drug targets" and "disease targets" through database screening to

estimate the therapeutic targets of VLZ for thrombocytopenia. After data integration,

we discovered 109 targets related to VLZ treatment. A total of 4356 targets related to

thrombocytopenia were obtained after duplicates were removed. Potential targets of

VLZ for treating thrombocytopenia included 63 common targets (**Fig. 6a**). The

STRING database and Cytoscape_v3.7.1 software were used to assess the 63 putative

therapeutic targets and create an interaction network (**Fig. 6b**). Then, we performed a

topology analysis to identify the core targets. The top 6 target proteins of MAPK1,

HTR1A, SRC, NFE2, ABL1, and MCL1 were regarded as core proteins (**Fig. 6c**).

Utilizing molecular docking stimulation, the direct connection between VLZ and the

primary targets was further investigated. Molecular affinity for the target was

indicated by a docking score of 4.52 kcal mol$^{-1}$, with a docking score >5 kcal mol$^{-1}$

being considered indicative of a strong binding affinity (Hsin et al., 2013). According

to the docking results, the binding scores of MAPK1 and SRC with VLZ were 8.97

and 5.31, respectively, which indicated strong affinity for VLZ (**Fig. 6d**). GO and

KEGG enrichment analyses were conducted to elucidate the potential underlying

mechanism of VLZ action on thrombocytopenia. The outcomes demonstrated that

VLZ treatment for thrombocytopenia was associated with a total of 58 GO terms (**Fig.**

**6e**). These GO annotations mainly involve positive regulation of MAP kinase activity,

positive regulation of ERK1 and ERK2 cascade, stem cell differentiation and positive

regulation of smooth muscle cell migration, growth factor binding, transcription factor

binding, and ATP binding. In summary, the top 10 CC, BP, and the most VLZ-related

process-enriched MF terms were sorted independently in descending order of P values.

(**Fig. 6f**). KEGG enrichment analysis revealed that VLZ regulatory action in VLZ was

significantly enriched in the MAPK signaling pathway, RAS signaling pathway,

PI3K-Akt signaling pathway, and Rap1 signaling pathway. The 20 most enriched

pathways were arranged in descending order of P values (**Fig. 6g**). Therefore, these

results suggest that VLZ might protect against thrombocytopenia via activation of the

SRC and MAPK (RAS/MEK/ERK) signaling pathways.

**Fig.6 Network pharmacology and molecular docking analysis of VLZ activity in**

**thrombocytopenia. a** Venn diagram of the common targets of VLZ and thrombocytopenia. **b** The

VLZ-targets-Thrombocytopenia network that was constructed by Cytoscape_v3.7.1 software. **c**

Protein-protein interaction (PPI) network based on the core targets of VLZ against

thrombocytopenia through the screening conditions of Degree >12, BC > 0.01105525, CC >

0.52777778. **d** Molecular docking shows the binding ability between VLZ and its core targets

(MAPK1 and SRC). **e** Visualization of cellular component, biological process, and molecular

function enrichment analysis. **f** The top 10 of CC, BP, and MF terms showing the greatest

enrichment with VLZ-related processes are listed in ascending order of *P* values. **g** The top 20 of

KEGG pathways enrichment analysis for the mechanisms of VLZ against thrombocytopenia. A

higher richness factor indicates more enrichment. The number of genes enriched in each pathway

is represented by the size of the bubble. The *p* value range is shown by the color of the bubble.

**3.7 The SRC/MAPK signaling pathway is activated during VLZ-facilitated**

**megakaryopoiesis**

Given the emphasis of the in-silico prediction results emphasized the involvement of

the MAPK signaling pathway and RAS signaling pathway in a latent regulatory

mechanism of VLZ, we proceeded to investigate the expression of the core proteins

within this pathway, which are diverse and embody VLZ-prompted pathway

regulation: (p-)SRC, RAS, (p-)MEK, and (p-)ERK1/2 (**Fig. 7a-d**). After 5 days, the

phosphorylation levels of SRC, MEK, ERK1/2 and RAS were significantly changed

in VLZ (2.5, 5 and 10 μM)-treated Meg-01 cells. To produce proplatelets and

differentiate and mature megakaryocytes, the transcription factors NF-E2, EGR1,

FOS and RUNX1 link upstream molecular signals with downstream transcript

expression (Noetzli et al., 2019; Kim et al., 2019). After VLZ treatment at 2.5 μM, 5

μM and 10 μM, we further confirmed that VLZ treatment led to a dose-dependent

increase in the expression of NF-E2, EGR1, FOS and RUNX1 at the translational

level (**Fig. 7e-h**). Additionally, immunofluorescence research verified that VLZ

boosted the expression of NF-E2 and RUNX1 in Meg-01 cells (**Fig. 7i, j**). Finally, we

used a blocking technique to determine whether they controlled the megakaryocyte

differentiation that VLZ generated. The ERK1/2-specific inhibitor SCH772984 was

used to block the phosphorylation of ERK1/2. Flow cytometry analysis demonstrated

that VLZ-induced acceleration of megakaryocyte differentiation was interrupted by

SCH772984 (3 µM) therapy (**Fig. 7k**). This result supports the molecular mechanism

by which VLZ activates the SRC/RAS/MEK/ERK1/2 cascade to promote the mRNA

expression and nuclear translocation of related transcription factors to induce MK

maturation and platelet production.

**Fig.7 VLZ induces MK maturation and differentiation by activating the**

**SRC/RAS/MEK/ERK signaling pathway. a-h** The expression of SRC, RAS, MEK, ERK,

NF-E2, EGR1, FOS and RUNX1 were detected by western blotting after Meg-01 cells were

treated with VLZ (2.5, 5 and 10 µM) for 5 days. **i, j** Immunofluorescence analysis of the

expression of NF-E2 and RUNX1 in Meg-01 cells after VLZ (2.5, 5 and 10 µM) intervention for 5

days. Cells were stained with DAPI for nuclei (blue) and antibodies for NF-E2 (green). Bars

represent 100 μm. **k** Flow cytometry analysis of the percentage of CD41$^+$ CD42b$^+$ complexes

surface expression on HEL and Meg-01 cells treated with VLZ (10 μM), SCH772984 (3 μM) and

VLZ (10 μM) + SCH772984 (3 μM) treated for 5 days. The histogram shows the percentage of

CD41$^+$ CD42b$^+$ cells for each group. The data represent the mean ± SD of three independent

experiments. *$P \leq 0.05$, **$P \leq 0.01$, and ***$P \leq 0.001$, ns: no significance, vs the control group.

**3.8 VLZ stimulated MK polyploidization and maturation via activation of the**

**5-HT1A receptor**

Despite VLZ being an agonist of 5-HTR1A, to further validate whether the target also

acted on MKs, we explored whether VLZ interacts specifically with 5-HTR1A to

promote MK differentiation. As a preliminary study, the expression of 5-HTR1A in

Meg-01 cells was evaluated. We also demonstrated that VLZ doses increased the

expression of 5-HTR1A (**Fig. 8a, d**). A DARTS assay was then employed to observe

the binding performance of VLZ to 5-HTR1A on Meg-01 cells. The DARTS results

demonstrated that VLZ administration increased the stability of 5-HTR1A against

protease-induced degradation in a concentration-dependent manner (**Fig. 8b, c**). To

obtain further evidence for the targeting of VLZ to 5-HTR1A, WAY-100635 (2.5 μM,

5-HT1A receptor antagonist) was applied with VLZ. In this case, the observed effects

in vitro were mediated via a 5-HTR1A-dependent mechanism, as WAY-100635

administration blocked the accelerative effects of VLZ in Meg-01 cells (**Fig. 8e-k**).

Once activation of 5-HTR1A induced by VLZ was blocked, the expression of RAS

and phosphorylation of SRC and ERK1/2 were significantly inhibited (**Fig. 8l**). We

indicated that the agonistic action of 5-HTR1A induced by VLZ was responsible for

the activation of RAS and phosphorylation of ERK1/2. Therefore, we further

demonstrated that VLZ activates 5-HTR1A on MKs to regulate the

SRC/RAS/MEK/ERK signaling pathway, encouraging the nuclear translocation and

mRNA expression of relevant transcription factors to trigger MK maturation and

platelet formation.

**Fig.8 VLZ stimulates megakaryocytopoiesis via the 5-HT1A receptor. a** Representative immunoblot images and biochemical quantification of 5-HTR1A after treatment with VLZ (2.5, 5 and 10 μM) in Meg-01 cells for 5 days. **b** The DARTS assay for target validation. 5-HTR1A protein stability was increased upon VLZ (200 μM) treatment in Meg-01 lysates. Pronase was added using several dilutions (1:500, 1:1000, or 1500) from 50 μg/mL stock for 10 min at 40 °C. **c** The DARTS assay demonstrated the dose-dependent binding of VLZ to 5-HTR1A in Meg-01

cells. Treatment with pronase (1:1000) was conducted for 10 min at 40 °C. **d** Immunofluorescence

analysis of the expression of 5-HTR1A in Meg-01 cells after VLZ (2.5, 5 and 10 μM) intervention

for 5 days. Cells were stained with DAPI for nuclei (blue) and antibodies for 5-HTR1A (green).

Bars represent 100 μm. **e-k** Meg-01 cells were treated with VLZ (10 μM), WAY-100635 (2.5 μM),

VLZ (10 μM) + WAY-100635 (2.5 μM) for 5 days. (e) Representative images, bars represent 25

μm. (f) Giemsa staining of Meg-01 cells, bars represent 100 μm. (g) Phalloidin staining of Meg-01

cells, bars represent 100 μm. (h, i) Flow cytometry analysis the expression of CD41/CD42b and

the DNA ploidy. (j, k) The histogram shows the percentage of $CD41^+CD42b^+$ cells and DNA

ploidy for each group. **l** Western blot analysis of 5-HTR1A, RAS and ERK expression after

Meg-01 cells were treated with VLZ (10 μM), WAY-100635 (2.5 μM), and VLZ (10 μM) +

WAY-100635 (2.5 μM) for 5 days. The histogram shows the expression of 5-HTR1A, RAS and

ERK in each group. The data represent the mean ± SD of three independent experiments. $*P \leq$

$0.05$, $**P \leq 0.01$, and $***P \leq 0.001$, ns: no significance, vs the control group.

## 4 Discussions

Thrombocytopenia caused by long-term radiotherapy and chemotherapy exists in

cancer treatment (Tkaczynski et al., 2018; Kiang et al., 2014). This complication

seriously affects the prognosis and endangers the life of patients (Mouthon et al.,

1999). Thrombocytopenia remains a challenging disease despite multiple advances in

drug research radiotherapies, and chemotherapies Currently, there are not entirely

reliable and secure medications for the rapid treatment of thrombocytopenia.

Although TPO-RA is a classical clinical treatment, it is also ineffective in individuals

with TPO resistance or c-Mpl gene loss-of-function mutations(Ghanima et al., 2019;

Poston and Gernsheimer, 2019; van den Oudenrijn et al., 2000). In addition, there are

few agonists targeting TPOR, so the research and development of novel

non-TPOR-dependent targets and agonists is currently a key strategy for the treatment

of thrombocytopenia (Jenkins et al., 2007; Nakamura et al., 2006). Several studies

have demonstrated the mechanisms underlying 5-HT-regulated hematopoietic lineages.

It is crucial and necessary to investigate this TPO-independent signaling pathway.

Here, we identified 5-HTR1A as an attractive target, VLZ could directly bind to

5-HTR1A to enhance megakaryocyte differentiation and platelet formation by

facilitating the activation of the SRC and MAPK signaling pathways at the same time

(**Fig. 9**).

**Fig.9 Schematic representation of the mechanism by which VLZ regulates megakaryocyte**

**differentiation and thrombopoiesis**

Machine learning (ML) has unique advantages in compound activity prediction and

screening (Bannigan et al., 2021). In this study, we relied on a random forest-based

drug screening model to characterize the hematopoietic activity of VLZ. Excitingly,

VLZ demonstrated an activity prediction score worthy of further investigation, and its

excellent hematopoietic activity was validated in subsequent experiments. In a safe

therapeutic range, it promoted MK differentiation and maturation in a

concentration-dependent manner in vitro. Furthermore, because of its advantages of in

vivo complexity and in vitro high-throughput screening simplicity, zebrafish is a

commonly recognized vertebrate model for research in many fields, including

developmental biology, toxicology, and drug discovery (Kong et al., 2020). Previous

studies showed the construction and application of megakaryocyte production in Tg

(itga2b: eGFP) zebrafish (Wang et al., 2022). At the same time, VLZ significantly

promoted the hematopoietic system of transgenic zebrafish and increased platelet
labeled fluorescence. To verify its therapeutic effect on thrombocytopenia, we
constructed a mouse model of thrombocytopenia by whole-body radiation. After
administration of different doses of VLZ for 12 days, its concentration-dependently
increased platelet counts from day 10 onward, shortening the time to normalization of
platelets. Under normal physiological conditions, it increased platelets in
nonirradiated mice in a manner similar to the zebrafish model. In addition, we found
that VLZ also had a significant effect on the recovery of red blood cells. In addition,
to determine whether VLZ restores platelet function in radiation injury and whether
its inhibitory effect on platelet SERT affects generated platelets, we explored platelet
function in vitro and in vivo. We found that VLZ restored hemostasis thrombosis and
platelet function in radiation model mice. As previously mentioned, SSRI drugs can
inhibit the platelet transporter SERT and affect its function (Scharf, 2012). However,
VLZ appeared to produce SERT-independent enhancement of platelet function in
irradiated mice. Of course, it is possible that the doses and periods of our experiments
did not have a significant influence on platelet inhibition. To determine which
hematopoietic tissues are responsible for the findings, we subsequently analyzed the
bone marrow (BM) and spleen. The results from H&E staining and flow cytometry
demonstrated that VLZ therapy stimulated the differentiation of BM hematopoietic
progenitor cells and megakaryocytes, thereby producing and activating platelets.
Similarly, VLZ increased splenic MK numbers, promoting extra splenial
hematopoiesis. These data suggest that VLZ treats thrombocytopenia by enhancing
bone marrow and splenic megakaryopoiesis and thrombosis. Simultaneously, VLZ
compensated for the extra slight inhibition by promoting the production of new
platelets in vivo. Admittedly, the overall effect of VLZ is to gradually restore the
functional system of platelets damaged by radiation to normal. This suggests that
VLZ may enhance platelet function in thrombocytopenic mice independently of
SERT-mediated 5-HT transport, which needs to be further investigated.
Progenitor cells in the bone marrow undergo a process known as megakaryopoiesis in
which they grow into megakaryocytes, which then create the platelets required for

healthy hemostasis (Crispino, 2022). Network pharmacology and molecular docking

analysis predicted that VLZ may regulate megakaryocyte differentiation through the

SRC and MAPK signaling pathways. Tyrosine-protein kinase proto-oncogene SRC,

including other SRC family kinases (SFK), is crucial for controlling signal

transduction by various cell surface receptors in response to alterations in the cellular

environment (De Kock and Freson, 2020). SRC tyrosine kinases are signaling

mediators that occur before MAPK activation and gene expression that is induced by

cytokines (Singer et al., 2011). Serine/threonine kinases known as mitogen-activated

protein kinases (MAPKs) regulate important cellular responses in eukaryotic

organisms and play a role in proliferation, migration, differentiation, and death (Roux

and Blenis, 2004). Extracellular signal-related kinases (ERKs) (also known as

p44/p42, ERK1/2), ERK5, p38MAPKs, and c-Jun amino-terminal kinases are the four

primary protein subgroups that make up the MAPK family (Anderson et al., 1990).

Further analysis of GO and KEGG enrichment led us to target the RAS-MEK/ERK

signaling pathway. These intracellular signaling pathways play a pivotal role in many

essential cellular processes, including proliferation and differentiation. Likewise, by

additional validation, we showed that VLZ can activate SRC, RAS, MEK, and ERK

and that it may have an impact on the signaling pathways linked to differentiation.

Additionally, cytokines tightly control MK differentiation and thrombosis. Studies

have revealed that NF-E2 is necessary for platelet release, proplatelet production, and

megakaryocyte differentiation and maturation (Machlus et al., 2016; Chen et al.,

2007). EGR1 is an early growth response transcription factor (TF) that regulates a

wide array of genes and plays a major role in MKs, including cell proliferation,

differentiation, and apoptosis (Kanaji et al., 2018; Crist et al., 2013). Several

megakaryocytic cell surface indicators and other auxiliary genes require FOS for

expression (Limb et al., 2009; Ahmad et al., 2017). The transcription factor RUNX1

(previously called AML-1 and CBFα2) is an essential regulator of definitive

hematopoietic stem cell (HSC) ontogeny, HSC self-renewal, MK maturation, and

lymphocyte differentiation (Huang et al., 2012; Raghuwanshi et al., 2020). The results

demonstrated that VLZ activated these TFs in a concentration-dependent manner. To

further investigate whether the MAPK signaling pathway was responsible for

VLZ-induced MK differentiation, we blocked this signaling pathway with its inhibitor

SCH772984. As anticipated, we discovered that blocking the MEK/ERK signaling

pathway eliminated the ability of VLZ to cause MK differentiation. Thus far, the

signaling pathway predicted by network pharmacology has been verified. VLZ

induces the maturation and differentiation of MKs by promoting the expression of

related transcription factors by regulating the SRC/MAPK signaling pathway.

As previously mentioned, VLZ is a 5-HT1A receptor agonist (Wang et al., 2015).

Previous studies have shown that the 5-HT tryptamine receptor signalingergic system

is involved in the process from embryonic to stem progenitor cells and subsequent

differentiation. However, most research has discovered and explored the effects of

5-HT2R on the hematopoietic system. The 5-HT receptor family is very large, and the

interreceptor interaction and signal networks are complex. The regulation of different

systems is both similar and different. To determine whether 5-HTR1A is involved in

the induction of MK differentiation, we first validated the expression of 5-HTR1A on

MKs. We first found that 5-HTR1A was expressed on Meg-01 cells and that VLZ

could directly bind to the receptor. To further determine whether 5-HTR1A activates

the MAPK signaling pathway and thus induces MK differentiation, we blocked the

receptor with its antagonist WAY-100635. As expected, we found that 5-HTR1A

inhibition completely abolished the induction of MK differentiation by VLZ and was

accompanied by a decrease in the expression of the downstream signaling pathways

RAS and ERK. Collectively, the data suggest that 5-HTR1A mediates the MK

differentiation and maturation of the VLZ and that this effect is accomplished by

5-HTR1A through regulation of the MAPK (RAS/ERK) signaling pathway. In

conclusion, we showed that VLZ stimulates MK differentiation and platelet formation

through the 5-HTR1A/SRC/MAPK pathway. Our findings highlight the importance of

5-HT receptor signaling for promoting hematopoiesis and provide new insights into

the underlying mechanism of VLZ.

Taken together, our study elucidated the effects and mechanisms of VLZ in the

treatment of thrombocytopenia through a combination of computational analysis and

experimental validation. We revealed a previously unknown interplay in which the expression of 5-HTR1A on MKs and the regulation of the SRC/RAS/MEK/ERK signaling pathway are involved in the effects of VLZ. Our study shows the therapeutic effects and potential mechanisms of VLZ in thrombocytopenia and enriches the study of the mechanisms of 5-HT receptor signalingergic regulation of the hematopoietic spectrum.

## Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (82074129, 82374073 and 82004073), Sichuan Science and Technology Program (2022ZYD0087, 2022YFS0607, 22ZYZYTS0191, 2022YFS0614 and 2022JDJQ0061), Joint Project of Luzhou Municipal People's Government and Southwest Medical University (2020LZXNYDZ03 and 2020LZXNYDP01).

## References

Ahmad HM, Muiwo P, Muthuswami R ,Bhattacharya A (2017) **FosB regulates expression of miR-22 during PMA induced differentiation of K562 cells to megakaryocytes** *Biochimie* **133:** 1-6.
https://doi.org/10.1016/j.biochi.2016.11.005

Amireault P, Hatia S, Bayard E, Bernex F, Collet C, Callebert J, Launay JM, Hermine O, Schneider E, Mallet J, Dy M ,Côté F (2011) **Ineffective erythropoiesis with reduced red blood cell survival in serotonin-deficient mice** *Proc Natl Acad Sci U S A* **108:** 13141-13146.
https://doi.org/10.1073/pnas.1103964108

Anderson NG, Maller JL, Tonks NK ,Sturgill TW (1990) **Requirement for integration of signals from two distinct phosphorylation pathways for activation of MAP kinase** *Nature* **343:** 651-653.
https://doi.org/10.1038/343651a0

Bannigan P, Aldeghi M, Bao Z, Häse F, Aspuru-Guzik A ,Allen C (2021) **Machine learning directed drug formulation development** *Adv Drug Deliv Rev* **175:** 113806.
https://doi.org/10.1016/j.addr.2021.05.016

Barnes NM, Ahern GP, Becamel C, Bockaert J, Camilleri M, Chaumont-Dubel S,

Claeysen S, Cunningham KA, Fone KC, Gershon M, Di Giovanni G, Goodfellow NM, Halberstadt AL, Hartley RM, Hassaine G, Herrick-Davis K, Hovius R, Lacivita E, Lambe EK, Leopoldo M, Levy FO, Lummis SCR, Marin P, Maroteaux L, Mccreary AC, Nelson DL, Neumaier JF, Newman-Tancredi A, Nury H, Roberts A, Roth BL, Roumier A, Sanger GJ, Teitler M, Sharp T, Villalón CM, Vogel H, Watts SW ,Hoyer D (2021) **International Union of Basic and Clinical Pharmacology. CX. Classification of Receptors for 5-hydroxytryptamine; Pharmacology and Function** *Pharmacol Rev* 73**:** 310-520.
https://doi.org/10.1124/pr.118.015552

Chen Z, Hu M ,Shivdasani RA (2007) **Expression analysis of primary mouse megakaryocyte differentiation and its application in identifying stage-specific molecular markers and a novel transcriptional target of NF-E2** *Blood* **109:** 1451-1459.
https://doi.org/10.1182/blood-2006-08-038901

Crispino JD (2022) **Introduction to a review series on megakaryopoiesis and platelet production** *Blood* **139:** 3227.
https://doi.org/10.1182/blood.2022016023

Crist SA, Elzey BD, Ahmann MT ,Ratliff TL (2013) **Early growth response-1 (EGR-1) and nuclear factor of activated T cells (NFAT) cooperate to mediate CD40L expression in megakaryocytes and platelets** *J Biol Chem* **288:** 33985-33996.
https://doi.org/10.1074/jbc.M113.511881

De Kock L ,Freson K (2020) **The (Patho)Biology of SRC Kinase in Platelets and Megakaryocytes** *Medicina (Kaunas)* 56.
https://doi.org/10.3390/medicina56120633

Della Rocca GJ, Mukhin YV, Garnovskaya MN, Daaka Y, Clark GJ, Luttrell LM, Lefkowitz RJ ,Raymond JR (1999) **Serotonin 5-HT1A receptor-mediated Erk activation requires calcium/calmodulin-dependent receptor endocytosis** *J Biol Chem* **274:** 4749-4753.
https://doi.org/10.1074/jbc.274.8.4749

Duross AN, Landry MR, Thomas CR, Jr., Neufeld MJ ,Sun C (2021) **Fucoidan-coated nanoparticles target radiation-induced P-selectin to enhance chemoradiotherapy in murine colorectal cancer** *Cancer Lett* **500:** 208-219.
https://doi.org/10.1016/j.canlet.2020.11.021

Erspamer V ,Asero B (1952) **Identification of enteramine, the specific hormone of the enterochromaffin cell system, as 5-hydroxytryptamine** *Nature* **169:** 800-801.
https://doi.org/10.1038/169800b0

904 Gauer JS, Duval C, Xu RG, Macrae FL, Mcpherson HR, Tiede C, Tomlinson D,
905 Watson SP ,Ariëns RAS (2023) **Fibrin-glycoprotein VI interaction increases**
906 **platelet procoagulant activity and impacts clot structure** *J Thromb Haemost* **21:**
907 667-681.
908 https://doi.org/10.1016/j.jtha.2022.09.004
909
910 Ghanima W, Cooper N, Rodeghiero F, Godeau B ,Bussel JB (2019) **Thrombopoietin**
911 **receptor agonists: ten years later** *Haematologica* **104:** 1112-1123.
912 https://doi.org/10.3324/haematol.2018.212845
913
914 Hsin KY, Ghosh S ,Kitano H (2013) **Combining machine learning systems and**
915 **multiple docking simulation packages to improve docking prediction reliability**
916 **for network pharmacology** *PLoS One* **8:** e83922.
917 https://doi.org/10.1371/journal.pone.0083922
918
919 Huang H, Woo AJ, Waldon Z, Schindler Y, Moran TB, Zhu HH, Feng GS, Steen
920 H ,Cantor AB (2012) **A Src family kinase-Shp2 axis controls RUNX1 activity in**
921 **megakaryocyte and T-lymphocyte differentiation** *Genes Dev* **26:** 1587-1601.
922 https://doi.org/10.1101/gad.192054.112
923
924 Jenkins JM, Williams D, Deng Y, Uhl J, Kitchen V, Collins D ,Erickson-Miller CL
925 (2007) **Phase 1 clinical study of eltrombopag, an oral, nonpeptide thrombopoietin**
926 **receptor agonist** *Blood* **109:** 4739-4741.
927 https://doi.org/10.1182/blood-2006-11-057968
928
929 Jimenez K, Khare V, Evstatiev R, Kulnigg-Dabsch S, Jambrich M, Strobl H ,Gasche
930 C (2015) **Increased expression of HIF2α during iron deficiency-associated**
931 **megakaryocytic differentiation** *J Thromb Haemost* **13:** 1113-1127.
932 https://doi.org/10.1111/jth.12884
933
934 Johansson MW, Han ST, Gunderson KA, Busse WW, Jarjour NN ,Mosher DF (2012)
935 **Platelet activation, P-selectin, and eosinophil β1-integrin activation in asthma** *Am*
936 *J Respir Crit Care Med* **185:** 498-507.
937 https://doi.org/10.1164/rccm.201109-1712OC
938
939 Kanaji T, Vo MN, Kanaji S, Zarpellon A, Shapiro R, Morodomi Y, Yuzuriha A, Eto K,
940 Belani R, Do MH, Yang XL, Ruggeri ZM ,Schimmel P (2018) **Tyrosyl-tRNA**
941 **synthetase stimulates thrombopoietin-independent hematopoiesis accelerating**
942 **recovery from thrombocytopenia** *Proc Natl Acad Sci U S A* **115:** E8228-e8235.
943 https://doi.org/10.1073/pnas.1807000115
944
945 Kiang JG, Zhai M, Liao PJ, Bolduc DL, Elliott TB ,Gorbunov NV (2014) **Pegylated**
946 **G-CSF inhibits blood cell depletion, increases platelets, blocks splenomegaly, and**
947 **improves survival after whole-body ionizing irradiation but not after irradiation**

948 **combined with burn** *Oxid Med Cell Longev* **2014:** 481392.
949 https://doi.org/10.1155/2014/481392
950
951 Kim M, Civin CI ,Kingsbury TJ (2019) **MicroRNAs as regulators and effectors of**
952 **hematopoietic transcription factors** *Wiley Interdiscip Rev RNA* **10:** e1537.
953 https://doi.org/10.1002/wrna.1537
954
955 Kong L, Chen Q, Campbell F, Snaar-Jagalska E ,Kros A (2020) **Light-Triggered**
956 **Cancer Cell Specific Targeting and Liposomal Drug Delivery in a Zebrafish**
957 **Xenograft Model** *Adv Healthc Mater* **9:** e1901489.
958 https://doi.org/10.1002/adhm.201901489
959
960 Lebois M, Dowling MR, Gangatirkar P, Hodgkin PD, Kile BT, Alexander
961 WS ,Josefsson EC (2016) **Regulation of platelet lifespan in the presence and**
962 **absence of thrombopoietin signaling** *J Thromb Haemost* **14:** 1882-1887.
963 https://doi.org/10.1111/jth.13397
964
965 Lien TS, Chan H, Sun DS, Wu JC, Lin YY, Lin GL ,Chang HH (2021) **Exposure of**
966 **Platelets to Dengue Virus and Envelope Protein Domain III Induces Nlrp3**
967 **Inflammasome-Dependent Platelet Cell Death and Thrombocytopenia in Mice**
968 *Front Immunol* **12:** 616394.
969 https://doi.org/10.3389/fimmu.2021.616394
970
971 Limb JK, Yoon S, Lee KE, Kim BH, Lee S, Bae YS, Jhon GJ ,Kim J (2009)
972 **Regulation of megakaryocytic differentiation of K562 cells by FosB, a member of**
973 **the Fos family of AP-1 transcription factors** *Cell Mol Life Sci* **66:** 1962-1973.
974 https://doi.org/10.1007/s00018-009-8775-5
975
976 Lin J, Zeng J, Liu S, Shen X, Jiang N, Wu YS, Li H, Wang L ,Wu JM (2021) **DMAG,**
977 **a novel countermeasure for the treatment of thrombocytopenia** *Mol Med* **27:** 149.
978 https://doi.org/10.1186/s10020-021-00404-1
979
980 Liu T, Gu J, Yuan Y, Yang Q, Zheng PF, Shan C, Wang F, Li H, Xie XQ, Chen
981 XH ,Ouyang Q (2022) **Discovery of a pyrano[2,3-b]pyridine derivative YX-2102**
982 **as a cannabinoid receptor 2 agonist for alleviating lung fibrosis** *J Transl Med* **20:**
983 565.
984 https://doi.org/10.1186/s12967-022-03773-1
985
986 Lucas D (2021) **Structural organization of the bone marrow and its role in**
987 **hematopoiesis** *Curr Opin Hematol* **28:** 36-42.
988 https://doi.org/10.1097/moh.0000000000000621
989
990 Lv J, Wang L, Gao Y, Ding YQ ,Liu F (2017) **5-hydroxytryptamine synthesized in**
991 **the aorta-gonad-mesonephros regulates hematopoietic stem and progenitor cell**

survival *J Exp Med* **214:** 529-545.
https://doi.org/10.1084/jem.20150906

Machlus KR ,Italiano JE, Jr. (2013) **The incredible journey: From megakaryocyte development to platelet formation** *J Cell Biol* **201:** 785-796.
https://doi.org/10.1083/jcb.201304054

Machlus KR, Johnson KE, Kulenthirarajan R, Forward JA, Tippy MD, Soussou TS, El-Husayni SH, Wu SK, Wang S, Watnick RS, Italiano JE, Jr. ,Battinelli EM (2016) **CCL5 derived from platelets increases megakaryocyte proplatelet formation** *Blood* **127:** 921-926.
https://doi.org/10.1182/blood-2015-05-644583

Marini I, Uzun G, Jamal K ,Bakchoul T (2022) **Treatment of drug-induced immune thrombocytopenias** *Haematologica* **107:** 1264-1277.
https://doi.org/10.3324/haematol.2021.279484

Mathew J, Pang CK, Luo M ,Leong WH (2018) **Classification of Imbalanced Data by Oversampling in Kernel Space of Support Vector Machines** *IEEE Trans Neural Netw Learn Syst* **29:** 4065-4076.
https://doi.org/10.1109/tnnls.2017.2751612

Matsuda M, Imaoka T, Vomachka AJ, Gudelsky GA, Hou Z, Mistry M, Bailey JP, Nieport KM, Walther DJ, Bader M ,Horseman ND (2004) **Serotonin regulates mammary gland development via an autocrine-paracrine loop** *Dev Cell* **6:** 193-203.
https://doi.org/10.1016/s1534-5807(04)00022-x

Mccormack PL (2015) **Vilazodone: a review in major depressive disorder in adults** *Drugs* **75:** 1915-1923.
https://doi.org/10.1007/s40265-015-0490-y

Moiseiwitsch JR ,Lauder JM (1995) **Serotonin regulates mouse cranial neural crest migration** *Proc Natl Acad Sci U S A* **92:** 7182-7186.
https://doi.org/10.1073/pnas.92.16.7182

Mouthon MA, Van Der Meeren A, Gaugler MH, Visser TP, Squiban C, Gourmelon P ,Wagemaker G (1999) **Thrombopoietin promotes hematopoietic recovery and survival after high-dose whole body irradiation** *Int J Radiat Oncol Biol Phys* **43:** 867-875.
https://doi.org/10.1016/s0360-3016(98)00477-5

Nakamura T, Miyakawa Y, Miyamura A, Yamane A, Suzuki H, Ito M, Ohnishi Y, Ishiwata N, Ikeda Y ,Tsuruzoe N (2006) **A novel nonpeptidyl human c-Mpl**

activator stimulates human megakaryopoiesis and thrombopoiesis *Blood* **107:** 4300-4307.

https://doi.org/10.1182/blood-2005-11-4433

Ninos JM, Jefferies LC, Cogle CR ,Kerr WG (2006) **The thrombopoietin receptor, c-Mpl, is a selective surface marker for human hematopoietic stem cells** *J Transl Med* **4:** 9.

https://doi.org/10.1186/1479-5876-4-9

Noetzli LJ, French SL ,Machlus KR (2019) **New Insights Into the Differentiation of Megakaryocytes From Hematopoietic Progenitors** *Arterioscler Thromb Vasc Biol* **39:** 1288-1300.

https://doi.org/10.1161/atvbaha.119.312129

Poston JN ,Gernsheimer TB (2019) **Glucocorticoids promote response to thrombopoietin-receptor agonists in refractory ITP: a case series** *Int J Hematol* **110:** 255-259.

https://doi.org/10.1007/s12185-019-02638-6

Provan D ,Semple JW (2022) **Recent advances in the mechanisms and treatment of immune thrombocytopenia** *EBioMedicine* **76:** 103820.

https://doi.org/10.1016/j.ebiom.2022.103820

Raghuwanshi S, Dahariya S, Sharma DS, Kovuru N, Sahu I ,Gutti RK (2020) **RUNX1 and TGF-β signaling cross talk regulates Ca(2+) ion channels expression and activity during megakaryocyte development** *Febs j* **287:** 5411-5438.

https://doi.org/10.1111/febs.15329

Rosignoli S ,Paiardini A (2022) **DockingPie: a consensus docking plugin for PyMOL** *Bioinformatics* **38:** 4233-4234.

https://doi.org/10.1093/bioinformatics/btac452

Roux PP ,Blenis J (2004) **ERK and p38 MAPK-activated protein kinases: a family of protein kinases with diverse biological functions** *Microbiol Mol Biol Rev* **68:** 320-344.

https://doi.org/10.1128/mmbr.68.2.320-344.2004

Ruan GY, Ye LX, Lin JS, Lin HY, Yu LR, Wang CY, Mao XD, Zhang SH ,Sun PM (2023) **An integrated approach of network pharmacology, molecular docking, and experimental verification uncovers kaempferol as the effective modulator of HSD17B1 for treatment of endometrial cancer** *J Transl Med* **21:** 204.

https://doi.org/10.1186/s12967-023-04048-z

Scharf RE (2012) **Drugs that affect platelet function** *Semin Thromb Hemost* **38:**

865-883.
https://doi.org/10.1055/s-0032-1328881

Short C, Lim HK, Tan J ,O'neill HC (2019) **Targeting the Spleen as an Alternative Site for Hematopoiesis** *Bioessays* **41:** e1800234.
https://doi.org/10.1002/bies.201800234

Singer CA, Lontay B, Unruh H, Halayko AJ ,Gerthoffer WT (2011) **Src mediates cytokine-stimulated gene expression in airway myocytes through ERK MAPK** *Cell Commun Signal* **9:** 14.
https://doi.org/10.1186/1478-811x-9-14

Skurikhin EG, Dygai AM, Suslov NI, Provalova NV, Zyuz'kov GN ,Gol'dberg ED (2001) **Role of the central nervous system in hemopoiesis regulation during experimental neuroses** *Bull Exp Biol Med* **131:** 33-37.
https://doi.org/10.1023/a:1017522411119

Spohn SN ,Mawe GM (2017) **Non-conventional features of peripheral serotonin signalling - the gut and beyond** *Nat Rev Gastroenterol Hepatol* **14:** 412-420.
https://doi.org/10.1038/nrgastro.2017.51

Tkaczynski E, Arulselvan A, Tkaczynski J, Avery S, Xiao L, Torok-Storb B, Abrams K, Rao NV, Johnson G, Kennedy TP, Poncz M ,Lambert MP (2018) **2-O, 3-O desulfated heparin mitigates murine chemotherapy- and radiation-induced thrombocytopenia** *Blood Adv* **2:** 754-761.
https://doi.org/10.1182/bloodadvances.2017013672

Van Den Oudenrijn S, Bruin M, Folman CC, Peters M, Faulkner LB, De Haas M ,Von Dem Borne AE (2000) **Mutations in the thrombopoietin receptor, Mpl, in children with congenital amegakaryocytic thrombocytopenia** *Br J Haematol* **110:** 441-448.
https://doi.org/10.1046/j.1365-2141.2000.02175.x

Van Dijk J, Bompard G, Cau J, Kunishima S, Rabeharivelo G, Mateos-Langerak J, Cazevieille C, Cavelier P, Boizet-Bonhoure B, Delsert C ,Morin N (2018) **Microtubule polyglutamylation and acetylation drive microtubule dynamics critical for platelet formation** *BMC Biol* **16:** 116.
https://doi.org/10.1186/s12915-018-0584-6

Wang L, Liu S, Luo J, Mo Q, Ran M, Zhang T, Li X, Zou W, Mei Q, Chen J, Yang J, Zeng J, Huang F, Wu A, Zhang C ,Wu J (2022) **Targeting TPO-independent strategy in the discovery of a novel megakaryocytopoiesis inducer DMAG for treatment of thrombocytopenia** *Haematologica*.
https://doi.org/10.3324/haematol.2022.282209

Wang SM, Han C, Lee SJ, Patkar AA, Masand PS ,Pae CU (2015) **Vilazodone for the treatment of major depressive disorder: focusing on its clinical studies and mechanism of action** *Psychiatry Investig* **12:** 155-163.
https://doi.org/10.4306/pi.2015.12.2.155

Xu Z, Xu YJ, Hao YN, Ren LJ, Zhang ZB, Xu X, Cao BY, Dai KS, Zhu L, Fang Q, Kong Y ,Mao XL (2017) **A novel STAT3 inhibitor negatively modulates platelet activation and aggregation** *Acta Pharmacol Sin* **38:** 651-659.
https://doi.org/10.1038/aps.2016.155

Yang M, Li K, Ng PC, Chuen CK, Lau TK, Cheng YS, Liu YS, Li CK, Yuen PM, James AE, Lee SM ,Fok TF (2007) **Promoting effects of serotonin on hematopoiesis: ex vivo expansion of cord blood CD34+ stem/progenitor cells, proliferation of bone marrow stromal cells, and antiapoptosis** *Stem Cells* **25:** 1800-1806.
https://doi.org/10.1634/stemcells.2007-0048

Ye JY, Liang EY, Cheng YS, Chan GC, Ding Y, Meng F, Ng MH, Chong BH, Lian Q ,Yang M (2014) **Serotonin enhances megakaryopoiesis and proplatelet formation via p-Erk1/2 and F-actin reorganization** *Stem Cells* **32:** 2973-2982.
https://doi.org/10.1002/stem.1777

Zhang T, Mo Q, Jiang N, Wu Y, Yang X, Chen W, Li Q, Yang S, Yang J, Zeng J, Huang F, Huang Q, Luo J, Wu J ,Wang L (2023) **The combination of machine learning and transcriptomics reveals a novel megakaryopoiesis inducer, MO-A, that promotes thrombopoiesis by activating FGF1/FGFR1/PI3K/Akt/NF-κB signaling** *Eur J Pharmacol* **944:** 175604.
https://doi.org/10.1016/j.ejphar.2023.175604

Zheng C, Yang R, Han Z, Zhou B, Liang L ,Lu M (2008) **TPO-independent megakaryocytopoiesis** *Crit Rev Oncol Hematol* **65:** 212-222.
https://doi.org/10.1016/j.critrevonc.2007.11.003

Zhou XH, Cheng ZP, Lu M, Lin WY, Luo LL, Ming ZY ,Hu Y (2023) **Adiponectin receptor agonist AdipoRon modulates human and mouse platelet function** *Acta Pharmacol Sin* **44:** 356-366.
https://doi.org/10.1038/s41401-022-00943-1

## Ethics approval and consent to participate

This study was performed in line with the principles of the Declaration of Helsinki. The experimental animal ethics committee of Southwest Medical University was followed during all procedures on mice (License NO.20230716-002).

## Consent for publication

All authors agreed to publication.

## Availability of data and materials

All data of this study are included in this article or the supplementary materials. All data are available from the corresponding author upon request.

## Competing interests

The authors declare no conflicts of interest.

## Author information

These authors contributed equally to this work: Ling Zhou, Cheng-yang Ni and Rui-xue Liao.

### contributions

JMW and LZ conceived and designed the experiments and supervised all the research. DLQ, LW, and XGZ provided useful suggestions on the methodology. JSL and TAY constructed the drug screening model. LZ, CYN and RXL performed *in vitro* experiments. XX and MR constructed the transgenic zebrafish model. LZ, XQT, FHH, MH and RL constructed the mice model and carried out *in vivo* experiments. LZ and YWW performed the network pharmacological analysis. LZ, CYN and RXL analyzed the data and wrote the original manuscript. JMW, JSL and YWW revised the manuscript. All authors had full access to the data and approved the final version of the manuscript.

### Corresponding authors

Correspondence to Jian-ming Wu, Jie-si Luo or Yi-wei Wang.

**Supplementary information**

**Supplementary Figure 1**

Results of the CCK-8 assay and LDH assay.

**Supplementary Figure 2**

Peripheral blood counts and Flow cytometry analysis after treated with VLZ in normal mice.

**Supplementary Figure 3**

WBC counts.

**Supplementary Figure 4**

Toxicity evaluation of VLZ *in vivo*.

MK blast

Mature MK

Pro-MK

itga2b:eGFP

Transgenic zebrafish

Proplatelet MK

Platelet

4 Gy (IR)

Vascular

Thrombocytopenic mice

Vilazodone

5-HTR1A

Gα i/o

γ

β

Phosohorylation

SRC

RAS

Phosorylation

MEK

Phosorylation

ERK1/2

FOS

NF-E2

RUNX1

EGR1

Trascriptional activation of MKs differentiation and platelet production