Quantifying the strength of viral fitness tradeoffs between hosts: A meta-analysis of pleiotropic fitness effects

Xuechun 'May' Wang[1], Julia Muller[2], Mya McDowell[3] and David A. Rasmussen[1,4,*]

[1] Department of Entomology and Plant Pathology, North Carolina State University, Raleigh, NC, USA.
[2] Gillings School of Global Public Health, University of North Carolina Chapel Hill, Chapel Hill, NC, USA.
[3] Department of Biological Sciences, North Carolina State University, Raleigh, NC, USA.
[4] Bioinformatics Research Center, North Carolina State University, Raleigh, NC, USA.

*Corresponding author: David A. Rasmussn (drasmus@ncsu.edu)

Abstract

The range of hosts a given virus can infect is widely presumed to be limited by tradeoffs in fitness between alternative hosts. These fitness tradeoffs may arise naturally due to antagonistic pleiotropy if mutations that increase fitness in one host tend to decrease fitness in alternate hosts. While antagonistic pleiotropy underlies certain tradeoffs, there is a growing recognition that positive pleiotropy may be more common than previously appreciated. With positive pleiotropy, mutations have concordant fitness effects such that a beneficial mutation can simultaneously increase fitness in different hosts, providing a genetic mechanism by which selection can overcome fitness tradeoffs. How readily evolution can overcome fitness tradeoffs therefore depends on the overall distribution of mutational fitness effects between hosts, including the relative frequency of antagonistic versus positive pleiotropy. We therefore conducted a systematic meta-analysis of the pleiotropic fitness effects of viral mutations reported in different hosts. Our analysis indicates that the relative frequency of antagonistic versus positive pleiotropy may simply reflect the underlying frequency of beneficial and deleterious mutations in individual hosts. Given a mutation is beneficial in one host, the probability that it is deleterious in another host is roughly equal to the probability that any mutation is deleterious, suggesting there is no natural tendency towards antagonistic pleiotropy beyond that expected by chance. While both antagonistic and positive pleiotropy are common,

fitness effects are overall positively correlated between hosts and unconditionally beneficial mutations are not uncommon. The widespread prevalence of positive pleiotropy suggests that many fitness tradeoffs may be readily overcome by evolution given the right selection pressures.

Introduction

Viruses regularly jump between hosts and exhibit a tremendous capacity to adapt to novel host environments (1,2). Consequently, new viral pathogens regularly emerge on novel hosts and expand their host range (3). Nevertheless, many observations suggest that the host range of individual viruses is restricted, often to a set of closely related hosts (4–6). Likewise, when viruses are experimentally passaged on a single host, fitness tends to decline rapidly in alternative hosts not encountered during passaging (7–10). This seemingly paradoxical pattern of tremendous adaptive potential yet relatively restricted host ranges suggests that fitness tradeoffs may constrain the ability of viruses to adapt to multiple hosts simultaneously, thereby limiting the emergence of generalist viruses.

Evolutionary theory tends to emphasize the role of antagonistic pleiotropy in generating fitness tradeoffs between hosts (6,11,12). Broadly defined, pleiotropy refers to an individual gene or mutation affecting multiple phenotypic traits or the same trait in different environments (13,14). Antagonistic pleiotropy results when a given mutation has opposite effects on two traits, such as a viral mutation being beneficial in one host but deleterious in another. By contrast, positive pleiotropy results when a mutation has concordant effects across traits or environments. In viruses, there is substantial evidence for individual mutations having antagonistic or opposite fitness effects between hosts (11,15,16). For example, mutations in viral glycoproteins frequently exhibit antagonistic pleiotropy due to the need for these proteins to bind highly host-specific receptors during cell entry (17–20). Antagonistic pleiotropy therefore provides a simple genetic explanation for fitness tradeoffs: if most mutations have antagonistic effects then gaining the ability to infect a new host will come at the cost of becoming less well adapted to alternate hosts.

If all mutations had antagonistic effects on fitness between, then fitness tradeoffs would be absolute or "hard" in the sense that there would be no way for evolution to resolve tradeoffs since increasing fitness in one host would inevitably decrease fitness in an alternate host. Yet fitness tradeoffs are not always absolute and viruses often evolve to overcome tradeoffs. For

example, generalists with high fitness across multiple hosts often evolve when viruses are alternately transferred between different hosts, even though fitness on alternate hosts tends to decline when viruses are passaged exclusively on a single host (7–9,21,22). Moreover, viruses adapted to more than one host sometimes have increased fitness relative to specialists adapted to a single host, leading to what has been referred to as fitness trade-ups or no-cost generalists (see 23,24). These fitness trade-ups can arise due to unconditionally beneficial mutations increasing fitness in more than one host simultaneously, a form of positive pleiotropy which may be more common than previously appreciated (16,25–27). Taken together, these observations suggest that many fitness tradeoffs may be "soft" in the sense that evolution could resolve the tradeoff given appropriate genetic variation and selection pressures. The strength of a fitness tradeoff in terms of how easy or difficult it will be for evolution to overcome will therefore largely depend on the relative frequency of antagonistic versus positive pleiotropy.

Given the central importance of pleiotropy to fitness tradeoffs and host range evolution, surprisingly little work has been done to quantify the overall distribution of fitness effects of mutations between hosts. We therefore conducted a systematic meta-analysis of published studies in which the fitness effects of individual viral mutations were estimated in two or more hosts, allowing us to quantify both the magnitude and sign of pleiotropy, including the frequency of antagonistic versus positive pleiotropy. We also included several viral and host features in our meta-analysis, allowing us to explore whether any features can predict the pleiotropic effects of mutations between different hosts. We find that fitness effects are overall positively correlated between hosts with positive pleiotropy being slightly more frequent than antagonistic pleiotropy. Perhaps more surprisingly, antagonistic pleiotropy is no more common than would be expected by chance given the frequency of beneficial versus deleterious mutations in individual hosts. We end by discussing the implications of these findings on the strength of fitness tradeoffs between hosts more generally.

Methods

Literature search

We searched for published studies reporting mutational fitness effects in viruses using Google Scholar and Web of Science with the search terms "virus* OR viral AND mutation* AND fitness". However, the vast majority of studies returned by this search only measured viral fitness in one host. We therefore added at least one of the following terms to our original search term:

"pleiotropy OR pleiotropic", "host range", "host alternation", "tradeoff". Systematic searches were performed originally in January 2021 and then again in April 2023. Study relevance was initially determined based on title and abstract, which resulted in 180 relevant publications. Relevant studies cited by these initially identified papers were subsequently added. After examining the full text of each study, 26 remained that provided relevant data and met all inclusion criteria. Supplemental Table S1 provides a complete list of all included studies. Our full dataset containing all extracted fitness effects and metadata is available at https://github.com/davidrasm/PleiotropyMetaAnalysis.

Study inclusion criteria

We only considered published studies which reported the fitness effects of individual mutations in DNA and RNA viruses. To be included, the fitness effects of a viral mutation had to be reported in at least two host environments. Different genotypes and ecotypes of the same host species were considered to be different hosts. Fitness effects determined both *in vivo* and using *in vitro* cell/tissue culture experiments were included, but we did not include fitness measured in extracellular environments. We only considered fitness effects determined in isogenic (single-mutant) backgrounds and did not include the fitness of any genotype with more than a single mutation relative to the wildtype. Because it is often unclear whether mutations were naturally occurring or experimentally derived, we included both and do not differentiate between the two. We only included studies that reported fitness in terms of viral replication or growth rates and excluded studies that used less direct proxies of fitness such as host virulence or disease symptoms.

Quantifying and standardizing fitness effects

In order for the fitness effect of a mutation in a given host to be included, the fitness of the mutant genotype (w_{mut}) needed to be compared with the fitness of a wildtype/reference genotype (w_{ref}). We included fitness measurements from competition assays as well as experiments that determined the fitness of each genotype in isolation. If fitness effects were not reported as numerical values, we extracted data from figures using WebPlotDigitizer 4.4 (28) or, in one case, from gel band intensities using ImageJ (Rasband and contributors, National Institutes of Health, USA).

In order to meaningfully compare mutational fitness effects between hosts and across studies, we converted all reported values to a relative fitness ratio:

$$\omega = w_{mut} / w_{ref}.$$

We chose to standardize fitness effects as ratios rather than differences because fitness differences depend on comparing rates in units of time that are often not transparent, complicating direct comparisons of fitness (29,30). Since that magnitude of fitness effects varied considerably between studies, we further $\log_2$ transform ω.

However, many studies report fitness in terms of selection coefficients (s) instead of the growth rates w_{mut} and w_{ref}. In this case, there is a straightforward relationship with growth rates: $s = w_{mut} - w_{ref}$ (31,32). We therefore simply convert selection coefficients to fitness ratios using the reported fitness of either genotype, e.g. $\omega = (s + w_{ref}) / w_{ref}$. Alternatively, selection coefficients can be defined in terms of growth rates per discrete generation:

$$s_T = \ln(\lambda_{mut}/\lambda_{ref}) \approx \lambda_{mut}/\lambda_{ref} - 1,$$

where λ_{mut} and λ_{ref} are the per-generation, multiplicative growth rates of the mutant and reference respectively. In this case, s_T can be converted back to a fitness ratio using the relation $\omega = s_T + 1$ (30,32).

Converting viral titers/abundance to fitness

Often no direct measure of fitness is reported but can be computed from growth curves showing changes in viral titers or abundance through time. In this case, we computed growth rates w from the change in viral abundance through time:

$$w = \log(N_{t+1} / N_t) = \log(N_{t+1}) - \log(N_t),$$

where N_t and N_{t+1} are the viral titers at two consecutive time points. If viral titres were reported at multiple time points, we computed the growth rate over each time interval and then recorded the average growth rate. In the later case, we only considered time points in the exponential growth phase up to the point at which viral titers peaked.

Viral mutational features

We recorded metadata about each mutation including features that might explain observed fitness effects. This included the mutation substitution type (synonymous vs. nonsynonymous), virus genome type (ssDNA, (+)ssRNA, (-)ssRNA, dsRNA), viral protein functional class (non-structural, structural, accessory), viral envelope status (present/absent), host type (plant, animal, bacteria), viral genome size and whether or not viral protein is involved in host-cell binding. Hereafter we refer to all these as mutational features for simplicity. Additionally, we recorded the median estimated divergence time between each unique pair of hosts reported on timetree.org (33).

Quantifying pleiotropy

We differentiate between the magnitude and the sign of pleiotropy, following the terminology used to classify different types of epistasis (14,34). The sign of pleiotropy refers to whether a mutation has concordant (positive pleiotropy) or opposite (antagonistic pleiotropy) fitness effects between pairs of hosts. Note that our definition of positive pleiotropy encompasses all mutations with equal sign effects, including unconditionally beneficial and unconditionally deleterious mutations. The magnitude of pleiotropy refers to the absolute difference in fitness effects between two hosts: $m = |\omega_{Host1} - \omega_{Host2}|$. The sign and the magnitude of pleiotropy were computed for every unique pair of hosts in which fitness effects for a given mutation were available.

Statistical analysis

The relationship between mutational fitness effects and mutational features were explored using different linear regression models in *statsmodels* version 0.14 (35). We used a mixed effect linear model to test for significant associations between mutational features (fixed effects) and fitness effects in single hosts. To account for uncontrolled differences in how fitness was measured between studies, we allowed mean fitness effects to randomly vary between studies by including random intercepts for each study in the mixed model. A similar mixed linear model was used to model the relationship between mutational features and the magnitude of pleiotropy. We used logistic regression to test for significant associations between mutational features and the sign of pleiotropy (positive or antagonistic). For the latter, we used standard as opposed to mixed logistic models as the mixed models did not converge. Finally, we tested whether the observed distribution of pleiotropic effects grouped by category was different from what would be expected given the observed distribution of fitness effects in single hosts using a Pearson's chi-squared test.

Results

We identified 26 studies in which the fitness effects of individual viral mutations were experimentally determined in at least two host environments (Supplemental Table S1). In all, our data set includes fitness effects for 153 mutations in 20 unique viruses and 34 unique hosts. The resulting data set allows us to make 681 pairwise comparisons of fitness effects between hosts in order to quantify the overall distribution of pleiotropic effects. In order to standardize fitness effects across studies, we quantify the relative fitness effect of a mutation in each host as $\omega = log_2(w_{mut} / w_{ref})$, where w_{mut} and w_{ref} are the fitness of the mutant and wildtype virus, respectively. A neutral mutation therefore has a relative fitness effect $\omega = 0$, a mutation that reduces fitness by half has $\omega = -1$ and a mutation that doubles fitness will have $\omega = 1$.

Distribution of mutational fitness effects in single hosts

As expected from previous work characterizing the distribution of fitness effects (DFE) of viral mutations in single hosts (29,36,37), most non-lethal mutations are nearly neutral with a prominent peak at $\omega = 0$, but the distribution is skewed towards deleterious effects (Figure 1). Out of 457 total fitness effects, 66.8% are deleterious in individual hosts. Averaged over all hosts, the mean fitness effect is -0.205 (95% CI: -0.306:-0.102), which translates to a 13.6% decrease in fitness on a linear scale. Mutations in bacteriophages are on average substantially more deleterious than mutations in plant and animal viruses (Figure 1A). As a result, mutations in single-stranded DNA viruses, which are largely bacteriophages, are more deleterious than mutations in positive (+) or negative (-) single-stranded RNA viruses (Figure 1B). Likewise, mutations in structural proteins are substantially more deleterious than mutations in non-structural and accessory proteins (Figure 1C). Fitness effects did not differ significantly between nonsynonymous and synonymous mutations (Figure 1D), nor between enveloped and non-enveloped viruses (Figure 1E). There was a slight tendency for mutations in viruses with larger genomes to be more beneficial (Figure 1F).

Figure 1: Distribution of mutational fitness effects in single hosts. Fitness effects ω are plotted on a log$_2$ scale. The fitness effects are grouped by: (A) host type, (B) viral genome type, (C) viral protein functional class, (D) codon effect, and (E) viral envelope status. (F) The relationship between viral genome size and mutational fitness effects.

To account for potential confounding effects among mutational features as well as uncontrolled differences between studies, we estimated the impact of each feature on mutational fitness effects using a mixed effect linear model. Viral features were treated as fixed effects while allowing mean fitness effects (intercepts) to randomly vary by study. Only two features were estimated to significantly impact fitness effects: host type with bacteriophages having more

deleterious fitness effects and viral genome length with mutations becoming slightly more beneficial in viruses with larger genomes (Table 1). The intraclass correlation coefficient (ICC) among studies is 0.341, indicating fitness effects measured within each study tended to be relatively similar due to within-study correlations.

Table 1: Mutational fitness effects in single hosts grouped by viral features. Regression coefficients, P-values and 95% CIs are estimated under a mixed effect linear model.

Variable	N	Group mean	Estimated coefficient	P-value	95% CI
Intercept	457	-	-3.639	0.65	-8.043 : 0.766
Synonymous	121	-0.37	-	-	-
Nonsynonymous	336	-0.14	-0.063	0.644	-0.329 : 0.204
Accessory	83	0.25	-	-	-
Non-structural	156	-0.13	-0.19	0.225	-0.497 : 0.117
Structural	218	-0.42	0.368	0.108	-0.081 : 0.816
dsRNA	36	-0.69	-	-	-
ssDNA	112	-1.01	0.958	0.462	-1.593 : 3.509
(+)ssRNA	243	0.12	0.14	0.889	-1.834 : 2.115
(-)ssRNA	46	0.26	-1.299	0.211	-3.336 : 0.738
Animal host	119	0.13	-	**-**	-
Bacteria host	139	-0.98	-1.761	**0.024**	-3.288 : -0.235
Plant host	199	0.14	-0.659	0.478	-2.480 : 1.162
Enveloped	144	-0.06	-	-	-
Non-enveloped	313	-0.27	1.289	0.236	-0.841 : 3.419
Viral genome size	457	-	0.329	**0.032**	0.028 : 0.630

Joint distribution of fitness effects between hosts

We compared the fitness effects of each mutation between every unique pair of hosts for which fitness data was available. These pairwise comparisons of fitness effects allowed us to characterize the pleiotropic effects of mutations in terms of the pairwise or joint distribution of fitness effects between hosts (Figure 2). We further characterize the pleiotropic effects of mutations in terms of both the magnitude and the sign of pleiotropy in the sections below.

Figure 2: Joint distribution of mutational fitness effects between pairs of hosts. The underlying distribution is identical in all plots but mutations are colored by: (A) host type, (B) viral genome type, (C) codon effect, (D) protein functional class, (E) viral envelope status and (F) whether the mutation occurs in a protein involved in host cell binding. Fitness effects ω in each host are $\log_2$ transformed.

Magnitude of pleiotropy

We define the magnitude of pleiotropy as the absolute difference in fitness effects between hosts, such that the magnitude is larger the more different the fitness effects are between two hosts, regardless of the sign of the fitness effects. Because we expect fitness effects to differ more between more phylogenetically divergent hosts (1,38), we first considered the relationship between host divergence times and the magnitude of pleiotropy (Figure 3). When considered among all pairs of hosts, divergence times do not have a significant effect on the magnitude of pleiotropy (β_{div} = -0.005; 95% CI = -0.052 : 0.042; p=0.832). However, when we considered interactions between host type and divergence times, the magnitude of pleiotropy increases strongly and significantly with divergence times for plants ($\beta_{div\,X\,plant}$ = 0.985: 95% CI = 0.786 : 1.184; p<0.001), weakly but not significantly for bacteria ($\beta_{div\,X\,bacteria}$ = 0.109; 95% CI = -0.013 : 0.23; p=0.079), but not for animals ($\beta_{div\,X\,animal}$ = 0.016, 95% CI = -0.029 : 0.06; p=0.49).

Figure 3: Magnitude of pleiotropic fitness effects between pairs of hosts regressed against host divergence times in millions of years. Colored lines show the estimated regression coefficients (slope) for each host type.

We also considered how the magnitude of pleiotropy depends on several other mutational features (Table 2). The magnitude is larger between plant hosts than between animal or bacteria hosts, indicating that fitness effects tend to vary more between plant hosts than other types of hosts (Figure 4A). The magnitude is also larger for (-)ssRNA and (+)ssRNA viruses than for ssDNA viruses (Figure 4B). The magnitude does not significantly differ between synonymous

and nonsynonymous mutations (Figure 4C), but the magnitude is slightly larger for mutations in non-structural and accessory proteins than in structural proteins (Figure 4D). The magnitude is significantly larger for mutations in non-enveloped viruses than enveloped viruses and also increases slightly with viral genome size (Figure 4E-F).

After controlling for confounding effects and uncontrolled differences between studies using a mixed effect model, no feature was estimated to have a significant effect on the magnitude of pleiotropy. However, when we included interactions between host type and host divergence times in the model, divergence times between plant hosts have a strong and significant effect on the magnitude of pleiotropy (Table 2). Thus of all the features we considered only divergence times between plant hosts strongly predict the magnitude of pleiotropy. Under the mixed effect model, the intraclass correlation coefficient among studies is low (ICC=0.046), indicating the magnitude of pleiotropic effects measured within each study is not highly correlated within studies.

Table 2: Magnitude of pleiotropic fitness effects grouped by viral features. Regression coefficients, P-values and 95% CIs are estimated under a mixed effect linear model.

Variable	N	Group mean	Estimated coefficient	P-value	95% CI
Intercept	681	-	-1.702	0.268	-4.710 : 1.306
Synonymous	183	0.92	0.13	0.245	-0.089 : 0.349
Nonsynonymous	498	0.93	-	-	-
Accessory	180	1.1	-	-	-
Non-structural	299	0.92	0.113	0.31	-0.105 : 0.331
Structural	202	0.76	0.287	0.117	-0.072 : 0.645
dsRNA	18	0.73	-	-	-
ssDNA	124	0.67	0.598	0.437	-0.909 : 2.105
(+)ssRNA	516	0.96	0.143	0.746	-0.722 : 1.007
(-)ssRNA	23	1.42	0.605	0.146	-0.211 : 1.421
Enveloped	129	0.72	-	-	-
Non-enveloped	552	0.97	0.021	0.566	-0.051 : 0.093
Divergence x Animal host	121	-	0.021	0.566	-0.051 : 0.093
Divergence x Bacteria host	133	-	0.181	0.057	-0.006 : 0.369
Divergence x Plant host	427	-	0.891	**<0.001**	0.649 : 1.134
Viral genome size	681	-	0.16	0.146	-0.056 : 0.375

Figure 4: Magnitude of pleiotropic fitness effects computed as the absolute difference in fitness effects between hosts. Fitness effects in each host were first $\log_2$ transformed, such that a magnitude of one reflects a 2X difference in fitness effects between hosts. Box plots show the median and quartiles of each categorical group while whiskers show the full range of observed values.

Sign of pleiotropy

Across all pairwise comparisons of fitness effects, positive pleiotropy is slightly more frequent (391/681 = 57.4%) than antagonistic pleiotropy. Likewise, mutational fitness effects are overall positively correlated between hosts (Pearson correlation coefficient ρ = 0.32). However, our definition of positive pleiotropy includes mutations that are both unconditionally beneficial and unconditionally deleterious. If we only consider pairwise comparisons with positive pleiotropic effects, the majority are unconditionally deleterious (297/391 = 75.9%). Nevertheless, 13.8% of

all pairwise comparisons are unconditionally beneficial, suggesting that mutations that increase fitness in more than one host are not uncommon.

We next tested whether the observed frequency of each type of sign pleiotropy (+/+, +/-, -/-) differed from what would be expected based on the number of beneficial and deleterious mutations observed in single hosts (Table 3). A Pearson chi-square test indicates the observed frequencies are consistent with what would be expected by chance if the pairwise fitness effects were randomly drawn from the distribution of fitness effects in single hosts (χ^2 = 7.12, p-value = 0.008). In other words, antagonistic and positive pleiotropy arise at roughly the frequency we would expect given the relative number of beneficial versus deleterious mutations observed in individual hosts. Given a mutation is beneficial in one host, it might be expected that the likelihood of the mutation being deleterious in another host would increase with the size of the beneficial fitness effect in the first host. However, the odds of a mutation being beneficial in both hosts actually increases with the size of the beneficial fitness effect in the first host (β = 1.17; 95% CI = 1.07 : 1.29; p=0.001).

Table 3: Contingency table with the observed versus expected number of mutations in each pleiotropic class. All mutations with antagonistic pleiotropic effects (+/- and -/+) are grouped as the ordering of hosts is arbitrary.

		Host 2 Fitness Effect	
		Negative (-)	Positive (+)
Host 1 Fitness Effect	Positive (+)	Obsv = 290 (42.6%) Exp = 319 (46.9%)	Obsv = 94 (13.8%) Exp = 78 (11.5%)
	Negative (-)	Obsv = 297 (43.6%) Exp = 282 (41.4%)	Obsv = 290 (42.6%) Exp = 319 (46.9%)

We further explored whether certain types of mutations were more likely to have positive rather than antagonistic pleiotropic effects using logistic regression (Table 4). The odds of a mutation having positive pleiotropic effects are higher in ssDNA viruses, and by extension bacteriophages, but these mutations were nearly always unconditionally deleterious (Figure 2A-B). Synonymous mutations are no more likely to have positive pleiotropic effects than nonsynonymous mutations (Figure 2C). Although not statistically significant, mutations in structural proteins are more likely to have positive pleiotropic effects, but again most mutations in structural proteins are unconditionally deleterious (Figure 2D). Finally, mutations in

non-enveloped viruses are significantly less likely to exhibit positive pleiotropy than enveloped viruses (Figure 2E).

Table 4: Odds of a mutation having a positive pleiotropic effect. Odds ratios, p-values and 95% CI are estimated under a multivariate logistic regression model.

Variable	Frequency	Odds ratio	P-value	95% CI
Intercept	-	10.83	0.38	-
Synonymous	92/183	-	-	-
Nonsynonymous	284/498	0.82	0.36	0.54 : 1.25
Accessory	81/180	-	-	-
Non-structural	138/299	0.71	0.13	0.46 : 1.1
Structural	157/202	1.22	0.55	0.64 : 2.33
dsRNA	16/18	-	-	-
ssDNA	114/124	3.32	0.4	0.20 : 55.4
(+)ssRNA	236/516	0.29	0.19	0.04 : 1.80
(-)ssRNA	10/23	0.1	**0.006**	0.02 : 0.52
Enveloped	81/129	-	-	-
Non-enveloped	295/552	0.38	**0.03**	0.15 : 0.92
Viral genome size (kb)	-	0.98	0.9	0.67 : 1.41

Among mutations exhibiting positive pleiotropy, we further examined whether any features increase the odds of mutations being unconditionally beneficial rather than deleterious (Table 5). The odds of mutation being unconditionally beneficial are slightly higher for nonsynonymous than synonymous mutations. Mutations in (-)ssRNA viruses also have significantly higher odds of being unconditionally beneficial than in viruses with other genome types, but this was based on a small number of observations. On the other hand, mutations in both structural and non-structural proteins are more likely to be unconditionally deleterious than mutations in accessory proteins, although only significantly for non-structural proteins. To better understand what types of proteins tend to increase the odds of unconditional deleterious effects, we further divided proteins by their more specific molecular function (Supp. Figure S1) and whether or not the protein is involved in host cell-binding (Figure 2F). While no clear pattern emerged in terms of protein molecular function, mutations in structural proteins like glycoproteins that bind to host cells are significantly more likely to be unconditionally deleterious than mutations in proteins that do not bind to host cells.

Table 5: Odds of a mutation being unconditionally beneficial versus deleterious. Odds ratios, p-values and 95% CI are estimated under a multivariate logistic regression model.

Variable	Frequency	Odds ratio	P-value	95% CI
Intercept	-	0.05	0.56	-
Synonymous	11/92	-	-	-
Nonsynonymous	68/284	2.21	**0.048**	1.01 : 4.84
Accessory	31/81	-	-	-
Non-structural	32/138	0.34	**0.009**	0.15 : 0.77
Structural	16/157	0.44	0.203	0.12 : 1.56
dsRNA	1/16	-	-	-
ssDNA	7/114	5.81	0.47	0.05 : 710.5
(+)ssRNA	65/236	2.43	0.51	0.16 : 34.85
(-)ssRNA	6/10	19.18	**0.016**	1.73 : 212.1
Enveloped	27/81	-	-	-
Non-enveloped	52/295	0.34	0.18	0.07 : 1.65
Non-host cell binding	69/250	-	-	-
Host cell binding	10/126	0.08	**0.002**	0.02 : 0.41
Viral genome size (kb)	-	1.22	0.58	0.60 : 2.49

Discussion

Viral host ranges are widely believed to be constrained by fitness tradeoffs arising from antagonistic pleiotropy (6,11,12). But to our knowledge, no previous work has systematically surveyed the fitness effects of mutations in different hosts to quantify the relative frequency of antagonistic versus positive pleiotropy. Within individual hosts, our meta-analysis confirmed that the distribution of fitness effects (DFE) is skewed towards deleterious mutations with individual mutations reducing fitness by 13.6% on average. This is remarkably similar to previous estimates which estimated mutations reduce fitness 10.3 - 13.2% on average (30). Considering the pairwise or joint DFE between hosts, fitness effects are fairly strongly positively correlated between hosts and positive pleiotropy is actually slightly more common than antagonistic pleiotropy. While antagonistic pleiotropy may nevertheless constrain the ability of viruses to adapt to certain hosts, our study calls into question whether antagonistic pleiotropy by itself is frequent and strong enough to constrain viral host ranges more generally.

Perhaps our most surprising result is that antagonistic pleiotropy is no more frequent than would be expected by chance given the frequency of beneficial and deleterious mutations in individual

hosts. Because the DFE is skewed towards deleterious mutations in individual hosts, between hosts we expect a majority of mutations to be unconditionally deleterious, a moderate fraction to have antagonistic effects and very few to be unconditionally beneficial in both hosts. The observed distribution of pleiotropic fitness effects is remarkably similar to this expectation, suggesting that the sign of a mutation's fitness effect in one host is essentially independent of its fitness effect in another. This is consistent with earlier work on individual viruses showing the fitness effects of new mutations are positively correlated between hosts overall but the sign of fitness effects are highly unpredictable between hosts (16,25). This may also explain why we found few features of mutations, viruses or hosts that could reliably predict whether or not mutations will have antagonistic effects.

Given that there appears to be no natural tendency towards mutations having antagonistic effects between hosts, antagonistic pleiotropy may be less ubiquitous than previously thought. This may at first seem surprising; afterall viruses are obligate cellular parasites requiring many finely tuned molecular interactions between viral and host factors. However, even if fitness effects are host-specific, these effects may only vary by degree between hosts, impacting the magnitude of pleiotropy but not the sign. Many viral proteins may also interact with similar host factors in different hosts, especially if the hosts are closely related (39,40). As might be expected based on these considerations, we found that fitness effects are more similar between pairs of more phylogenetically related hosts, at least in bacteria and plant hosts. Furthermore, mutations in certain viral proteins may even be expected to have concordant fitness effects between hosts. For example, improving the processivity of RNA polymerases or the stability of structural proteins may increase fitness across a wide range of hosts (11,41). Likewise, in other microbial systems, positive pleiotropy appears more common than previously thought, including in bacteria growing on different carbon sources (42–44) and in different antibiotic environments (45–47). Thus, rather than being viewed as surprising, reports of positive pleiotropy should often be expected.

The more pertinent question may be whether the observed distribution of pleiotropic effects can prevent the evolution of generalist viruses adapted to multiple hosts. On longer timescales, purifying selection will remove most unconditionally deleterious mutations, such that it is ultimately the availability of beneficial mutations on the timescales relevant to adaptation in each host that determines whether generalists evolve. Since it is commonly observed that viruses do adapt to multiple hosts when rapidly passaged between alternate hosts (7–9,21,22), it appears

that the supply of unconditional beneficial mutations is often sufficient for generalists to evolve. Indeed, our meta-analysis revealed that unconditionally beneficial mutations are not especially rare (13.8% of all pairwise comparisons). However when viruses infrequently encounter an alternate host, antagonistic pleiotropy may be sufficiently common to constrain the evolution of generalists (48,49). In this case, beneficial mutations will be selected in the more frequently encountered host irrespective of their fitness effects in the alternate host. Thus, even if antagonistic pleiotropy is no more common than expected by chance, mutations fixed in the more common host will on average be deleterious in the rare host. The resulting accumulation of deleterious mutations can lead to the evolution of specialists adapted only to the more common host, as proposed under the mutation accumulation hypothesis (50,51). Thus, whether or not antagonistic pleiotropy is sufficient to constrain the evolution of generalist viruses may strongly depend on the frequency at which different hosts are encountered.

Interestingly, many studies report cases of both antagonistic pleiotropy constraining and positive pleiotropy promoting adaptation to different hosts, often in the same host-virus pair (10,11,26,52,53). Theory should therefore be able to explain why pleiotropy sometimes constrains adaptation and yet other times promotes the evolution of multi-host generalists. We propose a simple conceptual model for how pleiotropy may both constrain and promote evolution based on general observations of the DFE that should apply across many host-pathosystems (Figure 5). Similar to other recent work, we view fitness tradeoffs along a continuum from hard to soft based on how easy it would be for evolution to resolve or overcome a tradeoff (47,54). Hard or "rigid" tradeoffs reflect absolute constraints arising from conflicting structural or biophysical requirements in different hosts. In this case, viral populations may be evolving along a Pareto optimality front where fitness cannot be increased in one host without compromising fitness in an alternate host (55,56). At such a Pareto front, the biophysical constraints underlying the tradeoff may skew the distribution of fitness effects towards mutations with antagonistic and unconditionally deleterious effects (Figure 5A). Selection is therefore unable to resolve such tradeoffs because mutations that simultaneously increase fitness in both hosts are not accessible.

Figure 5: Conceptual model illustrating how the distribution of pleiotropic effects can lead to both hard and soft fitness tradeoffs between hosts. (A) Hard tradeoffs arise when absolute biophysical constraints, represented as a hypothetical Pareto front (curved line), prevent fitness from improving in one host without declining in the other host. In this case, the joint DFE (orange points) is shifted towards mutations with antagonistic or unconditionally deleterious effects. (B) Soft tradeoffs arise when a population lies behind a Pareto front, shifting the joint DFE (blue points) towards unconditionally beneficial mutations.

In contrast to hard tradeoffs, soft or "plastic" tradeoffs can easily be overcome by evolution. Soft tradeoffs can arise when a population lies behind a hypothetical Pareto front, possibly due to a lack of prior selection in one or both environments. In this case, the distribution of fitness effects may be shifted towards positive pleiotropy such that unconditionally beneficial mutations are accessible, allowing a population to rapidly adapt to both hosts without paying any apparent cost (Figure 5B). This provides a general mechanism whereby viruses could overcome fitness tradeoffs that is strongly supported by our knowledge of the DFE. In populations that are poorly adapted to their environment, the entire DFE shifts towards beneficial mutations which increase in both frequency and effect sizes (36,57–59). Likewise, if we consider two environments, the joint DFE may shift towards unconditionally beneficial mutations as a result of the marginal DFE in each host shifting towards beneficial effects (60).

The overall distribution of pleiotropic fitness effect we observe here, where fitness effects are overall positively correlated between hosts with many unconditionally beneficial mutations, implies that most fitness tradeoffs between hosts are soft and could thus be easily overcome. However, one caveat to this conclusion is that, because our meta-analysis combines fitness effects from many different viruses and hosts, it may not be representative of the DFE in any

virus-host pair *per se*, but rather reflect a mixture of the different distributions shown in Figure 5. Thus, the general occurrence of unconditionally beneficial mutations does not necessarily imply that beneficial mutations will be accessible in any specific host-virus system. Nevertheless, we conjecture that the overall distribution of pleiotropic fitness effects observed here is likely representative of the situation many viruses experience as they adapt to novel hosts. To viruses, novel hosts represent low-quality, stressful environments which tends to skew the DFE towards beneficial mutations (25,61). Many of these mutations may be beneficial across multiple hosts, allowing for fitness trade ups rather than tradeoffs. We therefore expect many tradeoffs between hosts to be soft whereas hard tradeoffs may only arise in situations where historical selection pressures have nearly optimized fitness in both hosts, as seen in some vector-borne viruses (15,18,62).

Finally, while theoretical models of fitness tradeoffs generally try to explain the evolutionary success of either specialists or generalists, in nature both specialist and generalist viruses co-exist and host ranges vary in breadth from single hosts to thousands of hosts spanning multiple kingdoms (63–65). This variability in host range breadth is only explicable if we consider the different selection pressures viruses experience alongside genetic factors like the DFE. For example, a plant virus vectored by a herbivorous insect may repeatedly experience selection to adapt to any plant host on which its vector next alights (66). Meanwhile, a phage specialized to infect an abundant bacterial host may experience effectively no selection to adapt to a less common alternate host. Thus, while the widespread prevalence of positive pleiotropy suggests many tradeoffs may be soft, the right historical selection pressures still need to be in place for viruses to expand their host range. A more fully developed theory of viral host range evolution would therefore consider how both biophysical constraints as well as past and current selection pressures shape the distribution of pleiotropic fitness effects and the strength of fitness tradeoffs between hosts.

References

1. Longdon B, Brockhurst MA, Russell CA, Welch JJ, Jiggins FM. The Evolution and Genetics of Virus Host Shifts. PLoS Pathog. 2014 Nov 6;10(11):e1004395–8.

2. Geoghegan JL, Duchêne S, Holmes EC. Comparative analysis estimates the relative frequencies of co-divergence and cross-species transmission within viral families. PLoS Pathog. 2017 Feb;13(2):e1006215.

3. Lloyd-Smith JO, George D, Pepin KM, Pitzer VE, Pulliam JRC, Dobson AP, et al. Epidemic dynamics at the human-animal interface. Science. 2009 Dec 4;326(5958):1362–7.

4. Dawson WO, Hilf ME. Host-range determinants of plant viruses. Annu Rev Plant Biol [Internet]. 1992; Available from: https://www.annualreviews.org/doi/pdf/10.1146/annurev.pp.43.060192.002523?casa_token =zr35YJ-PzJMAAAAA:h_qOIynFE4jJbMNkLV6uKH_0yCagzonTDp9kcoxOeahfAnX0ImoyQ 4ZgdBjJLdpGuSv6Alqak01bBQ

5. Longdon B, Hadfield JD, Webster CL, Obbard DJ, Jiggins FM. Host Phylogeny Determines Viral Persistence and Replication in Novel Hosts. PLoS Pathog. 2011 Sep 22;7(9):e1002260–9.

6. Bedhomme S, Hillung J, Elena SF. ScienceDirect Emerging viruses: why they are not jacks of all trades? Curr Opin Virol. 2015 Feb 1;10:1–6.

7. Weaver SC, Brault AC, Kang W, Holland JJ. Genetic and fitness changes accompanying adaptation of an arbovirus to vertebrate and invertebrate cells. J Virol. 1999 May;73(5):4316–26.

8. Kassen R. The experimental evolution of specialists, generalists, and the maintenance of diversity. J Evol Biol. 2002 Mar 25;15(2):173–90.

9. Bedhomme S, Lafforgue G, Elena SF. Multihost Experimental Evolution of a Plant RNA Virus Reveals Local Adaptation and Host-Specific Mutations. Mol Biol Evol. 2012 Apr 29;29(5):1481–92.

10. Visher E, Uricchio L, Bartlett L, DeNamur N, Yarcan A, Alhassani D, et al. The evolution of host specialization in an insect pathogen. Evolution. 2022 Oct;76(10):2375–88.

11. Duffy S, Turner PE, Burch CL. Pleiotropic costs of niche expansion in the RNA bacteriophage phi 6. Genetics. 2000 Dec;156(4):1465–70.

12. Elena SF, Agudelo-Romero P, Lalić J. The Evolution of Viruses in Multi-Host Fitness Landscapes. Open Virol J. 2009 Jan 1;3(1):1–6.

13. Wagner GP, Zhang J. The pleiotropic structure of the genotype–phenotype map: the evolvability of complex organisms. Nat Rev Genet. 2011 Feb 18;12(3):204–13.

14. Remold S. Understanding specialism when the jack of all trades can be the master of all. Proceedings of the Royal Society B: Biological Sciences. 2012 Nov 12;279(1749):4861–9.

15. Coffey LL, Vasilakis N, Brault AC, Powers AM, Tripet F, Weaver SC. Arbovirus evolution in vivo is constrained by host alternation. Proc Natl Acad Sci U S A. 2008 May 13;105(19):6970–5.

16. Lalić J, Cuevas JM, Elena SF. Effect of Host Species on the Distribution of Mutational Fitness Effects for an RNA Virus. PLoS Genetics. 2011 Nov 17;7(11):e1002378–12.

17. Rogers GN, Paulson JC, Daniels RS, Skehel JJ, Wilson IA, Wiley DC. Single amino acid substitutions in influenza haemagglutinin change receptor binding specificity. Nature. 1983;304(5921):76–8.

18. Greene Ivorlyne P., Wang Eryu, Deardorff Eleanor R., Milleron Rania, Domingo Esteban, Weaver Scott C. Effect of Alternating Passage on Adaptation of Sindbis Virus to Vertebrate and Invertebrate Cells. J Virol. 2005 Nov 15;79(22):14253–60.

19. Tsetsarkin KA, Vanlandingham DL, McGee CE, Higgs S. A single mutation in chikungunya virus affects vector specificity and epidemic potential. PLoS Pathog. 2007 Dec;3(12):e201.

20. Urbanowicz RA, McClure CP, Sakuntabhai A, Sall AA, Kobinger G, Müller MA, et al. Human Adaptation of Ebola Virus during the West African Outbreak. Cell. 2016 Nov 3;167(4):1079–85.e5.

21. Coffey LL, Vignuzzi M. Host Alternation of Chikungunya Virus Increases Fitness while Restricting Population Diversity and Adaptability to Novel Selective Pressures. J Virol. 2010 Dec 22;85(2):1025–35.

22. Smith-Tsurkan SD, Wilke CO, Novella IS. Incongruent fitness landscapes, not tradeoffs, dominate the adaptation of vesicular stomatitis virus to novel host types. J Gen Virol. 2010;91(Pt 6):1484–93.

23. Burmeister AR, Turner PE. Trading-off and trading-up in the world of bacteria-phage evolution. Curr Biol. 2020 Oct 5;30(19):R1120–4.

24. McGee LW, Aitchison EW, Caudle SB, Morrison AJ, Zheng L, Yang W, et al. Payoffs, not tradeoffs, in the adaptation of a virus to ostensibly conflicting selective pressures. PLoS Genet. 2014 Oct;10(10):e1004611.

25. Vale PF, Choisy M, Froissart R, Sanjuán R, Gandon S. The distribution of mutational fitness effects of phage φX174 on different hosts. Evolution. 2012 Nov;66(11):3495–507.

26. Moreno-Pérez MG, García-Luque I, Fraile A, García-Arenal F. Mutations That Determine Resistance Breaking in a Plant RNA Virus Have Pleiotropic Effects on Its Fitness That Depend on the Host Environment and on the Type, Single or Mixed, of Infection. J Virol. 2016 Sep 29;90(20):9128–37.

27. Ruark-Seward CL, Bonville B, Kennedy G, Rasmussen DA. Evolutionary dynamics of Tomato spotted wilt virus within and between alternate plant hosts and thrips. Sci Rep. 2020 Sep 25;10(1):15797.

28. Rohatgi A. Webplotdigitizer: Version 4.4. URL https://automeris io/WebPlotDigitizer.

29. Sanjuán R, Moya A, Elena SF. The distribution of fitness effects caused by single-nucleotide substitutions in an RNA virus. Proc Natl Acad Sci U S A. 2004 Jun 1;101(22):8396–401.

30. Sanjuán R. Mutational fitness effects in RNA and single-stranded DNA viruses: common patterns revealed by site-directed mutagenesis studies. Philos Trans R Soc Lond B Biol Sci. 2010 Jun 27;365(1548):1975–82.

31. Fisher SRA. The Genetical Theory of Natural Selection. Clarendon Press; 1930. 272 p.

32. Chevin LM. On measuring selection in experimental evolution. Biol Lett. 2011 Mar 11;7(2):210–3.

33. Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, et al. TimeTree 5: An Expanded Resource for Species Divergence Times. Mol Biol Evol [Internet]. 2022 Aug 6;39(8). Available from: http://dx.doi.org/10.1093/molbev/msac174

34. Weinreich DM, Watson RA, Chao L. Perspective: sign epistasis and genetic costraint on evolutionary trajectories. Evolution [Internet]. 2005; Available from: https://academic.oup.com/evolut/article-abstract/59/6/1165/6755653

35. Seabold S, Perktold J. Statsmodels: Econometric and statistical modeling with python. In: Proceedings of the 9th Python in Science Conference [Internet]. SciPy; 2010 [cited 2023 Oct 31]. Available from: https://pdfs.semanticscholar.org/3a27/6417e5350e29cb6bf04ea5a4785601d5a215.pdf

36. Eyre-Walker A, Keightley PD. The distribution of fitness effects of new mutations. Nat Rev Genet. 2007 Aug;8(8):610–8.

37. Domingo-Calap P, Cuevas JM, Sanjuán R. The fitness effects of random mutations in single-stranded DNA and RNA bacteriophages. PLoS Genet. 2009 Nov;5(11):e1000742.

38. Longdon B, Day JP, Alves JM, Smith SCL, Houslay TM, McGonigle JE, et al. Host shifts result in parallel genetic changes when viruses evolve in closely related species. PLoS Pathog. 2018 Apr 12;14(4):e1006951–14.

39. Long JS, Mistry B, Haslam SM, Barclay WS. Host and viral determinants of influenza A virus species specificity. Nature Publishing Group. 2018 Nov 27;23:1–15.

40. Damas J, Hughes GM, Keough KC, Painter CA, Persky NS, Corbo M, et al. Broad host range of SARS-CoV-2 predicted by comparative and structural analysis of ACE2 in vertebrates. Proc Natl Acad Sci U S A. 2020 Sep 8;117(36):22311–22.

41. Manz B, Schwemmle M, Brunotte L. Adaptation of Avian Influenza A Virus Polymerase in Mammals To Overcome the Host Species Barrier. J Virol. 2013 Jun 10;87(13):7200–9.

42. Ostrowski EA, Rozen DE, Lenski RE. PLEIOTROPIC EFFECTS OF BENEFICIAL MUTATIONS IN ESCHERICHIA COLI. Evolution. 2005;59(11):2343–52.

43. Leiby N, Marx CJ. Metabolic Erosion Primarily Through Mutation Accumulation, and Not Tradeoffs, Drives Limited Evolution of Substrate Specificity in Escherichia coli. PLoS Biol. 2014 Feb 18;12(2):e1001789–10.

44. Sane M, Miranda JJ, Agashe D. Antagonistic pleiotropy for carbon use is rare in new mutations. Evolution. 2018 Oct;72(10):2202–13.

45. Chevereau G, Dravecká M, Batur T, Guvenek A, Ayhan DH, Toprak E, et al. Quantifying the Determinants of Evolutionary Dynamics Leading to Drug Resistance. PLoS Biol. 2015 Nov 18;13(11):e1002299–18.

46. Reding-Roman C, Hewlett M, Duxbury S, Gori F, Gudelj I, Beardmore R. The unconstrained evolution of fast and efficient antibiotic-resistant bacterial genomes. Nature Ecology & Evolution. 2017 Jan 30;1(3):e1001490–11.

47. Ardell SM, Kryazhimskiy S. The population genetics of collateral resistance and sensitivity. Elife [Internet]. 2021 Dec 10;10. Available from: http://dx.doi.org/10.7554/eLife.73250

48. Bono LM, Jr LBS, Pfennig DW, Burch CL. The emergence of performance trade-offs during local adaptation: insights from experimental evolution. Mol Ecol. 2017 Jan 13;26(7):1720–33.

49. Visher E, Boots M. The problem of mediocre generalists: population genetics and eco-evolutionary perspectives on host breadth evolution in pathogens. Proc Biol Sci. 2020 Aug 26;287(1933):20201230.

50. Kawecki TJ. Accumulation of Deleterious Mutations and the Evolutionary Cost of Being a Generalist. Am Nat. 1994 Nov 1;144(5):833–8.

51. Whitlock MC. The red queen beats the jack-of-all-trades: The limitations on the evolution of phenotypic plasticity and niche breadth. Am Nat. 1996 Nov;148:S65–77.

52. Remold SK, Rambaut A, Turner PE. Evolutionary genomics of host adaptation in vesicular stomatitis virus. Mol Biol Evol. 2008 Jun;25(6):1138–47.

53. Novella IS, Presloid JB, Smith SD, Wilke CO. Specific and nonspecific host adaptation during arboviral experimental evolution. J Mol Microbiol Biotechnol. 2011;21(1-2):71–81.

54. Tikhonov M, Kachru S, Fisher DS. A model for the interplay between plastic tradeoffs and evolution in changing environments. Proc Natl Acad Sci U S A. 2020 Apr 21;117(16):8934–40.

55. Shoval O, Sheftel H, Shinar G, Hart Y, Ramote O, Mayo A, et al. Evolutionary trade-offs, Pareto optimality, and the geometry of phenotype space. Science. 2012 Jun 1;336(6085):1157–60.

56. Li Y, Petrov DA, Sherlock G. Single nucleotide mapping of trait space reveals Pareto fronts that constrain adaptation. Nature Ecology & Evolution. 2019 Oct 23;3(11):1–15.

57. MacLean RC, Perron GG, Gardner A. Diminishing returns from beneficial mutations and pervasive epistasis shape the fitness landscape for rifampicin resistance in Pseudomonas aeruginosa. Genetics. 2010 Dec;186(4):1345–54.

58. Chou HH, Chiu HC, Delaney NF, Segrè D, Marx CJ. Diminishing returns epistasis among beneficial mutations decelerates adaptation. Science. 2011 Jun 3;332(6034):1190–2.

59. Kryazhimskiy S, Rice DP, Jerison ER, Desai MM. Microbial evolution. Global epistasis makes adaptation predictable despite sequence-level stochasticity. Science. 2014 Jun 26;344(6191):1519–22.

60. Wei X, Zhang J. Environment-dependent pleiotropic effects of mutations on the maximum growth rate r and carrying capacity K of population growth. PLoS Biol. 2019 Jan;17(1):e3000121.

61. Martin G, Lenormand T. The fitness effect of mutations across environments: a survey in light of fitness landscape models. Evolution. 2006 Dec;60(12):2413–27.

62. Cooper LA, Scott TW. Differential Evolution of Eastern Equine Encephalitis Virus Populations in Response to Host Cell Type. Genetics. 2001 Apr 1;157(4):1403–12.

63. Wilson DS, Yoshimura J. On the coexistence of specialists and generalists. Am Nat. 1994

Oct;144(4):692–707.

64. Parrella G, Gognalons P, Gebre-Selassie K, Vovlas C, Marchoux G. An update of the host range of tomato spotted wilt virus. J Plant Pathol. 2003 Dec;85(4):227–64.

65. Andika IB, Wei S, Cao C, Salaipeth L, Kondo H, Sun L. Phytopathogenic fungus hosts a plant virus: A naturally occurring cross-kingdom viral infection. Proc Natl Acad Sci U S A. 2017 Nov 14;114(46):12267–72.

66. Lefeuvre P, Martin DP, Elena SF, Shepherd DN, Roumagnac P, Varsani A. Evolution and ecology of plant viruses. Nat Rev Microbiol. 2019 Jul 16;17(10):632–44.