Unveiling genome plasticity and a novel phage in *Mycoplasma felis*: Genomic investigations of

four feline isolates

Sara M. Klose[1,2*], Alistair R. Legione[1*], Rhys N. Bushell[3], Glenn F. Browning[1] and Paola K. Vaz[1]

*S.M.K. and A.R.L. have contributed equally to this work.

Affiliation:

1. Asia-Pacific Centre for Animal Health, Melbourne Veterinary School, Faculty of Science, The

University of Melbourne, VIC, Australia

2. Institute of Molecular Medicine and Experimental Immunology, Faculty of Medicine, University

of Bonn, NRW, Germany

3. Department of Veterinary Clinical Sciences, Melbourne Veterinary School, Faculty of Science, The

University of Melbourne, VIC, Australia

Corresponding Author: Alistair Legione, legionea@unimelb.edu.au

Keywords: mycoplasma felis, whole genome sequencing, genome plasticity, evolution, phage

Abstract:

We present the complete genomes of four clinical isolates of *Mycoplasma felis* from three domestic cats, assembled with the aid of short and long read sequencing methods. The completed genomes encode a median of 759 open reading frames (min, 743, max 777) and a median 98.2% average nucleotide identity to that of the available equid origin reference strain. Comparative genomic analysis revealed the occurrence of multiple horizontal gene transfer (HGT) events and genome reassortment. This resulted in the acquisition or loss of numerous genes within the Australian felid genomes, encoding putative proteins involved in DNA transfer, metabolism, DNA replication, host cell interaction, and restriction modification systems. Additionally, a novel mycoplasma phage was detected in one Australian felid *M. felis* isolate via genomic analysis and visualised via cryo transmission electron microscopy. This research identifies complex genomic dynamics and evolutionary adaptation of mycoplasmas to different host environments. Furthermore, the sequences obtained in this work are important for the development of new diagnostic tools, and identification of future infection control and treatment options for respiratory disease complex in cats.

Data summary:

All genome data for this study have been deposited in GenBank under BioProject PRJNA906261. Genome assemblies, as well as Illumina and Oxford Nanopore sequence reads for each isolate can be found under their respective BioSamples: SAMN32182834 (isolate 047), SAMN32182835 (isolate 219), SAMN32182836 (isolate 329), and SAMN32182837 (isolate 632). The authors confirm all supporting data and protocols have been provided within the article.

Impact statement:

Mycoplasma felis is commonly associated with clinical cases of conjunctivitis and feline respiratory disease complex in cats, the leading cause of euthanasia in shelters. In the absence of vaccines, infection control is currently limited to the prolonged use of antimicrobials. Prior to this study there was only one complete genome assembly of *M. felis*, which was isolated from a horse. The outcomes from this study marks the first high quality hybrid assembled genomes assembled from cats. This work adds four new genomes from clinical cases, as well as the identification and validation of the presence of a novel phage that utilises the mycoplasma translation table. The

52 genome data presented here can assist future projects investigating improved diagnostics and

53 development of new treatment options.

Introduction:

Mycoplasmas are the smallest free-living bacteria with the smallest bacterial genomes identified to date (0.58 – 1.38 Mb). Mycoplasmas are thought to approach the minimal gene sets essential for independent life, making them ideal model organisms for studies on the fundamental requirements of a living cell [1]. Mycoplasmas evolved from Gram positive bacterial ancestors 600 million years ago [2], and have undergone rapid genome reduction during evolution [3, 4]. They are cell wall-less, Gram negative pleomorphic bacteria and can be found to infect and cause both acute and chronic disease in a diverse array of animal and plant species [5].

Mycoplasmas are known for their rapid evolution and genetic plasticity, mainly driven by intra-species spontaneous mutation or horizontal gene transfer (HGT) between and within species, facilitating swift adaptation to environmental changes and enhancing survival and/or virulence [6-9]. Studies examining the genomic stability of *Mycoplasma synoviae* revealed that it not only undergoes frequent mutations in birds [10], but also manifests rapid thermoadaptive evolution *in vitro* and *in vivo* [11], presumably to regain fitness and pathogenicity [12, 13]. HGT is observed to occur more frequently between species that share an ecological niche irrespective of their phylogenetic divergence, such as transfer of surface VlhA lipoprotein involved in phase variation from *Mycoplasma gallisepticum* to *M. synoviae* infecting poultry [14], and transfer of several surface lipoproteins involved in host colonisation from *Mycoplasma capricolum* subsp. *capricolum* and *Mycoplasma mycoides* subsp. *mycoides* to *Mycoplasma agalactiae*, infecting ruminants [15].

There have been at least four species of mycoplasmas isolated from cats (*Mycoplasma felis*, *Mycoplasma gatae, Mycoplasma arginini*, and *Mycoplasma feliminutum*), in most cases as common commensal microflora, but they have been associated with conjunctivitis and, increasingly, feline respiratory disease [16-19]. Feline respiratory disease complex (FRDC) in cats is the leading cause of euthanasia in animal shelters [20] and is commonly associated with several viral (feline herpesvirus 1 and feline calicivirus) and bacterial pathogens, including *M. felis* [21, 22]. *M. felis* infections are common in most cat populations and their ability to establish chronic persistence in an animal enhances opportunities for recurring opportunistic infections. Vaccines against *M. felis* are not currently available and treatment is limited to the prolonged use of antimicrobials in the tetracycline and fluoroquinolone classes [23, 24]. Currently there are three isolates with full

genomes published for this species: strain Myco-2 (GenBank accession number: AP022325.1), an
isolate from a case of respiratory disease in a horse in Japan [25] which has been completely
assembled, and multiple assemblies of isolates 16-040612-4 (CP104192, CP104193, CP104194,
CP11027) and 16-057065-4 (CP103988, CP103993, CP110269), which were each isolated from
feline bronchoalveolar lavage in Canada [26]. Additionally, there is an incomplete ATCC strain (ATCC
23391) assembled into scaffolds (GenBank Biosample: SAMN02841218).
92
This report describes four high quality, complete, *M. felis* genomes, the first that have been
generated using hybrid assembly methods. These genomes also represent the first complete
assembly of feline isolates of this species. Additionally, we have identified and visualised a novel
mycoplasma phage, which utilises mycoplasma codons. This report represents a significant
expansion on our limited understanding of the genomic structure and plasticity of *M. felis*. Such
work may lead to enhanced development of diagnostic tests and vaccines, as has been seen in
other mycoplasma species.
100

Methods:

Microbial isolation
Swabs were taken for bacteriological culturing from clinical cases of respiratory disease (3 cases)
and an unresolved post-surgical infection (1 case) (Table 1). Swabs were spread onto aerobic Sheep
Blood Agar (SBA) plates and incubated at 37°C. Following observation and presumptive
identification of small mycoplasma-like colonies, single colonies were transferred to Mycoplasma
Broth (MB) containing 10% swine serum (Sigma-Australia) and 0.01% nicotinamide adenine
dinucleotide (NAD) (Sigma-Australia) based on a formulation of Frey's medium with minor
modification [27, 28]. Acidification of MB media was used to confirm the growth of *Mycoplasma*
spp. and the cultures were stored at -80°C for future investigations.
111

DNA extraction, sequencing, and genome assembly
Bacterial isolates were cultured axenically in MB for 18 hours at 37°C prior to DNA extraction.
Bacteria were pelleted and media removed, followed by resuspension in extraction kit specific lysis
buffers. The DNA for short read sequencing was extracted using the DNeasy Blood and tissue kit
(QIAGEN) following the manufacturer's protocol. Following this, 100 ng of extracted DNA was used

to prepare sequencing libraries using Illumina's Nextera Flex DNA library prep kit. Sequencing was

performed on the NovaSeq platform at Charles River Laboratories, Victoria, Australia, to generate

150 bp paired end reads. Two batches of Oxford Nanopore Technologies (ONT) long read

sequencing were performed. The DNA was extracted using the Promega Wizard® DNA Purification

Kit (Batch 1) or the Promega Wizard High Molecular Weight DNA Purification Kit (Batch 2). In both

instances DNA libraries were generated using the ONT Rapid Barcoding (SQK-RBK004) kit.

Sequencing was performed on a MinION Mk1b device fitted with a FLO-MIN106D flow cell (R9.4.1

chemistry). Base calling, demultiplexing, and adapter removal utilised Guppy (version

6.2.1+6588110a6) high accuracy base caller and a minimum quality score of 9 (default settings).

The Illumina reads were trimmed and adaptors removed using fastp v0.20.1 [29] using default

settings. The ONT reads from the two batches were merged per isolate and processed through

Filtlong v0.2.0 [30] to remove all reads shorter than 1000 bp and retain 90% of input data

(removing short and low quality reads). Subsequently, the processed short and long reads were

used as input into Unicycler v0.5.0 [31] in 'normal' mode, with a modified start gene database to

include *M. felis*. The assemblies where then polished using Polypolish v0.5.0 [32], incorporating

BWA-MEM v0.7.17-r1188 [33]. Full genome alignments were generated using the progressive

MAUVE algorithm [34] to identify regions of reassortment and gene loss or gain.

Genome annotation and sequencing analysis

Genomes were annotated using the Prokaryotic Genome Annotation Pipeline v2022-12-

13.build6494 [35] for primary analysis. A Tenericutes specific protein database was downloaded

from UniProt containing both Swiss-Prot (4464 sequences) and TrEMBL (343,598 sequences)

annotations and converted to a diamond format protein database [36]. This database was used as

input to Pydeel (https://github.com/alegione/pydeel) to determine the coding ratio of our

annotated genomes compared to *M. felis* genomes on GenBank. This tool identifies the coding ratio

of annotated genes compared to their best hit within the provided database to identify the

proportion of annotated genes that may have erroneous insertions or deletions as a by-product of

the use of long read sequencing. From here, annotated genomes were processed through Panaroo

v1.3.0 [37] along with previously sequenced *M. felis* genomes (Myco-2 and ATCC23391) to identify

the core genome. The core genome results were visualised using Pagoo [38]. The annotated

genomes from this study were investigated for pathogenicity islands, and integrative conjugative

elements using Island Viewer 4 [39], and ICE finder [40], respectively. Antimicrobial resistance
genes and virulence factors were analysed by generating a Diamond database of the CARD [41] and
full virulence factor databases [42], and using Diamond BLASTp to identify best hits with a cut off
bit score of 50. The UniProt webserver was used to identify homologs of specific open reading
frames (ORFs) in other genomes (https://www.uniprot.org).

Phage isolation and visualisation
A phage-like contig was identified in the annotation results of one of the assemblies (MF329). In
order to confirm if the detection of bacteriophage genomic sequence corresponds to integrated
prophage or to replicative virions, bioinformatic methods were first utilised. The probable terminal
ends were investigated through short and long read mapping to the contig using BWA-MEM and
minimap2 v2.26-r1175 [43], respectively, and the contig's starting point adjusted based on the
depth of coverage. The tool PhageTerm [44] was also utilised to investigate the potential phage
start/end point. To investigate whether this contig was a prophage the long reads were mapped,
using minimap2, to the updated phage contig. Any reads that had overhangs at the 5' and 3' end of
the contig were extracted and subsequently mapped to the MF329 assembled genome to
determine if there was a common point of overlap.
To determine the presence of productive bacteriophage virions, transmission electron microscopy
was performed on cultured MF329. Isolate MF329 was cultured in 50 mL of axenic mycoplasma
media as described above for 20 hours. The bacterial culture was centrifuged for 15 mins at 10,000
x g at 4°C and the media fraction separated from the bacterial cell pellet. The bacterial cells were
gently lysed by resuspension in 150 uL of sterile water, followed by centrifugation at 10,000 x g at
4°C for 5 mins to remove cellular debris. This was repeated twice, with supernatant fraction
captured each time for ultracentrifugation. The culture media and the cell lysate supernatant
fractions were separately ultra-centrifugated at 100,000 x g for 90 mins at 4°C, and pellets
resuspended in 0.2 M HEPES buffer, pH 7.7. Samples were inspected and visualised by negative
staining at the Ian Holmes Imaging Center (Melbourne, Australia) using a FEI Talos L120C cryoTEM
(Talos L120C) microscope.

Results:

Genome characterisation and comparisons

All four isolates were successfully assembled into circularised, complete genomes. In addition to

these four genomes, a novel phage genome of *M. felis*, was identified that utilised the Mycoplasma

translation table. Metrics for long and short read sequencing output are available in Table 2. The

assembly statistics for the four isolates are available in Table 3. Briefly, the circularised isolates

were sufficiently larger than the reference genome (Myco-2; 841,695 bp), with a median length of

940,935 bp (range: 905,741 − 948,716) and a median of 759 coding domain sequences (CDS)

(range: 743 − 777) compared to the reference genome which has 740. The NCBI Prokaryotic

Genome Annotation Pipeline incorporates a CheckM [45] completeness analysis and an average

nucleotide identity taxonomic classification step to identify any potential issues with an assembled

genome. For all four cases CheckM completeness mirrored the reference genome Myco-2 (99.21%

completeness), and all assemblies were classified as *Mycoplasma felis* with high confidence (Table

3). The coding ratio of annotated assemblies were investigated to identify the frequency of

potential indel errors. Two published *M. felis* genomes (Myco-2 and ATCC23391) had a mean

coding ratio of 1.0 (Standard deviation ±0.02) and 0.96 (±0.32), respectively. Our completed

genomes, MF47, MF219, MF329, and MF632 had mean coding ratios of 1.0 (±0.28), 0.99 (±0.30),

1.0 (±0.27), and 1.0 (±0.26), respectively. Seven recently sequenced *M. felis* isolates that utilised

only Oxford Nanopore sequencing had a substantial number of indel errors in their annotations

based on coding ratio analysis (mean coding ratio between 0.44 and 0.47) and were not used in

further comparisons for this reason (Figure 1).

Two isolates, MF047 and MF632 had an additional tRNA compared to the remaining isolates, for

tRNA-Trp and an unknown tRNA respectively. The newly sequenced isolates had 1-3 secretory

systems (in all cases these were annotated as Type IV secretory systems), whilst Myco-2 has no

annotated secretory systems. The newly annotated sequences also and an increased number of

transposases, typically annotated as IS1634 or IS256, and IS3 in a few cases. By comparison Myco-2

only had IS3 and IS1634 family transposase annotations.

A progressive MAUVE alignment of the four assemblies in comparison to Myco-2 (Figure 2)

identified two significant genome inversions in the reference sequence compared to the isolates

assembled here, as well as more localised inversions between each of the isolates. These

inversions, along with genomic regions present in individual isolates, were typically flanked by

transposases. Comparison of ANIs between the four new assemblies, and previously published *M.*
felis genomes (Table 4) found that all four Australian isolates' best match was to another isolate
from this study besides MF329 (MF047 and MF219, 99.8%; MF632 and MF329, 98.3%, MF329 and
ATCC23391 98.4%), noting that MF047 and MF219 originated from the same animal two months
apart.

The core genome of *M. felis* as determined by Panaroo (genes present in >99% of genomes)
contained 595 genes, with 301 shell genes (genes present in 15% - 95% of genomes). Collapsing
paralogues increased the core genes by one and reduced the shell genes to 219. Principal
component analysis (PCA) highlighted clustering of the two published genomes, as well as
clustering of isolates that were taken from the same individual (Figure 3). All genes, besides those
annotated as hypothetical proteins, identified in newly sequenced isolates but not present in the
reference sequence are present in Table 5, including the homologs found in other genomes. The
variations between the genomes were mostly observed in sequences encoding putative proteins
involved in DNA transfer, metabolism, DNA replication, mycoplasma-host cell interaction, and
restriction modification systems (Figure 4).

Genomic islands, virulence genes, antibiotic resistance genes
Genomic islands were detected in three out of the four sequenced genomes (Table 7). The average
length of these islands was 22,145 bp (±5818 bp) and each contained an average of 16 (±5)
annotated genes. In comparison, the Myco-2 strain of *M. felis* had two predicted genomic islands of
3052 bp and 10,027 bp length. Further inspection of these predicted genomic islands identified
matching regions present in isolates in alternative locations. For example, MF329 had a 24 kb
(311,409 – 335,810 nucleotide position [nt]) predicted genomic island containing 19 genes
including a type IV secretion system and an insertion sequence (IS1634) at its midpoint. Extraction
and remapping of this genomic island found that it was repeated three times within a wider 122 kb
region (213,995 – 335,810 nt) of the MF329 genome. Of these repeated regions, only the predicted
genomic island repeat contained the IS element. This same region occurred without the IS element
present in MF047 (twice), MF219 (once), and MF632 (once). In the case of MF632, the region was
also a predicted as a genomic island, but in this instance was flanked with restriction endonucleases
and either predicted transposases (3') or an IS256 element (5').

Diamond BLAST searches for antibiotic resistance genes and virulence genes resulted in
identification of few high identity hits (Table 6 and Supplementary Tables 2 and 3). Only one hit to
the virulence factor database had an amino acid identity greater than 90%, predicted to encode for
the adherence virulence gene elongation factor Tu in *Mycoplasma synovia* (90.1% average identity)
and was present in all genomes. Below that, the only five genes had hits with >50% amino acid
identity. The closest hits to these genes were oppF (59.9% identity; *Mycoplasma mycoides* subsp.
Mycoides; Immune modulation), eno (58.9%; *Streptococcus pneumoniae*; Exoenzyme), gap (57.9%;
Mycoplasma genitalium; Adherence), dnaK (53.9%; *Chlamydia trachomatis*; Adherence), and sugC
(50.4%; *Mycobacterium marinum*; Nutritional/Metabolic factor). No genes had hits with >50%
amino acid identity with antibiotic resistance genes in the CARD database.
253
Phage/Prophage detection and identification
A 33.6 kb contig (vB Mfe PM329; *Mycoplasma felis* strain PM329) was assembled and identified as
a likely phage or prophage within the sequenced *M. felis* MF329 isolate. It had an average depth of
coverage of 40,847.7 reads, with 82.6% of Illumina MF329 reads mapping to phage genome.
Interrogation of the nanopore read dataset to determine genome integration status detected only
63/18134 (0.35%) reads mapping to both phage and bacterial genome. Of these, most were found
to map across the bacterial genome at mostly single-read depth, with the exception of five reads
which clustered mid-genome in the region of 456,500 nt. If this were to be indicative of a true
insertion site it would likely disrupt ORF MF329_000403, encoding a putative 16S rRNA
pseudouridine synthase gene. The phage genome has a 27.1% G+C content and encodes 36
predicted ORFs using the mycoplasma translation table. It shared 94.8% nt global pairwise identity
with that of two nanopore contigs in the publicly available unassembled *M. felis* contig dataset
(NCBI Biosample: SAMN30346842), and 96.8% and 97.4% aa pairwise identity between their
encoded putative recombinase and portal proteins, respectively. Interrogation of the read datasets
of the other three isolates sequenced in this study identified only 4/9515 (0.04%; MF047),
19/19128 (0.1%; MF632) and 22/65383 (0.03%; MF219) mapped nanopore reads, within the range
of barcode error, and no Illumina reads mapped to the phage genome, indicating those isolates
were not carrying the phage. Phylogenetic analyses via VipTree [46] and vContact2 [47] was not
able to assign vB Mfe PM329 to any known phage genus or family, however gene arrangement was
found to be similar to those of the MAgV1 from *M. agalactiae* and MAgV1-like prophage sequences
described in mycoplasma species of the Hominis phylogenetic host group. In particular, both

protein and gene arrangement similarity were greatest to those in prophage encoded within *M.*
mustelae and *M. molare*, including a central position of the recombinase gene. BLASTX analysis of
predicted ORFs detected protein pairwise identities of 48% to 69% between core conserved ORFs
encoded within the *M. mustelae* and *M. molare* prophage sequences [48] (**Error! Reference source**
not found.).

Confirmation of productive virus formation was performed by cryoTEM on media and cell lysate
supernatant fractions, revealing the presence of non-enveloped contractile-tailed polyhedral viral
particles of approximately 100 nm in length, with a 50 nm diameter head and tails of varying
lengths observed. Higher numbers of virions were observed within the cellular lysate supernatant
fraction than in the culture media fraction, with clusters of virions seemingly embedded within
bacterial membrane debris observed (Figure 5). These long tailed polyhedral viral structures are
consistent with those possessed by siphoviruses and myoviruses found within the recently ratified
Caudoviricetes class of bacteriophage. The contractile tail and polyhedral head structure is
consistent across all visually elucidated mycoplasma phages, with the main differences
corresponding with head and tail lengths. *M. pulmonis* Phage P1 has a 28 nm isometric head and
short tail, Br1 has 77 nm head diameter and a 104 nm long tail [49], and *M. hyorhinis* Hr1 phage, is
a short-tailed (14nM) phage with a 34 nm head [50].

294 Discussion

In this study, genomic characterisation of four new *M. felis* isolates was performed; three derived
from clinical cases of respiratory disease and the remaining from an unresolved post-surgical
infection. Comparisons that included several published datasets detected genomic variations and
inversions between the Australian felid isolates and the only assembled *M. felis* reference equid
isolate, prominently flanked by transposases which is indicative of HGT and genome reassortment.
This includes differences detected in the number of copies of several encoded putative transposase
families, such as IS1634, IS256 and IS3 family transposases (Figure 4). Insertion sequences (ISs) are
transposable elements often present in multiple copies in a genome and involved in chromosomal
rearrangements by deletion, insertion, or inversion. The substantial similarity between various
isoforms of ISs enables them to facilitate DNA homologous recombination [51]. IS families encoded
in mycoplasma genomes play an important role in genome plasticity with several reported

instances of IS exchange between strains or species [15, 52-55]. This is in contrast to early research
that suggested some ISs, such as IS1634 which was in abundance in our sequenced genomes, were
unique to specific mycoplasma species [56]. The dissimilarity of the Australian felid genomes
obtained from three different cats, based on gene presence/absence and the number of coding
sequences, underscores the genetic diversity present amongst sequenced felid isolates (Figure 3) as
a result of the acquisition or loss of numerous genes within their genomes. Multiple gene
insertions, relative to the equid reference genome, were identified within the cluster of Australian
field isolates (Table 5). This includes genes involved in DNA transfer, replication, metabolism,
restriction modification (RM) systems, as well as various proteins of unknown function (Figure 4).

Gene gains compared to the equid reference strain

Of particular interest are those gene acquisitions involved in DNA transfer, such as the detection of
several copies encoding the putative type IV secretory system (T4SS) conjugative DNA transfer
family protein within the Australian felid genomes, with none detected in the publicly available
equid and felid genomes. T4SS conjugation systems translocate DNA through a process dependent
on cell to cell contact and have been identified as one of the core genes within Integrative and
Conjugative Elements (ICEs) in mycoplasmas, which play a significant role in genome plasticity of
these bacteria [57, 58]. ICEs have been shown to be self-transmissible in _M. agalactiae_ providing
conjugative properties from ICE-positive cells to ICE-negative cells, while chromosomal transfers
(CTs) occur in the opposite direction at the same time [59]. Protein sequence analysis of the
putative T4SS identified in the Australian feline isolates showed 43.2% identity to the ORF5 of ICE in
Mycoplasma cynos. Whilst _M. cynos_ predominantly infects the respiratory tract of dogs, it has also
been detected in cats in association with conjunctivitis and upper respiratory tract disease [60].
Taken together, it is plausible that a potential HGT event may have occurred to transfer the T4SS
conjugative element of ICE from an _M. cynos_-like species to _M. felis_ in a shared feline host. This
T4SS was part of a defined region of genes that, along with 15 hypothetical proteins and a DUF87
domain-containing protein, was repeated three times in MF329. DUF87 domain-containing proteins
are associated with conjugative elements in other organisms such as TraJ in the virulence plasmid
of _Salmonella enterica_ serovar Typhimurium [61], adding further weight to the likelihood of this
being an ICE, despite the bioinformatics tools utilised here not detecting it as one.

Several genes involved in metabolism were newly detected in felid isolates sequenced in this study.

A type 2 glycerol-3-phosphate oxidase (GlpO) was identified within the Australian genomes, while

in the equid isolate, only its FAD-dependent oxidoreductase domain was detected, accompanied by

two transposases located upstream. GlpO converts glycerol-3-phosphate to dihydroxyacetone

phosphate (DHAP) releasing hydrogen peroxide as a by-product which has been shown to play a

role in *in vitro* cytotoxicity of *M. gallisepticum* and *Mycoplasma pneumoniae* [62, 63], and virulence

of *M. mycoides* subsp. *mycoides* small colony and mosquito-associated *Spiroplasma* species [64,

65]. The significant reduction or complete loss of H_2O_2 production in highly passaged *M. bovis, M.*

agalactiae, M. gallisepticum, and attenuated vaccine strains of *M. gallisepticum* compared to wild

type strains suggests that glycerol metabolism capabilities can rapidly become redundant [62, 66,

67]. It is possible that the sequences coding for GlpO was lost in the equid *M. felis* isolate possibly

due to different levels of available glycerol as the nutrient source, or an altered pathogenicity in

horses compared to cats. Likewise, variations in the gene counts coding for nutrient transport

systems were observed, including ATP-binding cassette (ABC) transporters and the PEP-dependent

phosphotransferase transport system (PTS), as well as numerous other metabolic enzymes amongst

the Australian and equid isolates. A number of these nutrient transport systems, and many cell

surface enzymes are multifunctional in mycoplasmas playing critical roles in pathogenesis and

survival in the host [13, 68-73]. The differences observed in the PTS and ABC transporters of the

Australian felid isolates, in contrast to the equid isolate, may also be indicative of rapid metabolic

gene gains and losses occurring in response to the varying nutrient availabilities within their felid

and equid ecological niche.

In the Australian felid genomes, five new coding sequences related to replication were identified

which were absent in the equid origin genome, including genes responsible for putative anti-

termination protein NusB, Hsp70 family protein, C5 methylase, 16S rRNA methyltransferase RsmG,

and HNH endonuclease. The presence of these additional proteins suggests a potential difference in

bacterial growth within the feline host as compared to the equine host, potentially due to distinct

replication strategies. Future experiments comparing growth patterns of these isolates in felid or

equid derived cells are indicated. The potential origins of these additional sequences have

frequently been detected in mycoplasmas isolated from mink, swine, and dogs, which share a

common ecological environment with cats, or pigeons which are a common prey for cats. The HGT

events resulting in the acquisition of these genes by the Australian *M. felis* felid isolates may have

been facilitated by the close ecological proximity or may have occurred within the feline

gastrointestinal tract (GI) following the ingestion of the infected prey. HGT among diverse

microorganisms within the mammalian GI tract is frequently observed, possibly due to the physical

proximity it provides, thereby facilitating genetic exchange [74]. Furthermore, the conjugation

mechanism favoured for HGT within the GI tract protects DNA from nucleases and heavy metals,

and thus results in a successful genetic exchange [75].

Two additional restriction endonucleases (Eco47II and Sau3AI) were detected in the Australian felid

genomes, and the loss of a DpnII family type II restriction endonuclease. Restriction modification

(RM) systems play a pivotal role in regulating the bacterial HGT by protecting the genome against

incoming mobile genetic elements, or by facilitating recombination of the invading DNA [76]. A

strong correlation has been reported between the presence of active Type II RM systems and the

presence of mobile genetic elements and ICEs within *M. agalactiae* genomes, indicating the

facilitative role of Type II RM systems in HGT in mycoplasmas [77]. Furthermore, the diversification

of RM target recognition sites can regulate HGT by promoting gene exchanges among those with

RM systems that recognise the same target motifs, and by reducing gene transfer between lineages

that possess distinct RM systems [78]. Taken together, it is plausible that following a host switch

into horses, *M. felis* may have diversified their RM systems to enhance the level of genetic

exchange from phylogenetically distant bacteria, especially in cases where genetic exchanges were

previously restricted.

Gene losses compared to the equid reference strain

A sequence encoding a putative ComEC/Rec2 family competence protein has been observed to be

absent in the Australian felid isolates. This protein is one of the components of DNA translocation

via transformation enabling DNA transport by forming a channel across the cytoplasmic membrane

[79]. While ComEC component has been reported essential for DNA transport in Gram positive

bacteria, deletion of homologous genes did not affect DNA uptake in cell wall-deficient bacteria

[80]. Furthermore, the competence domain seems to be missing in many mycoplasmas possibly

due to the reductive evolution of these bacteria [81]. Characterisation of two mycoplasma genomes

recovered from gut microbiota of a deep-sea isopod detected *comEC* gene adjacent to genes

encoding enzymes producing dTMP from thymidine or dUMP, suggesting that the role of ComEC

protein might be importing extracellular DNA as a nutrient source [82]. The presence of ComEC in

the *M. felis* equid isolate, but its absence in the Australian felid isolates suggests that during the

evolutionary process the DNA translocation approach via a membrane channel might have been

made redundant and perhaps replaced with T4SS conjugation-mediated tactics as discussed above.

The Australian felid genomes exhibited a total absence of four coding sequences involved in host

cell interaction, namely those encoding putative variable surface lipoprotein, fibronectin type III

domain-containing protein, FIVAR domain-containing protein, and GA module-containing protein,

in contrast to the equid genome. The FIVAR (Found In Various Architectures Region) and GA

domains serve as essential constituents of the extracellular matrix-binding protein (Embp),

facilitating biofilm formation and attachment to fibronectin in some bacteria such as

Staphylococcus epidermidis and *S. aureus* [83, 84]. The presence of more than 30% similarity in

protein sequences between the FIVAR domain and GA protein in the equid *M. felis* reference Myco-

2 isolate, in comparison to the domains found in Embp of *S. aureus*, raises the possibility that the

equid isolate might have retained the potential for biofilm formation and/or fibronectin

attachment during the genome reductive evolutionary transition from Gram-positive bacteria.

Conversely, the absence of these proteins, in addition to the surface lipoprotein and fibronectin

type III domain-containing protein, suggests that these capacities may have become obsolete in the

Australian *M. felis* isolates infecting feline hosts, possibly indicating a different mode of host cell

interaction following the host switch.

Rapid evolution within two months in one host:

Two of the Australian *M. felis* isolates sequenced in this study, MF047 and MF219, were obtained

from separate bronchoalveolar lavages (BAL) of a feline patient presenting clinical signs consistent

with pneumonia. MF047 was isolated from the BAL taken at initial consultation, whereafter

treatment with 25 mg of doxycycline was administered twice a day orally for approximately two

months. Due to an unsatisfactory response to treatment, a second BAL was taken and MF219 was

isolated. While no genome reassortment was detected between the MF047 and MF219 genomes,

minor differences were observed in the counts of some genes encoding proteins involved in RM

systems, DNA transfer, and methylation (Figure 4). These variations may suggest a rapid host

adaptation of *M. felis* after only two months. The growing utilisation of whole genome sequencing

in mycoplasma diagnostics is revealing links between genetic mutations and antimicrobial

resistance. Genomic comparison of MF047 and MF219 revealed the presence of five single

nucleotide polymorphisms (SNPs) within the gene encoding ribosomal RNA small subunit

methyltransferase A (RsmA/KsgA), along with one SNP in the gene responsible for 16S rRNA and

another SNP in the gene responsible for 23S rRNA. In other studies, mutations within the 16S RNA

methyltransferase family have been reported to confer aminoglycoside resistance in *M. bovis* and

M. gallisepticum [85, 86], while mutations in the 16S rRNA or 23S rRNA genes have been shown to

be associated with resistance to tetracyclines in *M. pneumoniae*, *M. hominis*, *M. genitalium*, and *M.*

bovis [85, 87-89]. Both MF047 and MF219 contained five protein coding regions with low

confidence matches to tetracycline resistance genes, with the highest bit score of the five

associated with a predicted elongation factor G protein with an amino acid identity/similarity to

tet(T) from *Steptococcus pyogenes* of 25.8%/47.7%. It is important to emphasise that the

application of novel genome sequencing techniques for mycoplasma diagnostics, particularly in the

context of antimicrobial resistance, is still in its early stages, and the availability of data for

comprehensive analysis covering all mycoplasma species (particularly animal mycoplasmas) is

currently limited. Future research involving susceptibility testing of isolates MF047 and MF219 to

tetracyclines is necessary to determine whether there is a correlation with the observed genomic

variations and tetracycline resistance.

Phage in mycoplasmas

In other bacteria, phage have been demonstrated to provide critical virulence factors to their hosts,

required for the pathogenic activity of diverse bacteria (e.g. cholera, diptheriae, botulism, shiga

toxin-producing *E. coli*, etc). Approximately 20 mycoplasma-infecting phage/prophage have been

reported across diverse ruminant, avian, canine, swine, human and rodent mycoplasmas [48, 90].

Of these, most are unclassified, only three have had their virion structure visualised (P1, Hr1 and

Br1), and only one has been classified by the International Committee for the Taxonomy of Viruses

(ICTV). That is, the temperate *M. pulmonis* phage P1 (12 Kb genome), as the sole member of the

floating genus *Delislevirus* [91]. The well-described MAV1 temperate phage found in *M. arthritidis*

(16 Kb genome) has not yet (at time of publication) been classified within the new system. In other

mycoplasmas, phages have demonstrated dynamic mobility within the bacterial chromosome, loss

of integrated phage with clonal passage (MFV1 in *M. fermentans*), and variable expression of

unique phage-encoded membrane surface proteins (MFV1 and MAV1, in *M. fermentans* and *M.*

arthritidis, respectively) [92] with a potential role in bacterial adaptation and survival. Detection of

a similar contig within *M. felis* 16-057065-4 (NCBI Biosample: SAMN30346842) from Canada

indicates that this phage is likely carried by *M. felis* strains globally, although in that study it was not
identified as a phage/prophage but as *M. felis* bacterial genomic DNA. No known selective
advantage or disadvantage to the presence of this phage within *M. felis* was determined as part of
this study due to the low number of available *M. felis* genomes. Future epidemiological studies to
investigate the potential role of phage carriage in feline mycoplasma-associated disease cases is
indicated.

<u>Conclusions</u>
This study presents a comprehensive analysis of the first high-quality, complete *M. felis* genomes
isolated from cats, along with the characterisation of a novel mycoplasma phage. It provides
valuable insights into the intricate genomic dynamics of *M. felis*, showcasing a complex interplay of
gene gains, losses, horizontal gene transfer events, and adaptations to new host environments.
These findings suggest divergent evolutionary paths in different hosts and offer compelling avenues
for further research into the evolutionary mechanisms and ecological factors that influence the
genomic diversity of mycoplasmas.

Conflicts of interest:

The author(s) declare that there are no conflicts of interest.

Funding information:

This study was supported by the Feline Health Research Fund. P.K.V. was supported by a University
of Melbourne, Melbourne Postdoctoral Fellowship.

Author contributions:

R.N.B. performed the bacterial isolation and initial classification; A.R.L., S.M.K. and P.K.V. prepared
the DNA samples, performed the sequencing and analysis; P.K.V. isolated the phage for TEM; P.K.V.,
S.M.K., G.F.B. and A.R.L. conceptualised the study design; P.K.V., S.M.K. and A.R.L. acquired the
funding; S.M.K., A.R.L. and P.K.V. drafted the manuscript, and R.N.B. and G.F.B. revised the
manuscript.

Acknowledgments:

The authors would like to gratefully acknowledge the small animal medicine team of the U-Vet Werribee Animal Hospital at The University of Melbourne, for providing the samples taken from clinical cases, and Eric Hanssen at the Ian Holmes Imaging Centre for the cryoTEM imaging of vB Mfe PM329 phage.

503 **Table 1.** *Mycoplasma felis* isolates sequenced in this study.

Isolate	Month/Year	Animal ID	Clinical case	Isolation site	Other notes	Genbank #
047	3/2015	Cat A	Pneumonia	Respiratory, BAL	Heavy growth from BAL, treated with Doxycycline	CP114890
219	5/2015	Cat A	Pneumonia	Respiratory, BAL	Follow-up isolation from Cat A	CP114889
329	5/2014	Cat B	FRDC	Respiratory, BAL	Chronic bronchitis. PCR negative for herpesvirus, calicivirus and chlamydia	CP115656 OR753717 (phage)
632	12/2010	Cat C	Surgery	Fat tissue	Isolated from non-healing surgical wound	CP114888

504 BAL, Bronchioalveolar lavage; PCR, polymerase chain reaction; FRDC, feline respiratory disease complex

506 **Table 2.** DNA sequencing results summary after trimming and quality filtering

	MF047	MF219	MF329	MF632
Illumina reads				
Total paired reads	8,977,712	6,647,214	6,481,506	6,807,888
Median length	151	151	151	151
>Q20 (%)	96.93	95.7	96.83	96.99
>Q30 (%)	91.31	88.43	91.12	91.41
GC (%)	24.64	24.67	26.35	24.85
Nanopore reads^				
Total reads	65,383	9,515	103,977	19,128
Median length	7,049	6,563	3,717	5,058.5
N50	16,885	22,105	12,331	16,399
>Q20 (%)	48.38	51.69	41.64	49.27
>Q30 (%)	15.12	16.77	11.64	15.74
GC (%)	24.9	24.79	27.19	25.08

507 ^Results presented are from two Nanopore sequencing runs combined

509 **Table 3.** Assembly and annotation summary of four isolates from this study, compared to the complete reference sequence for Mycoplasma felis (Myco-2)

Isolate	Myco-2	MF047	MF219	MF329	MF632
Accession	NZAP022325	CP114890	CP114889	CP115656	CP114888
Assembly information					
Length (bp)	841,695	948,716	945,056	936,813	905,741
CheckM completeness (%)	99.21	99.21	99.21	99.21	99.21
PGAP ANI*	98.4%	98.5%	98.6%	98.6%	98.5%
Median sequencing depth					
Short reads	-	1762	1372	467	1499
Long reads	-	695	110	88	181
Genomic features					
CDS	740	759	777	759	743
ncRNA	2	2	2	2	2
rRNA	9	9	9	9	9
Regulatory	2	2	2	2	2
tRNA	29	30	29	29	30
tmRNA	1	1	1	1	1
Genes of interest					
Secretory Systems	0	2	1	3	1
Transposase	22	40	52	27	38

510 * Average nucleotide identity to Mycoplasma felis ATCC 23991 according to internal PGAP ANI taxonomic classification

Figure 1. Coding ratio frequency counts across *Mycoplasma felis* genomes from this study and those currently available
on GenBank. Coding ratio determined by dividing amino acid length of annotated proteins to best hit match in custom
UniProt Mycoplasma database. Coding ratio axis is log transformed for visualisation. Title names for NCBI origin data
includes accession number and strain name (e.g. NZ_CP104194, and 16-040612-4). For Oxford Nanopore only data
from NCBI, depth of coverage is added from NCBI (e.g. 58x).

Figure 2. A progressive Mauve alignment of the *Mycoplasma felis* reference genome (Myco-2) and the genomes sequenced in this project. Locally co-linear blocks are

separated by colour within genomes, and are the same colour between genomes, with links to identify placement and inversion differences. Solid blocks overlaying the

genomes represent the presence of transposases, type IV secretion systems, and genomic islands, whilst solid blocks of blue that break up the colour gradient within the

genomes are representative of LCBs not present in linked genomes

Table 4. Average nucleotide identity (ANI) results of Australian *M. felis* isolate genomes compared to equine reference (Myco-2) and felid contig data sets (Genbank

Accessions: MF047: CP114890; MF219: CP114889; MF329: CP115656; MF632: CP114888; Myco-2: AP022325; 16-057065-4: CP103988; 16-040612-4:CP104192;

ATCC23391: JNKA01000000).

	MF047	MF219	MF329	MF632	Myco-2	16-057065-4	16-040612-4
MF219	99.79	-	-	-	-	-	-
MF329	98.21	98.16	-	-	-	-	-
MF632	98.21	98.17	98.33	-	-	-	-
Myco-2	98.14	98.06	98.18	98.17	-	-	-
16-057065-4	97.77	97.80	98.25	97.77	97.77	-	-
16-040612-4	97.65	97.65	97.84	97.80	97.72	97.65	-
ATCC23391	98.31	98.29	98.39	98.27	98.33	97.95	97.95

Figure 3. *Mycoplasma felis* core genome and gene presence/absence analysis conducted using Panaroo and plotted using Pagoo. **A)** The frequency of clustered genes within the six genomes used in the analysis. **B)** A principal component analysis of gene presence/absence using Bray Curtis dissimilarity identifies clustering of the previously published genome sequences **C)** Clusters of genes present (blue) or absent (yellow) in the six genomes included in analysis.

533 **Table 5.** Coding sequences detected in the Australian *M. felis* isolate genomes but absent in the equine reference (Myco2), and their possible species of origin based on

534 amino acid sequence identity.

Coding sequence	Possible species of origin	% Amino acid pairwise identity	Predominant Host
Type IV secretory system conjugative DNA transfer family protein	*M. cynos*	43.2	Dog
IS256 family transposase	*M. penetrans*	50.9	Human
Type 2 glycerol-3-phosphate oxidase (GlpO)	*M. canis*	87.2	Dog
PTS fructose transporter subunit IIA	*M. flocculare*	52.6	Swine
DHHA1 domain-containing protein	*M. canis*	84.8	Dog
M13 family peptidase	*M. canis*	65.4	Dog
1-phosphofructokinase family	*M. canis*	52.6	Dog
Anti-termination protein NusB	*M. mustelae*	61.6	Mink
Hsp70 family protein	*M. flocculare*	85.4	Swine
C5 methylase (MAV1virus-like) protein	*M. columboralis*	70.2	Pigeon
16S rRNA methyltransferase (RsmG)	*M. mustelae*	63.3	Mink
HNH endonuclease	*M. columbina*	38.0	Pigeon
Eco47II family restriction endonuclease	*M. mycoides* subsp. *mycoides* SC	49.6	Cattle
Sau3AI family type II restriction endonuclease	*M. crocodyli*	52.7	Crocodile

536 **Table 6.** Quantity and range of amino acid identity and similarity of putative virulence and antimicrobial resistance genes (AMR) based on significant BLASTP hits (bit scores

537 >50) compared to the virulence factor database and CARD database, respectively.

Genome	Putative virulence genes			Putative AMR genes		
	Total	Median Identity (min − max)	Median Similarity (min − max)	Total	Median Identity (min − max)	Median Similarity (min − max)
Myco-2	48	31.7 (24.4 - 90.1)	52.6 (44.4 - 95.9)	24	29.8 (22.5 - 40.1)	51.7 (39.1 - 69.6)

Genome	Putative virulence genes			Putative AMR genes		
	Total	Median Identity (min − max)	Median Similarity (min − max)	Total	Median Identity (min − max)	Median Similarity (min − max)
MF047	49	31.6 (24.6 - 90.1)	52.7 (41.8 - 95.9)	25	29.4 (22.6 - 40.3)	50.6 (39.2 - 69.6)
MF219	49	31.6 (24.6 - 90.1)	52.4 (41.8 - 95.9)	25	29.6 (22.6 - 40.3)	50.6 (39.2 - 69.6)
MF329	50	32 (24.4 - 90.1)	52.2 (41.5 - 95.9)	25	29.7 (22.5 - 40.2)	50.6 (39.1 - 69.6)
MF632	51	31.6 (24.4 - 90.1)	52.4 (41.5 - 95.9)	26	29.7 (22.4 - 40.2)	51 (38.9 - 69.6)

540 **Table 7.** The location of predicted genomic islands in isolated *M. felis* genomes from this study, as well as the reference genome Myco-2. Genomic Islands determined with

541 the Island Finder 4 web tool. None were detected in MF047 isolate.

Strain	Genomic Island	Start − End (Relative to genome)	Length	Total number of genes (non-hypothetical)
Myco-2	1 [b,d,e]	133250 - 143277	10027	11 (5)
	2 [b,c,d,e]	723039 - 726091	3052	8 (6)
MF047	-	-	-	-
MF219	1 [a,b,d,*]	428340 - 453515	25175	15 (7)
	2 [a,b,d,e]	582071 - 602042	19971	11 (6)
MF329	1 [b,c,#]	311409 - 335810	24401	19 (15)

Strain	Genomic Island	Start – End (Relative to genome)	Length	Total number of genes (non-hypothetical)
MF632	1[a,b,d,*]	376254 - 403316	27062	19 (8)
	2[b,#]	511813 - 537933	26120	22 (15)
	3[a,b,c,d]	716455 - 726594	10139	8 (1)

542 [a, b, c, d, e] – Predicted genomic island is present with some variation, such as the presence/absence of a transposase, in Myco-2 (a), MF047 (b), MF219 (c), MF329 (d), or

543 MF632 (e) but not predicted to be a genomic island in those cases.

544 [*, #] - Predicted genomic islands with structural similarity across different genomes

Figure 4. Presence and absence of genes within Australian felid *M. felis* isolates compared to equid *M. felis* reference Myco-2 genome, clustered by putative protein

functions. Only genes that differ between isolates or differ from reference are listed. Counts indicate number of gene copies detected within each isolate.

Figure 5. A) Genome of bacteriophage identified in *Mycoplasma felis* strain MF329 compared to MAgV1-like prophage sequences found in *M. mustelae*, *M. molare* and *M. agalactiae*. Genome annotations depicting conserved phage/prophage gene sequences are indicated in black. H = Helicase; Pol = DNA polymerase; D = DNA primase; X = Xer recombinase; C = prohead protein; P = portal protein; T = terminase; HP = hypothetical protein. **B)** and **C)** Cryo-transmission electron photomicrographs of bacteriophage detected in cell lysate of *M. felis* isolate MF329 indicating the phage structure consists of a contractile-tail and polyhedral head.

References:

1.   **Glass JI, Assad-Garcia N, Alperovich N, Yooseph S, Lewis MR et al.** Essential genes of a minimal bacterium. *Proceedings of the National Academy of Sciences of the United States of America* 2006;103(2):425-430.

2.   **Woese CR, Stackebrandt E, Ludwig W**. What are mycoplasmas: The relationship of tempo and mode in bacterial evolution. *Journal of Molecular Evolution* 1985;21(4):305-316.

3.   **Fadiel A, Eichenbaum KD, El Semary N, Epperson B**. Mycoplasma genomics: tailoring the genome for minimal life requirements through reductive evolution. *Frontiers in Bioscience-Landmark* 2007;12(6):2020-2028.

4.   **Rocha EPC, Blanchard A**. Genomic repeats, genome plasticity and the dynamics of Mycoplasma evolution. *Nucleic Acids Research* 2002;30(9):2031-2042.

5.   **Shil PK, Wawegama NK, Browning GF, Noormohammadi AH, Marenda MS**. Mycoplasmas. In: J.F. Prescott, A.N. Rycroft, J.D. Boyce, J.I. MacInnes, Immerseel FV et al. (editors). *Pathogenesis of Bacterial Infections in Animals* 2022. pp. 667-700.

6.   **Krause DC, Leith DK, Wilson RM, Baseman JB**. Identification of *Mycoplasma pneumoniae* proteins associated with hemadsorption and virulence. *Infect Immun* 1982;35(3):809-817.

7.   **Delaney NF, Balenger S, Bonneaud C, Marx CJ, Hill GE et al.** Ultrafast evolution and loss of CRISPRs following a host shift in a novel wildlife pathogen, *Mycoplasma gallisepticum*. *PLOS Genetics* 2012;8(2):e1002511.

8.   **Momynaliev KT, Govorun VM**. Mechanisms of genetic instability in Mollicutes (Mycoplasmas). *Russian Journal of Genetics* 2001;37(9):979-992.

9.   **Citti C, Dordet-Frisoni E, Nouvel LX, Kuo CH, Baranowski E**. Horizontal Gene Transfers in Mycoplasmas (Mollicutes). *Current Issues in Molecular Biology* 2018;29(1):3-22.

10.   **Klose SM, Olaogun OM, Disint JF, Shil P, Gyuranecz M et al.** Genomic diversity of a globally used, live attenuated mycoplasma vaccine. *Microbiology Spectrum* 2022;10(6):e02845-02822.

11.   **Klose SM, Shil P, Underwood GJ, Morrow CJ, Marenda MS et al.** Obg plays a significant role in temperature sensitivity of Mycoplasma synoviae live attenuated vaccine strain MS-H. *Veterinary Microbiology* 2023:109818.

12.   **Klose SM, Omotainse OS, Zare S, Vaz PK, Armat P et al.** Virulence factors of Mycoplasma synoviae: Three genes influencing colonization, immunogenicity, and transmissibility. *Frontiers in Microbiology* 2022;13:1042212.

13.   **Klose SM, De Souza DP, Disint JF, Andrews DM, Underwood GJ et al.** Reversion of mutations in a live mycoplasma vaccine alters its metabolism. *Vaccine* 2023;41(21):3358-3366.

14.   **Noormohammadi A, Markham P, Duffy M, Whithear K, Browning G**. Multigene families encoding the major hemagglutinins in phylogenetically distinct mycoplasmas. *Infection and Immunity* 1998;66(7):3470-3475.

15.   **Sirand-Pugnet P, Lartigue C, Marenda M, Jacob D, Barré A et al.** Being pathogenic, plastic, and sexual while living with a nearly minimal bacterial genome. *PLOS Genetics* 2007;3(5):e75.

16.   **Randolph J, Moise N, Scarlett J, Shin S, Blue J et al.** Prevalence of mycoplasmal and ureaplasmal recovery from tracheobronchial lavages and of mycoplasmal recovery from pharyngeal swab specimens in cats with or without pulmonary disease. *American journal of veterinary research* 1993;54(6):897-900.

17.   **Foster S, Barrs V, Martin P, Malik R**. Pneumonia associated with *Mycoplasma* spp in three cats. *Australian veterinary journal* 1998;76(7):460-464.

18.   **Sykes JE**. Feline hemotropic mycoplasmas. *Journal of Veterinary Emergency and Critical Care* 2010;20(1):62-69.

19.   **Le Boedec K**. A systematic review and meta-analysis of the association between Mycoplasma spp and upper and lower respiratory tract disease in cats. *Journal of the American Veterinary Medical Association* 2017;250(4):397-407.

20.   **Cohn LA**. Feline respiratory disease complex. *Veterinary Clinics: Small Animal Practice* 2011;41(6):1273-1289.

21.   **Nguyen D, Barrs VR, Kelman M, Ward MP**. Feline upper respiratory tract infection and disease in Australia. *Journal of Feline Medicine and Surgery* 2019;21(10):973-978.

22. **Fernandez M, Manzanilla EG, Lloret A, León M, Thibault J-C**. Prevalence of feline
herpesvirus-1, feline calicivirus, *Chlamydophila felis* and *Mycoplasma felis* DNA and associated risk
factors in cats in Spain with upper respiratory tract disease, conjunctivitis and/or gingivostomatitis.
Journal of Feline Medicine and Surgery 2017;19(4):461-469.

23. **Lee-Fowler T**. Feline respiratory disease: What is the role of Mycoplasma species? *Journal of*
Feline Medicine and Surgery 2014;16(7):563-571.

24. **Tonozzi CC**. *Feline Respiratory Disease Complex.* MSD Veterinary Manual 2022.

25. **Kinoshita Y, Niwa H, Uchida-Fujii E, Nukada T**. Complete genome sequence of mycoplasma
felis Strain Myco-2, isolated from an equine tracheal wash sample in Japan. *Microbiology Resource*
Announcements 2020;9(9):10.1128/mra. 00057-00020.

26. **Framst I, D`Andrea C, Baquero M, Maboni G**. Development of a long-read next generation
sequencing workflow for improved characterization of fastidious respiratory mycoplasmas.
Microbiology 2022;168(11).

27. **Frey M**. A medium for the isolation of avian mycoplasmas. *American Journal of Veterinary*
Research 1968;29:2163-2171.

28. **Klose SM, Omotainse OS, Zare S, Vaz PK, Armat P et al.** Virulence factors of Mycoplasma
synoviae: Three genes influencing colonization, immunogenicity, and transmissibility. *Front*
Microbiol, Original Research 2022;13.

29. **Chen S, Zhou Y, Chen Y, Gu J**. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 2018;34(17):i884-i890.

30. **Wick R**. *Filtlong v. 0.2. 0.* 2021.

31. **Wick RR, Judd LM, Gorrie CL, Holt KE**. Unicycler: Resolving bacterial genome assemblies
from short and long sequencing reads. *PLOS Computational Biology* 2017;13(6):e1005595.

32. **Wick RR, Holt KE**. Polypolish: Short-read polishing of long-read bacterial genome
assemblies. *PLOS Computational Biology* 2022;18(1):e1009802.

33. **Li H**. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv*
preprint arXiv:13033997 2013.

34. **Darling AE, Mau B, Perna NT**. Progressive Mauve: Multiple alignment of genomes with gene
flux and rearrangement. *arXiv preprint arXiv:09105780* 2009.

35. **Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP et al.** NCBI prokaryotic
genome annotation pipeline. *Nucleic Acids Research* 2016;44(14):6614-6624.

36. **Buchfink B, Reuter K, Drost H-G**. Sensitive protein alignments at tree-of-life scale using
DIAMOND. *Nature Methods* 2021;18(4):366-368.

37. **Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G et al.** Producing polished
prokaryotic pangenomes with the Panaroo pipeline. *Genome Biology* 2020;21(1):180.

38. **Ferrés I, Iraola G**. An object-oriented framework for evolutionary pangenome analysis. *Cell*
Reports Methods 2021;1(5):100085.

39. **Dhillon BK, Chiu TA, Laird MR, Langille MG, Brinkman FS**. IslandViewer update: improved
genomic island discovery and visualization. *Nucleic Acids Research* 2013;41(W1):W129-W132.

40. **Liu M, Li X, Xie Y, Bi D, Sun J et al.** ICEberg 2.0: an updated database of bacterial integrative
and conjugative elements. *Nucleic Acids Research* 2018;47(D1):D660-D665.

41. **Alcock BP, Huynh W, Chalil R, Smith KW, Raphenya AR et al.** CARD 2023: expanded
curation, support for machine learning, and resistome prediction at the Comprehensive Antibiotic
Resistance Database. *Nucleic Acids Research* 2023;51(D1):D690-D699.

42. **Liu B, Zheng D, Zhou S, Chen L, Yang J**. VFDB 2022: a general classification scheme for
bacterial virulence factors. *Nucleic acids research* 2022;50(D1):D912-D917.

43.   **Li H**. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*
2018;34(18):3094-3100.
44.   **Garneau JR, Depardieu F, Fortier L-C, Bikard D, Monot M**. PhageTerm: a tool for fast and
accurate determination of phage termini and packaging mechanism using next-generation
sequencing data. *Scientific Reports* 2017;7(1):8292.
45.   **Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW**. CheckM: assessing the
quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome*
Research 2015;25(7):1043-1055.
46.   **Nishimura Y, Yoshida T, Kuronishi M, Uehara H, Ogata H et al.** ViPTree: the viral proteomic
tree server. *Bioinformatics* 2017;33(15):2379-2380.
47.   **Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S et al.** Taxonomic assignment of
uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. *Nature Biotechnology*
2019;37(6):632-639.
48.   **Citti C, Baranowski E, Dordet-Frisoni E, Faucher M, Nouvel L-X**. Genomic Islands in
Mycoplasmas. *Genes* 2020;11(8):836.
49.   **Gourlay RN, Wyld SG, Garwes DJ**. Some properties of mycoplasma virus Br1. *Archives of*
Virology 1983;75(1):1-15.
50.   **Gourlay RN, Wyld SG, Poulton ME**. Some characteristics of Mycoplasma virus Hr1, isolated
from and infectingMycoplasma hyorhinis. *Archives of Virology* 1983;77(1):81-85.
51.   **Mahillon J, Chandler M**. Insertion sequences. *Microbiology and Molecular Biology Reviews*
1998;62(3):725-774.
52.   **Frey J**. Insertion Sequence Analysis. In: Miles R, Nicholas R (editors). *Mycoplasma Protocols*.
Totowa, NJ: Humana Press; 1998. pp. 197-205.
53.   **Lysnyansky I, Calcutt MJ, Ben-Barak I, Ron Y, Levisohn S et al.** Molecular characterization of
newly identified IS 3, IS 4 and IS 30 insertion sequence-like elements in Mycoplasma bovis and their
possible roles in genome plasticity. *FEMS microbiology letters* 2009;294(2):172-182.
54.   **Thomas A, Linden A, Mainil J, Bischof DF, Frey J et al.** Mycoplasma bovis shares insertion
sequences with Mycoplasma agalactiae and Mycoplasma mycoides subsp. mycoides SC:
Evolutionary and developmental aspects. *FEMS Microbiology Letters* 2005;245(2):249-255.
55.   **Loreto ELS, Ortiz MF, Porto JIR**. Insertion sequences as variability generators in the
Mycoplasma hyopneumoniae and M. synoviae genomes. *Genetics and Molecular Biology*
2007;30:283-289.
56.   **Vilei EM, Nicolet J, Frey J**. IS*1634*, a novel insertion element creating long, variable-length
direct repeats which is specific for *Mycoplasma mycoides* subsp. *mycoides* small-colony type.
Journal of Bacteriology 1999;181(4):1319-1323.
57.   **Baranowski E, Dordet-Frisoni E, Sagné E, Hygonenq M-C, Pretre G et al.** The Integrative
Conjugative Element (ICE) of *Mycoplasma agalactiae*: Key Elements Involved in Horizontal
Dissemination and Influence of Coresident ICEs. *mBio* 2018;9(4):10.1128/mbio.00873-00818.
58.   **Alvarez-Martinez CE, Christie PJ**. Biological Diversity of Prokaryotic Type IV Secretion
Systems. *Microbiology and Molecular Biology Reviews* 2009;73(4):775-808.
59.   **Dordet-Frisoni E, Sagné E, Baranowski E, Breton M, Nouvel LX et al.** Chromosomal transfers
in mycoplasmas: when minimal genomes go mobile. *MBio* 2014;5(6):10.1128/mbio. 01958-01914.
60.   **Hartmann AD, Hawley J, Werckenthin C, Lappin MR, Hartmann K**. Detection of bacterial
and viral organisms from the conjunctiva of cats with conjunctivitis and upper respiratory tract
disease. *Journal of Feline Medicine & Surgery* 2010;12(10):775-782.
61.   **Ahmer BMM, Tran M, Heffron F**. The virulence plasmid of *Salmonella typhimurium* is self-
transmissible. *Journal of Bacteriology* 1999;181(4):1364-1368.

62.   **Szczepanek SM, Boccaccio M, Pflaum K, Liao X, Geary SJ**. Hydrogen Peroxide Production
from Glycerol Metabolism Is Dispensable for Virulence of Mycoplasma gallisepticum in the Tracheas
of Chickens. *Infection and Immunity* 2014;82(12):4915-4920.
63.   **Hames C, Halbedel S, Hoppert M, Frey J, Stülke J**. Glycerol metabolism is important for
cytotoxicity of Mycoplasma pneumoniae. *Journal of Bacteriology* 2009;191(3):747-753.
64.   **Chang T-H, Lo W-S, Ku C, Chen L-L, Kuo C-H**. Molecular evolution of the substrate utilization
strategies and putative virulence factors in mosquito-associated Spiroplasma species. *Genome*
Biology and Evolution 2014;6(3):500-509.
65.   **Pilo P, Vilei EM, Peterhans E, Bonvin-Klotz L, Stoffel MH et al.** A metabolic enzyme as a
primary virulence factor of Mycoplasma mycoides subsp. mycoides small colony. *Journal of*
bacteriology 2005;187(19):6824-6831.
66.   **Khan LA, Miles RJ, Nicholas RAJ**. Hydrogen Peroxide Production by Mycoplasma bovis and
Mycoplasma agalactiae and Effect of In Vitro Passage on a Mycoplasma bovis Strain Producing High
Levels of H2O2. *Veterinary Research Communications* 2005;29(3):181-188.
67.   **Szczepanek S, Tulman E, Gorton T, Liao X, Lu Z et al.** Comparative genomic analyses of
attenuated strains of Mycoplasma gallisepticum. *Infection and immunity* 2010;78(4):1760-1771.
68.   **Tseng C-W, Chiu C-J, Kanci A, Citti C, Rosengarten R et al.** The oppD gene and putative
peptidase genes may be required for virulence in Mycoplasma gallisepticum. *Infection and*
Immunity 2017;85(6):10.1128/iai. 00023-00017.
69.   **Masukagami Y, Nijagal B, Mahdizadeh S, Tseng C-W, Dayalan S et al.** A combined
metabolomic and bioinformatic approach to investigate the function of transport proteins of the
important pathogen Mycoplasma bovis. *Veterinary microbiology* 2019;234:8-16.
70.   **Masukagami Y, De Souza D, Dayalan S, Bowen C, O'callaghan S et al.** Comparative
metabolomics of Mycoplasma bovis and Mycoplasma gallisepticum reveals fundamental
differences in active metabolic pathways and suggests novel gene annotations. *MSystems*
2017;2(5):10.1128/msystems. 00055-00017.
71.   **Mahdizadeh S, Masukagami Y, Tseng C-W, Markham PF, De Souza DP et al.** A *Mycoplasma*
gallisepticum glycerol ABC transporter involved in pathogenicity. *Applied and Environmental*
Microbiology 2021;87(11):e03112-03120.
72.   **Chen H, Yu S, Shen X, Chen D, Qiu X et al.** The Mycoplasma gallisepticum α-enolase is cell
surface-exposed and mediates adherence by binding to chicken plasminogen. *Microbial*
Pathogenesis 2011;51(4):285-290.
73.   **Vilei EM, Frey J**. Genetic and biochemical characterization of glycerol uptake in Mycoplasma
mycoides subsp. mycoides sc: Its impact on h2o2production and virulence. *Clinical Diagnostic*
Laboratory Immunology 2001;8(1):85-92.
74.   **Shterzer N, Mizrahi I**. The animal gut as a melting pot for horizontal gene transfer. *Canadian*
Journal of Microbiology 2015;61(9):603-605.
75.   **Norman A, Hansen LH, Sørensen SJ**. Conjugative plasmids: vessels of the communal gene
pool. *Philosophical Transactions of the Royal Society B: Biological Sciences* 2009;364(1527):2275-
2289.
76.   **Oliveira PH, Touchon M, Rocha EP**. The interplay of restriction-modification systems with
mobile genetic elements and their prokaryotic hosts. *Nucleic acids research* 2014;42(16):10618-
10631.
77.   **Dordet-Frisoni E, Vandecasteele C, Contarin R, Sagné E, Baranowski E et al.** Impacts of
Mycoplasma agalactiae restriction-modification systems on pan-epigenome dynamics and genome
plasticity. *Microbial Genomics* 2022;8(5).

78.   **Oliveira PH, Touchon M, Rocha EPC**. Regulation of genetic flux between bacteria by
restriction–modification systems. *Proceedings of the National Academy of Sciences*
2016;113(20):5658-5663.
79.   **Chen I, Dubnau D**. DNA uptake during bacterial transformation. *Nature Reviews*
Microbiology 2004;2(3):241-249.
80.   **Kapteijn R, Shitut S, Aschmann D, Zhang L, de Beer M et al.** Endocytosis-like DNA uptake by
cell wall-deficient bacteria. *Nature Communications* 2022;13(1):5524.
81.   **Pimentel ZT, Zhang Y**. Evolution of the natural transformation protein, ComEC, in bacteria.
Frontiers in Microbiology 2018:2980.
82.   **Wang Y, Huang JM, Wang SL, Gao ZM, Zhang AQ et al.** Genomic characterization of
symbiotic mycoplasmas from the stomach of deep-sea isopod Bathynomus sp. *Environmental*
Microbiology 2016;18(8):2646-2659.
83.   **Christner M, Franke GC, Schommer NN, Wendt U, Wegert K et al.** The giant extracellular
matrix-binding protein of *Staphylococcus epidermidis* mediates biofilm accumulation and
attachment to fibronectin. *Molecular Microbiology* 2010;75(1):187-207.
84.   **Speziale P, Pietrocola G, Foster TJ, Geoghegan JA**. Protein-based biofilm matrices in
Staphylococci. *Frontiers in cellular and infection microbiology* 2014;4:171.
85.   **Ledger L, Eidt J, Cai HY**. Identification of antimicrobial resistance-associated genes through
whole genome sequencing of *Mycoplasma bovis* isolates with different antimicrobial resistances.
Pathogens 2020;9(7):588.
86.   **Mahdizadeh S, Sansom FM, Lee S-W, Browning GF, Marenda MS**. Targeted mutagenesis of
Mycoplasma gallisepticum using its endogenous CRISPR/Cas system. *Veterinary Microbiology*
2020;250:108868.
87.   **Pereyre S, Goret J, Bébéar C**. *Mycoplasma pneumoniae*: current knowledge on macrolide
resistance and treatment. *Frontiers in Microbiology* 2016;7:974.
88.   **Dégrange S, Renaudin H, Charron A, Pereyre S, Bébéar C et al.** Reduced susceptibility to
tetracyclines is associated *in vitro* with the presence of 16S rRNA mutations in *Mycoplasma hominis*
and *Mycoplasma pneumoniae*. *Journal of antimicrobial chemotherapy* 2008;61(6):1390-1392.
89.   **Shimada Y, Deguchi T, Nakane K, Yasuda M, Yokoi S et al.** Macrolide resistance–associated
23S rRNA mutation in *Mycoplasma genitalium*, Japan. *Emerging infectious diseases*
2011;17(6):1148.
90.   **Kovács ÁB, Wehmann E, Sváb D, Bekő K, Grózner D et al.** Novel prophage-like sequences in
Mycoplasma anserisalpingitidis. *Infection, Genetics and Evolution* 2021;92:104886.
91.   **Turner D, Shkoporov AN, Lood C, Millard AD, Dutilh BE et al.** Abolishment of morphology-
based taxa and change to binomial species names: 2022 taxonomy update of the ICTV bacterial
viruses subcommittee. *Archives of Virology* 2023;168(2):74.
92.   **Röske K, Calcutt MJ, Wise KS**. The Mycoplasma fermentans prophage φMFV1: genome
organization, mobility and variable expression of an encoded surface protein. *Molecular*
Microbiology 2004;52(6):1703-1720.
788