Nationwide genome surveillance of carbapenem-resistant *Pseudomonas*

***aeruginosa* in Japan**

Hirokazu Yano[1*], Wataru Hayashi[1], Sayoko Kawakami[1], Sadao Aoki[1], Eiko Anzai[1], Norikazu

Kitamura[1], Aki Hirabayashi[1], Toshiki Kajihara[1], Shizuo Kayama[1], Yo Sugawara[1], Koji Yahara[1],

Motoyuki Sugai[1*]

[1]Antimicrobial Resistance Research Center, National Institute of Infectious Diseases. 4-2-1

Aobacho, Higashimurayama, Tokyo 189-0002, Japan

*Corresponding author: Antimicrobial Resistance Research Center, National Institute of

Infectious Diseases. 4-2-1 Aobacho, Higashimurayama, Tokyo 189-0002, Japan.

Email address: h-yano@niid.go.jp (H Yano); sugai@niid.go.jp (M Sugai)

ABSTRACT

Japan is a representative country with a low prevalence rate of carbapenem-resistant

Pseudomonas aeruginosa (CRPA). A comprehensive overview of the genotype and phenotype

patterns of CRPA in the Asia-Pacific region is lacking. Herein, we conducted genome sequencing

and standardized quantitative antimicrobial susceptibility testing for 382 meropenem-resistant

CRPA isolates that were collected from 78 hospitals across Japan from 2019 to 2020. CRPA

exhibited susceptibility rates of 52.9%, 26.4%, and 88.0% against piperacillin-tazobactam,

ciprofloxacin, and amikacin, respectively, whereas 27.7% of CRPA isolates were classified as

difficult-to-treat resistance *P. aeruginosa*. Only 4.1% of CRPA isolates carried carbapenemase:

GES-5 (2), IMP-1 (5), IMP-10 (4), IMP-7 (3), IMP-34 (1), and IMP-98 (1). Regarding chromosomal

mutations, 87.1% of CRPA isolates possessed inactivating or other resistance mutations in *oprD*,

and 28.8% showed mutations in the regulatory genes (*mexR, nalC,* and *nalD*) for the MexAB-

OprM efflux pump. Additionally, 4.7% of CRPA isolates carried a known resistance mutation in

the PBP3-encoding gene *ftsI*. Of the 148 sequence types detected, ST274 (9.7%) was

predominant, followed by ST235 (7.6%) and ST253 (4.5%). The findings from this study and

other surveillance studies collectively demonstrate that CRPA exhibits marked genetic diversity

and that its multidrug resistance in Japan is less perceived than in other regions. This study

contributes a valuable dataset that addresses a gap in genotype/phenotype information

regarding CRPA in the Asia-Pacific region, where the epidemiological background markedly

differs between regions.

INTRODUCTION

Pseudomonas aeruginosa is a common opportunistic pathogen associated with

pneumonia, urinary tract infections, and bacteremia in clinical settings worldwide. Currently, it

ranks among the top five bacterial pathogens associated with human death globally (1). It also

intrinsically shows reduced susceptibility to many antibiotics due to its possession of four types

of RND family efflux pump (MexAB-OprM, MexCD-OprJ, MexEF-OprN, and MexXY-OprM),

chromosomally encoded AmpC β-lactamase (also called *Pseudomonas*-derived class C β-

lactamase: PDC), and other antimicrobial-inactivating enzymes (2, 3). Furthermore, *P.*

aeruginosa can acquire resistance mutations at multiple genomic locations and antimicrobial

resistance genes via mobile genetic elements, such as plasmids and integrative and conjugative

elements (ICEs), after which it elicits high levels of resistance to multiple antimicrobials, thereby

limiting treatment options for infected patients (3–5).

Carbapenems represent a crucial class of therapeutic options for treating *P. aeruginosa*

infections. In 2017, the World Health Organization (WHO) listed carbapenem-resistant *P.*

aeruginosa (CRPA) as one of the bacterial priority pathogens classified in "Priority 1: CRITICAL,"

for which new effective antibiotics are urgently required (6). A recent cohort study revealed that

patients infected with carbapenemase-producing CRPA have a 9% higher 30-day mortality rate

than patients infected with carbapenemase-nonproducing CRPA (7).

Carbapenem resistance of *P. aeruginosa* often involves inactivating mutation in several

chromosomal genes: inactivation of OprD porin leads to inefficient penetration of the outer

membrane by carbapenems (8), and inactivating mutation in two direct or one indirect

transcriptional regulators (repressors) MexR, NalD, and NalC induces increased production of

the MexAB-OprM efflux pump, thereby increasing β-lactam minimum inhibitory concentration

(MIC) (9–11). Furthermore, an experimental evolution study showed that nonsynonymous

substitutions in *ftsI*, an essential gene encoding PBP3, contribute to carbapenem resistance (12).

Notably, the extent to which these mutations have become prevalent in *P. aeruginosa* genomes

in clinical settings remains largely unknown.

Certain sequence types (STs) of *P. aeruginosa* are widely recognized as global high-risk

clones due to their high mortality rates and propensity to acquire multidrug resistance (MDR)

and to transmit worldwide (13). The top ten global high-risk clones are ST111, ST175, ST233,

ST235, ST244, ST277, ST298, ST308, ST357, and ST654 (13, 14). International consortiums have

made efforts to build a large *P. aeruginosa* dataset to obtain quantitative information for CRPA

prevalence, its susceptibility profiles, and the possession rate of carbapenemases (7, 15–17);

here, both class A group 2f β-lactamase and class B group 3 β-lactamase (so called metallo-β-

lactamases, MBLs) are included for simplicity. Globally disseminated carbapenemase types in *P.*

aeruginosa include VIM, IMP, NDM, GES, and KPC; however, the carbapenemase possession rate

and the relative frequencies of carbapenemase types differ between geographic regions (7, 16,

17). Notably, in the Asia-Pacific region, the Antimicrobial Testing Leadership and Surveillance

(ATLAS) program took the initiative of *P. aeruginosa* surveillance, revealing distinct patterns of

resistance rate and predominant β-lactamase levels in the *P. aeruginosa* population between 12

countries/regions, but the ATLAS did not cover genome surveillance (16).

Japan exhibits characteristic features in the epidemiology of antimicrobial-resistant

bacterial infection in the Asia-Pacific region. Regarding carbapenem resistance, the prevalence

of meropenem-resistant *E. coli* in bloodstream infections in Japan as of 2019 was 2 per 100,000

test patients, which is lower than, for example, 79 per 100,000 test patients observed in

Singapore, which is located in Southeast Asia, according to the WHO Global Antimicrobial

Resistance and Use Surveillance System (18). The established national phenotypic antimicrobial

resistance surveillance program—the Japan Nosocomial Infections Surveillance (JANIS) (19,

20)—reported that the carbapenem (meropenem) resistance rate of *P. aeruginosa* clinical

isolates in Japan, as of 2020, was 10.5%. This was rather low among representative countries

where antimicrobial resistance surveillance for *P. aeruginosa* was once or is currently being

performed; the resistance rate was 29.3% in India (ATLAS 2015–2019), 19.3% in China (CHINET

2020), 6.5% in Australia (ATLAS 2015–2019), 28.5% in Poland (EARS-net 2020), 15.7% in the

Czech Republic (EARS-net 2020), 13.9% in Germany (EARS-net 2020), 12.6% in France (EARS-net

2020), and 3.7% in Finland (EARS-net 2020) (16, 21, 22). Previous nationwide surveillance

studies of *P. aeruginosa* in Japan primarily focused on amikacin-nonsusceptible MDR *P.*

aeruginosa or isolates from respiratory system origin only (23–25). Therefore, a comprehensive

overview of the resistance phenotype, STs, carbapenemase presence, and chromosomal

resistance mutations of more common CRPA in Japanese clinical settings remains lacking.

To obtain an overview of both genomic features and resistance phenotypes of bacterial

pathogens in Japan, we initiated the national bacterial genome surveillance—Japan

Antimicrobial-Resistant Bacterial Surveillance (JARBS)—in 2019 (26). Through the JARBS-PA

project performed in 2019 and 2020, 382 meropemen-resistant CRPA isolates were collected

from 78 hospitals across Japan. Herein, we aimed to clarify the genotype and susceptibility

patterns of the CRPA isolates by conducting genome sequencing and wide-range MIC

remeasurements using the same susceptibility panel.

RESULTS

Nationwide sampling. From 2019 to 2020, more than 600 *P. aeruginosa* isolates were

transferred from Japanese hospitals to the Antimicrobial Resistance Research Center (AMR-RC)

at the National Institute of Infectious Diseases (NIID) based on the following criterion:

meropenem or imipenem MIC ≥8μg/ml in initial testing at the isolation site. All the collected

isolates were tested for antipseudomonal agent MICs using the Beckman MicroScan system at

AMR-RC. In total, 382 *P. aeruginosa* isolates showing meropenem MIC ≥8μg/ml in the tests at

AMR-RC were defined as CRPA and focused on in this study. The 382 CRPA isolates originated

from nonoverlapping patients in 78 hospitals (the number of collected isolates per island was as

follows: n = 30 from Hokkaido Island, n = 307 from Honshu Island, n = 7 from Shikoku Island, n =

32 from Kyusyu Island, and n = 6 from Okinawa Island). Source specimens were classified into

one of the following JANIS categories (19, 20): Oral/Endotracheal/Respiratory (n = 189),

Urinary/Genital (n = 76), Blood/Fluid (n = 40), Digestive (n = 27), and Other (n = 50). Table S1

provides more detailed source categories and metadata for each isolate.

Antimicrobial susceptibility profiles and STs of CRPA in Japan. Table 1 summarizes the

susceptibilities for antipseudomonal agents. Although the collected CRPA isolates were often

nonsusceptible to antipseudomonal agents other than meropenem, the percentage of

susceptible isolates remained 52.9% for piperacillin-tazobactam, 26.4% for ciprofloxacin, and

88.0% for amikacin (Table 1). Among the CRPA isolates, 27.7% (106/382) were deemed difficult-

to-treat resistance *P. aeruginosa* (DTR-*P. aeruginosa*) in Tamma *et al.*'s definition (27), and

85.8% (328/382) were MDR in Magiorakos *et al.*'s definition (28). Only three CRPA isolates were

resistant to colistin; two of the three colistin-resistant isolates (Table S1) were however DTR-

P.aeruginosa and elicited resistance to at least one agent from each of the seven antimicrobial

agent categories listed in Table 1.

The MLST profiles of 382 CRPA isolates were investigated in the PubMLST database (29).

Including newly assigned ST (ST4091–ST4125), 148 STs were identified (Fig. 1A). The most

frequently encountered clonal group (CG) was CG274 (comprising only ST274: type III secretion

system effector protein genotype $exoS^+/exoU^-$, 9.7%), followed by CG235 (ST235: $exoS^-/exoU^+$,

7.6%), CG253 (ST253: $exoS^-/exoU^+$, 4.5%), and CG155 (ST155 and ST4122: $exoS^+/exoU^-$, 3.4%).

Many minor STs, each contributing about 1%, occupied 49.0% of the total CRPA in Japan,

suggesting that clonal expansion is not common. All the top ten global high-risk clones (13)

were detected, accounting for 17.0% of the total CRPA in Japan (Table S1). A recent study on

CRPA (7) proposed globally most common CGs (10 CGs in Fig. 1B). Those globally common CGs

comprised 34.8%. Despite being among the top two most common CGs in the USA and Central

South America, CG111 remains a minor CG in Japan (Fig. 1B). Surprisingly, the dominant clone

CG463 (ST463), prevalent in Chinese clinical settings (7, 30), was entirely absent in the 382 CRPA

isolates collected in this study.

The proportions of the top eight most common CGs in Japan are shown per phenotype

group in Fig. 1C. In DTR-*P. aeruginosa*, the proportion of two major CGs, CG274 and CG235,

increased from 9.7% (37/382) in total CRPA to 11.3% (12/106) and from 7.6% (29/382) to 13.2%

(14/106), respectively (Fig. 1C), suggesting their propensity to accumulate MDR. However, when

focused on amikacin-nonsusceptible DTR-P. *aeruginosa* isolates (n = 25), the proportion of

CG274 decreased to 4.0% (1/25), while the proportion of CG235 increased to 36% (9/25),

highlighting the difference in their potential to acquire amikacin resistance.

Resistance mutations and carbapenemases in CRPA. The molecular mechanisms of

meropenem resistance are primarily classified into four types: porin inactivation, efflux pump

overproduction, target alteration, and horizontal acquisition of β-lactamase (2). Those

mechanisms are respectively associated with inactivation of the *oprD* gene, inactivation of

either *mexR*, *nalC*, or *nalD* genes encoding transcriptional regulators (repressors) for the

MexAB-OprM efflux pump (9–11), nonsynonymous substitutions in *ftsI* (12), and acquisition of

plasmid or ICE-borne carbapenemase genes (4, 5). We searched for the presence of inactivating

mutations (frameshift mutations due to insertion/deletion, nonsense mutations, and

inactivation of start codon or stop codon) in *oprD*, *mexR*, *nalC*, and *nalD* as well as resistance

mutations registered in the NCBI AMRFinderPluS database, as these likely contribute to

increasing meropenem MIC.

Database searches revealed that only 4.2% (16/382) of CRPA in Japan possessed

acquired carbapenemases. Known carbapenemase and MBL detected were GES-5 (n = 2), IMP-1

(n = 6), IMP-10 (n = 4), IMP-7 (n = 3), and IMP-34 (n = 1). No OXA-type carbapenemases were

detected. ST274 CRPA isolates rarely (2.7%, 1/37) carried carbapenemase, contrasting with the

34.5% (10/29) possession rate of ST235 isolates. One ST235 isolate encoded a novel variant of

IMP-type MBL, which was assigned a new allele name IMP-98 by NCBI.

The prevalence of inactivating mutations or known nonsynonymous resistance

mutations in chromosomal genes in 382 CRPA isolates was 87.1% (333/382) for *oprD*, 14.1%

(54/382) for *mexR*, 9.1% (35/382) for *nalD*, and 6.3% (24/382) for *nalC*. Collectively, 28.8%

(110/382) of CRPA carried an inactivating mutation in at least one of the three repressors for

mexAB. The most common mutation type in *oprD* was IN/DELs causing frameshift (237/382),

followed by nonsense mutations (90/382); these include, for instance, W277stop (24/90),

W339stop (9/90), and W138stop (9/90). In addition, 4.7% (18/382) of CRPA isolates carried

known resistance mutations in *ftsI* (P527S, F533L, R504C, or V471G) (12).

A simultaneous occurrence of resistance mutations and/or carbapenemase was

observed (Fig. 2). While 26.4% (101/382) of CRPA isolates had an inactivating mutation in both

oprD and at least one regulator for *mexAB*, 4.5% (17/382) had both *oprD* and *ftsI* mutations,

with 3.1% (12/382) exhibiting the *oprD* mutation and carbapenemase. The acquisition of

carbapenemase alone occurred only in 0.7% (4/382) of the CRPA. Notably, 9.1% (35/382) did

not carry identifiable inactivating mutations in the four focused genes, known *ftsI* resistance

mutations, or carbapenemase.

IMP-98 differed from IMP-1 by only one site at the nucleotide level and likely originated

from the bla_{IMP-1} gene cassette in the integron by C637T nonsynonymous substitution, leading

to P213S amino acid change (Fig. 3A). To deduce the origin of bla_{IMP-98}, we searched for the

bla_{IMP-98}-flanking sequences in the NCBI blast nr-database (2022.12.22). The top hit was an

integron gene cassette array found in a genome of an *E. coli* isolate from India (GenBank ID:

LC169568.1, Fig. 3B). This hit contains bla_{IMP-1} instead of bla_{IMP-98} in the gene cassette array. The

second hit was an integron region carrying bla_{GES-1} found in the chromosome of the *P.*

aeruginosa ST308 isolate from Mexico (GenBank ID: CP012755.1). This region was a class 1

integron as the contig contained specific motifs of *attI1* (5′-

<u>GTTATGG</u>AGCAGCAACGAT<u>GTTACGC</u>AGCAGGGCAGTCG<u>CCCTAAA</u>ACAAAG<u>TTAGGC</u>-3′: 7-bp core

sites are underlined) (30). Thus, bla_{IMP-98} might have originated from an ancestral integron like

that in LC169568.1.

To test whether IMP-98 is functional, bla_{IMP-98} was cloned and introduced into the drug-

susceptible *E. coli* strain DH5α, after which the susceptibility for various β-lactams was

investigated (Table 2). MICs of β-lactams except for aztreonam were considerably higher against

E. coli DH5α carrying bla_{IMP-98} than the control strain, confirming that IMP-98 retains the wide

substrate range of MBL.

Other resistance determinants. Among 46 amikacin-nonsusceptible CRPA isolates, 28.3%

(13/46) possessed known amikacin resistance genes *aac(6′)* variants, resistance mutations such

as *fusA1* (Q455R) and *amgS* (V121G) (31), or inactivating mutation in *mexZ* encoding a repressor

for the MexXY efflux pump contributing to aminoglycoside resistance (32) (Table S1). However,

73% (33/46) of amikacin-nonsusceptible CRPA did not possess identifiable amikacin resistance

genes or mutations. Among 281 ciprofloxacin nonsusceptible CRPA isolates, 74.3% (209/281)

possessed at least one of the quinolone resistance gene *crpP* (33) (166/281) and resistance

mutations in *gyrA* (T83I, D87N, or D87G; 99/281), *gyrB* (E468D, S466F, or S466Y; 13/281), or

parE (A473V; 4/281) (Table S1). Three colistin-resistant isolates did not possess any known

colistin resistance mutations or genes.

DISCUSSION

Before the global launch of national action plans on antimicrobial resistance, the endemic

spread of ST235 clones was occasionally reported in Japan (34–36). This was in part because the

Ministry of Health, Labor and Welfare of Japan monitors the occurrence of amikacin-

nonsusceptible, carbapenem and fluoroquinolone-resistant *P. aeruginosa* as a part of the

National Epidemiological Surveillance of Infectious Diseases program (37). However, national

surveillance of clinically more common CRPAs remained lacking until the present study. As a

country with a low prevalence rate of meropenem-resistant CRPA, Japan is expected to have a

low carbapenemase possession rate of *P. aeruginosa*. This study demonstrates that the

carbapenemase possession rate of meropenem-resistant CRPA in Japan is 4.2%, which is among

the lowest in the regions where surveillance studies have been performed to date: the

carbapenemase possession rate of meropenem-resistant CRPA was 32.9% in the Asia-Pacific

region in 2015–2019 (16), 1.9% for the USA in 2018–2019 (7), 69% for South and Central

America in 2018–2019 (7), 32% for China in 2018–2019 (7), and 21.9% for Europe in 2017–2019

(17).

The ciprofloxacin and amikacin susceptibility rates regarding CRPA were 26.4% and 88.0%,

respectively, in Japan. These values exceed 20.6% and 60.6% of pooled CRPA values reported in

the Asia-Pacific region (16), and are then equivalent to and higher than 26.2% and 63.4% of the

global CRPA population, respectively (17). Furthermore, the piperacillin-tazobactam

susceptibility rate in Japan was 52.9%, which contrasts with the 25.3% susceptibility rate in the

global CRPA population (17). This difference might reflect the difference in the carbapenemase

possession rate in CRPA between Japan and other regions. Therefore, in Japan, MDR is not

heavily perceived in the *P. aeruginosa* population in clinical settings compared with other Asia-

Pacific regions or the world average.

The most frequent ST in the meropenem-resistant CRPA in Japan was ST274 but not

ST235 (*n* = 37 vs. *n* = 29). ST274 and CG274 members have frequently been detected in clinical

settings worldwide (14) and are also known for chronic infection and intrabody evolution

involving MDR in patients with cystic fibrosis in Europe (38, 39). We do not know whether the

ST274 isolates collected in this study are associated with a chronic lung infection, but ST274 was

rather overrepresented in the "Oral/Endotracheal/Respiratory" source group (odds ratio = 1.84,

P = 0.054 in one-tailed Fisher's exact test). Since patients with cystic fibrosis are rare in Japan,

ST274 might generally be fitter in the respiratory systems than other STs.

Inactivating mutation in *oprD* was the most frequent meropenem-resistance mechanism

(found in 87.1%), as noted in previous genome surveillance performed in other regions; 69% of

meropemen-resistant CRPA possessed *oprD*-inactivating mutations and 22% possessed acquired

carbapenemase (7). We show that in Japan, not only *oprD*-inactivating mutations but also efflux

pump regulator inactivation and *ftsI* resistance mutations are more common carbapenem-

resistance mechanisms than the horizontal acquisition of carbapenemases. Therefore, future

surveillance of *P. aeruginosa* should prioritize the study of chromosomal mutations.

Since several countries/regions normally categorized in the Asia-Pacific region are

geographically isolated by the presence of sea, we expected to observe distinct features of the

CRPA genotype in Japan and other Asia-Pacific regions. The striking feature was the

predominance of CG463/ST463 in China (7) and its absence in Japan (Fig. 1B). Among the

carbapenemase types, IMP was the most frequent (87.5%; 14/16) in Japan. This trend contrasts

with the predominance of NDM in *P. aeruginosa* borne-carbapenemases in India (16), KPC in

China (7), and VIM in Europe, Africa/Middle East, and Latin America (17).

Surprisingly, 9.2% (35/382) of the CRPA isolates did not possess apparent inactivating

mutations, known *ftsI* resistance mutations, or carbapenemase. Furthermore, known resistance

mutations could not account for the mechanisms of amikacin nonsusceptibility in 73% of

amikacin-nonsusceptible isolates. These CRPA isolates must possess unreported resistance

genes or mutations in their genomes. A comparative genomics approach is required to clarify

this claim in the future to fully understand and control CRPA spread.

MATERIALS AND METHODS

Strains, media, and plasmids. CRPA candidates participating in our genome surveillance

program (the Japan Antimicrobial-Resistant Bacterial Surveillance [JARBS]) and associated with

isolation source and date information were transferred from hospitals to the NIID AMR center.

Single colony isolation was performed if necessary, using BD BBL™ BTB Lactose-Contained Agar

Medium (also called Drigalski agar) (Becton, Dickinson and Company, Franklin Lakes, NJ, USA).

BD BBL™ Muller–Hinton II (Cation adjusted) Broth (Becton, Dickinson and Company) was used

to culture *P. aeruginosa* strains in liquid. Table S1 contains the *P. aeruginosa* strains and

associated metadata.

E. coli strain DH5α [F⁻, Φ80d*lacZ*ΔM15, Δ(*lacZYA-argF*)U169, *deoR*, *recA*1, *endA*1,

*hsdR*17(r_K^-, m_K^+), *phoA*, *supE*44, λ⁻, *thi*-1, *gyrA*96, *relA*1] was used for DNA cloning and

functional analysis of *bla*$_{IMP-98}$. DH5α was cultured using BD Difco™ LB, Lennox Broth and agar

(Becton, Dickinson and Company). A gene cassette array of *bla*$_{OXA-10}$-*acc(6')-bla*$_{IMP-98}$ in *P.*

aeruginosa ST235 strain JBBCAEG-19-0032 was PCR-amplified using primers blaIMP_F: 5′-

TCGAGCTCGGTACCCCGCTACTTGAAGTGTTGACGC-3′ and blaIMP_R: 5′-

CTCTAGAGGATCCCCTTAGTTGCTTGGTTTTGATGGTTTTTTACTTTCG-3′, and KODone polymerase

(TOYOBO CO., LTD., Osaka, Japan). A low-copy number vector pMW218 (*nptII*, pSC101 replicon;

NIPPON GENE CO., LTD., Tokyo, Japan) was also PCR-amplified with primers pMW_mcs_F: 5′-

GGGTACCGAGCTCGAATTCGTA-3′ and pMW_mcs_R: 5′-GGGGATCCTCTAGAGTCGACCT-3′. The

two PCR products were joined using NEBuilder (New England Biolabs, Ipswich, MA, USA). The

resulting construct was named pHY1399. A segment on pHY1399 was PCR-amplified to delete

the *bla*$_{OXA-10}$-*acc(6')-Ib* region using 5′ phosphorylated primers P_Integ_del_F: 5′P-

CATGGCACCTTCTTGGTGGCTAACG-3′, and P_Integ_del_R: 5′P-GAGCGAACACGCAGTGATGCC-3′,

then the purified PCR product was ligated using Ligation High ver.2 (Toyobo) and introduced into

DH5α. The resulting recombinant plasmid obtained from one transformant was named

pHY1401.

Antimicrobial susceptibility test. To obtain MICs of antipseudomonal agents, CRPA strains were

propagated on Drigalski agar and incubated for 1 day at 35°C. Then, a single colony was

inoculated into MicroScan Neg MIC NF-1J panel using Prompt Inoculation System D (Beckman

Coulter Inc., Brea, CA, USA). Inoculated panels were incubated at 35°C for 18 h in MicroScan

Walkway40 Plus (Beckman Coulter Inc.) or MicroScan Walkway DxM 1096 (Beckman Coulter

Inc.). To cover a therapeutically achievable range, the antimicrobial susceptibility testing was

designed to measure major antimicrobials using five dilutions (26). To obtain β-lactam MIC

against two *E. coli* strains DH5α (pMW218) and DH5α (pHY1401), the strains were first

propagated on LB agar containing kanamycin (25 μg/ml) and incubated at 37°C overnight, then

the colonies were diluted and inoculated into the Neg MIC EN-2J panel containing β-lactams

listed in Table 2 and incubated in MicroScan Walkway40 Plus (Beckman Coulter Inc.).

Breakpoint criteria for susceptible (S), intermediate resistance (I), and resistance (R) followed

the CLSI 2022 guidelines M100-Ed32. The definition of DTR-*P. aeruginosa* followed Tamma *et*

al.'s criteria (27): nonsusceptible to all the eight antimicrobials belonging to either five

antipseudomonal agent categories, and is based on the following MIC criteria:

piperacillin/tazobactam ≥32/4 μg/ml, ceftazidime ≥16 μg/ml, cefepime ≥16 μg/ml, meropenem

≥4 μg/ml, imipenem ≥4 μg/ml, aztreonam ≥16 μg/ml, ciprofloxacin ≥1 μg/ml, and levofloxacin

≥2 μg/ml (I or R in CLSI 2022). The definition of MDR was modified from Magiorakos *et al.*'s

criteria (28) as follows: nonsusceptible to at least one agent in at least three categories except

for polymyxin listed in Table 1.

Genome sequencing. *P. aeruginosa* strains were cultured in Muller–Hinton II broth overnight at

35°C. Cells were collected in 1.5 mL tubes and resuspended in a lysis solution containing

lysozyme and RNaseA, then incubated at 37°C for 1 h. After adding proteinase K and SDS, the

reaction mixture was incubated at 55°C overnight. Genomic DNA was purified from the cell

lysates using Agencourt AMPure XP beads (Beckman Coulter Inc.) following the protocol

recommended by the manufacturer. NGS libraries were constructed using Enzymatics 5× WGS

fragmentation mix and WGS ligase reagents (Qiagen, Hilden, Germany), and then libraries were

sequenced on HiSeq X Five platform in Macrogen Japan (Tokyo, Japan). Low-quality reads were

removed from the original reads using fastp (40), after which the trimmed reads were

assembled using Shovill pipeline (https://github.com/tseemann/shovill/tree/master) set to

default options. Genomic assemblies fulfilled L90 < 150.

Data analysis. The PubMLST ST number was assigned by searches against the PubMLST

database (29). When hit ST was not present in PubMLST, MLST profiles of the isolates were

submitted to PubMLST, after which new ST numbers (ST4091-ST4125) were assigned to the new

MLST profiles. The relationship between MLST profiles was represented as a minimum spanning

tree by GrapeTree software (41). Acquired antimicrobial resistance genes in draft genome

assemblies were searched for using AMRFinderPlus (database ver. 2023-04-17.1) with the "--

organism Pseudomonas_aeruginosa" option, otherwise default setting (42). The

presence/absence and integrity of specific chromosomal genes (frameshift mutations, nonsense

mutations, and gene fusions caused by substitutions at stop/start codon positions) were

determined by tblastn searches against an in-house generated nucleotide database using blast

v2.12.0 (43), and the following retrieval of hit sequences and inspection of the alignments. The

RefSeq/GenBank accession numbers of blast queries used are as follows: ExoU,

WP_034024595.1; ExoS, NP_252530.1; FtsI (PBP3), WP_003094139.1; MexR, NP_249115.1;

NalC, NP_252410.1; NalD, NP_252264.1; OprD, NP_249649.1; MexZ, WP_003088626.1. The

alignments were generated using MAFFT v 7.490 with the einsi option (44). The integron

integrase gene and *attC* were searched for using IntegronFinder 2.0.2 with the default

parameter setting (45).

Data availability. Raw sequence reads generated in this study are deposited under DDBJ

Sequence Read Archive (DRA) accession number DRA017464. Draft genome assemblies are

available from Zenodo repository [DOI: 10.5281/zenodo.10081563] (46). PubMLST allele

profiles of 382 isolates and sequence alignments of *oprD*, *mexR*, *mexZ*, *nalC*, *nalD*, and *ftsI* are

available from Zenodo repository [DOI: 10.5281/zenodo.10081526] (47). The nucleotide

sequence of bla_{IMP-98} is deposited under GenBank/DDBJ/EMBL accession number LC740578.1.

ACKNOWLEDGMENTS

We thank all the hospitals contributing to the JARBS program. We thank Satoyo Wakai, Takahisa

Ishizuka, Liansheng Yu, Mikihisa Okuda, Akira Moriya, Emi Fujimura, Koichi Shimakawa, Rumi

Oki, Yoshie Taki, Mayumi Sasada, Mikako Nakazawa, Noriko Sakamoto, Elahi Shaheem, Chika

Arai, Yuko Kazumi, Akemi Suzuki, and Fumiko Hamamoto at AMR-RC NIID for their support of

isolate collection, susceptibility test, genomic DNA preparation, and JANIS data collection. We

thank Enago [https://www.enago.jp] for language editing. The computation is supported by

supercomputer SHIROKANE of Human Genome Center at the Institute of Medical Science

University of Tokyo, and NIG supercomputer at the Research Organization of Information and

Systems National Institute of Genetics. This research is supported by Japan Agency for Medical

Research and Development under grant number 23fk0108604 (to M.S.). The funder has no role

in study design, data collection, data analysis, and decision to publish results.

Author contributions are as follows: Conceptualization, H.Y., W.H., K.Y., and M.S.;

Investigation, H.Y., W.H., Sa.K., S.A., E.A., Sh.K., N.K., A.H., T.K., Y.S., and K.Y.; Formal analysis, H.Y.

and K.Y.; Data Curation, H.Y., N.K., and K.Y.; Methodology, Y.S.; Visualization, H.Y.; Writing -

Original Draft, H.Y.; Writing - Review & Editing, H.Y., W.H., N.K., A.H., S.K., Y.S., K.Y., and M.S.;

Project Administration and Funding Acquisition, M.S.. All authors reviewed and approved the

final manuscript.

ETHICAL STATEMENT

This study (JARBS) was reviewed and approved by the Institutional Review Board of the

National Institute of Infectious Diseases (approval number 1251). The use of JANIS data was

approved by Ministry of Health, Labor and Welfare of Japan (approval number 1553).

REFERENCES

1.   **GBD, 2019 Antimicrobial Resistance Collaborators**. 2022. Global mortality associated

with 33 bacterial pathogens in 2019: a systematic analysis for the Global Burden of

Disease Study 2019. Lancet **400**:2221–2248.

2.   **Poole K**. 2011. *Pseudomonas aeruginosa*: resistance to the max. Front Microbiol **2**:65.

3.   **Horcajada JP, Montero M, Oliver A, Sorlí L, Luque S, Gómez-Zorrilla S, Benito N, Grau S**.

2019. Epidemiology and treatment of multidrug-resistant and extensively drug-resistant

Pseudomonas aeruginosa Infections. Clin Microbiol Rev **32**:e00031–19.

4.   **Senda K, Arakawa Y, Nakashima K, Ito H, Ichiyama S, Shimokata K, Kato N, Ohta M**. 1996.

Multifocal outbreaks of metallo-β-lactamase-producing *Pseudomonas aeruginosa*

resistant to broad-spectrum beta-lactams, including carbapenems. Antimicrob Agents

Chemother **40**:349–353.

5.   **Botelho J, Roberts AP, León-Sampedro R, Grosso F, Peixe L**. 2018. Carbapenemases on

the move: it's good to be on ICEs. Mob DNA **9**:37.

6.   **World Health Organization**. 2017. WHO priority pathogens list for R&D of new antibiotics.

https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-

new-antibiotics-are-urgently-needed Accessed Nov 8 2023.

7.   **Reyes J, Komarow L, Chen L, Ge L, Hanson BM, Cober E, Herc E, Alenazi T, Kaye KS,**

Garcia-Diaz J, Li L, Kanj SS, Liu Z, Oñate JM, Salata RA, Marimuthu K, Gao H, Zong Z,

Valderrama-Beltrán SL, Yu Y, Tambyah P, Weston G, Salcedo S, Abbo LM, Xie Q, Ordoñez

K, Wang M, Stryjewski ME, Munita JM, Paterson DL, Evans S, Hill C, Baum K, Bonomo RA,

Kreiswirth BN, Villegas MV, Patel R, Arias CA, Chambers HF, Fowler VG, Doi Y, van Duin

D, Satlin MJ, Antibacterial Resistance Leadership Group and Multi-Drug Resistant

Organism Network Investigators. 2023. Global epidemiology and clinical outcomes of

carbapenem-resistant *Pseudomonas aeruginosa* and associated carbapenemases (POP): a

prospective cohort study. Lancet Microbe **4**:e159–e170.

8.   **Sanbongi Y, Shimizu A, Suzuki T, Nagaso H, Ida T, Maebashi K, Gotoh N**. 2009.

Classification of OprD sequence and correlation with antimicrobial activity of carbapenem

agents in *Pseudomonas aeruginosa* clinical isolates collected in Japan. Microbiol Immunol

53:361–367.

9.   **Srikumar R, Paul CJ, Poole K**. 2000. Influence of mutations in the *mexR* repressor gene on

expression of the MexA-MexB-oprM multidrug efflux system of *Pseudomonas aeruginosa*.

J Bacteriol **182**:1410–1414.

10. **Cao L, Srikumar R, Poole K**. 2004. MexAB-OprM hyperexpression in NalC-type multidrug-

resistant *Pseudomonas aeruginosa*: identification and characterization of the *nalC* gene

encoding a repressor of PA3720-PA3719. Mol Microbiol **53**:1423–1436.

11. **Morita Y, Cao L, Gould VC, Avison MB, Poole K**. 2006. *nalD* encodes a second repressor of

the *mexAB-oprM* multidrug efflux operon of *Pseudomonas aeruginosa*. J Bacteriol

188:8649–8654.

12. **Cabot G, Florit-Mendoza L, Sánchez-Diener I, Zamorano L, Oliver A**. 2018. Deciphering β-

lactamase-independent β-lactam resistance evolution trajectories in *Pseudomonas*

aeruginosa. J Antimicrob Chemother **73**:3322–3331.

13. **Del Barrio-Tofiño E, López-Causapé C, Oliver A**. 2020. *Pseudomonas aeruginosa* epidemic

high-risk clones and their association with horizontally-acquired β-lactamases: 2020

update. Int J Antimicrob Agents **56**:106196.

14. **Oliver A, Mulet X, López-Causapé C, Juan C**. 2015. The increasing threat of *Pseudomonas*

aeruginosa high-risk clones. Drug Resist Updat **21-22**:41–59.

15. **Karlowsky JA, Bouchillon SK, El Mahdy Kotb R, Mohamed N, Stone GG, Sahm DF**. 2022.

Carbapenem-resistant Enterobacterales and *Pseudomonas aeruginosa* causing infection in

Africa and the Middle East: a surveillance study from the ATLAS programme (2018-20).

JAC Antimicrob Resist **4**:dlac060.

16. **Lee YL, Ko WC, Hsueh PR**. 2022. Geographic patterns of carbapenem-resistant

Pseudomonas aeruginosa in the Asia-Pacific region: results from the antimicrobial testing

leadership and surveillance (ATLAS) program, 2015-2019. Antimicrob Agents Chemother

66:e0200021.

17. **Gales AC, Stone G, Sahm DF, Wise MG, Utt E**. 2023. Incidence of ESBLs and

carbapenemases among Enterobacterales and carbapenemases in *Pseudomonas*

aeruginosa isolates collected globally: results from ATLAS 2017-2019. J Antimicrob

Chemother dkad127.

18. **World Health Organization**. 2022. Global Antimicrobial Resistance and Use Surveillance

System (GLASS) Report: 2021 Report electronic Supplementary Material. URL:

https://www.who.int/publications/i/item/9789240027336 Accessed Nov 8 2023.

19. **Kajihara T, Yahara K, Hirabayashi A, Shibayama K, Sugai M**. 2021. Japan Nosocomial

Infections Surveillance (JANIS): current status, international collaboration, and future

directions for a comprehensive antimicrobial resistance surveillance system. Jpn J Infect

Dis **74**:87–96.

20. **Ministry of Health Labour and Welfare of Japan**. 2021. JANIS-Open-Report-Clinical-

Laboratory-Division. https://janis.mhlw.go.jp/english/report/index.html Accessed Nov 8

2023.

21. **Yang W, Ding L, Han R, Yin D, Wu S, Yang Y, Zhu D, Guo Y, Hu F, CHINT-Study-Group**.

2023. Current status and trends of antimicrobial resistance among clinical isolates in

China: a retrospective study of CHINET from 2018 to 2022

on behalf of the China Antimicrobial Surveillance Network (CHINET) Study Group. One Health

Advances **1**:8.

22. **European Centre for Disease Prevention and Control**. 2022. Antimicrobial resistance in

the EU/EEA (EARS-Net) - Annual Epidemiological Report for 2020.

https://www.ecdc.europa.eu/sites/default/files/documents/EUEEA_trend_tables_for_ind

ividual_isolates_2020.pdf Accessed Nov 8 2023.

23. **Tojo M, Tada T, Shimojima M, Tanaka M, Narahara K, Miyoshi-Akiyama T, Kirikae T,**

Ohmagari N. 2014. Dissemination in Japan of multidrug-resistant Pseudomonas

aeruginosa isolates producing IMP-type metallo-β-lactamases and AAC(6')-lae/AAC(6')-lb.

J Infect Chemother **20**:586–588.

24. **Miyoshi-Akiyama T, Tada T, Ohmagari N, Viet Hung N, Tharavichitkul P, Pokhrel BM,**

Gniadkowski M, Shimojima M, Kirikae T. 2017. Emergence and spread of epidemic

multidrug-resistant *Pseudomonas aeruginosa*. Genome Biol Evol **9**:3238–3245.

25. **Tokimatsu I, Matsumoto T, Tsukada H, Fujikura Y, Miki M, Morinaga Y, Sato J,**

Wakamura T, Kiyota H, Tateda K, Yanagisawa H, Sasaki T, Ikeda H, Horikawa H,

Takahashi H, Seki M, Mori Y, Takeda H, Kurai D, Hasegawa N, Uwamino Y, Kudo M,

Yamamoto M, Nagano Y, Nomura S, Tetsuka T, Hosokai M, Aoki N, Yamamoto Y, Iinuma

Y, Mikamo H, Suematsu H, Maruyama T, Kawabata A, Sugaki Y, Nakamura A, Fujikawa Y,

Fukumori T, Ukimura A, Kakeya H, Niki M, Yoshida K, Kobashi Y, Tokuyasu H, Yatera K,

Ikegami H, Fujita M, Matsumoto T, Yanagihara K, Matsuda J, Hiramatsu K, Shinzato T.

2023. Nationwide surveillance of bacterial respiratory pathogens conducted by the

surveillance committee of the Japanese Society of Chemotherapy, the Japanese

Association for Infectious Diseases, and the Japanese Society for Clinical Microbiology in

2019-2020: General view of the pathogens' antibacterial susceptibility. J Infect Chemother

29:731–743.

26. **Kayama S, Yahara K, Sugawara Y, Kawakami S, Kondo K, Zuo H, Kutsuno S, Kitamura N,**

Hirabayashi A, Kajihara T, Kurosu H, Yu L, Suzuki M, Hisatsune J, Sugai M. 2023. National

genomic surveillance integrating standardized quantitative susceptibility testing clarifies

antimicrobial resistance in Enterobacterales. Nat Comm **14**:1–13.

27. **Tamma PD, Aitken SL, Bonomo RA, Mathers AJ, van Duin D, Clancy CJ**. 2021. Infectious

Diseases Society of America Guidance on the Treatment of Extended-Spectrum β-

lactamase Producing Enterobacterales (ESBL-E), Carbapenem-Resistant Enterobacterales

(CRE), and *Pseudomonas aeruginosa* with Difficult-to-Treat Resistance (DTR-*P. aeruginosa*).

Clin Infect Dis **72**:1109–1116.

28. **Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S,**

Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens MJ,

Vatopoulos A, Weber JT, Monnet DL. 2012. Multidrug-resistant, extensively drug-

resistant and pandrug-resistant bacteria: an international expert proposal for interim

standard definitions for acquired resistance. Clin Microbiol Infect **18**:268–281.

29. **Jolley KA, Bray JE, Maiden MCJ**. 2018. Open-access bacterial population genomics:

BIGSdb software, the PubMLST.org website and their applications. Wellcome Open Res

3:124.

30. **Partridge SR, Recchia GD, Scaramuzzi C, Collis CM, Stokes HW, Hall RM**. 2000. Definition

of the *attI1* site of class 1 integrons. Microbiology (Reading) **146**:2855–2864.

31.   **Lau CH, Fraud S, Jones M, Peterson SN, Poole K**. 2013. Mutational activation of the

479   AmgRS two-component system in aminoglycoside-resistant *Pseudomonas aeruginosa*.

480   Antimicrob Agents Chemother **57**:2243–2251.

481   32.   **Morita Y, Tomida J, Kawamura Y**. 2012. Primary mechanisms mediating aminoglycoside

482   resistance in the multidrug-resistant *Pseudomonas aeruginosa* clinical isolate PA7.

483   Microbiology (Reading) **158**:1071–1083.

484   33.   **Chávez-Jacobo VM, Hernández-Ramírez KC, Romo-Rodríguez P, Pérez-Gallardo RV,**

485   **Campos-García J, Gutiérrez-Corona JF, García-Merinos JP, Meza-Carmen V, Silva-Sánchez**

486   **J, Ramírez-Díaz MI**. 2018. CrpP Is a Novel Ciprofloxacin-Modifying Enzyme Encoded by the

487   *Pseudomonas aeruginosa* pUM505 Plasmid. Antimicrob Agents Chemother **62**:e02629–17.

488   34.   **Shimizu W, Kayama S, Kouda S, Ogura Y, Kobayashi K, Shigemoto N, Shimada N, Yano R,**

489   **Hisatsune J, Kato F, Hayashi T, Sueda T, Ohge H, Sugai M**. 2015. Persistence and epidemic

490   propagation of a *Pseudomonas aeruginosa* sequence type 235 clone harboring an IS*26*

491   composite transposon carrying the *bla*$_{IMP-1}$ integron in Hiroshima, Japan, 2005 to 2012.

492   Antimicrob Agents Chemother **59**:2678–2687.

493   35.   **Hishinuma T, Tada T, Kuwahara-Arai K, Yamamoto N, Shimojima M, Kirikae T**. 2018.

494   Spread of GES-5 carbapenemase-producing *Pseudomonas aeruginosa* clinical isolates in

495   Japan due to clonal expansion of ST235. PLoS One **13**:e0207134.

496   36.   **Hishinuma T, Uchida H, Tohya M, Shimojima M, Tada T, Kirikae T**. 2020. Emergence and

497   spread of VIM-type metallo-β-lactamase-producing *Pseudomonas aeruginosa* clinical

498   isolates in Japan. J Glob Antimicrob Resist **23**:265–268.

499   37.   **Suzuki S**. 2021. A view on 20 years of antimicrobial resistance in Japan by two national

500   surveillance systems: the national epidemiological surveillance of infectious diseases and

501   Japan nosocomial infections surveillance. Antibiotics (Basel) **10**:1189.

502   38.   **Fernández-Olmos A, García-Castillo M, Alba JM, Morosini MI, Lamas A, Romero B, Galán**

503   **JC, del Campo R, Cantón R**. 2013. Population structure and antimicrobial susceptibility of

504   both nonpersistent and persistent Pseudomonas aeruginosa isolates recovered from

505   cystic fibrosis patients. J Clin Microbiol **51**:2761–2765.

506   39.   **López-Causapé C, Sommer LM, Cabot G, Rubio R, Ocampo-Sosa AA, Johansen HK,**

507   **Figuerola J, Cantón R, Kidd TJ, Molin S, Oliver A**. 2017. Evolution of the *Pseudomonas*

508   *aeruginosa* mutational resistome in an international Cystic Fibrosis clone. Sci Rep **7**:5555.

509   40.   **Chen S, Zhou Y, Chen Y, Gu J**. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

510   Bioinformatics **34**:i884–i890.

511   41.   **Zhou Z, Alikhan NF, Sergeant MJ, Luhmann N, Vaz C, Francisco AP, Carriço JA, Achtman**

512   **M**. 2018. GrapeTree: visualization of core genomic relationships among 100,000 bacterial

513   pathogens. Genome Res **28**:1395–1404.

514   42.   **Feldgarden M, Brover V, Gonzalez-Escalona N, Frye JG, Haendiges J, Haft DH, Hoffmann**

515   **M, Pettengill JB, Prasad AB, Tillman GE, Tyson GH, Klimke W**. 2021. AMRFinderPlus and

516   the Reference Gene Catalog facilitate examination of the genomic links among

517   antimicrobial resistance, stress response, and virulence. Sci Rep **11**:12728.

518   43.   **Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ**. 1990. Basic local alignment search

519   tool. J Mol Biol **215**:403–410.

520   44.   **Katoh K, Standley DM**. 2013. MAFFT multiple sequence alignment software version 7:

521   improvements in performance and usability. Mol Biol Evol **30**:772–780.

522   45.   **Néron B, Littner E, Haudiquet M, Perrin A, Cury J, Rocha EPC**. 2022. IntegronFinder 2.0:

523   identification and analysis of integrons across bacteria, with a focus on antibiotic

524   resistance in *Klebsiella*. Microorganisms **10**:700.

525   46.   **Antimicrobial Resistance Research Center, National Institute of Infectious Diseases**.

526   2023. Genome assemblies of 382 carbapenem-resistant *Pseudomonas aeruginosa* isolates

527   collected from Japanese hospitals in 2019–2020. Zenodo doi: 10.5281/zenodo.10081563

528   Accessed Nov 13 2023.

529   47.   **Yano H**. 2023. PubMLST allele profiles and sequence alignments of 382 carbapenem-

530   resistant *Pseudomonas aeruginosa* isolates collected from Japanese hospitals in

531   2019–2020. Zenodo doi: 10.5281/zenodo.10081526 Accessed Nov 13 2023.

SUPPLEMENTAL MATERIAL

Table S1. Susceptibility profile, genotype, and isolation source of 382 CRPA isolates.

FIGURE LEGENDS

FIG 1 (A) Minimum spanning tree of 382 meropenem-resistant CRPA isolates based on MLST

profiles. Each node represents ST, otherwise clonal group (CG). STs with one allele difference

were collapsed into one node as CG. Node size is proportional to the number of isolates. The

number on the branch denotes the number of different alleles at the 7 MLST loci between two

nodes. (B) Proportions of 10 globally most common CGs in Japan and other regions. Proportion

of CGs in the USA, China, and South and Central America are based on Reyes *et al.*'s report (7).

(C) Proportion of eight most common CGs in four phenotype groups in Japan. AMK non-S DTR,

AMK nonsusceptible DTR-*P. aeruginosa*.

FIG 2. Venn diagram showing the simultaneous occurrence of resistance mutations in

meropenem-resistant *P. aeruginosa*. Numbers in circles represent the numbers of isolates

having an acquired carbapenemase, a point mutation registered in AMRFinderPlus (42), or

inactivating mutations. The "Mex" category indicates the presence of a mutation in at least one

of three loci (*mexR*, *nalC,* and *nalD*) encoding a negative regulator for the *mexAB* operon. The

"PBP3" category indicates the presence of a mutation in *ftsI*. The acquired

carbapenemase/MBLs detected are GES-5, IMP-1, IMP-10, IMP-7, IMP-34, and IMP-98.

FIG 3 (A) Alignment of carboxyl-terminal regions of IMP variants. H, helix; e, sheet. IMP-98

differs from IMP-1 only at P213 indicated by an arrowhead. (B) Structure of class 1 integrons of *P.*

aeruginosa ST235 strain JBBCAEG-19-0032, *E. coli* strain MBL1-07, and *P. aeruginosa* ST308

strain Pa38. The *tniABQR* genes are homologues of Tn*402* family transposon genes. *bla*_{OXA-10} and

aac(6')-Ib overlapped by 23 bp. *intI* and *attC* were identified by IntegronFinder2.0 (45).

TABLES

TABLE 1. Antimicrobial susceptibilities of 382 meropenem-resistant CRPA isolates.

Antimicrobial agents	Percentage of susceptibility group (number of isolates)		
	S	I	R
Antipseudomonal carbapenems			
Imipenem	1.0 (4)	2.9 (11)	96.1 (367)
Meropenem	0.0 (0)	0.0 (0)	100 (382)
Doripenem	10.7 (41)	36.6 (136)	53.7 (205)

Antipseudomonal cephalosporins			
Ceftazidime	53.1 (203)	12.8 (49)	34.0 (130)
Cefepime	47.4 (181)	27.2 (104)	25.4 (97)
Antipseudomonal penicillin + inhibitor			
Piperacillin-tazobactam	52.9 (202)	34.8 (133)	22.8 (87)
Monobactam			
Aztreonam	18.3 (70)	29.8 (114)	51.8 (198)
Antipseudomonal fluoroquinolones			
Levofloxacin	27.5 (105)	20.7 (79)	51.8 (198)
Ciprofloxacin	26.4 (101)	35.6 (136)	38.0 (145)
Aminoglycosides			
Gentamycin	63.4 (242)	22.3 (85)	14.4 (55)
Tobramycin	88.7 (339)	2.1 (8)	9.2 (35)
Amikacin	88.0 (336)	4.2 (16)	7.1 (27)
Polymyxin			
Colistin	-	99.2 (379)	0.8 (3)

566   [a] Criteria for S/I/R follows CLSI M100-Ed32 (2022).

568   **TABLE 2.** β-lactam MICs against *E. coli* strain carrying bla_{IMP-98}.

Antimicrobials	DH5α(pMW218)	DH5α(pHY1401)[a]
Ampicillin	≤4	>16
Ampicillin-sulbactam	≤4/2	>16/8
Amoxicillin-clavulanate	≤8/4	>16/8
Piperacillin	≤4	32
Piperacilin-tazobactam	≤4	8
Cefazolin	≤1	>16
Cefotiam	≤8	16
Cefaclor	≤8	>16
Cefmetazole	≤4	>32
Cefotaxime	≤0.5	>2
Ceftriaxone	≤0.5	>2
Ceftazidime	≤1	>8
Cefpodoxime	≤1	>4
Cefditoren	≤1	>2
Cefoperazone-sulbactam	≤8/4	>32/16
Cefepime	≤1	8
Imipenem	≤0.5	2

Meropenem	≤0.25	2
Ertapenem	≤0.25	>1
Flomoxef	≤8	32
Aztreonam	≤1	≤1

569 [a] pHY1401 is a pMW218 derivative carrying bla_{IMP-98} and the putative promoter region of class 1

570 integron in strain JBBCAEG-19-0032.

A

```
                   179                                      ▼                           246
gb_CAA11471.1_IMP-1   179 KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVPSHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN 246
WP_268871851.1_IMP-98 179 KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVSSHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN 246
gb_BAB15941.1_IMP-6   179 KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVPGHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN 246
gb_BAB72069.1_IMP-10  179 KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVPSHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN 246
gb_BAP75826.1_IMP-34  179 KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVPSHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN 246
gb_AAK12087.1_IMP-7   179 KPYGLGNLGDANLEAWPKSAKLLVSKYGKAKLVVPSHSEVGDASLLKRTLEQAVKGLNESKKLSKPSN 246
gb_CAB94707.1_IMP-2   179 KPDGLGNLGDANLEAWPKSAKILMSKYVKAKLVVSSHSEIGDASLLKRTWEQAVKGLNESKKPSQPSN 246
gb_ALT53802.1_IMP-56  179 KPDGLGNLGDANLEAWPKSAKILMSKYGKAKLVVSGHSEIGNASLLQRTWEQAVKGLNESKKPLQPSS 246
Consensus_aa:             KP.GLGNLGDANlEAWPKSAKlLbSKYGKAKLVVstHSElGsASLLp.ThEQAVKGLNESKKP.pPSs
Consensus_ss:                     hhhhhhhhhhhhh      eeeee      hhhhhhhhhhhhhhhhhhhhhhhh
```

B

E. coli MBL1-07
LC169568.1
1 - 9,129
India

P. aeruginosa
JBBCAEG-19-0032 contig_133
1 - 5,316
(complementary)
Japan

P. aeruginosa Pa38
chromosome
CP021775.1
3,658,672 - 3,664,162
(complementary)
Mexico

○ *attI*   ● *attC*

80 85 90 95 100 (%)
blastn