Nationwide genome surveillance of carbapenem-resistant *Pseudomonas*

aeruginosa in Japan

Hirokazu Yano[1*], Wataru Hayashi[1], Sayoko Kawakami[1], Sadao Aoki[1], Eiko Anzai[1], Hui Zuo[1],

Norikazu Kitamura[1], Aki Hirabayashi[1], Toshiki Kajihara[1], Shizuo Kayama[1], Yo Sugawara[1], Koji

Yahara[1], Motoyuki Sugai[1*]

[1]Antimicrobial Resistance Research Center, National Institute of Infectious Diseases. 4-2-1

Aobacho, Higashimurayama, Tokyo 189-0002, Japan

*Corresponding author: Antimicrobial Resistance Research Center, National Institute of

Infectious Diseases. 4-2-1 Aobacho, Higashimurayama, Tokyo 189-0002, Japan.

Email address: h-yano@niid.go.jp (H Yano); sugai@niid.go.jp (M Sugai)

ABSTRACT

Japan is a country with an approximate 10 % prevalence rate of carbapenem-resistant

Pseudomonas aeruginosa (CRPA). Currently, a comprehensive overview of the genotype and

phenotype patterns of CRPA in Japan is lacking. Herein, we conducted genome sequencing and

quantitative antimicrobial susceptibility testing for 382 meropenem-resistant CRPA isolates that

were collected from 78 hospitals across Japan from 2019 to 2020. CRPA exhibited susceptibility

rates of 52.9%, 26.4%, and 88.0% against piperacillin-tazobactam, ciprofloxacin, and amikacin,

respectively, whereas 27.7% of CRPA isolates were classified as difficult-to-treat resistance *P.*

aeruginosa. Of the 148 sequence types detected, ST274 (9.7%) was predominant, followed by

ST235 (7.6%). The proportion of urine isolates in ST235 was higher than that in other STs (P =

0.0056, chi-square test). Only 4.1% of CRPA isolates carried the carbapenemase genes: bla_{GES} (2)

and bla_{IMP} (13). One ST235 isolate carried the novel bla_{IMP} variant bla_{IMP-98} in the chromosome.

Regarding chromosomal mutations, 87.1% of CRPA isolates possessed inactivating or other

resistance mutations in *oprD*, and 28.8% showed mutations in the regulatory genes (*mexR, nalC*,

and *nalD*) for the MexAB-OprM efflux pump. Additionally, 4.7% of CRPA isolates carried a

resistance mutation in the PBP3-encoding gene *ftsI*. The findings from this study and other

surveillance studies collectively demonstrate that CRPA exhibits marked genetic diversity and

that its multidrug resistance in Japan is less prevailed than in other regions. This study

contributes a valuable dataset that addresses a gap in genotype/phenotype information

regarding CRPA in the Asia–Pacific region, where the epidemiological background markedly

differs between regions.

INTRODUCTION

Pseudomonas aeruginosa is a common opportunistic pathogen associated with

pneumonia, urinary tract infections, and bacteremia in clinical settings worldwide. Currently, it

ranks among the top five bacterial pathogens associated with human death globally (1). It also

intrinsically shows reduced susceptibility to many antibiotics due to its possession of four types

of RND family efflux pump, chromosomally encoded AmpC β-lactamase (also called

Pseudomonas-derived class C β-lactamase: PDC), and other antimicrobial-inactivating enzymes

(2, 3). Furthermore, *P. aeruginosa* can acquire resistance mutations at multiple genomic

locations and antimicrobial resistance genes via mobile genetic elements, such as plasmids and

integrative and conjugative elements (ICEs), after which it elicits high levels of resistance to

multiple antimicrobials, thereby limiting treatment options for infected patients (3–5).

Carbapenems represent a crucial class of therapeutic options for treating *P. aeruginosa*

infections. In 2017, the World Health Organization (WHO) listed carbapenem-resistant *P.*

aeruginosa (CRPA) as one of the bacterial priority pathogens classified in "Priority 1: CRITICAL,"

for which new effective antibiotics are urgently required (6). Horizontally transferrable

carbapenemases represent a significant public health concern due to their insensitivity to the

conventionally used β-lactamase inhibitors (7). A recent cohort study demonstrated a 9% higher

30-day mortality rate among patients infected with carbapenemase-producing CRPA when

compared to those infected with carbapenemase-nonproducing CRPA (8). Currently, three

classes of β-lactamases, namely class A, class B (known as metallo-β-lactamases or MBLs), and

class D β-lactamases, are recognized as transferrable carbapenemase. Globally disseminated

carbapenemase types in *P. aeruginosa* include VIM, IMP, NDM, GES, and KPC (9). In addition to

the horizontal acquisition of carbapenemases, *P. aeruginosa* can develop carbapenem

resistance through the mutational inactivation of several chromosomally coded functions. These

include the inactivation of the OprD porin (10) and the inactivation of two direct or one indirect

transcriptional regulators (repressors, MexR, NalC, and NalD) of the MexAB-OprM efflux pump

(11–13). Furthermore, an experimental evolution study revealed that nonsynonymous

substitutions in *ftsI*, an essential gene encoding PBP3, contribute to carbapenem resistance (14).

Recently, international consortia have made efforts to build a large *P. aeruginosa* dataset

to obtain quantitative information for CRPA prevalence, its susceptibility profiles, and the

possession rate of carbapenemases (8, 15–17); Notably, in the Asia–Pacific region, the

Antimicrobial Testing Leadership and Surveillance (ATLAS) program took the initiative of *P.*

aeruginosa surveillance, revealing distinct patterns of resistance rate and predominant β-

lactamase levels in the *P. aeruginosa* population between 12 countries/regions, but the ATLAS

did not cover genome surveillance (16). Hence, information of the major types of resistance

mechanisms and dominant clones of *P. aeruginosa* in the Asia–Pacific region remain lacking.

Japan exhibits characteristic features in the epidemiology of antimicrobial-resistant

bacterial infection in the Asia–Pacific region. Regarding carbapenem resistance, the prevalence

of meropenem-resistant *E. coli* in bloodstream infections in Japan as of 2019 was 2 per 100,000

test patients, which is lower than, for example, 79 per 100,000 test patients observed in

Singapore, which is located in Southeast Asia, according to the WHO Global Antimicrobial

Resistance and Use Surveillance System (18). The established national phenotypic antimicrobial

resistance surveillance program—the Japan Nosocomial Infections Surveillance (JANIS) (19,

20)—reported that the carbapenem (meropenem) resistance rate of *P. aeruginosa* in

hospitalized patients in Japan, as of 2020, was 10.5%. This was average among representative

countries where antimicrobial resistance surveillance for *P. aeruginosa* was once or is currently

being performed; the resistance rate was 29.3% in India (ATLAS 2015–2019), 19.3% in China

(CHINET 2020), 6.5% in Australia (ATLAS 2015–2019), 28.5% in Poland (EARS-net 2020), 15.7% in

the Czech Republic (EARS-net 2020), 13.9% in Germany (EARS-net 2020), 12.6% in France (EARS-

net 2020), and 3.7% in Finland (EARS-net 2020) (16, 21, 22).

To obtain an overview of both genomic features and resistance phenotypes of bacterial

pathogens in Japan, we initiated the national bacterial genome surveillance—Japan

Antimicrobial-Resistant Bacterial Surveillance (JARBS)—in 2019 (23). Through the JARBS-PA

project performed in 2019 and 2020, 382 meropemen-resistant CRPA isolates were collected

from 78 hospitals across Japan. Herein, we report the genotype and susceptibility patterns of

the CRPA isolates in Japan based on genome surveillance and standardized MIC

remeasurements.

RESULTS

Nationwide sampling. From 2019 to 2020, more than 600 *P. aeruginosa* isolates were

transferred from Japanese hospitals to the Antimicrobial Resistance Research Center (AMR-RC)

at the National Institute of Infectious Diseases (NIID) based on the following criterion:

meropenem or imipenem MIC ≥8μg/ml in initial testing at the isolation site. All the collected

isolates were tested for antipseudomonal agent MICs using the Beckman MicroScan system at

AMR-RC. In total, 382 *P. aeruginosa* isolates showing meropenem MIC ≥8μg/ml in the tests at

AMR-RC were defined as CRPA and focused on in this study. The 382 CRPA isolates originated

from nonoverlapping patients in 78 hospitals (the number of collected isolates per island was as

follows: *n* = 30 from Hokkaido Island, *n* = 307 from Honshu Island, *n* = 7 from Shikoku Island, *n* =

32 from Kyusyu Island, and *n* = 6 from Okinawa Island). Source specimens were classified into

one of the following JANIS categories (19, 20): Oral/Endotracheal/Respiratory (*n* = 189),

Urinary/Genital (*n* = 76), Blood/Fluid (*n* = 40), Digestive (*n* = 27), and Other (*n* = 50). Table S1

provides more detailed source categories and metadata for each isolate. Draft genome

sequences of 382 CRPA strains were determined to identify the sequence type (ST), resistance

genes, and mutations.

Antimicrobial susceptibility profiles and STs of CRPA in Japan. Table 1 summarizes the

susceptibilities for antipseudomonal agents. Although the collected CRPA isolates were often

nonsusceptible to antipseudomonal agents other than meropenem, the percentage of

susceptible isolates remained 52.9% for piperacillin-tazobactam, 26.4% for ciprofloxacin, and

88.0% for amikacin based on breakpoints in CLSI M100-ed 32 in 2022 (Table 1). When focused

on isolates from urine (CLSI M100-ed 33 in 2023), the proportion of amikacin-susceptible

isolates was decreased to 75%.

Susceptibility data for individual isolates are provided in Table S1. Among the CRPA

isolates, 27.7% (106/382) were deemed difficult-to-treat resistance *P. aeruginosa* (DTR-*P.*

aeruginosa) in Tamma *et al.*'s definition (24), and 85.8% (328/382) were MDR in Magiorakos *et*

al.'s definition (25). The susceptibility profiles of DTR strains are summarized in Table S2. Based

on the CLSI M100-ed 32 breakpoints, tobramycin (80.2%) exhibited the highest susceptibility

rate of DTR isolates among the three aminoglycosides analyzed. Only three CRPA isolates were

resistant to colistin; two of the three colistin-resistant isolates (Table S1) were however DTR-*P.*

aeruginosa and elicited resistance to at least one agent from each of the seven antimicrobial

agent categories listed in Table 1.

The MLST profiles of 382 CRPA isolates were investigated in the PubMLST database (26).

Including newly assigned ST (ST4091–ST4125), 148 STs were identified (Fig. 1A). The most

frequently encountered clonal group (CG) was CG274 (comprising only ST274: type III secretion

system effector protein genotype $exoS^+/exoU^-$, 9.7%), followed by CG235 (ST235: $exoS^-/exoU^+$,

7.6%), CG253 (ST253: $exoS^-/exoU^+$, 4.5%), and CG155 (ST155 and ST4122: $exoS^+/exoU^-$, 3.4%).

Many minor STs, each contributing about 1%, occupied 49.0% of the total CRPA in Japan. We

further investigated the occurrence of nearly identical isolates among different patients in our

dataset by analyzing Mash distances between draft genomes (27). Among the 382 isolates

examined, 7.6% (29/382) showed a Mash distance of $<10^{-5}$ to at least one other isolate, which we

defined as nearly identical. We extended this analysis to an external dataset used in Reyes *et al*'s

CRPA surveillance across various regions (8). In the USA, the proportion of nearly identical CRPA

isolates was 0.6% (4/527), in China it was 19.2% (33/171), in Singapore it was 4.9% (2/41), and in

Central South America it was 3.5 % (6/171). Consequently, the contribution of outbreaks to the

CRPA dataset in Japan is lower than in China but higher than in the USA.

Furthermore, we identified all the top ten global high-risk clones (ST111, ST175, ST233,

ST235, ST244, ST277, ST298, ST308, ST357, and ST654) (28), which collectively accounted for

17.0% of the total CRPA isolates in Japan (Table S1). Reyes *et al* (8) proposed globally most

common CGs (10 CGs in Fig. 1B). Those globally common CGs comprised 34.8%. Despite being

among the top two most common CGs in the USA and Central South America, CG111 remains a

minor CG in Japan (Fig. 1B). Surprisingly, the dominant clone CG463 (ST463), prevalent in

Chinese clinical settings (7, 30), was entirely absent in the 382 CRPA isolates collected in this

study. The characteristic features of two major CG/STs in Japan, ST274 and ST235 are described

in more detail with genotype information later in the result section.

Resistance mutations and carbapenemases in CRPA. The molecular mechanisms of

meropenem resistance are primarily classified into four types: porin inactivation, efflux pump

overproduction, target alteration, and horizontal acquisition of β-lactamase (2). Those

mechanisms are respectively associated with inactivation of the *oprD* gene, inactivation of

either *mexR*, *nalC*, or *nalD* genes encoding transcriptional regulators (repressors) for the

MexAB-OprM efflux pump (11–13), nonsynonymous substitutions in *ftsI* (14), and acquisition of

plasmid or ICE-borne carbapenemase genes (4, 5). We searched for the presence of inactivating

mutations (frameshift mutations due to insertion/deletion, nonsense mutations, and

inactivation of start codon or stop codon) in *oprD*, *mexR*, *nalC*, and *nalD* as well as resistance

mutations registered in the NCBI AMRFinderPluS database, as these likely contribute to

increasing meropenem MIC.

Database searches revealed that only 4.2% (16/382) of CRPA in Japan possessed

acquired carbapenemases. Known carbapenemases detected were GES-5 (*n* = 2), IMP-1 (*n* = 6),

IMP-10 (*n* = 4), IMP-7 (*n* = 3), and IMP-34 (*n* = 1). No OXA-type carbapenemases were detected.

ST274 CRPA isolates rarely (2.7%, 1/37) carried carbapenemase, contrasting with the 34.5%

(10/29) possession rate of ST235 isolates. One ST235 isolate encoded a novel variant of IMP-

type MBL, which was assigned a new allele name IMP-98 by NCBI.

The prevalence of inactivating mutations or known nonsynonymous resistance

mutations in chromosomal genes in 382 CRPA isolates was 87.1% (333/382) for *oprD*, 14.1%

(54/382) for *mexR*, 9.1% (35/382) for *nalD*, and 6.3% (24/382) for *nalC*. Collectively, 28.8%

(110/382) of CRPA carried an inactivating mutation in at least one of the three repressors for

mexAB. The most common mutation type in *oprD* was IN/DELs causing frameshift (237/382),

followed by nonsense mutations (90/382); these include, for instance, W277stop (24/90),

W339stop (9/90), and W138stop (9/90). In addition, 4.7% (18/382) of CRPA isolates carried

known resistance mutations in *ftsI* (P527S, F533L, R504C, or V471G) (14).

A simultaneous occurrence of resistance mutations and/or carbapenemase was

observed (Fig. 2). While 26.4% (101/382) of CRPA isolates had an inactivating mutation in both

oprD and at least one regulator for *mexAB*, 4.5% (17/382) had both *oprD* and *ftsI* mutations,

with 3.1% (12/382) exhibiting the *oprD* mutation and carbapenemase. The acquisition of

carbapenemase alone occurred only in 0.7% (4/382) of the CRPA. Notably, 9.1% (35/382) did

not carry identifiable inactivating mutations in the four focused genes, known *ftsI* resistance

mutations, or carbapenemase.

bla$_{IMP-98}$. IMP-98 differed from IMP-1 by only one site at the nucleotide level and likely

originated from the *bla$_{IMP-1}$* gene by C637T nonsynonymous substitution, leading to P213S

amino acid change (Fig. 3A). To deduce the origin of the *bla$_{IMP-98}$* region, we determined

complete genome sequence of ST235 strain, JBBCAEG-19-0032, carrying *bla$_{IMP-98}$* using long

reads. This revealed that strain JBBCAEG-19-0032 genome consists of one chromosome, and

bla$_{IMP-98}$ is embedded in a class 1 integron as the upstream region of *bla$_{IMP-98}$* contained specific

motifs of *attI1* (5'-

<u>GTTATGG</u>AGCAGCAACGAT<u>GTTACGC</u>AGCAGGGCAGTCG<u>CCCTAA</u>AACAAAG<u>TTAGGC</u>-3': 7-bp core

sites are underlined) (29).

A comprehensive analysis of the *bla$_{IMP-98}$* context using the ISFinder database (30)

revealed that the class 1 integron is situated within a Tn*402*-related transposon, which is further

nested within a Tn*As3*-related transposon. The archetype Tn*As3* was originally identified in the

genus *Aeromonas* (Fig. 3B) (31, 32). However, the integron structure differs between Tn*As3* and

the Tn*As3*-related transposon in JBBCAEG-19-0032 in the gene cassette array composition and

the absence of conserved *attC/attI* motifs of class 1 integron in Tn*As3* (Fig. 3C).

Both transposon units in JBBCAEG-19-0032 possess intact terminal inverted repeats and

are flanked by 5-bp target site duplication (TSD): CCCTG for the Tn*402*-related transposon and

TTATA for the Tn*As3*-related transposon, indicating their insertions through transposition events.

Further comparison of the JBBCAEG-19-0032 chromosome with the PAO1 chromosome

revealed that the Tn*As3*-related transposon is nested within a genomic island containing distant

homologs of core genes of a recently defined mobile DNA element, the strand-biased

circularizing integrative element (SE) (33–35) (Fig. 3BD). The products of the SE core genes in

JBBCAEG-19-0032 (IntA: encoding tyrosine recombinase, Tfp: tyrosine recombinase fold protein,

IntB: large tyrosine recombinase, Srap: SE-associated recombination auxiliary protein) exhibited

low identities (22.6%–28.5%) in BLASTp comparisons to homologs of a characterized SE, SE-

6945, from the genus *Vibrio* (Fig. 3D) (33). However, IntA and IntB homologs of JBBCAEG-19-

0032 possess the catalytic RHRY motif of the tyrosine recombinase (36)(Supplemental material

Fig. S1), whereas Tfp and Srap homologs share secondary structures with known *Vibrio*

homologs (Fig. S2 and Fig. S3). The *in silico* removal of a 37,833-bp segment from the JBBCAEG-

19-0032 chromosome, with a putative 6-bp footprint normally formed on the right side of SE

after SE integration (33–35), regenerated an intact form of the PA2049-equivalent gene

encoding a hypothetical protein (Fig. 3E). This suggests that the 38-kb region was inserted via

site-specific recombination. Hence, we designated the 37.8-kb insert as putative SE, SE-

PaeJB0032. A BLASTn search against the NCBI-nr/nt database (2024-02-07) and a database

containing 382 Japanese CRPA strains using the left end (*intA* region) of the SE as a query did

not yield significant hits, indicating that SE-PaeJB0032 may have recently integrated into the *P.*

aeruginosa genome.

To test whether IMP-98 is functional, *bla*$_{IMP-98}$ was cloned and introduced into the drug-

susceptible *E. coli* strain DH5α, after which the susceptibility for various β-lactams was

investigated (Table 2). MICs of β-lactams except for aztreonam were considerably higher against

E. coli DH5α carrying bla_{IMP-98} than the control strain, confirming that IMP-98 retains the wide

substrate range of MBL.

Other resistance determinants. Among 46 amikacin-nonsusceptible CRPA isolates, 54.3%

(25/46) possessed known amikacin resistance genes *aac(6')* variants, resistance mutations such

as *fusA1* (Q455R) and *amgS* (V121G) (37), or inactivating mutation in *mexZ* encoding a repressor

for the MexXY efflux pump contributing to aminoglycoside resistance (38) (Table S1). However,

45.7% (21/46) of amikacin-nonsusceptible CRPA did not possess identifiable amikacin resistance

genes or mutations. Among 281 ciprofloxacin nonsusceptible CRPA isolates, 76.9% (216/281)

possessed at least one of the quinolone resistance gene *crpP* (39) (164/281) and resistance

mutations in *gyrA* (T83I, D87N, or D87G; 122/281), *gyrB* (E468D, S466F, or S466Y; 16/281), or

parC (S87L, S87W; 46/281), *parE* (A473V; 6/281) (Table S1). Three colistin-resistant isolates did

not possess any known colistin resistance mutations or genes.

Features of two major STs in Japan. Specifically, we aim to determine whether notable

phenotypic or genotypic features are associated with ST274 (*n* = 37) and ST235 (*n* = 29). Table 3

presents the observation frequencies of four source groups in ST274 and ST235, as well as other

STs. The relative frequencies among the three ST groups differed significantly (P = 2.5 x 10^{-2}, chi-

square test for Table 3, df = 6). The proportion of isolates liked to the

Oral/Endotracheal/Respiratory category was higher in ST274 (64.8%, 24/37) compared to the

rest (48.1%, 166/345), however, this difference was not significant (P = 7.8 x 10^{-2}, chi-square

test, df = 1). A significant difference was observed in the proportion of isolates in the

Urinary/Genital category between ST235 (41.3%, 12/29) and the rest (18.1%, 64/353) (P = 5.6 x

10^{-3}), suggesting that ST235 may have a competitive advantage over other STs in urine.

Table 4 provides a summary of resistance phenotypes, genes, and mutations observed in

the two major STs. Regarding nonsusceptibility, different proportions were observed among the

three ST groups for piperacillin-tazobactam, ceftadizime, ciprofloxacin, and amikacin (Table 3).

Notably, the proportion of amikacin-nonsusceptible isolates differed significantly between

ST235 (62.0%, 18/29) and the rest (7.9%, 28/353) (P < 2.2 x 10^{-16}, chi-square test, df = 1).

The proportion of DTR isolates also varied among STs (P = 4.2 x 10^{-2}, adjusted P-value in

Table 3). Particularly, the proportion of DTR isolates in ST235 (48.0%, 14/29) significantly

differed from the rest (26.1%, 92/352) (P = 1.8 x 10^{-2}). This trend was not significant in ST274

(32.4% in ST274 vs 27.2% in other STs) (P = 0.6).

The possession rates of chromosomal mutations in *oprD*, *mexAB* regulator genes, and

ftsI did not differ among STs (Table 4). However, the possession rates of carbapenemase genes

bla$_{IMP}$ and *bla*$_{GES}$ differed significantly among STs. The possession rates of *bla*$_{IMP}$ and *bla*$_{GES}$ in

ST235 were significantly higher than that in other STs: 34.5% in ST235 vs 1.7% in other STs (P =

1.4 x 10^{-15}). This trend was also observed for the possession rate of amikacin resistance genes:

58.6% in ST235 vs 2.0% in other STs (P < 2.2 x 10^{-16}). These characteristic features of ST235 were

absent in ST274.

Fluoroquinolone resistance determinants can be categorized into chromosomal

resistance mutations (*gyrA*, *gryB*, *parC*, *parD*), and transferrable quinolone resistance gene *crpP*

(39). Although both types appeared at different proportions among STs (Table 4), the

distribution trend differed between the two types. The proportion of isolates carrying *gyr* or *par*

mutations was higher in the two major STs (75.4%, 49/65) than in others (28.5%, 90/316) (P =

5.6 x 10^{-15}), while the proportion of isolates carrying *crpP* was higher in nonST274 and

nonST235 (63.0%, 199/316) than the two STs (34.8%, 23/66) (P = 4.6 x 10^{-5}). Surprisingly, *crpP*

was present in 57% (58/101) of ciprofloxacin-susceptible isolates. Thus, its significance should

be interpreted with caution. These observations validate the notion that ST235 exhibits an

extraordinary resistance phenotype among STs, functioning as an epidemic high-risk clone (28).

DISCUSSION

Since the adoption of a global action plan on antimicrobial resistance at the WHO

conference, regional surveillance on antimicrobial-resistant bacterial pathogens has

commenced worldwide (8, 15–17, 23). However, in the case of CRPA, most surveillance projects

have not provided both MIC values of antimicrobial agents and genome sequences for the

collected strains simultaneously, limiting data interpretation and reuse. Furthermore,

comprehensive surveillance in a single country with a 10% prevalence rate of CRPA, such as

Japan, has been lacking until this study. This study was aimed at addressing these gaps.

The ciprofloxacin and amikacin susceptibility rates regarding CRPA were 26.4% and 88.0%,

respectively, in Japan. These values exceed 20.6% and 60.6% of pooled CRPA values reported in

the Asia–Pacific region (CLSI breakpoint criteria before 2023) (16), and are then equivalent to

and higher than 26.2% and 63.4% of the global CRPA population, respectively (17).

Furthermore, the piperacillin-tazobactam susceptibility rate in Japan was 52.9%, which contrasts

with the 25.3% susceptibility rate in the global CRPA population (17). This difference might

reflect the difference in the carbapenemase possession rate in CRPA between Japan and other

regions. Therefore, in Japan, MDR has not yet heavily prevailed in the *P. aeruginosa* population

in clinical settings compared with other Asia–Pacific regions or the world average. Notably,

fluoroquinolone resistance among CRPA is a global problem and is common even in non-high-

risk clones.

Since several countries/regions normally categorized in the Asia–Pacific region are

geographically isolated by the presence of sea, we expected to observe distinct features of the

CRPA genotype in Japan and other Asia–Pacific regions. The striking feature was the

predominance of CG463/ST463 in China (8) and its absence in Japan (Fig. 1B). Among the

carbapenemase types, IMP was the most frequent (87.5%; 14/16) in Japan. This trend contrasts

with the predominance of NDM in *P. aeruginosa* borne-carbapenemases in India (16), KPC in

China (8), and VIM in Europe, Africa/Middle East, and Latin America (17). As a country with an

average prevalence rate of meropenem-resistant CRPA, Japan was expected to have a low

carbapenemase possession rate among *P. aeruginosa* strains. This study reveals that the

carbapenemase possession rate in Japan is 4.2%, which is among the lowest compared to other

regions where surveillance studies have been conducted. For instance, the carbapenemase

possession rate of meropenem-resistant CRPA was 32.9% in the Asia–Pacific region from 2015

to 2019 (16), 1.9% for the USA in 2018–2019 (8), 69% for South and Central America in 2018–

2019 (8), 32% for China in 2018–2019 (8), 57% in Singapore–Australia in 2018–2019 (8), and

21.9% for Europe in 2017–2019 (17).

The prevalence rate of carbapenemase is influenced by genetic, demographic, and

environmental factors. The genetic factors include the type of STs constituting the *P. aeruginosa*

population. As shown in Table 5, only so-called high-risk clones, represented by ST235, seem to

have a trend toward capturing transferrable carbapenemase. This could be associated with the

number of integrons possessed by the STs, as most reported MBL genes other than NDM are

present as integron gene cassettes (28). In Japan, the proportion of high-risk clones is 17%,

which is lower than that in China (30%, including KPC-producing ST463) and Singapore (60% in

Reyes's surveillance (8)). Therefore, potential carbapenemase gene recipients might be fewer in

Japan than in other regions. The demographic factor includes the low migration rate of

carbapenemase-producing CRPA to hospitals in Japan from high-carbapenemase prevalence

countries/regions. However, this factor is challenging to evaluate based on publicly available

information currently. Environmental factors include the hospital's environmental condition that

can prevent outbreaks and the resulting clonal expansion of carbapenemase-producing CRPA

within the country. The low occurrence of outbreaks is supported by the low frequency (7.6%)

of nearly identical isolates in the 382 CRPA dataset, as opposed to 19.2% in the Chinese CRPA

dataset. Previously, VIM-producing CRPA isolates were found in hospitals in Japan (40–42);

however, they were not detected in the 78 hospitals participating in the JARBS from 2019 to

2020. Furthermore, according to the National Epidemiological Surveillance of Infectious

Diseases by the Ministry of Health, Labor, and Welfare, the number of amikacin-nonsusceptible,

carbapenem, and fluoroquinolone-resistant *P. aeruginosa* has been decreasing over a decade in

Japan (43). Therefore, it is speculated that hospital environments are being managed to reduce

outbreaks and horizontal transmission of carbapenemases in Japan.

The complete sequence determination of a bla_{IMP-98}-carrying strain revealed that the *bla-*

containing-integron was embedded in a Tn*3*-family transposon of *Aeromonas* origin and nested

in a putative SE region in the chromosome, rather than in the plasmid. Previous studies have

identified *Pseudomonas* as one of the top five representative host genera of SEs; however, to

date, only one terminus-delineated SE from *P. aeruginosa*, SE-PaeBT2436, carrying the *tmexCD-*

topJ cluster conferring tetracycline and tigecycline resistance (34). Since homologs of SE core

genes could not be detected by nucleotide-based search in either the current NCBI database or

the CRPA dataset, the SE-PaeJB0032 type might have formed only recently, potentially capturing

bla_{IMP-98}, in other taxa such as *Aeromonas*, and then jumping into the chromosome of unique

ST235 clone.

This surveillance revealed that two STs, ST274 and ST235, members of the top 10

globally most common CG in CRPA, were also the most common in Japan. ST274 and CG274

members have frequently been detected in clinical settings worldwide (44) and are known for

chronic infection and intrabody evolution involving MDR in patients with cystic fibrosis in

Europe (45, 46). Whether the ST274 isolates collected in this study are associated with a chronic

lung infection remains unknown, but the "Oral/Endotracheal/Respiratory" source category was

most frequent in ST274 (Table 4). However, the proportion of this category did not significantly

differ from other STs (P = 0.078). Conversely, the other major ST, ST235, has more striking

features than ST274 in isolation source, nonsusceptibility, and resistance gene possession rate

(Table 3, Table 4). ST235 was proposed as an epidemic high-risk clone due to its worldwide

occurrence with an extremely drug-resistant phenotype (28, 44). However, statistical data

supporting this notion in a single surveillance has been lacking. Our data demonstrate that

ST235 has a trend to acquire resistance genes and resistance phenotypes more frequently than

other STs. The association of ST235 with urine is a novel finding, but the underlying mechanisms

remain to be investigated.

Inactivating mutations in *oprD* were the most frequent meropenem resistance

mechanism (found in 87.1%), as noted in previous genome surveillance performed in other

regions, where 69% of meropenem-resistant CRPA possessed *oprD*-inactivating mutations and

22% possessed acquired carbapenemase (8). We show that in Japan, not only *oprD*-inactivating

mutations but also efflux pump regulator inactivation and *ftsI* resistance mutations are more

common carbapenem resistance mechanisms than the horizontal acquisition of

carbapenemases. Therefore, future surveillance of *P. aeruginosa* should prioritize the study of

chromosomal mutations.

Surprisingly, 9.2% (35/382) of the CRPA isolates did not possess apparent inactivating

mutations, known *ftsI* resistance mutations, or carbapenemase. A recent multicenter study (9)

reported that 14% of modified carbapenemase-inactivation method (mCIM)-positive CRPA,

constituting 4.6% of the total examined CRPA, did not possess known carbapenemases.

Therefore, a fraction of the 9.2% of Japanese CRPA without a known resistance gene or

mutation may harbor enzymes capable of partially degrading meropenem. The potentially

relevant enzymes include variants of β-lactamase such as PDC and OXA (detected variants are

listed in Table S1), as well as Zn^{2+}-dependent imipenemase (47). However, pinpointing the

causal loci of meropenem resistance in the 9.2% of Japanese CRPA is challenging with current

knowledge, as these variants have been poorly characterized to date. Furthermore, known

resistance mutations could not account for the mechanisms of amikacin nonsusceptibility in

45.7% of amikacin-nonsusceptible isolates. These CRPA isolates must possess unreported

resistance genes or mutations in their genomes. A comparative genomics approach is required

to clarify this claim in the future to fully understand and control CRPA spread.

Limitations of this research include the inability to evaluate whether CRPA isolates are

associated with colonization or infection (the cause of disease) in patients when collecting

isolates from hospitals, and the lack of investigation into the effectiveness of new antimicrobial

agents such as cefiderocol, ceftolozane-tazobactam, and ceftazidime-avibactam against CRPA

isolates carrying chromosomal resistance mutations and carbapenemases. Further surveillance

studies will require new study designs to address these limitations.

Conclusions

The first nationwide surveillance for CRPA in Japan demonstrates Japan to be a low-prevalence

country for carbapenemase-producing CRPA. Although the CRPA largely consists of many minor

STs in Japan, the presence of globally common CG/ST and all top 10 high-risk clones was noted.

This underscores the need for continued and comprehensive surveillance. A high-risk clone

ST235 has a distinguished feature among STs to become DTR relatively easily and capture

carbapenemase and aminoglycoside resistance genes. The carbapenem resistance mechanisms

of CRPA cannot be fully explained by known resistance genes and mutations. The dataset built

in this study provides the national epidemiological features and serves as a foundation for

future studies on global epidemiology, therapeutic options, and infection control.

MATERIALS AND METHODS

Strains, media, and plasmids. CRPA candidates participating in our genome surveillance

program (the Japan Antimicrobial-Resistant Bacterial Surveillance [JARBS]) and associated with

isolation source and date information were transferred from hospitals to the NIID AMR center.

As part of this program, hospitals were requested to transfer all isolates showing an imipenem

or meropenem MIC ≥8 in the initial test, limited to one isolate per patient. A total of 668 viable

isolates were received at the AMR-RC between 2019 and 2020. Since the JARBS-PA project

commenced in mid-2019, the dataset primarily consists of isolates identified in 2020.

Subsequently, 273 isolates exhibiting a meropenem MIC < 8 upon remeasurements at AMR-RC

were excluded from the dataset to align with the definition of CRPA used in other surveillance

studies (8, 16, 17). Additionally, three isolates lacked linked isolation source data, and 10

isolates with poor draft genome assembly quality (L90 > 150) were excluded. The final CRPA

dataset comprises 382 isolates with linked isolation source data.

402   Single colony isolation was performed if necessary, using BD BBL™ BTB Lactose-

403   Contained Agar Medium (also called Drigalski agar) (Becton, Dickinson and Company, Franklin

404   Lakes, NJ, USA). BD BBL™ Muller–Hinton II (Cation adjusted) Broth (Becton, Dickinson and

405   Company) was used to culture *P. aeruginosa* strains in liquid. Table S1 contains the *P.*

406   *aeruginosa* strains and associated metadata.

407   *E. coli* strain DH5α [F$^-$, Φ80d*lacZ*ΔM15, Δ(*lacZYA-argF*)U169, *deoR*, *recA*1, *endA*1,

*hsdR*17(r$_K$$^-$, m$_K$$^+$), *phoA*, *supE*44, λ$^-$, *thi*-1, *gyrA*96, *relA*1] was used for DNA cloning and

functional analysis of *bla*$_{IMP-98}$. DH5α was cultured using BD Difco™ LB, Lennox Broth and agar

(Becton, Dickinson and Company). A gene cassette array of *bla*$_{OXA-10}$-*acc(6')-bla*$_{IMP-98}$ in *P.*

aeruginosa ST235 strain JBBCAEG-19-0032 was PCR-amplified using primers blaIMP_F: 5'-

TCGAGCTCGGTACCCCGCTACTTGAAGTGTTGACGC-3' and blaIMP_R: 5'-

CTCTAGAGGATCCCCTTAGTTGCTTGGTTTTGATGGTTTTTTACTTTCG-3', and KODone polymerase

(TOYOBO CO., LTD., Osaka, Japan). A low-copy number vector pMW218 (*nptII*, pSC101 replicon;

NIPPON GENE CO., LTD., Tokyo, Japan) was also PCR-amplified with primers pMW_mcs_F: 5'-

GGGTACCGAGCTCGAATTCGTA-3' and pMW_mcs_R: 5'-GGGGATCCTCTAGAGTCGACCT-3'. The two

PCR products were joined using NEBuilder (New England Biolabs, Ipswich, MA, USA). The

resulting construct was named pHY1399. A segment on pHY1399 was PCR-amplified to delete

the bla_{OXA-10}-$acc(6')$-Ib region using 5' phosphorylated primers P_Integ_del_F: 5'P-

CATGGCACCTTCTTGGTGGCTAACG-3', and P_Integ_del_R: 5'P-GAGCGAACACGCAGTGATGCC-3',

then the purified PCR product was ligated using Ligation High ver.2 (Toyobo) and introduced into

DH5α. The resulting recombinant plasmid obtained from one transformant was named

pHY1401.

Antimicrobial susceptibility test. To obtain MICs of antipseudomonal agents, CRPA strains were

propagated on Drigalski agar and incubated for 1 day at 35°C. Then, a single colony was

inoculated into MicroScan Neg MIC NF-1J panel using Prompt Inoculation System D (Beckman

Coulter Inc., Brea, CA, USA). Inoculated panels were incubated at 35°C for 18 h in MicroScan

Walkway40 Plus (Beckman Coulter Inc.) or MicroScan Walkway DxM 1096 (Beckman Coulter

Inc.). To cover a therapeutically achievable range, the antimicrobial susceptibility testing was

designed to measure major antimicrobials using four dilutions for amikacin and colistin, and five

dilutions for other antimicrobials.

432   To obtain β-lactam MIC against two *E. coli* strains DH5α (pMW218) and DH5α

(pHY1401), the strains were first propagated on LB agar containing kanamycin (25 μg/ml) and

incubated at 37°C overnight, then the colonies were diluted and inoculated into the Neg MIC

EN-2J panel containing β-lactams listed in Table 2 and incubated in MicroScan Walkway40 Plus

(Beckman Coulter Inc.). Breakpoint criteria for susceptible (S), intermediate resistance (I), and

resistance (R) followed the CLSI 2022 guidelines M100-Ed32. The definition of DTR-*P.*

aeruginosa followed Tamma *et al.*'s criteria (24): nonsusceptible to all the eight antimicrobials

belonging to either five antipseudomonal agent categories, and is based on the following MIC

criteria: piperacillin/tazobactam ≥32/4 µg/ml, ceftazidime ≥16 µg/ml, cefepime ≥16 µg/ml,

meropenem ≥4 µg/ml, imipenem ≥4 µg/ml, aztreonam ≥16 µg/ml, ciprofloxacin ≥1 µg/ml, and

levofloxacin ≥2 µg/ml (I or R in CLSI 2022). The definition of MDR was modified from

Magiorakos *et al.*'s criteria (25) as follows: nonsusceptible (CLSI 2022) to at least one agent in at

least three categories except for polymyxin listed in Table 1.

Notably, the equivalence of the Beckman MicroScan panel system to the broth dilution

method per antimicrobial, and received the following 510(k) numbers from The United States

Food and Drug Administration: imipenem, K162740; meropenem, K192355; doripenem,

K101425; levofloxacin, K193358; ciprofloxacin, K193536; ceftazidime, K202343; piperacillin-

tazobactam, K955910; amikacin, K862140; tobramycin, K862140; gentamycin, K862140;

aztreonam, K863776; cefepime, K962150; and colistin, N62-160/S1. Essential agreement

between the MicroScan Walk away method and the reference method for *P. aeruginosa* was

reported to be 91.3%–96.8% (48–51)

Genome sequencing. Draft genome was determined for all 382 CRPA isolates. For this purpose,

P. aeruginosa strains were cultured in Muller–Hinton II broth overnight at 35°C. Cells were

collected in 1.5 mL tubes and resuspended in a lysis solution containing lysozyme and RNaseA,

then incubated at 37°C for 1 h. After adding proteinase K and SDS, the reaction mixture was

incubated at 55°C overnight. Genomic DNA was purified from the cell lysates using Agencourt

AMPure XP beads (Beckman Coulter Inc.) following the protocol recommended by the

manufacturer. NGS libraries were constructed using Enzymatics 5× WGS fragmentation mix and

WGS ligase reagents (Qiagen, Hilden, Germany), and then libraries were sequenced on HiSeq X

Five platform in Macrogen Japan (Tokyo, Japan). Low-quality reads were removed from the

original reads using fastp (52), after which the trimmed reads were assembled using Shovill

pipeline (https://github.com/tseemann/shovill/tree/master) set to default options. Draft

genome assemblies fulfilled L90 < 150.

To determine the complete genome sequence of strain JBBCAEG-19-0032, genomic DNA

was extracted using the Monarch HMW DNA Extraction Kit for Tissues (New England Biolabs,

Ipswitch, MA, USA). Long-read sequencing was performed on the GridION platform (Oxford

Nanopore Technologies, Oxford, UK). The library was prepared using the Rapid Barcoding Kit,

and sequencing was performed using the R9.4.1 flow cell (Oxford Nanopore Technologies).

Basecalling was performed using the high-accuracy base calling model of Guppy v5.0.12 (Oxford

Nanopore Technologies). The complete sequence of JBBCAEG-19-0032 was obtained through

long-read-only assembly using Flye v. 2.9-b1768, set to the "--nano-raw" option (53), followed

by polishing with Illumina reads using Pilon v. 1.22 (54).

Data analysis. The PubMLST ST number was assigned by searches against the PubMLST

database (26). When hit ST was not present in PubMLST, MLST profiles of the isolates were

submitted to PubMLST, after which new ST numbers (ST4091-ST4125) were assigned to the new

MLST profiles. The relationship between MLST profiles was represented as a minimum spanning

tree by GrapeTree software (55). Acquired antimicrobial resistance genes in draft genome

assemblies were searched for using AMRFinderPlus (database ver. 2023-04-17.1) with the "--

organism Pseudomonas_aeruginosa" option, otherwise default setting (56). The

presence/absence and integrity of specific chromosomal genes (frameshift mutations, nonsense

mutations, and gene fusions caused by substitutions at stop/start codon positions) were

determined by tblastn searches against an in-house generated nucleotide database using blast

v2.12.0 (57), and the following retrieval of hit sequences and inspection of the alignments. The

RefSeq/GenBank accession numbers of blast queries used are as follows: ExoU,

WP_034024595.1; ExoS, NP_252530.1; FtsI (PBP3), WP_003094139.1; MexR, NP_249115.1;

NalC, NP_252410.1; NalD, NP_252264.1; OprD, NP_249649.1; MexZ, WP_003088626.1. The

alignments were generated using MAFFT v 7.490 with the einsi option (58). The integron

integrase gene and *attC* were searched for using IntegronFinder 2.0.2 with the default

parameter setting (59). Secondary structure predictions for SE-associated proteins were

conducted using PROMAL3D (60). Gene synteny was visualized using

GenomeMatcher(61).

The genetic distance between two draft genomes was evaluated based on Mash

distance (27) using Mash's "mash dist" function with default parameter setting (k =21, s = 1000).

Importantly, Mash distance serves as an approximation of 1- average nucleotide identity (ANI)

(27). When required, we used Shovill genome assemblies of global CRPA isolates as external

data (BioProject accession: PRJNA824880) (8). The Mash distance matrix of the 382 Japanese

CRPA isolates and subsets from Reyes *et al*'s CRPA isolates were provided in the Zenodo

repository.

Data availability. Raw sequence reads generated in this study are deposited under DDBJ

Sequence Read Archive accession number DRA017464 and DRA017994. Draft genome

assemblies and assembly quality statistics are available from Zenodo repository [DOI:

10.5281/zenodo.10693593] (62). PubMLST allele profiles of 382 isolates and sequence

alignments of *oprD*, *mexR*, *mexZ*, *nalC*, *nalD*, *ftsI*, and Mash distance matrices are available from

Zenodo repository [DOI: 10.5281/zenodo.10693794] (63). The nucleotide sequence of *bla*$_{\text{IMP-98}}$

is deposited under GenBank/DDBJ/EMBL accession number LC740578.1.

Complete sequence of strain JBBCAEG-19-0032 is deposited under GenBank/DDBJ/EMBL

accession number AP029374.

ACKNOWLEDGMENTS

We thank all the hospitals contributing to the JARBS program. We thank Satoyo Wakai, Takahisa

Ishizuka, Liansheng Yu, Mikihisa Okuda, Akira Moriya, Emi Fujimura, Koichi Shimakawa, Rumi

Oki, Yoshie Taki, Mayumi Sasada, Mikako Nakazawa, Noriko Sakamoto, Elahi Shaheem, Chika

Arai, Yuko Kazumi, Akemi Suzuki, and Fumiko Hamamoto at AMR-RC NIID for their support of

isolate collection, susceptibility test, genomic DNA preparation, and JANIS data collection. We

thank Enago [https://www.enago.jp] for language editing. The computation is supported by

supercomputer SHIROKANE of Human Genome Center at the Institute of Medical Science

University of Tokyo, and NIG supercomputer at the Research Organization of Information and

Systems National Institute of Genetics. This research is supported by Japan Agency for Medical

Research and Development under grant number 23fk0108604 (to M.S.). The funder has no role

in study design, data collection, data analysis, and decision to publish results.

522   Author contributions are as follows: Conceptualization, H.Y., W.H., K.Y., and M.S.;

Investigation, H.Y., W.H., Sa.K., S.A., E.A., H.Z., Sh.K., N.K., A.H., T.K., Y.S., and K.Y.; Formal

analysis, H.Y., H.Z., and K.Y.; Data Curation, H.Y., N.K., and K.Y.; Methodology, H.Z., Y.S.;

Visualization, H.Y.; Writing - Original Draft, H.Y.; Writing - Review & Editing, H.Y., W.H., N.K.,

A.H., S.K., H.Z., Y.S., K.Y., and M.S.; Project Administration and Funding Acquisition, M.S.. All

authors reviewed and approved the final manuscript.

ETHICAL STATEMENT

This study (JARBS) was reviewed and approved by the Institutional Review Board of the

National Institute of Infectious Diseases (approval number 1251). The use of JANIS data was

approved by Ministry of Health, Labor and Welfare of Japan (approval number 1553).

REFERENCES

1. **GBD 2019 Antimicrobial Resistance Collaborators**. 2022. Global mortality associated with

33 bacterial pathogens in 2019: a systematic analysis for the Global Burden of Disease

Study 2019. Lancet **400**:2221–2248.

2. **Poole K**. 2011. *Pseudomonas aeruginosa*: resistance to the max. Front Microbiol **2**:65.

3. **Horcajada JP, Montero M, Oliver A, Sorlí L, Luque S, Gómez-Zorrilla S, Benito N, Grau S**.

2019. Epidemiology and treatment of multidrug-resistant and extensively drug-resistant

Pseudomonas aeruginosa infections. Clin Microbiol Rev **32**:e00031–19.

4. **Senda K, Arakawa Y, Nakashima K, Ito H, Ichiyama S, Shimokata K, Kato N, Ohta M**.

1996. Multifocal outbreaks of metallo-β-lactamase-producing *Pseudomonas aeruginosa*

resistant to broad-spectrum beta-lactams, including carbapenems. Antimicrob Agents

Chemother **40**:349–353.

5. **Botelho J, Roberts AP, León-Sampedro R, Grosso F, Peixe L**. 2018. Carbapenemases on

the move: it's good to be on ICEs. Mob DNA **9**:37.

6. **World-Health-Organization**. 2017. WHO priority pathogens list for R&D of new

antibiotics. URL: https://www.who.int/news/item/27-02-2017-who-publishes-list-of-

bacteria-for-which-new-antibiotics-are-urgently-needed. Retrieved November 8, 2023

7. **Queenan AM, Bush K**. 2007. Carbapenemases: the versatile β-lactamases. Clin Microbiol

Rev **20**:440–458, table of contents.

8. **Reyes J, Komarow L, Chen L, Ge L, Hanson BM, Cober E, Herc E, Alenazi T, Kaye KS,**

Garcia-Diaz J, Li L, Kanj SS, Liu Z, Oñate JM, Salata RA, Marimuthu K, Gao H, Zong Z,

Valderrama-Beltrán SL, Yu Y, Tambyah P, Weston G, Salcedo S, Abbo LM, Xie Q, Ordoñez

K, Wang M, Stryjewski ME, Munita JM, Paterson DL, Evans S, Hill C, Baum K, Bonomo

RA, Kreiswirth BN, Villegas MV, Patel R, Arias CA, Chambers HF, Fowler VG, Doi Y, van

Duin D, Satlin MJ, Antibacterial-Resistance-Leadership-Group-and-Multi-Drug-Resistant-

Organism-Network-Investigators. 2023. Global epidemiology and clinical outcomes of

carbapenem-resistant *Pseudomonas aeruginosa* and associated carbapenemases (POP): a

prospective cohort study. Lancet Microbe **4**:e159–e170.

9. **Gill CM, Aktaþ E, Alfouzan W, Bourassa L, Brink A, Burnham CD, Canton R, Carmeli Y,**

Falcone M, Kiffer C, Marchese A, Martinez O, Pournaras S, Satlin M, Seifert H, Thabit AK,

Thomson KS, Villegas MV, Nicolau DP, ERACE-PA GSG. 2021. The ERACE-PA global

surveillance program: ceftolozane/tazobactam and ceftazidime/avibactam in vitro activity

against a global collection of carbapenem-resistant *Pseudomonas aeruginosa*. Eur J Clin

Microbiol Infect Dis **40**:2533–2541.

10. **Sanbongi Y, Shimizu A, Suzuki T, Nagaso H, Ida T, Maebashi K, Gotoh N**. 2009.

Classification of OprD sequence and correlation with antimicrobial activity of carbapenem

agents in *Pseudomonas aeruginosa* clinical isolates collected in Japan. Microbiol Immunol

53:361–367.

11.   **Srikumar R, Paul CJ, Poole K**. 2000. Influence of mutations in the *mexR* repressor gene on

expression of the MexA-MexB-oprM multidrug efflux system of *Pseudomonas aeruginosa*.

J Bacteriol **182**:1410–1414.

12.   **Cao L, Srikumar R, Poole K**. 2004. MexAB-OprM hyperexpression in NalC-type multidrug-

resistant *Pseudomonas aeruginosa*: identification and characterization of the *nalC* gene

encoding a repressor of PA3720-PA3719. Mol Microbiol **53**:1423–1436.

13.   **Morita Y, Cao L, Gould VC, Avison MB, Poole K**. 2006. *nalD* encodes a second repressor of

the *mexAB-oprM* multidrug efflux operon of *Pseudomonas aeruginosa*. J Bacteriol

188:8649–8654.

14.   **Cabot G, Florit-Mendoza L, Sánchez-Diener I, Zamorano L, Oliver A**. 2018. Deciphering β-

lactamase-independent β-lactam resistance evolution trajectories in *Pseudomonas*

aeruginosa. J Antimicrob Chemother **73**:3322–3331.

15.   **Karlowsky JA, Bouchillon SK, El Mahdy Kotb R, Mohamed N, Stone GG, Sahm DF**. 2022.

Carbapenem-resistant Enterobacterales and *Pseudomonas aeruginosa* causing infection in

Africa and the Middle East: a surveillance study from the ATLAS programme (2018-20).

JAC Antimicrob Resist **4**:dlac060.

16.   **Lee YL, Ko WC, Hsueh PR**. 2022. Geographic patterns of carbapenem-resistant

Pseudomonas aeruginosa in the Asia-Pacific region: results from the antimicrobial testing

leadership and surveillance (ATLAS) program, 2015-2019. Antimicrob Agents Chemother

66:e0200021.

17. **Gales AC, Stone G, Sahm DF, Wise MG, Utt E**. 2023. Incidence of ESBLs and

carbapenemases among Enterobacterales and carbapenemases in *Pseudomonas*

aeruginosa isolates collected globally: results from ATLAS 2017-2019. J Antimicrob

Chemother dkad127.

18. **World Health Organization**. 2022. Global antimicrobial resistance and use surveillance

system (GLASS) Report: 2021 report electronic supplementary material. URL:

https://www.who.int/publications/i/item/9789240027336. Retrieved November 8, 2023

19. **Kajihara T, Yahara K, Hirabayashi A, Shibayama K, Sugai M**. 2021. Japan nosocomial

infections surveillance (JANIS): current status, international collaboration, and future

directions for a comprehensive antimicrobial resistance surveillance system. Jpn J Infect

Dis **74**:87–96.

20. **Ministry of Health Labour and Welfare of Japan**. 2021. JANIS Open Report Clinical

Laboratory-Division. URL: https://janis.mhlw.go.jp/english/report/index.html. Retrieved

November 8, 2023

21. **Yang W, Ding L, Han R, Yin D, Wu S, Yang Y, Zhu D, Guo Y, Hu F, Xu Y, Zhang X, Zhang Z, Ji**

P, Kang M, He C, Wang C, Fu P, Xu Y, Huang Y, Chen Z, Sun Z, Ni Y, Sun J, Chu Y, Tian S,

Hu Z, Li J, Yu Y, Lin J, Shan B, Du Y, Guo S, Wang Y, Wei L, Wang X, Zhang H, Wang C, Hu

Y, Ai X, Zhuo C, Su D, Wang R, Fang H, Yu B, Gong P, Song M, Guo D, Zhao J, Liu W, Li Y,

Jin Y, Wang Y, Weng K, Zhang Y, Xu X, Yan C, Huang X, Yu H, Li Y, Wang S, Zhang L, Ma J,

Zhou S, Ke J, Zhu L, Meng J, Shen H, Zhou W, Li G, Jia W, Wu J, Lu Y, Li J, Liu J, Liao L, Gu

H, Jiang L, He W, Xue S, Feng J, Dou R, Yue C, Guo R, Jin Y, Ma X, Zheng Y, Hu F, Chen Y,

Meng Q, Xia Y, Sun S. 2023. Current status and trends of antimicrobial resistance among

clinical isolates in China: a retrospective study of CHINET from 2018 to 2022. One Health

Adv **1**:8.

22. **European Centre for Disease Prevention and Control**. 2022. Antimicrobial resistance in

the EU/EEA (EARS-Net) - Annual epidemiological report for 2020.

https://www.ecdc.europa.eu/sites/default/files/documents/EUEEA_trend_tables_for_indi

vidual_isolates_2020.pdf. Retrieved November 8, 2023.

23. **Kayama S, Yahara K, Sugawara Y, Kawakami S, Kondo K, Zuo H, Kutsuno S, Kitamura N,**

Hirabayashi A, Kajihara T, Kurosu H, Yu L, Suzuki M, Hisatsune J, Sugai M. 2023. National

genomic surveillance integrating standardized quantitative susceptibility testing clarifies

antimicrobial resistance in Enterobacterales. Nat Comm **14**:8046.

24. **Tamma PD, Aitken SL, Bonomo RA, Mathers AJ, van Duin D, Clancy CJ**. 2021. Infectious

Diseases Society of America guidance on the treatment of extended-spectrum β-

lactamase producing enterobacterales (ESBL-E), carbapenem-resistant enterobacterales

(CRE), and *Pseudomonas aeruginosa* with Difficult-to-treat resistance (DTR-*P. aeruginosa*).

Clin Infect Dis **72**:1109–1116.

25. **Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S,**

Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens

MJ, Vatopoulos A, Weber JT, Monnet DL. 2012. Multidrug-resistant, extensively drug-

resistant and pandrug-resistant bacteria: an international expert proposal for interim

standard definitions for acquired resistance. Clin Microbiol Infect **18**:268–281.

26.   **Jolley KA, Bray JE, Maiden MCJ**. 2018. Open-access bacterial population genomics:

BIGSdb software, the PubMLST.org website and their applications. Wellcome Open Res

3:124.

27.   **Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM**.

2016. Mash: fast genome and metagenome distance estimation using MinHash. Genome

Biol **17**:132.

28.   **Del Barrio-Tofiño E, López-Causapé C, Oliver A**. 2020. *Pseudomonas aeruginosa* epidemic

high-risk clones and their association with horizontally-acquired β-lactamases: 2020

update. Int J Antimicrob Agents **56**:106196.

29.   **Partridge SR, Recchia GD, Scaramuzzi C, Collis CM, Stokes HW, Hall RM**. 2000. Definition

of the *attI1* site of class 1 integrons. Microbiology (Reading) **146**:2855–2864.

30.   **Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M**. 2006. ISfinder: the reference

centre for bacterial insertion sequences. Nucleic Acids Res **34**:D32–6.

31.   **Minakhina S, Kholodii G, Mindlin S, Yurieva O, Nikiforov V**. 1999. Tn5053 family

transposons are res site hunters sensing plasmidal res sites occupied by cognate

resolvases. Mol Microbiol **33**:1059–1068.

32.   **Pfeiffer F**. 2015. Tn*As3.* URL: https://isfinder.biotoul.fr/scripts/ficheIS.php?name=TnAs3.

Retrieved February 1, 2024.

33.   **Nonaka L, Masuda M, Yano H**. 2022. Atypical integrative element with strand-biased

circularization activity assists interspecies antimicrobial resistance gene transfer from

Vibrio alfacsensis. PLOS ONE **17**:e0271627.

34.   **Idola D, Mori H, Nagata Y, Nonaka L, Yano H**. 2023. Host range of strand-biased

circularizing integrative elements: a new class of mobile DNA elements nesting in

Gammaproteobacteria. Mob DNA **14**:7.

35.   **Yano H, Suzuki M, Nonaka L**. 2024. Mobile class A β-lactamase gene bla$_{GMA-1}$. Microbiol

Spectr **12**:e0258923.

36.   **Nunes-Düby SE, Kwon HJ, Tirumalai RS, Ellenberger T, Landy A**. 1998. Similarities and

differences among 105 members of the Int family of site-specific recombinases. Nucleic

Acids Res **26**:391–406.

37.   **Lau CH, Fraud S, Jones M, Peterson SN, Poole K**. 2013. Mutational activation of the

AmgRS two-component system in aminoglycoside-resistant *Pseudomonas aeruginosa*.

Antimicrob Agents Chemother **57**:2243–2251.

38.   **Morita Y, Tomida J, Kawamura Y**. 2012. Primary mechanisms mediating aminoglycoside

resistance in the multidrug-resistant *Pseudomonas aeruginosa* clinical isolate PA7.

Microbiology (Reading) **158**:1071–1083.

39.   **Chávez-Jacobo VM, Hernández-Ramírez KC, Romo-Rodríguez P, Pérez-Gallardo RV,**

Campos-García J, Gutiérrez-Corona JF, García-Merinos JP, Meza-Carmen V, Silva-Sánchez

J, Ramírez-Díaz MI. 2018. CrpP is a novel ciprofloxacin-modifying enzyme encoded by the

Pseudomonas aeruginosa pUM505 plasmid. Antimicrob Agents Chemother **62**:e02629–17.

40.   **Kouda S, Ohara M, Onodera M, Fujiue Y, Sasaki M, Kohara T, Kashiyama S, Hayashida S,**

Harino T, Tsuji T, Itaha H, Gotoh N, Matsubara A, Usui T, Sugai M. 2009. Increased

prevalence and clonal dissemination of multidrug-resistant *Pseudomonas aeruginosa* with

the *bla*$_{IMP-1}$ gene cassette in Hiroshima. J Antimicrob Chemother **64**:46–51.

41. **Shimizu W, Kayama S, Kouda S, Ogura Y, Kobayashi K, Shigemoto N, Shimada N, Yano R,**

Hisatsune J, Kato F, Hayashi T, Sueda T, Ohge H, Sugai M. 2015. Persistence and epidemic

propagation of a *Pseudomonas aeruginosa* sequence type 235 clone harboring an IS*26*

composite transposon carrying the bla_{IMP-1} integron in Hiroshima, Japan, 2005 to 2012.

Antimicrob Agents Chemother **59**:2678–2687.

42. **Hishinuma T, Uchida H, Tohya M, Shimojima M, Tada T, Kirikae T**. 2020. Emergence and

spread of VIM-type metallo-β-lactamase-producing *Pseudomonas aeruginosa* clinical

isolates in Japan. J Glob Antimicrob Resist **23**:265–268.

43. **Suzuki S**. 2021. A view on 20 years of antimicrobial resistance in Japan by two national

surveillance systems: the national epidemiological surveillance of infectious diseases and

Japan nosocomial infections surveillance. Antibiotics (Basel) **10**:1189.

44. **Oliver A, Mulet X, López-Causapé C, Juan C**. 2015. The increasing threat of *Pseudomonas*

aeruginosa high-risk clones. Drug Resist Updat **21-22**:41–59.

45. **Fernández-Olmos A, García-Castillo M, Alba JM, Morosini MI, Lamas A, Romero B, Galán**

JC, del Campo R, Cantón R. 2013. Population structure and antimicrobial susceptibility of

both nonpersistent and persistent *Pseudomonas aeruginosa* isolates recovered from

cystic fibrosis patients. J Clin Microbiol **51**:2761–2765.

46. **López-Causapé C, Sommer LM, Cabot G, Rubio R, Ocampo-Sosa AA, Johansen HK,**

Figuerola J, Cantón R, Kidd TJ, Molin S, Oliver A. 2017. Evolution of the *Pseudomonas*

aeruginosa mutational resistome in an international cystic fibrosis clone. Sci Rep **7**:5555.

47.   **Fajardo A, Hernando-Amado S, Oliver A, Ball G, Filloux A, Martinez JL**. 2014.

Characterization of a novel Zn^{2+}-dependent intrinsic imipenemase from Pseudomonas

aeruginosa. J Antimicrob Chemother **69**:2972–2978.

48.   **Food and Drug Administration**. 2017. 510(k) Substantial equivalence determination

decision summary: K162740. URL:

https://www.accessdata.fda.gov/cdrh_docs/reviews/K162740.pdf. Retrieved Feb 1, 2024.

49.   **Food and Drug Administration**. 2019. 510(k) Substantial equivalence determinatinon

decision summary: K192355. URL:

https://www.accessdata.fda.gov/cdrh_docs/reviews/K192355.pdf. Retrieved February 1,

2024.

50.   **Food and Drug Administration**. 2020. 510(k) Substantial equivalence determination

decision summary: K202343. URL:

https://www.accessdata.fda.gov/cdrh_docs/reviews/K202343.pdf. Retrieved February 1,

2024.

51.   **Food and Drug Administration**. 2020. 510(k) Substantial equivalence determination

decision summary: K193536. URL:

https://www.accessdata.fda.gov/cdrh_docs/reviews/K193536.pdf. Retrieved February 1,

2024.

52.   **Chen S, Zhou Y, Chen Y, Gu J**. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**:i884–i890.

53.   **Kolmogorov M, Yuan J, Lin Y, Pevzner PA**. 2019. Assembly of long, error-prone reads

using repeat graphs. Nat Biotechnol **37**:540–546.

54. **Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,**

Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive

microbial variant detection and genome assembly improvement. PLOS ONE **9**:e112963.

55. **Zhou Z, Alikhan NF, Sergeant MJ, Luhmann N, Vaz C, Francisco AP, Carriço JA, Achtman**

M. 2018. GrapeTree: visualization of core genomic relationships among 100,000 bacterial

pathogens. Genome Res **28**:1395–1404.

56. **Feldgarden M, Brover V, Gonzalez-Escalona N, Frye JG, Haendiges J, Haft DH, Hoffmann**

M, Pettengill JB, Prasad AB, Tillman GE, Tyson GH, Klimke W. 2021. AMRFinderPlus and

the reference gene catalog facilitate examination of the genomic links among

antimicrobial resistance, stress response, and virulence. Sci Rep **11**:12728.

57. **Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ**. 1990. Basic local alignment search

tool. J Mol Biol **215**:403–410.

58. **Katoh K, Standley DM**. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol **30**:772–780.

59. **Néron B, Littner E, Haudiquet M, Perrin A, Cury J, Rocha EPC**. 2022. IntegronFinder 2.0:

identification and analysis of integrons across bacteria, with a focus on antibiotic

resistance in *Klebsiella*. Microorganisms **10**:700.

60. **Pei J, Kim BH, Grishin NV**. 2008. PROMALS3D: a tool for multiple protein sequence and

structure alignments. Nucleic Acids Res **36**:2295–2300.

61. **Ohtsubo Y, Ikeda-Ohtsubo W, Nagata Y, Tsuda M**. 2008. GenomeMatcher: a graphical

user interface for DNA sequence comparison. BMC Bioinformatics **9**:376.

62.   **Antimicrobial Resistance Research Center National Institute of Infectious Diseases**. 2024.

Genome assemblies of 382 carbapenem-resistant *Pseudomonas aeruginosa* isolates

collected from Japanese hospitals in 2019–2020. Zenodo

https://doi.org/10.5281/zenodo.10693593. Retrieved February 22, 2024.

63.   **Yano H**. 2024. PubMLST allele profiles and sequence alignments of 382 carbapenem-

resistant *Pseudomonas aeruginosa* isolates collected from Japanese hospitals in 2019–

2020. Zenodo https://doi.org/10.5281/zenodo.10693794. Retrieved February 22, 2024.

SUPPLEMENTAL MATERIAL

Table S1. Susceptibility profile, genotype, and isolation source of 382 CRPA isolates.

Table S2. Summary of susceptibility profile of DTR isolates.

FIG S1. Secondary structure alignment of IntA (JBP_33480 product) and IntB (JBP_33500

product) of JBBCAEG-19-0032 with known tyrosine recombinases. R-H-R-Y motif was highlighted

by yellow background. The alignment was generated using PROMAL3D. e: sheet; h, helix.

FIG S2. Secondary structure alignment of JBP_33490 product of JBBCAEG-19-0032 with known

Tfp homologs. The alignment was generated using PROMAL3D. e: sheet; h, helix.

FIG S3. Secondary structure alignment of JBP_33510 of JBBCAEG-19-0032 with known Srap

homologs. The alignment was generated using PROMAL3D. e: sheet; h, helix.

FIGURE LEGENDS

FIG 1 (A) Minimum spanning tree of 382 meropenem-resistant CRPA isolates based on MLST

profiles. Each node represents sequence type (ST), otherwise clonal group (CG). STs with one

allele difference were collapsed into one node as CG. Node size is proportional to the number of

isolates. The number on the branch denotes the number of different alleles at the 7 MLST loci

between two nodes. (B) Proportions of 10 globally most common CGs in Japan and other

regions. Proportion of CGs in the USA, China, and South and Central America are based on

Reyes *et al.*'s report (8).

FIG 2. Venn diagram showing the simultaneous occurrence of resistance mutations in

meropenem-resistant *P. aeruginosa*. Numbers in circles represent the numbers of isolates

having an acquired carbapenemase, a point mutation registered in AMRFinderPlus (56), or

inactivating mutations. The "Mex" category indicates the presence of a mutation in at least one

of three loci (*mexR*, *nalC*, and *nalD*) encoding a negative regulator for the *mexAB* operon. The

"PBP3" category indicates the presence of a mutation in *ftsI*. The acquired

carbapenemase/MBLs detected are GES-5, IMP-1, IMP-10, IMP-7, IMP-34, and IMP-98.

A

```
gb_CAA11471.1_IMP-1       179  KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVPSHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN  246
WP_268871851.1_IMP-98     179  KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVSSHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN  246
gb_BAB15941.1_IMP-6       179  KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVPGHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN  246
gb_BAB72069.1_IMP-10      179  KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVPSHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN  246
gb_BAP75826.1_IMP-34      179  KPYGLGNLGDANIEAWPKSAKLLKSKYGKAKLVVPSHSEVGDASLLKLTLEQAVKGLNESKKPSKPSN  246
gb_AAK12087.1_IMP-7       179  KPYGLGNLGDANLEAWPKSAKLLVSKYGKAKLVVPSHSEVGDASLLKRTLEQAVKGLNESKKLSKPSN  246
gb_CAB94707.1_IMP-2       179  KPDGLGNLGDANLEAWPKSAKILMSKYVKAKLVVSSHSEIGDASLLKRTWEQAVKGLNESKKPSQPSN  246
gb_ALT53802.1_IMP-56      179  KPDGLGNLGDANLEAWPKSAKILMSKYGKAKLVVSGHSEIGNASLLQRTWEQAVKGLNESKKPLQPSS  246
Consensus_aa:                  KP.GLGNLGDANlEAWPKSAKIlLbSKYGKAKLVVstHSElGsASLLp.ThEQAVKGLNESKKP.pPSs
Consensus_ss:                      hhhhhhhhhhhhh      eeeee      hhhhhhhhhhhhhhhhhhhhhhh
```

777
778
FIG 3 (A) Alignment of carboxyl-terminal regions of IMP variants. H, helix; e, sheet. IMP-98

differs from IMP-1 only at P213 indicated by an arrowhead. (B) Genetic context of *bla*IMP-98. *tni*

genes in green are associated with the transposition of the Tn*402*-related element. *tniA**

indicates disrupted *tniA*. *tnpA* and *tnpR* genes are associated with the transposition of the

Tn*As3*-related transposon. The left four genes in the insertion in JBBCAEG-19-0032 are SE core

genes: *intA*, encoding tyrosine recombinase; *tfp*, tyrosine recombinase fold protein; *intB*, large

tyrosine recombinase; *srap*, SE-associated recombination auxiliary protein (34). bla_{OXA-10} and

aac(6')-Ib in integon overlapped by 23 bp. *intI* and *attC* were identified by IntegronFinder2.0

(59). (C) Comparison between Tn*As3* and Tn*As3*-realted elements in the JBBCAEG-19-0032

chromosome. Integron-associated genes are shown in light green. *attI* and *attC* sites were not

found in Tn*As3*. *mer* genes in purple are mercury resistance genes.

(D) Similarity of SE core genes between SE-PaeJB0032 and SE-6945 (33). The number indicates

percentage identity in BLASTp. (E) Comparison of border regions between SE-PaeJB0032 and

chromosome (*attL*, *attR*) with unoccupied integration site (*attB*) in PAO1. Sequences in orange

are putative SE regions. The underlined sequence is a putative 6-bp footprint generated by SE

integration (33, 34). Sequences are obtained from the following accession numbers: JBBCAEG-

19-0032 chromosome, AP029374; PAO1 chromosome, NC_002516.2; pSEA1, AP024167; Tn*As3*,

CP00654.1.

TABLE 1. Antimicrobial susceptibilities of 382 meropenem-resistant CRPA isolates.

Antimicrobial agents	Percentage of susceptibility group (number of isolates)[a]		
	S	I	R
Antipseudomonal carbapenems			
Imipenem	1.0 (4)	2.9 (11)	96.1 (367)
Meropenem	0.0 (0)	0.0 (0)	100 (382)
Doripenem	10.7 (41)	36.6 (136)	53.7 (205)
Antipseudomonal cephalosporins			
Ceftazidime	53.1 (203)	12.8 (49)	34.0 (130)
Cefepime	47.4 (181)	27.2 (104)	25.4 (97)
Antipseudomonal penicillin + inhibitor			
Piperacillin-tazobactam (CLSI 2022)	52.9 (202)	24.3 (93)	22.8 (87)
Piperacillin-tazobactam (CLSI 2023)	52.9 (202)	11.0 (42)	35.3 (138)
Monobactam			
Aztreonam	18.3 (70)	29.8 (114)	51.8 (198)
Antipseudomonal fluoroquinolones			
Levofloxacin	27.5 (105)	20.7 (79)	51.8 (198)
Ciprofloxacin	26.4 (101)	35.6 (136)	38.0 (145)
Aminoglycosides			
Gentamycin (CLSI 2022)	63.4 (242)	22.3 (85)	14.4 (55)
Tobramycin (CLSI 2022)	88.7 (339)	2.1 (8)	9.2 (35)
Tobramycin (CLSI 2023)	51.6 (197)	31.7 (121)	16.8 (64)
Amikacin (CLSI 2022)	88.0 (336)	4.2 (16)	7.1 (27)
Amikacin (CLSI 2023)[b]	75.0 (54)	12.5 (9)	12.5 (9)
Polymyxin			
Colistin	-	99.2 (379)	0.8 (3)

[a] Unless mentioned, criteria for S/I/R are identical between CLSI M100-Ed32 (2022) and M100-

Ed33 (2023).

[b] Amikacin MIC against urine-associated isolates only (n = 72, linked to the source categories:

urine, urine collected by catheter, urine obtained from indwelling catheter, catheterized urine,

and urethral discharge in Table S1).

TABLE 2. β-lactam MICs against *E. coli* strain carrying *bla*IMP-98.

Antimicrobials	DH5α(pMW218)	DH5α(pHY1401)[a]
Ampicillin	≤4	>16
Ampicillin-sulbactam	≤4/2	>16/8
Amoxicillin-clavulanate	≤8/4	>16/8
Piperacillin	≤4	32
Piperacilin-tazobactam	≤4	8
Cefazolin	≤1	>16
Cefotiam	≤8	16
Cefaclor	≤8	>16
Cefmetazole	≤4	>32
Cefotaxime	≤0.5	>2
Ceftriaxone	≤0.5	>2
Ceftazidime	≤1	>8
Cefpodoxime	≤1	>4
Cefditoren	≤1	>2
Cefoperazone-sulbactam	≤8/4	>32/16
Cefepime	≤1	8
Imipenem	≤0.5	2
Meropenem	≤0.25	2
Ertapenem	≤0.25	>1
Flomoxef	≤8	32
Aztreonam	≤1	≤1

808 [a] pHY1401 is a pMW218 derivative carrying *bla*IMP-98 and the putative promoter region of class 1

809 integron in strain JBBCAEG-19-0032.

810
811

812 **Table 3.** Isolation sources observed in two major STs.

Source category group	ST274	ST235	Other STs
Oral/Endotracheal/Respiratory	23	9	157
Urinary/Genital	4	12	60
Blood/Fluids	1	3	36
Other sources	9	5	63
Total	37	29	316

813 [a] More specific source categories are shown in Table S1.

816   **Table 4.** Observation frequency of specific resistance phenotype, genes, or mutations in two

817   major ST groups.

Phenotype or genotype	Percentage of relevant isolates in ST (number of isolates)			Adjusted P-value[a]
	ST274 (total 37)	ST235 (total 29)	Other STs (total 316)	
Non-susceptibilities[b]				
Piperacillin-tazobactam	70.2 (26)	58.6 (17)	45.9 (145)	0.02309
Ceftazidime	32.4 (12)	65.5 (19)	44.3 (140)	0.04234
Ciprofloxacin	70.2 (26)	93.1 (27)	72.1 (228)	0.05972
Amikacin	13.5 (5)	62.0 (18)	7.2 (23)	0.00035
DTR	32.4 (12)	48.2 (14)	25.3 (80)	0.04235
Resistance genes/mutations				
Carbapenem				
$bla_{IMP,}$ or bla_{GES}	2.7 (1)	34.5 (10)	1.6 (5)	0.00035
$oprD$[c]	89.1 (33)	96.6 (28)	86.1 (272)	0.35682
$mexR$, $nalC$, or $nalD$[c]	27.0 (10)	27.6 (8)	28.4 (90)	1.0
$ftsI$[c]	5.4 (2)	10.3 (3)	4.1 (13)	0.37338
Fluoroquinolone				
$gyrA$, $gryB$, $parD$, or $parC$[c]	59.5 (22)	93.1 (27)	28.5 (90)	0.00035
$crpP$	32.4 (12)	37.9 (11)	63.0 (199)	0.00084
Amikacin				
$acc(6')$-1b, $aac(6')$-II, $aac(6')$-31, $aacA8$, $aac(6')$-Ia, $aac(6')$-Ib3, $aph(3')$-XV, or $aac(6')$-Iae	0 (0)	58.6 (17)	2.2 (7)	0.00035

[a] Chi-square test P-value after multiple testing corrections (false discovery rate) by Benjamini &

Hochberg method. The independence of two factors (phenotype/genotype and ST) was

evaluated by using a 2 by 3 contingency table (df = 2) where the rows are the count of qualified

isolates and the count of nonqualified isolates. DTR: difficult-to-treat resistance.

[b] Breakpoint criteria follows CLSI M100-ed 32 (2022).

[c] Acquired mutation in conserved genes.
824
825
826