Actin dynamics switches two distinct modes of endosomal fusion in yolk sac

visceral endoderm cells

Seiichi Koike[1,2,3], Masashi Tachikawa[4], Motosuke Tsutsumi[5,6], Takuya Okada[1,2],

Tomomi Nemoto[5,6], Kazuko Keino-Masu[1,2], Masayuki Masu[1,2]

[1]Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1

Tennodai, Tsukuba, Ibaraki 305-8575, Japan

[2]Department of Molecular Neurobiology, Institute of Medicine, University of Tsukuba,

1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan

[3]Laboratory of Molecular and Cellular Biology, Graduate School of Science and

Engineering for Research, University of Toyama, 3190 Gofuku, Toyama 930-8555,

Toyama, Japan

[4]Graduate School of Nanobioscience, Yokohama City University, 22-2 Seto, Kanazawa-

ku, Yokohama, Kanagawa 236-0027, Japan

[5]Exploratory Research Center on Life and Living Systems (ExCELLS), National

Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787,

Japan

[6]National Institute for Physiological Sciences, National Institutes of Natural Sciences, 5-

1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan

Running title: Two distinct modes of endosome fusion

Keywords: late endosome, lysosome, fusion, actin, cofilin, yolk sac visceral endoderm,

mouse embryo

Correspondence should be addressed to mmasu@md.tsukuba.ac.jp and skoike@eng.u-

toyama.ac.jp

Abstract

Membranes undergo various patterns of deformation during vesicle fusion, but how this

membrane deformation is regulated and contributes to fusion remains unknown. In this

study, we developed a new method of observing the fusion of individual late endosomes

and lysosomes by using yolk sac visceral endoderm cells that have huge endocytic

vesicles. We found that there were two distinct fusion modes that were differently

regulated. In homotypic fusion, two late endosomes fused quickly, whereas in

heterotypic fusion they fused to lysosomes slowly. Mathematical modeling showed that

vesicle size is a critical determinant of these fusion types and that membrane fluctuation

forces can overcome the vesicle size effects. We found that actin filaments were bound

to late endosomes and forces derived from dynamic actin remodeling were necessary for

quick fusion during homotypic fusion. Furthermore, cofilin played a role in endocytic

fusion by regulating actin turnover. These data suggest that actin promotes vesicle

fusion for efficient membrane trafficking in visceral endoderm cells.

Introduction

Eukaryotic cells contain numerous organelles that are separated from the cytoplasm by a

lipid bilayer. Exchange of proteins, lipids, and metabolites between the organelles

occurs through vesicle budding and fusion. To date, theoretical and computer

biophysics have modeled several steps in the process of membrane fusion

(Chernomordik and Kozlov, 2008; Lu and Guo, 2019). The priming of membranes is

followed by the merger of the outer leaflet of the membranes to establish a hemifusion

stalk. The stalk then expands into a hemifusion diaphragm and the merger of the inner

leaflets leads to the formation of a fusion pore, which expands and allows content

mixing. Recently, the hemifusion stage was observed in live cells, providing strong

evidence of the hemifusion model (Zhao et al., 2016).

Studies utilizing time-lapse microscopic observation of live cells, however, have

revealed that membrane behavior after pore formation is unexpectedly divergent. For

example, in giant endosomal vesicles in cells overexpressing constitutively active Rab5

protein, two different types of membrane fusion were observed (Roberts et al., 1999). In

"explosive" fusion, which is commonly seen in cells, the fusion pore expanded quickly.

In contrast, in "bridge" fusion, the pore did not expand; instead, one vesicle gradually

shrank and disappeared over time as the result of the transfer of its vesicle content into

another vesicle. In homotypic vacuole fusion in *Saccharomyces cerevisiae,* vacuole

membranes fused at the vertices instead of a central pore expanding radially, and then a

boundary membrane between the two docked vacuoles was released into the vacuole

lumen (Wang et al., 2002). Because pore dynamics determine the rate of content mixing,

they have significant effects on the trafficking efficiency of cargo molecules. However,

how different modes of membrane deformation are induced and whether the distinct

fusion modes are exchangeable remain unknown.

Membrane deformation is generally controlled by membrane-associating proteins as

well as by several physical parameters including the vesicle volume, membrane area,

bending energy, and osmotic energy in cells (Farsad and De Camilli, 2003; Lipowsky,

1991). Actin and its regulating proteins, via interaction with the membrane, exert

physical forces on and cause fluctuation in the membrane, which eventually leads to

membrane deformation by modification of the physical parameters (Salbreux et al.,

2012). Actin produces forces in two different manners. First, assembly of branched

actin filaments, nucleated from the membrane surface by actin nucleators such as the

actin-related protein 2/3 (Arp2/3) complex and formins, provides pushing forces on the

membrane (Goley and Welch, 2006; Svitkina, 2018). Second, actin generates pulling

forces through myosin-driven sliding of actin filaments (Svitkina, 2018). Both forces

play roles in induction of membrane deformation during vesicle fusion (Ebrahim et al.,

2019; Ma et al., 2020; Tran et al., 2015; Uenishi et al., 2013). Many experiments have

been carried out to elucidate the roles of actin in vesicle fusion through manipulation of

actin dynamics by use of specific drugs or gene modifications. These experiments gave

rise to different results depending on the experimental conditions. For example,

polymerization or stabilization of actin enhanced membrane fusion in some experiments

but inhibited it in others (reviewed in Eitzen, 2003). These different results imply that

the regulation mechanisms of vesicle fusion by actin filaments are more complex than

previously assumed. Moreover, how the actin forces contribute to the fusion modes

remains unknown, as does how the force generated by actin polymerization or by actin-

dependent motor proteins contributes to membrane deformation during fusion.

In this study, we utilized mouse yolk sac visceral endoderm (VE) cells as a model

system for addressing the above questions. VE cells play a critical role in the

maternofetal exchange of nutrients before the establishment of a chorioallantoic

placenta: they vigorously endocytose maternal proteins, hydrolyze the proteins in

lysosomes, and supply the resultant products to the developing embryo (Bielinska et al.,

1999; Jollie 1990). Reflecting this high endocytic activity, VE cells are characterized by

huge endosomal vesicles (Bielinska et al., 1999; Kawamura et al., 2012; Koike et al.,

2009). These properties of VE cells enable us to observe endosomal dynamics including

vesicle fusion. Previously, by using a whole embryo culture system combined with

specific pharmacologic inhibitors for intracellular signaling molecules, we showed that

actin dynamics regulated by lysophosphatidic acid (LPA) receptors and the Rho-ROCK-

LIM kinase pathway are required for the formation of huge lysosomes (Koike et al.,

2009).

Here, we show that late endosomes of VE cells exhibit different modes of fusion with

target organelles depending on their targets: they undergo homotypic fusion with late

endosomes in an explosive fusion-like manner and heterotypic fusion with lysosomes in

a bridge fusion-like manner. Our mathematical modeling predicts that the vesicle size is

a critical determinant of the fusion types and that membrane perturbation overcomes the

size effects. We provide evidence that actin filaments are associated with late

endosomes and that actin dynamics generate forces promoting vesicle fusion of

endosomes for efficient membrane trafficking.

Results

Endocytic vesicles in mouse yolk sac VE cells

To visualize endosomal vesicles in yolk sac VE cells, an embryonic day 8.5 (E8.5)

whole embryo was labeled with a fluorescent membrane probe, FM1-43, and

subsequently stained with organelle markers. The embryo was put on a glass-bottom

dish, and VE cells were observed by use of confocal microscopy (Figure 1A). These

observations revealed that different types of endosomal vesicles were aligned from the

apical to the basal area in the supranuclear portion of VE cells, reflecting the repeated

fusion and enlargement of vesicles as they migrated deeper into the cell (Figure 1B and

C). In the most superficial portion (within 0.5–1 μm of the apical surface; designated as

zone 1, Figure 1–figure supplement 1A), fuzzy FM1-43 staining was observed (Figure

1B). The small vesicles in this zone were positive for EEA1 (Figure 1B), indicating that

they were early endosomes. Immediately beneath zone 1, many medium-sized round

vesicles were observed (zone 2; at a depth of 1–6 μm from the apical surface, Figure 1–

figure supplement 1A). Electroporation of EGFP-Rab7 revealed that these vesicles were

positive for Rab7, suggesting that they were late endosomes. In the deep portion (zone

3; at a depth of 6–16 μm from the apical surface, Figure 1–figure supplement 1A), large

amorphous lysosomes, positive for LysoTracker Red and LAMP1, were observed

(Figure 1B; Koike et al., 2009).

To examine whether these vesicles constitute endocytic pathways, we traced vesicle

trafficking in VE cells. After E8.5 whole embryos were pulse-labeled with Alexa 488-

transferrin for 5 min, fluorescently labeled endocytic vesicles were tracked over time.

The fluorescent signals were initially detected in small vesicles (area $< 2.0 \ \mu m^2$) in zone

1; 10–20 min after labeling, the signals were mainly detected in medium-sized vesicles

($2.0–25 \ \mu m^2$) in zone 2; 40 min after labeling, the signals were strongly observed in

large lysosomes ($> 25 \ \mu m^2$) in zone 3 (Figure 1D and E). The time course of the

numbers of labeled vesicles in each zone showed that endocytosed transferrin was

rapidly transported from early endosomes to lysosomes via late endosomes in VE cells,

as reported previously (Richardson et al., 2000). In addition, rhodamine-labeled dextran,

which was transported by pinocytosis pathways, was also transported to lysosomes in a

similar fashion (Figure 1–figure supplement 1B).

Two distinct modes of late endosomal fusion

To examine the dynamic changes of endocytic vesicles, we performed time-lapse

imaging of Alexa 488-transferrin- or FM 1-43-labeled vesicles. We focused on late

endosomes in zone 2 because these vesicles were large enough and suitable for

microscopic observation of membrane fusion processes, as described below.

First, from 5 min after pulse labeling, fusion of two neighboring late endosomes was

observed frequently (Figure 2A and B; Video S1). The fusion showed an explosive

mode, in which a membrane pore quickly expanded and two vesicles fused to become a

single vesicle (Figure 2B). The size distribution of the late endosomes that underwent

fusion ($7.3 \pm 6.6 \ \mu m^2$, n = 34) was similar to that of the total late endosomes in the

observed area at the start of recording ($5.7 \pm 6.4 \ \mu m^2$, n = 265; Figure 2C), indicating

that late endosomes of any size underwent homotypic fusion. The fusion was completed

within 20–50 s (the time required from membrane fusion to single spherical vesicle

formation was 31.9 ± 0.2 s, n = 37), and the time was weakly correlated with the vesicle

size (r = 0.60, Figure 2D).

Second, from 15 min after pulse labeling, some late endosomes shrank gradually and

disappeared from the focal plane (Figure 3A and Video S2). Because a previous paper

reported that late endosomes were engulfed by lysosomes in E6.5 mouse VE cells

(Kawamura et al., 2012), we examined the mechanism of the disappearance of late

endosomes in more detail using different approaches. First, we used differential labeling

of late endosomes and lysosomes. When whole embryos were incubated with

rhodamine-dextran for 20 min and subsequently pulse-labeled with Alexa 488-

transferrin for 5 min, lysosomes and late endosomes were labeled with dextran and

transferrin, respectively, at the time of observation (15 min after transferrin labeling). In

a diminishing late endosome, the fluorescence intensity of transferrin in the vesicle

increased initially and then decreased quickly (Figure 3B–D). At the same time, the

rhodamine-dextran in the adjacent lysosome entered the late endosome (Figure 3B–D),

indicating that the vesicular contents of the late endosomes and lysosomes were mixed.

Second, using electron microscopy, we observed pore formation between a late

endosome and a lysosome (Figure 3E). These data suggest that the disappearance of late

endosomes is a consequence of fusion with the underlying lysosomes, but not of

engulfment in E8.5 embryos. This type of fusion, so-called bridge fusion, described in a

previous study (Nicot et al., 2006; Roberts et al., 1999), occurred as a result of the slow

mixing of vesicular contents through a narrow fusion pore. Actually, heterotypic fusion

was slower than homotypic fusion and was completed within 90–250 s (154.8 ± 1.4 s, n

= 25), and the fusion time (the time required from membrane fusion to disappearance of

the vesicle) did not correlate with the vesicle size (Figure 3F). The size of the labeled

late endosomes at this time point (10.1 ± 8.1 μm^2, n = 256) was larger than that at 5 min

after labeling (6.7 ± 6.2 μm^2, n = 242), indicating that larger late endosomes tended to

undergo heterotypic fusion (Figure 3G, see also below). However, the size of the late

endosomes that underwent heterotypic fusion (10.2 ± 9.2 μm^2, n = 34) did not differ

from that of the total late endosomes at 15 min after labeling (Figure 3G), indicating

that the size of late endosomes is not the sole trigger for heterotypic fusion.

Vesicle size is an important factor for the transition of fusion modes

To examine whether the differences in membrane deformation of late endosomes

depend on their fusion targets, we adopted a mathematical modeling approach that

describes the membrane deformation processes after pore formation. According to

thermodynamics, the deformation of a physical object proceeds as its free energy

decreases. The free energy of a vesicle consists of two terms: bending energy and

osmotic energy. The bending energy, b, is the integral of the square of the mean

curvature all over the surface (Helfrich, 1973). The osmotic energy is the product of the

osmotic pressure difference between the inside and the outside of the vesicle, Δp, and

the vesicle volume, V. Thus, the free energy is given by

$$F = b - \Delta p \cdot V.$$

Note that the bending energy depends only on its shape; in other words, two membranes

with the same shape but with different sizes have the same bending energy. On the other

hand, the volume depends on both the size and the shape. Therefore, the vesicle size

intrinsically affects the balance of both energy sources and may be a critical factor for

membrane deformation during fusion.

We focused on a closed membrane having a constricted neck region and introduced

two shape parameters, the neck width, w, and the area ratio between both sides of the

neck, a (Figure 4A), so that the basic processes of the two fusion types are described by

the changes in w and a just after pore formation, $w = 0$. The relaxation of the neck in

explosive fusion is expressed by the increase in w. On the other hand, the absorption of

smaller vesicles by the larger one in bridge fusion is the decrease in a while w remains

small.

When calculating the free energy as a function of w and a, the deformation is

expressed as the process of descending the free-energy landscape in the space. Because

the shape giving the minimum free energy is most likely to be realized, we first

specified the minimum-free-energy shape and then calculated the free energy value for

each parameter set.

We considered two extreme conditions: the membrane size is sufficiently small or

large. In the former case, the osmotic energy is negligible and the free energy is

approximated to

220
$$F \simeq F^{\text{small}} = b.$$

221 The free-energy landscape on the shape parameter space (w, a) is shown in Figure 4B.

222 In the latter case, although the osmotic energy is dominant, the bending energy is also

223 not negligible because the bending energy at the neck region becomes considerably

224 large. Thus, the free energy for the large membrane case is

$$F \simeq F^{\text{large}} = B^{\text{neck}} - \Delta p \cdot V,$$

where B^{neck} is the bending energy at the neck. The free-energy landscape for a vesicle

with the projected area of $10[\mu m^2]$ is shown in Figure 4C.

These free-energy landscapes (Figure 4B and C) clearly differ from each other,

implying that they drive different deformation processes. In particular, the contour of

the F^{small} landscape around $w \sim 0$ is perpendicular to the w axis, whereas the contour of

the F^{large} landscape is parallel to the w axis. Typical orbits of the shape deformations

from the initial condition $(w, a) = (0.0, 0.3)$ generated by the gradients of the landscape

are shown as gray arrows in Figure 4B and C. The orbit in the F^{small} landscape is

characterized by a smooth increase in w; i.e., the neck expands quickly (Figure 4D),

resulting in explosive fusion. On the other hand, in the F^{large} landscape, a increases

through the orbit, whilst w remains small, which evokes bridge fusion (Figure 4E).

The deformation processes observed in our modeling seem to be qualitatively

consistent with our experimental results showing that the size of the fusion target

changes the fusion modes (Figures 2 and 3). Next, we made a rough estimate of how

much of the vesicle size the mathematical model expects to switch the types of fusion.

The typical bending energy of a (spherical) vesicle is $b \simeq 2 \times 10^{-18}[J]$ (Evans and

Rawicz, 1990). Referring to the density differences in the dominant ions of endosomes

and cytoplasm (Scott and Gruenberg, 2011), we assume that the total ion difference is in

the range of 1– 10 mM, and that the osmotic pressure difference is $\Delta p \simeq 2.5 \sim 25 [Pa]$.

From these assumptions, $0.22– 1.0 [\mu m^2]$ would be the vesicle size to switch fusion

modes. However, the size of the observed late endosomes (5–20 μm^2 in Figure 3F) is

much larger than the predicted size, suggesting that homotypic fusion between late

endosomes should show the bridge fusion mode.

To resolve these inconsistencies, we considered the fluctuation effects. Intracellular

membranes receive various fluctuations including thermal noise and active fluctuation

from the cytoskeletons and motor proteins (Biswas et al., 2017; Chen et al., 2009).

Although orbits descending the F^{large} landscape are energetically restricted to keep w

small, these fluctuations may force the orbits to deviate from the restricted path (the

dashed arrow in Figure 4C); i.e., they are expected to induce explosive fusion. Then, we

performed Monte-Carlo simulations to generate orbits descending the free-energy

landscape with various magnitudes of fluctuations and determined the probabilities of

orbits that deviate from the path (Figure 4F). When we consider only the thermal noise

with room temperature with the estimated osmotic pressure difference ($\Delta p \simeq$

$2.5– 25 [Pa]$), as for late-endosome-sized vesicles ($2.0– 25 [\mu m^2]$), the simulation

predicts that vesicles will show bridge fusion (the green-shaded region in Figure 4F).

However, if we take into account the fact that a larger fluctuation allows explosive

fusion (the blue-shaded region in Figure 4F), fluctuations generated by the cytoskeleton

and motor proteins, much larger than thermal noise, may induce explosive fusion.

Actin filaments are bound to late endosomes in VE cells

Previously we found that lysosome biogenesis was regulated by actin dynamics in VE

cells (Koike et al., 2009). To examine the relationship between actin filaments and

endocytic vesicles, actin filaments were visualized by use of Alexa 488-phalloidin.

Endocytic vesicles were visualized by immunostaining with anti-mouse IgG because

IgG is endocytosed from the maternal tissue *in utero* and can be stained without

incubation of embryos *in vitro* for the pulse labeling of endosomes. In VE cells, strong

signals were detected to be associated with the cell membrane (Figure 5A–C). In

addition, in zone 1, fuzzy cytoplasmic staining was observed (Figure 5A), which

appeared to correspond to the actin meshwork in the terminal web. In zone 2, small

punctates and short filaments were detected in the cytoplasm (Figure 5B) and were

bound to the surface of late endosomes (Figure 5B, B', and B'') and infrequently to

lysosomes in zone 3 (Figure 5C, C', and C''). Optical cross-sectional images and 3D

reconstruction showed that several rod-like actin filaments, extending from the apical

side of VE cells into the cytoplasm, surrounded and held the late endosomes (Figure 5,

D–F). These data indicate that most actin punctates observed in the x-y plane

correspond to the cross-sectional images of actin filaments extending in the apical-to-

basal direction. In addition, when a fusion protein of an actin-bundling protein, fascin-1,

with a red fluorescent protein was expressed in VE cells by electroporation, fascin-1

was highly colocalized with the actin punctates, suggesting that most of the actin

punctates surrounding the late endosomes were bundled (Figure 5G).

To investigate the endosome-associated actin filaments in more detail, we performed

high-resolution imaging analysis using a deconvolution technique, which had resolution

comparable to super-resolution microscopy. It demonstrated that fine actin filaments

extended radially from the endosomal membranes in the x-y plane, in addition to the

actin filaments extending in the apical-basal axis (Figure 5H).

To examine the dynamics of actin filaments associated with late endosomes, E7.5

whole embryos were electroporated with a pEGFP-actin vector, and the EGFP signals in

VE cells were observed by use of confocal microscopy 24 h after electroporation.

Because high levels of actin overexpression led to the deterioration of cell functions

(see white arrows in Figure 6–figure supplement 1A), cells expressing low levels of

EGFP-actin were selected for analysis: those selected cells were normal in morphology

and endocytosis (see white arrowheads in Figure 6–figure supplement 1A). Punctate

signals of EGFP-actin were seen on the late endosomal membrane similarly to those

observed in Alexa 488-phalloidin labeling. Time-lapse imaging showed that some new

spots had appeared, others had disappeared, and two actin spots were unified on the

membrane surface of endocytic vesicles (Figure 6A). To examine the turnover rate of

actin spots in VE cells, we performed fluorescence recovery after a photobleaching

(FRAP) assay. After EGFP-actin on a late endosome was photobleached, recovery of its

fluorescence was measured (Figure 6B). Recovery of actin spots on late endosomes

occurred more rapidly than the control recovery measured at the cell surface ($t_{1/2} = 29.0$

s; Figure 6C-E). Especially, the recovery of actin spots in the apical area of late

endosomes was faster than in the basal area ($t_{1/2} = 11.5$ s in the apical area vs. $t_{1/2} = 18.9$

s in the basal area; Figure 6C, E). These results indicated that actin turnover on the

surface of late endosomes was dynamically regulated, with a tendency to be more active

in the apical area than in the basal area. The time-lapse images showed that when late

endosomes moved in an x-y plane in zone 2, spots of EGFP-actin were always at the

rear-end of the endosomes along the moving direction (Figure 6F, n = 13). Because it is

known that Arp2/3-dependent actin polymerization on the membrane acts as the driving

force of the motility of intracellular pathogens (Welch and Way, 2013) and

endosomes/lysosomes in *Xenopus* egg extracts (Taunton et al., 2000), we examined the

localization of Arp2/3 in VE cells to test the possible role of Arp2/3 in endosome

migration. When a fusion protein of Arp3-EGFP was expressed in VE cells by

electroporation, the EGFP signals were distributed in dots on the late endosomes and

partially colocalized with phalloidin spots in the x-y plane in zone 2 (Figure 6–figure

supplement 1B). In addition, Arp3-EGFP signals were always seen at the rear-end of the

endosomes along the moving direction (Figure 6–figure supplement 1C). Taken

together, these data suggest that the radial branched filaments are temporally

polymerized to provide the propulsive force on the membrane for intracellular

movement.

Actin dynamics regulate homotypic fusion of late endosomes

To examine the roles of actin in late endosomal fusion, we tested whether perturbation

of actin dynamics caused defects in vesicle fusion. Treatment with cytochalasin D (a

cell-permeable inhibitor of actin polymerization) for 5 min significantly changed the

distribution of actin filaments. High-resolution imaging analysis using a deconvolution

technique showed that phalloidin-positive filamentous structures decreased and instead

actin aggregation was observed in the cytosol (Figure 7–figure supplement 1A). The

treatment reduced the frequency of homotypic fusion in a dose-dependent manner

(Figure 7A and B; Video S3). In addition, it altered the speed and mode of the fusion.

Treatment with 0.1 or 1.0 μM cytochalasin D slowed down homotypic fusion (Figure

7C and D; Figure 7–figure supplement 1B). After membrane fusion, two endosomes

remained in a peanut shape for a long time, indicating slower neck expansion than that

in the control cells (Figure 7–figure supplement 1B). FM 1-43 labeling revealed that the

membranes were rapidly fused, whereas it took a long time for the gourd-shaped

vesicles to form a large round vesicle (Figure 7–figure supplement 1B). Furthermore,

upon 1.0 μM cytochalasin D treatment, about one-third of homotypic fusion showed

bridge fusion (Figure 7C and 7E; Video S4). Therefore, cytochalasin D slowed down

the fusion processes and changed the fusion mode to bridge fusion at higher

concentrations. These data are consistent with our mathematical model showing that

larger fluctuation is necessary for explosive fusion in homotypic fusion of late

endosomes (Figure 4F). Treatment with jasplakinolide (a cell-permeable stabilizer of

actin filaments) also changed the actin dynamics. After the treatment, the actin

meshwork beneath the plasma membrane (zone 1) was denser (Figure 7–figure

supplement 2). Furthermore, actin filaments extending radially from the late endosomes

were thicker and the fluorescence intensity was increased, although the actin filaments

extending from the apical-basal axis were not changed (Figure 7–figure supplement 2).

The treatment reduced the frequency of homotypic fusion in a dose-dependent manner

(Figure 7A and B; Video S5), indicating that dynamic actin turnover is required for

homotypic fusion.

We next observed the distribution of actin filaments during homotypic fusion. When

late endosomes underwent fusion, EGFP-actin punctates were excluded from the

membrane docking sites and highly polymerized transiently at both ends of the fusing

vesicles (opposite to the docking region) during the rounding period after membrane

fusion (Figure 7F). The fluorescence intensity of EGFP-actin on endosomes was

quantitated in the proximal, lateral, and distal regions at the predocking, docking,

rounding, and postfusion stages. This analysis indicated that actin filament distribution

was spatially regulated during fusion (Figure 7G, dark gray bar), whereas FM 1-43

distribution, used as an unchanged control, was uniform on the membrane throughout

the rounding stages (Figure 7G, light gray bar). Furthermore, the total fluorescence

intensity of EGFP-actin on late endosomes significantly increased transiently after pore

formation (Figure 7H), indicating that actin filaments were polymerized temporarily,

not simply distributed differently. However, cytochalasin D treatment stabilized EGFP-

actin spots and thereby reduced movement of EGFP-actin spots and temporal

polymerization during homotypic fusion (Figure 7–figure supplement 3). Moreover,

accumulation of Arp3-EGFP in the distal region during the rounding stage was observed

similarly to EGFP-actin accumulation (Figure 7–figure supplement 4), and treatment

with CK-666, an Arp2/3 inhibitor, resulted in a decrease in the frequency of homotypic

fusion (Figure 7B) and slowed down homotypic fusion (Figure 7C and D). These results

suggest that branched actin filaments nucleated by the Arp2/3 complex on the distal

surface of late endosomes are necessary to transform the vesicle into a large round

sphere quickly after membrane fusion of late endosomes.

Homotypic and heterotypic fusions are differentially regulated by actin

We next examined the role of actin filaments in heterotypic fusion. Cytochalasin D

treatment led to a significant reduction in the frequency of heterotypic fusion in a dose-

dependent manner (Figure 8A–C and Video S6). In contrast, jasplakinolide treatment

did not inhibit heterotypic fusion and somehow increased the frequency of heterotypic

fusion with lysosomes at as early as 5 min after pulse labeling (Figure 8B and Video

S7). In addition, CK-666 treatment had no effects, suggesting that Arp2/3 was not

required for heterotypic fusion (Figure 8C). These data suggest that actin played distinct

roles in homotypic fusion and heterotypic fusion and that stable actin filaments are

implicated in driving heterotypic fusion.

Myosins are a superfamily of actin-based motor proteins and their members are

grouped into many classes according to their structures and functions (Hartman and

Spudich, 2012). To determine which class of myosins is important for vesicle fusion

and motility, we examined the effects of inhibitors for several myosin classes that are

known to be involved in vesicle transport. We tested inhibitors of myosin I

(pentachloropseudilin, PClP), nonmuscle myosin II (NM II) (blebbistatin), myosin V

(MyoVin-1), and myosin VI (2,4,6-triiodophenol, TIP; Bond et al., 2013). Heterotypic

fusion frequency was reduced to 20% by MyoVin-1 and to a lesser extent, but

significantly, by blebbistatin, but was not affected by PClP or TIP (Figure 8D). In

contrast, homotypic fusion frequency was significantly reduced by PClP, MyoVin-1,

and TIP (Figure 8–figure supplement 1). EGFP-myosin IIA signals overlapped with

phalloidin-positive spots on late endosomes (Figure 8E), whereas EGFP-myosin Va

signals were juxtaposed to phalloidin-positive spots on late endosomes (Figure 8E).

Cofilin-dependent actin dynamics are involved in late endosome fusion

Late endosomes undergo homotypic fusion almost exclusively at the beginning and then

switch to heterotypic fusion over time after endocytosis (Figure 9A). Given that actin

plays a different role in homotypic fusion and heterotypic fusion and that pharmacologic

changes in actin dynamics shift the fusion mode as shown above, it is suggested that

actin dynamics are a critical factor for the transition of fusion modes. Thus, we next

decided to look into the regulation of actin dynamics in two different fusion modes.

Previously, we found that lysosome biogenesis was regulated by actin dynamics

through cofilin downstream of the autotaxin–LPA receptor–Rho–ROCK–LIM kinase

pathway in VE cells (Koike et al., 2009). When VE cells were incubated with the S3

peptide, which contains the N-terminal 16 amino acid sequence of cofilin fused to a cell

permeation sequence and is used as a competitive inhibitor for LIM kinase and serves as

a cofilin activator (Maekawa et al., 1999; Aizawa et al., 2001), lysosomes were

fragmented in VE cells and the fluorescence intensity of phalloidin in VE cells was

reduced (Koike et al., 2009).

We thus focused on cofilin in endosomal fusion in VE cells. YFP-cofilin showed

small punctate staining that colocalized with phalloidin-positive spots in the x-y plane

and extended along the phalloidin signal (Figure 9B and C). When the fluorescence

intensity of Alexa 546-phalloidin- and YFP-cofilin-positive filaments was quantitated

along the apical-basal axis of late endosomes in zone 2, cofilin was slightly abundant in

the apical region, whereas the phalloidin signal was stronger in the basal region (Figure

9D). Note that the possibility that overexpression of YFP-cofilin changed the

distribution of actin filaments cannot be excluded. EGFP-actin on a late endosome

showed that treatment of E7.5 whole embryos with the S3 peptide (15 μg/ml) for 24 h

resulted in a decrease in radially extended actin filaments in the apical area of zone 2,

whereas the actin filaments extending along the apical-basal axis appeared unchanged

(Figure 9–figure supplement 1). In addition, a FRAP assay showed that the treatment of

S3 peptide induced faster turnover of actin filaments of late endosomes when compared

with treatment with the RV peptide (Aizawa et al., 2001), which contains the reverse

sequence of cofilin and is used as a negative control ($t_{1/2}$ = 10.9 s with the S3 peptide, n

= 9 vs $t_{1/2}$ = 18.3 s with the RV peptide, n = 9; Figure 9E). These results indicate that

cofilin distributes with actin filaments on late endosomes and regulates actin dynamics

on late endosomes. Finally, the roles of cofilin in late endosomal fusion were

investigated. After treatment with the S3 peptide for 24 h, Alexa 488-transferrin-labeled

vesicles were observed by use of time-lapse imaging. When compared with the RV

peptide, the S3 peptide strongly inhibited homotypic fusion (Figure 9F) but not

heterotypic fusion (Figure 9G). These data suggest that spatial differentiation of actin

dynamics along the apical-basal direction, partially contributed to by cofilin, may be a

critical factor for the transition of fusion modes (Figure 10).

Discussion

In this study, we developed a novel imaging method of analyzing the motile behavior of

late endosomes in VE cells and found that VE cells have two different types of vesicle

fusion regulated by distinct actin-mediated mechanisms.

Actin filaments are associated with endosomal vesicles

We found two types of actin filaments associated with late endosomes in VE cells: those

that extended along the apical-basal axis and those that extended radially from the

endosomes (Figure 5). Our findings resemble the previous observation that early

endosomes were distributed along with actin cables and that Arp2/3-dependent small

cloud-like patches protruded from the membrane (Morel et al., 2009). Actin is

reportedly distributed around vesicles in different patterns. In addition to the well-

known interaction of vesicles with actin rails in directional vesicle transport (Titus,

2018), actin was localized at the vertex ring in vacuole fusion in yeast (Wang et al.,

2002) and entirely surrounded phagosomes in macrophages (Liebl and Griffiths, 2009).

Advanced microscopy techniques have clarified the fine structure of actin, which is

assembled into polyhedron-like lattices around vesicles in exocrine glands (Ebrahim et

al., 2019).

We observed strong actin signals as patches on late endosomes in the focal planes on

confocal microscopy. Three-dimensional reconstruction showed that most of these

patches represent the cross-section of rod-shaped actin filaments extending from the

apical side into the cytoplasm in VE cells. In addition to these patches, some signals

seem to represent actin filaments extending radially from membranes. Furthermore, we

showed that Arp3-positive actin spots appeared in the site opposite to that of endosome

movement (Figure 6F, Figure 6–figure supplement 1C) like a comet tail (Goley and

Welch, 2006), that Arp3-positive actin spots were asymmetrically distributed and

temporarily polymerized in the distal region during endosomal fusion (Figure 7F and G;

Figure 7–figure supplement 4), and that Arp2/3 inhibition reduced homotypic fusion

(Figure 7B). These results suggest that radial actin filaments polymerized by Arp2/3

generate force for homotypic fusion in VE cells. This notion is compatible with the

findings of previous reports that the radial filaments were nucleated and extended with

Arp2/3-dependent branches on the lateral membrane of endosomes to push vesicles

(Goley and Welch, 2006; Svitkina, 2018). Given that nucleation and Arp2/3 activities

are regulated by nucleation-promoting factors (NPFs; Goley and Welch, 2006), the

activity of NPFs and membrane fusion processes are likely tightly linked to accurate

membrane fusion. Differential regulation of NPF activity may give rise to different

patterns of actin surrounding vesicles and membrane dynamics. Future investigation is

necessary to elucidate how the NPF activity is spatially and temporally regulated on

membranes.

Actin filaments distribute on endosomes in various patterns depending on the cell

types. However, why cells show different patterns of actin filament distribution on

vesicles and whether different distribution patterns of actin filaments have different

functions remain unclear. In this study, by taking advantage of the large size of

endosomes, we found that there are two different types of actin filament distribution on

the surface of late endosomes in VE cells and that these actin filaments regulate

different aspects of membrane fusion, movement, and endosomal distribution. Taken

together, these data suggest that different patterns of actin filament distribution on

vesicles have various effects on specific functions of vesicles.

Actin regulates membrane behavior after pore formation

In this study, we found that the membranes of late endosomes behave differently

depending on the fusion mode. In homotypic fusion, the fusion pore formed between

docked endosomal vesicles expanded quickly and the fusion was completed within 20−

30 s (explosive fusion, Figure 2). In contrast, in heterotypic fusion with lysosomes, the

pore opened very slowly and late endosomes gradually shrank in diameter and

disappeared over 1−2 min (bridge fusion, Figure 3). Using mathematical modeling, we

showed that vesicle size is an important factor determining the fusion mode. When the

size of the vesicle becomes larger, the free energy for neck relaxation decreases and the

pore is kept small. Our modeling predicted that the late endosomes are large enough to

allow bridge fusion to occur, although they exhibit explosive fusion *in vivo*. Similarly,

homotypic fusion in an explosive mode was observed in very large vesicles, such as

vacuoles in yeast, endosomal vesicles in coelomocytes of *C. elegans*, and secretory

granules in the salivary glands (Liu et al., 2016; Masedunskas et al., 2011; Wang et al.,

2002). These observations indicate that vesicle size is not the sole factor determining the

fusion mode. Lipid composition, a well-known parameter affecting its bending modulus,

is also involved, but the effect on fusion mode may be less dominant when compared

with the vesicle size effects (Simon et al., 1998; Faizi et al., 2019; Ridolfi et al., 2021).

One of the critical factors other than vesicle size that determine the fusion mode is

membrane fluctuation. We found that homotypic fusion was converted to bridge-like

fusion after cytochalasin D treatment (Figure 7 C), which suggests that actin dynamics

around vesicles cause fluctuation and affect the fusion modes. This finding can be

successfully explained by introducing minor modifications in our mathematical

modeling so that a larger fluctuation can shift the fusion mode from bridge to explosive

fusion.

VE cells vigorously endocytose maternal proteins and supply the digested products to

the developing embryo to support embryonic development. This seems to be why VE

cells have such huge endosomal and lysosomal vesicles. However, larger vesicle size

leads to the reduction in the vesicle fusion speed and subsequent nutrient trafficking

efficiency. If vesicle size were the only factor determining the fusion mode, this

disadvantage would occur, but if fluctuation is taken into account, it can be overcome,

as shown in our mathematical modeling, and the forces generated by actin are thought

to be responsible for increasing the rate of vesicular trafficking. This idea is supported

by previous reports that knockout mice of actin-related genes (e.g., *Arc* and *Myh9*)

showed abnormal VE cells and embryonic lethality (Conti et al., 2004; Liu et al., 2000).

In heterotypic fusion, late endosomes fused with lysosomes in a bridge-like manner.

This may be because lysosomes are too large in comparison with late endosomes and

actin accumulation on the lysosomal membrane is scarce, resulting in insufficient force

of actin for explosive fusion. Bridge fusion was previously observed in live cells only

when giant vesicles were artificially generated by gene modification, for example, in

endosomes enlarged by expression of constitutively-active Rab5 (Roberts et al., 1999)

and lysosomes enlarged in the intestinal cells of *ppk-3*-deleted *C. elegans* (Nicot et al.,

2006). These findings suggest that the enlargement of these vesicles exceeds the

intrinsic forces of actin to induce explosive fusion.

Like homotypic fusion, heterotypic fusion is also dependent on actin, as shown by

reduction in fusion frequency upon cytochalasin D treatment. Intriguingly, in

heterotypic fusion, NM II, myosin V, and stabilized actin filaments, but not Arp2/3,

were required, indicating a distinct contribution of actin to heterotypic fusion when

compared with homotypic fusion. We showed that, consistent with the well-known roles

of myosin V in trafficking of lysosomes, ER, melanosomes, and synaptic vesicles

(Hammer III and Sellers, 2011), myosin V facilitates the directional transport of late

endosomes towards lysosomes along stabilized actin filaments that serve as a rail

(Figure 10). Moreover, our results indicate that NM II, which is primarily responsible

for generating contractile force (Svitkina, 2018), partially contributes to heterotypic

fusion. It is well established that NM II facilitates the constriction of the membrane into

the plasma membrane after membrane pore formation in exocytosis (Ebrahim et al.,

2019) in cooperation with actin depolymerization by cofilin-1 (Miklavc et al., 2015). In

contrast, our data showed that actomyosin appeared not to form a lattice on the surface

of endosomes for the contraction and that cofilin did not show a robust influence on

heterotypic fusion. Rather, actin stabilization tended to promote heterotypic fusion

(Figure 8A and 8B). These data suggest that NM II may contribute to heterotypic fusion

through a different mechanism from that of exocytosis. NM II is also implicated in the

movement of the nucleus along with cell migration, which is akin to a "conveyor-belt"

by the sliding of the nuclear membrane along the actin filament (Gomes et al., 2005).

Further investigation is necessary to elucidate the mechanisms by which NM II

transports late endosomes along actin filaments.

Determination of fusion targets

Late endosomes, which fuse frequently with neighboring late endosomes, shift the

target of fusion towards lysosomes. What triggers the transition of the fusion targets?

We found that larger endosomes tend to fuse with lysosomes (Figure 3G). Thus, one

possibility is that the larger endosomes located in the basal position have a higher

opportunity to fuse with lysosomes than do the smaller endosomes in the more apical

portion. However, upon treatment with jasplakinolide, even smaller endosomes

distributing in the more apical area underwent heterotypic fusion with lysosomes

(Figure 8A and B), thus suggesting rather that the size and the position of vesicles are

not the sole factors and that the different dynamics of actin shift the fusion modes.

Furthermore, we showed weaker actin polymerization/depolymerization activity in the

basal area as compared with that in the apical area (Figure 6C), a decrease in the

frequency of homotypic fusion but not of heterotypic fusion induced by S3 peptide

treatment (Figure 9F and G), a decrease in radially extending actin filaments by S3

peptide treatment (Figure 9–figure supplement 1), and an increase in actin filaments on

endosomes in the basal area as compared with in the apical area (Figure 9D). These

findings support the notion that cofilin-dependent regulation of actin

polymerization/depolymerization may play a critical role in determination of the fusion

target. In parallel with VE cells, it was shown that cofilin was specifically distributed in

the apical area of epithelial cells and was highly active under the plasma membrane

(Morris et al., 2011; Vitriol et al., 2013). Cofilin regulates actin remodeling for

lamellipodia formation and endosomal trafficking (Harrington et al., 2011; Vitriol et al.,

2013). Furthermore, early endosomes located close to the plasma membrane fuse

frequently when compared with late endosomes located at the perinuclear region

(Gruenberg et al., 1989). Taken together, higher dynamics of actin filaments regulated

by cofilin at the cell peripheral area may increase the fusion frequency in many cells.

In this study, we developed a method of observing and manipulating late endosomes

in VE cells. This method enables, for the first time, the observation of fusion of single

vesicles with two different targets and is very useful to examine the molecular

mechanisms that determine the fusion targets of vesicles and regulate different fusion

modes.

Materials and Methods

Animal experiments

All the experiments using animals were approved by the Animal Care and Use

Committee of the University of Tsukuba and performed under its guidelines. Noon of

the day on which a vaginal plug was observed was taken as embryonic day 0.5 (E0.5).

Time-pregnant ICR mice were purchased from CLEA Japan and Japan SLC.

Immunohistochemistry and phalloidin staining

For whole-mount immunohistochemistry, E8.5 embryos were fixed with 4%

paraformaldehyde (PFA) in PBS at 4°C for 3 h and incubated with FITC-conjugated

anti-EEA1 (1:100, BD Bioscience) or Cy3-labeled anti-mouse IgG (1:500, Jackson

ImmunoResearch). For phalloidin staining, the embryos were fixed with 4% PFA in

PBS at 37°C for 15 min and incubated with 0.2 U/ml Alexa Fluor 488- or Alexa Fluor

546-conjugated phalloidin (Molecular Probes) for 60 min.

Ex vivo **whole embryo culture**

The whole embryo culture was performed as described previously (Koike et al., 2009).

In brief, embryos were dissected at E7.5 and cultured in 100% rat serum (Charles River)

supplemented with 2 mg/ml glucose in a culture bottle placed in a rotation drum culture

system (Ikemoto Rika) at 37°C under 5% O_2, 5% CO_2, and 90% N_2 for 24 h. In the

pharmacologic experiments, embryos were incubated with 15 µg/ml S3 and RV

peptides (a kind gift of Dr Kensaku Mizuno, Tohoku University).

Endocytic uptake assays and intracellular vesicle tracing

Whole embryos with intact yolk sacs dissected out at E8.5 or embryos cultured for 1 d

from E7.5 were transferred into DMEM (Sigma-Aldrich) supplemented with 0.1% BSA.

Embryos were incubated in 100 nM LysoTracker Red (Molecular Probes) or 12 µM FM

1-43 (Molecular Probes) in 0.1% BSA/DMEM at 37°C for 5 min. For tracing endocytic

vesicles, embryos were incubated with 20 µg/ml Alexa Fluor 488-labeled transferrin

(Molecular Probes) or 100 µg/ml dextran rhodamine B 70,000 MW (Molecular Probes)

at 37°C for 5 min. For differential labeling of late endosomes and lysosomes, whole

embryos were initially cultured in 100% rat serum supplemented with 2 mg/ml glucose

and 75 µg/ml dextran rhodamine B 70,000 MW at 37°C for 20 min and then incubated

with 20 µg/ml Alexa Fluor 488-labeled transferrin for 5 min. To test the effects of actin

inhibitors, embryos were incubated with 0.01–1 µM cytochalasin D (Sigma-Aldrich),

0.01–1 µM jasplakinolide (Calbiochem), 20 µM CK666 (Calbiochem), 10 µM

pentachloropseudilin (AOBIOUS), 10 µM blebbistatin (Sigma-Aldrich), 10 µM

MyoVin-1 (Calbiochem), and 10 µM 2,4,6-triiodophenol (Sigma-Aldrich) for 5 min,

and the same drugs were added to the medium during observation.

Image acquisition and analysis

An embryo was placed on a glass-bottom dish (No.1S thickness, Matsunami Glass), and

confocal fluorescent images were obtained by use of an LSM510 laser scanning

microscope (Carl Zeiss) equipped with a C-APOCHROMAT 63x/1.2 W Korr objective

lens (Carl Zeiss) and a pinhole set to 1.0 AU. Images of zone 2 and zone 3 were taken

by focusing on the z-axis where late endosomes and lysosomes occupied the largest area

in the observation field, respectively. To compare endosomal sizes, the area of all intact

endosomes in each image was measured by use of Image J. Time-lapse images were

obtained by collecting 40 images with time intervals of 5 or 10 s at room temperature.

The series of images of the confocal z-stacks were obtained with a lateral resolution of

1024 x 1024 pixels and a scanning depth of 6–8 µm with 13–17 layers and 0.5-µm

intervals. To quantify fusion frequency, the number of membrane fusion events in the

400-s time-lapse images was counted, and the number of fusions per minute a single

late endosome underwent was calculated. To compare the time required for single

fusion processes, fusion time was defined as follows. For homotypic fusion, fusion time

was defined as the time from the start of membrane fusion of two late endosomes until

the formation of a single, round vesicle. For heterotypic fusion, fusion time was defined

as the time from when the late endosomes began to shrink until they disappeared as the

result of absorption by a lysosome.

For high-resolution imaging, the samples on the glass-bottom dish were mounted with

ProLong Glass (Thermo Fisher Scientific). Images were captured by use of a TCS SP8

laser confocal microscope (Leica Microsystems) equipped with an HC PL APO CS2

100×/1.40 OIL objective (Leica). Alexa Fluor 488 and Alexa Fluor 594 dyes were

excited at 488 and 594 nm of a supercontinuum white light laser, respectively. The

emitted fluorescence was sequentially detected by use of HyD-SMD detectors through a

spectral separation module (505–550 nm for AlexaFluor 488 and 605–700 nm for Alexa

Fluor 594). The pinhole size was set at 0.40 Airy units (60.7 μm). The pixel size and z-

step were set at 45 nm and 182 nm, respectively. The acquired images were processed

using the Lightning deconvolution algorithm on LAS-X software (Leica). Three-

dimensional reconstructions of the images were carried out on NIS-Elements software

(Nikon) or Imaris software (Bitplane).

Electroporation of whole embryos

The electroporation of whole embryos was described previously (Koike et al., 2009).

An E7.5 embryo was inserted into a small hole of an agarose mold (2% agarose/PBS, 7

x 7 mm) placed in Hanks buffered saline solution (HBSS). The solution bathing the

embryos was then replaced with HBSS containing 2 mg/ml DNA. Three square electric

pulses (30 V, 50 ms duration, 1 pulse/s) were delivered to the agarose mold by use of an

electroporator (CUY21, Nepa Gene) and a 10-mm electrode (CUY650P10, Nepa Gene).

After electroporation, the embryo was rinsed with HBSS and cultured as described in

the previous section. Rab7, nonmuscle myosin IIA, and myosin Va subcloned into a

pEGFP-C2 expression vector (Clontech), Arp3 subcloned into a pEGFP-N1 expression

vector (Clontech), fascin-1 subcloned into a pmRFP-C1 expression vector (Clontech),

pEGFP-Actin (BD Biosciences), and YFP-cofilin (kind gifts of Dr Kensaku Mizuno)

subcloned into the pEGFP-C1 vector (Clontech) were used.

Photobleaching assay

Photobleaching analysis was performed with the LSM510 laser scanning microscope.

The EGFP-actin-electroporated embryo was observed by use of a 63x objective lens.

For the bleaching, an ROI was drawn around a late endosome (4 μm x 4 μm) or cell

boundary and was bleached by use of a 488-nm laser beam for 8 bleach iterations at

100% intensity, and fluorescence recovery into the bleached area was monitored. To

calculate the percentage of recovery after photobleaching, the intensity profiles for the

actin filaments in the bleached area, a background region, and the whole cells were

collected. The background was subtracted from all the data, and the intensity of the

whole cells was used to correct for fluorescence loss during image acquisition.

Electron microscopy

E8.5 mouse embryos were fixed with 2% glutaraldehyde and 2% PFA in 0.1 M

phosphate buffer (pH 7.4) overnight. After 3 washes with 0.1 M phosphate buffer, pH

7.4, the embryos were postfixed with 1.0% osmium tetroxide in 0.1 M phosphate buffer

for 2 h, dehydrated, and embedded in Epon 812 according to a standard procedure.

Ultrathin sections were stained with uranyl acetate and lead citrate and observed by use

of an electron microscope (H-7100, Hitachi).

Acknowledgments

This work was supported in part by KAKENHI (17024006) and the 21st Century COE

program from the MEXT, Japan, KAKENHI (JP22H04926), Grant-in-Aid for

Transformative Research Areas – Platforms for Advanced Technologies and Research

Resources "Advanced Bioimaging Support" from JSPS, Japan, and Takeda Science

Foundation. We thank Drs K. Mizuno, K. Ohashi, and N. Ohsumi, for reagents,

protocols, and advice; Y. Sato (Carl Zeiss, Japan) for help in 3D reconstruction; and H.

Bito and F. Miyamasu for useful comments on the manuscript.

Author contributions

Conceptualization, S.K., T.O., K.K.-M., and M.M.; Methodology, S.K., M.Ta., M.Ts.,

and T.N.; Software, M.Ta.; Formal analysis, S.K. and M.Ta.; Investigation, S.K., T.O.,

K.K.-M., M.Ts., T.N., and M.M.; Writing–Original Draft, S.K. and M.M.; Writing–

Review & Editing, S.K., M.Ta., T.O., K.K.-M., and M.M.; Visualization, S.K. and

M.Ta.; Supervision, M.M.; Funding acquisition, S.K. and M.M.

Declaration of interests

The authors declare no competing interests.

References

Aizawa, H., Wakatsuki, S., Ishii, A., Moriyama, K., Sasaki, Y., Ohashi, K., Sekine-

Aizawa, Y., Sehara-Fujisawa, A., Mizuno, K., Goshima, Y., et al. 2001.

Phosphorylation of cofilin by LIM-kinase is necessary for semaphorin 3A-induced

growth cone collapse. *Nat. Neurosci*. 4:367-373. https://doi.org/10.1038/86011.

Bielinska, M., Narita, N., and Wilson, D.B. 1999. Distinct roles for visceral endoderm

during embryonic mouse development. *Int. J. Dev. Biol*. 43:183-205.

Biswas, A., Alex, A., and Sinha, B. 2017. Mapping Cell Membrane Fluctuations

Reveals Their Active Regulation and Transient Heterogeneities. *Biophys. J.*

113:1768-1781. https://doi.org/10.1016/j.bpj.2017.08.041.

Bond, L. M., Tumbarello, D.A., Kendrick-Jones, J. and Buss, F. 2013. Small-molecule

inhibitors of myosin proteins. *Future Med. Chem.* 5:41-52.

https://doi.org/10.4155/fmc.12.185

Chen, C.H., Tsai, F.C., Wang, C.C., and Lee, C.H. 2009. Three-dimensional

characterization of active membrane waves on living cells. *Phys. Rev. Lett.* 103:

238101. https://doi.org/10.1103/PhysRevLett.103.238101.

Chernomordik, L.V., and Kozlov, M.M. 2008. Mechanics of membrane fusion. *Nat.*

Struct. Mol. Biol. 15:675-683. https://doi.org/10.1038/nsmb.1455.

Conti, M.A., Even-Ram, S., Liu, C., Yamada, K.M., and Adelstein, R.S. 2004. Defects

in cell adhesion and the visceral endoderm following ablation of nonmuscle myosin

heavy chain II-A in mice. *J. Biol. Chem.* 279:41263-41266.

https://doi.org/10.1074/jbc.C400352200.

Ebrahim, S., Chen, D., Weiss, M., Malec, L., Ng, Y., Rebustini, I., Krystofiak, E., Hu,

L., Liu, J., Masedunskas, A., et al. 2019. Dynamic polyhedral actomyosin lattices

remodel micron-scale curved membranes during exocytosis in live mice. *Nat. Cell*

Biol. 21:933-939. https://doi.org/10.1038/s41556-019-0365-7.

Eitzen, G. 2003. Actin remodeling to facilitate membrane fusion. *Biochim. Biophys.*

Acta 1641:175-181. https://doi.org/10.1016/s0167-4889(03)00087-9.

Evans, E., and Rawicz, W. 1990. Entropy-driven tension and bending elasticity in

condensed-fluid membranes. *Phys. Rev. Lett.* 64:2094-2097.

https://doi.org/10.1103/PhysRevLett.64.2094.

Faizi, H. A., Frey, S.L., Steinkuhler, J., Dimova, R. and Vlahovska, P.M. 2019. Bending

rigidity of charged lipid bilayer membranes. *Soft Matter* 15:6006-6013.

https://doi.org/10.1039/c9sm00772e.

Farsad, K., and De Camilli, P. 2003. Mechanisms of membrane deformation. *Curr.*

Opin. Cell Biol. 15:372-381. https://doi.org/10.1016/s0955-0674(03)00073-5.

Goley, E.D., and Welch, M.D. 2006. The ARP2/3 complex: an actin nucleator comes of

age. *Nat. Rev. Mol. Cell Biol*. 7:713-726. https://doi.org/10.1038/nrm2026.

Gomes, E.R., Jani, S. and Gundersen, G.G. 2005. Nuclear movement regulated by

Cdc42, MRCK, myosin, and actin flow establishes MTOC polarization in migrating

cells. *Cell* 121:451-463. https://doi.org/10.1016/j.cell.2005.02.022

Gruenberg, J., Griffiths, G., and Howell, K.E. 1989. Characterization of the early

endosome and putative endocytic carrier vesicles in vivo and with an assay of

vesicle fusion in vitro. *J. Cell Biol*. 108:1301-1316.

https://doi.org/10.1083/jcb.108.4.1301.

Hammer III, J.A. and Sellers, J.R. 2011. Walking to work: roles for class V myosins as

cargo transporters. *Nat. Rev. Mol. Cell Biol*. 13:13-26.

https://doi.org/10.1038/nrm3248.

Harrington, A.W., St Hillaire, C., Zweifel, L.S., Glebova, N.O., Philippidou, P.,

Halegoua, S., and Ginty, D.D. 2011. Recruitment of actin modifiers to TrkA

endosomes governs retrograde NGF signaling and survival. *Cell* 146:421-434.

https://doi.org/10.1016/j.cell.2011.07.008.

Hartman, M.A. and Spudich, J.A. 2012. The myosin superfamily at a glance. *J. Cell Sci.*

125:1627-1632. https://doi.org/10.1242/jcs.094300

Helfrich, W. 1973. Elastic properties of lipid bilayers: theory and possible experiments.

Z. Naturforsch. C 28:693-703. https://doi.org/10.1515/znc-1973-11-1209.

Jollie, W.P. 1990. Development, morphology, and function of the yolk-sac placenta of

laboratory rodents. *Teratology* 41:361-381. https://doi.org/10.1002/tera.1420410403.

Kawamura, N., Sun-Wada, G.H., Aoyama, M., Harada, A., Takasuga, S., Sasaki, T.,

and Wada, Y. 2012. Delivery of endosomes to lysosomes via microautophagy in the

visceral endoderm of mouse embryos. *Nat. Commun.* 3:1071.

https://doi.org/10.1038/ncomms2069.

Koike, S., Keino-Masu, K., Ohto, T., Sugiyama, F., Takahashi, S., and Masu, M. 2009.

Autotaxin/lysophospholipase D-mediated lysophosphatidic acid signaling is

required to form distinctive large lysosomes in the visceral endoderm cells of the

mouse yolk sac. *J. Biol. Chem.* 284:33561-33570.

https://doi.org/10.1074/jbc.M109.012716.

Liebl, D., and Griffiths, G. 2009. Transient assembly of F-actin by phagosomes delays

phagosome fusion with lysosomes in cargo-overloaded macrophages. *J. Cell Sci.*

122:2935-2945. https://doi.org/10.1242/jcs.048355.

Lipowsky, R. 1991. The conformation of membranes. *Nature* 349:475-481.

https://doi.org/10.1038/349475a0.

Liu, D., Bei, D., Parmar, H., and Matus, A. 2000. Activity-regulated, cytoskeleton-

associated protein (Arc) is essential for visceral endoderm organization during early

embryogenesis. *Mech. Dev.* 92:207-215. https://doi.org/10.1016/S0925-

4773(99)00340-8.

789 Liu, K., Jian, Y., Sun, X., Yang, C., Gao, Z., Zhang, Z., Liu, X., Li, Y., Xu, J., Jing, Y.,

790  et al. 2016. Negative regulation of phosphatidylinositol 3-phosphate levels in early-

791  to-late endosome conversion. *J. Cell Biol*. 212:181-198.

792  https://doi.org/10.1083/jcb.201506081.

793 Lu, T., and Guo, H.X. 2019. How the Membranes Fuse: From Spontaneous to Induced.

794  *Adv. Theor. Simul.* 2:1900032. https://doi.org/10.1002/adts.201900032.

795 Ma, W., Chang, J., Tong, J., Ho, U., Yau, B., Kebede, M.A., and Thorn, P. 2020.

796  Arp2/3 nucleates F-actin coating of fusing insulin granules in pancreatic beta cells to

797  control insulin secretion. *J. Cell Sci.* 133:jcs236794

798  https://doi.org/10.1242/jcs.236794.

799 Maekawa, M., Ishizaki, T., Boku, S., Watanabe, N., Fujita, A., Iwamatsu, A., Obinata,

800  T., Ohashi, K., Mizuno, K., and Narumiya, S. 1999. Signaling from rho to the actin

801  cytoskeleton through protein kinases ROCK and LIM-kinase. *Science* 285:895-898.

802  https://doi.org/10.1126/science.285.5429.895.

803 Masedunskas, A., Sramkova, M., Parente, L., Sales, K.U., Amornphimoltham, P.,

804  Bugge, T.H., and Weigert, R. 2011. Role for the actomyosin complex in regulated

805  exocytosis revealed by intravital microscopy. *Proc. Natl. Acad. Sci. USA*

806  108:13552-13557. https://doi.org/10.1073/pnas.1016778108.

807 Miklavc, P., Ehinger, K., Sultan, A., Felder, T., Paul, P., Gottschalk, K.E. and Frick, M.

808  2015. Actin depolymerisation and crosslinking join forces with myosin II to contract

809  actin coats on fused secretory vesicles. *J. Cell Sci.* 128:1193-203.

810  https://doi.org/10.1242/jcs.165571.

Morel, E., Parton, R.G., and Gruenberg, J. 2009. Annexin A2-Dependent

Polymerization of Actin Mediates Endosome Biogenesis. *Dev. Cell* 16:445-457.

https://doi.org/10.1016/j.devcel.2009.01.007.

Morris, K., Ihnatovych, I., Ionetz, E., Reed, J., Braundmeier, A., and Strakova, Z. 2011.

Cofilin and Slingshot Localization in the Epithelium of Uterine Endometrium

Changes During the Menstrual Cycle and in Endometriosis. *Reprod. Sci.* 18:1014-

1024. https://doi.org/10.1177/1933719111401663.

Nicot, A.S., Fares, H., Payrastre, B., Chisholm, A.D., Labouesse, M., and Laporte, J.

2006. The phosphoinositide kinase PIKfyve/Fab1p regulates terminal lysosome

maturation in Caenorhabditis elegans. *Mol. Biol. Cell* 17:3062-3074.

https://doi.org/10.1091/mbc.e05-12-1120.

Richardson, S.J., Southwell, B.R., and Jaworowski, A. 2000. The rat visceral yolk sac

internalizes maternal transferrin and secretes hydrolyzed products towards the fetus.

Comp. Biochem. Phys. B 125:29-36. https://doi.org/10.1016/S0305-0491(99)00150-

9.

Ridolfi, A., Caselli, L., Baldoni, M., Montis, C., Mercuri, F., Berti, D., Valle, F. and

Brucale, M. 2021. Stiffness of Fluid and Gel Phase Lipid Nanovesicles: Weighting

the Contributions of Membrane Bending Modulus and Luminal Pressurization.

Langmuir 37:12027-12037. https://doi.org/10.1021/acs.langmuir.1c01660.

Roberts, R.L., Barbieri, M.A., Pryse, K.M., Chua, M., and Stahl, P.D. 1999. Endosome

fusion in living cells overexpressing GFP-rab5. *J. Cell Sci.* 112:3667-3675.

https://doi.org/10.1242/jcs.112.21.3667.

Salbreux, G., Charras, G., and Paluch, E. 2012. Actin cortex mechanics and cellular

morphogenesis. *Trends Cell Biol*. 22:536-545.

https://doi.org/10.1016/j.tcb.2012.07.001.

Scott, C.C., and Gruenberg, J. 2011. Ion flux and the function of endosomes and

lysosomes: pH is just the start. *Bioessays* 33:103-110.

https://doi.org/10.1002/bies.201000108.

Simson, R., Wallraff, E., Faix, J., Niewohner, J., Gerisch, G. and Sackmann, E. 1998.

Membrane bending modulus and adhesion energy of wild-type and mutant cells of

Dictyostelium lacking talin or cortexillins. *Biophys. J*. 74:514-22.

https://doi.org/10.1016/S0006-3495(98)77808-7.

Svitkina, T.M. 2018. Ultrastructure of the actin cytoskeleton. *Currr. Opin. Cell Biol*.

54:1-8. https://doi.org/10.1016/j.ceb.2018.02.007.

Taunton, J., Rowning, B.A., Coughlin, M.L., Wu, M., Moon, R.T., Mitchison, T.J., and

Larabell, C.A. 2000. Actin-dependent propulsion of endosomes and lysosomes by

recruitment of N-WASP. *J. Cell Biol*. 148:519-530.

https://doi.org/10.1083/jcb.148.3.519.

Titus, M.A. 2018. Myosin-driven intracellular transport. *Cold Spring Harb. Perspect.*

Biol. 10:a021972. https://doi.org/10.1101/cshperspect.a021972

Tran, D.T., Masedunskas, A., Weigert, R., and Ten Hagen, K.G. 2015. Arp2/3-mediated

F-actin formation controls regulated exocytosis in vivo. *Nat. Commun*. 6:10098

https://doi.org/10.1038/ncomms10098.

Uenishi, E., Shibasaki, T., Takahashi, H., Seki, C., Hamaguchi, H., Yasuda, T., Tatebe,

M., Oiso, Y., Takenawa, T., and Seino, S. 2013. Actin Dynamics Regulated by the

Balance of Neuronal Wiskott-Aldrich Syndrome Protein (N-WASP) and Cofilin

857 Activities Determines the Biphasic Response of Glucose-induced Insulin Secretion.

858 *J. Biol. Chem.* 288:25851-25864. https://doi.org/10.1074/jbc.M113.464420.

859 Vitriol, E.A., Wise, A.L., Berginski, M.E., Bamburg, J.R., and Zheng, J.Q. 2013.

860 Instantaneous inactivation of cofilin reveals its function of F-actin disassembly in

861 lamellipodia. *Mol. Biol. Cell* 24:2238-2247. https://doi.org/10.1091/mbc.E13-03-

862 0156.

863 Wang, L., Seeley, E.S., Wickner, W., and Merz, A.J. 2002. Vacuole fusion at a ring of

864 vertex docking sites leaves membrane fragments within the organelle. *Cell* 108:357-

865 369. https://doi.org/10.1016/S0092-8674(02)00632-3.

866 Welch, M.D., and Way, M. 2013. Arp2/3-Mediated Actin-Based Motility: A Tail of

867 Pathogen Abuse. *Cell Host Microbe* 14:242-255.

868 https://doi.org/10.1016/j.chom.2013.08.011.

869 Zhao, W.D., Hamid, E., Shin, W., Wen, P.J., Krystofiak, E.S., Villarreal, S.A., Chiang,

870 H.C., Kachar, B., and Wu, L.G. 2016. Hemi-fused structure mediates and controls

871 fusion and fission in live cells. *Nature* 534:548-552.

872 https://doi.org/10.1038/nature18598.

Figure Legends

Figure 1. Endocytic vesicles in VE cells. (**A**) Schematic diagram of the imaging setup.

An E8.5 whole mouse embryo was observed with a laser confocal microscope. (**B**)

Confocal images of VE cells stained with FM 1-43, LysoTracker Red, or anti-EEA1

antibody, or electroporated with EGFP-Rab7. The yellow dotted lines indicate the cell

boundaries. The yellow arrows show some vesicles that were positive for LysoTracker

Red in zone 2. (**C**) For descriptive purposes, the supranuclear portion of VE cells is

divided into 3 zones. Zone 1, zone 2, and zone 3 contain early endosomes (EE), late

endosomes (LE), and lysosomes, respectively. (**D**) Confocal images of VE cells labeled

with Alexa 488-transferrin. Images taken 1, 10, 20, and 40 min after pulse labeling are

shown. The yellow dotted lines indicate the cell boundaries. (**E**) Time-course analysis of

the Alexa 488-transferrin-containing vesicles in VE cells. n = 4. The scale bars indicate

4 μm (B) and 10 μm (D).

Figure 2. Homotypic fusion between late endosomes. (**A**) Time-lapse imaging of

endocytic vesicles in VE cells. After 5 min of pulse-labeling with Alexa 488-transferrin,

homotypic fusion of late endosomes (arrows) was frequently observed in zone 2. In the

upper panel, time 0 indicates the start of time-lapse imaging. In the lower panel, time 0

refers to the start of the fusion of the endosomes indicated by arrows. (**B**) By labeling of

VE cells with FM1-43, the fusion process of the cell membranes in zone 2 was observed.

Time 0 refers to the start of the fusion of the endosomes, indicated by arrows. (**C**)

Histograms showing the size distribution of total late endosomes (black line, n = 265)

and the late endosomes that underwent homotypic fusion (gray line, n = 37). No

significant difference was observed between the 2 groups (Mann-Whitney U test). (**D**)

Correlation between the size of the late endosomes that underwent homotypic fusion

and the time required for completion of fusion (i.e., the time required from membrane

fusion to formation of a single round vesicle). The sizes of the larger of the fused

vesicles are plotted. Thirty-seven fusion events were measured. The scale bars indicate

5 μm (A, top) and 1.5 μm (A, bottom; B).

Figure 3. Heterotypic fusion between late endosomes and lysosomes. (**A**) Time-lapse

imaging of VE cells. After 15 min of pulse-labeling with Alexa 488-transferrin,

heterotypic fusion of late endosomes was observed: they shrank gradually and

disappeared from the focal plane (arrows). In the upper panel, time 0 indicates the start

of time-lapse imaging. In the lower panel, time 0 refers to the start of the fusion of the

endosomes indicated by an arrow. (**B**) Time-lapse imaging of late endosomes in VE

cells. Late endosomes and lysosomes were labeled with Alexa 488-transferrin (Tfn,

green) and rhodamine-dextran (Dex, red), respectively. The bottom figures show optical

sectioning through the dotted line in the upper figure. Time 0 indicates the start of time-

lapse imaging. (**C**, **D**) Time course of the fluorescence intensity. In a late endosome

undergoing fusion (ROI 1 in [B-C]), the fluorescence intensity of transferrin (green)

slightly increased and then quickly decreased after membrane fusion. The fluorescence

signal of dextran (red) in an underlying lysosome leaked into the fusing late endosome.

The white box in (C) indicates the area shown in (B). The gray line in (D) indicates the

diameter of the transferrin-positive late endosomes. The blue dotted lines in (D) indicate

the moment the fusion pore was formed in ROI 1. ROI 2 is a negative control in a

lysosome that did not undergo fusion. (**E**) Electron microscopic image showing pore

formation between a late endosome (LE) and a lysosome (Ly). The inset shows a

magnified picture of the boxed region. (**F**) Correlation between the size of late

endosomes that underwent heterotypic fusion and the time required for completion of

fusion (i.e., the time from the start of shrinkage to disappearance). Twenty-five fusion

events were measured. (**G**) Histograms showing the size distribution of late endosomes

at 5 min (gray line, n = 256) and 15 min (red line, n = 242) after labeling with Alexa

488-transferrin, as well as the late endosomes that underwent heterotypic fusion at 15

min (blue line, n = 34). The average size of the vesicles was larger at 15 min than at 5

min (**$P<0.01$, Mann-Whitney U test). However, the size distribution of the late

endosomes that underwent heterotypic fusion at 15 min did not differ from that of the

total late endosomes at 15 min. The scale bars indicate 5 μm (A, top), 1.5 μm (A,

bottom), 2 μm (B), 3 μm (C), and 1 μm (E).

Figure 4. Mathematical modeling analysis of membrane deformation after connection

of 2 vesicles. (**A**) Definition of shape parameters. (**B**) Free-energy landscape for small

vesicles. The gray arrow (D) indicates a typical orbit of membrane deformation from the

initial condition $(w, a) = (0.0, 0.3)$. (**C**) Free-energy landscape for large vesicles. The

solid gray arrow (E) indicates the typical orbit of membrane deformation, whereas the

dashed gray arrow (D') indicates the typical orbit of membrane deformation under

fluctuated conditions. The inset is a blowup of the region $w = 0 \sim 0.15$, $a = 0.25 \sim 0.35$.

(**D**, **E**) Time series of shape variation on the typical orbits indicated in (B) and (C),

respectively. Note that the exact time scale cannot be estimated in this analysis. (**F**)

Probability of the explosive fusions in the F^{large} landscape (the orbit of the dashed gray

arrow in [C]) for various vesicle sizes, osmotic pressure differences, and fluctuation

energies, estimated by Monte-Carlo simulations. $\Gamma = 2\pi\kappa \sqrt{A}/R_0$ with the bending

modulus $\kappa = 10[k_B T]$ and the minimum bending radius $R_0 = 100[nm]$. The green

shaded region indicates the condition: $\Delta p \simeq 2.5 \sim 25[Pa]$, $V = 4A\sqrt{A}/3\sqrt{\pi}$ with

$A = 2 \sim 25[\mu m^2]$ and the fluctuation energy$= 1[k_B T]$ (thermal fluctuation at room

temperature). The green shaded region corresponds to late-endosome-sized vesicles

$(2.0 \sim 25[\mu m^2])$ under estimated osmotic pressure difference $(\Delta p \simeq 2.5 \sim 25[Pa])$ and

the thermal noise at room temperature. The blue-shaded region indicates the vesicles

under larger fluctuations.

Figure 5. Actin filaments are associated with late endosomes in VE cells. (**A**–**C**)

Confocal images of E8.5 mouse embryos stained with Alexa 488-phalloidin and anti-

mouse IgG. Mouse IgG was incorporated into late endosomes and used as a marker for

endosomes. Punctate cytoplasmic staining of phalloidin (green) was associated with the

IgG-containing endocytic vesicles (red) in zone 2 (B–B'') and at lower frequencies with

the lysosomes in zone 3 (arrows, C–C''). (**D**) Optical sectioning of VE cells. The cross-

sections along the x-z axis of the area indicated by the dotted line in the upper panel (x-

y axis) are shown. The dotted line in the IgG panel represents the apical region of zone

2. Actin filaments, extending from the apical surface into the cytoplasm (white

arrowheads), were closely associated with the endocytic vesicles (red). (**E**, **F**) Three-

dimensional surface rendering of actin filaments and a late endosome. The side view (E)

and basal view (F) are shown. (**G**) Alexa 488-phalloidin staining of mRFP-fascin-1-

expressing VE cells. Actin punctates were highly colocalized with fascin-1, whereas

fine actin filaments extending radially in the x-y plane from endosomes were negative

for mRFP-fascin-1 (white arrow heads). The fluorescence intensity plot along the white

arrow in the middle panel shows colocalization of fascin-1 and phalloidin in the large

spot. (**H**) Deconvolution images of late endosomes and actin. Maximum intensity

projection images are shown. In addition to strong phalloidin-positive spots (white

arrow heads), which were observable by use of conventional confocal microscopy, actin

filaments extending from the endosomal surface (orange arrow heads) were visualized

in the x-y axis image. In the x-z axis image, a late endosome was surrounded by actin

filaments, which extended from the apical cell surface to the basal side in the cytoplasm.

The scale bars indicate 3 μm (A–D), 5 μm (G), and 1 μm (E–F, H).

Figure 6. Dynamic regulation of actin filaments surrounding late endosomes. (**A**)

Representative time-lapse images of EGFP-actin spots on a late endosome. VE cells

were electroporated with EGFP-actin at E7.5, and time-lapse analysis was performed on

the next day. The arrow indicates the fusion of actin spots. The arrowhead indicates the

appearance of a new actin spot. Time 0 refers to an arbitrary time point in time-lapse

imaging. (**B**) The turnover rate of actin was analyzed by use of a FRAP assay. The area

(4 μm x 4 μm) covering a single late endosome at the apical region of zone 2 of VE

cells expressing EGFP-actin was bleached, and the fluorescence recovery was observed

by use of time-lapse microscopy. The yellow boxes indicate the laser-irradiated area,

and the yellow arrowheads, actin spots recovered after photobleaching. The time laser

irradiation was ended is indicated as 0. (**C**, **D**) Graphs showing the fluorescence

recovery of EGFP-actin surrounding late endosomes at the apical region of zone 2 (light

blue in [C], n = 4) and the basal region of zone 2 (dark blue in [C], n = 3) or at the cell

surface of VE cells (D, n = 5). Fluorescence intensity was normalized to prebleaching

intensities. (**E**) Half-recovery time ($t_{1/2}$) of EGFP-actin fluorescence at the apical and

basal regions of zone 2 and at the cell surface. Data were analyzed by use of one-way

ANOVA with the Tukey multiple comparison test (*P<0.05; **P<0.01). (**F**) EGFP-

actin localization during vesicle movement. Endocytic vesicles (red) visualized by use

of rhodamine-dextran in zone 2 are shown. When a vesicle (arrow) moved in a certain

direction (upward in this figure), the actin patch (arrowhead) was observed at the rear

end of the movement direction. The dotted line indicates the initial position of the

vesicle. Time 0 refers to an arbitrary time point in time-lapse imaging. The scale bars

indicate 1 μm (A) and 3 μm (B, F).

1001

Figure 7. Actin dynamics regulate homotypic fusion in VE cells. (**A**) Confocal

microscopic images of endocytic vesicles in zone 2. E8.5 mouse embryos pretreated

with 0.1 μM cytochalasin D (CD) or 0.1 μM jasplakinolide (Jasp) were labeled with

Alexa 488-transferrin for 5 min. Time-lapse images were taken for 400 s from 5 min

after labeling. Arrows indicate homotypic fusion. Time 0 indicates the start of time-

lapse imaging. (**B**) Frequency of homotypic fusion calculated from the number of

fusions that occurred in 400 s. Data are represented as means ± SEMs of 4–9

independent experiments; individual data are shown by circular dots. Data were

analyzed by use of one-way ANOVA with the Tukey multiple comparison test

(**P<0.01; ***P<0.001). (**C**) Quantitation of types of homotypic fusion. Fusion types

are classified into normal explosive fusion, slow explosive fusion (more than 60 s), and

bridge fusion. Cytochalasin D treatment slowed down the fusion process in a dose-

dependent manner and induced bridge fusion at 1 μM. Independent experiment numbers

are shown on the top of the bars. (**D**) Quantitation of the time for completion of

homotypic fusion. Treatment with 0.1 μM or 1 μM cytochalasin D or with 20 μM

CK666, but not with 0.1 μM jasplakinolide, resulted in a significant increase in the

fusion time. Data are represented as means ± SEMs of 4–30 independent experiments;

individual data are shown by circular dots. Data were analyzed by use of one-way

ANOVA with the Tukey multiple comparison test (**P<0.01; ***P<0.001). (**E**)

Representative images of homotypic fusion of late endosomes in VE cells treated with 1

μM cytochalasin D. Endosomes were visualized by incubation with Alexa 488-

transferrin. The arrows indicate the vesicles undergoing homotypic fusion in the bridge

fusion mode. Time 0 refers to the start of fusion of the endosomes indicated by arrows.

(**F**) Time-lapse imaging of EGFP-actin localization during homotypic fusion of late

endosomes. Localization of EGFP-actin (green) and endocytic vesicles (red), visualized

by use of rhodamine-dextran, was observed. When a pair of endosomes (arrows) fused,

EGFP-actin was newly polymerized at the opposite positions of the docking site and its

fluorescence intensity was increased (arrowheads). Time 0 refers to the start of fusion of

the endosomes indicated by arrows. (**G**) Distribution of EGFP-actin during homotypic

fusion of late endosomes. The fluorescence intensities of EGFP-actin (dark gray, n = 5)

and FM 1-43 (light gray, n = 5) were measured during homotypic fusion. The

percentages of their signals in the proximal (docking site, shown as 1), lateral (shown as

2), and distal (shown as 3) regions at the predocking, docking, rounding, and postfusion

stages are shown. Data are represented as means ± SEMs (*P<0.05, unpaired t-test). (**H**)

The time course of changes in the fluorescence intensity of EGFP-actin around a pair of

late endosomes during homotypic fusion. Time 0 s indicates the moment the fusion pore

of two late endosomes was formed. The graph indicates the average of 5 independent

experiments. The scale bars indicate 10 µm (A), 2 µm (E), and 3 µm (F).

1040

Figure 8. Actin stabilization induces heterotypic fusion of late endosomes with

lysosomes. (**A**) Confocal microscopic images of endocytic vesicles in zone 2. E8.5

mouse embryos pretreated with 0.1 µM cytochalasin D (CD) or with 0.01 µM

jasplakinolide (Jasp) were labeled with Alexa 488-transferrin for 5 min. Time-lapse

images were taken for 400 s from 15 min after labeling. The arrows indicate heterotypic

fusion with lysosomes. Time 0 indicates the start of time-lapse imaging. (**B**) Frequency

of heterotypic fusion calculated from the number of fusions that occurred in 400 s from

5 min after labeling. Data are represented as means ± SEMs of 4–7 independent

experiments; individual data are shown by circular dots. Data were analyzed by use of

one-way ANOVA with the Tukey multiple comparison test (**$P<0.01$, ***$P<0.001$).

(**C**) Quantitation of heterotypic fusion frequency at 15 min after labeling of VE cells

pretreated with cytochalasin D (CD), jasplakinolide (Jasp), or CK666. Data are

represented as means ± SEMs of 4–8 independent experiments; individual data are

shown by circular dots. Data were analyzed by use of one-way ANOVA with the Tukey

multiple comparison test (*$P<0.05$, **$P<0.01$). (**D**) Effects of myosins on heterotypic

fusion. Frequency of heterotypic fusion was calculated at 15 min after labeling. Data are

represented as means ± SEMs of 5–8 independent experiments; individual data are

shown by circular dots. Data were analyzed by use of one-way ANOVA with the Tukey

multiple comparison test (*$P<0.05$, ***$P<0.001$). (**E**) Colocalization of EGFP-myosin

IIA or EGFP-myosin Va with phalloidin in zone 2 of VE cells. The insets show the

magnified images of the boxed area. Following electroporation of the EGFP-fused

myosin expression plasmids and 24-h culture, whole embryos were stained with

Alexa546-phalloidin. EGFP-myosin IIA signals overlapped with phalloidin-positive

spots on late endosomes, whereas most of the EGFP-myosin Va signals were juxtaposed

with phalloidin-positive spots on late endosomes (see the schemas and fluorescence

intensity plots). The scale bars indicate 10 μm (A), 3 μm (E), and 1 μm (E, insert).

1067

Figure 9. Transition from homotypic to heterotypic fusion. (**A**) Time course of

frequencies of homotypic (blue line) and heterotypic (red line) fusion. The graph

indicates the means of 6 independent experiments. (**B**) Distribution of cofilin in zone 2

of VE cells. After whole embryos were electroporated with YFP-cofilin and cultured for

1 d, they were stained with Alexa 546-phalloidin. The lower graph plotting the

fluorescence intensity along the white arrow in the upper image indicates colocalization

of actin filaments and cofilin. (**C**) Optical sectioning images (right) of VE cells along

the dotted line in the x-y axial panel (left). Embryos, which were electroporated with

YFP-cofilin and cultured for 1 d, were stained with Alexa 546-phalloidin. The actin

filaments, extending from the apical to the basal region, were highly colocalized with

cofilin. Note that the white arrows and blue arrowheads indicate an actin filament that

surrounds late endosomes and the cell membrane, respectively. (**D**) Quantification of

the fluorescence intensity of phalloidin and cofilin along the apical-basal axis. The

intensity of spots colocalized with phalloidin and cofilin around late endosomes

observed in the x-y images was quantified along the z-axis. Fluorescence intensity was

normalized to the intensities in the most apical plane (8 late endosomes from 3 VE

cells). Data were analyzed by use of one-way ANOVA with the Tukey multiple

comparison test and represented as means ± SEMs (**P<0.01, ***P<0.001). (**E**) FRAP

assay to evaluate the dynamics of actin at the apical region of zone 2 in VE cell

pretreated with the S3 peptide. After treatment with the S3 or RV peptide overnight, the

area (4 μm x 4 μm) covering a single late endosome in VE cells expressing EGFP-actin

was bleached and the fluorescence recovery was observed by time-lapse microscopy.

Three independent experiments were performed. (**F**, **G**) Frequency of homotypic (F)

and heterotypic fusion (G) in embryos pretreated with 15 μg/ml S3 or RV peptides.

Frequencies of homotypic and heterotypic fusions were calculated according to the

numbers of fusions in 400 s from 5 min (F) and 15 min (G) after labeling with Alexa

488-transferrin. Data are represented as means ± SEMs of 3–4 independent

experiments; individual data are shown by circular dots. Data were analyzed by use of

one-way ANOVA with the Tukey multiple comparison test (*P<0.05). The scale bars

indicate 8 μm (B) and 2 μm (C). n.s. indicates no significant difference.

1098

Figure 10. Summary of late endosome fusion in VE cells. In VE cells, late endosomes (LE) show two different types of fusion with different targets. In homotypic fusion, once a membrane pore is formed between two late endosomes, the pore quickly expands and a single fused vesicle is formed. In contrast, in heterotypic fusion, the pore does not expand; instead, the late endosome gradually shrinks and disappears over time as the result of a transfer of its vesicle content into lysosomes. Two different types of actin are associated with the late endosomal membrane in VE cells: the actin filaments that extend along the apical-to-basal axis and the filaments that are radially polymerized from the membrane. Cofilin, an actin-binding protein that severs and depolymerizes actin filaments, activates actin turnover. In the apical region, late endosomes receive squeezing forces via active actin modification by cofilin and Arp2/3, which leads to homotypic fusion in an explosive mode. In contrast, in the basal region, actin is more static and late endosomes receive sliding forces via myosins.

Figure 1–figure supplement 1. Vesicular transport of dextran to lysosomes. (**A**) Electron microscopic image showing the position of early endosomes, late endosomes (LE), and a lysosome (Ly) in VE cells. N indicates the nucleus. The scale bar indicates 2 μm. (**B**) Confocal microscopic images of rhodamine-dextran-labeled endocytic vesicles in the yolk sac VE cells of E8.5 mouse embryos. Images in zones 1 to 3, taken at 1, 10, 20, and 40 min after labeling, are shown. The scale bar indicates 10 μm.

Figure 4–figure supplement 1. Mathematical modeling of membrane deformation during membrane fusion. (**A**) Definition of shape parameters for calculating the bending

energy of a part of small vesicles. (**B**) Definition of shape parameters for calculating the

volume of a part of large vesicles. (**C**) Calculated bending energy of a part of small

vesicles using Euler-Lagrange equations (red points) and its fitting curve with a

hyperbolic function (blue line). (**D**, **E**) Definition of shape parameters for calculating the

bending energy of the constricted neck region of large vesicles. (D) is the entire

structure and (E) is the close-up of the neck region.

1128

Figure 6–figure supplement 1. EGFP-actin electroporation and colocalization of Arp3-

EGFP with actin filaments surrounding late endosomes. (**A**) VE cells electroporated

with EGFP-actin were incubated with 100 µg/ml rhodamine-dextran for 5 min. VE cells

expressing low levels of EGFP-actin (white arrowheads) endocytosed rhodamine-

dextran normally and showed normal morphology of endosomal vesicles, whereas VE

cells expressing high levels of EGFP-actin (white arrows) did not endocytose normally.

Thus, cells expressing low levels of EGFP-actin were selected for functional analysis.

(**B**) Arp3-EGFP and Alexa 546-phalloidin were highly colocalized in VE cells. After

electroporation of the Arp3-EGFP expression plasmid and 24-h culture, the whole

embryos were stained with Alexa 546-phalloidin. In the x-y axis images, phalloidin-

positive actin spots were partially positive for Arp3-EGFP. Phalloidin-positive and

Arp3-EGFP-negative spots are shown by white arrowheads. The intensity plot along the

border of a late endosome (indicated by a white circular arrow in the upper panel)

shows partial colocalization between actin and Arp3. (**C**) Localization of Arp3-EGFP

during vesicle movement in the x-y direction. When a vesicle (indicated by the white

arrowheads) moved in a certain direction (upward in this figure), an Arp3 patch

(indicated by the orange arrowheads) was observed at the rear end of the movement

direction. Time 0 refers to an arbitrary time point in time-lapse imaging. The scale bars

indicate 30 μm (A), 5 μm (B), and 1 μm (C).

1148

Figure 7–figure supplement 1. Distribution of actin filaments and homotypic fusion in

cytochalasin D-treated VE cells. (**A**) Images of late endosomes and actin filaments at apical or

basal regions in zone 2 of VE cells pretreated with 1 μM cytochalasin D for 5 min. E8.5

embryos were subsequently labeled with Alexa 594-transferrin and Alexa 488-phalloidin.

Images on the right side show the maximum intensity projection (MIP) images along the x-y

axis (upper) and x-z axis (bottom) of a single late endosome in the boxed area in the left side

figure. After cytochalasin D treatment, filamentous structures observed on the control late

endosomes (both filaments extending radially and in the apical-basal direction) decreased, and

instead large clusters (white arrowheads) were observed in the cytosol, which appeared to be

due to aggregation of actin filaments. (**B**) E8.5 mouse embryos pretreated with 0.1 μM

cytochalasin D (CD) for 5 min were labeled with Alexa 488-transferrin or FM 1-43 for 5 min.

The fluorescent signal was observed 5 min after labeling. Homotypic fusion of late

endosomes after cytochalasin D treatment was slow when compared with that of the control

nontreated cells. Time is given in s. Time 0 s indicates the moment the fusion pore of 2 late

endosomes was formed. The scale bars indicate 2 μm (A, left), 0.6 μm (A, right), and 3 μm

(B).

1165

Figure 7–figure supplement 2. Distribution of actin filaments in jasplakinolide-treated

VE cells. Images of late endosomes and actin of VE cells pretreated with 1 μM

jasplakinolide for 5 min. E8.5 embryos were subsequently labeled with Alexa594-

transferrin and Alexa488-phalloidin. Images on the right side show the maximum

intensity projection (MIP) images along the x-y axis (upper) and x-z axis (bottom) of a

single late endosome in the boxed area in the figure on the left. After jasplakinolide

treatment, the fluorescence intensity of the phalloidin-positive mesh in zone 1 was

significantly increased (see the top portion of the x-z-axis image). In addition, at the

apical region of zone 2, actin filaments extending radially from the endosomal surface

(orange arrowheads) were thicker when compared with those of the control cells (Figure

5H), and the fluorescence intensity of the filaments was increased. In contrast, actin

filaments extending along the apical-basal axis were not changed (shown by white

arrowheads). The scale bar indicates 2 μm (left side) and 0.5 μm (right side).

1179

Figure 7–figure supplement 3. Distribution of actin during homotypic fusion in

cytochalasin D-treated VE cells. Time-lapse imaging of EGFP-actin during homotypic

fusion of late endosomes in E8.5 mouse embryos pretreated with 1 μM cytochalasin D

for 5 min. Localization of EGFP-actin (green) and endocytic vesicles (red), visualized

by use of Alexa 594-transferrin, was observed. EGFP-actin spots appeared to be rigid

and static. When a pair of endosomes (arrows) fused, drastic changes in the distribution

and fluorescence intensity of EGFP-actin signals as had been observed in the control

VE cells were not seen. Time 0 refers to the start of fusion of the endosomes indicated

by arrows. The scale bar indicates 3 μm.

1189

Figure 7–figure supplement 4. Distribution of Arp3-EGFP during fusion. Time-lapse

imaging of Arp3-EGFP localization during homotypic fusion of late endosomes. After

expression of Arp3-EGFP in VE cells, localization of EGFP was observed. When a pair

of endosomes (white arrowheads) fused, Arp3 was highly polymerized at the opposite

position of the docking site and its fluorescence intensity was increased (orange

arrowheads). Time 0 indicates the moment when the fusion pore of 2 late endosomes

was formed. Time is given in s. The scale bars indicate 1 μm.

1197

Figure 8–figure supplement 1. Effects of myosin inhibitors on homotypic fusion in VE cells.

E8.5 mouse embryos pretreated with 10 μM pentachloropseudilin (PClP), 10 μM blebbistatin,

10 μM MyoVin-1, or 10 μM 2,4,6-triiodophenol (TIP) for 5 min were labeled with Alexa

488-transferrin for 5 min. Frequency of homotypic fusion was calculated at 5 min after

labeling. Data are represented as means ± SEMs of 4–9 independent experiments; individual

data are shown by circular dots. Data were analyzed by use of one-way ANOVA with the

Tukey multiple comparison test (***P<0.001).

1205

Figure 9–figure supplement 1. Distribution of actin filaments in S3 peptide-treated VE

cells. Images of late endosomes and actin of VE cells pretreated with the S3 peptide (15

μg/ml) for 24 h and subsequently labeled with Alexa 594-transferrin and Alexa 488-

phalloidin. Images on the right side show the maximum intensity projection (MIP)

images along the x-y axis (upper) and x-z axis (bottom) of a single late endosome in the

boxed area in the figure on the left. Actin filaments extending radially from the

endosomal surface (orange arrowheads) were shorter and the fluorescence intensity was

decreased by S3 peptide treatment, whereas actin filaments extending from the apical

cell surface to the basal side on the surface of late endosomes were not changed (white

arrowheads). The scale bars indicate 2 μm (left) and 0.6 μm (right).

1216

A E8.5 mouse embryo

objective lens

C

EE — zone 1
LE — zone 2
Lysosome — zone 3

Nucleus

B

	FM 1-43	EEA1	EGFP-Rab7	LysoTracker
zone 1				
zone 2				
zone 3				

D

	1 min	10 min	20 min	40 min
zone 1				
zone 2				
zone 3				

E

zone 1
small vesicles
(<2 μm²)

zone 2
medium vesicles
(2-25 μm²)

zone 3
large vesicles
(>25 μm²)

Vesicle number vs Time (min)

C Fused (Ave. 7.3±6.6 μm²), Total (Ave. 5.7±6.4 μm²), n.s.

D r=0.60

A

$A = a_1 + a_2 \qquad w = W\sqrt{\dfrac{4\pi}{A}} \qquad a = \dfrac{a_1}{A}$

Actin

Arp2/3

Myosin V

Cofilin

Homotypic fusion
(Explosive)

Heterotypic fusion
(Bridge)

LE

LE

LE

LE

Lysosome

Nucleus

VE cell