Fluvoxamine maleate ameliorates Alzheimer disease pathology by mitigating amyloid-beta load and neuroinflammation in 5XFAD mice

Sukhleen Kaur[1, 2], Kuhu Sharma[1, 2], Ankita Sharma[1], Kamalpreet Kaur Sandha[1, 2], Syed Mudassir Ali[1], Riyaz Ahmed[3], P. Ramajayan[1], Parvinder Pal Singh[2, 3], Zabeer Ahmed[1, 2], Ajay Kumar[1, 2]

[1]Pharmacology Division, CSIR-Indian Institute of Integrative Medicine, Jammu-180016, India.

[2]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad-201002, India.

[3]Natural Products and Medicinal Chemistry Division, CSIR-Indian Institute of Integrative Medicine, Jammu-180016, India

* Correspondence

Ajay Kumar

Email: ajaykumar@iiim.res.in

ajaykumar@iiim.ac.in

Abstract

Background

Alzheimer pathology (AD) is accompanied by the deposition of amyloid beta (Aβ) and chronic neuroinflammation, where NLRP3 inflammasome is particularly involved. In this study, we found that the OCD drug fluvoxamine maleate (FXN) can potently ameliorate AD pathology in 5XFAD mice by autophagy-mediated clearance of Aβ and inhibition of NLRP3 inflammasome.

Methods

We used mice primary astrocytes to establish the mechanism of action of FXN against NLRP3 inflammasome by using various techniques like ELISA, Western blotting, confocal microscopy, Immunofluorescence, etc. The validation of the anti-AD activity of FXN was done in transgenic 5XFAD mice after two months of treatment followed by behavior analysis and studying inflammatory and autophagy proteins along with immunohistochemistry analysis for Aβ load in the hippocampi.

Results

Our data showed that FXN induces autophagy to inhibit NF-κB and NLRP3 inflammasome at a low concentration of 78 nM apart from directly inhibiting NLRP3 inflammasome in primary astrocytes. FXN activated the PRKAA2 pathway through CAMKK2 signaling, which led to the induction of autophagy in primary astrocytes. FXN inhibited the ATP-mediated NLRP3 inflammasome through autophagic degradation of NF-κB and thus caused the downregulation of pro-IL-1β and NLRP3. The anti-NLRP3

inflammasome effect of FXN was reversed when autophagy was inhibited either by

genetic knockdown of the PRKAA2 pathway or by bafilomycin A1.

Furthermore, FXN treatment led to improved AD pathology in 5XFAD mice, which

displayed a significant improvement in multiple behavior parameters like working

memory and neuromuscular coordination and they behaved more like wild-type animals.

We found that FXN improved behavior in 5XFAD mice by clearing the Aβ deposits from

the hippocampi along with a significant reduction in multiple inflammatory proteins,

including NF-κB, GFAP, IBA1, IL-1β, TNF-α, and IL-6 associated with NF-κB and

NLRP3 inflammasome in the brain. Moreover, these changes were accompanied by

increased expression of autophagic proteins.

Conclusion

Our data suggest that to ameliorate AD pathology, FXN simultaneously targets two key

pathological features of AD that is Aβ deposits and neuroinflammation. Being an

approved drug, FXN can be pushed as a potential drug candidate for human studies

against AD.

Key Words: Fluvoxamine maleate, Alzheimer disease, NLRP3 inflammasome;

Autophagy; Amyloid beta, Neuroinflammation, NF-κB, 5XFAD.

Background

Deposition of amyloid beta (Aβ) and neuroinflammation are essential pathological features of Alzheimer disease (AD). Aβ contributes to neuroinflammation through multiple mechanisms. It activates NLRP3 inflammasome either directly by acting as a DAMP for sensor NLRP3 and indirectly by activating TLR4 mediated induction of NF-κB pathway [1-6]. It can also activate NLRP3 inflammasome by inhibiting PRKAA2 and autophagy [7]. Inflammation, in turn, also promotes the Aβ pathology through various pathways. NLRP3-mediated activation of IL-1β inhibits the microglial cells from engulfing the Aβ plaques and thus contributes to its enhanced deposition [8-10]. ASC specks secreted by immune cells directly bind to the Aβ and augment the accumulation of Aβ aggregates [11]. NF-κB can also increase the activity of BACE1 to enhance the production of Aβ [12]. Therefore, it's a kind of vicious cycle which helps in the progression of AD. Many studies have shown that autophagy is a common link between Aβ deposition and activation of NLRP3 inflammasome [13, 14]. Autophagy, by virtue of its ability to recycle the cellular contents, can help clear protein aggregates and limit the activation of NLRP3 inflammasome by reducing the availability of proteins linked with inflammasome. Therefore, autophagy has emerged as one of the most important mechanisms that regulate NLRP3 inflammasome [15, 16]. Some recent studies have shown that induction of autophagy by pharmacological or non-pharmacological means can help mitigate NLRP3 inflammasome mediated or other types of inflammation [17-20]. Therefore, targeting NLRP3 inflammasome and autophagy together can be a good strategy, particularly when most of the monotargeted therapies have not shown any significant clinical advantage for treating AD.

In this study, as a part of our program related to the repurposing of drugs, we identified fluvoxamine (FXN) as a potent inhibitor of NLRP3 inflammasome and NF-κB pathway. We show here that the anti-inflammatory properties of FXN are regulated by PRKAA2-mediated autophagy. We attempted to utilize the ability of FXN to simultaneously target inflammation and autophagy in mitigating AD pathology in transgenic 5XFAD mice. We found that FXN can clear Aβ from the hippocampus of mice at a minimal dose 5 mg/kg along with a significant decline in neuroinflammation. These molecular changes in the brain helped improve working memory and overall brain health in 5XFAD mice. Our data projects FXN as a lead drug that should be explored for its possible development for the treatment of Alzheimer disease.

Methods

Chemicals and antibodies

Chemicals- RPMI-1640 (Sigma-Aldrich-R6504),Dulbecco's modified eagle medium (Sigma-Aldrich-D1152), Penicillin (Sigma-Aldrich-P3032), Streptomycin (Sigma-Aldrich-S6501), Sodium bicarbonate (Sigma-Aldrich-S5761), Fetal Bovine Serum (FBS) (GIBCO-10270106), Phosphate buffer saline (PBS) (Sigma-Aldrich-D5652), EDTA (Invitrogen-15575038), Lipopolysaccharide (LPS) (Sigma-Aldrich-L3129), Adenosine 5'-triphosphate (ATP) (Sigma-Aldrich-A6419), Acrylamide (MP Biomedical-193982), Glycine (MP Biomedical-194825),Triton X-100 (Sigma-Aldrich-T8787), Albumin Bovine Fraction V (MP Biomedical-160069), Phenylmethylsulfonyl fluoride (PMSF) (MP Biomedical-195381), Skimmed milk (Himedia-GRM1254), Strataclean resin (Agilent-

400714-61), Bafilomycin A1 (Sigma-Aldrich-B1793), Rapamycin (Sigma-Aldrich-R8781),DAPI (Sigma-Aldrich-D9542), glycerol (Sigma-Aldrich-G5516), Tween 20 (Sigma-Aldrich-P7949), HEPES (Sigma-Aldrich-H3375), Paraformaldehyde (Sigma-Aldrich-P6148), MTT (Sigma-Aldrich-M5655), Trizma (Sigma-Aldrich-T6066), SDS (Sigma-Aldrich-L3771), Uric acid sodium salt (Sigma-Aldrich-U2875), Suberic acid (Sigma-Aldrich-S1885), OPTI-MEM media (Gibco-11058-021), β-Amyloid (Anaspec-AS-60479).

Antibodies- NLRP3 (CST-15101S), ASC (Santa Cruz Biotechnology-SC-22514), CASP1 (Santa Cruz Biotechnology- SC-56036), anti-mIL-1β (R & D biotechnology-AF-401-NA), pCAMKK2 (CST-12818S), PRKAA2 (CST-2532S), pPRKAA2 (CST-2535S), pULK1 (CST-14202S), ULK1 (Santa Cruz Biotechnology-SC-33182), FIP200 (CST-12436S),BECN1 (Santa Cruz Biotechnology-SC-48341), ATG5 (Santa Cruz Biotechnology–SC-133158), ATG7 (Santa Cruz Biotechnology-SC-33211), ATG13 (CST- D4P1K), NF-κB (Santa Cruz Biotechnology-SC-372), pNF-κB (Santa Cruz Biotechnology-SC-136548), MTOR (CST-2972S), pMTOR (CST-5536S),LAMP1 (Santa Cruz Biotechnology-SC-20011), HRP-linked anti-goat IgG (Santa Cruz Biotechnology-SC-2354), siRNA PRKAA2 (CST-6620S), GFAP (CST-80788S), IBA1 (CST-17198S), HRP-linked anti-rabbit IgG (CST-7074S), HRP-linked anti-mouse IgG (CST-7076S), anti-mouse IgG Alexa flour 488 (CST-4408S), anti-mouse IgG Alexa flour 555 (CST-4409S), anti-rabbit IgG Alexa flour 488 (CST-4412S), anti-rabbit IgG Alexa flour 555 (CST-4413S), anti-ACTB (Sigma-Aldrich-A3854), ANTI-LC3B-II (Sigma-Aldrich-L7543), Anti-p62/SQSTM1 (Sigma-Aldrich-P0067).

Kits and reagents- PVDF Membrane (Millipore-ISEQ00010), Precision plus protein

markers (Bio-Rad-161-0375), ECL-kit (Millipore-WBKLS0500), Bradford reagent (Bio-

Rad-500-0006), FuGENE HD (Promega-E2313), mouse TNF-α ELISA kit (Invitrogen-

88-7324-88), mouse IL-6 ELISA kit (Invitrogen-88-7064-88), mouse IL- 1β ELISA kit

(Invitrogen-88-7013-88), mouse IL- 18 ELISA kit (Invitrogen-88-50618-88), Human Aβ42

ELISA kit (Invitrogen-KHB3441)

Synthesis of fluvoxamine maleate

The synthesis of fluvoxamine maleate was done by a well-established method (Figure

1A) (WO2014178064, US 6,433,225 B1). The scheme for synthesis of fluvoxamine

maleate (Scheme 1) and data related to its characterization is given in the

supplementary material file.

Cell culture

Primary astrocyte culture

The brain was isolated from four to five days old pups of C57BL/6 mice and cortical

hemisphere was treated with trypsin. Following centrifugation, the cells were passed

through cell strainer and cultured in low glucose DMEM media and supplemented with

10% FBS. After every 2 days, fresh media was added to the cells. After one week, the

shaking was carried out using rocker to separate out the astrocytes from the top layer of

the microglial cells. Thereafter, astrocytes were trypsinized and cultured for future

experiments under 95% humidity and 5% CO_2 at 37°C in an incubator chamber.

Cell line maintenance

The N9 microglial cells were cultured in RPMI-1640 media supplemented with 10%

Fetal bovine serum (FBS), penicillin (70 mg/L), streptomycin (100 mg/L) and $NaHCO_3$

(3.7 g/L), under constant supply of 5% CO_2 and 95% humidity at 37°C.

NLRP3 inflammasome activation in primary astrocytes

In order to activate the NLRP3 inflammasome in primary astrocytes, N9 microglial cells

were seeded in 24 well plates and allowed to grow for 24 h. The cells were then treated

with LPS (1 µg/ml) and incubated for next 24 h. The LPS activated microglial

conditioning media was collected and primary astrocytes were pretreated with the

conditioned medium for 24 h. Next day, primary astrocytes were primed with LPS (1

µg/ml) for 4 h and treated with different concentrations of FXN and standard MCC 950

(100 nM) for 1 h in serum free media, followed by ATP (5 mM) stimulation for 30 min.

Enzyme-linked Immunosorbent Assay (ELISA)

After activation of NLRP3 inflammasome and drug treatments (as mentioned in the

previous section), the supernatants were collected and analyzed to assess the levels of

IL-1β and IL-18 using ELISA following the manufacturer's instructions (Invitrogen). For

detection of LPS induced pro inflammatory cytokines, TNF-α and IL-6, primary

astrocytes were seeded in 24 well plate and pretreated with N9 conditioning media.

Then the astrocytes were incubated with LPS for 24 h and the levels of pro-

inflammatory cytokines were examined in the supernatants. The levels of cytokines

were normalized by dividing the values with total protein content present in the sample.

For *in vivo* studies, brain cortex was homogenized in RIPA buffer and centrifuged at

12000 rpm for 20 min at 4°C. The supernatants were collected and analyzed for IL-1β,

170   TNF-α and IL-6 cytokine levels. After completion of 2 months oral dosing, blood was

171   collected from retro-orbital plexus and the Aβ$_{42}$ levels in plasma of 5X FAD mice were

172   analyzed using ELISA following the manufacturer's guidelines (Invitrogen).

ASC Oligomerization

After drug treatments under NLRP3 inflammasome activation condition, primary

astrocytes were lysed in the cold buffer consisting of 1% cocktail, sodium orthovanadate

(1 mM), KCL (150 mM), PMSF (0.1 mM), HEPES-KOH (20 mM) and 1% of NP-40. The

lysed cells were centrifuged at 330g for 10 min at 4°C and the supernatant was

collected to analyze the protein expression by western blotting. The cell pellets were

washed with PBS and 500 μL of cold PBS was added to the cells. Suberic acid (2 mM)

was added and incubated with the pellets for 30 min at 37°C for cross-linking of ASC

proteins. Afterwards the pellets were centrifuged at 330g for 10 min at 4°C and 2X

Laemmli buffer was added to the cell pellets. Cell lysates were heated at 95°C for 5 min

and proteins were analyzed by western blotting.

Western blotting

For *in vitro* protein analysis by western blotting, primary astrocytes were seeded in 60

mm petri dishes and treated with FXN and standard MCC 950 under NLRP3

inflammasome activation condition. Cytokines secreted in the media were concentrated

using Strataclean resin according to the instructions provided by manufacturer and cells

were lysed with RIPA buffer [PMSF (2mM), Na3OV4 (0.5mM), NaF (50mM) and 1%

cocktail] and centrifuged at 12,000 rpm for 20 min at 4°C. For *in vivo* studies,

hippocampi from 5X FAD mice brain were homogenized and the lysed samples were

centrifuged. Supernatants were collected and the protein content was estimated using

Bradford's reagent. For in vitro studies 60 μg protein and for *in vivo* studies 20 μg

protein was loaded for separation using SDS PAGE and proteins were transferred to

PVDF membrane for 2 h at 4°C. The blots were blocked using 5% BSA or 5% skimmed

milk for 1 h at RT and incubated overnight with primary antibody at 4°C. Afterward blots

were incubated with HRP-conjugated secondary antibody for 2 h at RT.

Chemiluminescent HRP substrate (Millipore) was used for detection of protein bands

and visualized by Chemidoc system (Syngene G:BOX chemi XT 4).

Autophagy flux measurement

201 Autophagy flux was measured by treating primary astrocytes with different

202 concentrations of FXN in the presence and absence of bafilomycin A1 for 24 h. The

203 protein expression of LC3B-II and SQSTM1 were analyzed by western blotting.

Confocal microscopy

For confocal microscopic examination of the proteins, primary astrocytes were seeded

in 6-well plate over coverslip. After different treatments, cells were washed thrice with

PBS and fixed with 4% paraformaldehyde for 15 min. The cells were permeabilized with

0.2% Triton X-100 for 7 min and blocked using blocking buffer (2% BSA and 0.2% Triton

X-100) for 1 h. The cells were incubated overnight with primary antibody at 4°C. Next

day, the cells were incubated with secondary antibody (Alexa flour 555 or Alexa flour

488) for 1 h at RT and washed thrice with PBS. After counterstaining the nuclei with

DAPI, the coverslips were mounted using the mounting media and the images were

acquired using Yokogawa CQ1 Benchtop High-Content Analysis System at 40x or 60x.

Aβ clearance assay

Aβ$_{42}$-HiLyte flour488 peptide was prepared following the manufacturer's protocol (Anaspec Inc.). Primary astrocytes seeded on coverslips were treated with FXN or rapamycin. After 12 h, 2 µg/ml fluorochrome tagged (Hilyte flour488) Aβ$_{42}$ protein was added to cells for another 12 h. Bafilomycin A1 was given 3 h prior to experiment termination. Cells were washed with PBS and fixed with 4% PFA, afterwards cells were permeabilized with Triton-X 100, and the nuclei were counterstained with DAPI. Slides were prepared and images were taken and analyzed in the Yokogawa CQ1 Benchtop High-Content Analysis System at 40x or 60x.

Transfection of primary astrocytes with PRKAA2 *siRNA*

Primary astrocytes were grown in 6-well plate for confocal microscopic analysis and 60-mm dishes for western blotting. For transfection of PRKAA2, cells were incubated with OPTI-MEM media and PRKAA2 siRNA in FuGENE HD was added for 24 hours. Transfected primary astrocytes were incubated with microglial conditioning media for 24 h, followed by FXN (78 nM) treatment under NLRP3 inflammasome activation condition and the samples were analyzed by western blotting and confocal microscopy at 60x (Yokogawa CQ1 Benchtop High-Content Analysis System).

Drug Formulation

For in vivo studies, FXN was formulated in 5% of DMSO, 30% PEG400, 20% PEG200 and 45% distilled water. LPS and ATP were dissolved in PBS.

Animals and Ethical Clearance

Six months old 5XFAD transgenic mice were used in the study and randomly divided

into three groups (n = 5). C57BL6/J mice were used as Control wild type (WT). They

were housed under a 12-hour light/dark cycle in a temperature (65–75 °F; ˜18-23 °C)

and humidity-controlled (40–60 %) environment, supplied with free access to the food

and water (ad libitum). Prior to initiation of study all the animals were acclimatized for

one week under standard laboratory conditions, animals were drug naive with no prior

procedures performed. All testing were performed from 1 to 4 p.m. Mice were

randomized in groups based on their body weights for all behavioral assays and testing

were performed by an experimenter blinded to the treatment groups. Total study

duration was 2 months. All experiment protocols were approved by the Institutional

Animal Ethics Committee (IAEC) (IAEC approval no.- 321/82/2/2023), and followed the

Committee for Control and Supervision of Experiments on Animals (CCSEA; Ministry of

Environment and Forest, Government of India) guidelines for animal care.

Open-field test

The open field arena box (60 cm x 45 cm x 25 cm) made up of white colored non-

reactive plastic was used for all the assessments. The mice were acclimatized to the

testing room 30 minutes prior to experiment. Individual mice were placed in the center

of the open field arena and allowed to freely explore it for 5 minutes. The arena box was

cleaned with a 70% ethanol after every trial. The locomotor activity and exploratory

behavior of mice were recorded using a video camera connected to AnyMaze software.

Mice were assessed in the open field and the following parameters were recorded

automatically: Total distance travelled (cm), average speed (m/s), time spent in the

center area (sec), and time spent in the corners (sec).

Radial arm maze test

The eight-armed radial arm maze (UGO Basile) was used to assess spatial memory behavior. The mice were habituated to the maze for three days before the experiment. Each mouse was trained to reach to the baited arm having a butter cookie as food reward and flag as a visual cue. Training trial was of 3 days. On the memory retention trial day the mouse was placed on the end of one arm (entry arm) and allowed to navigate the baited arm. The movement of mice was tracked with the help of automated AnyMaze software connected with a video tracking camera.

Neuromuscular coordination test

The neuromuscular coordination of mice was recorded by rotarod. For this assessment mice were trained for three consecutive days with three trials per day. The maximum time allowed on the rotarod during the experiment was 300 seconds. The rotarod was initially set at a speed of 4 rpm at the start of the test and accelerated to 40 rpm over 300 seconds. The latency to fall from the rotarod was recorded for each mouse, and the mean latency to fall for each group was calculated.

Immunohistochemical analysis of 5XFAD mice brains

After the treatment completion, the hippocampi were isolated from the mice brains and the 10 μm thick tissue-sections were prepared. For IHC, the slides were deparaffinized and histochemical staining was carried out. The sections were incubated with primary antibody (overnight at 4°C under humid conditions after blocking using 2% BSA solution in TBS. For analysis of total Aβ, sections were incubated with anti-Aβ antibody (1-42) D3E10 followed by secondary antibody (Alexa fluor 488) and for localizing reactive

astrocytes, anti-GFAP was used followed by secondary antibody (Alexa fluor 555). The

alteration in autophagy levels were assessed by examination of LC3B levels in the

tissue sections of mice hippocampi. The sections were immuno-stained with LC3B

followed by secondary antibody (Alexa fluor 488). The nuclei were counterstained using

DAPI. The images were acquired and quantified using Confocal Quantitative Image

Cytometer at 40X (Yokogawa CQ1).

Statistical analysis

Statistical analyses were analyzed using Graph pad prism 9 software. The data shown

here are Mean±SD of three independent experiments. The response of independent

mice from the group was noted for animal experiments and the mean was calculated.

The statistical analysis of data was calculated using one-way ANOVA, followed by post-

hoc Bonferroni test. The p-value<0.05 was considered to be statistically significant with

values assigned as ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$ and ns= not

significant.

Results

FXN displayed a potent anti-NLRP3 inflammasome activity in primary astrocytes

NLRP3 inflammasome being an important player in neuroinflammation, we treated the

LPS-primed primary astrocytes with FXN before induction with ATP to analyze the anti-

NLRP3 inflammasome activity of FXN. The ELISA results showed that FXN could inhibit

NLRP3 inflammasome even at very low concentrations of nine nanomolar, which was

evidenced by the measurement of released IL-1β and IL-18 as inflammasome activation

products. FXN inhibited the release of IL-1β and IL-18 with IC50 values of 16 nM &

25.96 nM, respectively (Figure 1B and C). Based on the concentration-dependent

inhibitory activity of FXN, we chose 78 nM concentration for further detailed studies. We

also confirmed the inhibition release of cleaved IL-1β and CASP1 through western

blotting of the proteins collected from the supernatant of primary astrocytes treated with

FXN (Figure 1D). The analysis of whole cell lysates revealed reduced levels of pro-IL-1β

and NLRP3, while pro-CASP1 was unchanged (Figure 1D and Supplementary Figure

S1A). Further, the oligomerization of ASC is an important step in the activation of

NLRP3 inflammasome. Therefore, we analyzed the effect of FXN on ASC by two

methods, including confocal microscopy and western blotting. Both experiments showed

similar results with the potent inhibitory effect of FXN on ASC oligomerization (Figure 1E

and F and Supplementary Figures S1B and C). We also analyzed the co-localization of

CASP1 and NLRP3 in the presence of FXN. Data revealed that FXN and MCC950

(used as a standard) had a similar inhibitory effect on the co-localization of CASP1 and

NLRP3 (Figure 1G and Supplementary Figure S1D).

FXN inhibited the NLRP3 inflammasome through NF-κB pathway.

In the initial experiments, we treated the LPS-primed astrocytes with FXN to know

whether it could inhibit the assembly of the NLRP3 inflammasome complex. We also

wanted to know if it can inhibit the NLRP3 inflammasome at the transcriptional level by

inhibiting the NF-κB. Therefore, we treated the cells with FXN before priming them with

LPS. We found that FXN could significantly inhibit NLRP3 inflammasome at a

concentration of 78 nM contrary to nine nanomolar, observed during FXN treatment

post-priming. The ELISA measurement of IL-1β showed an IC50 value of 131.1 nM

(Figure 2A). We confirmed the inhibitory effect of FXN on NF-κB by direct and indirect

methods. The dwindling levels of NF-κB dependent proinflammatory cytokines TNF-α

and IL-6 after treatment with FXN at 78 nM indicated the inhibition of NF-κB in

astrocytes (Figure 2B & C). Direct evidence for inhibition of NF-κB pathway came from

western blot analysis of NF-κB in the cytosolic and nuclear fraction of astrocytes treated

with FXN. The data showed that the movement of NF-κB, p65 from the cytoplasm to the

nucleus is hindered in the cells treated with FXN, wherein the levels of NF-κB in the

cytoplasm were significantly higher than in the untreated control. On the contrary, its

level in the nucleus was significantly reduced (Figure 2D-F).

FXN induced the autophagy under the inflammatory conditions in primary

astrocytes

After confirming the inhibitory effect of FXN on NLRP3 inflammasome, we wanted to

know its inhibition mechanism. Therefore, we focused on autophagy as a major

mechanism regulating inflammasome activity. We found that FXN could induce

autophagy at a low concentration of 39 nM; however, its effect was enhanced with the

increasing concentration (Figure 3A and Supplementary Figure S2A). Further, to confirm

the completion of autophagy, we calculated the autophagy flux by using LC3B-II and

SQSTM1 expression after treatment of astrocytes with FXN (78 nM) in the presence

and absence of end-stage autophagy inhibitory bafilomycin A1. We found an autophagy

flux at all the tested time points through 24 h of astrocyte treatment with FXN (Figure 3B

and Supplementary Figure S2B). Induction of autophagy by FXN was also confirmed by

confocal microscopy analysis of co-localization of LC3B-II and LAMP1 in primary

astrocytes. The cells treated with FXN and rapamycin showed a significant amount of

co-localization of these proteins, while in the presence of bafilomycin A1, FXN-treated

cells did not show any observable co-localization of LC3B-II and LAMP1 (Figure 3C and

D). After confirming the induction of autophagy by FXN, we checked if it can induce

autophagy in primary astrocytes under inflammatory conditions where the cells have

been challenged with LPS and ATP. We found that the treatment of astrocytes with FXN

before ATP induced the expression of LC3B-II, while SQSTM1 was downregulated, thus

confirming the induction of autophagy (Figure 3E and Supplementary Figure S2C). We

further analyzed the expression of various proteins involved in the induction of

autophagy. We found that FXN induced autophagy through CAMKK2 mediated

induction of PRKAA2 pathway leading to activation of various proteins including

pPRKAA2 (Thr 172), pULK (Ser 317), ATG13, FIP 200, BECN1, ATG5 and ATG7 and

downregulation of pMTOR (Figure 3F and supplementary Figures S2C).

FXN inhibited NLRP3 inflammasome by inducing autophagy

For the confirmation of the involvement of autophagy in FXN-mediated inhibition of

NLRP3 inflammasome, we inhibited the autophagy through genetic and

pharmacological methods. We pretreated the primary astrocytes with *siPRKAA2* and

analyzed the expression of autophagic and NLRP3 inflammasome-related proteins. The

inhibition of PRKAA2 led to the reversal of the autophagic effect of FXN, which was

reflected in the form of a marked decrease in the expression of LC3B-II and BECN1,

while SQSTM1 expression was significantly increased (Figure 4A and Supplementary

Figure S3A). Similarly, the inhibition of NLRP3 inflammasome by FXN was also

reversed in the cells pretreated with *siPRKAA2*. The cleaved IL-1β levels were

significantly increased, whereas the reduction in expression of NLRP3 was also

reversed in the cells treated with FXN in the presence of *siPRKAA2* (Figure 4A and

Supplementary Figure S3A). We also confirmed these findings by using confocal

microscopy, where the co-localization of CASP1 and NLRP3 was restored to the level

present in cells primed with LPS and ATP when they were treated with FXN along with

knockdown of PRKAA2 by using siRNA (Figure 4B and Supplementary Figure S3B).

The knockdown of PRKAA2 was confirmed by using western blotting, where the

expression of PRKAA2 and pPRKAA2 was found to be significantly reduced (Figure

4C). We found that a similar kind of reversal of the inhibitory activity of FXN against

NLRP3 inflammasome could be achieved if the autophagy is inhibited by using the end-

stage pharmacological inhibitor bafilomycin A1. The cleavage of IL-1β was almost

completely inhibited, while the expression of NLRP3, NF-κB (p65) along with autophagy

markers LC3B-II, SQSTM1 was restored when the cells were treated with FXN in the

presence of bafilomycin A1 (Figure 4D and Supplementary Figure S3C)

FXN cleared Aβ_{42} in primary astrocytes by inducing autophagy

To confirm whether the autophagy induced by FXN can help in the clearance of Aβ_{42}, we

treated the cells for a total of 24 h with FXN, including 12 h treatment with hilyte fluor

488 tagged Aβ_{42}. The confocal analysis revealed that the cells treated with FXN and

rapamycin were almost completely devoid of green fluorescence indicating clearance of

Aβ_{42} in comparison to control cells. Further, 3 h treatment of cells with autophagy

inhibitor bafilomycin A1 prior to termination of the experiment completely repressed the

clearance of Aβ_{42}, as indicated by the same level of green fluorescence as that of control

cells (Figure 5A and C). The involvement of autophagy in the clearance of Aβ_{42} by FXN

was further confirmed when we inhibited the autophagy by using *siRNA* against

PRKAA2. The clearance of Aβ_{42} by FXN in the presence of *siPRKAA2* was similarly

stopped as we observed with bafilomycin. A1 (Figure 5B and D).

FXN improved working memory, exploratory behavior and neuromuscular coordination in 5XFAD mice

After confirming the mechanism of inhibition of NLRP3 inflammasome by FXN, we

wanted to know if its dual pharmacological activity can improve Alzheimer disease

pathology. Therefore, we treated the 6-month-old 5XFAD mice for two months and

analyzed multiple behavior parameters. In the radial arm maze test, the untreated

5XFAD mice displayed significant deterioration of neuronal functions in comparison to

wild-type control (Figure 6A). However, the groups treated with FXN (5 and 10 mg/kg)

took significantly less time and traveled a reduced distance in finding the baited arm and

spent more time in the baited arm apart from fewer entries into non-baited arms in

comparison to untreated 5XFAD mice, indicating improvement in the working memory

(Figure 6B-E). In the open field test, untreated 5XFAD mice spent significantly more

time in the center than in corners and showed little interest in exploring the surroundings

(Figure 6F). However, the mice treated with FXN covered a significantly higher distance

at greater speed and showed normal behavior by spending more time in the corners

than in the center (Figure 6G-K). We also observed a significant improvement in the

neuromuscular coordination in the mice treated with FXN, which showed delayed

latency to fall than the untreated 5XFAD mice (Figure 6L). In all the behavior

parameters studied, the mice treated with FXN behaved more like wild-type animals

than the untreated 5XFAD control group.

FXN reduced the Aβ_{42} in the hippocampus and plasma of 5XFAD mice

Improvement in neuronal health of 5XFAD mice after treatment with FXN led us to

investigate the underlying cause for such change. Therefore, through

immunohistochemistry (IHC), we checked the levels of $A\beta_{42}$ in the hippocampus after

two months of treatment with FXN. We found a significant decline in the levels of $A\beta_{42}$ in

the hippocampus of both groups of mice treated at 5 and 10 mg/kg, respectively

compared to the control group, as indicated by reduced green fluorescence (Figure 7A).

The average number of $A\beta_{42}$ plaques, skeletal length of $A\beta_{42}$ plaques, branch count and

circumference of $A\beta_{42}$ plaques were significantly reduced in FXN treated groups as

compared to 5XFAD control group (Figure 7B-E). $A\beta_{42}$ is strongly related to activation of

astrocytes, we therefore checked if reduced levels of $A\beta_{42}$ had any impact on the

reactive phenotype of astrocytes. The analysis of GFAP in the vicinity of $A\beta_{42}$ revealed

that as the levels of $A\beta_{42}$ were depleted after treatment with FXN, the expression of

GFAP reduced proportionately, as indicated by reduced red color fluorescence (Figure

7A and F). Further, the reduction of $A\beta_{42}$ levels in the brain also led to reduced

circulatory levels of $A\beta_{42}$ in the blood plasma of 5XFAD mice treated with FXN in

comparison to untreated control mice, as observed through ELISA (Figure 7G).

FXN ameliorated the AD pathology in 5XFAD mice through autophagy-mediated

reduction in amyloid beta levels and neuroinflammation

Based on the effect of FXN on in vitro and in vivo clearance of $A\beta_{42}$, we hypothesized

that autophagy induced by FXN is responsible for the reduced level of $A\beta_{42}$ in the

hippocampi of 5XFAD mice brains. Therefore, to test the hypothesis, we analyzed the

expression of LC3B-II in the hippocampi and found a highly significant increase in its

levels in the mice treated with FXN (Figure 8A). Further, analysis of expression of

various autophagic proteins including pCAMKK2 (Ser 511), pPRKAA2 (Thr 172),

pMTOR (Ser 2448), LC3-IIB, pULK1 (Ser 317), BECN1, and ATG5 revealed a

significant upregulation of these proteins in the hippocampi of mice treated with FXN in

comparison to untreated mice, while the effect of FXN at 10 mg/kg was more

pronounced as compared to the dose of 5 mg/kg (Figure 8B and Supplementary

Figures S4A).

The analysis of pro-inflammatory proteins in the hippocampi also supported the anti-

inflammatory effect of FXN. We found that treatment of mice with FXN significantly

reduced the levels of both NF-κB, p50 and p65 subunits confirming its effect on the

most important inflammatory pathway. Further analysis of NLRP3 inflammasome

proteins showed significant down-regulation of NLRP3, which significantly reduced ASC

oligomerization and cleaved CASP1 and IL-1β in the mice treated with FXN (Figure 8C

and D and Supplementary Figures S4B). The reduction of inflammation in the

hippocampi was indicated by reduced levels of microglial marker protein IBA1 (Figure

8C and Supplementary Figures S4B). We also observed a significant decline in the

levels of pro-inflammatory cytokines TNF-α, IL-6 and IL-1β, which were measured in the

cortex region of the brain by using ELISA (Figure 8E-G). These data emphasized a

potent effect of FXN on the pathology of AD in 5XFAD mice.

Discussion

Alzheimer disease progresses with a complex pathology. However, most cases of AD

are accompanied by the presence of $A\beta_{42}$ plaques and neuroinflammation where

NLRP3 inflammasome plays a significant role [2, 21-23]. There are multiple types of

damage associated molecular patterns (DAMPs) that can upregulate NLRP3

inflammasome in the brain, including Aβ, TAU NFTs, and ATP [24-27]. All these DAMPs

contribute to chronic inflammation and AD pathology. We initiated this study focusing on

finding NLRP3 inflammasome inhibitors that can be used to mitigate AD pathology. Our

preliminary studies found that fluvoxamine maleate (FXN) is a potent inhibitor of NLRP3

inflammasome with an inhibitory concentration in the nanomolar range in ATP-induced

primary astrocytes. US FDA had approved FXN for the treatment of obsessive-

compulsive disorder (OCD) in 1994. Its proposed mechanism against OCD is through

the inhibition of serotonin reuptake transporter (5-HT) [28-30]. However, our studies

revealed new targets of FXN that can be effectively exploited against AD. Additionally,

being an approved drug for CNS disease, FXN gave us an additional advantage to be

used for AD studies without being worried about its safety and crossing of the blood-

brain barrier (BBB). For detailed studies on the effect of FXN on neuroinflammation, we

continued working on primary astrocytes because of their emerging role in AD and other

neurodegenerative diseases. We found that FXN downregulated the expression of

NLRP3 and pro-IL1β along with reduced oligomerization of ASC leading to poor

assembly of NLRP3 inflammasome and thus cleavage of CASP1. These data indicated

that FXN works on both the steps of NLRP3 inflammasome activation to inhibit its

assembly. This assumption was confirmed when we observed a significant reduction in

the nuclear translocation of NF-κB, p65 and inhibition of pro-inflammatory cytokines

TNF-α and IL-6, which depend on it. These data emphasized that FXN has a wider

anti-inflammatory spectrum that targets multiple inflammation-related pathways. These

findings are important as inflammation is the key driver in neuronal cell death during AD

[31, 32].

After confirming the anti-inflammatory activity of FXN, we intended to know how the

upstream mechanisms of two important molecular pathways were inhibited by FXN.

Therefore, we explored the possible involvement of autophagy in the FXN mediated

downregulation of inflammation. Autophagy can regulate inflammation through multiple

ways including recycling of proteins, and organelles like mitochondria, clearance of

protein aggregates and other DAMPs, invading microorganisms and PAMPs associated

with them, etc. [33-35]. We found that FXN is a very potent inducer of autophagy, which

works in the same concentration range, which was effective against inflammation. We

further observed that it can also induce autophagy under the inflammatory conditions

generated by using LPS and ATP. The treatment of astrocyte with FXN under

inflammatory conditions showed a significant decline in the expression of NF-κB and

other proteins associated with activation of NLRP3 inflammasome. This was an

interesting observation. However, we wanted to be sure about the involvement of

autophagy in the anti-inflammatory effects of FXN. Therefore, we inhibited the

autophagy by both pharmacological and genetic means and found that all the anti-

inflammatory effects of FXN were reversed, which was evidenced by re-instatement of

expression level of inflammatory proteins to the pre FXN treatment levels.

Apart from being an important regulator of inflammation, the key physiological role of

autophagy is to remove protein aggregates [35]. The failure of autophagy to clear

proteins aggregates is strongly linked to number of pathological conditions including AD,

where it helps to maintain the physiological levels of Aβ along with other clearance

mechanisms [36]. We found that FXN could also help in the clearance of Aβ by inducing

autophagy in primary astrocytes. This was further confirmed when the clearance of Aβ

was completely reversed in the presence of *siPRKAA2*.

These findings encouraged us to validate the efficacy of FXN to ameliorate the AD

pathology in transgenic 5XFAD mice. Therefore, we treated the 5XFAD mice for two

months with FXN at two different doses. At the end of treatment, the analysis of different

behavior parameters revealed that the mice treated with both doses of FXN had

significantly improved brain health. The data from different parameters studied during

the radial arm maze test showed a marked improvement in the working memory of

mice. The hippocampus is the most important part of the brain that processes working

memory and is severely affected during AD, as was observed in untreated control mice.

Further, the exploratory behavior of FXN-treated mice was also significantly improved

and it appeared more like non-demented animals. They spent more time in corners,

traveled more distances at higher speeds, etc. The neuromuscular coordination was

also found to be similarly improved in mice treated with FXN. All these parameters

clearly indicated the improvement of brain health after treatment with FXN. These data

made us to explore molecular mechanisms behind this improvement in brain health,

particularly memory behavior. Therefore, we isolated the hippocampi of these mice and

analyzed the expression of various proteins related to autophagy and inflammation. We

found that the mice treated with FXN had a significantly increased expression of

proteins involved in autophagy induction including pCAMKK2 (Ser 511) and pPRKAA2

(Thr 172), and downregulation of pMTOR (Ser 2448). On the similar lines the

expression of autophagy initiation and execution proteins pULK1(Ser 317), BECN1,

LC3B-II, and ATG5 was found to be significantly increased. The most important impact

of autophagy induction by FXN in the hippocampus was the clearance of $A\beta_{42}$. We

found only traces of $A\beta_{42}$ in the hippocampus of FXN treated mice in comparison to

untreated control. Interestingly, reduced $A\beta_{42}$ levels were directly related to low-reactive

phenotype of astrocytes in the hippocampi as evidenced by reduced expression of

GFAP. Furthermore, the expression of microglial marker IBA1 was also found to be

significantly down regulated in mice treated with FXN. These data indicate that FXN

through autophagy not only cleared $A\beta_{42}$ but it also helped mitigation of overall

inflammatory microenvironment in the brain. These data were further supported by

down regulation of NF-κB and other proteins associated with the activation of NLRP3

inflammasome in the hippocampus. Moreover, these effects of the FXN were not

localized to hippocampi as the analysis of proinflammatory cytokines TNF-α, IL-6, and

IL-1β in the cortex region of the brain clearly established the strong effect of FXN

against neuroinflammation. These data directly correspond to the improvement of

memory behavior.

Conclusion

The analysis of whole data indicates that FXN exerts a potent autophagic effect through

activation of PRKAA2 pathway, which helps in the clearance of $A\beta_{42}$ and inhibition of

neuroinflammation. In AD, deposition of $A\beta_{42}$ and neuroinflammation together inflict

severe damage to neuronal tissue in the brain. Interestingly, FXN exerted these anti-

Alzheimer effects at the minimal dose of 5 mg/kg, which corresponds to approximately

25 mg/kg of human equivalent dose, which is a minimum prescribed dose of FXN for

OCD. Therefore, FXN through its ability to work against both these key pathological

hallmarks of AD at a minimal dose can raise a new hope and warrants its further

development against AD.

List of abbreviations

Aβ- Amyloid-beta

AD- Alzheimer's disease

PRKAA2- Protein kinase AMP-activated catalytic subunit alpha 2

ATG- Autophagy related

ANOVA- Analysis of variance

ATP- Adenosine triphosphate

ASC- Apoptosis-associated speck-like protein containing a Caspase recruitment domain

BECN1- Beclin 1

BSA- Bovine serum albumin

Baf. A1- Bafilomycin A1

CAMKK- Calcium/calmodulin-dependent protein kinase kinase

CASP1- Caspase-1

NF-κB- Nuclear factor kappa B subunit

DAMPs- Damage-associated molecular patterns

DAPI- 4,6-diaminido-2-phenylindole

DMEM- Dulbecco's modified eagle medium

DMSO- Dimethyl Sulfoxide

ECL- Enhanced chemiluminescence

EDTA- Ethylenediamine tetra acetic acid

ELISA- Enzyme-linked immunosorbent assay

FBS- Fetal bovine serum

FDA- Federal drug administration

Fluvoxamine maleate- FXN

GFAP- Glial fibrillary acidic protein

HEPES- 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

HRP- Horseradish peroxidase

IBA1- Ionized calcium-binding adapter molecule 1

IHC- Immunohistochemistry

LAMP-1- Lysosomal-associated membrane protein 1

LC3- Light chain 3

LPS- Lipopolysaccharide

MTOR- Mammalian target of rapamycin

NFT- Neurofibrillary tangles

NLR- NOD-like receptor

NLRP3- NLR family pyrin domain containing 3

OCD- Obsessive-compulsive Disorder

PBS- Phosphate buffer saline

PEG- Polyethylene glycol

PFA- Paraformaldehyde

PMSF- Phenylmethylsulfonyl fluoride

PVDF- Polyvinylidene difluoride

PYD- Pyrin domain

Rap- Rapamycin

RIPA- Radioimmunoprecipitation assay

SDS- Sodium dodecyl sulfate

SDS-PAGE- SDS poly acrylamide gel electrophoresis

SQSTM1- Sequestosome 1

ULK-1- unc-51-like kinase 1

Declarations

Ethics approval and consent to participate

All experiments involving mice were ethically approved by Institutional Animal Ethics Committee (IAEC) of CSIR-IIIM, Jammu, India. Detailed information regarding approval of *in-vivo* and *ex-vivo* experiments is given in Materials and Methods section.

Consent for publication

Not applicable

Availability of data and materials

All data generated or analyzed during this study are included in this published article and its supplementary information files.

Acknowledgements

We are thankful to University Grants Commission (UGC), India for providing the research fellowship to Ms. Sukhleen Kaur.

Authors' contributions

SK did most of the in vitro experiments along with some in vivo experiments. KS performed animal behavior studies. KKS and AS performed ELISA experiments. RA and PPS synthesized and characterized fluvoxamine maleate. SMA performed confocal

microscope studies. PR helped in genotyping of 5XFAD mice. ZA helped in designing the study. SK and AK designed the study, analyzed the data and wrote the manuscript.

Disclosure statement

The authors declare that they have no competing interests.

Funding

The funding for this project was provided by CSIR-Indian Institute of Integrative Medicine (CSIR-IIIM) through the project MLP-6002 (WP-4).

References:

1. Graff-Radford J, Yong KXX, Apostolova LG, Bouwman FH, Carrillo M, Dickerson BC, Rabinovici GD, Schott JM, Jones DT, Murray ME: **New insights into atypical Alzheimer's disease in the era of biomarkers.** *The Lancet Neurology* 2021, **20:**222-234.

2. Soria Lopez JA, Gonzalez HM, Leger GC: **Alzheimer's disease.** *Handbook of clinical neurology* 2019, **167:**231-255.

3. Minter MR, Taylor JM, Crack PJ: **The contribution of neuroinflammation to amyloid toxicity in Alzheimer's disease.** *Journal of neurochemistry* 2016, **136:**457-474.

4.   Al-Ghraiybah NF, Wang J, Alkhalifa AE, Roberts AB, Raj R, Yang E, Kaddoumi A:

Glial Cell-Mediated Neuroinflammation in Alzheimer's Disease. *International*

Journal of Molecular Sciences 2022, **23:**10572.

5.   Leng F, Edison P: **Neuroinflammation and microglial activation in Alzheimer**

disease: where do we go from here? *Nature Reviews Neurology* 2021, **17:**157-

172.

6.   Liu Y, Dai Y, Li Q, Chen C, Chen H, Song Y, Hua F, Zhang Z: **Beta-amyloid**

activates NLRP3 inflammasome via TLR4 in mouse microglia. *Neuroscience*

letters 2020, **736:**135279.

7.   Jung ES, Suh K, Han J, Kim H, Kang HS, Choi WS, Mook-Jung I: **Amyloid-β**

activates   NLRP3   inflammasomes   by   affecting   microglial

immunometabolism through the Syk-AMPK pathway. *Aging Cell* 2022,

21:e13623.

8.   Cai Z, Hussain MD, Yan L-J: **Microglia, neuroinflammation, and beta-amyloid**

protein in Alzheimer's disease. *International Journal of Neuroscience* 2014,

124:307-321.

9.   Thakur S, Dhapola R, Sarma P, Medhi B, Reddy DH: **Neuroinflammation in**

Alzheimer's   disease: current   progress   in   molecular   signaling   and

therapeutics. *Inflammation* 2023, **46:**1-17.

10.   Liang T, Zhang Y, Wu S, Chen Q, Wang L: **The role of NLRP3 inflammasome**

in Alzheimer's disease and potential therapeutic targets. *Frontiers in*

Pharmacology 2022, **13:**845185.

11.   Venegas C, Kumar S, Franklin BS, Dierkes T, Brinkschulte R, Tejera D, Vieira-

Saecker A, Schwartz S, Santarelli F, Kummer MP: **Microglia-derived ASC**

specks cross-seed amyloid-β in Alzheimer's disease. *Nature* 2017, **552:**355-

361.

12.   Cole SL, Vassar R: **The Alzheimer's disease β-secretase enzyme, BACE1.**

Molecular neurodegeneration 2007, **2:**1-25.

13.   Zhang Z, Yang X, Song Y-Q, Tu J: **Autophagy in Alzheimer's disease**

pathogenesis: Therapeutic potential and future perspectives. *Ageing*

research reviews 2021, **72:**101464.

14.   Wu A-G, Zhou X-G, Qiao G, Yu L, Tang Y, Yan L, Qiu W-Q, Pan R, Yu C-L, Law

BY-K: **Targeting microglial autophagic degradation in NLRP3**

inflammasome-mediated neurodegenerative diseases. *Ageing research*

reviews 2021, **65:**101202.

15.   Houtman J, Freitag K, Gimber N, Schmoranzer J, Heppner FL, Jendrach M:

Beclin1-driven autophagy modulates the inflammatory response of

microglia via NLRP 3. *The EMBO journal* 2019, **38:**e99430.

16.   Cho M-H, Cho K, Kang H-J, Jeon E-Y, Kim H-S, Kwon H-J, Kim H-M, Kim D-H,

Yoon S-Y: **Autophagy in microglia degrades extracellular β-amyloid fibrils**

and regulates the NLRP3 inflammasome. *Autophagy* 2014, **10:**1761-1775.

17.   Hou P, Tian T, Jia P, Lin Y, Li Z, Wang Y, Ye Y, Li C, Guo D: **The regulation of**

NLRP3 inflammasome activation by CCDC50-mediated autophagy.

Autophagy 2023, **19:**365-366.

18. La Rosa F, Zoia CP, Bazzini C, Bolognini A, Saresella M, Conti E, Ferrarese C,

Piancone F, Marventano I, Galimberti D: **Modulation of MAPK-and PI3/AKT-**

dependent autophagy signaling by stavudine (D4T) in PBMC of Alzheimer's

disease patients. *Cells* 2022, **11:**2180.

19. Zhang D, Zhang Y, Pan J, Cao J, Sun X, Li X, Zhang L, Qin C: **Degradation of**

NLRP3 by p62-dependent-autophagy improves cognitive function in

Alzheimer's disease by maintaining the phagocytic function of microglia.

CNS Neuroscience & Therapeutics 2023.

20. Deng Y, Wang S-Y, Wang Q-G, Xu Z-H, Peng Q, Chen S-Y, Zhu L, Zhang Y-D,

Duan R: **AVE 0991 Suppresses Astrocyte-Mediated Neuroinflammation of**

Alzheimer's Disease by Enhancing Autophagy. *Journal of Inflammation*

Research 2023**:**391-406.

21. Jorfi M, Maaser-Hecker A, Tanzi RE: **The neuroimmune axis of Alzheimer's**

disease. *Genome Medicine* 2023, **15:**1-25.

22. Veitch DP, Weiner MW, Aisen PS, Beckett LA, Cairns NJ, Green RC, Harvey D,

Jack Jr CR, Jagust W, Morris JC: **Understanding disease progression and**

improving Alzheimer's disease clinical trials: Recent highlights from the

Alzheimer's Disease Neuroimaging Initiative. *Alzheimer's & Dementia* 2019,

15:106-152.

23. Ferrari C, Sorbi S: **The complexity of Alzheimer's disease: an evolving**

puzzle. *Physiological reviews* 2021, **101:**1047-1081.

24. Elliott EI, Sutterwala FS: **Initiation and perpetuation of NLRP 3 inflammasome**

activation and assembly. *Immunological reviews* 2015, **265:**35-52.

25.   Moonen S, Koper MJ, Van Schoor E, Schaeverbeke JM, Vandenberghe R, von

717   Arnim CA, Tousseyn T, De Strooper B, Thal DR: **Pyroptosis in Alzheimer's**

718   **disease: cell type-specific activation in microglia, astrocytes and neurons.**

719   *Acta Neuropathologica* 2023, **145:**175-195.

720   26.   Severini C, Barbato C, Di Certo MG, Gabanella F, Petrella C, Di Stadio A, de

721   Vincentiis M, Polimeni A, Ralli M, Greco A: **Alzheimer's Disease: New**

722   **Concepts on the Role of Autoimmunity and NLRP3 Inflammasome in the**

723   **Pathogenesis of the Disease.** *Current neuropharmacology* 2021, **19:**498-512.

724   27.   Sharif H, Wang L, Wang WL, Magupalli VG, Andreeva L, Qiao Q, Hauenstein AV,

725   Wu Z, Núñez G, Mao Y: **Structural mechanism for NEK7-licensed activation**

726   **of NLRP3 inflammasome.** *Nature* 2019, **570:**338-343.

727   28.   Goodman WK, Ward H, Kablinger A, Murphy T: **Fluvoxamine in the treatment**

728   **of obsessive-compulsive disorder and related conditions.** *Journal of Clinical*

729   *Psychiatry* 1997, **58:**32-49.

730   29.   Mundo E, Maina G, Uslenghi C, Group MS: **Multicentre, double-blind,**

731   **comparison of fluvoxamine and clomipramine in the treatment of**

732   **obsessive-compulsive disorder.** *International clinical psychopharmacology*

733   2000, **15:**69-76.

734   30.   Koran LM, Bromberg D, Hornfeldt CS, Shepski JC, Wang S, Hollander E:

735   **Extended-release fluvoxamine and improvements in quality of life in**

736   **patients with obsessive-compulsive disorder.** *Comprehensive psychiatry*

737   2010, **51:**373-379.

31. Rajesh Y, Kanneganti T-D: **Innate immune cell death in neuroinflammation**
and Alzheimer's disease. *Cells* 2022, **11:**1885.

32. Maccioni RB, Rojo LE, Fernandez JA, Kuljis RO: **The role of**
neuroimmunomodulation in Alzheimer's disease. *Annals of the New York*
Academy of Sciences 2009, **1153:**240-246.

33. Yu L, Chen Y, Tooze SA: **Autophagy pathway: Cellular and molecular**
mechanisms. *Autophagy* 2018, **14:**207-215.

34. Saitoh T, Fujita N, Jang MH, Uematsu S, Yang B-G, Satoh T, Omori H, Noda T,
Yamamoto N, Komatsu M: **Loss of the autophagy protein Atg16L1 enhances**
endotoxin-induced IL-1β production. *Nature* 2008, **456:**264-268.

35. Nakahira K, Haspel JA, Rathinam VA, Lee S-J, Dolinay T, Lam HC, Englert JA,
Rabinovitch M, Cernadas M, Kim HP: **Autophagy proteins regulate innate**
immune responses by inhibiting the release of mitochondrial DNA
mediated by the NALP3 inflammasome. *Nature immunology* 2011, **12:**222-
230.

36. Lee J-H, Yang D-S, Goulbourne CN, Im E, Stavrides P, Pensalfini A, Chan H,
Bouchet-Marquis C, Bleiwas C, Berg MJ: **Faulty autolysosome acidification in**
Alzheimer's disease mouse models induces autophagic build-up of Aβ in
neurons, yielding senile plaques. *Nature neuroscience* 2022, **25:**688-701.

Figure legends

Figure 1. FXN displayed a potent anti-NLRP3 inflammasome activity in primary
astrocytes.

(A) Structure of Fluvoxamine maleate. (B) and (C) Graphs representing IC_{50} of FXN for

IL-1β and IL-18, respectively. The IC50 values were calculated in primary astrocytes

after treatment with FXN in the serum free media for 1 h and ATP (5 mM) for 30

minutes. (D) Immunoblots depicting the expression of NLRP3 inflammasome complex

proteins including IL-1β, CASP1 and NLRP3. Densitometric analysis of the western

blots is provided in Supplementary data (Fig. S1A). (E) Representative confocal images

indicating the ASC speck formation in primary astrocytes, which generally form after

NLRP3 inflammasome activation, and the graph showing average number of specks is

provided in the Supplementary data (Fig. S1B). (F) Western blots showing the ASC

oligomerization. Densitometric analysis of immunoblots is given in Supplementary data

(Fig. S1C). (G) Representative images showing co-localization of NLRP3 (green) and

CASP1 (red) in FXN (78 nM) treated primary astrocytes. The mean fluorescence

intensities of NLRP3 and CASP1 is provided in the Supplementary data (Fig S1D). The

above-mentioned findings revealed that FXN significantly decreased the release of IL-

1β and IL-18 and inhibited the expression of proteins involved in the NLRP3 complex

formation including CASP1, NLRP3 and ASC. The data provided are Mean±SD of three

independent experiments and the statistical analysis was measured using one-way

ANOVA, followed by post-hoc Bonferroni test. The p-value<0.05 was considered to be

statistically significant with values assigned as ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$,

*$p < 0.05$ and ns= not significant.

Figure 2. FXN inhibited the NLRP3 inflammasome through NF-κB pathway.

The effect of FXN on NF-κB pathway was analyzed by treating the primary astrocytes

with FXN, prior to LPS priming and ATP activation of NLRP3 inflammasome. The

analysis of pro-inflammatory cytokines regulated by NF-κB pathway was done through ELISA. (A) Graph representing IC50 value of FXN for IL-1β, (B) and (C) Graphs representing the TNF-α and IL-6 levels, respectively, after FXN treatment (78 nM). (D) Analysis of nuclear translocation of NF-κB (p65) through western blotting. LMNA was used as a nuclear marker and ACTB was used as cytosolic marker. (E) and (F) showing densitometric analysis of nuclear and cytosolic fractions of NF-κB (p65), respectively. The data indicates the significant reduction in the NF-κB mediated secretion of TNF-α and IL-6 by FXN treatment via inhibiting the nuclear translocation of NF-κB (p65). The data shown here are Mean±SD of three independent experiments and the statistical analysis was calculated using one-way ANOVA, followed by post-hoc Bonferroni test. The p-value<0.05 was considered to be statistically significant with values assigned as ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$ and ns= not significant.

Figure 3. FXN induced the autophagy under the inflammatory conditions in primary astrocytes

(A) Immunoblots showing concentration dependent effect of FXN and rapamycin on the expression of LC3B-II and SQSTM1 after 24 h treatment in primary astrocytes. Densitometric analysis is given in Supplementary data (Fig. S2A). (B) Immunoblots showing the time dependent effect of FXN (78 nM) on autophagic flux in the presence and absence of bafilomycin A1 in primary astrocytes. Densitometric analysis is provided in Supplementary (Fig. S2B). (C) Representative images showing the effect of FXN (78 nM) on co-localization of LAMP1 (green) and LC3B (red) in primary astrocytes. The increased colocalization of LAMP1 and LC3B after FXN treatment indicates the induction of autophagy. (D) Graph representing the mean fluorescence intensities of

LAMP1 and LC3B. (E) Immunoblots showing the effect of FXN (78 nM) on LC3B-II and

SQSTM1 under NLRP3 inflammasome activation conditions in primary astrocytes.

Densitometry of immunoblots is provided in Supplementary Data (Fig. S2C). (F)

Immunoblots showing the effect of FXN on pMTOR (Ser 2448) and other autophagic

proteins involved in the initiation, elongation, maturation and fusion of autophagosome

[pCAMKK2 (Ser 511), pAMPK (Thr 172), pULK1 (Ser 317), ATG 13, FIP 200, BECN1,

ATG5 and ATG7] in primary astrocytes. The data provided are Mean±SD of three

independent experiments and the statistical analysis was performed using one-way

ANOVA, followed by post-hoc Bonferroni test. The p-value<0.05 was considered to be

statistically significant with values assigned as ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$,

*$p < 0.05$ and ns= not significant.

Figure 4. FXN inhibited NLRP3 inflammasome by inducing autophagy

To investigate the involvement of autophagy in the FXN mediated inhibition of NLRP3

inflammasome, autophagy was genetically and pharmacologically inhibited by

siPRKAA2 and bafilomycin. A1 respectively in primary astrocytes under inflammasome

activation conditions. (A) Western blots indicating the effect of PRKAA2 knockdown on

the expression levels of pPRKAA2 (Thr 172), BECN1, LC3B-II, SQSTM1, NF-κB (p65),

IL-1β and NLRP3. Densitometric analysis is given in Supplementary data (Fig. S3A). (B)

Representative images indicating the colocalization of NLRP3 and CASP1 under

aforementioned conditions (genetic knockdown of PRKAA2 by *siPRKAA2*) after FXN

treatment. The mean fluorescence intensities of NLRP3 and CASP1 are provided in the

Supplementary data (Fig. S3B). (C) Immunoblots depicting the PRKAA2 expression in

primary astrocytes following treatment with the *siPRKAA2,* in comparison to mock

siRNA and their densitometric analysis. (D) Immunoblots showing the effect of FXN on

expression levels of IL-1β, NLRP3, NF-κB (p65), LC3B-II and SQSTM1 after

pharmacological inhibition of autophagy using bafilomycin A1. Densitometric analysis is

given in Supplementary data (Fig. S3C). The data represent Mean±SD of three

independent experiments and the statistical analysis of data was analyzed using one-

way ANOVA, followed by post-hoc Bonferroni test. The p-value<0.05 was considered to

be statistically significant with values assigned as ****p < 0.0001, ***p < 0.001, **p<

0.01, *p< 0.05 and ns= not significant.

Figure 5. FXN cleared Aβ$_{42}$ in primary astrocytes by inducing autophagy

(A) Representative images showing the effect of FXN, in the presence Baf. A1, on the

intracellular deposition of Aβ in primary astrocytes treated with fluorescently tagged

(HiLyte 488) Aβ peptide. Rapamycin was used as a standard autophagy inducer. (B)

Representative images showing the effect of FXN on intracellular deposition of Aβ in

primary astrocytes transfected with *siPRKAA2*. (C) Graph representing the mean

fluorescence intensity of images shown in Fig. 5A. (D) Graphical representation of mean

fluorescence intensity of images shown in Fig. 5B. The decrease in the fluorescence

intensity of Aβ in FXN treated cells shows its potential to clear deposited Aβ and the

reversal in the fluorescence intensities of Aβ where the primary astrocytes were treated

with Baf. A1 or transfected with *siPRKAA2,* indicates the involvement of autophagy in

FXN mediated Aβ clearance. The data shown here are Mean±SD of three independent

experiments and the statistical analysis of data was calculated using one-way ANOVA,

followed by post-hoc Bonferroni test. The p-value<0.05 was considered to be

statistically significant with values assigned as ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$,

*$p < 0.05$ and ns= not significant.

Figure 6. FXN improved working memory, exploratory behavior and

neuromuscular coordination in 5XFAD mice

The effect of FXN (5 and 10 mg/kg) on cognitive behavior was assessed post 2 months

dosing in Alzheimer (5XFAD) mice model (n=5 per group). Vehicle 5XFAD control mice

were compared with wild-type C57BL/6J mice of same age. Radial Arm Maze Test: (A)

Representative images showing track plots of radial arm maze post five days training.

The different parameters were assessed such as (B) Latency to first entry to baited arm,

(C) Time spent in baited arm, (D) Entries to non-baited arms and (E) Distance travelled

by mice. (F) Images depicting the track plots obtained from open field test. Various

parameters were investigated such as (G) Distance travelled by mice, (H) Mobility time,

(I) Time spent in the Centre zone, (J) Corner zone time and (K) Maximum speed. The

latency to fall time of mice from rotarod was assessed by using rotarod test. (L) Graph

representing the latency to fall time. Automated camera was used to capture the

movement of mice in radial arm maze and open field test and all the activities were

measured by using AnyMaze software. The statistical analysis of data was measured

using one-way ANOVA, followed by post-hoc Bonferroni test. The p-value <0.05 was

considered to be statistically significant with values assigned as ****$p < 0.0001$, ***$p <$

0.001, **$p < 0.01$, *$p < 0.05$ and ns= not significant.

Figure 7. FXN reduced the Aβ_{42} in the hippocampus and plasma of 5XFAD mice

Immunostaining post FXN treatment (5 and 10 mg/kg for 2 months), the formalin-fixed

paraffin embedded (FFPE) tissues of brain hippocampi were immuno-stained with

D3E10 for detection of total Aβ (green), GFAP for activated astrocytes (red) and the

nuclei were counterstained with DAPI (blue). (A) Representative images showing co-

localized Aβ plaques and reactive astrocytes in hippocampi of FXN treated mice (n=5

per group). Graphs showing (B) Average number of Aβ plaques (C) Average skeletal

length of Aβ plaques (D) Average branch count of Aβ plaques (E) Average

circumference of Aβ plaques (G) The plasma levels of Aβ-42 in mice measured using

ELISA. The Aβ load and astrogliosis in FXN treated 5XFAD mice was found to

significantly reduced. Confocal microscopic images were examined using cell pathfinder

software. The statistical analysis was performed using one-way ANOVA, followed by

post-hoc Bonferroni test. The p-value <0.05 was considered to be statistically significant

with values assigned as ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$ and ns= not

significant.

Figure 8. FXN ameliorated the AD pathology in 5XFAD mice through autophagy-

mediated reduction in amyloid beta levels and neuroinflammation

(A) Representative confocal images showing LC3B (green) in hippocampi sections of

FXN treated mice (5 and 10 mg/kg) (n=5 per group) and the graph representing the

average number of LC3B puncta from total of 20 images obtained from each group. (B)

Immunoblots showing CAMKK2 mediated autophagy induction via PRKAA2 pathway,

indicated by significant increase in the expression of pCAMKK2 (Ser 511), pPRKAA2

(Thr 172), LC3B-II, pULK (Ser 317), BECN1 and ATG5, and a decreased expression of

pMTOR (Ser 2448) in hippocampi of FXN treated mice. Densitometric analysis of the

western blots is provided in Supplementary data (Fig. S4A). (C) Immunoblots depicting the decreased expression of neuroinflammatory markers NF-κB (p50), NF-κB (p65), NLRP3, ASC oligomer, ASC dimer and ASC monomer, CASP1, IL-1β and IBA1 in hippocampi of FXN treated mice. (D) Densitometric analysis showing decreased levels of ASC oligomer, dimer and monomer forms in FXN treated mice. The densitometry of other western blots is provided in Supplementary data (Fig. S4B). Graphs depicting the levels of pro-inflammatory cytokines in brain cortex of FXN treated 5XFAD mice (E) TNF-α, (F) IL-6 and (G) IL-1β. The samples were compared by using one-way ANOVA, followed by post-hoc Bonferroni test. The p-value <0.05 was considered to be statistically significant with values assigned as ****$p < 0.0001$, ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$ and ns= not significant.

Fig. 1

Fig. 2

Fig. 3

A

Time 24 h				-					
Rap (nM)	-	250	-	39	-	-	-	-	-
FXN (nM)	-	-	19		78	156	312	625	1250

LC3B-I
LC3B-II

SQSTM1

ACTB

B

Time (h)	0	3	6	12	24	0	3	6	12	24
Baf. A1 (20 nM)	-	-	-	-	-	+	+	+	+	+
FXN (78 nM)	-	+	+	+	+	-	+	+	+	+

LC3B-I
LC3B-II

SQSTM1

ACTB

C

DAPI   LC3B   LAMP1   Merge

Control

Rap

FXN

Baf. A1

Baf. A1+FXN

D

E

FXN (nM)	-	-	-	78
ATP (5 mM)	-	-	+	+
LPS (1 µg/ml)	-	+	+	+

LC3B-I
LCB-II

SQSTM1

ACTB

F

FXN (nM)	-	-	-	78
ATP (5 mM)	-	-	+	+
LPS (1 µg/ml)	-	+	+	+

pCAMKK2
(Ser 511)

pPRKAA2
(Thr 172)

ACTB

pMTOR
(Ser 2448)

MTOR

pULK1
(Ser 317)

ULK1

ATG13

FIP200

ACTB

BECN1

ATG5

ATG7

ACTB

Fig. 4

A

FXN (78 nM)	-	-	+	-	-	+
ATP (5 mM)	-	+	+	-	+	+
LPS (1 µg/ml)	-	+	+	-	+	+
siPRKAA2	-	-	-	+	+	+
sicontrol	+	+	+	-	-	-

pPRKAA2 (Thr 172)

PRKAA2

BECN1

LC3B-I
LC3B-II

SQSTM1

pNF-κB (p65) (Ser 536)

IL-1β

Pro-IL-1β

NLRP3

ACTB

C

siPRKAA2	-	+
sicontrol	+	-

pPRKAA2 (Thr 172)

PRKAA2

ACTB

pPRKAA2: PRKAA2

****

siPRKAA2	-	+
sicontrol	+	-

D

FXN (78 nM)	-	-	+	-	-	+
ATP (5 mM)	-	+	+	-	+	+
LPS (1 µg/ml)	-	+	+	-	+	+
Baf. A1 (20 nM)	-	-	-	+	+	+

IL-1β

Pro-IL-1β

NLRP3

pNF-κB (p65) (Ser 536)

LC3B-I
LC3B-II

SQSTM1

ACTB

B

	Control	LPS+ATP	FXN (78 nM)	Control *siPRKAA2*	LPS+ATP+ *siPRKAA2*	FXN+*siPRKAA2*

Merge

CASP1

NLRP3

DAPI

● **DAPI**   ● **CASP1**   ● **NLRP3**

Fig. 5

Fig. 6

Fig. 7

A

	DAPI	Aβ₄₂ (D3E10)	GFAP	Merge	Zoomed Inset
5XFAD Control					
FXN 5 mg/Kg					
FXN 10 mg/Kg					

Fig. 8