Regulatory networks of KRAB zinc finger genes and transposable elements changed during human brain evolution and disease

Yao-Chung Chen[1,2*], Arnaud Maupas[3], Katja Nowick[1,2*]

[1] Human Biology and Primate Evolution, Institute for Biology, Freie Universität Berlin, Berlin, Germany

[2] Institute of Bioinformatics, Freie Universität Berlin, Berlin, Germany

[3] Institut de Biologie de l'École Normale Supérieure, Ecole Normale supérieure de Paris, Paris, France

[*]**Corresponding author/e-address**: yao-chung.chen@fu-berlin.de, katja.nowick@fu-berlin.de

Abstract

Evidence indicates that transposable elements (TEs) can contribute to the evolution of new traits, despite often being considered deleterious. In vertebrates, some KRAB-ZNF proteins repress TEs, offering genomic protection. Notably, KRAB-ZNF genes evolve rapidly and exhibit diverse expression patterns in primate brains, where TEs remain active. This prompts questions about their interactions in primate brains and potential roles in human brain evolution and disease. For a systematic comparative analysis of TE interactions with other genes, we developed the tool TEKRABber, and focused on strong and experimentally validated cases. Our bipartite network analysis revealed significantly more interactions between KRAB-ZNF genes and TEs in humans than in other primates, especially with recently evolved TEs (≤ 44.2 million years old). Notably, ZNF528, under positive selection in humans, shows numerous human-specific TE interactions. Most negative interactions in our network, indicative of repression by KRAB-ZNF proteins, entail Alu TEs, while links to other TEs are

generally positive. In Alzheimer's patients, interactions within a subnetwork involving 21 KRAB-ZNF and Alu pairs appear diminished or lost. Our findings suggest that KRAB-ZNF and TE interactions vary across TE families, have increased throughout human evolution, and may influence susceptibility to Alzheimer's disease.

Keywords: primate evolution, Alzheimer's disease, KRAB zinc finger, transposable elements, regulatory network, bipartite network.

Graphical Abstract

Introduction

Transposable elements (TEs) are repetitive DNA sequences capable of migrating and inserting into new locations within the host genome. When transposed, TEs may induce phenotypic changes in the organism (Bourque et al., 2018; Schrader & Schmitz, 2019). One well-known textbook example is the development of an industrial melanization mutant phenotype in the British peppered moth, *Biston betularia*, which has been linked to increased expression of the *cort* transcript due to a TE insertion in its first intron (Hof et al., 2016). Another classic example for primate evolution is the retrovirus insertion in the salivary amylase gene, *AMY1C*, with this sequence being essential for primates to evolve as one of the few mammals capable of expressing amylase in saliva (Ting et al., 1992). The other recent example is the evolutionary loss of tails in humans and apes, which is associated with the insertion of an Alu retrotransposon into the intron region of the *TBXT* gene. This insertion creates an alternative splice site with an ancestral Alu retrotransposon in the sequence, leading to the excision of the middle exon and the formation of an isoform (Xia et al., 2024).

TEs comprise one third to one half of the mammalian genome and are not randomly distributed (Platt et al., 2018; Qu et al., 2023). In fact, some TE insertions can be deleterious such as the insertion of an L1 element initiating colorectal cancer in humans (Scott et al., 2016). Therefore, a variety of host factors are employed to regulate TE expression, such as small RNAs, chromatin and DNA modification pathways, and KRAB zinc finger (KRAB-ZNF) proteins (Colonna Romano & Fanti, 2022). KRAB-ZNF proteins are the largest family of transcription factors in higher vertebrates (Ecco et al., 2017; Huntley et al., 2006), characterized by fast evolution and contributing to gene expression differences between primates (Nowick et al., 2009, 2013). KRAB-ZNF proteins bind to the interspersed DNA sequences of TEs and

repress their expression upon recruiting the cofactor KAP1 (Groner et al., 2010). Fast evolution was also reported for TEs, which led to hypothesizing an evolutionary arms-race model, in which mutated TEs have the chance to escape the repression from KRAB-ZNF proteins until the KRAB-ZNF proteins evolve again a suitable recognition ability (Imbeault et al., 2017; Jacobs et al., 2014). For instance, in primates, coevolution between a family of TEs, retroelements in endogenous retroviruses (ERVs), and the evolving tandem repeats in KRAB-ZNF genes has been observed (Thomas & Schneider, 2011).

Studying the expression of KRAB-ZNF genes and TEs in the evolution of the human brain has crucial relevance, since some TEs are actively transcribed in the development of the human brain (Bodea et al., 2018), including L1 subfamilies, and primate-specific Alu subfamilies (Hancks & Kazazian, 2010; Larsen et al., 2018). Furthermore, TE-derived promoters play a role in gene transcription within the human brain, which in turn suggests the significance of TEs in gene regulation during neurodevelopment (Playfoot et al., 2021). Interestingly, a substantial number of differentially expressed primate-specific KRAB-ZNF genes were detected in the adult prefrontal cortex, comparing humans to chimpanzees (Nowick et al., 2009) and newly evolved KRAB-ZNF genes were predominantly detected in the developing human brain (Zhang et al., 2011). Therefore, analyzing the interplay between TEs and KRAB-ZNF genes in the context of primate brain evolution could provide valuable insights into the complex mechanisms shaping human brain development and function.

The dysregulation of TE expression has also been linked to neurodegenerative diseases, including Alzheimer's disease (AD), the primary cause of dementia (Ravel-Godreuil et al., 2021). One of the biomarkers of AD, the Tau protein, appears to induce the expression of at least some TEs (Guo et al., 2018), and the presence of TE

products in the cytosol and endosomes induces neuroinflammation in AD patients (Evering et al., 2023). Previous studies have suggested that cognitive skills are related to zinc finger genes and proteins. For instance, patients carrying the schizophrenia-risk allele of *ZNF804A* showed differences in their reading and spelling performance (Becker et al., 2012). Lead poisoning causes lead ions to compete with zinc ions in zinc finger proteins, affecting proteins such as DNMT1, which are related to the progression of AD (Ordemann & Austin, 2016). Additionally, the single nucleotide polymorphism of a KRAB-ZNF gene, *ZNF224*, is associated with AD neuropathology and cognitive functions (Shulman et al., 2010).

Taken together, the study of the regulatory networks involving KRAB-ZNF genes and TEs is expected to provide insights into the evolution of the human brain and the development of neurodegenerative diseases. Here, we present results from two independent RNA-seq datasets: one comparing different brain regions between humans and multiple non-human primates (NHPs) (**Primate Brain Data**), and the other comparing human control samples with AD samples (**Mayo Data**). These datasets collectively encompass a total of 514 samples from different species, brain regions, and disease states. Since it is very challenging to quantify and normalize expression of TEs for across species comparison, we developed the TEKRABber R package for the systematic and comparative analysis of TE subfamilies (abbreviated as **TEs** in the following content) and KRAB-ZNFs relationships. Our work reveals an intricate network of TEs and KRAB-ZNF genes in the brain that changed during evolution and is modified in AD brains.

Results

TEKRABber: A software for cross-species comparative analysis of orthologs, TEs, and their co-expression

While expression of orthologous genes can be compared between closely related species relatively easily, this task is more challenging for TEs, because the same TE subfamilies can be located in different regions on chromosomes and have different sequence lengths. Additionally, differences in the copy number of TEs between species can further complicate these comparisons. Therefore, efficient tools are needed to estimate the pairwise correlations between orthologous genes and TEs for investigating the interplay between them. To efficiently compare the gene regulatory networks of KRAB-ZNFs and TEs across species, we developed an R Bioconductor software called TEKRABber (DOI: 10.18129/B9.bioc.TEKRABber). The name is derived from the idea that TEs are often repressed by KRAB-ZNF proteins. In a broader scope, TEKRABber addresses two primary challenges: comparing TE expression across species and efficiently calculating pairwise correlations between selected orthologous genes and TEs.

TEKRABber is designed to handle various types of transcriptomic read counts and offers two distinct modes of analysis (**Figure 1A**). In the first mode, tailored for interspecies comparison, we utilized the **Primate Brain Data** as a demonstration. Initially, it retrieves annotations from Ensembl (Harrison et al., 2024) for orthologs and from RepeatMasker (Smit et al., 2013) for TEs to estimate normalizing factors, ensuring comparable expression levels between species. Subsequently, users can employ the output data object to conduct differential expression (DE) analysis and identify one-to-one correlations based on selected parameters. Notably, TEKRABber (from version 1.8, Bioc3.19) includes a parallel computing option, significantly

enhancing computational efficiency based on the number of cores a device can provide. The second mode is designed for comparing different conditions, such as control and disease states within the same species. In this scenario, we used the **Mayo Data** as an example. Users can bypass the interspecies normalization steps and directly generate data objects for differential expression and correlation analyses. We used TEKRABber to comparatively investigate correlations between KRAB-ZNFs and TEs, but it can also be used for the analysis of functional relationships between TEs and any other genes. Furthermore, TEKRABber offers an interactive user interface, providing users with an initial overview of their results before delving into the details (**Figure 1B**).

In our study, we used TEKRABber to explore the functional connections between KRAB-ZNFs and TEs in the context of human brain evolution and Alzheimer's disease (AD). We conducted an analysis of KRAB-ZNF genes and TEs expression patterns and networks using two independent RNA-seq datasets: **Primate Brain Data** and **Mayo Data** (**Figure 1A**, **Tables S1-S3**). The Primate Brain Data contains genome-wide expression information from 33 brain regions classified into seven groups (Khrameeva et al., 2020) of four primate species, while the Mayo Data includes data from two brain regions (temporal cortex and cerebellum) of AD patients and controls. In brief, expression of all genes and TEs was quantified and normalized across samples of each dataset, and subsequently differential expression and correlations between KRAB-ZNFs and TEs were calculated (**Figure 1A**).

Figure 1. TEKRABber and the overview of the analysis workflow. (A) Two independent RNA-seq datasets, **Primate Brain Data** and **Mayo Data**, were analyzed in this study. **(1)** Transcriptomic data were first preprocessed with removing adapters and low-quality reads and then mapped to their reference genome using STAR to generate BAM files. **(2)** TEtranscript was used to quantify the expression of genes and TEs. **(3)** Expression profiles were normalized across different species. **(4)** DE analysis and pairwise correlations were calculated. Steps **(3)** and **(4)** were developed together in an R Bioconductor package, TEKRABber. **(B)** The user interface of TEKRABber features a dashboard layout that allows users to explore one-to-one gene-TE interaction, including correlation and differential expression results. (Details are given in the Materials and Methods section)

Dynamics of expression across species and brain regions: Strong species differences especially of evolutionary young KRAB-ZNF genes and TEs

We obtained normalized expression values of KRAB-ZNF genes and TEs and assessed their variance across different species using t-SNE clustering (**Figure 2A**). The data labeled by species revealed distinct boundaries of variance, demonstrating clustering based on species. Specifically, humans and macaques formed their own clusters, while chimpanzees and bonobos were grouped in the same cluster, which

agrees with phylogenetic distances among species. This finding indicated that clear differences in KRAB-ZNF genes and TE expression can be detected across species. For example, there were 12 up-regulated and 42 down-regulated KRAB-ZNF genes, along with 31 up-regulated and 22 down-regulated TEs in humans compared to chimpanzees in the primary and secondary cortices (**Figure 2B;** see **Table S2** for information on Brodmann areas included in the group "primary and secondary cortices"). In contrast to the clear species differences, expression patterns of KRAB-ZNFs and TEs differed less across brain regions within the same species.

According to the arms-race model, we hypothesize that the expression of TEs and KRAB-ZNFs that are evolutionarily similar in age will be significantly correlated. Thus, to investigate whether expression patterns differ between evolutionary old and young KRAB-ZNF genes and TEs, respectively, we dated these genomic elements and classified them into two groups based on their inferred evolutionary age. The **old** group consists of 237 KRAB-ZNF genes and 842 TEs that evolved prior to the emerge of Simiiformes (> 44.2 million years ago (mya)), while the **young** group consists of 100 KRAB-ZNF genes and 309 TEs that evolved less than 44.2 mya (**Figure S1** and **S2**). We first found that the young KRAB-ZNFs and young TEs exhibited significantly lower expression levels across all brain regions, regardless of species (**Figure 2C** and **S3**). Next, there were proportionally more young KRAB-ZNF genes and young TEs differentially expressed between humans and chimpanzees compared to old KRAB-ZNF genes and old TEs (**Figure 2D**). Comparing humans to NHPs, there are more proportions of young KRAB-ZNFs and young TEs that are differentially expressed which implies although in lower expression levels, young TEs and young KRAB-ZNFs are still more dynamically changed between different primates.

We further investigated if there are KRAB-ZNFs and TEs that are specifically

changed in humans compared to the three NHPs. We detected 36 KRAB-ZNF

genes and 18 such TEs in the primary and secondary cortices, with a trend towards

more downregulation of KRAB-ZNFs and upregulation of TEs in humans, such as

ZNF337 and *ZNF394* (**Figure 2E**). Unlike this trend, some KRAB-ZNFs linked to

cognitive disorders show human specific upregulation, e.g. *ZNF778*, a candidate gene

for autism spectrum disorder and cognitive impairment (Willemsen et al., 2010), and

ZNF267, which is up-regulated in the prefrontal cortex of AD patients (Patel et al.,

2021). On the other hand, the LTR12B subfamily is one of the most down-regulated

TEs in humans compared to NHPs. LTR12-related endogenous retroviruses

subfamilies had been reported to be repressed by ZNF676 and ZNF728 in early human

development (Iouranova et al., 2022) which we also observed.

Figure 2. Expression of KRAB-ZNF genes and TEs in Primate Brain Data. (A) t-SNE plots of the expression of KRAB-ZNF genes and TEs from all 422 samples including human, chimpanzee, bonobos, and macaques labeled by species and different brain regions. **(B)** Differentially expressed KRAB-ZNF genes and TEs comparing human and chimpanzee in primary and secondary cortices. **(C)** Expression of KRAB-ZNF genes and TEs in primary and secondary cortices across species. Both KRAB-ZNF genes and TEs were grouped in two groups based on their inferred evolutionary age, old (> 44.2 mya) and young (≤ 44.2mya). Young KRAB-ZNFs and young TEs have lower expression levels (Wilcoxon Rank Sum Test, $p < 0.05$). Expressions of all brain regions can be found in **Figure S3**. **(D)** Percentage of differentially expressed KRAB-ZNF genes and TEs in humans compared to chimpanzees in primary and secondary cortices and limbic and association cortices. **(E)** Human-specific DE (human-specifically-changed) KRAB-ZNF genes and TEs in primary and secondary cortices compared to NHPs. Gray indicates no expression information. The colors for age inferences in **(C)**, **(D)**, and **(E)** are the same: blue for evolutionary old and yellow for evolutionary young KRAB-ZNFs and TEs, respectively.

Changes in correlations between TEs and KRAB-ZNF genes: Increased connectivity in the human brain co-expression network with an enrichment for evolutionary young correlations

To systematically analyze the functional relationships between TEs and KRAB-ZNF genes, we conducted pairwise Pearson's correlation analysis using normalized expression levels in seven clustered brain regions (**Table S2** and Figure 1 in Khrameeva et al., 2020). We first analyzed the human samples. There were 324

KRAB-ZNFs and 895 TEs expressed in Primate Brain Data. In humans, we found 100,987 positive and 26,810 negative significant correlations between TEs and KRAB-ZNFs in the primary and secondary cortices and 38,295 positive and 11,475 negative significant correlations in the limbic and association cortices (FDR < 0.01), while the other clusters have fewer or no correlations detected (**Table S4**). We will thus mainly focus on the primary and secondary cortices and the limbic and association cortices for our subsequent analyses.

The numbers of correlations between TEs and KRAB-ZNFs are significantly more than expected by chance, as gauged by repeating the correlation analysis with randomly picked genes and TEs ($p < 0.001$; see Methods; **Figure 3A** and **S4**), indicating that functional relationships between KRAB-ZNFs and TEs can be detected in the data. The high number of positive correlations might be surprising, given that KRAB-ZNFs are considered to repress TEs. However, the dataset contains in general more positive correlations, even when choosing random genes, and KRAB-ZNFs still have more negative correlations to TEs than random genes.

To remove weak correlations, we set a threshold with absolute correlation coefficients greater than 0.4 and FDR below 0.01 (Benjamini-Hochberg correction) (**Figure 3B**). In addition, we sought for independent experimental confirmation of our detected relationships by overlapping the correlations with ChIP-exo data obtained from KRAB-ZNF proteins in human embryonic stem cells (Imbeault et al., 2017). Note that although ChIP-exo was performed in different cell types than the brain samples we analyzed, the overlap was significant by calculating their Jaccard similarity ($p < 0.001$, **Figure 3C**). This result supports that our correlations have significantly captured the interplay between TEs and KRAB-ZNFs. Focusing on the ChIP-exo confirmed correlations, we obtained 869 correlations in the primary and secondary cortices and

399 correlations in limbic and association cortices. Of those, 201 positive correlations and 166 negative correlations overlapped between these two brain region groups (**Figure 3D**).

To better understand the complexity of the interactions between TEs and KRAB-ZNFs, we represented the one-to-one pairwise correlations of TE and KRAB-ZNF (denoted as **TE:KRAB-ZNF** in the following content) in a bipartite network. In this bipartite network, nodes belong to one of two classes, TEs or KRAB-ZNFs, and only links between nodes of different classes are present. Our bipartite network of human primary and secondary cortices reveals 354 nodes connected by 869 links (**Figure 3E**) and the human limbic and association cortices has 247 nodes with 399 links (**Figure S5**). To identify the hubs in the network, we first calculated their normalized degree distributions, considering the unequal numbers of TEs and KRAB-ZNFs. Our analysis revealed that KRAB-ZNF nodes connect to more TEs than TE nodes connect to KRAB-ZNFs, indicating that KRAB-ZNF nodes are more likely to act as hubs with higher connectivity (**Figure 3F**). For investigating the structure of the network, we calculated the bipartite modularities based on the correlation coefficients. Our finding revealed that the network of the primary and secondary cortices is clustered into five modules (**Figure 3E**), while the network of the limbic and association cortices clusters into 13 modules (**Figure S5**).

We divided TE:KRAB-ZNF into evolutionary young and old, calling a correlation as **young** when one of the nodes is evolutionary young and **old** when both nodes are evolutionary old. Taking also the correlation coefficients into account, all links can be classified into four categories: positive-old (P-O), positive-young (P-Y), negative-old (N-O) and negative-young (N-Y). We found that the links are non-randomly distributed in the network. In particular, Alu, ERV1, ERVK, ERVL, L1, TcMar-Tigger, and SVA

have significantly different interactions with KRAB-ZNF than the other TE subfamilies (Chi-squared test, *p*< 0.001). For example, in a module with Alu subfamilies (**Figure 3G**), we observed that most of the correlations specifically belong to N-Y, while SVA had many P-Y correlations with KRAB-ZNFs (**Figure 3H**).

Figure 3. TE:KRAB-ZNF in human primary and secondary cortices. (A) and **(B)** demonstrate the workflow for checking significant TE:KRAB-ZNF using the primary and secondary cortices as an example. **(A)** We used random select gene sets and KRAB-ZNFs to calculate correlations with TEs. The violet dots indicate the correlation counts of TE:KRAB-ZNF based on comparing all correlations, positive correlations and negative correlations. They are significantly higher than random gene sets (boxplots below, 1000 iterations, *p*<0.001). **(B)** Finding overlaps between TE:KRAB-ZNF and the KRAB-ZNF protein ChIP-exo data (Imbeault et al., 2017). The overlapped counts are labeled on the y-axis and x axis is the correlation coefficient. Note that we use absolute coefficient values for negative correlations. Samples under the yellow area are selected **(C)** Jaccard similarity is used to demonstrate the correlations results were significantly overlapped (higher) with ChIP-exo data than random selected TE and KRAB-ZNF (*p* < 0.001). The points indicated the overlap with actual correlations and ChIP-exo data. The boxplots indicate the overlap between randomly selected TE and KRAB-ZNF pairs with ChIP-exo data. **(D)** Subsets of the number of positive and negative TE:KRAB-ZNF in the primary and secondary cortices (c1_p and c1_n) and the limbic and association cortices (c2_p and c2_n). **(E)** TE:KRAB-ZNF network in the primary and secondary cortices with five modules. Nodes are colored in five colors representing five modules and nodes in white do not belong to any module. Young links are in orange and old links are in blue. **(F)** Distribution of the normalized degree counts in TE and KRAB-ZNF nodes in TE:KRAB-ZNF network. **(G)** This is the subnetwork colored in pink from **(E)**, showing this module mainly consisted of Alu subfamilies. **(H)** The log count of correlations classified by TE subfamilies and the categories of link, including positive-old (P-O), positive-young (P-Y), negative-old (N-O) and negative young (N-Y). Red asterisks indicate that the class distribution of the TE subfamily is significantly different (Chi-squared test, *p*<0.001). The right-hand side barplot shows the exact count of Alu subfamilies from the first row in the heatmap.

Next, we compared the TE:KRAB-ZNF between species and only considered the 178 KRAB-ZNFs and 836 TEs, which were expressed in all four species. Since the original study (Khrameeva et al., 2020) included four human individuals but only three individuals per NHP species, we performed a leave-one-out analysis of the human

samples. For a fair comparison across species, we required that a correlation between KRAB-ZNFs and TEs in humans needed to be significantly detected in all human leave-one-out combinations (**Figure 4A** and **Table S5**). We then repeated our test of whether KRAB-ZNFs are more likely to correlate with TEs compared to randomly selected genes, and found that human KRAB-ZNF still have a significantly higher number of correlations with TEs. In contrast, in NHPs, KRAB-ZNFs did not have more correlations to TEs than randomly selected genes (**Figure S6**).

We subsequently determined human-specific and conserved TE:KRAB-ZNF interactions by assessing whether an interaction seen in humans also existed in any NHP. Remarkably, the results we obtained show that humans have a higher number of correlations and connectivity compared to NHPs (**Figure 4B**). This finding is not confounded by lack in gene/TE annotations nor sample size, given that we only included KRAB-ZNF genes and TEs expressed in all four species, and that we used the same number of individuals per species; being even more conservative by requiring that all four permutations of human triples show a significant correlation.

Interestingly, we found that some correlations had opposite signs between species. For example, 276 TE:KRAB-ZNF with positive correlations in humans were negatively correlated in bonobos (**Figure 4B**). Although not significant, out of the 276 TE:KRAB-ZNF, 200 in chimpanzees and 132 in rhesus macaques are also negative. These results represent human-specific changes in the sign of the correlation. Constructing a network of those 276 TE:KRAB-ZNF we detected that *ZNF112* and *ZNF528* are two hubs connected to TEs with mostly old links and young links, respectively (**Figure 4C**). We asked whether sequence differences exist between the orthologous zinc finger proteins that might explain this functional change. For ZNF112, there were 21 zinc finger domains. However, among 14 amino acid differences, none

of them affected a position within the zinc finger domains (-1, 3, and 6) that directly contacts the DNA. In the orthologous ZNF528 between human and bonobo, we discovered an amino acid difference in one position that directly contacts nucleotides of the DNA (-1 position of the 15th zinc finger domain) (**Figure 4D**). While in bonobo's ZNF528 there is Histidine at this position, it is a Glutamine in human ZNF528. For other KRAB-ZNF proteins it has been demonstrated that replacing DNA-contacting amino acids to Alanine or Glutamine reduces their repressor potency. Therefore, we speculate that variations in the zinc finger domain of ZNF528, specifically at positions Gln-588 and His-588, may explain why human ZNF528 is not as negatively correlated with TEs as bonobo ZNF528 (**Figure 4D**). Interestingly, this changed position represents a very rare human polymorphism (rs373201614), which seems to be under positive selection in humans including Denisovans and Neanderthals (Harrison et al., 2024).

The distribution of young and old links was not random in the 276 TE:KRAB-ZNF bipartite network (**Figure 4C**). For instance, TcMar-Tc2 and Gypsy had only old interaction with KRAB-ZNFs, while most of the ERVK correlations were classified as young (**Figure 4E**). Last, we checked the species-specific correlations based on TE subfamilies. For example, there are 24,063 positive and 2,455 negative correlations in humans that were not detected in NHPs (the first column and the fifth column in **Figure 4B**). Interestingly, these human-specific correlations were all evolutionary young links and many of them represented negative correlations involving TEs of the Alu subfamilies (**Figure 4F**).

Figure 4. Comparison of TE:KRAB-ZNF in human and NHPs. (A) Workflow for selecting TE:KRAB-ZNF comparing between species. First, there were 178 TEs and 836 KRAB-ZNFs detected in all four species. Second, the leave-one-out test in the human sample was performed for a fair comparison since humans had four repeats and NHPs only had three. (adjusted *p* < 0.01 and absolute coefficient > 0.4) **(B)** Number of positive and negative correlations in human and NHPs. Red brackets indicated the change of correlation between two species. For example, there are 276 human positive correlations which are negatively correlated in bonobos (suffix n: negative, p: positive; hs: *Homo sapiens*, pt: *Pan Troglodytes*, pp: *Pan paniscus*, mm: *Macaca mulatta*). **(C)** 276 TE:KRAB-ZNF network that were all positive correlations in humans but negatively correlated in bonobos. This network demonstrates two hubs, *ZNF528* and *ZNF112*, connecting to multiple TE subfamilies. Node size of TEs refers to the relative abundance of connections to the hubs. Details of this network can be found in **Figure S6**. **(D)** ZNF528 protein sequence difference in zinc finger domain (ZF) comparing humans with bonobos. Comparing humans and bonobos at the -1 finger position, humans have Glutamine (Q) while bonobos have Histidine (H). The lower part of the illustration indicates that the zinc finger domain binds to the DNA sequence using the -1, 3, and 6 finger positions. **(E)** Number of different TE subfamily nodes which form evolutionary old and young correlations in **(C)** network comparing humans to bonobos. **(F)** Distribution counts of human-specific correlations categorized based on TE subfamilies showing only young links. N-Y: negative and young correlations; P-Y: positive and young correlations.

Alterations in TE and KRAB-ZNF expression in brains of Alzheimer patients: Variance in TE and KRAB-ZNF gene expression reflects the brain regions differences

Several KRAB-ZNFs and TEs with human-specific expression patterns or correlations have been associated with AD. For example, *ZNF267*, which was upregulated in the human cortex compared to NHPs (**Figure 2E**), is a clear transcriptomic signature for the diagnosis of AD (Fehlbaum-Beurdeley et al., 2012), and AluYa5 expression leads to genetic dysregulation in AD (Kim et al., 2016). We thus hypothesized that the regulatory network of KRAB-ZNFs and TEs might be severely altered in the brains of AD patients. We utilized the **Mayo Data** to test this hypothesis (**Table 1**).

To investigate if there were differences in the expression patterns of TEs and KRAB-ZNFs comparing different brain regions and disease status, we conducted t-SNE analysis using their expression levels from the temporal cortex and cerebellum of human control individuals and AD patients. Results showed that variances were primarily influenced by brain regions rather than AD-status (**Figure 5A**). Similar to our previous results with the Primate Brain Data (**Figure 2C**), we found that evolutionary young KRAB-ZNF genes and young TEs were expressed at lower levels than their older counterparts (**Figure 5B**). Comparing expression levels between AD and control samples, we obtained for the temporal cortex 6 KRAB-ZNF genes and 4 TEs that were up-regulated in AD and 6 TEs that were down-regulated in AD (**Figure 5C**). In the cerebellum, there were 6 up-regulated and 1 down-regulated TEs, and 4 down-regulated and 10 up-regulated KRAB-ZNF genes (**Figure 5D**).

Figure 5. Expression of TEs and KRAB-ZNF genes in Mayo Data. (A) Variations in the expression of KRAB-ZNF genes and TEs using t-SNE analysis. cbe: cerebellum; tcx: temporal cortex. **(B)** Distributions of the expression of old (> 44.2 mya) and young (≤ 44.2 mya) KRAB-ZNF genes and TEs expression. **(C)** and **(D)** Differentially expressed KRAB-ZNF genes and TEs (absolute log2FoldChange > 0.5, *p* < 0.05) in temporal cortex (tcx) and cerebellum (cbe). The expression of KRAB-ZNF genes and TEs in cerebellum used the same log expression scale.

Human-specific correlations limited to the healthy human temporal cortex: 21 TE:KRAB-ZNFs not detected in AD

To select significant correlations (TE:KRAB-ZNF) between control and AD samples in the temporal cortex and cerebellum, we employed the same filtering criteria as described in the Primate Brain Data analysis (**Figure 3A-C**), requiring an adjusted *p*-value less than 0.01, absolute correlation coefficients higher than 0.4, and TE:KRAB-ZNF pairs detected in ChIP-exo data (Imbeault et al., 2017). The overlaps of TE:KRAB-ZNF pairs are depicted in **Figure 6A**, demonstrating a higher number of correlations in the control group. Next, we selected 21 TE:KRAB-ZNF, which are detected from the temporal cortex both in human Primate Brain Data and the healthy controls in Mayo

Data, but not detected in any NHPs in Primate Brain Data. These correlations represented a subset of TE:KRAB-ZNF, which were specific to healthy adult human brain samples but not detected in AD progression (**Figure 6B**). Among these 21 TE:KRAB-ZNF, there are 14 evolutionary young and 7 evolutionary old interactions, and we found that Alu subfamilies accounted for 11 of these 21 interactions (**Figure 6C**). We further investigated why these TE:KRAB-ZNF were not detected in AD samples and found that most of the correlations were not significant based on our defined threshold. For instance, AluYc:*ZNF182* exhibited a positive correlation in both control and AD samples in the temporal cortex. However, according to our selection criteria, this TE:KRAB-ZNF pair was not deemed significant in the AD sample (FDR=0.34) (**Figure 6D**). The sole exception, showing opposite direction of correlation between groups, was L1MA6:*ZNF211*, which displayed a negative correlation in the control group but had a non-significant very weak positive correlation in the AD group (**Figure 6D**). This indicates weakening and loss of some correlations between TEs and KRAB-ZNFs in AD.

Subsequently, we conducted a bipartite network analysis using the 21 TE:KRAB-ZNF pairs specific to the healthy human samples in the temporal cortex. Integrating evolutionary age inference, we identified a module of young Alu families in the temporal cortex characterized by negative correlations to KRAB-ZNFs (**Figure 6E** and **6F**). Interestingly, the negative young TE:KRAB-ZNF pairs in this Alu module are not significantly detected in AD brains, suggesting that the regulation of the involved TEs is lost or at least impaired in the disease condition.

Figure 6. TE:KRAB-ZNF analysis in Mayo Data. (A) Overlaps of TE:KRAB-ZNF between control and AD condition in temporal cortex and cerebellum (denoted as cbe_control, tcx_control, cbe_AD and tcx_AD). **(B)** 21 human-control-specific TE:KRAB-ZNFs were selected from the intersection of human-specific TE:KRAB-ZNFs from Primate Brain Data (not detected in the other NHPs) and control-specific TE:KRAB-ZNFs from Mayo Data in temporal cortex (not detected in AD samples). **(C)** Distribution of TE families counts involve 21 control-specific TE:KRAB-ZNF in the temporal cortex. **(D)** Comparison of the expression and correlation results of AluYc:*ZNF182* and L1MA6:*ZNF211* in the temporal cortex. **(E)** and **(F)** are the bipartite network of 21 TE:KRAB-ZNF in the temporal cortex. **(E)** Coloring TE nodes in green

and KRAB-ZNF nodes in violet. Evolutionary young links are in orange and evolutionary old links are in blue. Orange border specified that this TE or KRAB-ZNF evolved recently. **(F)** There are two modules in the network colored in pink and gray based on their bipartite modularity. Brown links indicate negative correlations and green links are positive correlations.

Discussion

In this study, we conducted systematic examinations of KRAB-ZNFs and TEs expression and correlation patterns in the brain and compared them between humans and NHPs as well as between healthy human samples and AD. We demonstrated high divergence in the expression of KRAB-ZNF genes and TEs between species, suggesting a prominent evolutionary signature in the regulation of KRAB-ZNF genes and TEs in primates (**Figure 2B**). We also found that evolutionary younger members of both, KRAB-ZNF genes and TEs, exhibit significantly lower expression levels compared to older members (**Figure 2C**), proposing that newly emerged genetic elements might be subjected to more stringent regulatory mechanisms, potentially avoiding disruption at the organismal level. To represent the complexity of interactions between KRAB-ZNF and TEs in bipartite networks, we showed that humans had higher connectivity in the co-expressed TE:KRAB-ZNF network compared to the networks of NHPs (**Figure 4B**). This observation is in line with previous findings that humans also have higher connectivity in the transcription factor networks of the brain compared to NHPs (Bakken et al., 2016; Berto & Nowick, 2018; Nowick et al., 2009). A substantial proportion of young TE:KRAB-ZNF seems to be human-specific, pointing to recent additions to the TE:KRAB-ZNF network in the human lineage and presumably an increase in the complexity of gene regulation (**Figure 4F**). The network of the 21 TE:KRAB-ZNF that were only discovered in the healthy human temporal cortex but not in other NHPs (**Figure 6E** and **6F**) indicates potential loss of control of some TE expression in AD brains. These collective findings suggest an important role of the TE:KRAB-ZNF network in shaping the evolution and in maintaining the functionality of

the healthy human brain, and particularly highlight the co-expression involving evolutionary young KRAB-ZNFs and young TEs.

Given the generally acknowledged role of KRAB-ZNF proteins as repressors on TE expressions, we expected to observe many negative correlations between KRAB-ZNFs and TEs. Indeed, many evolutionary young and human specific correlations involving especially TEs of the Alu subfamily are negative. However, we also observed numerous positive correlations between KRAB-ZNFs and TEs. Keeping in mind that correlations do not indicate direction of the causation and can also be caused by indirect relationships and other factors, several plausible explanations could be offered for these positive interactions. Firstly, some TEs can function as *cis*-acting regulatory elements capable of influencing the expression of nearby genes, which have been mentioned to be tissue-specifically co-opted TEs. (Coronado-Zamora & González, 2023; Sentmanat & Elgin, 2012; Wolf et al., 2020). Secondly, it is conceivable that the expression levels of KRAB-ZNF genes themselves are influenced by the presence and activity of TEs within the genome (Pontis et al., 2019; Senft & Macfarlan, 2021). Another perspective is that primate-specific KRAB-ZNFs can bind to gene promoters, regulating gene expression in the primate brain without necessarily targeting TEs (Farmiloe et al., 2020); observed correlations could then stem from indirect regulation. We should note, however, that 19.43% (446/2,296) of positively correlated pairs overlap significantly with the ChIP-exo results, i.e exist in the presence of a KRAB-ZNF protein binding to the TE (**Figure 3C**). Furthermore, we discover all the human-specific correlations are evolutionary young correlations and different TE subfamilies interact with KRAB-ZNFs differently, for example Alu subfamilies are the only TE subfamily which have more negative correlations than positive ones and LTR subfamilies have only positive correlations (**Figure 4F**). Therefore, we encourage future research

focusing on primate evolution to take these factors into account.

TEs have also been observed to be dysregulated in certain diseases. Here, we observed that the variances in expression levels of TEs and KRAB-ZNFs were more distinct across various brain regions than between control and AD samples (**Figure 5A**). This finding points out the challenge of discerning differences between control and AD samples solely through the consideration of differential expression. This is also illustrated by the detection of only 6 and 10 differentially expressed KRAB-ZNFs and TEs, respectively (**Figure 5C**). Upon combining expression data with correlation results and an evolutionary perspective, we determined 21 human-specific TE:KRAB-ZNFs that were not detectable in AD brains (**Figure 6B, 6E,** and **6F**), suggesting considerable alterations of connections in the context of a human-specific disease. By comparing healthy and disease samples, we discovered evolutionary young Alu subfamilies that are down-regulated in AD patients, such as AluYh9 and AluSp subfamily, which have been found to be negatively associated with Tau pathologic burden in the human dorsolateral prefrontal cortex (Guo et al., 2018). The AluY and AluSx subfamily derived Alu-mediated in-frame deletions of the exons 9 and 10 in mutated forms of *PSNE1*, which was associated with early-onset AD (Le Guennec et al., 2017). Alu subfamilies have further been shown to regulate the expression of *ACE* in neurons from Alu insertions (Wu et al., 2013) and to play a role in A-to-I RNA editing during neurogenesis in the central nervous system and mitochondrial homeostasis (Larsen et al., 2018). Thus, we suggest the dysregulation of modules of Alu subfamilies with their connected KRAB-ZNFs in the regulatory network (**Figure 6F**) plays a role in AD progression.

With respect to the arms-race hypothesis (Jacobs et al., 2014), we observed a higher number of negative correlations between young TEs and young KRAB-ZNF

genes in humans compared to NHPs. This suggests increased repression from young

KRAB-ZNFs on young TEs in the human brain (**Figure 4B** and **4F**). However, our

findings only weakly support the arms-race hypothesis. Firstly, we noted that young

TEs exhibit lower expression levels than old TEs (**Figure 2C** and **5B**), which might not

be expected if they had recently escaped repression. Secondly, discerning the specific

time point at which the TE:KRAB-ZNF relationship becomes a potential arms-race is

challenging. Additionally, some young TEs were also negatively correlated with old

KRAB-ZNF genes, leading to weak assortativity regarding age inference. We interpret

these results as suggesting that the interplay between the evolution of TEs and KRAB-

ZNFs is more complex than a straightforward one-to-one relationship.

In response to the need for systematically comparing expression profiles of

orthologous genes and TEs across diverse species or conditions, we developed

TEKRABber, an R Bioconductor package which equips users to effortlessly adapt its

comprehensive pipeline for their analyses. Our pipeline takes care of normalizing

expression data across species from transcriptomic data by extending the usage of the

scaling-based normalization method (Zhou et al., 2019). To optimize computational

efficiency and parallel computing during the calculations of pairwise correlations, we

harnessed the capabilities of RCpp (Eddelbuettel & François, 2011) and doParallel

(Corporation & Weston, 2011). This approach allows for effective scaling,

accommodating larger datasets and leveraging additional computational resources as

needed. Taking these advantages, users can adapt the TEKRABber's pipeline to

analyze selected orthologous genes and TEs from their expression data within an

acceptable time frame. Taking one of the examples in this study, calculating pairwise

correlations of 178 orthologs with 836 TEs (neglecting sample number) can be

executed in approximately 5 minutes on standard hardware (Intel Core i7 2.6GHz,

16GB memory). Nonetheless, it is crucial to recognize that our present approach is constrained by certain limitations. These constraints primarily stem from variations in gene lengths and positions of the same TE across individual genomes, leaving some room for improving the normalization of TE expression levels. We only used the available reference genomes for our analysis; however, it becomes increasingly clear that substantial individual differences in the presence of TEs exist, which are not covered by a single reference genome of a species. Currently, there are emerging methods designed to detect TE expression, including tools for advancements in sequencing resolution like long-read sequencing (Marx, 2023), de novo TE annotations (Orozco-Arias et al., 2023; Storer et al., 2022) and locus-specific expression detection in single-cell RNA-seq analysis (Rodríguez-Quiroz & Valdebenito-Maturana, 2022). These developments hold promise for achieving a more precise depiction of the variations between samples and the reference, ultimately enhancing our understanding of TE-associated expression patterns. Hence, we acknowledge the significant potential of integrating improved TE orthologous information into our method.

Conclusion

In summary, our findings emphasize the intricate network of interactions between KRAB-ZNFs and TEs in both human evolution and a neurodegenerative disease. To achieve a comprehensive understanding of KRAB-ZNFs and TEs functions, it is not enough to only examine expression levels, but network analysis as facilitated by TEKRABber needs to be leveraged. In primate evolution, the human brain exhibits a notably denser TE:KRAB-ZNF network compared to NHPs, particularly for more recently evolved TEs and KRAB-ZNFs. The healthy human brain TE:KRAB-ZNF network contains a distinct module composed exclusively of Alu subfamilies, which is

an evolutionary novelty not observed in AD brains. These insights emphasize the nuanced dynamics of TE:KRAB-ZNF interactions and their relevance in both evolutionary and disease contexts.

Materials and Methods

Primate Brain Data

For comparing humans with NHPs, we used published RNA-seq data (GSE127898) including 422 brain samples from biological replicates that consisted of 4 humans, 3 chimpanzees, 3 bonobos, and 3 macaques (**Table 1**). Samples from each individual were from 33 different brain regions, and total RNA was sequenced on the Illumina HiSeq 4000 system with a 150 bp paired-end sequencing protocol. More details about samples and the preparation steps can be found in **Table S1** and (Khrameeva et al., 2020). RNA-seq FASTQ data on Gene Expression Omnibus (Barrett et al., 2012) were retrieved using NCBI SRA Toolkit v3.0.3.

Mayo Data

The Mayo RNA-seq study (Allen et al., 2016) was utilized to compare human control and Alzheimer's disease samples. Total RNA was sequenced from both control and AD samples collected from the temporal cortex and cerebellum. The reprocessed FASTQ files, which include only control and AD samples in the temporal cortex and cerebellum, were downloaded via Synapse consortium studies (accession: syn5550404). The number of samples can be found in **Table 1** and more detailed, including biological sex, age, and Braak staging is in **Table S3.**

Table1. Datasets

Dataset	Categories (Biological replicates)	Total number of samples
Primate Brain Data (GSE127898)	Human (4)	132
	Chimpanzee (3)	96
	Macaque (3)	96
	Bonobo (3)	98
Mayo Data (syn5550404)	Control-temporal cortex	23
	AD temporal cortex	24
	Control-cerebellum	23
	AD cerebellum	22

Transcriptome analysis

Adapters and low-quality reads were removed from the FASTQ files using fastp v0.12.4 (Chen et al., 2018) with default parameters. The selected FASTQ files from both datasets were then mapped to their respective references downloaded from UCSC Table Browser, including hg38, panTro6, panPan3, and rheMac10 (Karolchik, 2004), using STAR v2.7.10b (Dobin et al., 2013) with the parameters `--outFilterMultimapNmax 100 --withAnchorMultimapNmax 100`. These parameters were chosen to increase the likelihood of capturing TE mapping by allowing for multiple alignments of reads and more loci anchors for mapping. The resulting BAM files were used to quantify counts of TEs and genes using TEtranscripts v2.2.3 (Jin et al., 2015) with the `--sortByPos` parameter to determine the expression levels of genes and TEs. Gene and TE indices were created using UCSC gene annotations and the RepeatMasker track, enabling reads to match with these intervals. If a read was overlapped with both a gene exon and a TE, it would be checked to determine whether it had a unique alignment or multiple locations in the genome. If an annotation existed, the uniquely aligned read was assigned to the gene. Otherwise, it was assigned to the TE. For reads with ambiguous mappings, they were evenly weighted across TE or gene annotations using the Expectation Maximization (EM) algorithm (Dempster et al., 1977). Differentially expressed genes and TEs were quantified using DESeq2 v1.4.4 (Love et al., 2014) utilizing an absolute log2FoldChange threshold greater than 1.5

(adjusted *p* < 0.05). However, the Mayo Data adopted an absolute log2FoldChange threshold greater than 0.5 due to a lower number of detected differentially expressed genes and TEs. Heatmaps and upset plots were visualized using ComplexHeatmap v2.2.0 (Gu, 2022; Gu et al., 2016).

Inferences about the evolutionary age of KRAB-ZNFs and TEs

A total of 337 KRAB-ZNF genes were obtained by referencing the comprehensive KRAB-ZNF catalog (Huntley et al., 2006). The annotation provided by GenTree (Shao et al., 2019) was utilized to infer the evolutionary age of KRAB-ZNF genes. GenTree employed a synteny-based pipeline to date primate-specific protein-coding genes and presented a branch view to depict the origin of these genes. To complement the age inference of KRAB-ZNF genes, which lacked direct dating annotations in GenTree, we referred to the annotations of primate orthologs in 27 primates, including humans (Jovanovic et al., 2021). By incorporating this complementary age information, we obtained the age inference of all 337 KRAB-ZNF genes for our downstream analysis (**Figure S1**). The evolutionary age inference of TEs was taken from Dfam (Storer et al., 2021) by extracting the species information for each TE subfamily (**Figure S2**).

Development of pipeline software, TEKRABber

To compare the expression of orthologous genes and TEs across species, we concatenated steps including normalization, differential expression, and correlation analysis into an R Bioconductor package, TEKRABber. The name was derived from the idea that TEs are bound ("grabbed") by KRAB-ZNF proteins. TEKRABber adapted the scale-based normalization method (Zhou et al., 2019) to normalize orthologous genes for comparison between two species. In brief, the conserved orthologous gene

lengths from two species were selected from Ensembl data (Harrison et al., 2024) and combined with the expression data to find an optimal scaling factor that can normalize the data to achieve a minimization of deviation between empirical and nominal type I error in a hypothesis testing framework. The normalization step for TEs followed a similar concept. However, instead of using orthologous genes, we used the subset of TEs including LTR, LINE, SINE, SVAs, and DNA transposons (as defined by RepeatMasker (Smit et al., 2013)), for which homologues can be found in other species, for scaling to normalize the expression of TEs. After normalization, differentially expressed orthologous genes and TEs were analyzed using DESeq2 v1.4.4 (Love et al., 2014). The one-to-one correlations between selected orthologous genes and TEs from each species were estimated. In this analysis, we used Pearson's correlations with an adjusted *p-value* using the Benjamini-Hochberg correction (Benjamini & Hochberg, 1995) to obtain significant correlations.

Correlation analysis

To obtain significant correlations of KRAB-ZNFs and TEs for down-stream analysis, we used a workflow including the following steps. We first apply Pearson's correlations on one-to-one gene and TE, using their normalized expression levels. Then, we tested whether KRAB-ZNFs statistically had more correlations with TEs than other random selected genes (1000 iterations, p-value < 0.001). Next, we investigate the relationships between the number of correlations and correlation coefficient values. We decide a threshold with an absolute value of 0.4 coefficient with adjusted p-value less than 0.01 from the distribution. Lastly, we cross-validated our results by comparing the pairs found with experimentally determined TE and KRAB-ZNF relationships using ChIP-exo data from human embryonic stem cell line (Imbeault et al., 2017) and testing

different correlations strengths. For notation, we employ TE:KRAB-ZNF to signify a correlation between a TE and a KRAB-ZNF gene. For AluYc:*ZNF441* indicates a significant correlation between the expression of AluYc and *ZNF441*.

Constructing TE:KRAB-ZNF regulatory network

Nodes and edges were first selected from the TE: KRAB-ZNF results and processed into a dataframe object. The network structure was created using RCy3 v2.16 (Tanja Muetze, 2017) to import into Cytoscape v3.10.1 (Shannon et al., 2003) for visualization and editing. To visualize the regulatory networks, we applied the yFiles Organic Layout for an undirected graph by assigning nodes as objects that had repulsive or attractive forces between them (https://www.yworks.com).

Network analysis

TE:KRAB-ZNF networks were analyzed as bipartite networks, consisting of two classes of nodes, KRAB-ZNF genes and TEs, with connections existing only between the two classes and no edges between elements from the same class. Node properties were first checked using a bipartite module in NetworkX software v2.8.4 (Hagberg et al., 2008). These values include bipartite degree centrality, bipartite strength centrality, and bipartite betweenness centrality for each KRAB-ZNF and TE node. The top 5% scores were selected as hubs (bipartite degree and bipartite strength centrality) and gatekeepers (bipartite betweenness centrality). For cluster detection, leading eigenvector community methods (Csárdi et al., 2023) were used to specify unipartite community structures, and then bipartite modularity (Barber, 2007) was calculated with CONDOR v1.1.1 (Platig et al., 2016). To conduct enrichment analysis among clusters, a Fisher exact test using 1 million simulations with a *p*-value < 0.05 was used.

Software and Code Availability

TEKRABber: https://bioconductor.org/packages/release/bioc/html/TEKRABber.html

Source code: https://github.com/ferygood/primateBrain_TEKRABZNF

Acknowledgement

1. We extend our sincere thanks to the data provider and all members of the research team involved in The MayoRNAseq study. We also appreciate the invaluable data resources provided (https://doi.org/10.7303/syn5550404), which have supported the progression of this research.

2. Our sincere thanks go to Vladimir M. Jovanovic, Rebecca S. Saager, Jeong-Eun Költzow, Fatemeh Zebardast, Melanie Sarfert, Marula Mathew, and Vanessa H. Schulmann for their contributions through critical reading and feedback.

Author Contributions

Conceptualization, KN and YC; Methodology, YC and AM; Visualization, YC and AM; Writing, KN, YC and AM; Data Curation, YC; Software Development, YC; Formal Analysis, YC and AM; Supervision, KN; Project Administration, KN; Funding Acquisition, KN. All authors edited and reviewed the manuscript.

Funding

This work supported by funding from the Deutsche Forschungsgemeinschaft awarded to KN (NO 920/8-1) and by the SPP "Die Genomischen Grundlagen Evolutionärer Innovationen (GEvol)" (NO 920/10-1).

References

Allen, M., Carrasquillo, M. M., Funk, C., Heavner, B. D., Zou, F., Younkin, C. S., Burgess, J. D., Chai, H.-S., Crook, J., Eddy, J. A., Li, H., Logsdon, B., Peters, M. A., Dang, K. K., Wang, X., Serie, D., Wang, C., Nguyen, T., Lincoln, S., … Ertekin-Taner, N. (2016). Human whole genome genotype and transcriptome data for Alzheimer's and other neurodegenerative diseases. *Scientific Data*, *3*(1), 160089. https://doi.org/10.1038/sdata.2016.89

Bakken, T. E., Miller, J. A., Ding, S.-L., Sunkin, S. M., Smith, K. A., Ng, L., Szafer, A., Dalley, R. A., Royall, J. J., Lemon, T., Shapouri, S., Aiona, K., Arnold, J., Bennett, J. L., Bertagnolli, D., Bickley, K., Boe, A., Brouner, K., Butler, S., … Lein, E. S. (2016). A comprehensive transcriptional map of primate brain development. *Nature*, *535*(7612), 367–375. https://doi.org/10.1038/nature18637

Barber, M. J. (2007). Modularity and community detection in bipartite networks. *Physical Review E*, *76*(6), 066102. https://doi.org/10.1103/PhysRevE.76.066102

Barrett, T., Wilhite, S. E., Ledoux, P., Evangelista, C., Kim, I. F., Tomashevsky, M., Marshall, K. A., Phillippy, K. H., Sherman, P. M., Holko, M., Yefanov, A., Lee, H., Zhang, N., Robertson, C. L., Serova, N., Davis, S., & Soboleva, A. (2012). NCBI GEO: Archive for functional genomics data sets—update. *Nucleic Acids Research*, *41*(D1), D991–D995. https://doi.org/10.1093/nar/gks1193

Becker, J., Czamara, D., Hoffmann, P., Landerl, K., Blomert, L., Brandeis, D., Vaessen, A., Maurer, U., Moll, K., Ludwig, K. U., Müller-Myhsok, B., Nöthen, M. M., Schulte-Körne, G., & Schumacher, J. (2012). Evidence for the involvement of ZNF804A in cognitive processes of relevance to reading and spelling. *Translational Psychiatry*, *2*(7), e136–e136. https://doi.org/10.1038/tp.2012.62

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, *57*(1), 289–300.

Berto, S., & Nowick, K. (2018). Species-Specific Changes in a Primate Transcription Factor Network Provide Insights into the Molecular Evolution of the Primate Prefrontal Cortex. *Genome Biology and Evolution*, *10*(8), 2023–2036. https://doi.org/10.1093/gbe/evy149

Bodea, G. O., McKelvey, E. G. Z., & Faulkner, G. J. (2018). Retrotransposon-induced mosaicism in the neural genome. *Open Biology*, *8*(7), 180074.

https://doi.org/10.1098/rsob.180074

Bourque, G., Burns, K. H., Gehring, M., Gorbunova, V., Seluanov, A., Hammell, M., Imbeault, M., Izsvák, Z., Levin, H. L., Macfarlan, T. S., Mager, D. L., & Feschotte, C. (2018). Ten things you should know about transposable elements. *Genome Biology*, *19*(1), 199. https://doi.org/10.1186/s13059-018-1577-z

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, *34*(17), i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Colonna Romano, N., & Fanti, L. (2022). Transposable Elements: Major Players in Shaping Genomic and Evolutionary Patterns. *Cells*, *11*(6), 1048. https://doi.org/10.3390/cells11061048

Coronado-Zamora, M., & González, J. (2023). Transposons contribute to the functional diversification of the head, gut, and ovary transcriptomes across *Drosophila* natural strains. *Genome Research*, genome;gr.277565.122v1. https://doi.org/10.1101/gr.277565.122

Corporation, M., & Weston, S. (2011). *doParallel: Foreach Parallel Adaptor for the "parallel" Package* (p. 1.0.17) [Dataset]. https://doi.org/10.32614/CRAN.package.doParallel

Csárdi, G., Nepusz, T., Müller, K., Horvát, S., Traag, V., Zanini, F., & Noom, D. (2023). *igraph for R: R interface of the igraph library for graph theory and network analysis* (v1.5.1) [Computer software]. Zenodo. https://doi.org/10.5281/ZENODO.7682609

Dempster, A. P., Laird, N. M., & Rubin, D. B. (1977). Maximum Likelihood from Incomplete Data Via the *EM* Algorithm. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, *39*(1), 1–22. https://doi.org/10.1111/j.2517-6161.1977.tb01600.x

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics (Oxford, England)*, *29*(1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

Ecco, G., Imbeault, M., & Trono, D. (2017). KRAB zinc finger proteins. *Development*, *144*(15), 2719–2729. https://doi.org/10.1242/dev.132605

Eddelbuettel, D., & François, R. (2011). **Rcpp**: Seamless *R* and *C++* Integration. *Journal of Statistical Software*, *40*(8). https://doi.org/10.18637/jss.v040.i08

Evering, T. H., Marston, J. L., Gan, L., & Nixon, D. F. (2023). Transposable elements and Alzheimer's disease pathogenesis. *Trends in Neurosciences*, *46*(3), 170–172. https://doi.org/10.1016/j.tins.2022.12.003

Farmiloe, G., Lodewijk, G. A., Robben, S. F., Van Bree, E. J., & Jacobs, F. M. J. (2020). Widespread correlation of KRAB zinc finger protein binding with brain-developmental gene expression patterns. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *375*(1795), 20190333. https://doi.org/10.1098/rstb.2019.0333

Fehlbaum-Beurdeley, P., Sol, O., Désiré, L., Touchon, J., Dantoine, T., Vercelletto, M., Gabelle, A., Jarrige, A.-C., Haddad, R., Lemarié, J. C., Zhou, W., Hampel, H., Einstein, R., & Vellas, B. (2012). Validation of AclarusDx™, a Blood-Based Transcriptomic Signature for the Diagnosis of Alzheimer's Disease. *Journal of Alzheimer's Disease*, *32*(1), 169–181. https://doi.org/10.3233/JAD-2012-120637

Groner, A. C., Meylan, S., Ciuffi, A., Zangger, N., Ambrosini, G., Dénervaud, N., Bucher, P., & Trono, D. (2010). KRAB–Zinc Finger Proteins and KAP1 Can Mediate Long-Range Transcriptional Repression through Heterochromatin Spreading. *PLoS Genetics*, *6*(3), e1000869. https://doi.org/10.1371/journal.pgen.1000869

Gu, Z. (2022). Complex heatmap visualization. *iMeta*, *1*(3), e43. https://doi.org/10.1002/imt2.43

Gu, Z., Eils, R., & Schlesner, M. (2016). Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics*, *32*(18), 2847–2849. https://doi.org/10.1093/bioinformatics/btw313

Guo, C., Jeong, H.-H., Hsieh, Y.-C., Klein, H.-U., Bennett, D. A., De Jager, P. L., Liu, Z., & Shulman, J. M. (2018). Tau Activates Transposable Elements in Alzheimer's Disease. *Cell Reports*, *23*(10), 2874–2880. https://doi.org/10.1016/j.celrep.2018.05.004

Hagberg, A. A., Schult, D. A., & Swart, P. J. (2008). Exploring Network Structure, Dynamics, and Function using NetworkX. In G. Varoquaux, T. Vaught, & J. Millman (Eds.), *Proceedings of the 7th Python in Science Conference* (pp. 11–15).

Hancks, D. C., & Kazazian, H. H. (2010). SVA retrotransposons: Evolution and genetic instability. *Seminars in Cancer Biology*, *20*(4), 234–245. https://doi.org/10.1016/j.semcancer.2010.04.001

Harrison, P. W., Amode, M. R., Austine-Orimoloye, O., Azov, A. G., Barba, M., Barnes, I., Becker, A., Bennett, R., Berry, A., Bhai, J., Bhurji, S. K., Boddu, S.,

Branco Lins, P. R., Brooks, L., Ramaraju, S. B., Campbell, L. I., Martinez, M. C., Charkhchi, M., Chougule, K., … Yates, A. D. (2024). Ensembl 2024. *Nucleic Acids Research*, *52*(D1), D891–D899. https://doi.org/10.1093/nar/gkad1049

Hof, A. E. V., Campagne, P., Rigden, D. J., Yung, C. J., Lingley, J., Quail, M. A., Hall, N., Darby, A. C., & Saccheri, I. J. (2016). The industrial melanism mutation in British peppered moths is a transposable element. *Nature*, *534*(7605), 102–105. https://doi.org/10.1038/nature17951

Huntley, S., Baggott, D. M., Hamilton, A. T., Tran-Gyamfi, M., Yang, S., Kim, J., Gordon, L., Branscomb, E., & Stubbs, L. (2006). A comprehensive catalog of human KRAB-associated zinc finger genes: Insights into the evolutionary history of a large family of transcriptional repressors. *Genome Research*, *16*(5), 669–677. https://doi.org/10.1101/gr.4842106

Imbeault, M., Helleboid, P.-Y., & Trono, D. (2017). KRAB zinc-finger proteins contribute to the evolution of gene regulatory networks. *Nature*, *543*(7646), 550–554. https://doi.org/10.1038/nature21683

Iouranova, A., Grun, D., Rossy, T., Duc, J., Coudray, A., Imbeault, M., de Tribolet-Hardy, J., Turelli, P., Persat, A., & Trono, D. (2022). KRAB zinc finger protein ZNF676 controls the transcriptional influence of LTR12-related endogenous retrovirus sequences. *Mobile DNA*, *13*(1), 4. https://doi.org/10.1186/s13100-021-00260-0

Jacobs, F. M. J., Greenberg, D., Nguyen, N., Haeussler, M., Ewing, A. D., Katzman, S., Paten, B., Salama, S. R., & Haussler, D. (2014). An evolutionary arms race between KRAB zinc-finger genes ZNF91/93 and SVA/L1 retrotransposons. *Nature*, *516*(7530), 242–245. https://doi.org/10.1038/nature13760

Jin, Y., Tam, O. H., Paniagua, E., & Hammell, M. (2015). TEtranscripts: A package for including transposable elements in differential expression analysis of RNA-seq datasets. *Bioinformatics (Oxford, England)*, *31*(22), 3593–3599. https://doi.org/10.1093/bioinformatics/btv422

Jovanovic, V. M., Sarfert, M., Reyna-Blanco, C. S., Indrischek, H., Valdivia, D. I., Shelest, E., & Nowick, K. (2021). Positive Selection in Gene Regulatory Factors Suggests Adaptive Pleiotropic Changes During Human Evolution. *Frontiers in Genetics*, *12*, 662239. https://doi.org/10.3389/fgene.2021.662239

Karolchik, D. (2004). The UCSC Table Browser data retrieval tool. *Nucleic Acids Research*, *32*(90001), 493D – 496. https://doi.org/10.1093/nar/gkh103

Khrameeva, E., Kurochkin, I., Han, D., Guijarro, P., Kanton, S., Santel, M., Qian, Z., Rong, S., Mazin, P., Sabirov, M., Bulat, M., Efimova, O., Tkachev, A., Guo, S., Sherwood, C. C., Camp, J. G., Pääbo, S., Treutlein, B., & Khaitovich, P. (2020). Single-cell-resolution transcriptome map of human, chimpanzee, bonobo, and macaque brains. *Genome Research*, *30*(5), Article 5. https://doi.org/10.1101/gr.256958.119

Kim, S., Cho, C.-S., Han, K., & Lee, J. (2016). Structural Variation of *Alu* Element and Human Disease. *Genomics & Informatics*, *14*(3), 70. https://doi.org/10.5808/GI.2016.14.3.70

Larsen, P. A., Hunnicutt, K. E., Larsen, R. J., Yoder, A. D., & Saunders, A. M. (2018). Warning SINEs: Alu elements, evolution of the human brain, and the spectrum of neurological disease. *Chromosome Research*, *26*(1–2), 93–111. https://doi.org/10.1007/s10577-018-9573-4

Le Guennec, K., Veugelen, S., Quenez, O., Szaruga, M., Rousseau, S., Nicolas, G., Wallon, D., Fluchere, F., Frébourg, T., De Strooper, B., Campion, D., Chávez-Gutiérrez, L., & Rovelet-Lecrux, A. (2017). Deletion of exons 9 and 10 of the Presenilin 1 gene in a patient with Early-onset Alzheimer Disease generates longer amyloid seeds. *Neurobiology of Disease*, *104*, 97–103. https://doi.org/10.1016/j.nbd.2017.04.020

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Marx, V. (2023). Method of the year: Long-read sequencing. *Nature Methods*, *20*(1), 6–11. https://doi.org/10.1038/s41592-022-01730-w

Nowick, K., Carneiro, M., & Faria, R. (2013). A prominent role of KRAB-ZNF transcription factors in mammalian speciation? *Trends in Genetics*, *29*(3), 130–139. https://doi.org/10.1016/j.tig.2012.11.007

Nowick, K., Gernat, T., Almaas, E., & Stubbs, L. (2009). Differences in human and chimpanzee gene expression patterns define an evolving network of transcription factors in brain. *Proceedings of the National Academy of Sciences*, *106*(52), 22358–22363. https://doi.org/10.1073/pnas.0911376106

Ordemann, J. M., & Austin, R. N. (2016). Lead neurotoxicity: Exploring the potential impact of lead substitution in zinc-finger proteins on mental health. *Metallomics*, *8*(6), 579–588. https://doi.org/10.1039/C5MT00300H

Orozco-Arias, S., Sierra, P., Durbin, R., & González, J. (2023). *MCHelper automatically curates transposable element libraries across species* [Preprint]. Genomics. https://doi.org/10.1101/2023.10.17.562682

Patel, S., Howard, D., Chowdhury, N., Derieux, C., Wellslager, B., Yilmaz, Ö., & French, L. (2021). Characterization of Human Genes Modulated by Porphyromonas gingivalis Highlights the Ribosome, Hypothalamus, and Cholinergic Neurons. *Frontiers in Immunology*, *12*, 646259. https://doi.org/10.3389/fimmu.2021.646259

Platig, J., Castaldi, P. J., DeMeo, D., & Quackenbush, J. (2016). Bipartite Community Structure of eQTLs. *PLOS Computational Biology*, *12*(9), e1005033. https://doi.org/10.1371/journal.pcbi.1005033

Platt, R. N., Vandewege, M. W., & Ray, D. A. (2018). Mammalian transposable elements and their impacts on genome evolution. *Chromosome Research*, *26*(1–2), 25–43. https://doi.org/10.1007/s10577-017-9570-z

Playfoot, C. J., Duc, J., Sheppard, S., Dind, S., Coudray, A., Planet, E., & Trono, D. (2021). Transposable elements and their KZFP controllers are drivers of transcriptional innovation in the developing human brain. *Genome Research*, *31*(9), 1531–1545. https://doi.org/10.1101/gr.275133.120

Pontis, J., Planet, E., Offner, S., Turelli, P., Duc, J., Coudray, A., Theunissen, T. W., Jaenisch, R., & Trono, D. (2019). Hominoid-Specific Transposable Elements and KZFPs Facilitate Human Embryonic Genome Activation and Control Transcription in Naive Human ESCs. *Cell Stem Cell*, *24*(5), 724-735.e5. https://doi.org/10.1016/j.stem.2019.03.012

Qu, Y., Izsvák, Z., & Wang, J. (2023). Retrotransposon: A versatile player in human preimplantation development and health. *Life Medicine*, *2*(1), lnac041. https://doi.org/10.1093/lifemedi/lnac041

Ravel-Godreuil, C., Znaidi, R., Bonnifet, T., Joshi, R. L., & Fuchs, J. (2021). Transposable elements as new players in neurodegenerative diseases. *FEBS Letters*, *595*(22), 2733–2755. https://doi.org/10.1002/1873-3468.14205

Rodríguez-Quiroz, R., & Valdebenito-Maturana, B. (2022). SoloTE for improved analysis of transposable elements in single-cell RNA-Seq data using locus-specific expression. *Communications Biology*, *5*(1), 1063. https://doi.org/10.1038/s42003-022-04020-5

Schrader, L., & Schmitz, J. (2019). The impact of transposable elements in adaptive evolution. *Molecular Ecology*, *28*(6), 1537–1549. https://doi.org/10.1111/mec.14794

Scott, E. C., Gardner, E. J., Masood, A., Chuang, N. T., Vertino, P. M., & Devine, S. E. (2016). A hot L1 retrotransposon evades somatic repression and initiates human colorectal cancer. *Genome Research*, *26*(6), 745–755. https://doi.org/10.1101/gr.201814.115

Senft, A. D., & Macfarlan, T. S. (2021). Transposable elements shape the evolution of mammalian development. *Nature Reviews Genetics*, *22*(11), 691–711. https://doi.org/10.1038/s41576-021-00385-1

Sentmanat, M. F., & Elgin, S. C. R. (2012). Ectopic assembly of heterochromatin in *Drosophila melanogaster* triggered by transposable elements. *Proceedings of the National Academy of Sciences*, *109*(35), 14104–14109. https://doi.org/10.1073/pnas.1207036109

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N., Schwikowski, B., & Ideker, T. (2003). Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Research*, *13*(11), 2498–2504. https://doi.org/10.1101/gr.1239303

Shao, Y., Chen, C., Shen, H., He, B. Z., Yu, D., Jiang, S., Zhao, S., Gao, Z., Zhu, Z., Chen, X., Fu, Y., Chen, H., Gao, G., Long, M., & Zhang, Y. E. (2019). GenTree, an integrated resource for analyzing the evolution and function of primate-specific coding genes. *Genome Research*, *29*(4), 682–696. https://doi.org/10.1101/gr.238733.118

Shulman, J. M., Chibnik, L. B., Aubin, C., Schneider, J. A., Bennett, D. A., & De Jager, P. L. (2010). Intermediate Phenotypes Identify Divergent Pathways to Alzheimer's Disease. *PLoS ONE*, *5*(6), e11244. https://doi.org/10.1371/journal.pone.0011244

Smit, AFA, Hubley, R & Green, & P. (2013). *RepeatMasker Open-4.0*. http://www.repeatmasker.org

Storer, J., Hubley, R., Rosen, J., & Smit, A. (2022). Methodologies for the De novo Discovery of Transposable Element Families. *Genes*, *13*(4), 709. https://doi.org/10.3390/genes13040709

Storer, J., Hubley, R., Rosen, J., Wheeler, T. J., & Smit, A. F. (2021). The Dfam community resource of transposable element families, sequence models, and genome annotations. *Mobile DNA*, *12*(1), 2. https://doi.org/10.1186/s13100-020-00230-y

Tanja Muetze, G. K. (2017). *RCy3* [Computer software]. Bioconductor.

https://doi.org/10.18129/B9.BIOC.RCY3

Thomas, J. H., & Schneider, S. (2011). Coevolution of retroelements and tandem zinc finger genes. *Genome Research*, *21*(11), 1800–1812. https://doi.org/10.1101/gr.121749.111

Ting, C. N., Rosenberg, M. P., Snow, C. M., Samuelson, L. C., & Meisler, M. H. (1992). Endogenous retroviral sequences are required for tissue-specific expression of a human salivary amylase gene. *Genes & Development*, *6*(8), 1457–1465. https://doi.org/10.1101/gad.6.8.1457

Willemsen, M. H., Fernandez, B. A., Bacino, C. A., Gerkes, E., De Brouwer, A. P., Pfundt, R., Sikkema-Raddatz, B., Scherer, S. W., Marshall, C. R., Potocki, L., Van Bokhoven, H., & Kleefstra, T. (2010). Identification of ANKRD11 and ZNF778 as candidate genes for autism and variable cognitive impairment in the novel 16q24.3 microdeletion syndrome. *European Journal of Human Genetics*, *18*(4), 429–435. https://doi.org/10.1038/ejhg.2009.192

Wolf, G., De Iaco, A., Sun, M.-A., Bruno, M., Tinkham, M., Hoang, D., Mitra, A., Ralls, S., Trono, D., & Macfarlan, T. S. (2020). KRAB-zinc finger protein gene expansion in response to active retrotransposons in the murine lineage. *eLife*, *9*, e56337. https://doi.org/10.7554/eLife.56337

Wu, S.-J., Hsieh, T.-J., Kuo, M.-C., Tsai, M.-L., Tsai, K.-L., Chen, C.-H., & Yang, Y.-H. (2013). Functional regulation of Alu element of human angiotensin-converting enzyme gene in neuron cells. *Neurobiology of Aging*, *34*(7), 1921.e1-1921.e7. https://doi.org/10.1016/j.neurobiolaging.2013.01.003

Xia, B., Zhang, W., Zhao, G., Zhang, X., Bai, J., Brosh, R., Wudzinska, A., Huang, E., Ashe, H., Ellis, G., Pour, M., Zhao, Y., Coelho, C., Zhu, Y., Miller, A., Dasen, J. S., Maurano, M. T., Kim, S. Y., Boeke, J. D., & Yanai, I. (2024). On the genetic basis of tail-loss evolution in humans and apes. *Nature*, *626*(8001), 1042–1048. https://doi.org/10.1038/s41586-024-07095-8

Zhang, Y. E., Landback, P., Vibranovski, M. D., & Long, M. (2011). Accelerated Recruitment of New Brain Development Genes into the Human Genome. *PLoS Biology*, *9*(10), e1001179. https://doi.org/10.1371/journal.pbio.1001179

Zhou, Y., Zhu, J., Tong, T., Wang, J., Lin, B., & Zhang, J. (2019). A statistical normalization method and differential expression analysis for RNA-seq data between different species. *BMC Bioinformatics*, *20*(1), 163. https://doi.org/10.1186/s12859-019-2745-1