Vole genomics links determinate and indeterminate growth of teeth

AUTHORS:

Zachary T. Calamari[1,2,3,4,*], zachary.calamari@baruch.cuny.edu

Andrew Song[1,5], ajs557@cornell.edu

Emily Cohen[1,6], ec4744@nyu.edu

Muspika Akter[1], muspika.akter@baruchmail.cuny.edu

Rishi Das Roy[7], rishi.dasroy@helsinki.fi

Outi Hallikas[7], outi.hallikas@helsinki.fi

Mona M. Christensen[7], mona.christensen@helsinki.fi

Pengyang Li[3,8], pengyang.li@cshs.org

Pauline Marangoni[3,8], pauline.marangoni@cshs.org

Jukka Jernvall[7,9], jernvall@fastmail.fm

Ophir D. Klein[3,8,*] ophir.klein@cshs.org

[1]Baruch College, City University of New York, One Bernard Baruch Way, New York, NY 10010, USA

[2]The Graduate Center, City University of New York, 365 Fifth Ave, New York, NY 10016, USA

[3]Program in Craniofacial Biology and Department of Orofacial Sciences, University of California, San Francisco, San Francisco, CA 94158, USA

[4]Division of Paleontology, American Museum of Natural History, Central Park West at 79th Street, New York, NY, 10024, USA

[5]Cornell University, 616 Thurston Ave, Ithaca, NY 14853, USA

[6]New York University College of Dentistry, 345 E 34[th] St, New York, NY 10010

[7]Institute of Biotechnology, University of Helsinki, FI-00014 Helsinki, Finland

[8]Department of Pediatrics, Cedars-Sinai Guerin Children's, 8700 Beverly Blvd., Suite 2416, Los

Angeles, CA 90048, USA

[9]Department of Geosciences and Geography, University of Helsinki, FI-00014 Helsinki, Finland

*Corresponding authors

ABSTRACT

Continuously growing teeth are an important innovation in mammalian evolution, yet genetic

regulation of continuous growth by stem cells remains incompletely understood. Dental stem

cells responsible for tooth crown growth are lost at the onset of tooth root formation. Genetic

signaling that initiates this loss is difficult to study with the ever-growing incisor and rooted

molars of mice, the most common mammalian dental model species, because signals for root

formation overlap with signals that pattern tooth size and shape (i.e., cusp patterns). Different

species of voles (Cricetidae, Rodentia, Glires) have evolved rooted and unrooted molars that

have similar size and shape, providing alternative models for studying roots. We assembled a *de*

novo genome of *Myodes glareolus*, a vole with high-crowned, rooted molars, and performed

genomic and transcriptomic analyses in a broad phylogenetic context of Glires (rodents and

lagomorphs) to assess differential selection and evolution in tooth forming genes. We identified

15 dental genes with changing synteny relationships and six dental genes undergoing positive

selection across Glires, two of which were undergoing positive selection in species with unrooted

molars, *Dspp* and *Aqp1*. Decreased expression of both genes in prairie voles with unrooted

molars compared to bank voles supports the presence of positive selection and may underlie

differences in root formation. Bulk transcriptomics analyses of embryonic molar development in

bank voles also demonstrated conserved patterns of dental gene expression compared to mice,

with species-specific variation likely related to developmental timing and morphological

differences between mouse and vole molars. Our results support ongoing evolution of dental

genes across Glires, revealing the complex evolutionary background of convergent evolution for

ever-growing molars.

Keywords: Evolution, selection, Glires, molar, root, dental, development, genome, rodent, tooth

DECLARATIONS

Ethics approval: The University of California, San Francisco (UCSF) Institutional Animal Care

and Use Program and the Finnish national animal experimentation board approved protocols for

humane euthanasia and collection of tissues for animals used in this study under protocols

AN189916 (UCSF) and KEK16-021, KEK19-019, and KEK17-030 (University of Helsinki).

Availability of data and materials: The datasets supporting the conclusions of this article are

available in the GenBank repository under the BioProject PRJNA1050237 (genome accession

number JBBHLL000000000) and in the article's additional files.

Competing interests: The authors declare that they have no competing interests.

Funding: This research was supported by National Science Foundation grants CNS-0958379,

CNS-0855217, OAC-1126113, and OAC-2215760 through the City University of New York

High Performance Computing Center at the College of Staten Island; OAC-1925590 through the

MENDEL high performance computing cluster at the American Museum of Natural History;

Academy of Finland to JJ; Doctoral Programme in Biomedicine, University of Helsinki to

MMC; and National Institutes of Health NIDCR R01-DE027620 and R35-DE026602 to ODK.

Authors' contributions: ZTC and ODK designed the study. ZTC and PM performed animal

husbandry. ZTC performed and oversaw tissue sampling, sequencing, genome assembly and

annotation for *Myodes glareolus*. ZTC, AS, EC, and MA performed genome computational

analyses. PL performed qPCR analyses. OH, MMC, RDR, and JJ designed and implemented

RNA sequencing experiments. ZTC wrote and all authors contributed to and approved the

manuscript.

Acknowledgements: The authors thank A. Joo, N. Ahituv, G. Amato, A. Narechania, S. Singh,

A. Scott, and A. Paasch for advice on methods and access to cluster computing resources.

INTRODUCTION

Hypselodonty, or the presence of unrooted and thus ever-growing teeth, has evolved

multiple times in mammals. Glires—the clade containing rodents, rabbits, and their relatives—

have hypselodont incisors (1), and multiple Glires have also evolved hypselodont molars (Fig.

1). At least in rodents, molar hypselodonty evolved considerably later than hypsodont molars,

which are high crowned but rooted, which in turn evolved later than hypselodont incisors. In

Glires, molars appear to increase in crown height from brachydonty (low-crowned, rooted),

through hypsodonty (high-crowned, rooted), toward hypselodonty (high-crowned, unrooted) (2).

Mice (*Mus musculus*), the primary mammalian model species of dental research, have

hypselodont incisors but retain brachydont molars. Because of this, mice cannot provide

information about the hypsodont teeth that likely preceded hypselodonty.

Mammalian teeth sit in bony sockets, held in place by soft tissue (periodontal ligament)

attached to cementum-covered tooth roots (3). Ligamentous tooth attachment may have arisen

along with a reduction in the rate of tooth replacements, providing greater flexibility for

repositioning the teeth as the dentary grows (4,3). Consequently, the limited replacement of

mammalian teeth (two sets of teeth in most mammals and one in Glires) may have spurred the

evolution of hypsodont and hypselodont teeth, both with high crowns that compensate for tooth

wear from gritty or phytolith-heavy diets (5,6), and resulted in further modification of the

anchoring roots. The convergent evolution of unrooted molars in Glires presents an opportunity

to identify whether consistent developmental and genomic changes underlie the formation of

hypselodont teeth in different species, in turn revealing the conserved mechanisms that produce

tooth roots. Furthermore, the relatively recent evolution of molar hypselodonty, starting in the

Middle Miocene (approximately 16-12 Ma) (2), should provide molecular evidence for the steps

required to make a continuously growing organ.

Dental development proceeds from the tooth germ, composed of epithelium and

mesenchyme, through phases known as the bud, cap, and bell (7). Multipotent enamel epithelium

differentiates into the cells that form the tooth crown (8–11). As development progresses in

rooted teeth, the epithelium at the tooth apex transitions first to a tissue called Hertwig's

epithelial root sheath, and eventually cementum-covered roots (9,10). Studies have identified

numerous candidate genes and pathways with various roles during root development, such as

Fgf10, which decreases in expression at the beginning of root formation (12–18). Although

research on mouse molars has identified genetic signals related to root formation, a number of

the key genes studied have broad developmental roles, such as *Wnt* family members (14), or

overlap considerably with genes also involved in patterning the size and shape of the tooth

(17,19–22). This overlap between shape and root expression patterns confounds our ability to

identify a clear signal initiating root formation.

Evolutionary novelties such as high-crowned hypsodont and hypselodont molars can

arise from differences in gene expression and regulation (23–26). Evolutionarily conserved gene

expression levels produce conserved phenotypes, and changes in gene regulatory networks have

long been linked to morphological evolution (27,28). The order of genes along a chromosome

(synteny) can affect gene expression and regulation, as regulatory sequences are often located

near their target genes (cis-regulatory elements) (29–31). Genome rearrangements that place

genes near new regulatory elements may change the expression and selective environment of

those genes; these small-scale rearrangements of genes may be common in mammals (32–34).

Likewise, regions of chromosomes that form topologically associated domains may experience

similar selective pressures, including selection against rearrangement (35,36). Genes involved in

molar development are not syntenic in the mouse genome nor are genes with organ-specific

expression (37), and thus the regulatory or selection effects of co-localization need not apply to

all dental genes at once. Changes in genome architecture between Glires species thus may result

in different selective and expression environments for dental genes that could result in the

evolution of hypselodont molars.

To establish a model rodent species with hypsodont molars for close comparison to

hypselodont molars, we sequenced and annotated a highly-complete *de novo* genome of *Myodes*

glareolus, the bank vole. The bank vole is increasingly used in medical and environmental

research, ranging from studying zoonotic diseases (38) to immune responses (39,40), and even

assessing environmental remediation efforts through heavy metals that accumulate in vole teeth

(41,42), thus our efforts may be of use beyond dental research. The bank vole's hypsodont

molars bridge the gap between low-crowned mouse and hypselodont prairie vole (*Microtus*

ochrogaster) molars, reducing the effects of morphological differences on root formation

signaling. We performed a suite of genomic and transcriptomic tests of our new bank vole

genome in a broad phylogenetic context to test the hypothesis that dental genes are undergoing

positive selection and exhibit different expression patterns in species with unrooted, hypselodont

molars. We predicted that genes without conserved syntenic relationships in these species would

be more likely to have sites under positive selection or significantly different expression. Our

analyses revealed loss of synteny and positive selection for dental genes in Glires with unrooted

molars compared to those with rooted molars. We also demonstrated strong conservation of

dental gene expression patterns between bank voles and mice, with key differences related to the

timing and patterning of tooth morphology.

RESULTS

Orthology assessment and loss of synteny

To identify which sequences in our bank vole (*Myodes glareolus*) genome and annotation

had the same evolutionary history as dental genes identified in other Glires and assess genome

rearrangements, we performed orthology and synteny analyses in a broad phylogenetic context.

OrthoFinder identified 20,547 orthogroups representing 97.9% of the genes across all 24

analyzed genomes (including the human outgroup). Of the orthogroups, 6,158 had all species

present. In our *de novo* bank vole genome, there were 27,824 annotated genes, of which 84.2%

were assigned to an orthogroup. Bank vole genes were present in 16,250 orthogroups. On

average, the genomes included in the OrthoFinder analysis had 19,814 genes, with 98.2% of

those assigned to orthogroups.

The completeness and large scaffold N50 (4.6 Megabases) of our bank vole assembly

supported its inclusion in generating a Glires synteny network. Using the infomap clustering

algorithm, we produced 19,694 microsynteny clusters from this overall synteny network. We did

not expect dental genes to share the same microsynteny cluster, and instead examined whether

each gene was in the same microsynteny cluster in species with rooted or unrooted molars. We

identified 15 hierarchical orthogroups in which synteny was not conserved for at least half of the

Glires with unrooted molars (Fig. 2). The genes form two groups (Fig. 2A), group 1, lacking

synteny across Glires, and group 2, lacking synteny mainly in species with unrooted molars.

Most of these genes also are missing from the orthogroups; only *Mmp20*, *Irx6*, *Aqp3*, *Sema3b*,

and *Col4a1* were well represented in their orthogroups but not in their synteny networks (full

comparisons of orthology and synteny are in Additional information 1). Overall, these genes

represent multiple categories of "keystone" dental genes. Null mutations in keystone dental

genes affect embryonic dental development (43): "shape" genes cause morphological errors;

"eruption" genes prevent tooth eruption; "progression" genes stop the developmental sequence;

"tissue" genes cause defects in tissues; "developmental process" genes are annotated with the

"GO:0032502" gene ontology term; "dispensable" genes, while dynamically expressed in

developing teeth, have no documented effect on phenotype; and "double" genes function

redundantly with a paralog and only produce a phenotype when both genes are mutated. The

group "other" is composed of the remaining protein coding genes (43). Most genes lacking

conserved synteny in species with unrooted molars are in the "dispensable" category (Fig. 2D),

thus the relationship between differences in these genes and tooth phenotypes is unclear, at least

during embryonic development.

Multiple dental genes under positive selection

We hypothesized that dental genes are undergoing positive selection in species with

unrooted molars. Our positive selection analyses in PAML (phylogenetic analysis by maximum

likelihood (44)) identified 6 dental gene orthogroups undergoing site-specific positive selection

across Glires (Table 1). Four orthogroups with site-specific positive selection lacked synteny

among Glires with unrooted molars: *Col4a1*, *Dspp*, *Runx3,* and the four-gene orthogroup with

sequences similar to *Runx3* (Fig. 2A). We then assessed genes for site-specific positive selection

in species with unrooted molars compared to species with rooted molars (branch-and-site-

specific positive selection (45)), focusing on those genes with site-specific positive selection or

evidence for loss of synteny. Two genes, *Dspp* and *Aqp1* were undergoing this branch-and-site

specific positive selection. Both genes had a single highly supported site (posterior probability >

0.95) under positive selection in species with unrooted molars based on the Bayes Empirical

Bayes method for identifying sites under selection implemented in PAML (46). *Dspp* also had

multiple sites with moderate support (posterior probability > 0.75). The overall selection patterns

on each gene differed. Maximum likelihood estimates of selection for *Dspp* showed the

percentage of sites under purifying and neutral selection on all branches were nearly equal (47%

and 44%, respectively). Percentages of sites under positive selection in the species with unrooted

molars (foreground branches) were nearly evenly divided as well, with 5% of sites from

branches where the species with rooted molars (background branches) were undergoing

purifying selection and 4% of sites from branches where the species with rooted molars were

under neutral selection. For *Aqp1*, nearly all sites were under purifying selection on all branches

(91%), and few sites were under neutral selection on all branches (7%). Few sites were

undergoing positive selection in the foreground branches and their distribution also was unevenly

split between sites under purifying and neutral selection on background branches (0.6% and

0.04%, respectively). The complete list of dental genes with hierarchical orthogroups,

microsynteny clusters, and positive selection test results are available in Additional file 1.

Because genes under positive selection are often expressed at lower levels than genes

under purifying selection (47–50), we also compared expression levels of *Dspp* and *Aqp1* in first

molars (M1) at postnatal days 1, 15, and 21 (P1, P15, and P21) in bank voles (rooted molars) and

prairie voles (unrooted molars) using quantitative PCR. Prairie vole molars expressed *Aqp1* at

significantly lower levels than bank vole molars across all three ages (Fig. 3). Prairie vole P1

molars expressed significantly lower levels of *Dspp* than bank vole molars; at P15 and P21, their

molars expressed *Dspp* at lower, but not statistically significantly different, levels than their bank

vole equivalent. For both genes, the prairie vole had consistent expression levels across three

biological replicates, while the bank vole had greater variation in expression levels across

replicates.

Few changes of secondary structure at positively selected sites

To detect whether substitutions at sites under positive selection influenced protein

structure and evolution, we analyzed ancestral states and secondary structure across Glires. We

first reconstructed ancestral sequences along the internal nodes of the Glires phylogeny for the

genes undergoing branch-and-site specific positive selection to assess potential secondary

structural changes in their protein sequences. At the best-supported site in *Dspp* (position 209 in

the gapped alignment, Additional file 2), there were three major amino acid changes. The

ancestral Glires sequence started with an asparagine (N) in this position. Two of the three species

with unrooted molars represented in the *Dspp* dataset had amino acid substitutions at this

position, with *Oryctolagus cuniculus* substituting a leucine (L) and *Dipodomys ordii* substituting

an aspartic acid (D) at this position (Fig. 4A). All muroids (the clade including the voles in

family Cricetidae and mice and rats in family Muridae) in our phylogeny substituted histidine

(H) for the asparagine at this position. The secondary structure predicted at this position was a

coil for most sequences but a helix for the *D. ordii* sequence (Fig. 5). *Aqp1* sequences varied

greatly at the position under putative positive selection in species with unrooted molars (position

294 in the gapped alignment, Additional file 3). The ancestral state reconstruction showed twelve

changes of the amino acid at this position across Glires (Fig. 4B), yet these changes did not

affect the predicted secondary structure of the protein near this residue, which was a coil for all

sequences tested. All secondary structure predictions are available in Additional file 4.

Bank vole molar gene expression is similar to that of other Glires

We also assessed differential gene expression between mouse and bank vole molars

across early development to study the effects of morphology on expression levels of dental

genes. Our gene expression analysis focused on keystone dental gene categories. Our bank vole

genome was like the mouse and rat genomes in terms of the numbers and expression patterns of

genes annotated from these keystone categories (Table 2). Ordination of gene expression results

from the bank vole and mouse data at embryonic day 13, 14, and 16 (E13, E14, E16) (43) by

principal components analysis showed a distinct separation between the mouse and bank vole

along the first principal component (PC1) of the 500 most variable genes (Fig. 6A). PC1

explained 82.81% of the variance in these genes; there are distinct, species-specific expression

patterns in these tissues. Along PC2 (7.47% of variance explained), E13 and E14 samples differ

from the E16 samples, although the difference in time points is much greater in bank voles.

Ordination of just the keystone dental genes showed clear separations between tissues based on

species and age (Fig. 6B). Within this focused set of genes, however, PC1 and PC2 explain less

variance (44.8% and 28.84% respectively), and have a less clear relationship to species and age.

There are two distinct, parallel trajectories for the mouse and bank vole. Although within each

species there is separation by age along PC1 and PC2, mouse E16 and bank vole E13 occupy a

similar position along PC1, and mouse E13 and bank vole E16 occupy a similar position along

PC2.

Examining individual genes underlying the differences between mouse and vole molars,

we note several upregulated genes in our vole molars are broadly expressed in developing molars

of other vole species (51,52). Relative to the mouse molars, vole molars overexpressed genes

related to forming tooth cusps, including *Bmp2*, *Shh*, *p21* (also known as *Cdkn1a*), and *Msx2*, a

difference explained by the faster patterning and larger number of cusps in the vole molar

compared to the mouse molar (52). Another gene upregulated in the patterning stage vole molar

is *Fgf10*, which is associated with delayed root formation later in vole molar development (9).

Nevertheless, developing bank vole molars at E13, E14, and E16 expressed keystone

dental genes in overall proportions like those observed at analogous stages of mouse and rat

molar development (Fig. 7). Permutation tests within each bank vole sample showed that log

counts for the set of genes related to the progression of dental development were significantly

higher than those in the tissue, dispensable, developmental process, and "other" categories at E14

and E16. The progression gene counts in E13 molars were higher for all of these except the

dispensable category. Shape category genes also were significantly higher than "other" category

genes in the E14 tissue. Overall, even though we observed conserved expression patterns of

dental genes at the system level, individual genes involved in cusp patterning and morphology

differed between the mouse and the vole.

DISCUSSION

Our two goals in sequencing the genome of *Myodes glareolus* were to support the

development of a comparative system for studying tooth root development and to investigate the

evolution of dental genes in Glires, a clade in which ever-growing molars have evolved multiple

times (1). Our new *M. glareolus* assembly and annotation captured nearly all of the single-copy

orthologs for Euarchontoglires and provided scaffolds with sufficient length for synteny

analyses. It was well represented in ortholog groups and microsynteny clusters across Glires. We

tested the hypothesis that dental genes are undergoing site-specific positive selection in species

with unrooted molars (branch-and-site specific positive selection (45)). We predicted that lack of

conserved syntenic relationships in species with unrooted molars could place dental genes in

regulatory and selective environments that promote changes among genes relevant to tooth root

formation. Our analyses identified 15 dental genes without conserved syntenic relationships

across Glires and two dental genes, *Dspp* and *Aqp1*, under positive selection in species with

unrooted molars. We also demonstrated conserved patterns of gene expression among dental

keystone genes between bank voles and mice during early embryonic development, and

deviations from these conserved patterns likely related to differences in molar morphology

between the two species.

We identified 15 genes which were not syntenic in at least half of the species with

unrooted molars, and six genes undergoing site-specific positive selection across all Glires.

Although four of the orthogroups with site-specific positive selection lacked synteny in species

with unrooted molars, only *Col4a1* was well represented among these species in its orthogroup.

The two genes undergoing branch-and-site-specific positive selection in species with unrooted

molars, *Dspp* and *Aqp1*, both maintained their synteny relationships across the Glires studied.

Although we predicted loss of synteny for dental genes in Glires with unrooted molars could

result in sequence evolution by placing genes in new selective contexts, our analyses did not

support a strong relationship between non-syntenic genes and branch-and-site-specific positive

selection. Maximum likelihood estimates of selection on each site for the genes with branch-

specific positive selection revealed different overall selective pressures on *Dspp* and *Aqp1*; *Dspp*

sites on background branches (i.e., branches with species that have rooted molars) were under a

mix of purifying and neutral selection, while nearly all *Aqp1* background branch sites were under

purifying selection. These selection regimes suggest there is greater conservation for *Aqp1*

function across Glires than for *Dspp* function. Gene duplication can result in functional

redundancy and evolution toward a novel function in some genes (53–56), which may explain

positive selection in *Aqp1*, as there are other aquaporin family genes present. Although *Dspp* has

no paralogs, it overlaps functionally with other SIBLING family proteins (e.g., *Opn*, *Dmp1*)

(57,58).

Aqp1 and *Dspp* play different functional roles during dental development. Under the

keystone dental development gene framework, *Aqp1* is a "dispensable" gene: developing teeth

express it, but tooth phenotypes do not change in its absence. *Aqp1* is expressed in endothelia of

microvessels in the developing tooth (59,60). *Dspp* may be particularly relevant for the

formation of an unrooted phenotype if its expression domain or function have been modified in

species with unrooted molars. *Dspp* is a "tissue" category keystone dental gene, meaning the

main effects of a null mutation occur during the tissue differentiation stage of dental

development (43). Null mutations of *Dspp* cause dentin defects in a condition called

dentinogenesis imperfecta (61,62); in some patients, teeth form short, brittle roots (62,63). *Dspp*

knockout mice also exhibit the shortened root phenotype, among a variety of other defects in

both endochondral and intramembranous bone, due to the disruption of collagen and bone

mineralization (64–66).

Our ancestral sequence reconstructions and estimated secondary protein structures

allowed us to assess whether nonsynonymous substitutions at sites under positive selection

resulted in structural differences, thus potentially affecting protein function. Although unrooted

molars are a convergent phenotype across Glires, the sites under positive selection did not

converge on the same amino acid substitution in species with unrooted molars, and *Aqp1*

appeared particularly labile at this residue. The non-synonymous substitutions at these sites often

resulted in changes of properties of the amino acid in the sequence, for example in *Dspp*, polar

asparagine was replaced with non-polar leucine in *O. cuniculus*. Only one of these substitutions

changed the predicted secondary structure. Nevertheless, single amino acid substitutions do

produce dental phenotypes for both *Dspp* (67) and *Aqp1* (68), thus we cannot rule out functional

changes in these genes in species with unrooted molars.

Although the exact relationship between gene expression and sequence divergence

remains unclear (69), studies of genome evolution across small numbers of mammal species

show correlations between gene sequence divergence and levels of expression (70). In particular,

highly-expressed genes are more likely to experience purifying selection (47–50), while lowly-

expressed genes and tissue-specific genes may experience positive selection (48). The decreased

expression of *Dspp* and *Aqp1* in prairie vole M1 compared to that of the bank vole M1 thus

supports our finding of positive selection in these genes in species with unrooted molars. If all

species with unrooted molars also exhibit decreased expression levels of *Dspp* and *Aqp1*, it could

suggest a strong link between lower levels of the genes and the unrooted phenotype.

Without analyses of functional variation caused by positive selection at these coding

sites, or spatial sampling to determine where these genes may be expressed during development,

we are limited from exploring the specific effects of *Dspp* and *Aqp1* on root formation.

Nevertheless, we found evidence for evolution of these genes in Glires with unrooted molars,

and *Dspp* especially has clinical relevance for tooth root formation. Future studies should explore

the spatial distribution of *Dspp* expression, which could be relevant to functional changes in

Glires with unrooted molars. If positive selection and corresponding amino acid changes

identified in *Dspp* here modify its expression domain or its interaction with yet-unidentified root

formation co-factors, it may serially reproduce the unrooted incisor phenotype in molars.

Our RNA sequencing results supported the bank vole as a suitable system for studying

dental development. Although molar morphology differs considerably across mammals,

candidate-gene approaches have identified numerous conserved genes involved in tooth

development and morphological patterning (71). Studies of single genes or gene families have

identified shape-specifying roles common to multiple species (52,72–74), and high-throughput

sequencing of mouse and rat molars demonstrate that both species express sets of dental

development genes in similar proportions during early stages of tooth development (43). The

similarity of our high-throughput RNA sequencing results (Fig. 7) to the mouse and rat results in

previous studies suggest overall expression patterns of keystone dental development genes

within each stage are conserved across Glires. Our principal component analyses and differential

expression analyses measuring changes between mouse and bank vole molars, however, showed

that several dental genes' expression levels differed significantly by species and age. Previous

research has documented organ expression patterns that are conserved across species early in

development and diverge over time, with some major organs displaying heterochronic shifts in

some species (75). If the major source of variation in keystone dental gene expression patterns

between mice and bank vole molars were solely attributable to species, we might expect to see

clear separation between the species along the first or second principal component (PC1 or PC2),

like that observed in PC1 of the 500 most variable genes (Fig. 6). If molar development follows

the diverging expression patterns observed in other organs, we might expect just the earliest age

classes to align on one, or multiple, PCs. Instead, we found two trajectories that were nearly

parallel across PC1 and PC2 and multiple keystone dental genes that were significantly

differentially expressed with respect to species and age. This variation between species is likely

driven by the larger number of cusps in the vole molar, and corresponding upregulation of genes

regulating cusp formation. The overall acceleration of patterning in vole molars likely explains

the significance of the age variable in our expression results, causing a heterochronic shift in the

expression patterns.

Our analyses were limited by the small number of rodent species with sufficiently

annotated genomes to be included in synteny and positive selection analyses. This limitation left

us with a small phylogeny for our ancestral state reconstructions, which thus did not encompass

the full diversity of Glires tooth roots, and potentially weakened model-based genomic analyses.

Although positive selection analyses using the Bayes Empirical Bayes criterion are robust to

smaller sample sizes (46), incomplete sampling can affect estimations of ancestral characteristics

(76). Innovations in paleoproteomics also offer the opportunity to compare fossil species' dental

gene sequences directly to living and estimated ancestral sequences (77,78). By incorporating

data for extinct Glires in both morphological and molecular analyses, we can further elucidate

links between dental gene evolution and unrooted teeth.

CONCLUSIONS

Our genomics and transcriptomics analyses, based on our newly sequenced, high-quality

draft bank vole genome assembly and annotation, showed that bank vole early tooth

development is comparable to other commonly used rodent models in dental development

research. We identified 6 dental gene orthogroups that were undergoing site-specific positive

selection across Glires and two genes, *Dspp* and *Aqp1*, that were undergoing site-specific

positive selection in Glires with unrooted molars. *Dspp* appears particularly relevant to root

formation, as loss-of-function mutations cause a dentin production defect that can result in

shortened tooth roots. Future research must explore the functional role that *Dspp* plays in tooth

root formation in Glires and other clades. The rodent dentary is an exciting system for

understanding tooth development; it provides an easily manipulated set of tissues that can be

produced quickly and features a lifelong population of stem cells in the incisor with genomic

mechanisms that are potentially replicated across other teeth in species with unrooted molars.

Our results identify candidate genes for future analyses, and our draft bank vole genome and

annotation improve the utility of this species for comparative dental research that can uncover

the genetic mechanisms of tooth root formation.

METHODS

Tissue collection and sequencing

To assemble the bank vole genome, we sequenced tissues from a single adult male

specimen housed in a colony at the UCSF Mission Center Animal Facility. We euthanized the

animal according to UCSF IACUC protocol AN189916 and harvested muscle, kidney, heart, and

liver tissue, which were immediately frozen at -80°C. Tissues were sent to a third-party

sequencing service, where they were combined and homogenized to achieve appropriate mass

for high molecular weight DNA extraction. We targeted 60x coverage with 150 base pair (bp)

reads using 10X Chromium linked-read chemistry (79,80) sequenced on the Illumina platform.

We also targeted 10x coverage with Pacific Biosciences SMRT long-read chemistry. For genome

annotation and gene expression analyses, we collected seven biological replicates each of first

molars at embryonic days 13-16 (E13, E14, E15, E16), second molars at E16, and jaw tissues at

E14 under University of Helsinki protocols KEK16-021, KEK19-019, and KEK17-030 and

stored them in RNAlater at -80°C for RNA sequencing, following a tissue harvesting protocol

established for mice and rats (43). We extracted RNA from these tissues using a guanidium

thiocyanate and phenol-chloroform protocol combined with an RNeasy column purification kit

(Qiagen) based on the keystone dental gene protocol (43). Single-end 84 bp RNA sequencing

was performed using the Illumina NextSeq 500 platform.

Genome assembly and quality control

We first assembled only the 10X Chromium linked reads using the default settings in

Supernova 2.1.1. (79,80). We selected the "pseudohaplotype" (pseudohap) output format, which

randomly selects between potential alleles when there are two possible contigs assembled for the

same region. This option produces two assemblies, each with a single resolved length of the

genome sequence (79–81). We used our lower-coverage, long-read data for gap filling and

additional scaffolding. First, we estimated the genome's length using the raw sequence data in

GenomeScope (82), which predicted a length of 2.6 gigabases. We then performed error

correction of the long reads using Canu (83), removing reads shorter than 500 base pairs (bp) and

disregarding overlaps between reads shorter than 350 bp. We kept only those reads with

minimum coverage of 3x for scaffolding. Following long read error correction, we used Cobbler

and RAILS (84) with a minimum alignment length of 200 bases to accept matches for gap filling

and scaffolding of both pseudohap assemblies.

For quality control, we assessed both unscaffolded and long-read scaffolded pseudohap

assemblies by standard assembly length statistics with QUAST (85) and presence of single-copy

orthologs with BUSCO v3 (86). Both scaffolded assemblies were approximately 2.44 Gigabases

long, with an N50 (the length of the shortest scaffold at 50% of the total assembly length) of 4.6

Megabases; we refer to them as Pseudohap1+LR and Pseudohap2+LR. The Pseudohap1+LR

assembly had 17,528 scaffolds over 1000 bp long, and the Pseudohap2+LR assembly had 17,518

scaffolds over 1000 bp long (Table 3). BUSCO searched for universal single-copy orthologs

shared by Euarchontoglires, recovering 89.4% of these genes in the scaffolded Pseudohap1+LR

assembly and 92.8% of the single-copy orthologs in the scaffolded Pseudohap2+LR assembly

(Fig. 8). The two assemblies were similar length and contiguity, but we based annotation and

downstream analyses on Pseudohap2+LR because it recovered more single-copy orthologs.

Genome annotation

We annotated the genome using multiple lines of evidence in three rounds of the

MAKER pipeline (87–89). For evidence from gene transcripts, we assembled a *de novo*

transcriptome assembly of the single-end RNA sequences pooled from all molar and jaw tissues

using Trinity (90). We also included cDNA sequences from the *Mus musculus* assembly

GRCm38 to provide additional transcript evidence from a close relative with a deeply annotated

genome. We used SwissProt's curated protein database to identify protein homology in the

genome. Two libraries of repeats provided information for repeat masking: the Dfam Rodentia

repeat library (91–93) and a custom library specific to the bank vole estimated with a protocol

modified from Campbell et al. (88). The custom library features miniature inverted-repeat

transposable elements identified with default settings in MiteFinder (94), long terminal repeat

retrotransposons extracted with the GenomeTools LTRharvest and LTRdigest functions (95)

based on the eukaryotic genomic tRNA database, and *de novo* repeats identified with

RepeatModeler (96). We combined elements identified by these programs into a single repeat

library, then removed any elements that matched to a custom SwissProt curated protein database

excluding known transposons. The custom repeat library is available in Additional file 5. We

trained a custom gene prediction model for MAKER as well. The first iteration of the model

came from BUSCO's implementation of augustus (97). Between each round of MAKER

annotation, we further updated the gene prediction model with augustus.

MAKER considered only contigs between 10,000-300,000 bp long during annotation.

Our second and third iterations of MAKER used the same settings but excluded the

"Est2genome" and "protein2genome" functions, as recommended in the MAKER tutorial. We

included a SNAP (98) gene prediction model based on the output of the first round of annotation

during the second and third iterations of MAKER annotation. Annotation quality (i.e., agreement

between different lines of evidence and the MAKER annotation) was assessed visually in

JBrowse after each iteration and using *compare_annotations_3.2.pl* (99), which calculates the

number of coding and non-coding sequences in the annotation in addition to basic statistics about

sequence lengths. Our MAKER annotation covered 2.41 Gb of the scaffolded Pseudohap2

assembly in 4,125 scaffolds. These scaffolds contained 27,824 coding genes (mRNA) and 15,320

non-coding RNA sequences. The average gene length was 12,705 bp. Most annotations (91.4%)

had an annotation edit distance (AED) of 0.5 or better. AED is a measure of congruency between

the different types of evidence for an annotation, where scores closer to zero represent better-

annotated genes (100).

Orthology and synteny analyses

We analyzed orthology and synteny of the bank vole genome to understand gene and

genome evolution related to dental development across Glires with rooted and unrooted molars.

We obtained genomes from Ensembl for 23 Glires species and one phylogenetic outgoup, *Homo*

sapiens (Table 4). These genomes all had an N50 over 1 Mb, which improves synteny

assessment (101). We first analyzed all 24 genomes for groups of orthologous genes

(orthogroups) in OrthoFinder (102), providing a tree topology based on the Ensembl Compara

reference tree (Fig. 1) to guide orthology detection. Because we would not analyze the human

outgroup in downstream analyses, we implemented the OrthoFinder option that splits

orthogroups at the root of Glires (hierarchical orthogroups), thus any group of orthologs studied

here represents only genes with shared, orthologous evolutionary history within Glires. We

selected MAFFT (103) for multiple sequence alignment and fastme (104) for phylogenetic tree

searches within OrthoFinder. We retained the gene trees estimated for each orthogroup for

downstream analyses.

Although dental development genes are spread throughout the genome, we were

interested in whether each gene remained in the same local arrangement across species of Glires.

We prepared each genome annotation and sequence file for synteny analysis using the

reformatting functions of Synima (105) to extract each peptide sequence associated with a gene

coding sequence in the Ensembl annotation. Collinear synteny blocks estimated by MCScanX

(106) formed the basis for synteny network analyses using the SynNet pipeline (107–109). We

inferred networks from the top five hits for each gene, requiring any network to have a minimum

of 5 collinear genes and no more than 15 genes between a collinear block, settings that perform

well for analyzing mammal genomes (109). Using the infomap algorithm, we clustered the

synteny blocks into microsynteny networks, from which we extracted network clusters

corresponding to the list of keystone dental genes (43). For each dental gene hierarchical

orthogroup, we assessed whether genes of species with unrooted molars were missing from the

synteny networks that contained other Glires species' sequences, representing loss of synteny for

those species.

Positive selection analysis

We aligned protein sequences for each dental gene orthogroup with clustal omega (110)

using default settings. Based on universal translation tables, we obtained codon-based nucleotide

alignments with pal2nal (111), removing sites in which any species had an indel (i.e., ungapped)

and formatting the output for analysis in PAML (44). We pruned and unrooted the orthogroup

gene trees from OrthoFinder to contain only tips representing the genes in each synteny network

or orthogroup under analysis in PAML. We tested whether any of the genes were undergoing

positive selection using a likelihood ratio test comparing site-specific models of "nearly neutral"

and positive selection. In these models, ω, the ratio of nonsynonymous to synonymous

nucleotide substitutions (also known as dN/dS), can vary at each codon site. In the "nearly

neutral" model, ω can take values between 0 and 1, while the positive selection model allows

sites to assume ω values greater than 1 (46,112). We estimated κ (the ratio of transitions to

transversions) and ω from initial values of 1 and 0.5, respectively, for both tests.

Dental genes with significant site-specific positive selection or those lacking synteny in

species with unrooted molars formed the basis for our second set of positive selection tests using

a branch-and-site model of positive selection. This model allows ω to vary not only among

codon sites, but also between "foreground" and "background" lineages (46). We marked the

species with unrooted molars as foreground lineages, then ran the model twice: once with ω

unconstrained to detect sites undergoing positive selection only on foreground branches, and a

second time and with ω fixed to 1, or neutral selection. A likelihood ratio test of the two models

determined whether the lineage-specific positive selection model was more likely than a neutral

model, and Bayes Empirical Bayes analyses (46) produced posterior probabilities to identify

sites under positive selection.

Genes under positive selection also tend to have lower expression levels (48), thus we

compared expression of the genes with branch-and-site specific positive selection between the

prairie (unrooted molars) and the bank vole (rooted molars) to provide further support for

selective differences. We collected three biological replicates of first molars from both species at

three postnatal stages (P1, P15, and P21) and immediately preserved them at -80°C in lysis

buffer (Buffer RLT; Qiagen) supplemented with 40 µM dithiothreitol. RNA was extracted from

homogenized tissues using a RNeasy column purification kit (Qiagen). We assessed

concentration and purity of extracted RNA using a NanoDrop 2000 spectrophotometer

(ThermoFisher Scientific). Using 1 μg of RNA, we synthesized cDNA using a high-capacity

cDNA reverse transcription kit (ThermoFisher Scientific). We used 1 μL diluted cDNA (1:3 in

ddH$_2$O) and iTaq Universal SYBR Green Supermix (Bio-rad) in the Bio-rad CFX96 real-time

PCR detection system for qPCR experiments, producing three technical replicates for each

biological replicate. We normalized cycle threshold (CT) values of genes of interest to GAPDH

expression levels and calculated relative expression levels as $2^{-\Delta\Delta CT}$. A two-tailed unpaired t-test

calculated in Prism 9 measured whether expression of these genes significantly differed between

bank voles and prairie voles. The oligonucleotide primers for each species and gene are in

Additional file 6.

Sequence and secondary structure evolution

We performed ancestral sequence reconstruction on the codon sequences of the genes

that had evidence of branch-and-site specific positive selection to understand how the sequence

has changed through time. The gapped clustal omega alignments were the basis for ancestral

sequence reconstruction on the Glires species tree (Fig. 1) using pagan2 (113). For each gene, we

plotted amino acid substitutions at the site with potential positive selection. Finally, we predicted

secondary structures (i.e., helices, beta sheets, and coils) for each unrooted species' protein

sequence and the reconstructed ancestral sequence prior to the change at the site under positive

selection using the PSIPRED 4.0 protein analysis workbench (114,115). Comparing these

predictions across the phylogeny, we assessed how these substitutions at the site under selection

may affect the structure of each protein.

Developmental gene expression

We performed quality control and filtering of the short reads for the seven replicates of

first molar tissues at E13, E14, and E16 using the nf-core/rnaseq v. 3.11.2 workflow (116) for

comparability to previous mouse and rat analyses (43). RNAseq reads were evaluated and

adapter sequences were filtered using FastQC v. 0.11.9 (117) and Cutadapt v. 3.4 (118), and

ribosomal RNA was removed using SortMeRNA v. 4.3.4 (119). We then aligned trimmed

sequences to our bank vole annotation using Salmon v. 1.10.1 (120). Counts were then

normalized by gene length. We categorized gene count data into functional groups based on their

established roles in tooth bud development (43) using the one-to-one orthology list between our

bank vole genome and the mouse GRCm39.103 genome annotation generated from our

OrthoFinder output. Using the rlog function of DESeq2 (121), we normalized gene counts within

each functional group on a log2 scale. A permutation test assessed whether the mean counts of

the progression, shape, and double functional groups were significantly different from genes in

the tissue, dispensable, and "other" groups (which are potentially relevant later in development)

based on 10,000 resampling replicates of the dataset (43).

We also assessed differential expression between the bank vole first molar and published

mouse M1 data at the same three time points (GEO accession GSE142199 (43)), combining the

data based on the one-to-one orthology relationships used in the functional permutation analysis.

Using the mouse E13 molar as the reference level, we modeled expression as a response to

species (mouse or vole), embryonic day (E13, E14, or E16), and the interaction between species

and day. We considered as significant any gene with a log fold change greater than 1, log fold

change standard error less than 0.5, and false discovery rate adjusted p value less than 0.05.

TABLES

Table 1 – Genes undergoing site-specific and branch-and-site-specific positive selection

Gene	*Mus* transcript	*Myodes* transcript	Site	Branch-and-site
Aqp1	ENSMUST00000004774	Mglareolus_00011822	Yes	Yes
Col4a1	ENSMUST00000033898	Mglareolus_00032740	Yes	No
Dspp	ENSMUST00000112771	Mglareolus_00014030	Yes	Yes
Fgf20	ENSMUST00000034014	Mglareolus_00013079	Yes	No
Runx3	ENSMUST00000056977	Mglareolus_00033992	Yes	No
similar to *Runx3*	–	–	Yes	–*

598 Table 1 Legend: *HOG only contained four genes with one unrooted species' sequence, could

599 not be tested for branch-and-site specific selection.

601 **Table 2 – P-values of permutation tests between keystone gene categories in bank vole M1**

602 **at embryonic days 13, 14, and 16**

		Tissue	Dispensable	Dev. Process	Other
E13	**Progression**	*0.0310*	0.0942	*0.0436*	*0.0402*
	Shape	0.6431	0.9041	0.2289	0.0995
	Double	0.1292	0.1521	0.0716	0.0655
E14	**Progression**	*0.0136*	*0.0383*	*0.0437*	*0.0401*
	Shape	0.3115	0.4725	0.0922	*0.0454*
	Double	0.1288	0.0945	0.0709	0.0630
E16	**Progression**	*0.0140*	*0.0401*	*0.0303*	*0.0274*
	Shape	0.3770	1	0.1831	0.0662
	Double	0.1343	0.1099	0.0638	0.0596

603 Table 2 Legend: Italicized values are statistically significant ($p < 0.05$)

605   **Table 3 – QUAST assembly statistics for *de novo* bank vole (*Myodes glareolus*) genome**

606   **assemblies**

	Pseudohap1	Pseudohap1+LR	Pseudohap2	Pseudohap2+LR*
Largest contig	27939478	32658832	27937749	32657565
Total length	2434151515	2441426554	2434099357	2441472313
GC (%)	41.88	41.89	41.88	41.89
N50	4187179	4579815	4187179	4558134
N75	1689669	1818134	1687188	1810460
L50	170	153	170	154
L75	388	357	388	358
Ns per 100 kbp	1151.99	1030.75	1151.96	1030.48

607   Table 3 Legend: *assembly used for annotation and downstream analyses in this paper.

609   **Table 4 – Genomes used in orthology, synteny, and positive selection analyses**

Species	Assembly	Citation
Myodes glareolus	CUNY_Mgla_1.0	This paper
Cavia porcellus*	Cavpor3.0	[122]
Cavia aperea*	CavAp1.0	[123]
Marmota marmota	marMar2.1	[124]
Microtus ochrogaster*	MicOch1.0	[125]
Mus musculus	GRCm39	[126]
Oryctolagus cuniculus*	OryCun2.0	[122]
Dipodomys ordii*	Dord_2.0	[122]
Jaculus jaculus	JacJac1.0	[127]

Rattus norvegicus	Rnor_6.0	(128)
Mus pahari	PAHARI_EIJ_v1.1	(129)
Mus caroli	CAROLI_EIJ_v1.1	(129)
Mus spretus	SPRET_EiJ_v1	(130)
Mus spicilegus	MUSP714	(131)
Cricetulus griseus	CHOK1GS	(132)
Mesocricetus auratus	MesAur1.0	(133)
Peromyscus maniculatus	HU_Pman_2.1	(134)
Nannospalax galili	S.galili_v1.0	(135)
Octodon degus*	OctDeg1.0	(136)
Heterocephalus glaber (F)	HetGla_female_1.0	(137)
Chinchilla lanigera*	ChiLan1.0	(138)
Urocitellus parryi	ASM342692v1	(139)
Ictidomys tridecemlineatus	SpeTri2.0	(140)
Homo sapiens**	GRCh38	(141)

610   Table 4 Legend: *Species with unrooted molars; **Peptide annotation used as outgroup only in

611   OrthoFinder analysis.

613   FIGURES

Figure 1 – Species tree of Glires based on the Ensembl Compara species tree. Whether each

species has rooted or unrooted molars is indicated by colored circles at the tip of each branch.

Note that unrooted, or hypselodont, molars have evolved multiple times across Glires. This

topology was the basis for our orthology analysis.

Figure 2 – A Presence (colored boxes) or absence (gray boxes) of gene sequences for each

species in hierarchical orthogroups where fewer than half of the species with unrooted molars

had conserved synteny. Columns are ordered according to phylogenetic positions (top) and rows

are ordered by Euclidean distance clustering. Rows are split into two major groups: group 1, in

which synteny is not conserved across Glires, and group 2, in which synteny is not conserved

mainly in species with unrooted molars. * = One hierarchical orthogroup represented only four

gene sequences annotated based on similarity to *Runx3*. **B** An example of a synteny network for

genes in Group 1, displayed using the Fruchterman-Reingold layout algorithm in the R package

iGraph (142). Small circles represent genes in the synteny network that are not part of the

hierarchical orthogroup, large circles represent genes in the hierarchical orthogroup, and lines

between circles represent a syntenic relationship between two species. Circle color represents

whether species has rooted or unrooted molars following the same key in A. **C** An example

synteny network for genes in Group 2, displayed using the Fruchterman-Reingold layout

algorithm in the R package *iGraph* (142). Circles represent genes in the hierarchical orthogroup,

and lines between circles represent a syntenic relationship between two species. Circle color

represents whether species has rooted or unrooted molars following the same key in A. **D**

Treemaps representing the keystone gene categories for all hierarchical orthogroups, the Group 1

hierarchical orthogroups, and the Group 2 hierarchical orthogroups. Most genes in each group

are in the "dispensable" keystone gene category, which includes genes that are dynamically

expressed during dental development but have no documented effect on phenotypes.

Figure 3 – Quantitative PCR comparisons of *Dspp* and *Aqp1* expression between bank vole and prairie vole M1 at postnatal days 1, 15, and 21 (P1, P15, P21). Expression levels for both genes are lower in the prairie vole (unrooted molars), which supports the positive selection detected for these genes in species with unrooted molars.

646 **Figure 4** – Ancestral state reconstructions of the residue under positive selection in PAML tests.

647 Letters at tips and internal nodes represent IUPAC codes for amino acids and ***** denotes species

648 with unrooted molars. **A** *Dspp*; **B** *Aqp1*.

Figure 5 – PSIPRED secondary structure predictions for the three species with unrooted molars represented in the *Dspp* sequences. Letters correspond to the most recent ancestor of each tip species where the amino acid at the site under positive selection differed: A, the predicted ancestor of *O. cuniculus*; B, the predicted ancestor of *D. ordii*; and C, the predicted ancestor of *M. ochrogaster*. Structure predictions, the relative confidence of the prediction, and the amino acid sequence for each pair of extant species and ancestor are on the right.

Figure 6 – Principal component (PC) analyses of differentially expressed genes in mouse and

bank vole M1. **A** PC1 and PC2 of the 500 most variable genes, showing a clear differentiation

between species along PC1 and differentiation between age classes along PC2. **B** PC1 and PC2

of the keystone dental genes. Both PC1 and PC2 separate age classes within, but not between,

the species, likely due to differences in developmental timing and molar morphology between

mice and voles.

Figure 7 – Box and whisker plots showing normalized log base 2 expression levels for each

keystone gene category in bank vole M1 at embryonic days 13, 14, and 16. Horizontal bar and

diamond within each box represent the median and mean values. Individual datapoints are

displayed for smaller keystone gene categories. Gene expression profiles at these stages are

comparable to mouse and rat molars at analogous developmental stages, as seen in Hallikas et al.

2021.

Figure 8 – BUSCO single-copy ortholog recovery for each "pseudohaploid" version of our draft

bank vole genome assembly and these version after long-read scaffolding (denoted by "+ LR").

Each bar represents the cumulative proportion of the 6,192 single-copy orthologs for

Euarchontoglires identified by BUSCO represented by complete single-copy, complete-

duplicated, fragmented, and missing orthologs. The Pseudohap2 and Pseudohap2 + LR

assemblies had the best single-copy ortholog recovery.

ADDITIONAL FILES

Additional file 1 [.xlsx] Dental gene results – Full table of orthology, synteny, and positive

selection test results for all dental genes assessed.

Additional file 2 [.txt] *Dspp* gapped alignment – Gapped codon-based alignment for *Dspp* in

fasta formatted sequences.

Additional file 3 [.txt] *Aqp1* gapped alignment – Gapped codon-based alignment for *Aqp1* in

fasta formatted sequences.

Additional file 4 [.pdf] Structure predictions – PSIPRED Secondary structure predictions for

each ancestral node and unrooted molar tip species for *Dspp* and *Aqp1*.

Additional file 5 [.txt] Custom repeat library – Custom repeat library of fasta formatted

sequences used in annotation of the draft *Myodes glareolus* genome. See Methods for description

of the process used to generate the library.

Additional file 6 [.pdf] Oligonucleotide primers – List of oligonucleotide primers for *Dspp*,

Aqp1, and *GAPDH* used in bank vole and prairie vole qPCR experiments.

REFERENCES

1. Renvoisé E, Michon F. An Evo-Devo perspective on ever-growing teeth in mammals and

dental stem cell maintenance. Front Physiol. 2014;5(324):1–12.

2.   Tapaltsyan V, Eronen JT, Lawing AM, Sharir A, Janis C, Jernvall J, et al. Continuously

growing rodent molars result from a predictable quantitative evolutionary change over 50

million years. Cell Rep. 2015;11(5):673–80.

3.   LeBlanc ARH, Brink KS, Whitney MR, Abdala F, Reisz RR. Dental ontogeny in extinct

synapsids reveals a complex evolutionary history of the mammalian tooth attachment

system. Proc R Soc B Biol Sci. 2018 Nov 7;285(1890):20181792.

4.   Saffar JL, Lasfargues JJ, Cherruau M. Alveolar bone and the alveolar process: the socket that

is never stable. Periodontol 2000. 1997;13(1):76–90.

5.   Davit-Béal T, Tucker AS, Sire JY. Loss of teeth and enamel in tetrapods: Fossil record,

genetic data and morphological adaptations. J Anat. 2009;214(4):477–501.

6.   Damuth J, Janis CM. On the relationship between hypsodonty and feeding ecology in

ungulate mammals, and its utility in palaeoecology. Biol Rev. 2011;86(3):733–58.

7.   Miletich I, Sharpe PT. Normal and abnormal dental development. Hum Mol Genet. 2003 Apr

2;12(suppl_1):R69–73.

8.   Harada H, Kettunen P, Jung HS, Mustonen T, Wang YA, Thesleff I. Localization of putative

stem cells in dental epithelium and their association with Notch and FGF signaling. J Cell

Biol. 1999;147(1):105–20.

9.   Tummers M, Thesleff I. Root or crown: a developmental choice orchestrated by the

differential regulation of the epithelial stem cell niche in the tooth of two rodent species.

Development. 2003;130(6):1049–57.

10. Thesleff I, Tummers M. Tooth organogenesis and regeneration. In: StemBook. Cambridge,

MA: Harvard Stem Cell Institute; 2008.

11. Krivanek J, Buchtova M, Fried K, Adameyko I. Plasticity of dental cell types in

development, regeneration, and evolution. J Dent Res. 2023 Jun 1;102(6):589–98.

12. Luan X, Ito Y, Diekwisch TGH. Evolution and development of Hertwig's epithelial root

sheath. Dev Dyn. 2006;235(5):1167–80.

13. Kumakami-Sakano M, Otsu K, Fujiwara N, Harada H. Regulatory mechanisms of Hertwig's

epithelial root sheath formation and anomaly correlated with root length. Exp Cell Res.

2014;325(2):78–82.

14. Wen Q, Jing J, Han X, Feng J, Yuan Y, Ma Y, et al. *Runx2* regulates mouse tooth root

development via activation of WNT inhibitor *NOTUM*. J Bone Miner Res.

2020;35(11):2252–64.

15. Yang S, Choi H, Kim TH, Jeong JK, Liu Y, Harada H, et al. Cell dynamics in Hertwig's

epithelial root sheath are regulated by β-catenin activity during tooth root development. J

Cell Physiol. 2021;236(7):5387–98.

16. Yamashiro T, Tummers M, Thesleff I. Expression of bone morphogenetic proteins and Msx

genes during root formation. J Dent Res. 2003;82(3):172–6.

17. Yokohama-Tamaki T, Ohshima H, Fujiwara N, Takada Y, Ichimori Y, Wakisaka S, et al.

Cessation of Fgf10 signaling, resulting in a defective dental epithelial stem cell

compartment, leads to the transition from crown to root formation. Development.

2006;133(7):1359–66.

18. Ota MS, Vivatbutsin P, Nakahara T, Eto K. Tooth root development and the cell-based

regenerative therapy. J Oral Tissue Eng. 2007;4(3):137–42.

19. Jernvall J, Thesleff I. Reiterative signaling and patterning during mammalian tooth

morphogenesis. Mech Dev. 2000;92:19–29.

20. Harada H, Toyono T, Toyoshima K, Yamasaki M, Itoh N, Kato S, et al. FGF10 maintains

stem cell compartment in developing mouse incisors. Dev Camb Engl. 2002;129(6):1533–

41.

21. Tapaltsyan V, Charles C, Hu J, Mindell D, Ahituv N, Wilson GM, et al. Identification of

novel *Fgf* enhancers and their role in dental evolution. Evol Dev. 2016;18(1):31–40.

22. Christensen MM, Hallikas O, Das Roy R, Väänänen V, Stenberg OE, Häkkinen TJ, et al.

The developmental basis for scaling of mammalian tooth size. Proc Natl Acad Sci. 2023

Jun 20;120(25):e2300374120.

23. Chen ZJ. Genetic and epigenetic mechanisms for gene expression and phenotypic cariation

in plant polyploids. Annu Rev Plant Biol. 2007;58(1):377–406.

24. Stranger BE, Forrest MS, Dunning M, Ingle CE, Beazley C, Thorne N, et al. Relative impact

of nucleotide and copy number variation on gene expression phenotypes. Science. 2007 Feb

9;315(5813):848–53.

25. Romero IG, Ruvinsky I, Gilad Y. Comparative studies of gene expression and the evolution

of gene regulation. Nat Rev Genet. 2012 Jul;13(7):505–16.

26. de Montaigu A, Giakountis A, Rubin M, Tóth R, Cremer F, Sokolova V, et al. Natural

diversity in daily rhythms of gene expression contributes to phenotypic variation. Proc Natl

Acad Sci. 2015 Jan 20;112(3):905–10.

27. Erwin DH, Davidson EH. The last common bilaterian ancestor. Development. 2002 Jul

1;129(13):3021–32.

28. Irie N, Kuratani S. Comparative transcriptome analysis reveals vertebrate phylotypic period

during organogenesis. Nat Commun. 2011;2:248.

29. Koonin EV. Evolution of genome architecture. Int J Biochem Cell Biol. 2009 Feb

1;41(2):298–306.

30. Wray GA. The evolutionary significance of cis-regulatory mutations. Nat Rev Genet. 2007

Mar;8(3):206–16.

31. Acemel RD, Maeso I, Gómez-Skarmeta JL. Topologically associated domains: a successful

scaffold for the evolution of gene regulation in animals. WIREs Dev Biol. 2017;6(3):e265.

32. Coghlan A, Eichler EE, Oliver SG, Paterson AH, Stein L. Chromosome evolution in

eukaryotes: a multi-kingdom perspective. Trends Genet. 2005 Dec 1;21(12):673–82.

33. Swenson KM, Blanchette M. Large-scale mammalian genome rearrangements coincide with

chromatin interactions. Bioinformatics. 2019 Jul 15;35(14):i117–26.

34. Long HS, Greenaway S, Powell G, Mallon AM, Lindgren CM, Simon MM. Making sense of

the linear genome, gene function and TADs. Epigenetics Chromatin. 2022 Jan 29;15(1):4.

35. Harmston N, Ing-Simmons E, Tan G, Perry M, Merkenschlager M, Lenhard B.

Topologically associating domains are ancient features that coincide with Metazoan clusters

of extreme noncoding conservation. Nat Commun. 2017 Sep 5;8(1):441.

36. Szabo Q, Bantignies F, Cavalli G. Principles of genome folding into topologically

associating domains. Sci Adv. 2019 Apr 10;5(4):eaaw1668.

37. Das Roy R, Hallikas O, Christensen MM, Renvoisé E, Jernvall J. Chromosomal

neighbourhoods allow identification of organ specific changes in gene expression. PLOS

Comput Biol. 2021 Sep 10;17(9):e1008947.

38. Torelli F, Zander S, Ellerbrok H, Kochs G, Ulrich RG, Klotz C, et al. Recombinant IFN-γ

from the bank vole *Myodes glareolus*: a novel tool for research on rodent reservoirs of

zoonotic pathogens. Sci Rep. 2018;8(1):1–11.

39. Kloch A, Babik W, Bajer A, Siński E, Radwan J. Effects of an MHC-DRB genotype and

allele number on the load of gut parasites in the bank vole *Myodes glareolus*. Mol Ecol.

2010;19(SUPPL. 1):255–65.

40. Migalska M, Sebastian A, Konczal M, Kotlík P, Radwan J. *De novo* transcriptome assembly

facilitates characterisation of fast-evolving gene families, MHC class I in the bank vole

(*Myodes glareolus)*. Heredity. 2017;118(4):348–57.

41. Appleton J, Lee KM, Sawicka Kapusta K, Damek M, Cooke M. The heavy metal content of

the teeth of the bank vole (*Clethrionomys glareolus*) as an exposure marker of

environmental pollution in Poland. Environ Pollut. 2000;110:441–9.

42. Gdula-Argasińska J, Appleton J, Sawicka-Kapusta K, Spence B. Further investigation of the

heavy metal content of the teeth of the bank vole as an exposure indicator of environmental

pollution in Poland. Environ Pollut. 2004;131(1):71–9.

43. Hallikas O, Das Roy R, Christensen MM, Renvoisé E, Sulic AM, Jernvall J. System-level

analyses of keystone genes required for mammalian tooth development. J Exp Zoolog B

Mol Dev Evol. 2021;336(1):7–17.

44. Yang Z. PAML 4: Phylogenetic analysis by maximum likelihood. Mol Biol Evol. 2007 Aug

1;24(8):1586–91.

45. Zhang J, Nielsen R, Yang Z. Evaluation of an improved branch-site likelihood method for

detecting positive selection at the molecular level. Mol Biol Evol. 2005 Dec;22(12):2472–9.

46. Yang Z, Wong WSW, Nielsen R. Bayes Empirical Bayes inference of amino acid sites under

positive selection. Mol Biol Evol. 2005 Apr 1;22(4):1107–18.

47. Drummond DA, Bloom JD, Adami C, Wilke CO, Arnold FH. Why highly expressed proteins

evolve slowly. Proc Natl Acad Sci. 2005 Oct 4;102(40):14338–43.

48. Kosiol C, Vinař T, Fonseca RR da, Hubisz MJ, Bustamante CD, Nielsen R, et al. Patterns of

positive selection in six mammalian genomes. PLOS Genet. 2008 Aug 1;4(8):e1000144.

49. Martincorena I, Luscombe NM. Non-random mutation: The evolution of targeted

hypermutation and hypomutation. BioEssays. 2013;35(2):123–30.

50. Martincorena I, Roshan A, Gerstung M, Ellis P, Van Loo P, McLaren S, et al. High burden

and pervasive positive selection of somatic mutations in normal human skin. Science. 2015

May 22;348(6237):880–6.

51. Keränen SVE, Åberg T, Kettunen P, Thesleff I, Jernvall J. Association of developmental

regulatory genes with the development of different molar tooth shapes in two species of

rodents. Dev Genes Evol. 1998;208(9):477–86.

52. Jernvall J, Keränen SVE, Thesleff I. Evolutionary modification of development in

mammalian teeth: Quantifying gene expression patterns and topography. Proc Natl Acad

Sci. 2000;97(26):14444–8.

53. Hughes AL. The evolution of functionally novel proteins after gene duplication. Proc R Soc

Lond B Biol Sci. 1997 Jan;256(1346):119–24.

54. Wagner A. Selection and gene duplication: a view from the genome. Genome Biol. 2002 Apr

15;3(5):reviews1012.1.

55. David KT, Oaks JR, Halanych KM. Patterns of gene evolution following duplications and

speciations in vertebrates. PeerJ. 2020 Mar 31;8:e8813.

56. Copley SD. Evolution of new enzymes by gene duplication and divergence. FEBS J.

2020;287(7):1262–83.

57. Fisher LW. DMP1 and DSPP: Evidence for duplication and convergent evolution of two

SIBLING proteins. Cells Tissues Organs. 2011 Aug;194(2–4):113–8.

58. Bouleftour W, Juignet L, Bouet G, Granito RN, Vanden-Bossche A, Laroche N, et al. The

role of the SIBLING, Bone Sialoprotein in skeletal biology — Contribution of mouse

experimental genetics. Matrix Biol. 2016 May 1;52–54:60–77.

59. Felszeghy S, Módis L, Németh P, Nagy G, Zelles T, Agre P, et al. Expression of aquaporin

isoforms during human and mouse tooth development. Arch Oral Biol. 2004 Apr

1;49(4):247–57.

60. Yoshii T, Harada F, Saito I, Nozawa-Inoue K, Kawano Y, Maeda T. Immunoexpression of

aquaporin-1 in the rat periodontal ligament during experimental tooth movement. Biomed

Res. 2012;33(4):225–33.

61. Zhang X, Zhao J, Li C, Gao S, Qiu C, Liu P, et al. *DSPP* mutation in dentinogenesis

imperfecta Shields type II. Nat Genet. 2001 Feb;27(2):151–2.

62. de La Dure-Molla M, Philippe Fournier B, Berdal A. Isolated dentinogenesis imperfecta and

dentin dysplasia: revision of the classification. Eur J Hum Genet. 2015 Apr;23(4):445–51.

63. Shields ED, Bixler D, El-Kafrawy AM. A proposed classification for heritable human

dentine defects with a description of a new entity. Arch Oral Biol. 1973 Apr 1;18(4):543-

IN7.

64. Sreenath T, Thyagarajan T, Hall B, Longenecker G, D'Souza R, Hong S, et al. Dentin

Sialophosphoprotein knockout mouse teeth display widened predentin zone and develop

defective dentin mineralization similar to human dentinogenesis imperfecta type III. J Biol

Chem. 2003 Jul 4;278(27):24874–80.

65. Verdelis K, Ling Y, Sreenath T, Haruyama N, MacDougall M, van der Meulen MCH, et al.

DSPP effects on *in vivo* bone mineralization. Bone. 2008 Dec 1;43(6):983–90.

66. Chen Y, Zhang Y, Ramachandran A, George A. DSPP is essential for normal development

of the dental-craniofacial complex. J Dent Res. 2016 Mar 1;95(3):302–10.

67. von Marschall Z, Mok S, Phillips MD, McKnight DA, Fisher LW. Rough endoplasmic

reticulum trafficking errors by different classes of mutant dentin sialophosphoprotein

(DSPP) cause dominant negative effects in both dentinogenesis imperfecta and dentin

dysplasia by entrapping normal DSPP. J Bone Miner Res. 2012;27(6):1309–21.

68. Smith BL, Preston GM, Spring FA, Anstee DJ, Agre P. Human red cell aquaporin CHIP. I.

Molecular characterization of ABH and Colton blood group antigens. J Clin Invest. 1994

Sep 1;94(3):1043–9.

69. Jordan IK, Mariño-Ramírez L, Koonin EV. Evolutionary significance of gene expression

divergence. Gene. 2005 Jan 17;345(1):119–26.

70. Warnefors M, Kaessmann H. Evolution of the correlation between expression divergence

and protein divergence in mammals. Genome Biol Evol. 2013;5(7):1324–35.

71. Jernvall J, Thesleff I. Tooth shape formation and tooth renewal: evolving with the same

signals. Development. 2012;139(19):3487–97.

72. Mitsiadis TA. Role of Islet1 in the patterning of murine dentition. Development.

2003;130(18):4451–60.

73. Charles C, Pantalacci S, Peterkova R, Tafforeau P, Laudet V, Viriot L. Effect of *eda* loss of

function on upper jugal tooth morphology. Anat Rec. 2009;292(2):299–308.

74. Zurowski C, Jamniczky H, Graf D, Theodor J. Deletion/loss of bone morphogenetic protein

7 changes tooth morphology and function in *Mus musculus*: implications for dental

evolution in mammals. R Soc Open Sci. 2018 Jan 3;5(1):170761.

75. Cardoso-Moreira M, Halbert J, Valloton D, Velten B, Chen C, Shao Y, et al. Gene

expression across mammalian organ development. Nature. 2019 Jul;571(7766):505–9.

76. Finarelli JA, Flynn JJ. Ancestral state reconstruction of body size in the Caniformia

(Carnivora, Mammalia): the effects of incorporating data from the fossil record. Syst Biol.

2006;55(2):301–13.

77. Welker F, Collins MJ, Thomas JA, Wadsley M, Brace S, Cappellini E, et al. Ancient proteins

resolve the evolutionary history of Darwin's South American ungulates. Nature. 2015

Jun;522(7554):81–4.

78. Warinner C, Korzow Richter K, Collins MJ. Paleoproteomics. Chem Rev. 2022 Aug

24;122(16):13401–46.

79. Zheng GXY, Lau BT, Schnall-Levin M, Jarosz M, Bell JM, Hindson CM, et al. Haplotyping

germline and cancer genomes with high-throughput linked-read sequencing. Nat

Biotechnol. 2016 Feb;34:303.

80. Marks P, Garcia S, Martinez A, Belhocine K. Resolving the full spectrum of human genome

variation using linked-reads. 2017;

81. Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB. Direct determination of diploid

genome sequences. Genome Res. 2017;27(5):757–67.

82. Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, et al.

GenomeScope: fast reference-free genome profiling from short reads. Bioinformatics. 2017

Jul 15;33(14):2202–4.

83. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and

accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome

Res. 2017 May 1;27(5):722–36.

84. Warren RL. RAILS and Cobbler: Scaffolding and automated finishing of draft genomes

using long DNA sequences. J Open Source Softw. 2016 Nov 17;1(7):116.

85. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome

assemblies. Bioinformatics. 2013 Apr 15;29(8):1072–5.

86. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing

genome assembly and annotation completeness with single-copy orthologs. Bioinformatics.

2015 Oct 1;31(19):3210–2.

87. Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, et al. MAKER: an easy-to-use

annotation pipeline designed for emerging model organism genomes. Genome Res.

2008;18:188–96.

88. Campbell MS, Law M, Holt C, Stein JC, Moghe GD, Hufnagel DE, et al. MAKER-P: A tool

kit for the rapid creation, management, and quality control of plant genome annotations.

Plant Physiol. 2014 Feb 1;164(2):513–24.

89. Campbell MS, Holt C, Moore B, Yandell M. Genome annotation and curation using

MAKER and MAKER-P. Curr Protoc Bioinforma. 2014 Dec 12;48:4.11.1-4.11.39.

90. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Trinity:

reconstructing a full-length transcriptome without a genome from RNA-Seq data. Nat

Biotechnol. 2011 May 15;29(7):644–52.

91. Wheeler TJ, Clements J, Eddy SR, Hubley R, Jones TA, Jurka J, et al. Dfam: a database of

repetitive DNA based on profile hidden Markov models. Nucleic Acids Res. 2013

Jan;41(Database issue):D70-82.

92. Caballero J, Smit AFA, Hood L, Glusman G. Realistic artificial DNA sequences as negative

controls for computational genomics. Nucleic Acids Res. 2014 Jul;42(12):e99.

93. Hubley R, Finn RD, Clements J, Eddy SR, Jones TA, Bao W, et al. The Dfam database of

repetitive DNA families. Nucleic Acids Res. 2016 Jan 4;44(D1):D81–9.

94. Hu J, Zheng Y, Shang X. MiteFinder: A fast approach to identify miniature inverted-repeat

transposable elements on a genome-wide scale. In: 2017 IEEE International Conference on

Bioinformatics and Biomedicine (BIBM). 2017. p. 164–8.

95. Gremme G, Steinbiss S, Kurtz S. GenomeTools: A comprehensive software library for

efficient processing of structured genome annotations. IEEE/ACM Trans Comput Biol

Bioinform. 2013 May 1;10(03):645–56.

96. Smit A, Hubley R. RepeatModeler Open-1.0. 2008.

97. Keller O, Kollmar M, Stanke M, Waack S. A novel hybrid gene prediction method

employing protein multiple sequence alignments. Bioinformatics. 2011 Mar 15;27(6):757–

63.

98. Korf I. Gene finding in novel genomes. BMC Bioinformatics. 2004 May 14;5(1):59.

99. Campbell MS. compare_annotations_3.2.pl [Internet]. 2015. Available from:

https://github.com/mscampbell/Genome_annotation/blob/master/compare_annotations_3.2.

pl

100. Eilbeck K, Moore B, Holt C, Yandell M. Quantitative measures for the management and

comparison of annotated genomes. BMC Bioinformatics. 2009 Feb 23;10(1):67.

101. Liu D, Hunt M, Tsai IJ. Inferring synteny between genome assemblies: a systematic

evaluation. BMC Bioinformatics. 2018 Jan;19(1):26.

102. Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. Genome Biol. 2015

Aug 6;16(1):157.

103. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002

Jul;30(14):3059–66.

104. Lefort V, Desper R, Gascuel O. FastME 2.0: A comprehensive, accurate, and fast distance-

based phylogeny inference program. Mol Biol Evol. 2015 Oct 1;32(10):2798–800.

105. Farrer RA. Synima: a Synteny imaging tool for annotated genome assemblies. BMC

Bioinformatics. 2017 Nov 21;18(1):507.

106. Wang Y, Tang H, DeBarry JD, Tan X, Li J, Wang X, et al. MCScanX: a toolkit for

detection and evolutionary analysis of gene synteny and collinearity. Nucleic Acids Res.

2012 Apr;40(7):e49.

107. Zhao T, Schranz ME. Network approaches for plant phylogenomic synteny analysis. Curr

Opin Plant Biol. 2017 Apr 1;36:129–34.

108. Zhao T, Holmer R, de Bruijn S, Angenent GC, van den Burg HA, Schranz ME.

Phylogenomic synteny network analysis of MADS-Box transcription factor genes reveals

lineage-specific transpositions, ancient tandem duplications, and deep positional

conservation. Plant Cell. 2017 Jun 1;29(6):1278–92.

109. Zhao T, Schranz ME. Network-based microsynteny analysis identifies major differences

and genomic outliers in mammalian and angiosperm genomes. Proc Natl Acad Sci. 2019

Feb 5;116(6):2165–74.

110. Sievers F, Higgins DG. Clustal Omega. Curr Protoc Bioinforma. 2014;48(1):3.13.1-

3.13.16.

111. Suyama M, Torrents D, Bork P. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. Nucleic Acids Res. 2006 Jul

1;34(suppl_2):W609–12.

112. Wong WSW, Yang Z, Goldman N, Nielsen R. Accuracy and power of statistical methods

for detecting adaptive evolution in protein coding sequences and for identifying positively

selected sites. Genetics. 2004 Oct 1;168(2):1041–51.

113. Löytynoja A, Vilella AJ, Goldman N. Accurate extension of multiple sequence alignments

using a phylogeny-aware graph algorithm. Bioinformatics. 2012 Jul 1;28(13):1684–91.

114. Jones DT. Protein secondary structure prediction based on position-specific scoring

matrices. J Mol Biol. 1999 Sep 17;292(2):195–202.

115. Buchan DWA, Jones DT. The PSIPRED protein analysis workbench: 20 years on. Nucleic

Acids Res. 2019 Jul 2;47(W1):W402–7.

116. Ewels PA, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A, et al. The nf-core

framework for community-curated bioinformatics pipelines. Nat Biotechnol. 2020

Mar;38(3):276–8.

117. Andrews S. FastQC: a quality control tool for high throughput sequence data. [Internet].

2010. Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

118. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal. 2011 May 2;17(1):10–2.

119. Kopylova E, Noé L, Touzet H. SortMeRNA: fast and accurate filtering of ribosomal RNAs

in metatranscriptomic data. Bioinformatics. 2012 Dec 1;28(24):3211–7.

120. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-

aware quantification of transcript expression. Nat Methods. 2017 Apr;14(4):417–9.

121. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014 Dec 5;15(12):550.

122. Lindblad-Toh K, Garber M, Zuk O, Lin MF, Parker BJ, Washietl S, et al. A high-resolution

map of human evolutionary constraint using 29 mammals. Nature. 2011

Oct;478(7370):476–82.

123. Weyrich A, Schüllermann T, Heeger F, Jeschek M, Mazzoni CJ, Chen W, et al. Whole

genome sequencing and methylome analysis of the wild guinea pig. BMC Genomics. 2014

Nov 28;15(1):1036.

124. Gossmann TI, Ralser M. *Marmota marmota*. Trends Genet. 2020 May;36(5):383–4.

125. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Microtus ochrogaster. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/72443

126. Mouse Genome Sequencing Consortium, Waterston RH, Lindblad-Toh K, Birney E,

Rogers J, Abril JF, et al. Initial sequencing and comparative analysis of the mouse genome.

Nature. 2002 Dec 5;420(6915):520–62.

127. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Jaculus jaculus. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/72445

128. Gibbs RA, Weinstock GM, Metzker ML, Muzny DM, Sodergren EJ, Scherer S, et al.

Genome sequence of the Brown Norway rat yields insights into mammalian evolution.

Nature. 2004 Apr;428(6982):493–521.

129. Kolmogorov M, Armstrong J, Raney BJ, Streeter I, Dunn M, Yang F, et al. Chromosome

assembly of large and complex genomes using multiple references. Genome Res. 2018 Nov

1;28(11):1720–32.

130. Lilue J, Doran AG, Fiddes IT, Abrudan M, Armstrong J, Bennett R, et al. Sixteen diverse

laboratory mouse reference genomes define strain-specific haplotypes and novel functional

loci. Nat Genet. 2018 Nov;50(11):1574–83.

131. Couger MB, Arévalo L, Campbell P. A high quality genome for *Mus spicilegus*, a close

relative of house mice with unique social and ecological adaptations. G3

GenesGenomesGenetics. 2018 May 24;8(7):2145–52.

132. Chinese hamster CHOK1GS assembly and gene annotation. Horiz Eagle [Internet]. 2017;

Available from: https://www.ensembl.org/Cricetulus_griseus_chok1gshd/Info/Annotation

133. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Mesocricetus auratus. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/77669

134. Lassance JM, Hopi Hoekstra. Improved assembly of the deer mouse *Peromyscus*

maniculatus genome. Harv Univ Hughes Med Inst [Internet]. 2018; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/494228

135. Fang X, Nevo E, Han L, Levanon EY, Zhao J, Avivi A, et al. Genome-wide adaptive

complexes to underground stresses in blind mole rats *Spalax*. Nat Commun. 2014 Jun

3;5(1):3966.

136. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Octodon degu. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/74595

137. Keane M, Craig T, Alföldi J, Berlin AM, Johnson J, Seluanov A, et al. The Naked Mole Rat

Genome Resource: facilitating analyses of cancer and longevity-related adaptations.

Bioinforma Oxf Engl. 2014 Dec 15;30(24):3558–60.

138. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Chinchilla lanigera. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/68239

139. V. Federov, Dalen L, Olsen RA, Goropashnaya AV, Barnes BM. The genome of the Arctic

ground squirrel *Urocitellus parryii.* Inst Arct Biol [Internet]. 2018; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/477386

140. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Ictidomys tridecemlineatus. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/61725

141. Schneider VA, Graves-Lindsay T, Howe K, Bouk N, Chen HC, Kitts PA, et al. Evaluation

of GRCh38 and de novo haploid genome assemblies demonstrates the enduring quality of

the reference assembly. Genome Res. 2017 May 1;27(5):849–64.

142. Csárdi G, Nepusz T, Traag V, Horvát S, Zanini F, Noom D, et al. igraph: Network analysis

and visualization in R [Internet]. 2024. Available from: https://CRAN.R-

project.org/package=igraph

1053