Conserved and derived expression patterns and positive selection on dental genes reveal

complex evolutionary context of ever-growing rodent molars

AUTHORS:

Zachary T. Calamari[1,2,3,4,*], zachary.calamari@baruch.cuny.edu

Andrew Song[1,5], ajs557@cornell.edu

Emily Cohen[1,6], ec4744@nyu.edu

Muspika Akter[1], muspika.akter@baruchmail.cuny.edu

Rishi Das Roy[7], rishi.dasroy@helsinki.fi

Outi Hallikas[7], outi.hallikas@helsinki.fi

Mona M. Christensen[7], mona.christensen@helsinki.fi

Pengyang Li[3,8], pengyang.li@cshs.org

Pauline Marangoni[3,8], pauline.marangoni@cshs.org

Jukka Jernvall[7,9], jernvall@fastmail.fm

Ophir D. Klein[3,8,*] ophir.klein@cshs.org

[1]Baruch College, City University of New York, One Bernard Baruch Way, New York, NY

10010, USA

[2]The Graduate Center, City University of New York, 365 Fifth Ave, New York, NY 10016, USA

[3]Program in Craniofacial Biology and Department of Orofacial Sciences, University of

California, San Francisco, San Francisco, CA 94158, USA

[4]Division of Paleontology, American Museum of Natural History, Central Park West at 79[th]

Street, New York, NY, 10024, USA

[5]Cornell University, 616 Thurston Ave, Ithaca, NY 14853, USA

[6]New York University College of Dentistry, 345 E 34[th] St, New York, NY 10010

[7]Institute of Biotechnology, University of Helsinki, FI-00014 Helsinki, Finland

[8]Department of Pediatrics, Cedars-Sinai Guerin Children's, 8700 Beverly Blvd., Suite 2416, Los

Angeles, CA 90048, USA

[9]Department of Geosciences and Geography, University of Helsinki, FI-00014 Helsinki, Finland

*Corresponding authors

ABSTRACT

Background: Continuously growing teeth are an important innovation in mammalian evolution,

yet genetic regulation of continuous growth by stem cells remains incompletely understood.

Dental stem cells are lost at the onset of tooth root formation, but this loss of continuous crown

growth is difficult to study in the mouse because regulatory signaling overlaps with signals that

pattern tooth size and shape. Within the voles (Cricetidae, Rodentia, Glires), species have

evolved both rooted and unrooted molars that have similar size and shape. We assembled a *de*

novo genome of *Myodes glareolus*, a vole with high-crowned, rooted molars, and performed

genomic and transcriptomic analyses in a broad phylogenetic context of Glires (rodents and

lagomorphs) to assess differential selection and evolution in tooth forming genes.

Results: Our *de novo* genome recovered 91% of single-copy orthologs for Euarchontoglires and

had a total length of 2.44 Gigabases, enabling genomic and transcriptomic analyses. We

identified six dental genes undergoing positive selection across Glires and two genes undergoing

positive selection in species with unrooted molars, *Dspp* and *Aqp1*. Transcriptomics analyses

demonstrated conserved patterns of dental gene expression with species-specific variation likely

related to developmental timing and morphological differences between mouse and vole molars.

Conclusions: Our results support ongoing dental gene evolution in rodents with unrooted molars.

We identify candidate genes for further functional analyses, particularly *Dspp*, which plays an

important role in mineralizing tissues. Our expression results support conservation of dental

genes between voles and model species like mice, while revealing significant effects of overall

tooth morphology on gene expression.

3-10 Keywords: Evolution, selection, Glires, molar, root, dental, development, genome, rodent,

tooth

DECLARATIONS

Ethics approval: The University of California, San Francisco (UCSF) Institutional Animal Care

and Use Program and the Finnish national animal experimentation board approved protocols for

humane euthanasia and collection of tissues for animals used in this study under protocols

AN189916 (UCSF) and KEK16-021, KEK19-019, and KEK17-030 (University of Helsinki).

Consent for publication: Not applicable.

Availability of data and materials: The datasets supporting the conclusions of this article are

available in the GenBank repository under [GenBank reference number to be added upon

acceptance] and in the article's additional files.

Competing interests: The authors declare that they have no competing interests.

Funding: This research was supported by National Science Foundation grants CNS-0958379,

CNS-0855217, OAC-1126113, and OAC-2215760 through the City University of New York

High Performance Computing Center at the College of Staten Island; OAC-1925590 through the

MENDEL high performance computing cluster at the American Museum of Natural History;

Academy of Finland to JJ; Doctoral Programme in Biomedicine, University of Helsinki to

MMC; and National Institutes of Health NIDCR R01-DE027620 and R35-DE026602 to ODK.

Authors' contributions: ZTC and ODK designed the study. ZTC and PM performed animal

husbandry. ZTC performed and oversaw tissue sampling, sequencing, genome assembly and

annotation for *Myodes glareolus*. ZTC, AS, JR, EC, and MA performed genome computational

analyses. PL performed qPCR analyses. OH, MMC, RDR, and JJ designed and implemented

RNA sequencing experiments. ZTC wrote and all authors contributed to and approved the

manuscript.

Acknowledgements: The authors thank A. Joo, N. Ahituv, G. Amato, A. Narechania, S. Singh,

A. Scott, and A. Paasch for advice on methods and access to cluster computing resources.

BACKGROUND

Hypselodonty, or the presence of unrooted and thus ever-growing teeth, has evolved

multiple times in mammals. Glires—the clade containing rodents, rabbits, and their relatives—

have hypselodont incisors (1), and multiple Glires have also evolved hypselodont molars (Fig.

1). At least in rodents, molar hypselodonty evolved considerably later than hypsodont molars,

which are high crowned but rooted, which in turn evolved later than hypselodont incisors. In

Glires, molars appear to increase in crown height from low-crowned brachydont (low-crowned,

rooted), through hypsodonty (high-crowned, rooted), toward hypselodonty (high-crowned,

unrooted) (2). Mice (*Mus musculus*), the primary mammalian model species of dental research,

have highly derived hypselodont incisors while retaining brachydont molars. Because of this,

mice do not provide information about the hypsodont teeth that likely preceded hypselodonty.

Mammalian teeth sit in bony sockets, held in place by soft tissue (periodontal ligament)

attached to cementum-covered tooth roots (3). Ligamentous tooth attachment may have arisen

along with a reduction in the rate of tooth replacements, providing greater flexibility for

repositioning the teeth as the dentary grows (4,3). Consequently, the limited replacement of

mammalian teeth (two sets of teeth in most mammals and one in Glires) may have spurred the

evolution of hypsodont and hypselodont teeth, both with high crowns that compensate for tooth

wear from gritty or phytolith-heavy diets (5,6), and resulted in further modification of the

anchoring roots. The convergent evolution of unrooted molars in Glires presents an opportunity

to identify whether consistent developmental and genomic changes underlie the formation of

hypselodont teeth, in turn revealing the mechanisms that must remain unchanged to produce

tooth roots. Furthermore, the relatively recent evolution of molar hypselodonty, starting in the

Middle Miocene (approximately 16-12 Ma) (2), should provide molecular evidence for the steps

required to make a continuously growing organ.

Dental development proceeds from the tooth germ, composed of epithelium and

mesenchyme, through phases known as the bud, cap, and bell (7). Multipotent enamel epithelium

differentiates into the cells that form the tooth crown (8–11). As development progresses in

rooted teeth, the epithelium at the tooth apex transitions first to a tissue called Hertwig's

epithelial root sheath (HERS), and eventually cementum-covered roots (9,10). Studies have

identified numerous candidate genes and pathways with various roles during root development,

such as *Fgf10*, which decreases in expression at the beginning of root formation (12–18).

Although research on mouse molars has identified genetic signals of root formation, a number of

the key genes studied have broad developmental roles, such as *Wnt* family members (14), or

overlap considerably with genes also involved in patterning the size and shape of the tooth

(17,19–22). This overlap between shape and root expression patterns confounds our ability to

identify a clear signal initiating root formation.

Evolutionary novelties such as high-crowned hypsodont and hypselodont molars can

arise from differences in gene expression and regulation (23–26). Evolutionarily conserved gene

expression levels produce conserved phenotypes, and changes in gene regulatory networks have

long been linked to morphological evolution (27,28). The order of genes along a chromosome

(synteny) can affect gene expression and regulation, as regulatory sequences are often located

near their target genes (cis-regulatory elements) (29–31). Genome rearrangements that place

genes near new regulatory elements may result in changes of the expression levels and selective

environment of those genes; these small-scale rearrangements of genes may be common in

mammals (32–34). Genes involved in molar development are not syntenic in the mouse genome

nor are genes with organ-specific expression (35), and thus the regulatory effects of co-

localization need not apply to all dental genes at once. Changes in genome architecture between

Glires species thus may result in different selective and expression environments for dental genes

that could result in the evolution of hypselodont molars.

To establish a model rodent species with hypsodont molars for close comparison to

hypselodont molars, we sequenced and annotated a highly-complete *de novo* genome of *Myodes*

glareolus, the bank vole. The bank vole is increasingly used in medical and environmental

research, ranging from studying zoonotic diseases (36) to immune responses (37,38), and even

assessing environmental remediation efforts through heavy metals that accumulate in vole teeth

(39,40), thus our efforts may be of use beyond dental research. The bank vole's hypsodont

molars bridge the gap between low-crowned mouse and hypselodont prairie vole (*Microtus*

ochrogaster) molars. We performed a suite of genomic and transcriptomic tests of our new bank

vole genome in a broad phylogenetic context to test the hypothesis that dental genes are

undergoing positive selection and exhibit different expression patterns in species with unrooted,

hypselodont molars. We predicted that genes without conserved syntenic relationships in these

species would be more likely to have sites under positive selection or significantly different

expression. Our analyses revealed positive selection among two dental genes in Glires with

unrooted molars compared to those with rooted molars and demonstrated strong conservation of

dental gene expression patterns between bank voles and mice, with key differences related to the

timing and patterning of tooth morphology.

RESULTS

Orthology and synteny analyses

To identify which sequences in our bank vole (*Myodes glareolus*) genome and annotation

had the same evolutionary history as dental genes identified in other Glires and assess genome

rearrangements, we performed orthology and synteny analyses in a broad phylogenetic context.

OrthoFinder identified 20,547 orthogroups representing 97.9% of the genes across all 24

analyzed genomes (including the human outgroup). Of the orthogroups, 6,158 had all species

present. In our *de novo* bank vole genome, there were 27,824 annotated genes, of which 84.2%

were assigned to an orthogroup. Bank vole genes were present in 16,250 orthogroups. On

average, the genomes included in the OrthoFinder analysis had 19,814 genes, with 98.2% of

those assigned to orthogroups.

The completeness and large scaffold N50 (4.6 Megabases) of our bank vole assembly

supported its inclusion in generating a Glires synteny network. Using the infomap clustering

algorithm, we produced 19,694 microsynteny clusters from this overall synteny network. We did

not expect dental genes to share the same microsynteny cluster, and instead examined whether

each gene was in the same microsynteny cluster in species with rooted or unrooted molars.

Among the microsynteny clusters containing dental genes, 28 networks lacked synteny in at least

half the species with unrooted molars or did not have a one-to-one relationship with an

orthogroup (Fig. 2).

Positive selection analysis

We hypothesized that dental genes in species with unrooted molars are undergoing

positive selection. Our positive selection analyses in PAML (phylogenetic analysis by maximum

likelihood (41)) identified 6 dental gene orthogroups undergoing site-specific positive selection

across Glires (Table 1). One of these genes, *Col4a1*, also was largely not syntenic in species with

unrooted molars (Fig. 2). Another orthogroup consisted mainly of predicted sequences similar to

Runx3 but only had sequences from four species; both showed site-specific positive selection.

We then assessed 24 genes (those with site-specific positive selection or which lacked synteny in

at least half of the species with unrooted molars) for site-specific positive selection in species

with unrooted molars compared to species with rooted molars (branch-and-site specific positive

selection (42)). Two genes, *Dspp* and *Aqp1* were undergoing this branch-and-site specific

positive selection. Both genes had a single highly supported site (posterior probability > 0.95)

under positive selection in species with unrooted molars based on the Bayes Empirical Bayes

method for identifying sites under selection implemented in PAML (43). *Dspp* also had multiple

sites with moderate support (posterior probability > 0.75). The overall selection patterns on each

gene differed. Maximum likelihood estimates of selection for *Dspp* showed the percentage of

sites under purifying and neutral selection on all branches were nearly equal (47% and 44%,

respectively). Percentages of sites under positive selection in the species with unrooted molars

(foreground branches) were nearly evenly divided as well, with 5% of sites from branches where

the species with rooted molars (background branches) were undergoing purifying selection and

4% of sites from branches where the species with rooted molars were under neutral selection. For

Aqp1, nearly all sites were under purifying selection on all branches (91%), and few sites were

under neutral selection on all branches (7%). Few sites were undergoing positive selection in the

foreground branches and their distribution also was unevenly split between sites under purifying

and neutral selection on background branches (0.6% and 0.04%, respectively). The complete list

of dental genes with hierarchical orthogroups, microsynteny clusters, and positive selection test

results are available in Additional file 1.

201   Because genes under positive selection are often expressed at lower levels than genes

under purifying selection (44–47), we also compared expression levels of *Dspp* and *Aqp1* in

postnatal first molars (M1) at postnatal days 1, 15, and 21 (P1, P15, and P21) in bank voles

(rooted molars) and prairie voles (unrooted molars) using quantitative PCR. Prairie vole molars

expressed *Aqp1* at significantly lower levels in all three ages than bank vole molars (Fig. 3).

Prairie vole P1 molars expressed significantly lower levels of *Dspp* than bank vole molars; at

P15 and P21, their molars expressed *Dspp* at lower, but not statistically significantly different,

levels than their bank vole equivalent. For both genes, the prairie vole had consistent expression

levels across three biological replicates, while the bank vole had greater variation in expression

levels across replicates.

Sequence and secondary structure evolution

To detect whether substitutions at sites under positive selection influenced protein

structure and evolution, we analyzed ancestral states and secondary structure across Glires. We

first reconstructed ancestral sequences along the internal nodes of the Glires phylogeny for the

genes undergoing branch-and-site specific positive selection to assess potential secondary

structural changes in their protein sequences. At the best-supported site in *Dspp* (position 209 in

the gapped alignment, Additional file 2), there were three major amino acid changes. The

ancestral Glires sequence started with an asparagine (N) in this position. Two of the three species

with unrooted molars represented in the *Dspp* dataset had amino acid substitutions at this

position, with *Oryctolagus cuniculus* substituting a leucine (L) and *Dipodomys ordii* substituting

an aspartic acid (D) at this position (Fig. 4A). All muroids (the clade including the voles in

family Cricetidae and mice and rats in family Muridae) in our phylogeny substituted histidine

(H) for the asparagine at this position. The secondary structure predicted at this position was a

coil for most sequences but a helix for the *D. ordii* sequence (Fig. 5). *Aqp1* sequences varied

greatly at the position under putative positive selection in species with unrooted molars (position

294 in the gapped alignment, Additional file 3). The ancestral state reconstruction showed twelve

changes of the amino acid at this position across Glires (Fig. 4B), yet these changes did not

affect the predicted secondary structure of the protein near this residue, which was a coil for all

sequences tested. All secondary structure predictions are available in Additional file 4.

Developmental gene expression

We also assessed differential gene expression between mouse and bank vole molars

across early development to study the effects of morphology on expression levels of dental

genes. Our gene expression analysis focused on keystone dental gene categories, which are based

on the effects null mutations of each gene are reported to have during embryonic dental

development (48): "shape" genes cause morphological errors, "eruption" genes prevent tooth

eruption, "progression" genes stop the developmental sequence, "tissue" genes cause defects in

tissues, "developmental process" genes are annotated with the "GO:0032502" gene ontology

term, and "dispensable" genes, while dynamically expressed in developing teeth, have no

documented effect on phenotype. The group "other" is composed of the remaining protein

coding genes (48). Our bank vole genome was like the mouse and rat genomes in terms of the

numbers and expression patterns of genes annotated from these keystone categories (Table 2).

Ordination of gene expression results from the bank vole and mouse data at embryonic day 13,

14, and 16 (E13, E14, E16) (48) by principal components analysis showed a distinct separation

between the mouse and bank vole along the first principal component (PC1) of the 500 most

variable genes (Fig. 6A). PC1 explained 82.81% of the variance in these genes; there are distinct,

species-specific expression patterns in these tissues. Along PC2 (7.47% of variance explained),

E13 and E14 samples differ from the E16 samples, although the difference in time points is

much greater in bank voles. Ordination of just the keystone dental genes showed clear

separations between tissues based on species and age (Fig. 6B). Within this focused set of genes,

however, PC1 and PC2 explain less variance (44.8% and 28.84% respectively), and how species

and age relate to the PCs is less clear. There are two distinct, parallel trajectories for the mouse

and bank vole. Although within each species there is separation by age along PC1 and PC2,

mouse E16 and bank vole E13 occupy a similar position along PC1, and mouse E13 and bank

vole E16 occupy a similar position along PC2.

257   Examining individual genes underlying the differences between mouse and vole molars,

we note several upregulated genes in our vole molars are broadly expressed in developing molars

of other vole species (49,50). Relative to the mouse molars, vole molars overexpressed genes

related to forming tooth cusps, including *Bmp2*, *Shh*, *p21* (also known as *Cdkn1a*), and *Msx2*, a

difference explained by the faster patterning and larger number of cusps in the vole molar

compared to the mouse molar (50). Another gene upregulated in the patterning stage vole molar

is *Fgf10*, which is associated with delayed root formation later in vole molar development (9).

264   Nevertheless, developing bank vole molars at E13, E14, and E16 expressed keystone

dental genes in overall proportions like those observed at analogous stages of mouse and rat

molar development (Fig. 7). Permutation tests within each bank vole sample showed that log

counts for the set of genes related to the progression of dental development were significantly

higher than those in the tissue, dispensable, developmental process, and "other" categories at E14

and E16. The progression gene counts in E13 molars were higher for all of these except the

dispensable category. Shape category genes also were significantly higher than "other" category

genes in the E14 tissue. Overall, even though we observed conserved expression patterns of

dental genes at the system level, individual genes involved in cusp patterning and morphology

differed between the mouse and the vole.

DISCUSSION

Our two goals in sequencing the genome of *Myodes glareolus* were to support the

development of a comparative system for studying tooth root development and to investigate the

evolution of dental genes in Glires, a clade in which ever-growing molars have evolved multiple

times (1). Our new *M. glareolus* assembly and annotation captured nearly all of the single-copy

orthologs for Euarchontoglires and provided scaffolds with sufficient length for synteny

analyses. It was well represented in ortholog groups and microsynteny clusters across Glires. We

tested the hypothesis that dental genes are undergoing site-specific positive selection in species

with unrooted molars (branch-and-site specific positive selection (42)) and exhibit differential

expression patterns. We predicted that lack of conserved syntenic relationships in species with

unrooted molars could place dental genes in regulatory and selective environments that promote

changes among genes relevant to tooth root formation. Our analyses revealed that most dental

genes have conserved syntenic relationships across Glires, yet two dental genes, *Dspp* and *Aqp1*,

were undergoing positive selection in species with unrooted molars. We also demonstrated

conserved patterns of gene expression among dental keystone genes between bank voles and

mice during early embryonic development, and deviations from these conserved patterns likely

related to differences in molar morphology between the two species.

We identified 13 genes which were not syntenic in at least half of the species with

unrooted molars, and 6 genes undergoing site-specific positive selection across all Glires. Only

one gene, *Col4a1*, lacked synteny and had evidence of positive selection. The two genes

undergoing positive selection in species with unrooted molars, *Dspp* and *Aqp1*, both maintained

their synteny relationships across the Glires studied. Although we predicted loss of synteny for

dental genes in Glires with unrooted molars could result in sequence evolution by placing genes

in new selective contexts, our analyses did not support a relationship between non-syntenic genes

and positive selection. Maximum likelihood estimates of sites under different types of selection

for the genes with branch-specific positive selection did reveal different selective pressures on

Dspp and *Aqp1* overall; *Dspp* sites on background branches (i.e., branches with species that have

rooted molars) were under a mix of purifying and neutral selection, while nearly all *Aqp1*

background branch sites were under purifying selection. These selection regimes suggest there is

greater conservation for *Aqp1* function across Glires than for *Dspp* function. Gene duplication

can result in functional redundancy and evolution toward a novel function in some genes (51–

54), which may explain positive selection in *Aqp1*, as there are other aquaporin family genes

present. *Dspp* has no paralogs, but overlaps functionally with other SIBLING family proteins

(e.g., *Opn*, *Dmp1*) (55,56).

Aqp1 and *Dspp* play different functional roles during dental development. Under the

keystone dental development gene framework, *Aqp1* is a "dispensable" gene: developing teeth

express it, but tooth phenotypes do not change in its absence. *Aqp1* is expressed in endothelia of

microvessels in the developing tooth (57,58). *Dspp* may be particularly relevant for the

formation of an unrooted phenotype if its expression domain or function have been modified in

species with unrooted molars. *Dspp* is a "tissue" category keystone dental gene, meaning the

main effects of a null mutation occur during the tissue differentiation stage of dental

development (48). Null mutations of *Dspp* cause dentin defects in a condition called

dentinogenesis imperfecta (59,60); in some patients, teeth form short, brittle roots (60,61). *Dspp*

knockout mice also exhibit the shortened root phenotype, among a variety of other defects in

both endochondral and intramembranous bone, due to the disruption of collagen and bone

mineralization (62–64).

Our ancestral sequence reconstructions and estimated secondary protein structures

allowed us to assess whether nonsynonymous substitutions at sites under positive selection

resulted in structural differences, thus potentially affecting protein function. Although unrooted

molars are a convergent phenotype across Glires, the sites under positive selection did not

converge on the same amino acid substitution in species with unrooted molars, and *Aqp1*

appeared particularly labile at this residue. The non-synonymous substitutions at these sites often

resulted in changes of properties of the amino acid in the sequence, for example in *Dspp*, polar

asparagine was replaced with non-polar leucine in *O. cuniculus*. Only one of these substitutions

changed the predicted secondary structure. Nevertheless, single amino acid substitutions do

produce phenotypes for both *Dspp* (65) and *Aqp1* (66), thus we cannot rule out functional

changes in these genes in species with unrooted molars.

Although the exact relationship between gene expression and sequence divergence

remains unclear (67), studies of genome evolution across small numbers of mammal species

show correlations between gene sequence divergence and levels of expression (68). In particular,

highly-expressed genes are more likely to experience purifying selection (44–47), while lowly-

expressed genes and tissue-specific genes may experience positive selection (45). The decreased

expression of *Dspp* and *Aqp1* in prairie vole M1 compared to that of the bank vole M1 thus

supports our finding of positive selection in these genes in species with unrooted molars. If all

species with unrooted molars also exhibit decreased expression levels of *Dspp* and *Aqp1*, it could

suggest a strong link between lower levels of the genes and the unrooted phenotype.

Without analyses of functional variation caused by positive selection at these coding

sites, or spatial sampling to determine where these genes may be expressed during development,

we are limited from exploring the specific effects of *Dspp* and *Aqp1* on root formation.

Nevertheless, we found evidence for evolution of these genes in Glires with unrooted molars,

and *Dspp* especially has clinical relevance for tooth root formation. Future studies should explore

the spatial distribution of *Dspp* expression, which could be relevant to functional changes in

Glires with unrooted molars. If *Dspp* is relevant to the lack of root formation in hypselodont

Glires incisors, the positive selection identified here may modify its expression domain or its

interaction with yet-unidentified root formation co-factors, thus serially reproducing the unrooted

incisor phenotype in molars.

Our RNA sequencing results supported the bank vole as a suitable system for studying

dental development. Although molar morphology differs considerably across mammals,

candidate-gene approaches have identified numerous conserved genes involved in tooth

development and morphological patterning (69). Studies of single genes or gene families have

identified shape-specifying roles common to multiple species (50,70–72), and high-throughput

sequencing of mouse and rat molars demonstrate that both species express sets of dental

development genes in similar proportions during early stages of tooth development (48). The

similarity of our high-throughput RNA sequencing results to the mouse and rat results in

previous studies suggest overall expression patterns of keystone dental development genes

within each stage may be conserved in Glires. Our principal component analyses and differential

expression analyses measuring changes between mouse and bank vole molars, however, showed

that several dental genes' expression levels differed significantly by species and age. Previous

research has documented organ expression patterns that are conserved across species early in

development and diverge over time, with some major organs displaying heterochronic shifts in

some species (73). If the major source of variation in keystone dental gene expression patterns

between mice and bank vole molars were solely attributable to species, we might expect to see

clear separation between the species along the first or second principal component (PC1 or PC2),

like that observed in PC1 of the 500 most variable genes (Fig. 6). If molar development follows

the diverging expression patterns observed in other organs, we might expect just the earliest age

classes to align on one, or multiple, PCs. Instead, we found two trajectories that were nearly

parallel across PC1 and PC2 and multiple keystone dental genes that were significantly

differentially expressed with respect to species and age. This variation between species is likely

driven by the larger number of cusps in the vole molar, and corresponding upregulation of genes

regulating cusp formation. The overall acceleration of patterning in vole molars likely explains

the significance of the age variable in our expression results, causing a heterochronic shift in the

expression patterns.

Our analyses were limited by the small number of rodent species with sufficiently

annotated genomes to be included in synteny and positive selection analyses. This limitation left

us with a small phylogeny for our ancestral state reconstructions, which thus did not encompass

the full diversity of Glires tooth roots, and potentially weakened model-based genomic analyses.

Although positive selection analyses using the Bayes Empirical Bayes criterion are robust to

smaller sample sizes (43), including fossil species in ancestral state reconstructions can change

estimations of ancestral characteristics (74). Innovations in paleoproteomics also offer the

opportunity to compare fossil species' dental gene sequences directly to living and estimated

ancestral sequences (75,76). By incorporating data for extinct Glires in both morphological and

molecular analyses, we can further elucidate links between dental gene evolution and unrooted

teeth.

CONCLUSIONS

Our genomics and transcriptomics analyses, based on our newly sequenced, high-quality

draft bank vole genome assembly and annotation, showed that bank vole early tooth

development is comparable to other commonly used rodent models in dental development

research. We identified 6 dental gene orthogroups that were undergoing site-specific positive

selection across Glires and two genes, *Dspp* and *Aqp1*, that were undergoing site-specific

positive selection in Glires with unrooted molars. *Dspp* appears particularly relevant to root

formation, as loss-of-function mutations cause a dentin production defect that can result in

shortened tooth roots. Future research must explore the functional role that *Dspp* plays in tooth

root formation in Glires and other clades. The rodent dentary is an exciting system for

understanding tooth development; it provides an easily manipulated set of tissues that can be

produced quickly and features a lifelong population of stem cells in the incisor with genomic

mechanisms that are potentially replicated across other teeth in species with unrooted molars.

Our results identify candidate genes for future analyses, and our draft bank vole genome and

annotation improve the utility of this species for comparative dental research that can uncover

the genetic mechanisms of tooth root formation.

METHODS

Tissue collection and sequencing

To assemble the bank vole genome, we sequenced tissues from a single adult male

specimen housed in a colony at the UCSF Mission Center Animal Facility. We euthanized the

animal according to UCSF IACUC protocol AN189916 and harvested muscle, kidney, heart, and

liver tissue, which were immediately frozen at -80°C. Tissues were sent to a third-party

sequencing service, where they were combined and homogenized to achieve appropriate mass

for high molecular weight DNA extraction. We targeted 60x coverage with 150 base pair (bp)

reads using 10X Chromium linked-read chemistry (77,78) and sequenced on the Illumina

platform. We also targeted 10x coverage with Pacific Biosciences SMRT long-read chemistry.

For genome annotation and gene expression analyses, we collected seven biological replicates

each of first molars at embryonic days 13-16 (E13, E14, E15, E16), second molars at E16, and

jaw tissues at E14 under University of Helsinki protocols KEK16-021, KEK19-019, and KEK17-

030 and stored them in RNAlater at -80°C for RNA sequencing, following a tissue harvesting

protocol established for mice and rats (48). We extracted RNA from these tissues using a

guanidium thiocyanate and phenol-chloroform protocol combined with an RNeasy column

purification kit (Qiagen) based on the keystone dental gene protocol (48). Single-end 84 bp RNA

sequencing was performed using the Illumina NextSeq 500 platform.

Genome assembly and quality control

We first assembled only the 10X Chromium linked reads using the default settings in

Supernova 2.1.1. (77,78). We selected the "pseudohaplotype" (pseudohap) output format, which

randomly selects between potential alleles when there are two possible contigs assembled for the

same region. This option produces two assemblies, each with a single resolved length of the

genome sequence (77–79). We used our lower-coverage, long-read data for gap filling and

additional scaffolding. First, we estimated the genome's length using the raw sequence data in

GenomeScope (80), which predicted a length of 2.6 gigabases. We then performed error

correction of the long reads using Canu (81), removing reads shorter than 500 bp and

disregarding overlaps between reads of fewer than 350 bp. We kept only those reads with

minimum coverage of 3x for scaffolding. Following long read error correction, we used Cobbler

and RAILS (82) with a minimum alignment length of 200 bases to accept matches for gap filling

and scaffolding of both pseudohap assemblies.

For quality control, we assessed both unscaffolded and long-read scaffolded pseudohap

assemblies by standard assembly length statistics with QUAST (83) and presence of single-copy

orthologs with BUSCO v3 (84). Both scaffolded assemblies were approximately 2.44 Gigabases

long, with an N50 (the length of the shortest scaffold at 50% of the total assembly length) of 4.6

Megabases; we refer to them as Pseudohap1+LR and Pseudohap2+LR. The Pseudohap1+LR

assembly had 17,528 scaffolds over 1000bp (base pairs) long, and the Pseudohap2+LR assembly

had 17,518 scaffolds over 1000bp long (Table 3). BUSCO searched for universal single-copy

orthologs shared by Euarchontoglires, recovering 89.4% of these genes in the scaffolded

Pseudohap1+LR assembly and 92.8% of the single-copy orthologs in the scaffolded

Pseudohap2+LR assembly (Fig. 8). The two assemblies were similar length and contiguity, but

because the scaffolded Pseudohap2+LR assembly recovered more single-copy orthologs, we

based annotation and downstream analyses on it.

Genome annotation

We annotated the genome using three rounds of the MAKER pipeline (85–87). MAKER

combines multiple lines of evidence to annotate a genome. For evidence from gene transcripts,

we assembled a *de novo* transcriptome assembly based on the single-end RNA sequencing of all

molar and jaw tissues using Trinity (88). We also included cDNA sequences from the *Mus*

musculus assembly GRCm38. We used SwissProt's curated protein database to identify protein

homology in the genome. Two libraries of repeats provided information for repeat masking: the

Dfam Rodentia repeat library (89–91) and a custom library specific to the bank vole estimated

based on the modified protocol of Campbell et al. (86). The custom library features miniature

inverted-repeat transposable elements identified with default settings in MiteFinder (92), long

terminal repeat retrotransposons extracted with the GenomeTools LTRharvest and LTRdigest

functions (93) based on the eukaryotic genomic tRNA database, and *de novo* repeats identified

with RepeatModeler (94). We combined elements identified by these programs into a single

repeat library, then removed any elements that matched to a custom SwissProt curated protein

database with known transposons excluded; this custom repeat library is available in Additional

file 5. We trained a custom gene prediction model for MAKER as well. The first iteration of the

model came from BUSCO's implementation of augustus (95). Between each round of MAKER

annotation, we updated the gene prediction model with augustus.

MAKER considered only contigs between 10,000-300,000 bp long during annotation.

Our second and third iterations of MAKER used the same settings but excluded the

"Est2genome" and "protein2genome" functions, as recommended in the MAKER tutorial. We

included a SNAP (96) gene prediction model based on the output of the first round of annotation

during the second and third iterations of MAKER annotation. Annotation quality (i.e., agreement

between different lines of evidence and the MAKER annotation) was assessed visually in

JBrowse after each iteration and using *compare_annotations_3.2.pl* (97), which calculates the

number of coding and non-coding sequences in the annotation in addition to basic statistics about

sequence lengths. Our MAKER annotation covered 2.41 Gb of the scaffolded Pseudohap2

assembly in 4,125 scaffolds. These scaffolds contained 27,824 coding genes (mRNA) and 15,320

non-coding RNA sequences. The average gene length was 12,705 bp. Most annotations (91.4%)

had an annotation edit distance (AED) of 0.5 or better. AED is a measure of congruency between

the different types of evidence for an annotation, where scores closer to zero represent better-

annotated genes (98).

Orthology and synteny analyses

We analyzed orthology and synteny of the bank vole genome to understand gene and

genome evolution related to dental development across Glires with rooted and unrooted molars.

We obtained genomes from Ensembl for 23 Glires species and one phylogenetic outgoup, *Homo*

sapiens (Table 4). These genomes all had an N50 over 1 Mb, which improves synteny

assessment (99). We first analyzed all 24 genomes for groups of orthologous genes (orthogroups)

in OrthoFinder (100), providing a tree topology based on the Ensembl reference tree (Fig. 1) to

guide orthology detection. Because we would not analyze the human outgroup in downstream

analyses, we implemented the OrthoFinder option that splits orthogroups at the root of Glires

(hierarchical orthogroups), thus any group of orthologs studied here represents only genes with

shared, orthologous evolutionary history within Glires. We selected MAFFT (101) for multiple

sequence alignment and fastme (102) for phylogenetic tree searches within OrthoFinder; we

retained the gene trees estimated for each orthogroup for downstream analyses.

Although dental development genes are spread throughout the genome, we were

interested in whether each gene remained in the same local arrangement across species of Glires.

We prepared each genome annotation and sequence file for synteny analysis using the

reformatting functions of Synima (103) to extract each peptide sequence associated with a gene

coding sequence in the Ensembl annotation. Collinear synteny blocks estimated by MCScanX

(104) formed the basis for microsynteny network analyses using the SynNet pipeline (105–107).

We inferred networks from the top five hits for each gene, requiring any network to have a

minimum of 5 collinear genes and no more than 15 genes between a collinear block, settings that

perform well for analyzing mammal genomes (107). Using the infomap algorithm, we clustered

the synteny blocks into microsynteny networks, from which we extracted network clusters

corresponding to the list of keystone dental genes (48). For each dental gene microsynteny

network, we assessed whether genes of species with unrooted molars were not syntenic with the

other Glires species' sequences.

Positive selection analysis

We aligned protein sequences for each dental gene orthogroup with clustal omega (108)

using default settings. Based on universal translation tables, we obtained codon-based nucleotide

alignments with pal2nal (109), removing sites in which any species had an indel (i.e., ungapped)

and formatting the output for analysis in PAML (41). We pruned and unrooted the orthogroup

gene trees from OrthoFinder to contain only tips representing the genes in each synteny network

or orthogroup under analysis in PAML. We tested whether any of the genes were undergoing

positive selection using a likelihood ratio test comparing site-specific models of "nearly neutral"

and positive selection. In these models, ω, the ratio of nonsynonymous to synonymous

nucleotide substitutions (also known as dN/dS), can vary at each codon site. In the "nearly

neutral" model, ω can take values between 0 and 1, while the positive selection model allows

sites to assume ω values greater than 1 (43,110). We allowed PAML to estimate κ (the ratio of

transitions to transversions) and ω from initial values of 1 and 0.5, respectively, for both tests.

Dental genes with significant site-specific positive selection or those for which over half

the unrooted species' sequences were not in the same synteny block as sequences for species

with rooted molars formed the basis for our second set of positive selection tests using a branch-

and-site model of positive selection. This model allows ω to vary not only among codon sites,

but also between "foreground" and "background" lineages (43). We marked the species with

unrooted molars as foreground lineages, then ran the model twice: once with ω unconstrained to

detect sites undergoing positive selection only on foreground branches, and a second time and

with ω fixed to 1, or neutral selection. A likelihood ratio test of the two models determined

whether the lineage-specific positive selection model was more likely than a neutral model, and

Bayes Empirical Bayes analyses (43) produced posterior probabilities to identify sites under

positive selection.

Genes under positive selection also tend to have lower expression levels (45), thus we

compared expression of the genes with branch-and-site specific positive selection between the

prairie (unrooted molars) and the bank vole (rooted molars) to provide further support for

selective differences. We collected three biological replicates of first molars from both species at

three postnatal stages (P1, P15, and P21) and immediately preserved them at -80°C in lysis

buffer (Buffer RLT; Qiagen) supplemented with 40 μM dithiothreitol. RNA was extracted from

homogenized tissues using a RNeasy column purification kit (Qiagen). We assessed

concentration and purity of extracted RNA using a NanoDrop 2000 spectrophotometer

(ThermoFisher Scientific). Using 1 μg of RNA, we synthesized cDNA using a high-capacity

cDNA reverse transcription kit (ThermoFisher Scientific). We used 1 μL diluted cDNA (1:3 in

ddH$_2$O) and iTaq Universal SYBR Green Supermix (Bio-rad) in the Bio-rad CFX96 real-time

PCR detection system for qPCR experiments, producing three technical replicates for each

biological replicate. We normalized cycle threshold (CT) values of genes of interest to GAPDH

expression levels and calculated relative expression levels as $2^{-\Delta\Delta CT}$. A two-tailed unpaired t-test

calculated in Prism 9 measured whether expression of these genes significantly differed between

bank voles and prairie voles. The oligonucleotide primers for each species and gene are in

Additional file 6.

Sequence and secondary structure evolution

We performed ancestral sequence reconstruction on the codon sequences of the genes

that had evidence of branch-and-site specific positive selection to understand how the sequence

has changed through time. The gapped clustal omega alignments were the basis for ancestral

sequence reconstruction on the Glires species tree (Fig. 1) using pagan2 (111). For each gene, we

plotted amino acid substitutions at the site with potential positive selection. Finally, we predicted

secondary structures (i.e., helices, beta sheets, and coils) for each unrooted species' protein

sequence and the reconstructed ancestral sequence prior to the change at the site under positive

selection using the PSIPRED 4.0 protein analysis workbench (112,113). Comparing these

predictions across the phylogeny, we assessed how these substitutions at the site under selection

may affect the structure of each protein.

Developmental gene expression

We used performed quality control and filtering of the short reads for the seven replicates

of first molar tissues at E13, E14, and E16 using the nf-core/rnaseq v. 3.11.2 workflow (114) for

comparability to previous mouse and rat analyses (48). RNAseq reads were evaluated and

adapter sequences were filtered using FastQC v. 0.11.9 (115) and Cutadapt v. 3.4 (116), and

ribosomal RNA was removed using SortMeRNA v. 4.3.4 (117). We then aligned trimmed

sequences to our bank vole annotation using Salmon v. 1.10.1 (118). Counts were then

normalized by gene length. We categorized gene count data into functional groups based on their

established roles in tooth bud development (48) using the one-to-one orthology list between our

bank vole genome and the mouse GRCm39.103 genome annotation generated from our

OrthoFinder output. Using the rlog function of DESeq2 (119), we normalized gene counts within

each functional group on a log2 scale. A permutation test assessed whether the mean counts of

the progression, shape, and double functional groups were significantly different from genes in

the tissue, dispensable, and "other" groups (which are potentially relevant later in development)

based on 10,000 resampling replicates of the dataset (48).

We also assessed differential expression between the bank vole first molar and published

mouse M1 data at the same three time points (GEO accession GSE142199 (48)), combining the

data based on the one-to-one orthology relationships used in the functional permutation analysis.

Using the mouse E13 molar as the reference level, we modeled expression as a response to

species (mouse or vole), embryonic day (E13, E14, or E16), and the interaction between species

and day. We considered as significant any gene with a log fold change greater than 1, log fold

change standard error less than 0.5, and false discovery rate adjusted p value less than 0.05.

596 TABLES

597 **Table 1 – Genes undergoing site-specific and branch-and-site-specific positive selection**

Gene	*Mus* transcript	*Myodes* transcript	Site	Branch-and-site
Aqp1	ENSMUST00000004774	Mglareolus_00011822	Yes	Yes
Col4a1	ENSMUST00000033898	Mglareolus_00032740	Yes	No
Dspp	ENSMUST00000112771	Mglareolus_00014030	Yes	Yes
Fgf20	ENSMUST00000034014	Mglareolus_00013079	Yes	No
Runx3	ENSMUST00000056977	Mglareolus_00033992	Yes	No
similar to *Runx3*	–	–	Yes	No*

598 Table 1 Legend: *HOG only contained four genes with one unrooted species' sequence, could

599 not be tested for branch-and-site specific selection.

601 **Table 2 – P-values of permutation tests between keystone gene categories in bank vole M1**

602 **at embryonic days 13, 14, and 16**

		Tissue	Dispensable	Dev. Process	Other
E13	**Progression**	*0.0310*	0.0942	*0.0436*	*0.0402*
	Shape	0.6431	0.9041	0.2289	0.0995
	Double	0.1292	0.1521	0.0716	0.0655
E14	**Progression**	*0.0136*	*0.0383*	*0.0437*	*0.0401*
	Shape	0.3115	0.4725	0.0922	*0.0454*
	Double	0.1288	0.0945	0.0709	0.0630
E16	**Progression**	*0.0140*	*0.0401*	*0.0303*	*0.0274*
	Shape	0.3770	1	0.1831	0.0662
	Double	0.1343	0.1099	0.0638	0.0596

603 Table 2 Legend: Italicized values are statistically significant ($p < 0.05$)

604 **Table 3 – QUAST assembly statistics for *de novo* bank vole (*Myodes glareolus*) genome**

605 **assemblies**

	Pseudohap1	Pseudohap1+LR	Pseudohap2	Pseudohap2+LR*
Largest contig	27939478	32658832	27937749	32657565
Total length	2434151515	2441426554	2434099357	2441472313
GC (%)	41.88	41.89	41.88	41.89
N50	4187179	4579815	4187179	4558134
N75	1689669	1818134	1687188	1810460
L50	170	153	170	154
L75	388	357	388	358
Ns per 100 kbp	1151.99	1030.75	1151.96	1030.48

606 Table 3 Legend: *assembly used for annotation and downstream analyses in this paper.

608 **Table 4 – Genomes used in orthology, synteny, and positive selection analyses**

Species	Assembly	Citation
Myodes glareolus	CUNY_Mgla_1.0	This paper
*Cavia porcellus**	Cavpor3.0	(120)
*Cavia aperea**	CavAp1.0	(121)
Marmota marmota	marMar2.1	(122)
*Microtus ochrogaster**	MicOch1.0	(123)
Mus musculus	GRCm39	(124)
*Oryctolagus cuniculus**	OryCun2.0	(120)
*Dipodomys ordii**	Dord_2.0	(120)
Jaculus jaculus	JacJac1.0	(125)

Rattus norvegicus	Rnor_6.0	(126)
Mus pahari	PAHARI_EIJ_v1.1	(127)
Mus caroli	CAROLI_EIJ_v1.1	(127)
Mus spretus	SPRET_EiJ_v1	(128)
Mus spicilegus	MUSP714	(129)
Cricetulus griseus	CHOK1GS	(130)
Mesocricetus auratus	MesAur1.0	(131)
Peromyscus maniculatus	HU_Pman_2.1	(132)
Nannospalax galili	S.galili_v1.0	(133)
*Octodon degus**	OctDeg1.0	(134)
Heterocephalus glaber (F)	HetGla_female_1.0	(135)
*Chinchilla lanigera**	ChiLan1.0	(136)
Urocitellus parryi	ASM342692v1	(137)
Ictidomys tridecemlineatus	SpeTri2.0	(138)
*Homo sapiens***	GRCh38	(139)

609 Table 4 Legend: *Species with unrooted molars; **Peptide annotation used as outgroup only in

610 OrthoFinder analysis.

Figure 1 – Species tree of Glires based on the Ensembl Compara species tree. Whether each species has rooted or unrooted molars is indicated at the tip of each branch. Note that unrooted, or hypselodont, molars have evolved multiple times across Glires. This topology was the basis for our orthology analysis.

Figure 2 – Heatmap showing the number of genes per species (inset gradient scale from 0 to

10+) in each synteny cluster. The figure shows clusters where species with unrooted molars had

no representation or did not have sequences in all microsynteny clusters associated with a single

gene, clusters where more than one gene mapped to the same cluster, or a single gene mapped to

multiple clusters. Microsynteny cluster number is noted on the right side of the heatmap (one

row per cluster), and corresponding genes are noted on the left with alternating bands showing

rows to which those genes mapped. * = genes where hierarchical orthogroup did not contain

genes for all 23 species. We found little evidence that non-syntenic genes in species with

unrooted molars are undergoing selection for novel functions.

655   **Figure 3** – Quantitative PCR comparisons of *Dspp* and *Aqp1* expression between bank vole and

656   prairie vole M1 at postnatal days 1, 15, and 21 (P1, P15, P21). Expression levels for both genes

657   are lower in the prairie vole (unrooted molars), which supports the positive selection detected for

658   these genes in species with unrooted molars.

668   **Figure 4** – Ancestral state reconstructions of the residue under positive selection in PAML tests.

669   Letters at tips and internal nodes represent IUPAC codes for amino acids. **A** *Dspp*; **B** *Aqp1*.

Figure 5 – PSIPRED secondary structure predictions for the three species with unrooted molars

represented in the *Dspp* sequences. Letters correspond to the most recent ancestor of each tip

species that had a different amino acid at the position under positive selection.

Figure 6 – Principal component (PC) analyses of differentially expressed genes in mouse and

bank vole M1. **A** PC1 and PC2 of the 500 most variable genes, showing a clear differentiation

between species along PC1 and differentiation between age classes along PC2. **B** PC1 and PC2

of the keystone dental genes. Both PC1 and PC2 separate age classes within, but not between,

the species, likely due to differences in developmental timing and molar morphology between

mice and voles.

Figure 7 – Box and whisker plots showing normalized log base 2 expression levels for each

keystone gene category in bank vole M1 at embryonic days 13, 14, and 16. Gene expression

profiles at these stages are comparable to mouse and rat molars at analogous developmental

stages, as seen in Hallikas et al. 2021.

Figure 8 – BUSCO single-copy ortholog recovery for each "pseudohaploid" version of our draft

bank vole genome assembly and these version after long-read scaffolding (denoted by "+ LR").

Each bar represents the cumulative proportion of the 6,192 single-copy orthologs for

Euarchontoglires identified by BUSCO represented by complete single-copy, complete-

duplicated, fragmented, and missing orthologs. The Pseudohap2 and Pseudohap2 + LR

assemblies had the best single-copy ortholog recovery.

ADDITIONAL FILES

Additional file 1 [.xlsx] Dental gene results – Full table of orthology, synteny, and positive

selection test results for all dental genes assessed.

Additional file 2 [.txt] *Dspp* gapped alignment – Gapped codon-based alignment for *Dspp* in

fasta formatted sequences.

Additional file 3 [.txt] *Aqp1* gapped alignment – Gapped codon-based alignment for *Aqp1* in

fasta formatted sequences.

Additional file 4 [.pdf] Structure predictions – PSIPRED Secondary structure predictions for

each ancestral node and unrooted molar tip species for *Dspp* and *Aqp1*.

Additional file 5 [.txt] Custom repeat library – Custom repeat library of fasta formatted

sequences used in annotation of the draft *Myodes glareolus* genome. See Methods for description

of the process used to generate the library.

Additional file 6 [.pdf] Oligonucleotide primers – List of oligonucleotide primers for *Dspp*,

Aqp1, and *GAPDH* used in bank vole and prairie vole qPCR experiments.

REFERENCES

1. Renvoisé E, Michon F. An Evo-Devo perspective on ever-growing teeth in mammals and

dental stem cell maintenance. Front Physiol. 2014;5 AUG(August):1–12.

2. Tapaltsyan V, Eronen JT, Lawing AM, Sharir A, Janis C, Jernvall J, et al. Continuously

growing rodent molars result from a predictable quantitative evolutionary change over 50

million years. Cell Rep. 2015;11(5):673–80.

722 3. LeBlanc ARH, Brink KS, Whitney MR, Abdala F, Reisz RR. Dental ontogeny in extinct

723   synapsids reveals a complex evolutionary history of the mammalian tooth attachment

724   system. Proc R Soc B Biol Sci. 2018 Nov 7;285(1890):20181792.

725 4. Saffar JL, Lasfargues JJ, Cherruau M. Alveolar bone and the alveolar process: the socket that

726   is never stable. Periodontol 2000. 1997;13(1):76–90.

727 5. Davit-Béal T, Tucker AS, Sire JY. Loss of teeth and enamel in tetrapods: Fossil record,

728   genetic data and morphological adaptations. J Anat. 2009;214(4):477–501.

729 6. Damuth J, Janis CM. On the relationship between hypsodonty and feeding ecology in

730   ungulate mammals, and its utility in palaeoecology. Biol Rev. 2011;86(3):733–58.

731 7. Miletich I, Sharpe PT. Normal and abnormal dental development. Hum Mol Genet. 2003 Apr

732   2;12(suppl_1):R69–73.

733 8. Harada H, Kettunen P, Jung HS, Mustonen T, Wang YA, Thesleff I. Localization of putative

734   stem cells in dental epithelium and their association with Notch and FGF signaling. J Cell

735   Biol. 1999;147(1):105–20.

736 9. Tummers M, Thesleff I. Root or crown: A developmental choice orchestrated by the

737   differential regulation of the epithelial stem cell niche in the tooth of two rodent species.

738   Development. 2003;130(6):1049–57.

739 10. Thesleff I, Tummers M. Tooth organogenesis and regeneration. In: StemBook. Cambridge,

740   MA: Harvard Stem Cell Institute; 2008.

11. Krivanek J, Buchtova M, Fried K, Adameyko I. Plasticity of dental cell types in

development, regeneration, and evolution. J Dent Res. 2023 Jun 1;102(6):589–98.

12. Luan X, Ito Y, Diekwisch TGH. Evolution and development of Hertwig's epithelial root

sheath. Dev Dyn. 2006;235(5):1167–80.

13. Kumakami-Sakano M, Otsu K, Fujiwara N, Harada H. Regulatory mechanisms of Hertwig's

epithelial root sheath formation and anomaly correlated with root length. Exp Cell Res.

2014;325(2):78–82.

14. Wen Q, Jing J, Han X, Feng J, Yuan Y, Ma Y, et al. *Runx2* regulates mouse tooth root

development via activation of WNT inhibitor *NOTUM*. J Bone Miner Res.

2020;35(11):2252–64.

15. Yang S, Choi H, Kim TH, Jeong JK, Liu Y, Harada H, et al. Cell dynamics in Hertwig's

epithelial root sheath are regulated by β-catenin activity during tooth root development. J

Cell Physiol. 2021;236(7):5387–98.

16. Yamashiro T, Tummers M, Thesleff I. Expression of bone morphogenetic proteins and Msx

genes during root formation. J Dent Res. 2003;82(3):172–6.

17. Yokohama-Tamaki T, Ohshima H, Fujiwara N, Takada Y, Ichimori Y, Wakisaka S, et al.

Cessation of Fgf10 signaling, resulting in a defective dental epithelial stem cell

compartment, leads to the transition from crown to root formation. Development.

2006;133(7):1359–66.

18. Ota MS, Vivatbutsin P, Nakahara T, Eto K. Tooth root development and the cell-based

regenerative therapy. J Oral Tissue Eng. 2007;4(3):137–42.

19. Jernvall J, Thesleff I. Reiterative signaling and patterning during mammalian tooth

morphogenesis. Mech Dev. 2000;92:19–29.

20. Harada H, Toyono T, Toyoshima K, Yamasaki M, Itoh N, Kato S, et al. FGF10 maintains

stem cell compartment in developing mouse incisors. Dev Camb Engl. 2002;129(6):1533–

41.

21. Tapaltsyan V, Charles C, Hu J, Mindell D, Ahituv N, Wilson GM, et al. Identification of

novel *Fgf* enhancers and their role in dental evolution. Evol Dev. 2016;18(1):31–40.

22. Christensen MM, Hallikas O, Das Roy R, Väänänen V, Stenberg OE, Häkkinen TJ, et al.

The developmental basis for scaling of mammalian tooth size. Proc Natl Acad Sci. 2023

Jun 20;120(25):e2300374120.

23. Chen ZJ. Genetic and epigenetic mechanisms for gene expression and phenotypic variation

in plant polyploids. Annu Rev Plant Biol. 2007;58(1):377–406.

24. Stranger BE, Forrest MS, Dunning M, Ingle CE, Beazley C, Thorne N, et al. Relative impact

of nucleotide and copy number variation on gene expression phenotypes. Science. 2007 Feb

9;315(5813):848–53.

25. Romero IG, Ruvinsky I, Gilad Y. Comparative studies of gene expression and the evolution

of gene regulation. Nat Rev Genet. 2012 Jul;13(7):505–16.

26. de Montaigu A, Giakountis A, Rubin M, Tóth R, Cremer F, Sokolova V, et al. Natural

diversity in daily rhythms of gene expression contributes to phenotypic variation. Proc Natl

Acad Sci. 2015 Jan 20;112(3):905–10.

27. Erwin DH, Davidson EH. The last common bilaterian ancestor. Development. 2002 Jul

1;129(13):3021–32.

28. Irie N, Kuratani S. Comparative transcriptome analysis reveals vertebrate phylotypic period

during organogenesis. Nat Commun. 2011;2:248.

29. Koonin EV. Evolution of genome architecture. Int J Biochem Cell Biol. 2009 Feb

1;41(2):298–306.

30. Wray GA. The evolutionary significance of cis-regulatory mutations. Nat Rev Genet. 2007

Mar;8(3):206–16.

31. Acemel RD, Maeso I, Gómez-Skarmeta JL. Topologically associated domains: A successful

scaffold for the evolution of gene regulation in animals. WIREs Dev Biol. 2017;6(3):e265.

32. Coghlan A, Eichler EE, Oliver SG, Paterson AH, Stein L. Chromosome evolution in

eukaryotes: A multi-kingdom perspective. Trends Genet. 2005 Dec 1;21(12):673–82.

33. Swenson KM, Blanchette M. Large-scale mammalian genome rearrangements coincide with

chromatin interactions. Bioinformatics. 2019 Jul 15;35(14):i117–26.

34. Long HS, Greenaway S, Powell G, Mallon AM, Lindgren CM, Simon MM. Making sense of

the linear genome, gene function and TADs. Epigenetics Chromatin. 2022 Jan 29;15(1):4.

35. Das Roy R, Hallikas O, Christensen MM, Renvoisé E, Jernvall J. Chromosomal

neighbourhoods allow identification of organ specific changes in gene expression. PLOS

Comput Biol. 2021 Sep 10;17(9):e1008947.

36. Torelli F, Zander S, Ellerbrok H, Kochs G, Ulrich RG, Klotz C, et al. Recombinant IFN-γ

from the bank vole *Myodes glareolus*: A novel tool for research on rodent reservoirs of

zoonotic pathogens. Sci Rep. 2018;8(1):1–11.

37. Kloch A, Babik W, Bajer A, Siński E, Radwan J. Effects of an MHC-DRB genotype and

allele number on the load of gut parasites in the bank vole *Myodes glareolus*. Mol Ecol.

2010;19(SUPPL. 1):255–65.

38. Migalska M, Sebastian A, Konczal M, Kotlík P, Radwan J. *De novo* transcriptome assembly

facilitates characterisation of fast-evolving gene families, MHC class I in the bank vole

(*Myodes glareolus*). Heredity. 2017;118(4):348–57.

39. Appleton J, Lee KM, Sawicka Kapusta K, Damek M, Cooke M. The heavy metal content of

the teeth of the bank vole (*Clethrionomys glareolus*) as an exposure marker of

environmental pollution in Poland. Environ Pollut. 2000;110:441–9.

40. Gdula-Argasińska J, Appleton J, Sawicka-Kapusta K, Spence B. Further investigation of the

heavy metal content of the teeth of the bank vole as an exposure indicator of environmental

pollution in Poland. Environ Pollut. 2004;131(1):71–9.

41. Yang Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Mol Biol Evol. 2007

Aug 1;24(8):1586–91.

42. Zhang J, Nielsen R, Yang Z. Evaluation of an improved branch-site likelihood method for

detecting positive selection at the molecular level. Mol Biol Evol. 2005 Dec;22(12):2472–9.

43. Yang Z, Wong WSW, Nielsen R. Bayes Empirical Bayes inference of amino acid sites under

positive selection. Mol Biol Evol. 2005 Apr 1;22(4):1107–18.

44. Drummond DA, Bloom JD, Adami C, Wilke CO, Arnold FH. Why highly expressed proteins

evolve slowly. Proc Natl Acad Sci. 2005 Oct 4;102(40):14338–43.

45. Kosiol C, Vinař T, Fonseca RR da, Hubisz MJ, Bustamante CD, Nielsen R, et al. Patterns of

positive selection in six mammalian genomes. PLOS Genet. 2008 Aug 1;4(8):e1000144.

46. Martincorena I, Luscombe NM. Non-random mutation: The evolution of targeted

hypermutation and hypomutation. BioEssays. 2013;35(2):123–30.

47. Martincorena I, Roshan A, Gerstung M, Ellis P, Van Loo P, McLaren S, et al. High burden

and pervasive positive selection of somatic mutations in normal human skin. Science. 2015

May 22;348(6237):880–6.

48. Hallikas O, Das Roy R, Christensen MM, Renvoisé E, Sulic AM, Jernvall J. System-level

analyses of keystone genes required for mammalian tooth development. J Exp Zoolog B

Mol Dev Evol. 2021;336(1):7–17.

49. Keränen SVE, Åberg T, Kettunen P, Thesleff I, Jernvall J. Association of developmental

regulatory genes with the development of different molar tooth shapes in two species of

rodents. Dev Genes Evol. 1998;208(9):477–86.

50. Jernvall J, Keränen SVE, Thesleff I. Evolutionary modification of development in

mammalian teeth: Quantifying gene expression patterns and topography. Proc Natl Acad

Sci. 2000;97(26):14444–8.

51. Hughes AL. The evolution of functionally novel proteins after gene duplication. Proc R Soc

Lond B Biol Sci. 1997 Jan;256(1346):119–24.

52. Wagner A. Selection and gene duplication: A view from the genome. Genome Biol. 2002

Apr 15;3(5):reviews1012.1.

53. David KT, Oaks JR, Halanych KM. Patterns of gene evolution following duplications and

speciations in vertebrates. PeerJ. 2020 Mar 31;8:e8813.

54. Copley SD. Evolution of new enzymes by gene duplication and divergence. FEBS J.

2020;287(7):1262–83.

55. Fisher LW. DMP1 and DSPP: Evidence for duplication and convergent evolution of two

SIBLING proteins. Cells Tissues Organs. 2011 Aug;194(2–4):113–8.

56. Bouleftour W, Juignet L, Bouet G, Granito RN, Vanden-Bossche A, Laroche N, et al. The

role of the SIBLING, bone sialoprotein in skeletal biology — Contribution of mouse

experimental genetics. Matrix Biol. 2016 May 1;52–54:60–77.

57. Felszeghy S, Módis L, Németh P, Nagy G, Zelles T, Agre P, et al. Expression of aquaporin

isoforms during human and mouse tooth development. Arch Oral Biol. 2004 Apr

1;49(4):247–57.

58. Yoshii T, Harada F, Saito I, Nozawa-Inoue K, Kawano Y, Maeda T. Immunoexpression of

aquaporin-1 in the rat periodontal ligament during experimental tooth movement. Biomed

Res. 2012;33(4):225–33.

59. Zhang X, Zhao J, Li C, Gao S, Qiu C, Liu P, et al. *DSPP* mutation in dentinogenesis

imperfecta Shields type II. Nat Genet. 2001 Feb;27(2):151–2.

60. de La Dure-Molla M, Philippe Fournier B, Berdal A. Isolated dentinogenesis imperfecta and

dentin dysplasia: Revision of the classification. Eur J Hum Genet. 2015 Apr;23(4):445–51.

61. Shields ED, Bixler D, El-Kafrawy AM. A proposed classification for heritable human

dentine defects with a description of a new entity. Arch Oral Biol. 1973 Apr 1;18(4):543-

IN7.

62. Sreenath T, Thyagarajan T, Hall B, Longenecker G, D'Souza R, Hong S, et al. Dentin

sialophosphoprotein knockout mouse teeth display widened predentin zone and develop

defective dentin mineralization similar to human dentinogenesis imperfecta type III. J Biol

Chem. 2003 Jul 4;278(27):24874–80.

63. Verdelis K, Ling Y, Sreenath T, Haruyama N, MacDougall M, van der Meulen MCH, et al.

DSPP effects on *in vivo* bone mineralization. Bone. 2008 Dec 1;43(6):983–90.

64. Chen Y, Zhang Y, Ramachandran A, George A. DSPP is essential for normal development

of the dental-craniofacial complex. J Dent Res. 2016 Mar 1;95(3):302–10.

65. von Marschall Z, Mok S, Phillips MD, McKnight DA, Fisher LW. Rough endoplasmic

reticulum trafficking errors by different classes of mutant dentin sialophosphoprotein

(DSPP) cause dominant negative effects in both dentinogenesis imperfecta and dentin

dysplasia by entrapping normal DSPP. J Bone Miner Res. 2012;27(6):1309–21.

66. Smith BL, Preston GM, Spring FA, Anstee DJ, Agre P. Human red cell aquaporin CHIP. I.

Molecular characterization of ABH and Colton blood group antigens. J Clin Invest. 1994

Sep 1;94(3):1043–9.

67. Jordan IK, Mariño-Ramírez L, Koonin EV. Evolutionary significance of gene expression

divergence. Gene. 2005 Jan 17;345(1):119–26.

68. Warnefors M, Kaessmann H. Evolution of the correlation between expression divergence

and protein divergence in mammals. Genome Biol Evol. 2013;5(7):1324–35.

69. Jernvall J, Thesleff I. Tooth shape formation and tooth renewal: Evolving with the same

signals. Development. 2012;139(19):3487–97.

70. Mitsiadis TA. Role of Islet1 in the patterning of murine dentition. Development.

2003;130(18):4451–60.

71. Charles C, Pantalacci S, Peterkova R, Tafforeau P, Laudet V, Viriot L. Effect of *eda* loss of

function on upper jugal tooth morphology. Anat Rec. 2009;292(2):299–308.

72. Zurowski C, Jamniczky H, Graf D, Theodor J. Deletion/loss of bone morphogenetic protein

7 changes tooth morphology and function in *Mus musculus*: Implications for dental

evolution in mammals. R Soc Open Sci. 2018 Jan 3;5(1):170761.

73. Cardoso-Moreira M, Halbert J, Valloton D, Velten B, Chen C, Shao Y, et al. Gene

expression across mammalian organ development. Nature. 2019 Jul;571(7766):505–9.

74. Finarelli JA, Flynn JJ. Ancestral state reconstruction of body size in the Caniformia

(Carnivora, Mammalia): The effects of incorporating data from the fossil record. Syst Biol.

2006;55(2):301–13.

75. Welker F, Collins MJ, Thomas JA, Wadsley M, Brace S, Cappellini E, et al. Ancient proteins

resolve the evolutionary history of Darwin's South American ungulates. Nature. 2015

Jun;522(7554):81–4.

76. Warinner C, Korzow Richter K, Collins MJ. Paleoproteomics. Chem Rev. 2022 Aug

24;122(16):13401–46.

77. Zheng GXY, Lau BT, Schnall-Levin M, Jarosz M, Bell JM, Hindson CM, et al. Haplotyping

germline and cancer genomes with high-throughput linked-read sequencing. Nat

Biotechnol. 2016 Feb;34:303.

78. Marks P, Garcia S, Martinez A, Belhocine K. Resolving the full spectrum of human genome

variation using linked-reads. 2017;

79. Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB. Direct determination of diploid

genome sequences. Genome Res. 2017;27(5):757–67.

80. Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, et al.

GenomeScope: Fast reference-free genome profiling from short reads. Bioinformatics. 2017

Jul 15;33(14):2202–4.

81. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: Scalable and

accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome

Res. 2017 May 1;27(5):722–36.

82. Warren RL. RAILS and Cobbler: Scaffolding and automated finishing of draft genomes

using long DNA sequences. J Open Source Softw. 2016 Nov 17;1(7):116.

83. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: Quality assessment tool for genome

assemblies. Bioinformatics. 2013 Apr 15;29(8):1072–5.

84. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: Assessing

genome assembly and annotation completeness with single-copy orthologs. Bioinformatics.

2015 Oct 1;31(19):3210–2.

85. Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, et al. MAKER: An easy-to-use

annotation pipeline designed for emerging model organism genomes. Genome Res.

2008;18:188–96.

86. Campbell MS, Law M, Holt C, Stein JC, Moghe GD, Hufnagel DE, et al. MAKER-P: A tool

kit for the rapid creation, management, and quality control of plant genome annotations.

Plant Physiol. 2014 Feb 1;164(2):513–24.

87. Campbell MS, Holt C, Moore B, Yandell M. Genome annotation and curation using

MAKER and MAKER-P. Curr Protoc Bioinforma. 2014 Dec 12;48:4.11.1-4.11.39.

88. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Trinity:

Reconstructing a full-length transcriptome without a genome from RNA-Seq data. Nat

Biotechnol. 2011 May 15;29(7):644–52.

89. Wheeler TJ, Clements J, Eddy SR, Hubley R, Jones TA, Jurka J, et al. Dfam: A database of

repetitive DNA based on profile hidden Markov models. Nucleic Acids Res. 2013

Jan;41(Database issue):D70-82.

90. Caballero J, Smit AFA, Hood L, Glusman G. Realistic artificial DNA sequences as negative

controls for computational genomics. Nucleic Acids Res. 2014 Jul;42(12):e99.

91. Hubley R, Finn RD, Clements J, Eddy SR, Jones TA, Bao W, et al. The Dfam database of

repetitive DNA families. Nucleic Acids Res. 2016 Jan 4;44(D1):D81–9.

92. Hu J, Zheng Y, Shang X. MiteFinder: A fast approach to identify miniature inverted-repeat

transposable elements on a genome-wide scale. In: 2017 IEEE International Conference on

Bioinformatics and Biomedicine (BIBM). 2017. p. 164–8.

93. Gremme G, Steinbiss S, Kurtz S. GenomeTools: A comprehensive software library for

efficient processing of structured genome annotations. IEEE/ACM Trans Comput Biol

Bioinform. 2013 May 1;10(03):645–56.

94. Smit A, Hubley R. RepeatModeler Open-1.0. 2008.

95. Keller O, Kollmar M, Stanke M, Waack S. A novel hybrid gene prediction method

employing protein multiple sequence alignments. Bioinformatics. 2011 Mar 15;27(6):757–

63.

96. Korf I. Gene finding in novel genomes. BMC Bioinformatics. 2004 May 14;5(1):59.

97. Campbell MS. compare_annotations_3.2.pl [Internet]. 2015. Available from:

https://github.com/mscampbell/Genome_annotation/blob/master/compare_annotations_3.2.

pl

98. Eilbeck K, Moore B, Holt C, Yandell M. Quantitative measures for the management and

comparison of annotated genomes. BMC Bioinformatics. 2009 Feb 23;10(1):67.

99. Liu D, Hunt M, Tsai IJ. Inferring synteny between genome assemblies: A systematic

evaluation. BMC Bioinformatics. 2018 Jan;19(1):26.

100. Emms DM, Kelly S. OrthoFinder: Solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. Genome Biol. 2015

Aug 6;16(1):157.

101. Katoh K, Misawa K, Kuma K ichi, Miyata T. MAFFT: A novel method for rapid multiple

sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002

Jul;30(14):3059–66.

102. Lefort V, Desper R, Gascuel O. FastME 2.0: A comprehensive, accurate, and fast distance-

based phylogeny inference program. Mol Biol Evol. 2015 Oct 1;32(10):2798–800.

103. Farrer RA. Synima: A synteny imaging tool for annotated genome assemblies. BMC

Bioinformatics. 2017 Nov 21;18(1):507.

104. Wang Y, Tang H, DeBarry JD, Tan X, Li J, Wang X, et al. MCScanX: A toolkit for

detection and evolutionary analysis of gene synteny and collinearity. Nucleic Acids Res.

2012 Apr;40(7):e49.

105. Zhao T, Schranz ME. Network approaches for plant phylogenomic synteny analysis. Curr

Opin Plant Biol. 2017 Apr 1;36:129–34.

106. Zhao T, Holmer R, de Bruijn S, Angenent GC, van den Burg HA, Schranz ME.

Phylogenomic synteny network analysis of MADS-Box transcription factor genes reveals

lineage-specific transpositions, ancient tandem duplications, and deep positional

conservation. Plant Cell. 2017 Jun 1;29(6):1278–92.

107. Zhao T, Schranz ME. Network-based microsynteny analysis identifies major differences

and genomic outliers in mammalian and angiosperm genomes. Proc Natl Acad Sci. 2019

Feb 5;116(6):2165–74.

108. Sievers F, Higgins DG. Clustal Omega. Curr Protoc Bioinforma. 2014;48(1):3.13.1-

3.13.16.

109. Suyama M, Torrents D, Bork P. PAL2NAL: Robust conversion of protein sequence

alignments into the corresponding codon alignments. Nucleic Acids Res. 2006 Jul

1;34(suppl_2):W609–12.

110. Wong WSW, Yang Z, Goldman N, Nielsen R. Accuracy and power of statistical methods

for detecting adaptive evolution in protein coding sequences and for identifying positively

selected sites. Genetics. 2004 Oct 1;168(2):1041–51.

111. Löytynoja A, Vilella AJ, Goldman N. Accurate extension of multiple sequence alignments

using a phylogeny-aware graph algorithm. Bioinformatics. 2012 Jul 1;28(13):1684–91.

112. Jones DT. Protein secondary structure prediction based on position-specific scoring

matrices. J Mol Biol. 1999 Sep 17;292(2):195–202.

113. Buchan DWA, Jones DT. The PSIPRED protein analysis workbench: 20 years on. Nucleic

Acids Res. 2019 Jul 2;47(W1):W402–7.

114. Ewels PA, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A, et al. The nf-core

framework for community-curated bioinformatics pipelines. Nat Biotechnol. 2020

Mar;38(3):276–8.

115. Andrews S. FastQC: A quality control tool for high throughput sequence data. [Internet].

2010. Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc

116. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal. 2011 May 2;17(1):10–2.

117. Kopylova E, Noé L, Touzet H. SortMeRNA: Fast and accurate filtering of ribosomal RNAs

in metatranscriptomic data. Bioinformatics. 2012 Dec 1;28(24):3211–7.

118. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-

aware quantification of transcript expression. Nat Methods. 2017 Apr;14(4):417–9.

119. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014 Dec 5;15(12):550.

120. Lindblad-Toh K, Garber M, Zuk O, Lin MF, Parker BJ, Washietl S, et al. A high-resolution

map of human evolutionary constraint using 29 mammals. Nature. 2011

Oct;478(7370):476–82.

121. Weyrich A, Schüllermann T, Heeger F, Jeschek M, Mazzoni CJ, Chen W, et al. Whole

genome sequencing and methylome analysis of the wild guinea pig. BMC Genomics. 2014

Nov 28;15(1):1036.

122. Gossmann TI, Ralser M. *Marmota marmota*. Trends Genet. 2020 May;36(5):383–4.

123. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Microtus ochrogaster. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/72443

124. Mouse Genome Sequencing Consortium, Waterston RH, Lindblad-Toh K, Birney E,

Rogers J, Abril JF, et al. Initial sequencing and comparative analysis of the mouse genome.

Nature. 2002 Dec 5;420(6915):520–62.

125. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Jaculus jaculus. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/72445

126. Gibbs RA, Weinstock GM, Metzker ML, Muzny DM, Sodergren EJ, Scherer S, et al.

Genome sequence of the Brown Norway rat yields insights into mammalian evolution.

Nature. 2004 Apr;428(6982):493–521.

127. Kolmogorov M, Armstrong J, Raney BJ, Streeter I, Dunn M, Yang F, et al. Chromosome

assembly of large and complex genomes using multiple references. Genome Res. 2018 Nov

1;28(11):1720–32.

128. Lilue J, Doran AG, Fiddes IT, Abrudan M, Armstrong J, Bennett R, et al. Sixteen diverse

laboratory mouse reference genomes define strain-specific haplotypes and novel functional

loci. Nat Genet. 2018 Nov;50(11):1574–83.

129. Couger MB, Arévalo L, Campbell P. A high quality genome for *Mus spicilegus*, a close

relative of house mice with unique social and ecological adaptations. G3

GenesGenomesGenetics. 2018 May 24;8(7):2145–52.

130. Chinese hamster CHOK1GS assembly and gene annotation. Horiz Eagle [Internet]. 2017;

Available from: https://www.ensembl.org/Cricetulus_griseus_chok1gshd/Info/Annotation

131. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Mesocricetus auratus. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/77669

132. Lassance JM, Hopi Hoekstra. Improved assembly of the deer mouse *Peromyscus*

maniculatus genome. Harv Univ Hughes Med Inst [Internet]. 2018; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/494228

133. Fang X, Nevo E, Han L, Levanon EY, Zhao J, Avivi A, et al. Genome-wide adaptive

complexes to underground stresses in blind mole rats *Spalax*. Nat Commun. 2014 Jun

3;5(1):3966.

134. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Octodon degu. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/74595

135. Keane M, Craig T, Alföldi J, Berlin AM, Johnson J, Seluanov A, et al. The naked mole rat

genome resource: Facilitating analyses of cancer and longevity-related adaptations.

Bioinforma Oxf Engl. 2014 Dec 15;30(24):3558–60.

136. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Chinchilla lanigera. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/68239

137. V. Federov, Dalen L, Olsen RA, Goropashnaya AV, Barnes BM. The genome of the Arctic

ground squirrel *Urocitellus parryii*. Inst Arct Biol [Internet]. 2018; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/477386

138. Di Palma F, Alföldi J, Johnson J, Berlin A, Gnerre S, Jaffe D, et al. The draft genome of

Ictidomys tridecemlineatus. Broad Inst [Internet]. 2012; Available from:

https://www.ncbi.nlm.nih.gov/bioproject/61725

139. Schneider VA, Graves-Lindsay T, Howe K, Bouk N, Chen HC, Kitts PA, et al. Evaluation

of GRCh38 and de novo haploid genome assemblies demonstrates the enduring quality of

the reference assembly. Genome Res. 2017 May 1;27(5):849–64.

1066