Maternal body condition affects the response of the gut microbiome to a widespread contaminant in larval spined toads

Tartu Sabrina*[1], Pollet Nicolas[2], Clavereau Isabelle [2], Bouchard Gauthier[2], Brischoux François[1]

[1]Centre d'Etudes Biologiques de Chizé (CEBC), UMR 7372 CNRS-La Rochelle Université, 79360 Villiers-en-Bois, France

[2]Université Paris-Saclay, CNRS, IRD, Evolution Génomes Comportement et Ecologie, Institut Diversité Ecologie et Evolution du Vivant, 12 route 128, 91190 Gif-sur-Yvette, France.

*Corresponding author: tartu.sabrina@gmail.com

Abstract

Glyphosate's primary metabolite, aminomethylphosphonic acid (AMPA) is the most detected

pollutant in surface waters. Recent studies have raised concerns regarding its toxicity, yet the

involved underlying mechanisms remain poorly understood. A disruption of the gut

microbiome, which plays a crucial role in host health, could be the mediator of most adverse

effects. We investigated the impact of AMPA exposure on the gut microbiome of spined toad

tadpoles (*Bufo spinosus*). We hypothesized that AMPA could alter the gut microbiota

composition and that these effects could depend on the microbiota source. We exposed tadpoles

to minute concentrations of AMPA and analyzed their faecal microbiota using 16S rRNA gene

sequencing as a proxy of the gut microbiota. AMPA exposure decreased the gut bacterial

biomass and affected bacterial community composition of tadpole's faeces. Furthermore, we

observed interactions between AMPA exposure and maternal body condition on the

Bacteroidota and Actinobacteriota phyla abundances. This suggests a maternal effect on early-

life microbial colonizers that could influence the response of the gut microbiome to AMPA.

These findings highlight the importance of considering the gut microbiome when studying the

effects of environmental contaminants. Further research is needed to elucidate the long-term

implications of this microbiome alteration for amphibian health.

Keywords: Gut microbiota; *Bufo spinosus*; Amphibian; AMPA; Aminomethylphosphonic
acid; Microbial vertical transfer

INTRODUCTION

Over the years, most of the highly toxic and persistent pesticides have been legally banned, such as the notorious DDT, and replaced with other, faster degrading and more species-specific, compounds. However, several current-used pesticides and/or their breakdown products still pose ecotoxicological issues (Gonçalves et al., 2021). Aminomethylphosphonic acid (AMPA, CAS No. 1066-51-9) is one of those breakdown products that may pose higher risks than its parents compounds (Grandcoin et al., 2017). AMPA has two primary sources, amino polyphosphates and glyphosate, but its origin in surface water and groundwater is mostly linked to the latter (Carles et al., 2019; Struger et al., 2015). Importantly, AMPA is detected much more frequently (20-50% more detected) and is more persistent in the environment (half-life 2-8 times longer) than glyphosate (Duke, 2020; Kolpin et al., 2006; Maggi et al., 2020; Ojelade et al., 2022).

In aquatic organisms, the negative effects of AMPA exposure are controversial, ranging from low toxicity to numerous adverse effects, including increased mortality, development delay, increased morphological abnormalities, genotoxicity, increased oxidative stress, cardiac defects, hepatic inflammatory response or changes in metabolic activity (Antunes et al., 2017; Barreto et al., 2023; Cheron et al., 2022; Cheron and Brischoux, 2023, 2020; Guilherme et al., 2014; Ivantsova et al., 2022; Levine et al., 2015; Martins-Gomes et al., 2022; Matozzo et al., 2018; Tartu et al., 2022; Zhang et al., 2021). However, despite the numerous adverse effects reported on non-target organisms, AMPA is not under particular scrutiny, and the underlying mechanisms involved are poorly identified.

The gut microbiome is considered as a key endpoint for ecotoxicological studies (Claus et al., 2016; Evariste et al., 2019). The trillions of microorganisms that have colonized an host compose its microbiota, with more than 90% of this commensal, mutualistic and symbiotic microbe community located in the hosts' gut (Sharpton, 2018). A large number of studies have

underlined the functional importance of the gut microbiota composition and structure in food

digestion, nutrient synthesis, host's physiology, development, behaviour, or immune system

performance (Clemente et al., 2012; Grond et al., 2018; Jiménez and Sommer, 2017); and a

dysbiosis (i.e. a modification in the composition and function of the gut microbiota) can alter

intestinal permeability, affect physiological performances and immune response, increasing

disease susceptibility (Gomaa, 2020; Grond et al., 2018; Warne et al., 2019; Xiong et al., 2019),

which could lead to unforeseen hazardous consequences for wild populations.

While several studies on vertebrates and invertebrates have tested the effects of glyphosate

exposure on gut microbiota composition (Blot et al., 2019; Cuzziol Boccioni et al., 2023; Ding

et al., 2021; Fréville et al., 2022; Iori et al., 2020; Lehman et al., 2023; Lozano et al., 2018;

Mesnage et al., 2021; Motta et al., 2018; Owagboriaye et al., 2021; Puigbò et al., 2022;

Ruuskanen et al., 2020; Walsh et al., 2023), only two have focused on the effects of AMPA on

invertebrate models, and reported slight effects on gut microbiota (Blot et al., 2019; Iori et al.,

2020). Considering the higher presence of AMPA in the environment and its ability to affect

microorganism growth by inhibiting bacterial cell wall biosynthesis (Atherton et al., 1982;

Azam and Jayaram, 2016; Carles et al., 2019; Coupe et al., 2012; Poiger et al., 2017), there is

still a lack of research linking the harmful effects of AMPA to a gut microbiome dysbiosis.

Amphibians face a higher overall extinction rate than other vertebrates (Harfoot et al., 2021;

Hoffmann et al., 2010). Intensive agriculture is a significant threat to which they are exposed,

contributing to habitat loss and pollution (Harfoot et al., 2021; Rollins-Smith, 2020; Stuart et

al., 2004; Wake and Vredenburg, 2008). Their reproductive migrations, water-dependent

breeding, aquatic larval stages, highly permeable skin and limited movements make amphibians

particularly vulnerable to environmental changes and highly relevant models to study the effects

of environmental contamination (Langlois, 2021; Stebbins and Cohen, 1995).

Previous studies conducted on the spined toad, *Bufo spinosus*, have associated AMPA exposure

to numerous adverse effects such as higher embryonic and larval mortality, increased oxidative

stress in tadpoles (Cheron et al., 2022; Cheron and Brischoux, 2023, 2020) and altered

colouration in adult males (Tartu et al., 2023). In addition, in a companion study using the same

individuals as the present one, we reported that environmentally relevant concentrations of

AMPA (0.4 µg/L) led to increased deformity rate upon hatching and increased development

length, with effects depending on the habitat of origin of the parents (agricultural *versus* forest,

Tartu et al., 2022). We therefore hypothesized that the previously observed adverse effects of

AMPA on spined toad tadpoles could originate from a gut microbiota dysbiosis. In addition,

because in vertebrates with external fertilization, the gut microbiota composition depends on

both horizontal (e.g. habitat, diet, conspecifics) and vertical transmission (e.g. parents)

(Comizzoli et al., 2021; Murphy et al., 2023; Scalvenzi et al., 2020), we further predicted that

AMPA-mediated effects on the gut microbiota composition of tadpoles would be dependent on

parent's body-condition (as a proxy of parental diet and quality) and habitat of origin

(agricultural or forest).

MATERIAL AND METHODS

Fieldwork

The 120 spined toad tadpoles used presently are a subset of the 240 individuals employed in a

previous companion study (Tartu et al., 2022). Study sites, parent captures, and housing

conditions have been described elsewhere (Tartu et al., 2022). Briefly, between 28/01/21 and

22/02/21, we searched for spined toad amplectant pairs at night in four different sites (two

agricultural sites and two sites surrounded by woodlands, see supporting information in (Tartu

et al., 2022)). We collected the ten first spotted couples to guarantee comparable inter-site

individual quality. We then returned breeding pairs to the laboratory until the female laid her

eggs (**Figure S1**). After oviposition, we measured the snout-vent length of both parents with a

calliper and weighed them on a precision scale. We calculated the scaled mass index (SMI)

developed by Peig and Green (2009) to assess parental body condition as SMI accurately

reflects amphibian energy stores in anuran species (Băncilă et al., 2010; Landler et al., 2023;

MacCracken and Stebbings, 2012; Zhelev and Minchev, 2023). We described how we

categorized males and females into the thin and fat groups in the "statistical analyses" section

below. Finally, we released the pair in their breeding site after body measurements.

Housing conditions and AMPA treatment

We obtained six segments of 34 eggs for each clutch and placed each segment in individual

aquariums with 2L of dechlorinated tap water. Among the six segments, we exposed three to

$0.4 \ \mu g \ L^{-1}$ ($\pm 0.01 \ \mu g \ L^{-1}$) of AMPA (AMPA group), and we kept the remaining three in

dechlorinated tap water as a control group. We obtained the AMPA solution by dissolving

commercial crystalline powder (Aminomethylphosphonic acid, 99% purity, ACROS

ORGANICS™) with dechlorinated tap water. As evidenced by data available from French

national surveys conducted between 2020 and 2023 on 380 samples obtained from 67 rivers of

the Deux-Sèvres Region, AMPA concentrations ranged from 0.02 to $3.4 \ \mu g \ L^{-1}$ (Naïades, 2023),

the selected AMPA concentration ($0.4 \ \mu g \ L^{-1}$) in the present study represents actual

concentrations found in surface water in our study area.

Upon hatching, we randomly isolated one tadpole and released the remaining tadpoles in the

breeding pond of their parents. We housed each selected tadpole individually (n=240 in total,

see **Figure 1, S1**) in a 2L aquarium with either dechlorinated tap water or AMPA according to

the treatment experienced during embryonic development. Consequently, tadpoles were

exposed to the same treatment during embryonic and larval development. We checked tadpoles

daily and monitored their development through six key Gosner stages (25, 30, 37, 41, 42 and

46 (Gosner, 1960)). The effects of AMPA exposure on tadpole development according to the

parent's origin (forest vs agricultural) have been published previously (Tartu et al., 2022).

From the egg stage until metamorphosis, the tadpoles were kept under simulated 12:12 h day

and night in a room at 17°C to avoid any basal metabolism and development variation. Water

was changed weekly, adding AMPA at 0.4 µg L^{-1} for the contaminated aquariums. Upon

hatching, we fed tadpoles with organic ground spinach *ad libitum*.

Faecal sampling for microbiome analyses

During pre-metamorphic larval development, spined toad tadpoles are the most active

swimming and feeding at Gosner stage 37 (Cheron et al., 2021), resulting in a larger production

of faeces. We therefore started sampling tadpoles' faeces for gut microbiota analyses once they

reached Gosner stage 37. To do so, we collected faeces accumulated in the bottom of the

aquarium since the last water change (4-6 days) using sterile pipettes for half of the individuals

(n = 120). We placed faeces in a sterile petri dish, then transferred them by pipetting with filter

tips to a sterile microtube and added twice their volume of DNA/RNA Shield™ (Zymo

Research). To increase genetic diversity, we then pooled the faeces from three sibling tadpoles

receiving the same treatment (control or AMPA) in one tube, we thus obtained 40 pools (**Figure**

1). DNA/RNA Shield™ preserves the nucleic acids integrity of samples at ambient

temperatures, we therefore kept the pooled faecal samples at room temperature until analyses.

Figure 1: Common-garden experiment conducted on spined toad *Bufo spinosus* tadpoles. The upper panel shows the experimental design we used, and the metrics obtained for each Gosner stage (GS). From each clutch, three segments were exposed to control conditions (dechlorinated tap water) and three segments to AMPA (0.4 µg L^{-1}). From each segment that produced at least one viable tadpole, one was kept and monitored until metamorphosis (Gosner stage 46). Lower panel: on a sub-sample of tadpoles (n=120), we collected faeces of individuals when they reached GS 37, we then pooled the faeces of three siblings that had received the same treatment in a single tube (n=20 control, n=20 AMPA) to conduct microbiota analyses.

DNA extraction, libraries and sequencing

According to the manufacturer's instructions, we extracted DNA using the ZymoBIOMICS DNA/RNA Miniprep kits. We added 1 µL of the Zymobiomics Spike-in control I (High microbial Load D6320) to each sample before lysis using a Precellys Evolution Touch equipped with a cryolys module to perform six steps of bead beating at 10,000 rpm during 10 s at 0°C, with 30 s pause between each step. We included a set of samples from artificial communities

as positive controls (Zymobiomics Microbial community standard (D6300) and log distribution

(D6310)) and negative controls (tap water used to rear tadpoles and nuclease-free water). We

controlled the quality and quantity of extracted DNA using spectrophotometry (Nanodrop) and

fluorometry (Qubit). We amplified the V1-V8 portion of the 16S rDNA gene by PCR using

tailed primers tBACT27F (5'-

TTTCTGTTGGTGCTGATATTGCAGAGTTTGATCCTGGCTCAG-3') and tBACT1391R

(5'- ACTTGCCTGTCGCTCTATCTTCGACGGGCGGTGWGTRCA-3'). We ran PCR in

triplicate reactions of 25 µL assembled under a PCR laminar flow cabinet using the following

conditions: 40 ng DNA, 0.3µM each primer, 0.5 mM dNTPs, 1 Unit of tiAmplus DNA

Polymerase HotStart (Roboklon), 1X tiAmplus Buffer containing 25 mM $MgSO_4$. The PCR

program was 93°C for 2 min, followed by ten cycles of 10s at 93°C, 30s at 57°C, 120s at 68°C

and 25 cycles of 10 s at 93°C, 30s at 65°C, 120s at 68°C and 7 min at 68°C. We monitored

amplification products using regular gel electrophoresis before pooling the triplicate PCR and

purification using magnetic beads (Macherey-Nagel NucleoMag cleanup). We then used the

PCR barcoding expansion pack (EXP-PBC096, Oxford Nanopore Technology) and the ligation

sequencing kit (SQK-LSK109) to prepare a sequencing library of which 150 ng were loaded on

the flow cell (R9.1) and sequenced using a MinIon device (Oxford Nanopore Technology). We

made two libraries sequenced on two different flow cells yielding 5,043,247 and 5,184,836

usable reads after demultiplexing. The sequenced datasets generated and/or analyzed during the

current study are available in the EMBL Nucleotide Sequence Database (ENA) at

https://www.ebi.ac.uk/ena/data/view/PRJEB71117.

16s rDNA sequence analysis

We mapped individual reads using minimap2 on the SILVA SSU database version 138, and

retained only reads mapping with a coverage of 75%, and an alignment length between 1000

and 2000 nt to a SILVA entry identified by at least two reads in a given sample. Abundance

analyses were made using OTUs identified by a prevalence of at least four with a minimum

read depth of three.

Analysis of spike control

We analysed four samples as negative controls to monitor the inputs from the dechlorinated tap

water used to raise tadpoles (two samples) or any contaminants arising from sample

manipulation, DNA extraction, PCR amplification and sequencing library construction (two

samples). As in all samples, we included spiked bacteria (D6320, ZymoBIOMICS) to these

controls that were analysed in two independent sequencing runs. We obtained 19,270 and

28,848 reads in the water samples; and 26,579 and 38,633 reads in the kit samples, from which

respectively 98.25; 98.73; 99.77; and 99.71% were identified as the spiked bacteria *Imtechella*

and *Allobacillus*. The maximum number of reads from another OTU was between 11 and 140,

and we found at least one read attributed to 25-78 phylogenetically unrelated OTUs. The ratios

of *Allobacillus* to *Imtechella* reads were 0.84-1.05, compared with the theoretical expectation

of 2.3. In conclusion, as expected we found that a vast majority of the reads from these spike

control identified the two bacteria from the spike. Yet, we observed a trace level of reads

derived from phylogenetically unrelated OTUs that artificially increased the number of

observed OTUs. We observed that two samples out of seven expected to be devoid of spike-in

contained one and three reads from *Allobacillus* but none from *Imtechella*. The sample

containing one read may represent a case of spillover since it is located next to a D6320 sample.

We computed a spike in OTU abundance from the mean of 100 rarefactions to account for

sample read depth variations.

Statistical analyses

210   First, we obtained an estimation of the ratios of absolute endogenous bacteria by analyzing the

211   abundance of spike-in controls according to the treatment (AMPA vs control) and their

212   interaction with the parental capture site (agricultural vs forest) and parental (maternal and

paternal) body condition (Hornung et al., 2019; Stämmler et al., 2016). We used linear mixed

effect models (LME) with clutch identity as a random factor to do so. Then, we evaluated the

similarity of bacterial communities between the different treatments and their interaction with

the parental capture site and parental body condition with unweighted (based on the presence

or absence of observed bacterial taxa) and weighted (based on the abundance of observed

bacterial taxa) UniFrac distances (Lozupone et al., 2006). To compare spike-in control

abundances and bacterial communities according to treatment while considering parent body

condition, we constructed discrete body condition classes. To do so, we used an objective

categorization method by grouping all the individuals with an SMI ≤ median SMI value as

"thin" and all individuals with an SMI > median SMI value as "fat". We used the non-

parametric Kruskal-Wallis (KW) test when comparing UniFrac distances according to more

than two variables. Post-hoc analyses were performed using Dunn's test with the Bonferroni

adjustment method. We then performed principal coordinate analysis (PCoA) based on

unweighted and weighted UniFrac distances using the '*Phyloseq*' and '*ade4*' packages (Dray

and Dufour, 2007; McMurdie and Holmes, 2012). Second, we extracted the abundances of the

nine most represented phyla and inserted them into linear mixed effect models (LME) with

clutch identity as a random factor. We used LME to specifically analyze which phyla were

influenced by the treatment and potential interaction with other variables (e.g. parental habitat

and body condition). At last, we conducted Linear discriminant analysis Effect Size (LEfSe)

analyses within the phyla and variables previously identified as sensitive to AMPA exposure to

determine which microbiome biomarkers characterized the observed differences (Segata et al.,

2011) by using the '*microbiomeMarker*' package (Cao et al., 2022). We performed all analyses

with R v.4.3.2 (R Core Team, 2019).

RESULTS

Faecal microbiota composition of spined toad tadpoles

238 We identified 664 Operational Taxonomic Units (OTUs) within the 40 faecal samples analyzed.

239 Proteobacteria (537 OTUs) was the most abundant phylum, followed by Bacteroidota (73

240 OTUs), Actinobacteriota (18 OTUs), Firmicutes (13 OTUs), Verrumicrobiota (7 OTUs),

241 Acidobacteriota (7 OTUs), Campilobacterota (4 OTUs), Desulfobacterota (3 OTUs),

242 Planctomycetota (2 OTUs).

Treatment effects and their interactions with habitat and parental body condition

We observed a higher abundance of spiked-in bacteria in faecal samples from AMPA-exposed

tadpoles as compared to controls (LME, estimate: 20.93 ± 6.39, p=0.004, **Figure 2**), which

indicates an endogenous lower biomass in faecal samples of AMPA treated tadpoles. The

interaction of treatment with parental habitat and body condition was unrelated to spike-in

control abundances (LME, p>0.560 for all tests). Although AMPA treatment did not overall

affect faecal microbiota community structure (unweighted UniFrac distances, Kruskal-Wallis

χ^2= 1.99, df = 2, p = 0.370), it affected its composition (weighted UniFrac distances (χ^2= 11.0,

df = 2, p=0.004, post-hoc test AMPA vs Control: p=0.003).

Figure 2: Exogenous spike-in bacterial relative abundance in tadpoles' faecal samples.
The greater abundance of exogenous spiked-in bacteria in samples from AMPA-treated
tadpoles reveals an overall lower endogenous bacterial biomass than in the control group.

Bacterial abundances were computed as the mean value from 100 rarefactions. Significant
differences are represented by *** (p=0.004).

The effect of the treatment on the faecal microbiota community was not exacerbated when

considering the habitat of the parents (unweighted UniFrac, $\chi^2=11.2$, df=9, p=0.262; weighted

UniFrac, $\chi^2= 15.4$, df = 9, p = 0.080). Although close to statistical significance for weighted

UniFrac distances, pairwise post-hoc comparisons did not reveal any trend (p>0.21 for all tests).

Yet, the AMPA effect depended on maternal body condition (**Figure 3**). More specifically, the

structure and composition of faecal microbial communities were affected by AMPA in tadpoles

produced by females in better condition. In comparison, this effect was not found in tadpoles

produced by females characterized by lower condition (unweighted UniFrac distances, $\chi^2=46.2$,

df = 9, p = 5.46×10^{-07}, post-hoc test: AMPA vs Control in fatter females: p = 0.006, thinner

females: p =0.198; weighted UniFrac, $\chi^2=34.1$, df = 9, p = 8.53×10^{-05}, post-hoc test: AMPA vs

Control in fatter females: p =0.0004, thinner females: p = 0.999, **Figure 3A-D**). Paternal body

condition alone or in interaction with the treatment did not significantly influence the tadpole

microbiome (p>0.05 for all tests, **Figure 4**).

Figure 3: **Interaction of treatment and maternal body condition on GS 37 *Bufo spinosus* tadpoles' faecal microbiota community.** The upper panels represent calculated unweighted (A) and weighted (B) UniFrac distances according to the treatment (control in blue vs AMPA in red) and maternal body condition. Lower panels represent principal coordinate analysis (PCoA) for each pool of siblings (AMPA, n = 20, control group, n=20) according to treatment and maternal body condition. Closer dots in the PCoA figure indicate a more similar microbial community. The percentage of variation explained by principal coordinates (PC) is shown on the axes. We calculated body condition from the scaled mass index. Condition categories were separated by median value ('thinner' ≤ median value; 'fatter'> median value). Significant differences are represented by *** (p<0.007 in A and B, we performed pairwise comparisons using Dunn's all-pairs test).

Figure 4: **Interaction of treatment and paternal body condition on GS 37 _Bufo spinosus_ tadpoles' faecal microbiota community.** The upper panels represent calculated unweighted (A) and weighted (B) UniFrac distances according to the treatment (control in blue vs AMPA in red) and paternal body condition. Lower panels represent principal coordinate analysis (PCoA) for each pool of siblings (AMPA, n = 20, control group, n=20) according to treatment and paternal body condition. Closer dots in the PCoA figure indicate a more similar microbial community. The percentage of variation explained by principal coordinates (PC) is shown on the axes. Body condition was calculated from the scaled mass index. Condition categories were separated by median value ('thinner' ≤ median value; 'fatter'> median value).

Interaction between treatment and maternal body condition on tadpole's microbiome

In AMPA-exposed tadpoles, the abundance of Bacteroidota decreased with increasing maternal body condition (**Figure 5A, Table 1**), whereas in control tadpoles, we observed a positive relationship between Bacteroidota abundance and maternal body condition (**Figure 5A, Table 1**).

Table 1: Relationships between the abundance of the nine major phyla sequenced in tadpole's faecal microbiome, according to AMPA exposure and maternal body condition. Values were obtained using linear mixed effects models with clutch identity as a random factor. Values in bold are significant at the 0.05 level.

Phylum abundance	Treatment				SMI				Treatment × SMI			
	Estimate ± SE	df	t	p	Estimate ± SE	df	t	p	Estimate ± SE	df	t	p
Proteobacteria	1727 ± 63057	18	0.03	0.978	432 ± 791	18	0.5464299	0.592	-148 ± 796	18	-0.19	0.855
Bacteroidota	**65703 ± 26942**	**18**	**2.44**	**0.025**	205 ± 251	18	0.8167347	0.425	**-846 ± 340**	**18**	**-2.48**	**0.023**
Actinobacteriota	**-742.2 ± 238.6**	**18**	**-3.11**	**0.006**	-3.52 ± 2.64	18	-1.334115	0.199	**9.70 ± 3.01**	**18**	**3.22**	**0.005**
Firmicutes	-91 ± 224	18	-0.41	0.690	1.01 ± 2.07	18	0.4872227	0.632	1.37 ± 2.83	18	0.48	0.635
Verrucomicrobiota	-1048 ± 624	18	-1.68	0.111	-6.7 ± 5.6	18	-1.201961	0.245	12.9 ± 7.9	18	1.63	0.120
Acidobacteriota	18.7 ± 21.5	18	0.87	0.397	0.11 ± 0.21	18	0.52	0.610	-0.25 ± 0.27	18	-0.92	0.369
Campylobacterota	-18.4 ± 19.3	18	-0.95	0.353	-0.29 ± 0.18	18	-1.6236091	0.122	0.26 ± 0.24	18	1.05	0.309
Desulfobacterota	-9.45 ± 18.6	18	-0.51	0.618	-0.24 ± 0.24	18	-1.0182636	0.322	0.07 ± 0.24	18	0.31	0.758
Planctomycetota	-133 ± 116	18	-1.15	0.267	-1.38 ± 1.04	18	-1.326659	0.201	1.55 ± 1.47	18	1.05	0.306

In contrast, Actinobacteriota abundance increased with increasing maternal body condition in AMPA-exposed tadpoles, while the Actinobacteriota/maternal body condition relationship was negative in control tadpoles (**Figure 5B, Table 1**). The abundance of the seven other phyla was unrelated to treatment, SMI and their interaction (**Table 1**).

Figure 5: Relationships between tadpoles' faecal phylum abundance and maternal body condition according to AMPA exposure. Bacteroidota (A) and Actinobacteriota (B) abundances (number of reads) were differently associated with maternal body condition as inferred by their scaled mass index (SMI) according to AMPA exposure. Each red triangle (AMPA exposed) and blue dot (control) represent a pool of faeces obtained from three siblings exposed to a similar treatment.

The effects of AMPA on microbiota composition were stronger in offspring produced by fatter females (**Figures 3, 5**) within the phyla Bacteroidota and Actinobacteriota (**Table 1**). We

therefore conducted LEfSe analyses within this subset (i.e. Bacteroidota and Actinobacteriota

in the 'fatter' female group, **Table 2**). Within the Bacteroidota phylum, ten markers were

identified for the AMPA exposed group (**Table 2**). The most features that explain the AMPA

group were class Bacteroidia, order Flavobacteriales, family *Weeksellaceae*, genus

Cloacibacterium, species *cloacibacterium* uncultured, and class Bacteroidia, order

Sphingobacteriales, family *KD3-93*, genus *KD3-93 uncultured*. Whereas within the

Actinobacteriota phylum, four markers were identified in the AMPA group (**Table 2**), all

explained by class Actinobacteria, order Streptomycetales, family *Streptomycetaceae*, genus

Streptomyces and species *streptomyces*.

Table 2: LEfSe analysis of taxonomic biomarkers of gut microbiota influenced by AMPA exposure in offspring produced by fatter females. LEfSe analysis identified the most differentially abundant taxa within a) Bacteroidota and b) Actinobacteriota according to AMPA exposure. We show LDA scores > 2; k=kingdom, p=phylum, c=class, o=order, f=family, g=genus, s=species, ef. LDA= effect size linear discriminant analysis.

Marker	Feature	Enriched group	ef. LDA	p-value
a) Bacteroidota (Fatter mothers)				
Marker1	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Flavobacteriales|f__Weeksellaceae	AMPA	3.383	**0.008**
Marker2	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Sphingobacteriales|f__KD3-93	AMPA	3.336	**0.041**
Marker3	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Sphingobacteriales|f__KD3-93|g__KD3-93_uncultured	AMPA	3.336	**0.041**
Marker4	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Sphingobacteriales|f__KD3-93|g__KD3-93_uncultured|s__KD3-93_uncultured_uncultured	AMPA	3.336	**0.041**
Marker5	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Flavobacteriales|f__Weeksellaceae|g__Cloacibacterium	AMPA	3.245	**0.049**
Marker6	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Flavobacteriales|f__Weeksellaceae|g__Cloacibacterium|s__Cloacibacterium_uncultured	AMPA	3.245	**0.049**
Marker7	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Flavobacteriales|f__Weeksellaceae|g__Weeksellaceae_uncultured	AMPA	2.820	**0.013**
Marker8	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Flavobacteriales|f__Weeksellaceae|g__Weeksellaceae_uncultured|s__Weeksellaceae_uncultured_unc	AMPA	2.820	**0.013**
Marker9	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Flavobacteriales|f__Flavobacteriaceae|g__Leptobacterium	AMPA	2.711	**0.028**
Marker10	k__Bacteria|p__Bacteroidota|c__Bacteroidia|o__Flavobacteriales|f__Flavobacteriaceae|g__Leptobacterium|s__Leptobacterium	AMPA	2.711	**0.028**
b) Actinobacteriota (Fatter mothers)				
Marker1	k__Bacteria|p__Actinobacteriota|c__Actinobacteria|o__Streptomycetales|f__Streptomycetaceae	AMPA	4.015	**0.014**
Marker2	k__Bacteria|p__Actinobacteriota|c__Actinobacteria|o__Streptomycetales|f__Streptomycetaceae|g__Streptomyces|s__Streptomyces	AMPA	4.012	**0.014**
Marker3	k__Bacteria|p__Actinobacteriota|c__Actinobacteria|o__Streptomycetales	AMPA	4.012	**0.014**
Marker4	k__Bacteria|p__Actinobacteriota|c__Actinobacteria|o__Streptomycetales|f__Streptomycetaceae|g__Streptomyces	AMPA	4.009	**0.014**

DISCUSSION

We here show that a minute concentration of AMPA (aminomethylphosphonic acid) - the primary metabolite of glyphosate and the main contaminant detected in surface waters worldwide - can affect gut microbiota biomass and community composition in larvae of a widespread amphibian species. Interestingly, this effect was, at least partly, mediated by maternal condition, as the effects of AMPA on tadpoles' microbiome were exacerbated in individuals produced by females with better body condition. AMPA did not affect the gut microbiome of tadpoles from leaner females, and paternal body condition was unrelated to the effects of AMPA. The interaction between AMPA and maternal body condition on tadpoles' gut microbiome was driven by contrasting alterations of the abundance of Bacteroidota and Actinobacteriota, which are respectively the second and the third most abundant phylum in spined toad larval faecal microbiome.

Possible vertical transmission of the gut microbiota

In amphibian species with aquatic larvae, gut microbiota majorly originates from the environment (i.e. water and diet), and the contribution of parental microbiomes was thought to be minor (Hernández-Gómez and Hua, 2023; Prest et al., 2018; Scalvenzi et al., 2020). In this study, amplectant pairs were captured in four different sites then were placed in tank filled with dechlorinated tap water until all females laid their eggs. The tap water was controlled, and analyses indicated that tap water did not contribute to significant amounts of bacterial DNA. We exposed egg strings to the same water and fed tadpoles with the same diet of organic ground spinach. Therefore, the only source of variation in early microbial egg colonizers were those present on the parents (skin and cloacal microbiomes), and we were able to show evidence of a maternal signature on the tadpole faecal microbiome.

Vertical transmission has been extensively studied in mammals, with maternal faecal microbes transferred to newborns during birth (Ferretti et al., 2018; Wampach et al., 2018; Wang et al., 2020). Although prenatal transfer is still debated in oviparous vertebrates, evidence of vertical transmission has been reported. For instance, bacterial colonization during egg formation has been suggested in Eastern Fence Lizard (*Sceloporus undulatus*) (Trevelline et al., 2018), and female *Sceloporus virgatus* lizards transfer beneficial microbes from their cloaca onto their eggs during oviposition with beneficial effects on the offspring (Bunker et al., 2021). Moreover, hatchling loggerhead sea turtles *Caretta caretta* harboured distinct microbial communities with respect to sand and eggshells, suggesting here also a maternal origin of their pioneer gut microbiome (Vecchioni et al., 2022). In addition, the faecal microbiome of neonate Rock pigeons (*Columba livia*) hatched in an incubator resembled the cloacal microbiome of females sampled from the same population (Dietz et al., 2020). Evidence is accumulating from pathogen transmission studies that *Salmonella enterica* contamination of chicken eggs does not occur from penetration through the shell but by the passage from the hen's intestinal tract to the reproductive tract, then from pathogen colonization into the forming egg on the vitelline membrane, in the egg white or the shell membranes (Gantois et al., 2009). Maternal intestinal microbiota could therefore colonize the egg yolk before shell deposition.

In contrast to amniotic vertebrates, amphibians produce jelly-coated eggs. Egg-jelly has various functions such as fertilization, insulation, gas exchange, and protection (Altig and McDiarmid, 2007; Beattie, 1980; Burggren, 1985; Olson and Chandler, 1999), yet some pathogenic bacteria can penetrate the thick jelly layer of the egg (Khalifa et al., 2021). Since vertical transmission is possible in shelled eggs, a maternal transmission in non-shelled eggs is even more likely, as observed in crustaceans (Giraud et al., 2022). But surprisingly, few researchers have investigated vertical transmission in amphibians without parental care (Hughey et al., 2017). In African clawed frogs (*Xenopus tropicalis*), the environment was identified as the primary driver

of egg bacterial communities by contributing around 70% of the bacteria in controlled

conditions. In the same study, the skin and feces of parents were identified as minor

contributors(Scalvenzi et al., 2020). In wild boreal toad populations (*Anaxyrus boreas*), a

quarter of the bacterial communities observed on eggs and a third of communities observed on

early-stage tadpoles were comprised of bacteria acquired from an unknown source (neither

water, upland soil nor sediments), and these strains were likely parentally transmitted (Prest et

al., 2018).

Although further studies would be needed to better understand the vertical transmission of the

gut microbiota in spined toads, substantial evidence points towards this direction. As observed

in other taxa, such as mammals and reptiles, vertically transmitted strains are likely to be more

ecologically relevant for the offspring compared with non-maternal strains (Ferretti et al.,

2018), and transmitted strains could confer fitness advantages to the progeny (Bunker et al.,

2021; Trevelline et al., 2018).

Mechanisms of transgenerational immune priming could provide complementary explanations

(Roth et al., 2018). There is complementary evidence for the transmission of innate immunity

compounds (i.e. antimicrobial skin peptides and mutualistic microbiota) from females to eggs

in amphibians (Walke et al., 2011). Yet, more studies are needed to investigate and quantify

immune responses transmitted to offsprings in amphibians.

Tadpole gut microbiome depends on the maternal body condition

Body condition can be a good indicator of fitness in numerous species (Bowers et al., 2014;

Bright Ross et al., 2021; Liu et al., 2020; Milner et al., 2003), including spined toads (Renoirt

et al., 2022). Associations between body condition and gut microbiota composition have been

well described in human and rodent models, and this is not surprising given the function of the

gut microbiome to produce metabolites involved in energy homeostasis and metabolic health

(Aron-Wisnewsky et al., 2021.; Fan and Pedersen, 2021; Moreno-Navarrete and Fernandez-

Real, 2019; Turnbaugh et al., 2006; Zwartjes et al., 2021); however, in the wild, these

relationships are more difficult to characterize.

For instance, no associations between the gut microbiome and body condition were reported in

fire salamanders *Salamandra Salamandra*, Seychelles warbler *Acrocephalus sechellensis*,

three-spined stickleback *Gasterosteus aculeatus* (Friberg et al., 2019; Wang et al., 2021;

Worsley et al., 2021). In contrast, in Eurasian perch *Perca fluviatilis*, lower microbial diversity

was related to improved condition; in great tit nestlings *Parus major*, a time-lagged association

was observed between gut microbiota composition, nestling weight and survival; in coyotes

Canis latrans, the consumption of anthropogenic food in urban individuals was associated with

increased microbiome diversity, higher abundances of *Streptococcus* and *Enterococcus* and

poorer average body condition (Bolnick et al., 2014; Davidson et al., 2021; Sugden et al., 2020),

and in wood frogs, *Rana sylvatica* egg microbiome manipulation accelerated larvae growth and

development rates (Warne et al., 2019). These contrasted relationships between gut microbiota

composition and body condition in wild species could result from environment-dependent

variations in feeding activity, diet composition, and body condition, and all these features can

also vary according to sex, age, and breeding cycle. Measuring microbiome-fitness

relationships at just one time point could therefore be misleading in free-ranging species.

In the present study, we observed an association between tadpole faecal microbiome

composition and maternal body condition which was affected by AMPA exposure. This

suggests that components of the maternal microbiome or determinants of microbiota

composition were transmitted to the eggs during oviposition, and that this specific microbiome

signature was more sensitive to AMPA exposure. We can hypothesize that a gut microbiota

composition more efficient in harvesting energy, as suggested by the higher maternal body

condition in that group, would consist of a gut bacterial assemblage with species more sensitive to AMPA. One may hypothesize that those females in better body condition originate from habitats preserved from AMPA exposition (i.e. forest sites, see Tartu et al. (2022)), and that their gut bacterial composition would be more sensitive to AMPA-exposure, however maternal body-condition was not related to habitat (agricultural vs. forested, LME estimate: -2.31 ± 3.69, p = 0.538). The observed relationships underline the dependence of the gut microbiome on interactions among other deterministic factors that were not accounted for in this study, such as host genetic and epigenetic background, age, diet or other environmental stressors (Chen et al., 2022; Shu et al., 2019; Song et al., 2021; Zhou et al., 2021). Nevertheless, we were able to identify a modification of the abundance of two major phyla in tadpoles gut microbiota, Bacteroidota and Actinobacteriota, which varied according to both maternal body condition and AMPA exposure.

Effects of agrochemicals according to gut microbiota composition

In line with our hypothesis, AMPA exposure affected tadpoles' gut microbiome by reducing bacterial biomass and changing community composition. In several taxa, including amphibians, a dysbiosis induced by decreased bacterial biomass is associated with deficient nutrient absorption and impaired immunity (Adamovsky et al., 2018; Gomaa, 2020; Jiménez and Sommer, 2017). In addition, the AMPA – microbiome relationship was exacerbated when considering transgenerational traits, such as maternal body condition, highlighting microbial colonizers' importance in susceptibility to pollutants. Specifically, in tadpoles from better-condition females, we observed a weaker Bacteroidota abundance and a more substantial Actinobacteriota abundance in the AMPA-exposed group.

As previously mentioned only a few studies conducted on invertebrates have investigated whether AMPA exposure would lead to a gut microbiota dysbiosis (Blot et al., 2019; Iori et al., 2020), yet, effects of glyphosate exposure similar to those observed in the present study have

been reported in Sprague-Dawley rats (Mesnage et al., 2021). For instance, glyphosate exposure decreased Bacteroidota abundance and concomitantly increased Firmicutes and Actinobacteria abundances in rats' caecum microbiome (Mesnage et al., 2021). The reported effects of glyphosate could as well be the consequence of AMPA, as glyphosate is degraded to AMPA in vertebrates and highly accumulates in the intestine, as observed in bird and fish models (Fréville et al., 2022; Yan et al., 2023).

The ability to cleave the C - P bond of AMPA and to use it as a phosphorus source is widespread in bacteria (Dick and Quinn, 1995; Fox and Mendz, 2006; Harkness, 1966; Studnik et al., 2015) including various species of the Actinobacteriota phylum as *Streptomyces* (Obojska et al., 1999; Obojska and Lejczak, 2003). The observed increase of Actinobacteriota is likely to result from the ability of *Streptomyces* and other species to utilize AMPA as a phosphorus source, which growth will therefore be promoted. In contrast, the decreased abundance of Bacteroidota in tadpole gut microbiota, and more specifically the orders Sphingobacteriales (family *KD3-93*) and Flavobacteriales (family *Weeksellaceae*, genus *Cloacibacterium*) could either result from a higher sensitivity of these orders to AMPA, a modification of the gut environment which became less favourable to their growth (e.g. pH) or resource competition with Actinobacteriota (Firrman et al., 2022). This gut microbiota alteration associated to AMPA exposure can have important implications for the host health.

For instance, Sphingobacteriales is a taxonomic class composed of a single order of environmental bacteria that are capable of producing sphingolipids (An et al., 2011; Olsen and Jantzen, 2001). Bacterial sphingolipids have the ability to migrate from epithelial cells to various host organs, thereby regulating its immune system and lipid metabolism (Bai et al., 2023). Flavobacteriales on the other hand, play several roles in various metabolic pathways, such as biotin metabolism, nitrogen metabolism, vitamins, amino-acid and fatty acid

biosynthesis (Rosas-Pérez et al., 2014; Yang et al., 2017; Zhou et al., 2022) and can bare positive effects on the host growth and development (Pan et al., 2023). At the genus level, *Cloacibacterium* sp. is able to degrade cellulose, and may have a critical role in transforming plant-derived complex dietary carbohydrates into essential short-chained fatty acids (SCFA) for herbivores organisms such as spined toad tadpoles (Flint et al., 2012; Fujimori, 2021; Hu et al., 2021; Martens et al., 2011; Zhang et al., 2018).

A disruption of essential SCFA production, which represent an essential energy source for the host, could lead to impaired metabolic functions, as observed in *Rana chensinensis* tadpoles (Shen et al., 2022), and altered metabolic processes could lead to delayed growth. In the crucian carp (*Carassius auratus*), for instance, glyphosate exposure resulted in a dysbiosis of Bacteroidota at the phylum level, and Bacteroidota abundance was negatively correlated with different metrics of growth performance (condition factor, fat ratio and specific growth rate) (Yan et al., 2022).

We have previously reported that AMPA exposure in tadpoles from the present study, increased deformity rate in individuals originating from AMPA-free forest and increased development length in AMPA-exposed individuals originating from agricultural sites (Tartu et al., 2022). While embryonic stages may be more sensitive to AMPA exposure in forest individuals (AMPA-preserved population) they might be more resilient to a gut microbiome dysbiosis as no further effects on fitness were observed (Tartu et al., 2022). In contrast, agricultural individuals (AMPA-exposed population) could be more resistant during embryonic stages, but still, a gut microbiome dysbiosis, could lead to SCFA alteration and thus a longer duration to develop (Tartu et al., 2022). These findings once more underline the part that may play the hosts genotype in shaping the consequences of a gut microbiota dysbiosis.

ACKNOWLEDGEMENT

The authors thank Matthias Renoirt and Marion Cheron for their help on the field and tadpole monitoring and Léa Lorrain-Soligon for providing an amplexus cropped picture, and two anonymous referees for their constructive comments.

FUNDING

Funding was provided by the CNRS, the Agence de l'Eau Loire-Bretagne, the Agence de l'Eau Adour-Garonne, the Région Nouvelle-Aquitaine (Aquastress 2018-1R20214, Amphitox 2019-1R20216), the ANSES (BiodiTox project # 2019/1/031), The Plan d'Action National ECOPHYTO (n°OFB-21-0941).

AUTHOR CONTRIBUTION

F.B. obtained the funding and supervised the study; F.B., N.P. and S.T. designed the study; S.T., G. B. and I.C. generated the libraries; N.P. performed bioinformatic analyses; S.T. collected, analyzed, interpreted the data, wrote the first version of the manuscript. All authors have revised and approved the submitted version of the manuscript.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

ETHICS STATEMENT

We followed all applicable institutional and national guidelines for the care and use of animals. This work was approved by the French authorities (COMETHEA ethic committee and Ministère de L'Enseignement Superieur, de la Recherche et de L'innovation) under permits APAFIS#13477–2018032614077834 and DREAL/2020D/8041.

REFERENCES

Adamovsky, O., Buerger, A.N., Wormington, A.M., Ector, N., Griffitt, R.J., Bisesi Jr., J.H., Martyniuk, C.J., 2018. The gut microbiome and aquatic toxicology: An emerging concept for environmental health. Environ. Toxicol. Chem. 37, 2758–2775. https://doi.org/10.1002/etc.4249

Altig, R., McDiarmid, R.W., 2007. Morphological diversity and evolution of egg and clutch structure in amphibians. Herpetol. Monogr. 21, 1–32.

An, D., Na, C., Bielawski, J., Hannun, Y.A., Kasper, D.L., 2011. Membrane sphingolipids as essential molecular signals for Bacteroides survival in the intestine. Proc. Natl. Acad. Sci. 108, 4666–4671. https://doi.org/10.1073/pnas.1001501107

Antunes, A.M., Rocha, T.L., Pires, F.S., de Freitas, M.A., Leite, V.R.M.C., Arana, S., Moreira, P.C., Sabóia-Morais, S.M.T., 2017. Gender-specific histopathological response in guppies Poecilia reticulata exposed to glyphosate or its metabolite aminomethylphosphonic acid. J. Appl. Toxicol. 37, 1098–1107. https://doi.org/10.1002/jat.3461

Aron-Wisnewsky, J., Warmbrunn, M.V., Nieuwdorp, M., Clément, K., 2021. Metabolism and Metabolic Disorders and the Microbiome: The Intestinal Microbiota Associated With Obesity, Lipid Metabolism, and Metabolic Health—Pathophysiology and Therapeutic Strategies. Gastroenterology, The Gut microbiome: Reaching the Promise through Discovery 160, 573–599. https://doi.org/10.1053/j.gastro.2020.10.057

Atherton, F.R., Hall, M.J., Hassall, C.H., Lambert, R.W., Lloyd, W.J., Ringrose, P.S., Westmacott, D., 1982. Antibacterial activity and mechanism of action of phosphonopeptides based on aminomethylphosphonic acid. Antimicrob. Agents Chemother. 22, 571–578. https://doi.org/10.1128/AAC.22.4.571

Azam, M.A., Jayaram, U., 2016. Inhibitors of alanine racemase enzyme: a review. J. Enzyme Inhib. Med. Chem. 31, 517–526. https://doi.org/10.3109/14756366.2015.1050010

Bai, X., Ya, R., Tang, X., Cai, M., 2023. Role and interaction of bacterial sphingolipids in human health. Front. Microbiol. 14, 1289819. https://doi.org/10.3389/fmicb.2023.1289819

Băncilă, R.I., Hartel, T., Plăiaşu, R., Smets, J., Cogălniceanu, D., 2010. Comparing three body condition indices in amphibians: a case study of yellow-bellied toad Bombina variegata. Amphib.-Reptil. 31, 558–562. https://doi.org/10.1163/017353710X518405

Barreto, L.S., Souza, T.L. de, Morais, T.P. de, Oliveira Ribeiro, C.A. de, 2023. Toxicity of glyphosate and aminomethylphosphonic acid (AMPA) to the early stages of development of Steindachneridion melanodermatum, an endangered endemic species of Southern Brazil. Environ. Toxicol. Pharmacol. 102, 104234. https://doi.org/10.1016/j.etap.2023.104234

Beattie, R.C., 1980. A physico-chemical investigation of the jelly capsules surrounding eggs of the Common frog (Rana temporaria temporaria). J. Zool. 190, 1–25. https://doi.org/10.1111/j.1469-7998.1980.tb01419.x

Blot, N., Veillat, L., Rouzé, R., Delatte, H., 2019. Glyphosate, but not its metabolite AMPA, alters the honeybee gut microbiota. PLOS ONE 14, e0215466. https://doi.org/10.1371/journal.pone.0215466

Bolnick, D.I., Snowberg, L.K., Hirsch, P.E., Lauber, C.L., Knight, R., Caporaso, J.G., Svanbäck, R., 2014. Individuals' diet diversity influences gut microbial diversity in two freshwater fish (threespine stickleback and Eurasian perch). Ecol. Lett. 17, 979–987. https://doi.org/10.1111/ele.12301

Bowers, E.K., Hodges, C.J., Forsman, A.M., Vogel, L.A., Masters, B.S., Johnson, B.G.P., Johnson, L.S., Thompson, C.F., Sakaluk, S.K., 2014. Neonatal body condition, immune responsiveness, and hematocrit predict longevity in a wild bird population. Ecology 95, 3027–3034. https://doi.org/10.1890/14-0418.1

Bright Ross, J.G., Newman, C., Buesching, C.D., Connolly, E., Nakagawa, S., Macdonald, D.W., 2021. A fat chance of survival: Body condition provides life-history dependent buffering of environmental change in a wild mammal population. Clim. Change Ecol. 2, 100022. https://doi.org/10.1016/j.ecochg.2021.100022

Bunker, M.E., Elliott, G., Heyer-Gray, H., Martin, M.O., Arnold, A.E., Weiss, S.L., 2021. Vertically
transmitted microbiome protects eggs from fungal infection and egg failure. Anim.
Microbiome 3, 43. https://doi.org/10.1186/s42523-021-00104-5
Burggren, W., 1985. Gas Exchange, Metabolism, and "Ventilation" in Gelatinous Frog Egg Masses.
Physiol. Zool. 58, 503–514. https://doi.org/10.1086/physzool.58.5.30158578
Cao, Y., Dong, Q., Wang, D., Zhang, P., Liu, Y., Niu, C., 2022. microbiomeMarker: an R/Bioconductor
package for microbiome marker identification and visualization. Bioinforma. Oxf. Engl. 38,
4027–4029. https://doi.org/10.1093/bioinformatics/btac438
Carles, L., Gardon, H., Joseph, L., Sanchís, J., Farré, M., Artigas, J., 2019. Meta-analysis of glyphosate
contamination in surface waters and dissipation by biofilms. Environ. Int. 124, 284–293.
https://doi.org/10.1016/j.envint.2018.12.064
Chen, Z., Chen, J.-Q., Liu, Y., Zhang, J., Chen, X.-H., Qu, Y.-F., 2022. Comparative study on gut
microbiota in three Anura frogs from a mountain stream. Ecol. Evol. 12, e8854.
https://doi.org/10.1002/ece3.8854
Cheron, M., Brischoux, F., 2023. Exposure to Low Concentrations of AMPA Influences Morphology
and Decreases Survival During Larval Development in a Widespread Amphibian Species. Arch.
Environ. Contam. Toxicol. 85, 92–103. https://doi.org/10.1007/s00244-023-01008-y
Cheron, M., Brischoux, F., 2020. Aminomethylphosphonic acid alters amphibian embryonic
development at environmental concentrations. Environ. Res. 190, 109944.
Cheron, M., Costantini, D., Angelier, F., Ribout, C., Brischoux, F., 2022. Aminomethylphosphonic acid
(AMPA) alters oxidative status during embryonic development in an amphibian species.
Chemosphere 287, 131882. https://doi.org/10.1016/j.chemosphere.2021.131882
Cheron, M., Raoelison, L., Kato, A., Ropert-Coudert, Y., Meyer, X., MacIntosh, A.J.J., Brischoux, F.,
2021. Ontogenetic changes in activity, locomotion and behavioural complexity in tadpoles.
Biol. J. Linn. Soc. 134, 165–176. https://doi.org/10.1093/biolinnean/blab077
Claus, S.P., Guillou, H., Ellero-Simatos, S., 2016. The gut microbiota: a major player in the toxicity of
environmental pollutants? Npj Biofilms Microbiomes 2, 16003.
https://doi.org/10.1038/npjbiofilms.2016.3
Clemente, J.C., Ursell, L.K., Parfrey, L.W., Knight, R., 2012. The Impact of the Gut Microbiota on
Human Health: An Integrative View. Cell 148, 1258–1270.
https://doi.org/10.1016/j.cell.2012.01.035
Comizzoli, P., Power, M.L., Bornbusch, S.L., Muletz-Wolz, C.R., 2021. Interactions between
reproductive biology and microbiomes in wild animal species. Anim. Microbiome 3, 87.
https://doi.org/10.1186/s42523-021-00156-7
Coupe, R.H., Kalkhoff, S.J., Capel, P.D., Gregoire, C., 2012. Fate and transport of glyphosate and
aminomethylphosphonic acid in surface waters of agricultural basins. Pest Manag. Sci. 68,
16–30. https://doi.org/10.1002/ps.2212
Cuzziol Boccioni, A.P., García-Effron, G., Peltzer, P.M., Lajmanovich, R.C., 2023. Effect of glyphosate
and ciprofloxacin exposure on enteric bacteria of tadpoles. Rev. Argent. Microbiol.
https://doi.org/10.1016/j.ram.2022.08.004
Davidson, G.L., Somers, S.E., Wiley, N., Johnson, C.N., Reichert, M.S., Ross, R.P., Stanton, C., Quinn,
J.L., 2021. A time-lagged association between the gut microbiome, nestling weight and
nestling survival in wild great tits. J. Anim. Ecol. 90, 989–1003. https://doi.org/10.1111/1365-
2656.13428
Dick, R.E., Quinn, J.P., 1995. Glyphosate-degrading isolates from environmental samples: occurrence
and pathways of degradation. Appl. Microbiol. Biotechnol. 43, 545–550.
https://doi.org/10.1007/BF00218464
Dietz, M.W., Salles, J.F., Hsu, B.-Y., Dijkstra, C., Groothuis, T.G.G., van der Velde, M., Verkuil, Y.I.,
Tieleman, B.I., 2020. Prenatal Transfer of Gut Bacteria in Rock Pigeon. Microorganisms 8, 61.
https://doi.org/10.3390/microorganisms8010061

Ding, W., Shangguan, Y., Zhu, Y., Sultan, Y., Feng, Y., Zhang, B., Liu, Y., Ma, J., Li, X., 2021. Negative
impacts of microcystin-LR and glyphosate on zebrafish intestine: Linked with gut microbiota
and microRNAs? Environ. Pollut. 286, 117685. https://doi.org/10.1016/j.envpol.2021.117685
Dray, A.B., Dufour, S., 2007. The ade4 package: implementing the duality diagram for ecologists. J.
Stat. Softw. 22, 1–20.
Duke, S.O., 2020. Glyphosate: environmental fate and impact. Weed Sci. 68, 201–207.
https://doi.org/10.1017/wsc.2019.28
Evariste, L., Barret, M., Mottier, A., Mouchet, F., Gauthier, L., Pinelli, E., 2019. Gut microbiota of
aquatic organisms: A key endpoint for ecotoxicological studies. Environ. Pollut. 248, 989–999.
https://doi.org/10.1016/j.envpol.2019.02.101
Fan, Y., Pedersen, O., 2021. Gut microbiota in human metabolic health and disease. Nat. Rev.
Microbiol. 19, 55–71. https://doi.org/10.1038/s41579-020-0433-9
Ferretti, P., Pasolli, E., Tett, A., Asnicar, F., Gorfer, V., Fedi, S., Armanini, F., Truong, D.T., Manara, S.,
Zolfo, M., Beghini, F., Bertorelli, R., De Sanctis, V., Bariletti, I., Canto, R., Clementi, R.,
Cologna, M., Crifò, T., Cusumano, G., Gottardi, S., Innamorati, C., Masè, C., Postai, D., Savoi,
D., Duranti, S., Lugli, G.A., Mancabelli, L., Turroni, F., Ferrario, C., Milani, C., Mangifesta, M.,
Anzalone, R., Viappiani, A., Yassour, M., Vlamakis, H., Xavier, R., Collado, C.M., Koren, O.,
Tateo, S., Soffiati, M., Pedrotti, A., Ventura, M., Huttenhower, C., Bork, P., Segata, N., 2018.
Mother-to-Infant Microbial Transmission from Different Body Sites Shapes the Developing
Infant Gut Microbiome. Cell Host Microbe 24, 133-145.e5.
https://doi.org/10.1016/j.chom.2018.06.005
Firrman, J., Liu, L., Mahalak, K., Tanes, C., Bittinger, K., Tu, V., Bobokalonov, J., Mattei, L., Zhang, H.,
Van den Abbeele, P., 2022. The impact of environmental pH on the gut microbiota
community structure and short chain fatty acid production. FEMS Microbiol. Ecol. 98,
fiac038. https://doi.org/10.1093/femsec/fiac038
Flint, H.J., Scott, K.P., Duncan, S.H., Louis, P., Forano, E., 2012. Microbial degradation of complex
carbohydrates in the gut. Gut Microbes 3, 289–306. https://doi.org/10.4161/gmic.19897
Fox, E.M., Mendz, G.L., 2006. Phosphonate degradation in microorganisms. Enzyme Microb.
Technol., Papers from the 1st International Conference on Environmental, Industrial and
Applied Microbiology (BioMicroWorld-2005) 40, 145–150.
https://doi.org/10.1016/j.enzmictec.2005.10.047
Fréville, M., Estienne, A., Ramé, C., Lefort, G., Chahnamian, M., Staub, C., Venturi, E., Lemarchand, J.,
Maximin, E., Hondelatte, A., Zemb, O., Canlet, C., Guabiraba, R., Froment, P., Dupont, J.,
2022. Chronic dietary exposure to a glyphosate-based herbicide results in total or partial
reversibility of plasma oxidative stress, cecal microbiota abundance and short-chain fatty
acid composition in broiler hens. Front. Physiol. 13.
Friberg, I.M., Taylor, J.D., Jackson, J.A., 2019. Diet in the Driving Seat: Natural Diet-Immunity-
Microbiome Interactions in Wild Fish. Front. Immunol. 10.
https://doi.org/10.3389/fimmu.2019.00243
Fujimori, S., 2021. Humans have intestinal bacteria that degrade the plant cell walls in herbivores.
World J. Gastroenterol. 27, 7784–7791. https://doi.org/10.3748/wjg.v27.i45.7784
Gantois, I., Ducatelle, R., Pasmans, F., Haesebrouck, F., Gast, R., Humphrey, T.J., Van Immerseel, F.,
2009. Mechanisms of egg contamination by Salmonella Enteritidis. FEMS Microbiol. Rev. 33,
718–738. https://doi.org/10.1111/j.1574-6976.2008.00161.x
Giraud, C., Callac, N., Boulo, V., Lam, J.-S., Pham, D., Selmaoui-Folcher, N., Wabete, N., 2022. The
Active Microbiota of the Eggs and the Nauplii of the Pacific Blue Shrimp Litopenaeus
stylirostris Partially Shaped by a Potential Vertical Transmission. Front. Microbiol. 13,
886752. https://doi.org/10.3389/fmicb.2022.886752
Gomaa, E.Z., 2020. Human gut microbiota/microbiome in health and diseases: a review. Antonie Van
Leeuwenhoek 113, 2019–2040. https://doi.org/10.1007/s10482-020-01474-7

Gonçalves, A.M.M., Rocha, C.P., Marques, J.C., Gonçalves, F.J.M., 2021. Enzymes as useful biomarkers to assess the response of freshwater communities to pesticide exposure – A review. Ecol. Indic. 122, 107303. https://doi.org/10.1016/j.ecolind.2020.107303

Gosner, K.L., 1960. A Simplified Table for Staging Anuran Embryos and Larvae with Notes on Identification. Herpetologica 16, 183–190.

Grandcoin, A., Piel, S., Baurès, E., 2017. AminoMethylPhosphonic acid (AMPA) in natural waters: Its sources, behavior and environmental fate. Water Res. 117, 187–197. https://doi.org/10.1016/j.watres.2017.03.055

Grond, K., Sandercock, B.K., Jumpponen, A., Zeglin, L.H., 2018. The avian gut microbiota: community, physiology and function in wild birds. J. Avian Biol. 49, e01788. https://doi.org/10.1111/jav.01788

Guilherme, S., Santos, M.A., Gaivão, I., Pacheco, M., 2014. DNA and chromosomal damage induced in fish (Anguilla anguilla L.) by aminomethylphosphonic acid (AMPA)—the major environmental breakdown product of glyphosate. Environ. Sci. Pollut. Res. 21, 8730–8739. https://doi.org/10.1007/s11356-014-2803-1

Harfoot, M.B.J., Johnston, A., Balmford, A., Burgess, N.D., Butchart, S.H.M., Dias, M.P., Hazin, C., Hilton-Taylor, C., Hoffmann, M., Isaac, N.J.B., Iversen, L.L., Outhwaite, C.L., Visconti, P., Geldmann, J., 2021. Using the IUCN Red List to map threats to terrestrial vertebrates at global scale. Nat. Ecol. Evol. 5, 1510–1519. https://doi.org/10.1038/s41559-021-01542-9

Harkness, D.R., 1966. Bacterial Growth on Aminoalkylphosphonic Acids. J. Bacteriol. 92, 623–627. https://doi.org/10.1128/jb.92.3.623-627.1966

Hernández-Gómez, O., Hua, J., 2023. From the organismal to biosphere levels: environmental impacts on the amphibian microbiota. FEMS Microbiol. Rev. 47, fuad002. https://doi.org/10.1093/femsre/fuad002

Hoffmann, M., Hilton-Taylor, C., Angulo, A., Böhm, M., Brooks, T.M., Butchart, S.H.M., Carpenter, K.E., Chanson, J., Collen, B., Cox, N.A., Darwall, W.R.T., Dulvy, N.K., Harrison, L.R., Katariya, V., Pollock, C.M., Quader, S., Richman, N.I., Rodrigues, A.S.L., Tognelli, M.F., Vié, J.-C., Aguiar, J.M., Allen, D.J., Allen, G.R., Amori, G., Ananjeva, N.B., Andreone, F., Andrew, P., Ortiz, A.L.A., Baillie, J.E.M., Baldi, R., Bell, B.D., Biju, S.D., Bird, J.P., Black-Decima, P., Blanc, J.J., Bolaños, F., Bolivar-G., W., Burfield, I.J., Burton, J.A., Capper, D.R., Castro, F., Catullo, G., Cavanagh, R.D., Channing, A., Chao, N.L., Chenery, A.M., Chiozza, F., Clausnitzer, V., Collar, N.J., Collett, L.C., Collette, B.B., Fernandez, C.F.C., Craig, M.T., Crosby, M.J., Cumberlidge, N., Cuttelod, A., Derocher, A.E., Diesmos, A.C., Donaldson, J.S., Duckworth, J.W., Dutson, G., Dutta, S.K., Emslie, R.H., Farjon, A., Fowler, S., Freyhof, J., Garshelis, D.L., Gerlach, J., Gower, D.J., Grant, T.D., Hammerson, G.A., Harris, R.B., Heaney, L.R., Hedges, S.B., Hero, J.-M., Hughes, B., Hussain, S.A., Icochea M., J., Inger, R.F., Ishii, N., Iskandar, D.T., Jenkins, R.K.B., Kaneko, Y., Kottelat, M., Kovacs, K.M., Kuzmin, S.L., La Marca, E., Lamoreux, J.F., Lau, M.W.N., Lavilla, E.O., Leus, K., Lewison, R.L., Lichtenstein, G., Livingstone, S.R., Lukoschek, V., Mallon, D.P., McGowan, P.J.K., McIvor, A., Moehlman, P.D., Molur, S., Alonso, A.M., Musick, J.A., Nowell, K., Nussbaum, R.A., Olech, W., Orlov, N.L., Papenfuss, T.J., Parra-Olea, G., Perrin, W.F., Polidoro, B.A., Pourkazemi, M., Racey, P.A., Ragle, J.S., Ram, M., Rathbun, G., Reynolds, R.P., Rhodin, A.G.J., Richards, S.J., Rodríguez, L.O., Ron, S.R., Rondinini, C., Rylands, A.B., Sadovy de Mitcheson, Y., Sanciangco, J.C., Sanders, K.L., Santos-Barrera, G., Schipper, J., Self-Sullivan, C., Shi, Y., Shoemaker, A., Short, F.T., Sillero-Zubiri, C., Silvano, D.L., Smith, K.G., Smith, A.T., Snoeks, J., Stattersfield, A.J., Symes, A.J., Taber, A.B., Talukdar, B.K., Temple, H.J., Timmins, R., Tobias, J.A., Tsytsulina, K., Tweddle, D., Ubeda, C., Valenti, S.V., Paul van Dijk, P., Veiga, L.M., Veloso, A., Wege, D.C., Wilkinson, M., Williamson, E.A., Xie, F., Young, B.E., Akçakaya, H.R., Bennun, L., Blackburn, T.M., Boitani, L., Dublin, H.T., da Fonseca, G.A.B., Gascon, C., Lacher, T.E., Mace, G.M., Mainka, S.A., McNeely, J.A., Mittermeier, R.A., Reid, G.M., Rodriguez, J.P., Rosenberg, A.A., Samways, M.J., Smart, J., Stein, B.A., Stuart, S.N., 2010. The Impact of Conservation on the Status of the World's Vertebrates. Science 330, 1503–1509. https://doi.org/10.1126/science.1194442

Hornung, B.V.H., Zwittink, R.D., Kuijper, E.J., 2019. Issues and current standards of controls in
microbiome research. FEMS Microbiol. Ecol. 95, fiz045.
https://doi.org/10.1093/femsec/fiz045

Hu, Z., Tong, Q., Chang, J., Yu, J., Li, S., Niu, H., Ma, D., 2021. Gut bacterial communities in the
freshwater snail Planorbella trivolvis and their modification by a non-herbivorous diet. PeerJ
9, e10716. https://doi.org/10.7717/peerj.10716

Hughey, M.C., Delia, J., Belden, L.K., 2017. Diversity and stability of egg-bacterial assemblages: The
role of paternal care in the glassfrog Hyalinobatrachium colymbiphyllum. Biotropica 49, 792–
802. https://doi.org/10.1111/btp.12461

Iori, S., Rovere, G.D., Ezzat, L., Smits, M., Ferraresso, S.S., Babbucci, M., Marin, M.G., Masiero, L.,
Fabrello, J., Garro, E., Carraro, L., Cardazzo, B., Patarnello, T., Matozzo, V., Bargelloni, L.,
Milan, M., 2020. The effects of glyphosate and AMPA on the mediterranean mussel Mytilus
galloprovincialis and its microbiota. Environ. Res. 182, 108984.
https://doi.org/10.1016/j.envres.2019.108984

Ivantsova, E., Wengrovitz, A.S., Souders, C.L., Martyniuk, C.J., 2022. Developmental and behavioral
toxicity assessment of glyphosate and its main metabolite aminomethylphosphonic acid
(AMPA) in zebrafish embryos/larvae. Environ. Toxicol. Pharmacol. 93, 103873.
https://doi.org/10.1016/j.etap.2022.103873

Jiménez, R.R., Sommer, S., 2017. The amphibian microbiome: natural range of variation, pathogenic
dysbiosis, and role in conservation. Biodivers. Conserv. 26, 763–786.
https://doi.org/10.1007/s10531-016-1272-x

Khalifa, A.Y.Z., AlMalki, M.A., Bekhet, G.M., 2021. Pathological and mortality findings associated with
Aeromonas hydrophila from frog eggs in Al-Ahsa region of Saudi Arabia. Aquac. Res. 52,
1227–1236. https://doi.org/10.1111/are.14981

Kolpin, D.W., Thurman, E.M., Lee, E.A., Meyer, M.T., Furlong, E.T., Glassmeyer, S.T., 2006. Urban
contributions of glyphosate and its degradate AMPA to streams in the United States. Sci.
Total Environ. 354, 191–197. https://doi.org/10.1016/j.scitotenv.2005.01.028

Landler, L., Burgstaller, S., Spießberger, M., Horvath, A., Zhelev, Z., Mollov, I., Sinsch, U., Nepita, J.,
Schwabel, F., Kuhn, W., Köbele, C., Sedlmeier, H., Amon, C., Mazgajska, J., Mazgajski, T.D.,
Sistani, A., Schluckebier, R., Andrä, E., Ott, M., Gollmann, G., 2023. A Unified Approach to
Analysis of Body Condition in Green Toads. Diversity 15, 43.
https://doi.org/10.3390/d15010043

Langlois, V.S., 2021. Amphibian Toxicology: A Rich But Underappreciated Model for Ecotoxicology
Research. Arch. Environ. Contam. Toxicol. 80, 661–662. https://doi.org/10.1007/s00244-021-
00844-0

Lehman, P.C., Cady, N., Ghimire, S., Shahi, S.K., Shrode, R.L., Lehmler, H.-J., Mangalam, A.K., 2023.
Low-dose glyphosate exposure alters gut microbiota composition and modulates gut
homeostasis. Environ. Toxicol. Pharmacol. 100, 104149.
https://doi.org/10.1016/j.etap.2023.104149

Levine, S.L., von Mérey, G., Minderhout, T., Manson, P., Sutton, P., 2015. Aminomethylphosphonic
acid has low chronic toxicity to Daphnia magna and Pimephales promelas. Environ. Toxicol.
Chem. 34, 1382–1389. https://doi.org/10.1002/etc.2940

Liu, G., Cain, K., Schwanz, L., 2020. Maternal Temperature, Corticosterone, and Body Condition as
Mediators of Maternal Effects in Jacky Dragons (Amphibolurus muricatus). Physiol. Biochem.
Zool. 93, 434–449. https://doi.org/10.1086/711955

Lozano, V.L., Defarge, N., Rocque, L.-M., Mesnage, R., Hennequin, D., Cassier, R., de Vendômois, J.S.,
Panoff, J.-M., Séralini, G.-E., Amiel, C., 2018. Sex-dependent impact of Roundup on the rat
gut microbiome. Toxicol. Rep. 5, 96–107. https://doi.org/10.1016/j.toxrep.2017.12.005

Lozupone, C., Hamady, M., Knight, R., 2006. UniFrac – An online tool for comparing microbial
community diversity in a phylogenetic context. BMC Bioinformatics 7, 371.
https://doi.org/10.1186/1471-2105-7-371

MacCracken, J.G., Stebbings, J.L., 2012. Test of a Body Condition Index with Amphibians. J. Herpetol.
46, 346–350.

Maggi, F., la Cecilia, D., Tang, F.H.M., McBratney, A., 2020. The global environmental hazard of
glyphosate use. Sci. Total Environ. 717, 137167.
https://doi.org/10.1016/j.scitotenv.2020.137167

Martens, E.C., Lowe, E.C., Chiang, H., Pudlo, N.A., Wu, M., McNulty, N.P., Abbott, D.W., Henrissat, B.,
Gilbert, H.J., Bolam, D.N., Gordon, J.I., 2011. Recognition and Degradation of Plant Cell Wall
Polysaccharides by Two Human Gut Symbionts. PLOS Biol. 9, e1001221.
https://doi.org/10.1371/journal.pbio.1001221

Martins-Gomes, C., Silva, T.L., Andreani, T., Silva, A.M., 2022. Glyphosate vs. Glyphosate-Based
Herbicides Exposure: A Review on Their Toxicity. J. Xenobiotics 12, 21–40.
https://doi.org/10.3390/jox12010003

Matozzo, V., Marin, M.G., Masiero, L., Tremonti, M., Biamonte, S., Viale, S., Finos, L., Lovato, G.,
Pastore, P., Bogialli, S., 2018. Effects of aminomethylphosphonic acid, the main breakdown
product of glyphosate, on cellular and biochemical parameters of the mussel Mytilus
galloprovincialis. Fish Shellfish Immunol. 83, 321–329.
https://doi.org/10.1016/j.fsi.2018.09.036

McMurdie, P.J., Holmes, S., 2012. Phyloseq: a bioconductor package for handling and analysis of
high-throughput phylogenetic sequence data. Pac. Symp. Biocomput. Pac. Symp. Biocomput.
235–246.

Mesnage, R., Panzacchi, S., Bourne, E., Mein, C.A., Perry, M.J., Hu, J., Chen, J., Mandrioli, D., Belpoggi,
F., Antoniou, M.N., 2021. Glyphosate and its formulations Roundup Bioflow and RangerPro
alter bacterial and fungal community composition in the rat caecum microbiome.
https://doi.org/10.1101/2021.11.19.468976

Milner, J.M., Stien, A., Irvine, R.J., Albon, S.D., Langvatn, R., Ropstad, E., 2003. Body condition in
Svalbard reindeer and the use of blood parameters as indicators of condition and fitness.
Can. J. Zool. 81, 1566–1578. https://doi.org/10.1139/z03-152

Moreno-Navarrete, J.M., Fernandez-Real, J.M., 2019. The gut microbiota modulates both browning
of white adipose tissue and the activity of brown adipose tissue. Rev. Endocr. Metab. Disord.
20, 387–397. https://doi.org/10.1007/s11154-019-09523-x

Motta, E.V.S., Raymann, K., Moran, N.A., 2018. Glyphosate perturbs the gut microbiota of honey
bees. Proc. Natl. Acad. Sci. 115, 10305–10310. https://doi.org/10.1073/pnas.1803880115

Murphy, K.M., Le, S.M., Wilson, A.E., Warner, D.A., 2023. The Microbiome as a Maternal Effect: A
Systematic Review on Vertical Transmission of Microbiota. Integr. Comp. Biol. 63, 597–609.
https://doi.org/10.1093/icb/icad031

Naïades, 2023. Naïades [WWW Document]. URL https://naiades.eaufrance.fr/acces-
donnees#/physicochimie/resultats?debut=30-11-2013&fin=30-11-2023¶metres=1907
(accessed 11.30.23).

Obojska, A., Lejczak, B., 2003. Utilisation of structurally diverse organophosphonates by
Streptomycetes. Appl. Microbiol. Biotechnol. 62, 557–563. https://doi.org/10.1007/s00253-
003-1281-z

Obojska, A., Lejczak, B., Kubrak, M., 1999. Degradation of phosphonates by streptomycete isolates.
Appl. Microbiol. Biotechnol. 51, 872–876. https://doi.org/10.1007/s002530051476

Ojelade, B.S., Durowoju, O.S., Adesoye, P.O., Gibb, S.W., Ekosse, G.-I., 2022. Review of Glyphosate-
Based Herbicide and Aminomethylphosphonic Acid (AMPA): Environmental and Health
Impacts. Appl. Sci. 12, 8789. https://doi.org/10.3390/app12178789

Olsen, I., Jantzen, E., 2001. Sphingolipids in Bacteria and Fungi. Anaerobe 7, 103–112.
https://doi.org/10.1006/anae.2001.0376

Olson, J.H., Chandler, D.E., 1999. Xenopus laevis Egg Jelly Contains Small Proteins That Are Essential
to Fertilization. Dev. Biol. 210, 401–410. https://doi.org/10.1006/dbio.1999.9281

Owagboriaye, F., Mesnage, R., Dedeke, G., Adegboyega, T., Aladesida, A., Adeleke, M., Owa, S.,
Antoniou, M.N., 2021. Impacts of a glyphosate-based herbicide on the gut microbiome of

836   three earthworm species (Alma millsoni, Eudrilus eugeniae and Libyodrilus violaceus): A pilot
837   study. Toxicol. Rep. 8, 753–758. https://doi.org/10.1016/j.toxrep.2021.03.021
838 Pan, X., Raaijmakers, J.M., Carrión, V.J., 2023. Importance of Bacteroidetes in host–microbe
839   interactions and ecosystem functioning. Trends Microbiol.
840 Peig, J., Green, A.J., 2009. New perspectives for estimating body condition from mass/length data:
841   the scaled mass index as an alternative method. Oikos 118, 1883–1891.
842   https://doi.org/10.1111/j.1600-0706.2009.17643.x
843 Poiger, T., Buerge, I.J., Bächli, A., Müller, M.D., Balmer, M.E., 2017. Occurrence of the herbicide
844   glyphosate and its metabolite AMPA in surface waters in Switzerland determined with on-
845   line solid phase extraction LC-MS/MS. Environ. Sci. Pollut. Res. 24, 1588–1596.
846   https://doi.org/10.1007/s11356-016-7835-2
847 Prest, T.L., Kimball, A.K., Kueneman, J.G., McKenzie, V.J., 2018. Host-associated bacterial community
848   succession during amphibian development. Mol. Ecol. 27, 1992–2006.
849   https://doi.org/10.1111/mec.14507
850 Puigbò, P., Leino, L.I., Rainio, M.J., Saikkonen, K., Saloniemi, I., Helander, M., 2022. Does Glyphosate
851   Affect the Human Microbiota? Life 12, 707. https://doi.org/10.3390/life12050707
852 R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Foundation for
853   Statistical Computing.
854 Renoirt, M., Angelier, F., Cheron, M., Brischoux, F., 2022. What are the contributions of maternal and
855   paternal traits to fecundity and offspring development? A case study in an amphibian
856   species, the spined toad Bufo spinosus. Curr. Zool. zoac072.
857   https://doi.org/10.1093/cz/zoac072
858 Rollins-Smith, L.A., 2020. Global Amphibian Declines, Disease, and the Ongoing Battle between
859   Batrachochytrium Fungi and the Immune System. Herpetologica 76, 178–188.
860   https://doi.org/10.1655/0018-0831-76.2.178
861 Rosas-Pérez, T., Rosenblueth, M., Rincón-Rosales, R., Mora, J., Martínez-Romero, E., 2014. Genome
862   Sequence of "Candidatus Walczuchella monophlebidarum" the Flavobacterial Endosymbiont
863   of Llaveia axin axin (Hemiptera: Coccoidea: Monophlebidae). Genome Biol. Evol. 6, 714–726.
864   https://doi.org/10.1093/gbe/evu049
865 Roth, O., Beemelmanns, A., Barribeau, S.M., Sadd, B.M., 2018. Recent advances in vertebrate and
866   invertebrate transgenerational immunity in the light of ecology and evolution. Heredity 121,
867   225–238. https://doi.org/10.1038/s41437-018-0101-2
868 Ruuskanen, S., Rainio, M.J., Gómez-Gallego, C., Selenius, O., Salminen, S., Collado, M.C., Saikkonen,
869   K., Saloniemi, I., Helander, M., 2020. Glyphosate-based herbicides influence antioxidants,
870   reproductive hormones and gut microbiome but not reproduction: A long-term experiment
871   in an avian model. Environ. Pollut. 266, 115108.
872   https://doi.org/10.1016/j.envpol.2020.115108
873 Scalvenzi, T., Clavereau, I., Pollet, N., 2020. Insights on the evolution of vertebrate microbiomes from
874   the analysis of the Xenopus frog microbiota across life stages. bioRxiv 2020.05.25.110734.
875   https://doi.org/10.1101/2020.05.25.110734
876 Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., Huttenhower, C., 2011.
877   Metagenomic biomarker discovery and explanation. Genome Biol. 12, R60.
878   https://doi.org/10.1186/gb-2011-12-6-r60
879 Sharpton, T.J., 2018. Role of the Gut Microbiome in Vertebrate Evolution. mSystems 3, e00174-17.
880   https://doi.org/10.1128/mSystems.00174-17
881 Shen, Y., Jiang, Z., Zhong, X., Wang, H., Liu, Y., Li, X., 2022. Manipulation of cadmium and diethylhexyl
882   phthalate on Rana chensinensis tadpoles affects the intestinal microbiota and fatty acid
883   metabolism. Sci. Total Environ. 821, 153455.
884   https://doi.org/10.1016/j.scitotenv.2022.153455
885 Shu, Y., Hong, P., Yu, Q., Wang, G., Zhang, J., Donde, O.O., Xiao, B., Wu, H., 2019. High-Throughput
886   Sequencing Analysis Reveals Correlations between Host Phylogeny, Gut Microbiota, and

887   Habitat of Wild Frogs from a Mountainous Area. Copeia 107, 131–137.
888   https://doi.org/10.1643/OT-18-040

889 Song, X., Zhang, J., Song, J., Zhai, Y., 2021. Decisive Effects of Life Stage on the Gut Microbiota
890   Discrepancy Between Two Wild Populations of Hibernating Asiatic Toads (Bufo gargarizans).
891   Front. Microbiol. 12.

892 Stämmler, F., Gläsner, J., Hiergeist, A., Holler, E., Weber, D., Oefner, P.J., Gessner, A., Spang, R., 2016.
893   Adjusting microbiome profiles for differences in microbial load by spike-in bacteria.
894   Microbiome 4, 28. https://doi.org/10.1186/s40168-016-0175-0

895 Stebbins, R.C., Cohen, N.W., 1995. A Natural History of Amphibians, Princeton University Press. ed.

896 Struger, J., Van Stempvoort, D.R., Brown, S.J., 2015. Sources of aminomethylphosphonic acid (AMPA)
897   in urban and rural catchments in Ontario, Canada: Glyphosate or phosphonates in
898   wastewater? Environ. Pollut. 204, 289–297. https://doi.org/10.1016/j.envpol.2015.03.038

899 Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S.L., Fischman, D.L., Waller, R.W.,
900   2004. Status and Trends of Amphibian Declines and Extinctions Worldwide. Science 306,
901   1783–1786. https://doi.org/10.1126/science.1103538

902 Studnik, H., Liebsch, S., Forlani, G., Wieczorek, D., Kafarski, P., Lipok, J., 2015. Amino
903   polyphosphonates – chemical features and practical uses, environmental durability and
904   biodegradation. New Biotechnol. 32, 1–6. https://doi.org/10.1016/j.nbt.2014.06.007

905 Sugden, S., Sanderson, D., Ford, K., Stein, L.Y., St. Clair, C.C., 2020. An altered microbiome in urban
906   coyotes mediates relationships between anthropogenic diet and poor health. Sci. Rep. 10,
907   22207. https://doi.org/10.1038/s41598-020-78891-1

908 Tartu, S., Lorrain-Soligon, L., Cheron, M., Dupoué, A., Brischoux, F., 2023. Colouration matters in dull
909   toads: ultraviolet adornment for ladies and agrochemicals fading effects. Oecologia 203,
910   491–502. https://doi.org/10.1007/s00442-023-05482-y

911 Tartu, S., Renoirt, M., Cheron, M., Gisselmann, L.-L., Catoire, S., Brischoux, F., 2022. Did decades of
912   glyphosate use have selected for resistant amphibians in agricultural habitats? Environ.
913   Pollut. 310, 119823. https://doi.org/10.1016/j.envpol.2022.119823

914 Trevelline, B.K., MacLeod, K.J., Knutie, S.A., Langkilde, T., Kohl, K.D., 2018. In ovo microbial
915   communities: a potential mechanism for the initial acquisition of gut microbiota among
916   oviparous birds and lizards. Biol. Lett. 14, 20180225. https://doi.org/10.1098/rsbl.2018.0225

917 Turnbaugh, P.J., Ley, R.E., Mahowald, M.A., Magrini, V., Mardis, E.R., Gordon, J.I., 2006. An obesity-
918   associated gut microbiome with increased capacity for energy harvest. Nature 444, 1027–
919   1031. https://doi.org/10.1038/nature05414

920 Vecchioni, L., Pace, A., Sucato, A., Berlinghieri, F., Cambera, I., Visconti, G., Hochscheid, S., Arculeo,
921   M., Alduina, R., 2022. Unveiling the egg microbiota of the loggerhead sea turtle Caretta
922   caretta in nesting beaches of the Mediterranean Sea. PLOS ONE 17, e0268345.
923   https://doi.org/10.1371/journal.pone.0268345

924 Wake, D.B., Vredenburg, V.T., 2008. Are we in the midst of the sixth mass extinction? A view from
925   the world of amphibians. Proc. Natl. Acad. Sci. https://doi.org/10.1073/pnas.0801921105

926 Walke, J.B., Harris, R.N., Reinert, L.K., Rollins-Smith, L.A., Woodhams, D.C., 2011. Social Immunity in
927   Amphibians: Evidence for Vertical Transmission of Innate Defenses. Biotropica 43, 396–400.
928   https://doi.org/10.1111/j.1744-7429.2011.00787.x

929 Walsh, L., Hill, C., Ross, R.P., 2023. Impact of glyphosate (RoundupTM) on the composition and
930   functionality of the gut microbiome. Gut Microbes 15, 2263935.
931   https://doi.org/10.1080/19490976.2023.2263935

932 Wampach, L., Heintz-Buschart, A., Fritz, J.V., Ramiro-Garcia, J., Habier, J., Herold, M., Narayanasamy,
933   S., Kaysen, A., Hogan, A.H., Bindl, L., Bottu, J., Halder, R., Sjöqvist, C., May, P., Andersson,
934   A.F., de Beaufort, C., Wilmes, P., 2018. Birth mode is associated with earliest strain-conferred
935   gut microbiome functions and immunostimulatory potential. Nat. Commun. 9, 5091.
936   https://doi.org/10.1038/s41467-018-07631-x

Wang, S., Ryan, C.A., Boyaval, P., Dempsey, E.M., Ross, R.P., Stanton, C., 2020. Maternal Vertical
Transmission Affecting Early-life Microbiota Development. Trends Microbiol. 28, 28–45.
https://doi.org/10.1016/j.tim.2019.07.010
Wang, Y., Smith, H.K., Goossens, E., Hertzog, L., Bletz, M.C., Bonte, D., Verheyen, K., Lens, L., Vences,
M., Pasmans, F., Martel, A., 2021. Diet diversity and environment determine the intestinal
microbiome and bacterial pathogen load of fire salamanders. Sci. Rep. 11, 20493.
https://doi.org/10.1038/s41598-021-98995-6
Warne, R.W., Kirschman, L., Zeglin, L., 2019. Manipulation of gut microbiota during critical
developmental windows affects host physiological performance and disease susceptibility
across ontogeny. J. Anim. Ecol. 88, 845–856. https://doi.org/10.1111/1365-2656.12973
Worsley, S.F., Davies, C.S., Mannarelli, M.-E., Hutchings, M.I., Komdeur, J., Burke, T., Dugdale, H.L.,
Richardson, D.S., 2021. Gut microbiome composition, not alpha diversity, is associated with
survival in a natural vertebrate population. Anim. Microbiome 3, 84.
https://doi.org/10.1186/s42523-021-00149-6
Xiong, J.-B., Nie, L., Chen, J., 2019. Current understanding on the roles of gut microbiota in fish
disease and immunity. Zool. Res. 40, 70–76. https://doi.org/10.24272/j.issn.2095-
8137.2018.069
Yan, B., Han, J., Sun, Y., Lei, L., Yuan, J., Qiao, Z., Men, J., Wang, X., Guo, Y., Wang, Q., Zhou, B., 2022.
Probiotics ameliorate growth retardation of glyphosate by regulating intestinal microbiota
and metabolites in crucian carp (Carassius auratus). Sci. Total Environ. 851, 158260.
https://doi.org/10.1016/j.scitotenv.2022.158260
Yan, B., Sun, Y., Fu, K., Zhang, Y., Lei, L., Men, J., Guo, Y., Wu, S., Han, J., Zhou, B., 2023. Effects of
glyphosate exposure on gut–liver axis: Metabolomic and mechanistic analysis in grass carp
(Ctenopharyngodon idellus). Sci. Total Environ. 902, 166062.
https://doi.org/10.1016/j.scitotenv.2023.166062
Yang, J.-A., Yang, S.-H., Kim, J., Kwon, K.K., Oh, H.-M., 2017. Comparative genome analysis of the
Flavobacteriales bacterium strain UJ101, isolated from the gut of Atergatis reticulatus. J.
Microbiol. 55, 583–591. https://doi.org/10.1007/s12275-017-7172-2
Zhang, T., Yang, Y., Liang, Y., Jiao, X., Zhao, C., 2018. Beneficial Effect of Intestinal Fermentation of
Natural Polysaccharides. Nutrients 10, 1055. https://doi.org/10.3390/nu10081055
Zhang, W., Wang, J., Song, J., Feng, Y., Zhang, S., Wang, N., Liu, S., Song, Z., Lian, K., Kang, W., 2021.
Effects of low-concentration glyphosate and aminomethyl phosphonic acid on zebrafish
embryo development. Ecotoxicol. Environ. Saf. 226, 112854.
https://doi.org/10.1016/j.ecoenv.2021.112854
Zhelev, Z.M., Minchev, D.S., 2023. Body condition based on scaled mass index in adult Pelophylax
ridibundus (Amphibia: Ranidae) inhabiting a polluted area in Southern Bulgaria. Biologia
(Bratisl.). https://doi.org/10.1007/s11756-023-01521-4
Zhou, C., Yang, S., Ka, W., Gao, P., Li, Y., Long, R., Wang, J., 2022. Association of Gut Microbiota With
Metabolism in Rainbow Trout Under Acute Heat Stress. Front. Microbiol. 13.
Zhou, J., Tesfamicael, K., Zhou, S.J., Rollins, L.A., Lopez, C.M.R., 2021. Genetic similarity enhances the
strength of the relationship between gut bacteria and host DNA methylation.
https://doi.org/10.1101/2021.07.10.451923
Zwartjes, M.S.Z., Gerdes, V.E.A., Nieuwdorp, M., 2021. The Role of Gut Microbiota and Its Produced
Metabolites in Obesity, Dyslipidemia, Adipocyte Dysfunction, and Its Interventions.
Metabolites 11, 531. https://doi.org/10.3390/metabo11080531
982