Gut microbiota DPP4-like enzymes are increased in type-2 diabetes and contribute to incretin inactivation

Marta Olivares[1], Paula Hernández-Calderón[2], Sonia Cárdenas-Brito[2], Rebeca Liébana-García[1], Yolanda Sanz[1*], Alfonso Benítez-Páez[1,2*]

[1] Microbiome, Nutrition and Health Research Unit. Institute of Agrochemistry and Food Technology, Spanish National Research Council, IATA-CSIC. 46980 Paterna-Valencia, Spain.

[2] Host-Microbe Interactions in Metabolic Health Laboratory. Principe Felipe Research Center (CIPF). 46012 Valencia, Spain.

Email addresses:

MO email: m.olivares@iata.csic.es

PH-C email: phernandez@cipf.es

SC-B email: sonicard@gmail.com

RL-G email: reliegar@iata.csic.es

YS email: yolsanz@iata.csic.es.

AB-P email: abenitez@cipf.es.

* Correspondence must be addressed to these authors.

Abstract

Background: The gut microbiota controls broad aspects of human metabolism and feeding behavior, but the basis for this control remains largely unclear. Given the key role of human dipeptidyl peptidase 4 (DPP4) in host metabolism, we investigated whether microbiota DPP4-like counterparts perform the same function.

Results: We identified novel functional homologs of human DPP4 in several bacterial species inhabiting the human gut, and specific associations between *Parabacteroides* and *Porphyromonas* DPP4-like genes and type 2 diabetes (T2D). We also found that the DPP4-like enzyme from the gut symbiont *Parabacteroides merdae* mimics the proteolytic activity of the human enzyme on peptide YY, neuropeptide Y, gastric inhibitory polypeptide (GIP) and glucagon-like peptide 1 (GLP-1) hormones *in vitro*. Importantly, administration of *E. coli* overexpressing the *P. merdae* DPP4-like enzyme to lipopolysaccharide-treated mice with impaired gut barrier function reduced active GIP and GLP-1 levels, which was attributed to increased DPP4 activity in the portal circulation and the cecal content. Finally, we observed that linagliptin, saxagliptin, sitagliptin and vildagliptin, antidiabetic drugs with DPP4 inhibitory activity, differentially inhibit the activity of the DPP4-like enzyme from *P. merdae*.

Conclusions: Our findings confirm that proteolytic enzymes produced by the gut microbiota are likely to contribute to the glucose metabolic dysfunction that underlies T2D by inactivating incretins, which might inspire the development of improved antidiabetic therapies.

Keywords: DPP4 enzyme, GLP-1, gliptins, incretins, gut microbiome, obesity, glucose

metabolism, type 2 diabetes.

Background

The causal relationship between obesity and comorbidities such as type 2 diabetes (T2D)

and the gut microbiota has long been established; however, our understanding of the

specific mechanisms driving this relationship remains limited [1]. Microbial mimicry of

host activities has been proposed as a mechanism by which pathogens manipulate host

cellular functions [2], and recent evidence indicates that mimicry is also a strategy that can

be used by commensal bacteria in the gut. For example, *Bacteroides vulgatus* produces the

metabolite N-acyl-3-hydroxypalmitoyl-glycine, which can structurally mimic long-chain N-

acyl amides that function as mammalian signaling molecules. Both metabolites activate the

GPR132/G2A receptor, which has been implicated in autoimmune disease and

atherosclerosis [3, 4]. Similarly, the caseinolytic protease ClpB from the gut commensal

Eschericia coli has been shown to translocate and mimic the action of α-melanocyte-

stimulating hormone to control host satiety [5, 6].

Dipeptidyl peptidase 4 (DPP4 or CD26) is a serine protease that rapidly inactivates the

incretin peptides glucagon-like peptide-1 (GLP-1) and gastric inhibitory polypeptide (GIP)

to modulate postprandial insulin secretion and glycemia [7]. Accordingly, administration of

DPP4 inhibitors is a pharmacological strategy to control T2D by prolonging the half-life of

GLP-1 and GIP [8]. Plasma DPP4 activity increases with obesity, which has been

postulated to contribute to reduced incretin activity in the setting of obesity and insulin

resistance [9, 10]. However, the generation of mice with loss of DPP4 activity in multiple

tissues and cell types has shown that not all cellular sources of DPP4 activity affect glucose

metabolism [11, 12]. For example, endothelial DPP4 cleaves inactive GLP-1 and GIP [11],

but adipocyte- and hepatocyte-derived DPP4 are dispensable for glucose control [12]. We

recently demonstrated the ability of the gut microbiota to produce DPP4-like activity in

germ-free and conventionally colonized mice, leading us to propose that gut microbes are

potential mediators of incretin inactivation [13]. Subsequent studies have identified the

primary sources of human DPP4 (hsDPP4) homologs in species of the genera *Bacteroides,*

Parabacteroides and *Porphyromonas* [14-16]. In addition, a peptidase activity

complementary to that of DPP4, termed Xaa-Pro dipeptidyl-peptidase (PepX), has been

identified in lactic acid bacteria commonly found in fermented foods and the human gut,

such as *Lactobacillus* spp., *Lactococcus* spp. and *Streptococcus* spp. [17, 18].

In line with our proposition that gut microbial DPP4 activity might influence host

metabolism [13], some species of *Bacteroides* have recently been reported to inactivate

GLP-1 in mice fed a high-fat diet [16]. Here, we investigated how bacterial DPP4, which is

prevalent in patients with T2D, impairs gut hormone function and evades the catalytic

inhibition of drug therapies for T2D. We identified novel gut bacterial species that harbor

functional homologs of hsDPP4 and associated bacterial DPP4 genes with T2D and

glycemia using metagenomic data from a large cohort of patients [19]. We also demonstrate

the potential of bacterial DPP4 to be secreted extracellularly and to hydrolyze human

incretins (GLP-1 and GIP) and neuropeptides (PYY and NPY) *in vitro*, which was

confirmed in a preclinical model. Finally, we demonstrate the differential interaction

between bacterial DPP4 and four prescribed anti-diabetic drugs with DPP4 inhibitory

activity. Our results shed light on the molecular mechanisms by which specific gut bacterial

species alter the host glucose metabolism and the efficacy of DPP4 inhibitors, providing

new opportunities to improve drug development for the treatment of T2D.

Results

DPP4-like genes are widespread across human gut bacteria

We used a bottom-up approach to search for bacterial DPP4-like genes in the metagenomes

of human subjects with metabolic disorders [20], where the presence of the enzymes may

have clinical implications. We retrieved DPP4-like genes encoded by different Gram-

negative bacteria belonging not only to the genus *Bacteroides*, but also to the genera

Alistipes, formerly *Prevotella*, *Barnesiella* and *Paraprevotella*. We constructed a

phylogenetic tree using amino acid sequence information from more than 1,600 non-

redundant bacterial peptidases, including PepX (belonging to the so-called S15 family) and

DPP4-like and eukaryotic proteins (family S9). After applying advanced phylogenetic

methods, we obtained a maximum likelihood tree explaining the evolutionary relationship

between the two peptidase families (Figure 1A). According to Akaike's Information

Criterion (AIC), the WAG+I+G model of protein evolution better explained the

relationship between the S15 and S9 peptidase families and determined both as independent

lineages. Notably, PepX proteins were found exclusively in Gram-positive bacteria whereas

DPP4-like proteins were found exclusively in Gram-negative bacteria. Taxonomic

assessment revealed that PepX peptidases are predominantly found in the families

Streptococcaceae and Lactobacillaceae, and their respective genera, within the Bacillota

phylum (formerly Firmicutes) (Figure 1A). The families Enterococcaceae,

Leuconostocaceae and Sporolactobacillaceae were also represented to a minor extent.

Contrastingly, DPP4-like proteins were more widely distributed across several families of

the Bacteroidota phylum (formerly Bacteroides), and were evident in several species from

the Flavobacteriaceae, Prevotellaceae, Porphyromonadaceae and Bacteroidaceae families

(Figure 1A). Furthermore, the DPP4-like family proteins showed greater diversity and

genetic distances between members, and proteins from Flavobacteriaceae members

appeared to have evolved independently of other Bacteroidota families. Overall, the amino

acid sequences and the evolutionary relationships between the S15 and S9 peptidase

families support a process of functional convergence [21]. Moreover, we found proteins

corresponding to the three eukaryote proteins (humans, rat and mouse), with identity values

of 25–35% (within the *twilight zone* of sequence similarity) (Figure 1B). The position of the

eukaryotic DPP4 enzymes in the tree seem to not support a phylogenetic relationship with

bacterial DPP4-like enzymes, suggesting that they did not diversify from a common

ancestor. Indeed, the closest relationship between human, rat and mouse DPP4 enzyme

sequences was established with DPP4-like sequences from environmental bacterial species

of the Balneolaceae family (genera *Gracilimonas* and *Aliifondinibius*), which are marine

species with a high frequency of horizontal gene transfer [22].

Analysis of the protein architecture of the bacterial DPP4-like proteins revealed a leader

peptide similar to that of the human counterpart (Supplementary material S1A-C), which is

involved in transmembrane translocation, suggesting extracellular secretion. Furthermore,

the amino acid profile of all the analyzed DPP4-like proteins (bacterial and eukaryote, N =

971) indicated that all proteins contain the same catalytic motif of S9 serine proteases

(GWSYGGY) according to the MEROPS database (Supplementary material S1D) [23].

Increased abundance of microbial DPP4-like genes is associated with T2D

To investigate the association between microbial DPP4 and T2D, we evaluated the

abundance of all retrieved DPP4-like bacterial genes in the metagenome of 220 subjects

with T2D from a previous study [19]. A GLMM analysis controlling for covariates

indicated that some DPP4-like genes were more abundant in patients with T2D than in non-

T2D controls (Figure 1C). The largest differences were detected for DPP4-like genes from

Parabacteroides sp. (DPP4_927, GLMM-adj = 0.40, p-adj = 0.009), *Chryseobacterium* sp.

YR203 (DPP4_609, GLMM-adj = 0.33, p-adj < 0.001), *Parabacteroides merdae*

(DPP4_931, GLMM-adj = 0.31, p-adj < 0.001), *P. timonensis* (DPP4_302, GLMM-adj =

0.29, p-adj = 0.006) and *Parabacteroides* sp. SN4 (DPP4_249, GLMM-adj = 0.27, p-adj =

0.023). Indeed, a genus-level (taxa grouped) enrichment approach revealed that DPP4-like

genes are encoded essentially by species of two genera in patients with T2D:

Parabacteroides (*Chi-squared* = 122.9, p = 5.0e-4) and *Porphyromonas* (*Chi-squared* =

25.8, p = 0.003). Contrastingly, DPP4-like genes encoded by species of the genera

Bacteroides and *Prevotella* were less abundant in patients with T2D (Figure 1C). We then

used a machine learning approach (sPLS-DA) to question whether the DPP4-like genes

could be used as a classification trait in T2D. Robust models were built using two and three

more informative features from the DPP4-like dataset, which revealed that the genes from

Parabacteroides timonensis (DPP4_302), *Parabacteroides* MULTISPECIES 99%

(DPP4_927) and *Porphyromonas catoniae* (DPP4_761) had high discriminatory power

when evaluated in combination (AUROC = 0.823, p = 6.7e-16) (Figure 1D).

Bacterial DPP4 is secreted like its human counterpart

To understand the extent to which the function of bacterial DPP4-like proteins is similar to

hsDPP4, we cloned the DPP4 gene from *P. merdae*, the member of the gut microbiome that

is more strongly associated with T2D and the *Parabacteroides* species that is more

abundant in the human gut [24]. We cloned the full-length DPP4-like gene (pmDPP4) and a

truncated version lacking the predicted leader sequence of 23 amino acids at the N-terminus

(pmDPP4Δ23) into a His-tag expression vector, and confirmed over-expression of the

recombinant proteins in *E. coli* BL21-DE3 cells (Supplementary material S2A). Mass

spectrometry analysis revealed that cell-free supernatants from *E. coli cultures* expressing

pmDPP4, but not those expressing pmDPP4Δ23, could cleave GLP-1, as reflected by the

expected shift in the molecular mass of GLP-1 from 3354 Da to 3147 Da resulting from the

cleavage of the His-Ala N-terminal dipeptide (Figure 2A-C). We next investigated the

subcellular localization of pmDPP4 and pmDPP4Δ23 proteins in bacterial cultures using

anti-His immunogold staining and transmission electron microscopy. Results showed that

full-length pmDPP4 tended to localize to the inner membrane and was also secreted and

embedded in the extracellular polysaccharide capsule [25], whereas pmDPP4Δ23 tended to

localize in the cytosol (Figure 2D-F).

To confirm the role of the pmDPP4 signal peptide in directing secretion and allowing

extracellular GLP-1 hydrolysis, we performed an *in vitro* enzymatic assay using the

synthetic dipeptide Gly-Pro p-nitroanilide (gpPNA) as a substrate. We also measured the

rate of cell lysis by titrating lactate dehydrogenase (LDH) activity during the preparation of

cell-free supernatants, to control for spontaneous DPP4-like activity. We were unable to

detect bacterial cell lysis during the reaction, and thus the increase in DPP4-like activity

observed in *E. coli* BL21-DE3 cells expressing full-length pmDPP4 (but not in cells

expressing pmDPP4Δ23) was likely mediated by secretion of the accumulated enzyme into

the extracellular medium (Supplementary material S2B). Thus, the leader peptide, which is

evolutionarily conserved in bacterial DPP4-like proteins, is likely required for protein

secretion and favors interaction with their substrates.

Parabacteroides merdae DPP4-like enzyme hydrolyzes human incretins and

neuropeptides in vitro

As our data suggested that the pmDPP4Δ23 protein was likely to be the mature form

produced and secreted by *P. merdae*, we tested additional functional aspects of the protein.

Gel filtration analysis indicated that pmDPP4Δ23 dimerizes in solution to a protein of ~130

kDa, similar to the quaternary structure of the human counterpart (Supplementary material

S2C) [26]. To assess whether the pmDPP4Δ23 protein was similarly active to the full-

length protein produced and secreted by *E. coli* BL21-DE3 cells, we determined its

hydrolytic activity (Supplementary material S2C,D). As a qualitative approach, we tested

whether pmDPP4Δ23 could cleave the incretin and neuropeptide targets of hsDPP4 in a

comparative assay with hsDPP4 by MALDI-TOF-MS (Supplementary material S2D).

Under equimolar conditions and equal substrate concentrations, the pmDPP4Δ23 protein

cleaved GLP-1, GIP, PYY, and NPY peptides as efficiently (and identically) as hsDPP4

(100% substrate hydrolysis) (Figure 3). Specifically, GLP-1 and GIP peptides were

completely processed by pmDPP4Δ23 at the first two N-terminal amino acids (His-Ala and

Tyr-Ala, respectively), consistent with the molecular mass shift (Figure 3A,B).

pmDPP4Δ23 also hydrolyzed the anorexigenic hormone PYY and the neuropeptide NPY

by removing the Tyr-Pro N-terminal dipeptide in both cases, as for hsDPP4 (Figure 3C,D).

Parabacteroides merdae DPP4-like enzyme inactivates GLP-1 and GIP in vivo

To investigate whether the DPP4-like enzyme cleaves and inactivates incretins *in vivo*, we

administered mice with *E. coli* BL21-DE3 carrying the plasmid encoding full-length

pmDPP4 or the empty vector. To mimic obesity-associated low-grade inflammation and

gut barrier defects, mice were injected intraperitoneally with lipopolysaccharide (LPS) to

stimulate metabolic endotoxemia, which would facilitate the translocation of bacterial

products from the intestinal lumen (Figure 4A). We observed significantly higher DPP4

activity in the cecal content and portal vein of the group receiving *E. coli* carrying the

bacterial DPP4 (Figure 4B,C). Supporting our hypothesis, the increase in DPP4 activity in

the portal circulation was associated with a significant reduction in the active forms of

GLP-1 and GIP (Figure 4D,E). We excluded the existence of differences in hormone

secretion, as no changes were found in total GLP-1, its precursor proglucagon or total GIP

(Figure 4F-H). There were also no differences in the total form of other hormones, such as

PYY, or in insulinemia, or glycemia (Figure 4I-K).

P. merdae DPP4-like enzyme is differentially inhibited by gliptins

Finally, we evaluated the inhibition of bacterial DPP4-like and hsDPP4 proteins by

currently approved gliptins: sitagliptin, vildagliptin, saxagliptin and linagliptin.

Lineweaver-Burk double-reciprocal plots of the kinetics showed that all inhibitors showed

classical competitive inhibition of hsDPP4, affecting the K_m but not the V_{max}

(Supplementary material S3). Overall, pmDPP4Δ23 had an estimated K_m of 1.63 mM,

representing on average a 3.97 fold-change (p < 0.001) over the observed hydrolysis rate

for hsDPP4 (Km = 0.33 mM) (Supplementary material S4). As expected, the enzymatic

activity hsDPP4 was inhibited by all gliptins, with sitagliptin, saxagliptin and linagliptin

having a more significant effect on its dipeptide hydrolysis function (Supplementary

material S4). Contrastingly, bacterial pmDPP4Δ23 was only inhibited by saxagliptin and

vildagliptin, which reduced its hydrolytic capacity on gpPNA (fold-change = 1.73, p =

0.036 and fold-change = 0.78, p = 0.076, respectively, at the highest inhibitor

concentration) (Supplementary material S4). The determination of EC_{50} values confirmed

the poor effect of gliptins on pmDPP4 function compared with hsDPP4 (Figure 5). The

EC_{50} for all gliptins against hsDPP4 was in the nanomolar range ($< 10^{-6}$) with linagliptin

showing the strongest inhibitory activity (3.68×10^{-9}). Conversely, pmDPP4Δ23 was only

inhibited by vildagliptin and saxagliptin, but always to a lesser extent than hsDPP4 (2-fold

and 2.4-fold less, respectively) (Table 1). Non-linear fitting of the dose-response data

showed an almost null effect of sitagliptin and linagliptin on pmDPP4Δ23, leaving its

hydrolytic activity intact (Figure 5).

Discussion

DPP4 activity, present on the surface of mammalian cells and as a soluble form in plasma,

is crucial in regulating host metabolism by cleaving and deactivating the incretins that

signal insulin release. The present study makes an advance by showing that the prevalence

of bacterial DPP4-like genes correlates with human metabolic status in several

metagenomic analyses. In addition, our *in vitro* and *in vivo* studies with the DPP4 of *P.*

merdae provide a mechanistic rationale for the findings of our metagenomic assessment

and argue for the detrimental impact of bacterial DPP4 on human health.

To best characterize the gut microbiota as a source of DPP4-like activity, we first searched

for one of the millions of genetic traits encoded in the human gut microbiome based on its

distant similarity to the human *DPP4* or *CD26* gene product. We show that human DPP4

has distant sequence-based homologs that are widely distributed in the genome of hundreds

of Gram-negative bacterial species, some of which are prevalent members of the human gut

microbiota. We demonstrate that the products of such bacterial genes are more similar to

eukaryote DPP4 proteins (e.g., human, rat, and mouse) than to the bacterial XaaPro/PepX

peptidases (so-called S15 family). The PepX family comprises a set of proteins often

referred to as DPP4 enzymes, which has led to confusion in distinguishing the true DPP4-

type enzymes probably because of their similar activity to trim X-A or X-P N-terminal

dipeptides in many substrates [23, 27]. In the latter respect, the conserved and consensus

GWSPGGF motif (GWSYGGY in eukaryotes) present in the bacterial DPP4 homolog has

led to the assignment of this peptidase to the S9B family of serine peptidases according to

the MEROPS database [23]. Our results are in agreement with recent findings showing the

presence of DPP4-like enzymes in several species of *Bacteroides* [14, 16].

We have also addressed one of the criticisms we received when our hypothesis paper was

published, which questioned the ability of bacteria to secrete DPP4 [28]. In this sense, our

new data suggest that bacterial DPP4 proteins contain N-terminal signal peptides that

potentially drive secretion, and are indeed similar to the transmembrane domains present in

the hsDPP4 amino acid sequence. Using various cell-based assays and microscopy

approaches, we show that the leader sequence of bacterial DPP4 proteins, exemplified by

the 23 amino acid N-terminal peptide from *P. merdae*, serves as a signal for secretion. We

conclude that, similar to hsDPP4, the bacterial DPP4-like protein tends to be located

extracellularly rather than intracellularly as for PepX peptidases [29, 30]. In addition,

bacterial DPP4-like proteins share other features with the human enzyme, such as

similarities in the amino acid sequence (~30%), domain architecture, quaternary structure

and catalytic motifs [26, 31].

Given the known role of hsDPP4 as a regulator of glucose homeostasis, we questioned the

value of DPP4 expression in the gut microbiome. Our metagenomic assessment suggests

that DPP4-like genes from *Parabacteroides* and *Porphyromonas* species may play an

important role in T2D, as they are enriched in patients with T2D. Our results are in accord

with the taxonomic evaluation performed in one of the largest cohorts of individuals with

T2D published to date, where *Parabacteroides* appears as one of the genera enriched in

T2D subjects [19]. Much less is known about the effect of *Porphyromonas* on host

metabolism, likely because its predominant habitat is the oral cavity. Nevertheless, some

studies have provided information on the metabolic deterioration caused by

Porphyromonas gingivalis, with recently reported DPP4-like activity [16], when

administered to mice [32]. We hypothesize that the mechanism underlying our results, as

well as those reported by others, lies in the functional overlap of DPP4 activity within

eukaryotic cells and gut microbes. Indeed, we show *in vitro* that the *P. merdae* DPP4-like

enzyme inactivates GLP-1 and GIP as efficiently as hsDPP4. In mice sensitized by LPS, we

confirm that the administration of *E. coli* overexpressing the full-length DPP4-like gene

from *P. merdae* reduces the active forms of GLP-1 and GIP, which is associated with

higher DPP4 activity in portal blood. Very recently, Wang et al. reported similar results

regarding the potential of the DPP4-like activity encoded by *Bacteroides thetaiotaomicron,*

Bacteroides fragilis and *Bacteroides vulgatus* to inactivate GLP-1 in mice fed a high-fat

diet for 12 weeks, but not in mice fed a control diet [16]. Notably, both studies agree that

the gut microbiota contributes to DPP4 activity and the decrease of active GLP-1 when the

gut barrier is disrupted, either by a high-fat diet or by LPS [16]. In addition to the

inactivation of incretins, we also provide evidence for the ability of bacterial DPP4-like to

process other gut peptides *in vitro*, including PYY and NPY, which would have further

implications for host behavior and mental health. For example, full-length PYY binds with

similar affinity to all Y receptors, but the predominant circulating form of PYY-

immunoreactivity, truncated PYY_{3-36}, preferentially binds to the Y2 receptor [33], a G

protein-coupled receptor implicated in appetite and feeding control, mood and mental

disorders [34]. By degrading NPY_{1-36} to NPY_{3-36}, DPP4 increases the affinity of NPY for

Y2 and Y5 receptors but reduces its activity at Y1 receptors, which are known to be

responsible for the antidepressant effects of NPY [35]. Accordingly, in mice fed an

obesogenic diet, increases in DPP4 activity in systemic circulation have been associated

with a depression-like behavior [36], an effect that is counteracted by sitagliptin [37].

Unfortunately, the lack of commercial kits to measure the cleaved forms of PYY and NPY

makes their *in vivo* modification by the microbiota a matter for future studies.

The existence of gut bacteria with DPP4-like activity rationalizes the evaluation of their

interaction with antidiabetic drugs with inhibitory DPP4 activity. Sitagliptin and

vildagliptin were the first drugs to be approved and are the most widely prescribed [8]. Our

results indicate that DPP4 from *P. merdae* was minimally inhibited by sitagliptin, in line

with previous studies on the DPP4-like enzyme from *B. thetaiotaomicron* [16]. By contrast,

vildagliptin and saxagliptin have an effect on the DPP4 activity of *P. merdae*, with EC_{50}

values in the nanomolar range. For vildagliptin, concordant results in mice showed that the

drug abolished DPP4 activity in the contents of the cecum and feces [38]. However, our

results contrast with the recently characterized DPP4-like enzyme from *B.*

thetaiotaomicron, which is inhibited in the nanomolar range (10^{-8} range) by saxagliptin and

in the micromolar range by sitagliptin (10^{-6} range) and linagliptin (10^{-5} range) [14]. These

different inhibition patterns could be explained by the extensive sequence diversity of

bacterial DPP4-like enzymes, resulting in a large repertoire of proteins with differing

abilities to bind gliptins and affect their catalytic activity. Therefore, characterizing the

function of DPP4-like enzymes from different gut bacterial species is crucial to elucidate

their impact on T2D progression and gliptin therapy. Indeed, the information obtained from

our inhibition studies could have significant consequences for the pharmacology of T2D

treatment. For instance, the partial inhibition of the *P. merdae* DPP4-like activity by

vildagliptin and saxagliptin suggests that bacterial enzymes can bind DPP4 inhibitors

regardless of the inhibition efficacy. Given the substantial prevalence of DPP4-like

enzymes in the intestinal lumen, this phenomenon could significantly affect the

bioavailability of drugs targeting the human enzyme [39]. Furthermore, we and others have

demonstrated that sitagliptin does not inhibit bacterial DPP4, highlighting the limitation of

this drug to prevent incretin inactivation from all possible enzymatic sources. This

observation likely explains the differential response to sitagliptin recently reported in

individuals with T2D [16].

Conclusions

We show that the gut microbiome encodes a large number of DPP4-like functional

homologs, the prevalence of which positively correlates with T2D status. The mechanism

underlying this association might be functional mimicry of bacterial DPP4-like enzymes, as

exemplified by the ability of *P. merdae* DPP4-like enzyme to inactivate incretins. Our

results of the characterization of the inhibitory activity of gliptins on bacterial DPP4-like

enzymes may serve as a starting point for redefining the efficacy of current DPP4

inhibitors, as we show that some DPP4 inhibitors (vildagliptin and saxagliptin) are

compromised by nonspecific binding to bacterial enzymes and others, such as sitagliptin,

by the null effect on bacterial DPP4-like enzymes.

Material and methods

Search strategy for bacterial DPP4 homologs in the gut microbiome

We searched for bacterial homologs of hsDPP4 in metagenome-assembled genomes from our previous study on people with overweight and the metabolic syndrome [20]. We used a Blast-based approach against human (SwissProt id: P27487), mouse (P28843) and rat (P14740) DPP4 amino acid sequences. Thresholds were set to at least 85% coverage in length and >25% sequence identity at the protein level.

Phylogeny tree of bacterial DPP4 and X-prolyl dipeptidyl peptidase families

The amino acid profile (*via hmmer3*) of the aligned DPP4-like (S9 peptidase family) sequences detected in the metagenome of overweight subjects with metabolic syndrome was used as bait to compile the most complete set of similar proteins present in the RefSeq NCBI database through PSI-Blast [40]. In similar manner, the PepX (S15 peptidase family) from *Lactobacillus acidophilus* NCFM (UniProt id Q5FJC6) was used to retrieve the most complete set of similar enzymes potentially present in bacterial genomes. Multiple PSI-Blast iterations were run to select aligned protein sequences with coverage of >95% of query length and ≥25% amino acid identity. Redundancy was removed to retain sequences at 99% identity. Finally, we obtained 925 non-redundant amino acid sequences of bacterial DPP4 homologs and 671 sequences for the PepX family of peptidases. Human (hsDPP4), mouse (mmDPP4) and rat (rnDPP4) sequences were merged with those previously obtained from bacterial metagenes as well as with the protein P19577 from *Dichelobacter nodosus* (member of the S17 peptidase family), the outlier for phylogeny reconstruction. In total, we performed a multiple sequence alignment of more than of >1,600 protein sequences, using an iterative process with refinement steps [41, 42]. Amino acid profiles and consensus

sequences for both peptidase families were computed using the Hidden Markov Model

approach and the *hmmer3* package [43, 44]. The best tree explaining the evolutionary

relationships between the two types of peptidases was built using ProtTest v3.0 [45], by

selecting the best of the 120 evolutionary models implemented, according to AIC. Tree

topology was optimized by using the maximum likelihood method. The amino acid

sequences derived from hsDPP4 and the consensus amino acid sequences obtained for S15

and S9 peptidase families using *hmmer3* were subjected to further analysis to depict protein

architecture and functional domains using the SMART server [46]. In a similar manner, the

SignalP v4.1 server was used to predict the presence of signal peptides in the amino acid

sequences (http://www.cbs.dtu.dk/services/SignalP-4.1/) [47], whereas the TMHMM server

was used to predict the presence of transmembrane helices

(http://www.cbs.dtu.dk/services/TMHMM/).

Cloning, over-expression and purification of bacterial DPP4

The *P. merdae* DPP4-like gene (pmDPP4) was amplified from DNA obtained from the

type strain *P. merdae* DSM19495[T] (DSMZ, Braunschweig, Germany), using the primers

DPP4-PMER-F gata<u>ccatggt</u>gaaacgattagggtttggtgcttta (with an NcoI restriction site,

underlined) and DPP4-PMER-R

gata<u>ggatcc</u>ttatta*gtgatggtgatggtgatg*caaattatcaaataaaaagttagacat (with a BamHI restriction

site, underlined) and a 6×His tag (italics) to produce a C-terminal tagged protein.

Additionally, an open reading frame (ORF) encoding an N-terminal 6×His-tagged pmDPP4

protein with no signal peptide (pmDPP4Δ23) was amplified from human fecal DNA

samples using the primers DPP4-PMER-F23

gata<u>ccatggg</u>a*catcaccatcaccatcac*agcaaacgtgtcgatttaaaagaaattaca and DPP4-PMER-R2

gataggatccttacaaattatcaaataaaaagttagacat. The PCR products of (~2.5 kb) were purified,

mixed, and 500 ng of DNA was digested with 5U NcoI and BamHI restriction enzymes

(New England Biolabs, Ipswich, MA) overnight at 37ºC. Digested fragments were cloned

into the BamHI/NcoI pre-digested pET15b plasmid using the Fast-Link™ DNA Ligation

Kit (Epicentre, Madison, WI), and *E. coli* DH5α cells were transformed with the ligation

products. Colonies were selected on LB agar with 100 µg/mL ampicillin (Sigma, Madrid,

Spain) and checked by PCR for the presence of the correct plasmids. The inserts were

analyzed by Sanger sequencing to ensure in-frame position of the ORFs and tags. The

plasmids encoding pmDPP4 and pmDPP4Δ23 ORFs were isolated from DH5α cells and

transferred to *E. coli* BL21-DE cells by heat-shock transformation. To over-express the

recombinant proteins, freshly transformed *E. coli* BL21-DE cells containing the respective

plasmids were grown on NZY Auto-induction LB medium (NZYTech, Lisbon, Portugal) at

37ºC for 20 h. Then, bacterial cells were recovered by centrifugation at 5,000 *g* for 25 min

at 4ºC and washed and resuspended in 10 mL of TBS buffer pH 8.0. Cells were disrupted

by a pre-incubation with 500 µg lysozyme (Sigma) at room temperature for 10 min,

followed by 4 cycles of bead-beating (FastPrep-24™ 5G bead beating grinder; MP

Biomedicals, Santa Ana, CA) of 30 sec, using 0.1 mm glass beads. The supernatant was

obtained by centrifugation of the cell lysate at 16,000 *g* for 20 min at 4ºC, and was then

incubated with 2 mL TBS pH 8.0 pre-equilibrated Talon® Metal Affinity resin during 2 h

at 4ºC with permanent mixing (Takara Bio, Kusatsu, Japan). The resin was washed with 10

volumes of TBS pH 8.0 buffer containing 0.1% Triton X-100 and 10 mM imidazol (both

from Sigma). Elution was performed with 10 resin volumes of TBS pH 8.0 containing 150

mM imidazol. Proteins were concentrated by filtration of the eluate in 15-mL Amicon®

Ultra 50k filters (Millipore, Burlington, MA). Subsequently, a concentration step was

performed in 0.5-mL Amicon® Ultra 50k filters, retaining a final volume of 30–50 μL.

Purity and concentration of the recombinant proteins was assessed by SDS-PAGE and the

Qubit Protein Assay Kit (Thermo Fisher Scientific, Waltham, MA). Gel filtration was used

to recover the recombinant protein fraction according to its expected size on an AKTA

Ettan LC FPLC System (Amersham Pharmacy Biotech, Amersham, UK), with a Superdex

200 5/150GL column and Ca/Mg-free sterile PBS pH 7.4 (MP Biomedicals, Aurora, OH)

as the mobile phase.

Immunogold staining and electron microscopy

E. coli BL21-DE cells containing the pET15b/pmDPP4 or pET15b/pmDPP4Δ23 plasmids

were cultured in LB medium and induced to express the recombinant proteins as described

above. *E. coli* BL21-DE cells harboring the pET15b were equally treated and used as a

negative control for immunogold staining. After the induction of protein expression

induction, 0.5 mL of cells were centrifuged at 3,000 *g* and washed with 1 volume of PBS.

Cells were pelleted and sent to the Electron Microscopy Core Facility at SCSIE (University

of Valencia) for fixation, embedding and immunogold staining using a 6×His monoclonal

antibody (Takara Bio) diluted 1/5000 as the primary antibody, and 10 nM anti-mouse IgG-

Gold antibody (Sigma) as the secondary antibody, diluted 1/20. Images were acquired on a

JEM-1010 transmission electron microscope (JEOL, Tokyo, Japan) operating at 100 KV

equipped with an AMT RX80 (8 Mpx) digital camera. Image analysis was performed at the

SCSIE.

Protein secretion assay

A colorimetric assay was developed to detect pmDPP4 secretion dependent on the presence

of the 23 amino acid leader peptide. *E. coli* BL21-DE3 cells containing the plasmids

pET15b/pmDPP4 or pET15b/pmDPP4Δ23 were grown overnight at 37ºC in LB medium

with 100 µg/mL ampicillin (Sigma). Overnight cultures were 1:100 refreshed in 5 mL

ampicillin-containing LB medium and grown for a further 3 h at 37ºC with shaking. For

recombinant protein over-expression, we added 0.1 mM IPTG (isopropyl β d-1-

thiogalactopyranoside) and 100 µg/mL ampicillin to cultures grown under the same

condition for a further 3 h. After protein induction, 1 mL of culture was recovered in a 1.5-

mL sterile microtube and cells were pelleted at 3000 *g* for 10 min at 4ºC. Pellets were

resuspended in 1 mL of cold PBS and incubated on ice for 5 min. From the suspension, 100

µL was removed to estimate the colony formation unit (CFU) count on LB-ampicillin

plates. The remaining cell suspensions were again pelleted and finally resuspended in 200

µL of cold PBS. The secretion assay consisted of adding 450 µL PBS to 50 µL cellular

suspension, mixing, and incubating at 37ºC with 600 rpm shaking. One hundred microliters

of the suspension was collected in new 1.5-mL sterile microtubes at 0, 30, 60, and 90 min

of incubation and were maintained on the ice up until the end of the collection. To obtain

cell-free supernatants, all aliquots were centrifuged at 6000 *g* for 10 min at 4ºC and

supernatants were recovered in 1.5-mL microtubes and stored on ice until processing. The

cell-free supernatants were used to determine DPP4 activity secreted into the medium

according to the protocol described below. Bacterial cell lysis was determined in cell-free

supernatants using the LDH Cytotoxicity Assay Kit (Thermo Fisher Scientific, Waltham,

MA). Both DPP4 and LDH activities obtained from cell-free supernatants were normalized

to the CFU counts.

Colorimetric measurement of DPP4 activity and inhibition assays

The hydrolytic activity of the bacterial DPP4 enzyme was measured using Gly-Pro p-

nitroanilide hydrochloride (gpPNA) (Sigma), as described for assessing hsDPP4 activity

[38]. A range of substrate concentrations from 2.5 to 0.31 mM gpPNA in 50 mM Tris-HCl

pH 8.3 was used for kinetic analysis. The standard curve was established with free para-

nitroaniline (PNA) (Sigma) ranging from 2 mM to 0.031 mM in 50 mM Tris-HCl pH 8.3.

Briefly, 20 µL of each cell-free supernatant was placed in triplicate in a flat 96-well plate.

Two hundred microliters of the respective PNA solutions were plated in duplicate to obtain

the standard curve. A total of 180 µL of substrate solution (gpPNA) was added to bacterial

supernatants and the enzymatic activity was monitored for 30 min at 37°C (380 nm) in a

VICTOR2 fluorometer/spectrophotometer (Perkin Elmer, Waltham, MA). The enzymatic

activity ([PNA generated (nM) / hydrolysis time (min)] / pmole enzyme) was quantified

between 10 and 30 min of incubation time (linear kinetics). Three independent experiments

with different purified protein batches were performed. The same protocol was followed for

the murine samples (cecal content and plasma from the portal vein). In the case of the cecal

content, the values of DPP4 activity were normalized by the amount of protein, quantified

by the Bradford method (Bio-Rad, Hercules, CA).

The commercially available FDA-approved human DPP4 inhibitors sitagliptin, saxagliptin,

linagliptin and vildagliptin (all from Sigma) were diluted in PBS to a 2 µM working

solution. The inhibitory activity of the four gliptins on hsDPP4 and pmDPP4Δ23 was

assessed at the ratios 1:10, 1:50, and 1:100 purified protein:inhibitor. Four independent

replicates with different purified protein batches were performed. Dose-response assays to

determine the EC_{50} were carried out using serial dilutions of the inhibitors from 1 μM to

3.91 nM, and the relative activity was calculated with respect to reactions lacking the

inhibitor.

Peptidase activity of DPP4-like enzymes on gut hormones

The DPP4 substrates GLP-1 7-37 fragment, GIP, NPY and PYY (all from Sigma) were

diluted to 2.5 μg/μL in Ca/Mg-free sterile PBS pH 7.4. Human DPP4 (Sigma) was used as

a control in PBS at 50 ng/μL (0.585 μM). The *in vitro* peptidase activity was assayed in a

10-μL reaction containing 5 μg substrate (100-150 μM) and 10 nM of hsDPP4 or

pmDPP4Δ23 recombinant proteins (molar ratio substrate:enzyme, 10000:1), mixed on ice

and incubated at 37ºC for 10 min with gentle orbital shaking (600 rpm). To test the activity

of the pmDPP4 secreted protein, the reaction was the same but the purified recombinant

protein was replaced with the same volume of *E. coli* BL21-DE cells carrying pET15b,

pET15b/pmDPP4, or pET15b/pmDPP4Δ23 plasmids incubated with 0.1 mM IPTG for 3 h

at 37ºC. The cells were washed and diluted in PBS (~5 × 10^6 cells per reaction). Inhibition

assays were performed under similar conditions but in the presence of 500 nM of

vildagliptin or sitagliptin. Hydrolysis reactions were inactivated by incubation at 90ºC for

10 min. For reactions containing cells, the cells were first collected by centrifugation and

then heat-inactivated. The inactivated reactions were stored on ice until processed for MS.

MALDI-TOF analysis

The MS analysis was performed at the SCSIE proteomic service (University of Valencia).

Briefly, for every *in vitro* reaction performed, 1 μL of each sample was spotted onto the

MALDI plate and, after the droplets were air-dried at room temperature, 1 μL of matrix (10

mg/mL alpha-cyano matrix in 70% acetonitrile, 0.1% trifluoroacetic acid) was added and

allowed to air-dry at room temperature. The resulting mixtures were analyzed on a 5800

MALDI TOF/TOF instrument (AB Sciex, Framingham, MA) in positive reflector and

linear mode. The raw mass spectra were released in ASCII format and were analyzed using

mMass v5.5 software (http://www.mmass.org/).

Bacterial DPP4-like in T2D human metagenomes

A total of 220 samples (metadata available) derived from one of the largest metagenome

projects on T2D (BioProject PRJNA422434) were evaluated for the distribution of bacterial

DPP4-like genes. For read mapping against the non-redundant DPP4-like peptide database

(N = 971), we used the *usearch* v8.0.1623 algorithm [48] with the following parameters: -

usearch_local, *-id 0.7*, *-strand both*, and *-top_hit_only*. Alignments were filtered to retain

those with more than 66% read length. Differential abundance of DPP4-like genes was

assessed using Generalized Linear Mixed Models (GLMM, *lme4::glmer* R function) with

prior filtering for low abundant features (*CoDaSeq::codaSeq.filter*, *min.occurrence*=0.25,

min.reads=1), execution of *zCompositions::cmultRepl* function supporting a Bayesian-

Multiplicative replacement of zero counts, and centered log-ratio (clr) transformation of

compositional microbial abundance data (*CoDaSeq::codaSeq.clr*). Changes in DPP4-like

gene abundance were assessed individually using sex, age, and body mass index (BMI) as

covariates in the GLMM model. Genus enrichment analysis on significantly abundant

DPP4-like genes in subjects with T2D (N = 135) compared to non-T2D individuals (N =

85) (from same cohort) was evaluated using the *Chi-square* test with Monte-Carlo

simulation. Lastly, the DPP4-like gene abundance from T2D (N = 135) and no-T2D (N =

85) samples was compiled in an integrative analysis supported by sparse Partial Least

Square-Discriminant Analysis (sPLS-DA) algorithms implemented in the *DIABLO* v6.10.8

package [49]. A tuned sPLS-DA model was built using 4-fold cross-validation (limiting

over-fitting) across multiple iterations (*nrepeat* = 20). Selection of reliable models

explaining variation was based on their performance (AUROC ≥ 0.70).

Animal study

Experiments were performed using 16 male C57BL/6J mice (7 weeks of age; Charles River

Laboratories, Écully, France). Mice were housed in individually-ventilated cages (4

mice/cage) in a Biosafety level 2 laboratory with a 12-h light/dark cycle and temperature-

controlled conditions (23 ± 2°C). Mice were acclimatized for ten days and had *ad libitum*

access to food and water. Mice were randomized to receive for five consecutive days *E.*

coli BL21-DE cells carrying the empty expression plasmid pET15b (3×10^8 UCF/day) (N

= 8) or carrying pET15b/pmDPP4 (3×10^8 UCF/day) (N = 8). Both groups were also

treated with LPS from *E. coli* O111:B4 (0.1 mg/kg, dissolved in sterile PBS) to induce

intestinal permeability, as previously described [50]. Mice were fasted on day 5. Mice then

received an oral load of Ensure Plus® (0.0124 kcal/g) to induce the release of gut hormones

[51]. After 12 min, mice were anesthetized with isoflurane to sample blood from the portal

vein, which was collected in EDTA-containing tubes with or without a DPP4 inhibitor

(Millipore, Burlington, MA). The plasma was recollected after centrifugations at 12,000 *g*

for 3 min. Mice were sacrificed by cervical dislocation to minimize potential confounding

effects, and mice of the different experimental groups were interspersed. The cecal content

was also collected and snap-frozen in liquid nitrogen. All samples were stored at -80°C. In

plasma samples containing the DPP4 inhibitor, the levels of active and total GLP-1 and

total GIP, PYY, insulin and proglucagon were measured using the Luminex™ Mouse

Metabolic Hormone Expanded kit (Merck & Co., Inc. Kenilworth, NJ). Also, active GIP

was measured using an ELISA kit (CrystalChem, Zaandam, The Netherlands). Glycemia

postmortem was quantified using the Glucose Colorimetric Detection Kit (ThermoFisher

Scientific). DPP4 activity was measured in an aliquot of plasma without DPP4 inhibitor

and in the cecal content and feces, as described above.

Statistical analysis

A normality test (Shapiro-Wilk) was conducted on data to select the most appropriate

statistical comparison test (parametric or non-parametric) between sample groups and

treatments. The *t*-test or Wilcoxon-Rank Sum test was applied to compare the *in vitro*

enzymatic activity of hsDPP4 and pmDPP4 and the biochemical and physiological data

from the animal study. A GLMM with covariate controls (e.g., age, sex and BMI) was used

to evaluate the differential abundance of DPP4-like genes in patients with T2D. The

sequencing project was also included as a covariate when multiple cohorts of metagenome

samples were assessed simultaneously. The sPLS-DA was used to quantify the power of

bacterial DPP4-like genes for the classification of T2D samples, using a 4-fold cross-

validation. All tests were applied in R v4.1.2. Estimation statistics and plots were obtained

using the *dabestr* v0.3.0 R package (Gardner-Altman plots) or GraphPad v9 (San Diego,

CA).

Declarations

Ethics approval:

The animal experiment was performed following European Union 2010/63/UE and Spanish RD53/201 guidelines, approved by the ethics committee of the University of Valencia (Animal Production Section, SCSIE, University of Valencia), and authorized (Ref 2022/VSC/PEA/0244) by the competent authority (Generalitat Valenciana).

Availability of data and material.

The raw metagenomic data was obtained through accessing the following bioprojects publicly available *via* European Nucleotide Archive (ENA): PRJNA422434 and PRJEB25727.

Competing interests

The authors express to have no conflict of interest during the execution of this research.

Funding

The authors acknowledge the following funding bodies that provided support for this study: European Commission 7[th] Framework Program through the MyNewGut project (Grant agreement No. 613979) and the Spanish Ministry of Science and Innovation (MICINN) (grant PID2020-119536RB-I00) to YS; Institute of Health Carlos III (ISCIII) via CP19/00132 grant to AB-P with funds from the European Social Fund (ESF/FSE), CIPF Predoctoral Training Fellowships 2022 funded by JANSSEN-CILAG – supporting the contract of PH-C, CIPF core funding for the contract of SC-B, an Early Career Grant from the Society for Endocrinology award to MO, Spanish Government MCIN/AEI for the Center of Excellence Accreditation Severo Ochoa (CEX2021-001189-S/ MCIN/AEI / 10.13039/501100011033).

Author contributions

MO, YS and AB-P, intellectually conceived the study. PH-C, SC-B and AB-P, performed the molecular biology experiments and the *in vitro* experiments. AB-P performed the bioinformatics and metagenomic analysis. MO and RL-G performed the animal experiments and sample analysis. MO, YS and AB-P drafted the manuscript. All authors read the final version. YS and AB-P are responsible for its content.

Acknowledgements

The authors thank the Computational Biology & Data Science Unit from Principe Felipe Research Center (CIPF) for providing access to its high-performance computing cluster, co-funded by European Regional Development Funds (ERDF/FEDER). Finally, the authors thank the technical staff of the proteomics, microscopy and animal facilities of SCSIE University of Valencia and, in particular, I. Noguera, for their assistance in protocols and procedures described for animal experimentation.

References

1. de Vos WM, Tilg H, Van Hul M, Cani PD: **Gut microbiome and health: mechanistic insights.** *Gut* 2022, **71:**1020-1032.

2. Stebbins CE, Galan JE: **Structural mimicry in bacterial virulence.** *Nature* 2001, **412:**701-705.

3. Cohen LJ, Esterhazy D, Kim SH, Lemetre C, Aguilar RR, Gordon EA, Pickard AJ, Cross JR, Emiliano AB, Han SM, et al: **Commensal bacteria make GPCR ligands that mimic human signalling molecules.** *Nature* 2017, **549:**48-53.

4. Cohen LJ, Kang HS, Chu J, Huang YH, Gordon EA, Reddy BV, Ternei MA, Craig JW, Brady SF: **Functional metagenomic discovery of bacterial effectors in the human microbiome and isolation of commendamide, a GPCR G2A/132 agonist.** *Proc Natl Acad Sci U S A* 2015, **112:**E4825-4834.

5. Breton J, Tennoune N, Lucas N, Francois M, Legrand R, Jacquemot J, Goichon A, Guerin C, Peltier J, Pestel-Caron M, et al: **Gut Commensal E. coli Proteins Activate Host Satiety Pathways following Nutrient-Induced Bacterial Growth.** *Cell Metab* 2016, **23:**324-334.

6. Tennoune N, Chan P, Breton J, Legrand R, Chabane YN, Akkermann K, Jarv A, Ouelaa W, Takagi K, Ghouzali I, et al: **Bacterial ClpB heat-shock protein, an antigen-mimetic of the anorexigenic peptide alpha-MSH, at the origin of eating disorders.** *Transl Psychiatry* 2014, **4:**e458.

7. Mulvihill EE, Drucker DJ: **Pharmacology, physiology, and mechanisms of action of dipeptidyl peptidase-4 inhibitors.** *Endocr Rev* 2014, **35:**992-1019.

8. Sneha P, Doss CG: **Gliptins in managing diabetes - Reviewing computational strategy.** *Life Sci* 2016, **166:**108-120.

9. Ahmed RH, Huri HZ, Muniandy S, Al-Hamodi Z, Al-Absi B, Alsalahi A, Razif MF: **Altered circulating concentrations of active glucagon-like peptide (GLP-1) and dipeptidyl peptidase 4 (DPP4) in obese subjects and their association with insulin resistance.** *Clin Biochem* 2017, **50:**746-749.

10. Lee SA, Kim YR, Yang EJ, Kwon EJ, Kim SH, Kang SH, Park DB, Oh BC, Kim J, Heo ST, et al: **CD26/DPP4 levels in peripheral blood and T cells in patients with type 2 diabetes mellitus.** *J Clin Endocrinol Metab* 2013, **98:**2553-2561.

11. Mulvihill EE, Varin EM, Gladanac B, Campbell JE, Ussher JR, Baggio LL, Yusta B, Ayala J, Burmeister MA, Matthews D, et al: **Cellular Sites and Mechanisms Linking Reduction of Dipeptidyl Peptidase-4 Activity to Control of Incretin Hormone Action and Glucose Homeostasis.** *Cell Metab* 2017, **25:**152-165.

12. Varin EM, Mulvihill EE, Beaudry JL, Pujadas G, Fuchs S, Tanti JF, Fazio S, Kaur K, Cao X, Baggio LL, et al: **Circulating Levels of Soluble Dipeptidyl Peptidase-4 Are Dissociated from Inflammation and Induced by Enzymatic DPP4 Inhibition.** *Cell Metab* 2019, **29:**320-334 e325.

13. Olivares M, Schuppel V, Hassan AM, Beaumont M, Neyrinck AM, Bindels LB, Benítez-Páez A, Sanz Y, Haller D, Holzer P, Delzenne NM: **The potential role of the dipeptidyl peptidase-4-like activity from the gut microbiota on the host health.** *Front Microbiol* 2018, **9:**1900.

14. Keller LJ, Nguyen TH, Liu LJ, Hurysz BM, Lakemeyer M, Guerra M, Gelsinger DJ, Chanin R, Ngo N, Lum KM, et al: **Chemoproteomic identification of a DPP4 homolog in Bacteroides thetaiotaomicron.** *Nat Chem Biol* 2023.

15. Mills RH, Dulai PS, Vazquez-Baeza Y, Sauceda C, Daniel N, Gerner RR, Batachari LE, Malfavon M, Zhu Q, Weldon K, et al: **Multi-omics analyses of the ulcerative colitis gut microbiome link Bacteroides vulgatus proteases with disease severity.** *Nat Microbiol* 2022, **7:**262-276.

16. Wang K, Zhang Z, Hang J, Liu J, Guo F, Ding Y, Li M, Nie Q, Lin J, Zhuo Y, et al:
Microbial-host-isozyme analyses reveal microbial DPP4 as a potential antidiabetic
target. *Science* 2023, **381:**eadd5787.
17. Anastasiou R, Papadelli M, Georgalaki MD, Kalantzopoulos G, Tsakalidou E: **Cloning and**
sequencing of the gene encoding X-prolyl-dipeptidyl aminopeptidase (PepX) from
Streptococcus thermophilus strain ACA-DC 4. *J Appl Microbiol* 2002, **93:**52-59.
18. Stressler T, Eisele T, Schlayer M, Lutz-Wahl S, Fischer L: **Characterization of the**
recombinant exopeptidases PepX and PepN from Lactobacillus helveticus ATCC
12046 important for food protein hydrolysis. *PLoS One* 2013, **8:**e70055.
19. Qin J, Li Y, Cai Z, Li S, Zhu J, Zhang F, Liang S, Zhang W, Guan Y, Shen D, et al: **A**
metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature* 2012,
490:55-60.
20. Benitez-Paez A, Kjolbaek L, Gomez Del Pulgar EM, Brahe LK, Astrup A, Matysik S,
Schott HF, Krautbauer S, Liebisch G, Boberska J, et al: **A Multi-omics Approach to**
Unraveling the Microbiome-Mediated Effects of Arabinoxylan Oligosaccharides in
Overweight Humans. *mSystems* 2019, **4**.
21. Galperin MY, Koonin EV: **Divergence and convergence in enzyme evolution.** *J Biol*
Chem 2012, **287:**21-28.
22. McDaniel LD, Young E, Delaney J, Ruhnau F, Ritchie KB, Paul JH: **High frequency of**
horizontal gene transfer in the oceans. *Science* 2010, **330:**50.
23. Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD: **The MEROPS**
database of proteolytic enzymes, their substrates and inhibitors in 2017 and a
comparison with peptidases in the PANTHER database. *Nucleic Acids Res* 2018,
46:D624-D632.
24. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen T, Pons N,
Levenez F, Yamada T, et al: **A human gut microbial gene catalogue established by**
metagenomic sequencing. *Nature* 2010, **464:**59-65.
25. Andreishcheva EN, Vann WF: **Escherichia coli BL21(DE3) chromosome contains a**
group II capsular gene cluster. *Gene* 2006, **384:**113-119.
26. Chung KM, Cheng JH, Suen CS, Huang CH, Tsai CH, Huang LH, Chen YR, Wang AH,
Jiaang WT, Hwang MJ, Chen X: **The dimeric transmembrane domain of prolyl**
dipeptidase DPP-IV contributes to its quaternary structure and enzymatic activities.
Protein Sci 2010, **19:**1627-1638.
27. Rawlings ND, Barrett AJ: **Evolutionary families of peptidases.** *Biochem J* 1993, **290 (Pt**
1):205-218.
28. Da Silva RR: **Commentary: The Potential Role of the Dipeptidyl Peptidase-4-Like**
Activity From the Gut Microbiota on the Host Health. *Front Microbiol* 2018, **9:**3313.
29. Hafeez Z, Cakir-Kiefer C, Lecomte X, Miclo L, Dary-Mourot A: **The X-prolyl dipeptidyl-**
peptidase PepX of Streptococcus thermophilus initially described as intracellular is
also responsible for peptidase extracellular activity. *J Dairy Sci* 2019, **102:**113-123.
30. Tsakalidou E, Anastasiou R, Papadimitriou K, Manolopoulou E, Kalantzopoulos G:
Purification and characterisation of an intracellular X-prolyl-dipeptidyl
aminopeptidase from Streptococcus thermophilus ACA-DC 4. *J Biotechnol* 1997,
59:203-211.
31. Durinx C, Lambeir AM, Bosmans E, Falmagne JB, Berghmans R, Haemers A, Scharpe S,
De Meester I: **Molecular characterization of dipeptidyl peptidase activity in serum:**
soluble CD26/dipeptidyl peptidase IV is responsible for the release of X-Pro
dipeptides. *Eur J Biochem* 2000, **267:**5608-5613.
32. Yoshida S, Hatasa M, Ohsugi Y, Tsuchiya Y, Liu A, Niimi H, Morita K, Shimohira T,
Sasaki N, Maekawa S, et al: **Porphyromonas gingivalis Administration Induces**
Gestational Obesity, Alters Gene Expression in the Liver and Brown Adipose Tissue

in Pregnant Mice, and Causes Underweight in Fetuses. *Front Cell Infect Microbiol*
2021, **11:**745117.

33. Murphy KG, Bloom SR: **Gut hormones and the regulation of energy homeostasis.**
Nature 2006, **444:**854-859.

34. Parker SL, Balasubramaniam A: **Neuropeptide Y Y2 receptor in health and disease.** *Br J*
Pharmacol 2008, **153:**420-431.

35. Redrobe JP, Dumont Y, Fournier A, Quirion R: **The neuropeptide Y (NPY) Y1 receptor**
subtype mediates NPY-induced antidepressant-like activity in the mouse forced
swimming test. *Neuropsychopharmacology* 2002, **26:**615-624.

36. Hassan AM, Mancano G, Kashofer K, Frohlich EE, Matak A, Mayerhofer R, Reichmann F,
Olivares M, Neyrinck AM, Delzenne NM, et al: **High-fat diet induces depression-like**
behaviour in mice associated with changes in microbiome, neuropeptide Y, and brain
metabolome. *Nutr Neurosci* 2019, **22:**877-893.

37. Magdy YM, El-Kharashi OA, Nabih ES, Shaker SM, Abd-Elaziz LF, Aboul-Fotouh S:
Potential involvement of JNK1 repression in the hepatic effect of sitagliptin and
metformin in rats subjected to high fat diet and chronic mild distress. *Biomed*
Pharmacother 2017, **85:**225-238.

38. Olivares M, Neyrinck AM, Potgens SA, Beaumont M, Salazar N, Cani PD, Bindels LB,
Delzenne NM: **The DPP-4 inhibitor vildagliptin impacts the gut microbiota and**
prevents disruption of intestinal homeostasis induced by a Western diet in mice.
Diabetologia 2018, **61:**1838-1848.

39. Schnapp G, Klein T, Hoevels Y, Bakker RA, Nar H: **Comparative Analysis of Binding**
Kinetics and Thermodynamics of Dipeptidyl Peptidase-4 Inhibitors and Their
Relationship to Structure. *J Med Chem* 2016, **59:**7466-7477.

40. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: **Gapped**
BLAST and PSI-BLAST: a new generation of protein database search programs.
Nucleic Acids Res 1997, **25:**3389-3402.

41. Edgar RC: **MUSCLE: a multiple sequence alignment method with reduced time and**
space complexity. *BMC Bioinformatics* 2004, **5:**113.

42. Edgar RC: **MUSCLE: multiple sequence alignment with high accuracy and high**
throughput. *Nucleic Acids Res* 2004, **32:**1792-1797.

43. Eddy SR: **Multiple alignment using hidden Markov models.** *Proc Int Conf Intell Syst*
Mol Biol 1995, **3:**114-120.

44. Eddy SR: **Hidden Markov models.** *Curr Opin Struct Biol* 1996, **6:**361-365.

45. Darriba D, Taboada GL, Doallo R, Posada D: **ProtTest 3: fast selection of best-fit models**
of protein evolution. *Bioinformatics* 2011, **27:**1164-1165.

46. Letunic I, Doerks T, Bork P: **SMART: recent updates, new developments and status in**
2015. *Nucleic Acids Res* 2014, **43:**D257-260.

47. Petersen TN, Brunak S, von Heijne G, Nielsen H: **SignalP 4.0: discriminating signal**
peptides from transmembrane regions. *Nat Methods* 2011, **8:**785-786.

48. Edgar RC: **Search and clustering orders of magnitude faster than BLAST.**
Bioinformatics 2010, **26:**2460-2461.

49. Singh A, Shannon CP, Gautier B, Rohart F, Vacher M, Tebbutt SJ, Le Cao KA: **DIABLO:**
an integrative approach for identifying key molecular drivers from multi-omics
assays. *Bioinformatics* 2019, **35:**3055-3062.

50. Guo S, Al-Sadi R, Said HM, Ma TY: **Lipopolysaccharide causes an increase in intestinal**
tight junction permeability in vitro and in vivo by inducing enterocyte membrane
expression and localization of TLR-4 and CD14. *Am J Pathol* 2013, **182:**375-387.

51. Vannan DT, Bomhof MR, Reimer RA: **Comparison of Glucose and Satiety Hormone**
Response to Oral Glucose vs. Two Mixed-Nutrient Meals in Rats. *Front Nutr* 2018,
5:89.

784 52. Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J,
785   Ludwig W, Glockner FO: **The SILVA and "All-species Living Tree Project (LTP)"**
786   **taxonomic frameworks.** *Nucleic Acids Res* 2014, **42:**D643-648.
787 53. Letunic I, Bork P: **Interactive tree of life (iTOL) v3: an online tool for the display and**
788   **annotation of phylogenetic and other trees.** *Nucleic Acids Res* 2016, **44:**W242-245.

     Figure legends

Figure 1. A, Taxonomic distribution and phylogenetic analysis of PepX and DPP4
peptidases. The phylogenetic relationship between both families of bacterial peptidases is
illustrated by a circular tree. A small unrooted tree is also shown at the top-right corner.
The tree includes the bait used in the PSI-Blast search for distant homologs and the human,
mouse and rat sequences of DPP4 enzymes for comparison. The outlier consisted of an S17
peptidase sequence from *Dichelobacter nodosus* (SwissProt ID P19577). The taxonomy
assignment of sequences at the family and genus level was performed using the SILVA
database taxonomy annotation (release 138) [52]. Branch colors are according to the
peptidase family grouping: green for XaaPro peptidases (S15), orange for DPP4 peptidases
(S9) and blue for the outlier (S17). Strip colors indicate the taxonomic distribution of
sequences accordingly. Labels for the most abundant taxonomy groups in terms of families
and genera (numbers within parenthesis) are shown. The tree was visualized using the
iTOL web server [53]. **B**, Distribution of amino acid identities of eukaryote proteins *versus*
bacterial DPP4-like sequences is shown as a boxplot. **C**, Metagenome abundance of DPP4-
like proteins for classification of type 2 diabetes (T2D). A volcano plot shows the
abundance variation of bacterial DPP4-like proteins in human gut metagenomes linked to
T2D. Generalized linear mixed models (GLMM) estimates and p-values obtained after
controlling for covariates (sex, age, body mass index) were used to identify proteins
associated with T2D (N = 135) *versus* no-T2D controls (N = 85). A 2×2 contingency
table-based *Chi*-squared test with Monte-Carlo simulation was computed to determine
protein enrichment at the genera level. The threshold for selection was set at $> 0.20 \parallel < -$
0.20 (light grey lines). Red points indicate an association with T2D samples, and blue ones
with no-T2D samples. Enrichment genera categories are shown as insets in the main plot.
D, Area under the receiver operating characteristic (AUROC) curve resulting from the
sparse partial least square discriminant analysis (sPLS-DA) to determine the performance
of the DPP4-like proteins to classify T2D samples. Two models with two and three
components (informative genetic traits) support the higher discriminant power of DPP4-like
gene abundance from *Parabacteroides* and *Porphyromonas* species (on top) to discern T2D
samples. AUROC and p-values are shown as insets in respective subpanels.

Figure 2. Secretory signal detection in the pmDPP4 protein. MALDI-TOF mass
spectrometry monitoring of human GLP-1 incretin inactivation in enzymatic reactions
using cell-free supernatants from *E. coli* BL21 cells carrying the pET15b (**A**),
pET15b/pmDPP4 (**B**) and pET15b/pmDPP4Δ23 (**C**) constructs upon induction. Active
form of GLP-1 (MW 3,355) was detected in all cell-mediated hydrolysis assays and the
expected inactive form (MW 3,147) resulting from N-terminal dipeptide cleavage was only
evident in cells carrying pET15b/pmDPP4 plasmid encoding full-length protein. Anti-His
transmission electron microscopy immunogold staining of *E. coli* BL21 cells carrying the
pET15b (**D**), pET15b/pmDPP4 (**E**) and pET15b/pmDPP4Δ23 (**F**) constructs upon
induction. Red arrows indicate predominant distribution of 6×His-tagged proteins for E and
F panels. Scale bars at 100 nm (bars below TEM images). MW: monoisotopic mass.

Figure 3. *In vitro* function of the pmDPP4 protein. MALDI-TOF mass spectrometry
spectra of hydrolytic reactions performed in the presence of pmDPP4Δ23 or hsDPP4
proteins. Panels are distributed in terms of substrate used and the enzyme added (**A-D**). The
substrates used for reactions are depicted on top of every panel series with the respective
sequence information and the theoretical molecular weight (MW, monoisotopic mass).
Delta molecular mass values were calculated from the monoisotopic masses of products
observed in the control hsDPP4 reactions. The x-axis indicates the molecular mass values
in terms of m/z, and the y-axis indicates relative intensity (R.I.) of the most abundant
peptide species.

Figure 4. *In vivo* function of the pmDPP4 protein. Scheme of the animal study to test *in*
vivo activity of bacterial DPP4-like enzyme. **A**, Study design, mice administered with *E.*
coli BL21-DE3 cells carrying the pET15b empty vector or the pET15b/pmDPP4 vector
(n=8 each). **B, C** DPP4 activity measured in cecal content (B) and portal vein (C) of mice.
In the portal vein, active GLP-1, active GIP, total GLP-1, proglucagon, total GIP and PYY
are shown in panels **D**, **E**, **F**, **G**, **H** and **I**, respectively. Insulinemia and glycemia is shown
in panels **J** and **K**. Data were analyzed using a *t*-test and significance was set at p<0.05.

Figure 5. Dose-response of gliptins on DPP4 enzymes. The EC_{50} parameter was calculated

for hsDPP4 and pmDPP4Δ23 to determine the inhibitory efficacy of the drug on the

hydrolytic activity of the enzymes. Serial dilutions of FDA- or EMA-approved gliptins

from 1×10^{-6} to 1×10^{-9} M were established to measure the production of para-nitroaniline.

Top panel shows dose-response curves for hsDPP4, and bottom panel shows dose-response

curves for bacterial pmDPP4Δ23. Data obtained were computed *via* non-linear fitting

regression to estimate the EC_{50}. Data were derived from four independent replicates using

different protein purification batches.

Supplementary material S1. Comparison of DPP4-like function-associated peptidases.

Domain architecture, signal peptide presence and transmembrane domain predictions were

examined on the human DPP4 enzyme (SwissProt id P27487) (**A**), the consensus amino

acid profile of bacterial DPP4-like proteins (**B**), and the consensus amino acid profile of

PepX peptidases (**C**) using the respective web servers (top labels). Individual functional

domains according to SMART and/or Pfam annotation present in full-length query proteins

are presented in classical architecture diagrams. SignalP prediction for the N-terminal 70

amino acids shows C- (raw cleavage site score, red lines), S- (signal peptide score, green

lines), and Y- (combined cleavage site score, blue lines) scores to determine the individual

probability of amino acids to be part of a signal peptide driving protein secretion or

potential membership of a transmembrane structure. TMHMM prediction for the N-

terminal 70 amino acids shows the probability of individual amino acids to be part of alpha

helices embedded in biological membranes, thus indicating localization in the

transmembrane (red lines), inside of cell (blue line) or outside of cell (pink lines). The

amino acid logo representation of the multiple sequence alignment (MSA) of DPP4-like

proteins (bacteria, mouse, rat, and human) for the catalytic motif of S9 peptidase family is

drawn in **D**. Amino acid logo was created using the positions between amino acids 935 and

941 of the MSA using the SeqLogo sever (http://www.cbs.dtu.dk/biotools/Seq2Logo/) and

based on the P-Weighted Kullback-Leibler score. The amino acid positions for the human

protein are also shown at the bottom. This regions shows the MEROPS [23] catalytic motif

(GWSYGGY) of these serine proteases (S9).

Supplementary material S2. A, SDS-PAGE analysis of *E. coli* BL21-DE3 cell extracts
expressing recombinant pmDPP4 and pmDPP4Δ23 proteins. **B**, Ninety minute kinetics of
pmDPP4 activity (left y-axis, in terms of mU/mL units) and LDH activity (right y-axis, in
terms of 490 nm - 680 nm absorbance) mediated by cell-free supernatants from *E. coli*
BL21 cells carrying pET15b/pmDPP4 or pET15b/pmDPP4Δ23 constructs upon induction.
C, Gel filtration assay of an ionic-exchange purified fraction of the pmDPP4Δ23 protein,
revealing a dimeric nature in solution according to the calculated MW from the standards
elution profile. **D**, Purified pmDPP4Δ23 and human DPP4 proteins used in the *in vitro*
assays for human hormone hydrolysis and gliptin inhibition. Gels were stained with
Coomassie blue to visualize protein bands. The Dual Xtra pre-stained protein standard
(Bio-Rad, Hercules, CA) was used as a molecular size reference. LDH, lactate
dehydrogenase.

Supplementary material S3. Gliptin inhibition kinetics on pmDPP4Δ23. Lineweaver-
Burk double-reciprocal (1/V and 1/S) plots showing pmDPP4Δ23 (bottom panels) and
hsDPP4 (top panels) gpPNA hydrolysis in presence of vildagliptin, sitagliptin, saxagliptin,
and linagliptin. Inhibitor concentration is shown as line legend color from lower (0 nM,
light red) to higher (100 nM, burgundy lines) concentrations. The apparent K_m was
calculated respectively from inhibitor-associated data points using linear regression to find
intersection point at the y-axis (1/S). V, reaction velocity (nmol PNA min^{-1} pmole enzyme^{-1}
); S, substrate concentration (mM gpPNA).

Supplementary material S4. K_m fold-change on pmDPP4Δ23 and hsDPP4 enzymes upon
gliptin inhibition. Data derived from kinetics assays shown in Supplementary material S5
are shown in this spreadsheet. Headers indicate enzyme source, inhibitor tested, K_m
estimated, standard deviation (SD), fold-change, and p-value obtained from comparisons
with respective reactions with no inhibitor.

A — BL21 cells + pET15b

Active human GLP-1 (7-37)
HAEGTFTSDVSSYLE-
GQAAKEFIAWLVKGRG
(MW 3355)

3354.7858
3376.7731

B — BL21 cells + pET15b/pmDPP4

Inactive human GLP-1 (7-37)
EGTFTSDVSSYLE-
GQAAKEFIAWLVKGRG
(MW 3147)
Relative intensity = 12%

3354.7117
3377.6929

C — BL21 cells + pET15b/pmDPP4Δ23

3354.7549
3376.7438

D — 10 nm

E — 10 nm

F — 10 nm

TEM ImmunoGold staining of DPP4 (S9 family)

A

IC$_{50}$	Vildagliptin	Sitagliptin	Saxagliptin	Linagliptin
	1.545e-007	5.007e-008	4.288e-008	3.684e-009

B

IC$_{50}$	Vildagliptin	Sitagliptin	Saxagliptin	Linagliptin
	4.571e-007	9.043e-005	1.452e-007	1.484e-005

Table 1. Estimation of EC_{50} in a gliptins dose-response assay.

	hsDPP4			pmDPP4		
Gliptin	EC_{50}	CI	R^2	EC_{50}	IC	R^2
Sitagliptin	5.01×10^{-8}	4.76×10^{-8} to 5.27×10^{-8}	0.995	9.04×10^{-5}	1.35×10^{-5} to 1	ND
Vildaliptin	1.54×10^{-7}	1.41×10^{-7} to 1.70×10^{-7}	0.980	4.57×10^{-7}	3.68×10^{-7} to 5.72×10^{-7}	0.833
Saxagliptin	4.29×10^{-8}	4.14×10^{-8} to 4.44×10^{-8}	0.998	1.45×10^{-7}	1.30×10^{-7} to 1.69×10^{-7}	0.971
Linagliptin	3.68×10^{-9}	3.30×10^{-9} to 4.10×10^{-9}	0.982	1.48×10^{-5}	4.87×10^{-6} to 1	ND

EC_{50}, Estimated at molar (M) concentration. CI, confidence interval at 95%. R^2, calculated as the fit of data into the non-linear function of the inhibitor concentration vs normalized response. Data obtained from independent experiments (N = 2). ND, data did not fit as non-linear function.