PUPpy: a primer design pipeline for substrain-level microbial detection and absolute quantification.

Hans Ghezzi[1], Michelle Y. Fan[2], Katharine M. Ng[3], Juan C. Burckhardt[3], Deanna M. Pepin[3], Xuan Lin[4], Ryan M. Ziels[4], Carolina Tropini[1,3,5,6]*

[1]Department of Bioinformatics, University of British Columbia, Vancouver, Canada

[2]Department of Cellular & Physiological Sciences, University of British Columbia, Vancouver, Canada

[3]Department of Microbiology & Immunology, University of British Columbia, Vancouver, Canada

[4]Civil Engineering, The University of British Columbia, Vancouver, BC, Canada

[5]School of Biomedical Engineering, University of British Columbia, Vancouver, Canada

[6]Humans and the Microbiome Program, Canadian Institute for Advanced Research (CIFAR), Toronto, Canada

*Corresponding Author: Carolina Tropini
Email: carolina.tropini@ubc.ca

Abstract

Characterizing microbial communities at high-resolution is crucial to unravel the complexity and diversity of microbial ecosystems. Advances in bulk sequencing assays such as 16S rRNA and shotgun sequencing have enabled unparalleled qualitative and quantitative microbiota investigations. However, these methods generally do not provide accurate resolution beyond the genus or species level and lack insights into absolute microbial abundance. Here, we introduce Phylogenetically Unique Primers in python (PUPpy), a fully automated pipeline to design microbe- and group-specific primers within a given microbial community. PUPpy-designed primers detect individual microbes and quantify absolute microbial abundance in a defined community below the species level, requiring only coding sequence files of the community members as input. We experimentally evaluated the performance of PUPpy-designed primers using two bacterial communities as benchmarks. Each community was comprised of 10 members, exhibiting a range of genetic similarities that spanned from different phyla to substrains. PUPpy-designed primers also enabled the detection of groups of bacteria in an undefined community, such as the detection of a gut bacterial family in a complex stool microbiota sample. Taxon-specific primers designed with PUPpy showed 100% specificity to their intended targets, without unintended amplification, independently of community composition and complexity. Lastly, we show absolute quantification of microbial abundance using PUPpy-designed primers in droplet digital PCR (ddPCR), benchmarked against 16S rRNA and shotgun sequencing. Our data shows that PUPpy-designed microbe-specific primers can be used to quantify substrain-level absolute counts, providing more resolved and accurate quantification in defined communities than 16S rRNA and shotgun sequencing.

Importance

Profiling microbial communities at high resolution and with absolute quantification is essential to uncover hidden ecological interactions within microbial ecosystems. Nevertheless, achieving resolved and quantitative investigations has been elusive due to methodological limitations in distinguishing and quantifying highly related microbes. Here, we describe PUPpy, an automated computational pipeline to design taxon-specific primers within defined microbial communities. Taxon-specific primers can be used to selectively detect and quantify individual microbes and larger taxa within a microbial community. PUPpy achieves substrain-level specificity without the need for computationally intensive databases and prioritises user-friendliness by enabling both terminal and graphical user interface (GUI) applications. Altogether, PUPpy enables fast, inexpensive, and highly accurate perspectives into microbial ecosystems, supporting the characterization of bacterial communities in both *in vitro* and complex microbiota settings.

Introduction

Microbial communities play a crucial role in numerous aspects of life on Earth, including nutrient cycling, disease development, and carbon fixation (1, 2). These processes are driven by a remarkable genetic and functional diversity spanning various taxonomic levels, from unrelated phyla to closely related species. Although functional diversity is greatest among unrelated microbes, it is also present in microorganisms with closely related genetic makeups. Importantly, subtle genetic variations at the strain level can also lead to extensive genomic variations (3, 4), such as changes in microbial metabolism (5–7), immunomodulation (8, 9), antibiotic resistance (10, 11), biofilm formation (12), and pathogenesis (13). For example, the impacts of different *Escherichia coli* strains on human health can vary significantly: Nissle 1917 (14) is commensal, while enterohemorrhagic *E. coli* (EHEC) O157:H7 and enteroaggregative *E. coli* (EAEC) O104:H4 (15) are pathogenic. Thus, profiling microbial communities with at least strain-level granularity is imperative to uncover hidden ecological interactions that may otherwise be missed at lower taxonomic resolution. However, detecting and quantifying closely related microbes remains a challenging task due to their largely indistinguishable genetic makeup, limiting microbial profiling to lower taxonomic levels.

Over the last decade, advances in both culture-dependent and independent approaches have enabled the high-throughput characterization of microbial communities at increasingly high resolution. In the case of culture-dependent methods, while significant progress in growing bacteria in controlled environments has enabled the identification of bacterial strains and species from mixed communities, the work-intensive nature of these methods has limited their widespread application (16, 17). On the other side, culture-independent methods, specifically amplicon (e.g., 16S rRNA) and shotgun sequencing have enabled increasingly cost-effective and high-throughput detection and quantification of microbes, including rare and unculturable ones (18). Although sequencing methods have been used as the gold standard for microbial measurements, they suffer from considerable limitations. Specifically, 16S rRNA sequencing usually cannot resolve taxonomies beyond the genus level, does not directly provide information about microbial functional potential, is sensitive to PCR biases caused by primer choice and variations in copy numbers across organisms, and depends on databases to accurately assign taxonomy (18, 19). While providing functional information and greater resolution, shotgun sequencing also presents several limitations, including difficulty in resolving strains of the same species due to high genetic similarity, biases introduced during sample preparation and sequencing, and the requirement for considerable computational resources and complex bioinformatic pipelines (18, 20). In addition, both 16S rRNA and shotgun sequencing can at best estimate microbial relative abundance in a community, as opposed to absolute counts.

Another method for detecting microbes beyond the species level and quantifying absolute counts is by selectively amplifying microbe-specific genes. This can be achieved through widely available polymerase chain reaction (PCR) techniques, including quantitative PCR (qPCR) and droplet digital PCR (ddPCR) (21, 22). Unlike sequencing, PCR methods are widely accessible, time efficient, and cost-effective, empowering researchers to achieve highly resolved and quantifiable microbial investigations without considerable computational and monetary resources. Nevertheless, designing primers to selectively amplify specific target organisms at any taxonomic resolution (e.g., genus, species, or strain level) remains a major challenge that has hindered the high throughput use of PCR assays in microbial investigations. This becomes especially challenging when surveying complex and undefined microbial communities (e.g., the fecal microbiome). In these communities, the genetic makeup of each microbe is often not known *a priori,* making it impossible to check for alternative primer binding sites. Currently, several tools exist to design primers specific to input genomic sequences, including SpeciesPrimer (23), find_differential_primers (fdp) (24), RUCS (25), and TOPSI (26). However, these tools have either been discontinued, require significant manual handling to generate configuration files, or cannot flexibly

110   design primers for any user-defined combination of microbes in a community, highlighting the need for a
111   high-throughput, user-friendly, and highly selective taxon-specific primer design tool.
112
Here, we introduce Phylogenetically Unique Primers in python (PUPpy), a fully automated pipeline
to design primers targeting individual microbes or groups of microbes within a given microbial
community. PUPpy only requires coding sequences (CDSs) of the microbes of interest as input to design
primers with substrain level specificity. In addition, PUPpy achieves exceptional specificity without the
need for external databases, meaning that it can also be run under low computational resources. We
benchmarked PUPpy-designed primers against two defined bacterial communities, each consisting of 10
members with varying degrees of genetic similarity, ranging from distinct phyla to substrains. We also
evaluated the ability of PUPpy-designed primers to detect members of a highly diverse and fastidious gut
bacterial family, the Muribaculaceae, in a complex and undefined conventional mouse microbiota. We
show that all taxon-specific primers designed with PUPpy selectively amplified the respective targets,
independently of community composition and complexity. Finally, we assessed the effectiveness of
PUPpy-designed primers for accurately quantifying the total microbial count in specific communities.
This assessment was conducted using ddPCR and compared with results obtained from 16S rRNA and
shotgun sequencing methods. Our data suggests that taxon-specific primers enable more resolved and
accurate quantification than short-read 16S rRNA and shotgun sequencing. Altogether, PUPpy represents
a versatile tool to quantify absolute microbial counts of individual strains, an essential parameter that has
been missing in microbiome studies. The versatility of PUPpy makes it suitable for a variety of
applications involving both defined and complex communities, ranging from environmental studies to
host-associated microbiota research.
132
133   **Results**
134
135   **Overview of the PUPpy pipeline**
136
PUPpy is a fully automated computational pipeline that allows the design of taxon-specific PCR primers
targeting user-defined microbial communities (Fig. 1A). This pipeline defines two primary applications of
taxon-specific primer design: microbe- and group-specific (Fig. 1B). Microbe-specific primers target
unique CDSs only present in a single member of the community. These primers allow users to target
individual microbes with full specificity, a crucial aspect in tracking and quantifying specific microbes in
communities with high degrees of genetic similarity. Conversely, group-specific primers are designed to
select genes that are shared across all user-defined targets (Fig. 1B). Importantly, these primers enable the
selective assessment of microbial dynamics of an entire taxon (e.g., all *Bacteroides*) in a sample. In both
microbe and group-specific applications, PUPpy designs taxon-specific primers by aligning every CDS in
the community and identifying unique (microbe-specific) or shared (group-specific) genes. Because
PUPpy only requires CDS files as input, it can run locally and does not require memory intensive
databases (e.g., the NCBI nr database). More details on the computational workflow are provided in the
Methods section. To demonstrate the applicability of PUPpy in microbiome research, we empirically
validated taxon-specific primers in 3 increasingly complex microbial communities. These communities
were composed of 1) a defined community of 10 phylogenetically distinct bacteria, 2) a defined
community of 10 bacteria from 3 increasingly related taxa, and 3) a complex microbiota from
conventional mouse fecal samples. By gradually increasing the genetic similarity and complexity of the
community, we were able to assess the resolution limits and effectiveness of the pipeline under diverse
microbial community scenarios.
156
157   **Microbe-specific primers show specificity to intended targets in a genetically diverse community**
158
The first community, hereafter referred to as the "GUT community", was composed of 10
phylogenetically distinct bacteria belonging to 9 families and 5 phyla (Bacillota, Bacteroidota,

Pseudomonadota, Actinomycetota, Verrucomicrobiota) (Table S1). These bacteria were chosen as being
phylogenetically distinct, common gut commensals, and representative of the 5 most abundant phyla of
the human gut microbiota. This community, while simple, accurately represents a defined microbial
consortium similar to that represented in several gnotobiotic animal models (27–29). Notably, each
member of this community possesses a significant number of unique genes and has been previously
characterized (27, 30, 31). To confirm the genetic diversity within the GUT community, we calculated
Average Nucleotide Identity (ANI) on the genomic assemblies for the 10 bacteria (see Methods). The
community included two *Bacteroides* species, *Bacteroides ovatus* and *Bacteroides thetaiotaomicron*,
which, as expected, shared a greater degree of genetic similarity, with 80% percentage identity and ~50%
percentage coverage (Fig. 2A). All other GUT members displayed greater genetic diversity, with <75%
percentage identity and <8.5% percentage coverage (Fig. 2A). Following ANI analysis, we ran PUPpy
with default parameters and generated microbe-specific primers for all 10 GUT community members (see
Table S2). Consistent with the degree of genetic diversity observed in the ANI analysis, PUPpy identified
fewer unique CDSs in more genetically similar organisms. PUPpy orders primer targets based on
descending number of unique genes found, enabling an immediate visual and quantitative evaluation of
community diversity (Fig. 2B). Specifically, compared to all other community members, PUPpy
identified ~27% of the CDSs as unique in *B. ovatus* and ~26% in *B. thetaiotaomicron,* the most related
microbes, as opposed to ~74% in *E. coli* BW25113 (Fig. 2B). Importantly, PUPpy considers all unique
CDSs to design microbe-specific primers, and multiple primers can be designed on the same CDS. Thus,
even in the organism with the fewest unique genes, *B. thetaiotaomicron*, PUPpy evaluated 1248 options
(i.e., ~26%) for primer design (Fig. 2B). Next, we empirically validated the specificity of the GUT
microbe-specific primers by PCR and gel electrophoresis using gDNA isolated from the 10 GUT
community members (see Methods). Each primer pair was tested against i) the respective gDNA positive
control, ii) the 10-member gDNA pool, iii) a 9-member gDNA pool including all taxa except the primer
target, and iv) a no-template control (Fig. 2C). Following this experimental validation, we found that all
microbe-specific primers were specific to the respective primer target, without unintended amplification
(Fig. 2D, S1). PUPpy can therefore be leveraged to design microbe-specific primers and selectively detect
individual microbes in a genetically diverse and defined microbial community.
189
190   **Taxon-specific primers selectively detect microbes down to the substrain level**
191
As microbial communities often harbor multiple species and strains within the same genus, we next asked
whether PUPpy could be used to design taxon-specific primers and detect highly related microbes. To test
this, we created a second 10-member bacterial community, hereafter referred to as the "Species-Strain-
Substrain (SSS) community" (Table S1). This community consisted of 3 *Enterocloster* species, 3 *B.*
thetaiotaomicron strains and 4 *E. coli* K-12 substrains. These bacteria were chosen to assess the
taxonomic resolution achievable by PUPpy, while also maintaining a degree of phylogenetic diversity
across taxa. To evaluate the genetic similarity in the community we performed ANI analysis, which
grouped the SSS community members into 3 distinct memberships consistent with their assigned
taxonomic classifications (Fig. 3A). As expected, the *Enterocloster* species displayed the lowest
similarity, with a minimum of ~78% identity and ~34% coverage. Conversely, the closely related *E. coli*
substrains were more genetically similar, sharing >99.95% identity and ~99% coverage (Fig. 3A). Next,
we ran PUPpy with default parameters to design both microbe-specific primers for each member and
group-specific primers for each taxon (Table S2). Group-specific primers are designed on genes shared
across all primer targets (e.g., all *Enterocloster* species) but missing in all unintended targets (e.g., all *B.*
thetaiotaomicron strains and *E. coli* substrains), enabling the selective detection of microbial taxa within a
microbial community. Consistent with the ANI analysis and taxonomic classification, PUPpy ordered the
10 members into 3 groups based on the number of unique CDSs found (Fig. 3B). Specifically, PUPpy
identified between ~19% and ~36% unique CDSs for the 3 *Enterocloster* species, between ~7% and
~17% for the *B. thetaiotaomicron* strains, and between 0% and ~1% for the *E. coli* K-12 substrains (Fig.
3B). Using default parameters, PUPpy did not identify any unique CDSs for *E. coli* MC4100 within the

SSS community, and thus no microbe-specific primers were designed or tested for this organism (Fig.
3B). The lack of unique genes is likely due the close genetic similarity to all 3 other *E. coli* members.
Increasing the percentage identity threshold in `puppy-align` may yield unique genes for *E. coli* MC4100
by decreasing the alignment stringency (see Methods for details). We validated the specificity of SSS
microbe-specific primers following the same experimental design as the GUT community (Fig. 3C). In
addition, we tested the specificity of group-specific primers by running each pair against i) the respective
primer gDNA targets, (condition (ii) was not run as it would provide redundant information), iii) the 9-
member SSS gDNA pool including all taxa except the target strain gDNAs, and iv) a no-template control
(Fig. 3C). All microbe- and group-specific primers tested showed specificity to their intended targets,
independent of taxonomic level and genetic similarity (Fig. 3C, 3D, S2). These data show that PUPpy
enables the selective detection of microbes down to the substrain level by identifying and designing PCR
primers for unique CDSs. In addition, this validation highlights the ability of PUPpy to identify conserved
CDSs across specific groups of bacteria, allowing the selective amplification of multiple organisms with a
single primer pair.
226
227 **Taxon-specific primers selectively amplify target microbes independently of community**
228 **composition and complexity**
229
Having validated the specificity of PUPpy-designed primers in defined microbial communities, we asked
whether the pipeline could also be applied to complex communities in which the exact membership is not
known. To investigate whether we could selectively identify the presence of specific taxa in this setting,
we leveraged a conventional mouse gut microbiota. The mouse microbiota has high abundance of the
family Muribaculaceae, a highly diverse and hard-to-culture bacterial family that is broadly prevalent in
warm-blooded animals (32–34). We have previously shown that mice treated with the osmotic laxative
polyethylene glycol (PEG) are depleted of Muribaculaceae (35) and we wanted to ascertain whether
group-specific primers to this family could identify the loss of this family without the need for
sequencing. We extracted gDNA from fecal samples of mice that were either untreated (Muribaculaceae-
positive) or PEG-treated (Muribaculaceae-depleted) (see Methods). We confirmed Muribaculaceae
presence or absence by performing 16S rRNA sequencing on the fecal gDNA samples, and then evaluated
PUPpy-designed primers against the Muribaculaceae family. Such experimental validation presents two
main challenges in the design of specific primers: 1) the overall composition of the community was
treated as unknown *a priori* and therefore we did not have a complete list of CDS files to input into
PUPpy and 2) we did not know which Muribaculaceae strains were present in the community of untreated
mice, and thus for which microbes to design primers. Therefore, to minimize the chances of off-target
amplification we selected a comprehensive list of 156 unintended targets and 11 intended Muribaculaceae
targets (Table S1). The unintended targets were selected among prevalent gut commensal bacteria to
represent members of the phyla Bacillota, Bacteroidota, Pseudomonadota, Actinomycetota,
Verrucomicrobiota, and Fusobacteriota. Using this 167-member input community we were able to
identify a gene shared across all Muribaculaceae but absent in all other 156 members. Importantly, since
Muribaculaceae is a highly diverse family (33), we designed 8 primer pairs to account for higher potential
representation of different sub-taxa in the complex community and pooled them to create a
Muribaculaceae-specific primer cocktail. We validated this mix against both Muribaculaceae-positive and
Muribaculaceae-depleted fecal samples and evaluated its specificity (Fig. 3E) (see Methods). Our results
show that Muribaculaceae-specific primers selectively detected Muribaculaceae in the positive samples
but did not in the depleted samples (Fig. S3), consistent with the results from 16S sequencing (Fig. 3F). In
addition to being specific, these Muribaculaceae-specific primers did not display any off-target
amplification of PCR amplicons with different sizes in the untreated samples (see Fig. S3). This is crucial
in PCR assays such as qPCR and ddPCR, where quantification may be biased by unintended
amplification. Altogether, we define a methodology involving the use of PUPpy to design taxon-specific
primers that can selectively amplify microbial targets even in a complex community.
262

263
264 **Taxon-specific primers enable high resolution absolute microbial quantification**
265
Beyond the detection of presence and absence of microbes, taxon-specific primers can also be used to
quantify absolute microbial counts at extremely high-resolution using qPCR or ddPCR (21, 23). To
benchmark microbial quantification with PUPpy, we compared quantification of PUPpy-designed primers
via ddPCR compared to the gold standard methods 16S rRNA and shotgun sequencing. We pooled 10
extracted gDNA samples for each member of the GUT and SSS communities, respectively, in a
theoretical 1:1 genomic copy ratio calculated from the respective microbial gDNA concentration and
genome size (see Methods). We then aliquoted the gDNA pools for analysis with 16S rRNA sequencing,
shotgun sequencing, and ddPCR with PUPpy primers (Fig. 4A) (see Methods). Using ddPCR we
calculated absolute microbial abundance and compared it to the relative abundance estimated by the
sequencing methods. In the GUT community, ddPCR and shotgun sequencing achieved similar
resolution, which, as expected, was markedly higher than in 16S rRNA sequencing due to the limited
variability in the V3-V4 regions targeted in this technique (Fig. 4B). Specifically, amplicon sequencing
classified only 4 members at the species level (*Akkermansia muciniphila, Eubacterium rectale*, *B. ovatus*
and *B. thetaiotaomicron*) and 6 members at the genus level (Fig. 4B). Conversely, shotgun sequencing
identified all 10 members at the strain level. This resolution level was enabled by the custom reference
database used in the analysis, which only included the GUT reference genomes (see Methods). This level
of resolution in shotgun sequencing would be unlikely using a comprehensive reference database. Similar
to shotgun sequencing, ddPCR using microbe-specific primers was also able to achieve strain level
resolution (Fig. 4B). However, unlike the sequencing methods, ddPCR provided faster and more
quantitative assessment of bacterial composition by measuring absolute microbial counts, which is
inherently more accurate than relative abundance.
287
Among the 3 methods, ddPCR also yielded the most resolved and accurate quantification in the
genetically related SSS members (Fig. 4C). ddPCR selectively detected and quantified absolute counts for
all SSS community members, except *E. coli* K-12 MC4100, for which we did not design primers given its
similarity within the community. Similar to the results for the GUT community, 16S rRNA sequencing
achieved the lowest resolution, only detecting the 3 major taxa and no individual strains (Fig. 4C).
Finally, shotgun sequencing also achieved high resolution and detected all 10 microbes, however, it was
unable to assign reads to each strain unambiguously. Specifically, in short-read shotgun sequencing, reads
that map equally well to several potential targets in the reference database are commonly assigned
randomly across multi-mapping microbes (36). This approach becomes problematic in accurately
quantifying closely related microbes, which share a substantial part of the genomic DNA. In the SSS
community, quantification of strains and substrains with the Bowtie2 (37) pipeline closely reflected the
theoretical input of 1:1 genomic copy ratio. However, we found this was an artefact of randomly
assigning multimapping reads, which coincidentally matched the expected input and would not extend to
scenarios where strains and substrains are present in uneven proportions. To further investigate this effect
for communities of arbitrary microbial concentrations, we generated *in silico* reads for the SSS members
at known input proportions and analysed these synthetic reads using both the heuristic alignment
approach with Bowtie2 (37) used in Fig. 4, and Kraken2 (38), a Lowest Common Ancestor (LCA)
approach (see Methods for details). Unlike the heuristic alignment approach, the LCA approach does not
attempt to randomly assign ambiguous reads to multimapping targets, and instead assigns them to the
LCA. Both methods accurately quantified the 3 *Enterocloster* species, supporting the use of shotgun
sequencing when microbes are sufficiently distinct for reads to be confidently assigned (Fig. 5A and 5B).
However, quantification accuracy in both methods gradually decreased as genetic similarity increased.
Notably, the observed abundance obtained with the LCA approach for all strains and substrains was
considerably lower than with Bowtie2, consistent with the random distribution of ambiguous reads by the
latter method (Fig. 5A and 5C). To further support this, we found that quantification with Kraken2 was
considerably more accurate at the LCA level as opposed to the microbe level for both strains and

substrains (Fig. 5C and 5D). Specifically, the ratio of observed vs expected relative abundance was ~0.80
for the *E. coli* LCA (Fig. 5C) while it dropped to between ~0.001 and ~0.05 for individual *E. coli* strains
(Fig. 5D). These data suggest that most reads cannot be confidently mapped to closely related strains and
can only be assigned to LCAs. These findings also confirm that shotgun sequencing was not actually able
to accurately quantify individual strains and substrains in the SSS community but instead yielded a
uniform abundance by randomly distributing ambiguous reads (Fig. 4C). Altogether, these results shine
light on the pitfalls of using short-read sequencing for high-resolution profiling and highlight the potential
of PUPpy taxon-specific primers to accurately detect and quantify microbes in complex communities.
322
323   **Discussion**
324
In this study, we introduced PUPpy, a new computational pipeline to design taxon-specific primers in
microbial communities (Fig. 1). We benchmarked applications of PUPpy in microbiome research to
detect and quantify absolute bacterial counts at high resolution in both defined and complex communities
(Fig. 2-4). Our results show that PUPpy-designed taxon-specific primers can detect microbes down to the
substrain level in defined bacterial communities (Fig. 2, S1, 3, and S2). Importantly, we also used PUPpy-
designed primers to selectively target taxa in complex communities, even when the full membership was
unknown (Fig. 3 and S3). Finally, our data suggest that microbe-specific primers can be used in ddPCR to
quantify absolute microbial counts, achieving greater accuracy and resolution than short-read 16S rRNA
and shotgun sequencing in defined bacterial communities (Fig. 4 and 5).
334
Unlike sequencing methods, using PUPpy enables microbes to be identified at the substrain level
rapidly and without specialised sequencing equipment, as it only requires PCR and gel electrophoresis
instrumentation that are widely available. PCR assays not only enable substrain level resolution but also
absolute microbial quantification when qPCR or ddPCR are used, further improving its scope of use. As a
software package, PUPpy was specifically developed to prioritize user-friendliness, with minimal manual
handling and command-line knowledge required. By not requiring configuration files (which are needed
for other primer design tools such as fdp (24)), PUPpy shortens the overall execution time and reduces the
chances of human error, while remaining customizable. In addition, unlike the primer design tools fdp
(24), TOPSI (26), and SpeciesPrimer (23), PUPpy does not use large databases for specificity checks,
which are time and memory-expensive tasks. Instead, PUPpy leverages an alignment-based strategy with
MMseqs2 (39), which enables both rapid homology search across all genes in the community and low
computational resources, making the pipeline suitable for both computing clusters and personal
computers. In the current study, we successfully designed taxon-specific primers on both a computing
cluster and personal computer in 2-10 minutes for each 10-member community and under 1 hour for a
167-member complex community with the cluster (see Methods for details). The increased time in more
complex communities reflects the fact that identifying unique genes, and thus designing microbe-specific
primers, is more computationally intensive in larger communities. Furthermore, designing microbe-
specific primers is more challenging in communities containing genetically related microbes, such as
multiple substrains. In these communities, homology searches may not identify any genes sufficiently
different to be binned as unique, for instance in the case of *E. coli* MC4100 in the SSS community (Fig.
3B). Here, finding coding sequences with the lowest possible sequence identity (and hence greatest
dissimilarity), is essential to design microbe-specific primers. PUPpy enables users to lower the alignment
stringency and thus increase the alignment identity required for genes not to be considered unique. For
example, increasing the alignment identity from the default 30% to 50% ensures that any gene aligning
with up to 50% identity is considered unique. Although decreasing the alignment stringency increases the
chances of finding unique genes, it also increases the risk of false positives. This parameter should thus be
manipulated with caution and only if necessary, such as in communities containing, or expected to
contain, multiple genetically related microbes. Importantly experimental validation is essential and
always recommended to ensure proper function of the primers.
364

Beyond identification of bacteria below the species level, we successfully detected a bacterial
family with high specificity within a complex microbiota, even without *a priori* knowledge of the
microbial membership. In general, designing selective primers for an undefined microbiota represents a
challenging task due to the unknown gene pool, limiting the applicability of *in silico* specificity checks
through both large databases (e.g., BLAST database) and the input community. Here, we designed a
Muribaculaceae-specific primer cocktail for a conventional mouse gut microbiome by using a
comprehensive list of 156 unintended targets and 11 intended Muribaculaceae targets. Primer mixes may
be beneficial for genetically diverse targets, such as Muribaculaceae members, as even few single
nucleotide variations may affect primer affinity and prevent amplification. By providing an extensive
input community of 167 members, PUPpy identified highly conserved genes among the target microbes,
decreasing the chances of false positive hits even in a complex community. Such conserved genes are
likely to be exclusively found in the target members or taxon, thus remaining highly specific
independently of the input community complexity. Therefore, the alignment strategy leveraged by PUPpy
may circumvent the need for specificity checks with large databases, while requiring lower computational
resources and achieving specificity in undefined microbiomes.
380
As research continues to uncover remarkable functional diversity among closely related microbes,
tools to achieve strain level microbial profiling have become necessary. In the last decades, short-read
sequencing has been the primary approach to characterize communities, enabling unparalleled high-
throughput microbial profiling (18). Nevertheless, short-read metagenomics can only achieve genus or
species-level resolution due to the limits imposed by sequencing read length (Fig. 4B and 4C). Alternative
approaches, specifically long-read sequencing, have been gaining traction by offering numerous
advantages, including greater resolution, better assembly of *de novo* genomes, and better detection of
structural variants (40–42). Despite these promising benefits, as well as advances in the use of reference
standards (43), and bias correction (44, 45), microbial profiling by long-read sequencing remains
restricted to compositional data as opposed to the preferred absolute quantification. PUPpy represents a
faster, cheaper, and more accessible solution to achieve both substrain level resolution and absolute
microbial quantification. While sequencing remains invaluable in large scale and exploratory
investigations, PUPpy may support highly quantitative investigations, especially in the context of
gnotobiotic studies.
395
396   In summary, PUPpy designs taxon-specific primers in diverse microbial communities and
397   represents a fast, user-friendly, and widely accessible solution to profiling the microbiota. As microbiome
398   research continues to uncover functional diversity beyond the species-level, the ability to detect microbes
399   quickly and accurately below the species level will support high-resolution characterizations of microbial
400   ecosystems that are required for the field to advance.
401
402
403

404
405 **MATERIALS AND METHODS**

406 **Resource Table:**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms		
Swiss-Webster Mice - Germ Free	Taconic, in-house colony	SW GF
Chemicals		
Chopped Meat Medium	Anaerobe Systems	AS-811
BD BBL™ Dehydrated Culture Media: Brain Heart Infusion	Fisher Scientific	B211059
Reinforced Clostridial Medium	Oxoid™	CM0149B
Mucin from porcine stomach (Type III)	Sigma-Aldrich	M1778-100G
L-cysteine	Thermo Scientific™	A1043518
Hemin chloride	MilliporeSigma™	37415GM
Vitamin K_1	Alfa Aesar	L10575
Agar	Fisher BioReagents™	BP1423500
Critical Commercial Assays		
DNeasy Blood & Tissue Kit	QIAGEN	69504
DNeasy 96 PowerSoil Pro QIAcube HT Kit	QIAGEN	47021
Quant-iT™ 1X dsDNA HS Assay	Invitrogen	Q33232
Nextera XT DNA Library Preparation Kit	Illumina	FC-131-1024
16S rRNA Library Preparation	Sequencing and Bioinformatics Consortium (SBC)	NA
2X DreamTaq Green PCR Master Mix	Fisher Bioreagents	K1082
Agarose	Fisher Bioreagents	BP160-500

ddPCR^{TM} 96-Well Plates	Bio-Rad	12001925
QX200^{TM} ddPCR^{TM} EvaGreen Supermix	Bio-Rad	1864034
DG32 Automated Droplet Generator Cartridges	Bio-Rad	1864108
Pipet Tips for AutoDG^{TM} System	Bio-Rad	1864120
ddPCR^{TM} Droplet Reader Oil	Bio-Rad	1863004
Automated Droplet Generation Oil for EvaGreen	Bio-Rad	1864112
PCR Plate Heat Seal, foil, pierceable	Bio-Rad	1814040
SYBR^{TM} Safe DNA Gel Stain	Invitrogen	S33102
Software and Algorithms		
MMseqs2	https://github.com/soedinglab/MMseqs2	V14.7e284 (39)
R software	https://www.r-project.org/	Version 4.1.2 (46)
ggplot2	https://www.tidyverse.org/	Version 3.3.5 (47)
tidyverse	https://www.tidyverse.org/	Version 1.3.1 (48)
pyani	https://github.com/widdowquinn/pyani	Version 0.2.9 (49)
FastQC	https://www.bioinformatics.babraham.ac.uk/projects/fastqc/	Version 0.11.9 (50)
MultiQC	https://github.com/ewels/MultiQC	Version 1.10.1 (51)
fastp	https://github.com/OpenGene/fastp	Version 0.23.2 (52)
Bowtie2	https://github.com/BenLangmead/bowtie2	Version 2.4.2 (37)
CoverM	https://github.com/wwood/CoverM	Version 0.6.1 (53)
Kraken2	https://github.com/DerrickWood/kraken2	Version 2.1.2 (38)
InSilicoSeq	https://github.com/HadrienG/InSilicoSeq	Version 1.5.4 (54)
primer3-py	https://github.com/libnano/p	Version 0.6.1 (55)

	rimer3-py	
pandas	https://github.com/pandas-dev/pandas	Version 1.5 (56)
biopython	https://github.com/biopython/biopython	Version 1.80 (57)
dask	https://github.com/dask/dask	Version 0.15.2 (58)
seaborn	https://github.com/mwaskom/seaborn	Version 0.13.0 (59)
matplotlib	https://github.com/matplotlib/matplotlib	Version 3.8.2 (60)
colorama	https://github.com/tartley/colorama	Version 0.4.1 (61)
QIIME2	https://qiime2.org/	Version 2023.2 (62)
DADA2	https://github.com/benjjneb/dada2	Version 1.20 (63)
Guppy	https://nanoporetech.com/	Version 4.5.3 (64)
Bonito	https://github.com/nanoporetech/bonito	Version 3.1 (65)
Flye	https://github.com/fenderglass/Flye	Version 2.7 (66)
Racon	https://github.com/isovic/racon	Version 1.4.22 (67)
Medaka	https://github.com/nanoporetech/medaka	Version 1.4.3 (68)
CheckM	https://github.com/Ecogenomics/CheckM	Version 1.0.18 (69)
Prokka	https://github.com/tseemann/prokka	Version 1.14.5 (70)
SILVA database	https://www.arb-silva.de/documentation/release-1381/	Version 138.1 (71)
QX Manager 2	https://www.bio-rad.com/en-ca/life-science/digital-pcr/qx-	Version 2.0 (Bio-Rad)

	software	
Other		
Standard diet 5K67 - JL Rat & Mouse/Auto 6F autoclavable diet	LabDiet https://www.labdiet.com/product/detail/5k67-labdiet-jl-rat-and-mouse-auto-6f	5K67
Automated Droplet Generator	Bio-Rad	1864101
PX1™ PCR Plate Sealer	Bio-Rad	1814000
C1000 Touch™ Thermal Cycler	Bio-Rad	1851196
QX200™ Droplet Reader	Bio-Rad	1864003
Deposited data		
https://borealisdata.ca/privateurl.xhtml?token=37d3e60b-62ca-40d8-9696-f77f9d7ccb57		

408
409
410 **Ethics statement:**
411
412 All animal experiments were performed according to protocol number (A19-0078) approved by the
413 Animal Care Committee at the University of British Columbia and in direct accordance with the
414 guidelines of the Canadian Council of Animal Care (CCAC).
415

416 **EXPERIMENTAL MODELS**
417
Bacterial strains and culture conditions
Metadata regarding all bacteria used in this study, including taxonomy, sources, and individual culture
conditions can be found in Table S1. All bacteria used in this study were cultured anaerobically with the
following atmosphere: 5% H_2, 5% CO_2, and 90% N_2 (Linde Canada, Delta, BC, Canada) at 37°C. All
media were pre-reduced in an anaerobic chamber (Coy Laboratories, Grass Lake, MI, USA) for at least
24 hours prior to use. Media names, abbreviations, ingredients, and recipes can be found in Table S3. All
microbes were streaked onto their respective media plates from glycerol stocks stored at -80°C, followed
by inoculation of single colonies into 3 mL of liquid media. Culture conditions and incubation times for
individual microbes can be found in Table S1.

PUPpy Pipeline Implementation
Technical specifications
All analyses involving PUPpy were performed on UBC ARC Sockeye, a high-performance computing
platform, running an Intel Xeon Silver 4110 (2.1GHz) processor with 16 cores, 192GB DDR4-2666 ECC
RAM and 2TB SATA hard disk drive. PUPpy workflows were replicated on a consumer laptop with
MacOS Ventura operating system, 10-core CPU with 8 performance cores, 16-core GPU, 16GB RAM
and 512 GB solid-state drive to validate the pipeline's viability under lower computational resources.

Executing the PUPpy pipeline
The PUPpy pipeline can be executed from terminal by first running `puppy-align`, followed by the
`puppy-primers` script. Alternatively, it is possible to execute both commands through a GUI by running
`puppy-GUI` and interacting with a button-based system. See below and the PUPpy GitHub repository
(https://github.com/Tropini-lab/PUPpy) for more detailed instructions and execution guidelines.

Sequence alignment of input coding sequence files
The first command of PUPpy, `puppy-align`, performs many-against-many local pairwise sequence
alignment with MMseqs2 (39) of all user-provided CDSs in a microbial community. This script requires a
directory containing the CDS files of the targets for which primers should be designed. Optionally, users
can provide a directory with the CDS files of non-targets, which will be exclusively used for specificity
checks of the intended targets. Importantly, all CDS files provided as input in the `puppy-align` script will
be aligned to ensure specificity of all taxon-specific primers. The actual design of primers occurs in the
`puppy-primers` scripts, thus the CDS file choice in `puppy-align` is exclusively meant to indicate which
microbes are present in the defined community of interest. CDS files provided by users must meet the
following criteria: 1) FASTA headers must be provided in Prokka, RAST or NCBI formats, 2) filenames
must contain the string "cds" prior to the extension, and 3) filenames must end with the file extension
".fna". The command `puppy-align` prepends microbe identifiers prior to the string "_cds" in the CDS
filenames to all FASTA headers, which is required for downstream parsing. For improved readability,
users are encouraged to name CDS files with easily interpretable and unique microbe identifiers prior to
the string "_cds" in the filename. Examples of acceptable CDS filenames include
"B_theta_VPI5482_cds.fna" and "B_thetaiotaomicron_VPI_5482_cds_from_genomic.fna". Following
file renaming, PUPpy concatenates all input CDS files (from the primer target and non-target directories)
into one file, which is used to create both the query and target databases. Next, PUPpy implements
`mmseqs search` from MMSeqs2 (39) (v14.7e284) with default parameters to align the query database
against the target database, yielding the output file "ResultDB.tsv" with alignments of every CDS against
every other CDS in the input community. The alignment percentage identity threshold of 'mmseqs search'
can be modified from its default value of '-p 0.3' in `puppy-align` to increase or decrease the alignment
stringency. Providing '-p' > 0.3 decreases the alignment stringency and thus increases the chances of
finding unique genes. This is not recommended unless the input community is highly related and no
unique genes can be found with default parameters, as it will increase the chances of unspecific primer
amplification.

Taxon-specific primer design
The second command of PUPpy, `puppy-primers`, designs taxon-specific primers for user-defined primer
targets of the input community (Fig. 1A). `puppy-primers` expects the following inputs: 1) the key output
file of puppy-align, "ResultDB.tsv", and 2) a directory containing the CDS files of the primer targets
PUPpy should design taxon-specific primers for (Fig. 1A). As default, `puppy-primers` designs microbe-
specific primers (Fig. 1B). For example, given a 3-member bacterial community composed of *Bacteroides
thetaiotaomicron*, *Bacteroides ovatus*, and *Escherichia coli*, a microbe-specific primer set would only
amplify *B. thetaiotaomicron* but no other microbe. Users can request PUPpy to design group-specific
primers by adding the flag `-p group` (Fig. 1B). For example, given the same 3-member community
above, a *Bacteroides*-specific group primer set would amplify both *B. ovatus* and *B. thetaiotaomicron*, but
not *E. coli*. Depending on the mode, `puppy-primers` parses the alignment file to identify candidate CDSs
to design microbe- or group-specific primers. CDSs that only align to themselves are considered for
microbe-specific primers, while CDSs present in all intended targets are considered for group-specific
primers. `puppy-primers` utilizes Primer3 (55) to design primers while giving flexibility to adjust key
primer design parameters. A comprehensive list of all modifiable primer design parameters, including all
default values, is available by running `puppy-primers -h`. The key output of PUPpy `puppy-primers` for
microbe- and group-specific primers are "UniquePrimerTable.tsv" and "GroupPrimerTable.tsv",
respectively. Both files contain information about the identified gene name, gene sequence, primer pair

485   penalty scores, amplicon size, forward and reverse primer sequence, length, GC content and melting
486   temperature for all the user-defined primer targets. Running `puppy-primers` on microbe-specific mode
487   also yields the barplot "UniqueGenesPlot.pdf", which shows the number of unique and non-unique genes
488   found for every primer target of the input community.

DNA Extraction and Quantification

490   Bacterial gDNA was extracted using the DNeasy Blood & Tissue Kit (QIAGEN) and quantified with 3
491   technical replicates using the Quant-iT™ 1X dsDNA HS (High-Sensitivity) Assay (Invitrogen), following
492   the manufacturer's instructions. Microbial gDNA measured concentrations in ng/μL were converted to
493   copies/μL using the formula below, where *X* is the amount of DNA in ng/μL, and *N* is the genome size in
494   base pairs (bp) as reported by the NCBI (Table S1).

$$genomic\ copies\ per\ \mu L = \frac{(X\ ng\ *\ 6.0221x10^{23}\ molecules/mole)}{(N\ *\ 660\ g/mole)\ *\ 1x10^9\ ng/g}$$

Primer design and specificity validation

498   All taxon-specific primers used in this study were designed with PUPpy (see Table S2) and ordered with
499   Thermo Fisher Scientific (https://www.thermofisher.com/order/custom-standard-oligo/). Custom
500   Standard DNA Oligos were ordered dry, desalted, and with a synthesis scale of at least 25 nmole.

GUT and SSS defined communities

Taxon-specific primers, and their respective key parameters, designed for the GUT and SSS communities
can be found in Table S2. Microbe-specific primers for both the GUT and SSS community were designed
separately by running `puppy-align` with default values, followed by `puppy-primers` with the parameters
`-s 125 175 -optm 62 -tmd 1.5`. Group-specific primers targeting the 3 major taxa of the SSS community
were designed in 3 distinct runs by running `puppy-align` with default values, followed by `puppy-
primers` with the parameters `-p group -pr <taxon_CDSs> -s 125 175 -optm 62 -tmd 1.5`, where
taxon_CDSs were the CDS files of the1) *Enterocloster* species, 2) *B. thetaiotaomicron* strains, and 3) *E.*
coli substrains. Primer specificity was experimentally validated by PCR and gel electrophoresis. Each
taxon-specific primer in the GUT and SSS communities was validated against 4 conditions: 1) the
gDNA(s) targeted by the primer pair 2) the 10-member pool gDNA, 3) the 10-member pool without the
target gDNA(s), and 4) a no-template control (water). The 10-member GUT and SSS communities were
made by pooling an equal ratio of genomic copies of the respective microbial gDNA samples (quantified
as above).

Muribaculaceae complex community

The Muribaculaceae-specific primer cocktail was created by pooling microbe-specific primer pairs
targeting individual *Muribaculum* members (see Table S2). To maximise the chances of specificity within
a complex community, microbe-specific primers targeting *Muribaculum* members were designed against
a comprehensive input community of 167 gut commensals (including the target Muribaculaceae)
available in the Tropini Strain Library (see Table S1). Although all Muribaculaceae-specific primers
target the same CDS across intended targets, distinct primers were designed to account for base pair
differences that may have affected primer annealing. PUPpy was used to identify a gene shared across all
Muribaculaceae members and individual primers were manually designed to maximise specificity across
all target members. The specificity of the Muribaculaceae-specific primer cocktail within a complex
community was validated using a Muribaculaceae-depleted and -positive conventional mouse model (35).
Conventional mice were treated with 15% (w/v) osmotic laxative polyethylene glycol (PEG) for 7

days. Following PEG administration, fecal samples from both treated (Muribaculaceae-depleted) and
untreated (Muribaculaceae-positive) mice were collected and total DNA was extracted from faecal
contents using DNeasy 96 PowerSoil Pro QIAcube HT Kit (QIAGEN, Germantown, MD, USA),
following the manufacturer's protocol. Muribaculaceae presence or absence in both samples was
confirmed by 16S rRNA sequencing. The Muribaculaceae-specific primer cocktail was experimentally
validated by PCR and gel electrophoresis against the following 4 conditions: i) 11 distinct *Muribaculum*
gDNA samples, ii) no-template control (water), iii) Muribaculaceae-depleted samples, and iv)
Muribaculaceae-positive.
538
All PCR reactions used to experimentally validate primers contained 7.5 μL of 2X DreamTaq
Green PCR Master Mix (Thermo Fisher^{TM}), 5 μL of NF-H$_2$O, 0.75 μL of forward and reverse primers
(final concentration of 340 nM) and 1 μL of DNA template. PCR amplification was performed on a
C1000 Touch^{TM} Thermal Cycler (Bio-Rad) with the following program: 98°C for 2 min, 40 cycles of
(98°C for 30 s, *specific_annealing_T* for 30 s, 72°C for 15 s), 72°C for 5 min and 4°C hold. The
annealing temperature of each primer pair (*specific_annealing_T)* can be found in Table S2. Extension
time was decreased from 1 minute to 30 seconds to avoid unspecific amplification. Following
amplification, 8 μL of PCR product was run on a 1.5% agarose gel stained with SYBR^{TM} Safe DNA Gel
Stain (Invitrogen) at 90V for 60 minutes. Primer specificity was assessed by visually inspecting gels for
the presence or absence of bands (Fig. S1, S2 and S3).

Shotgun and 16S rRNA Sequencing and Analyses

Library preparation and sequencing for the defined communities were performed at Sequencing +
Bioinformatics Consortium (SBC) at the University of British Columbia (Vancouver, Canada). Extracted
DNA was quantified using Qubit fluorometry. Shotgun metagenomic libraries were then prepared using
the Illumina DNA prep library preparation kit (Illumina, San Diego, CA, USA). These libraries were
sequenced on a NextSeq Mid output flow cell, generating paired-end 150 bp reads. For 16S, the V3 and
V4 regions of the 16S rRNA gene were PCR amplified using the following primers (72): F: 5′-
TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG and R: 5′-
GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC. These
amplicons were then converted to sequencing libraries using an 8-cycle indexing PCR with Nextera XT
primers (Illumina, San Diego, CA, USA). These libraries were sequenced on a MiSeq v2 flow cell
(Illumina, San Diego, CA, USA) to generate paired-end 250 bp reads. Raw base call data (bcl) were
converted into fastq format using the bcl2fastq conversion software from Illumina. All computational
analyses were performed on the UBC ARC Sockeye high-performance computing platform.
16S library preparation (73) for the Muribaculaceae-depleted and -positive samples was performed at
Gut4Health, BC Children's Hospital Research Institute, Vancouver, BC, Canada. The V4 region was
amplified using the following sequences: F: 5′-TATGGTAATTGTGTGYCAGCMGCCGCGGTAA and
R: 5′-AGTCAGTCAGCCGGACTACNVGGGTWTCTAAT. Amplicon libraries were purified,
normalized, and pooled using the SequalPrep^{TM} normalization plate (Applied Biosystems). Library
concentrations were verified using a Qubit TM dsDNA high sensitivity assay kit (Invitrogen) and KAPA
Library Quantification Kit (Roche) following manufacturer details. The purified pooled libraries were
submitted to the Bioinformatics + Sequencing Consortium at UBC which verifies the DNA quality and
quantity using an Agilent high sensitivity DNA kit (Agilent) on an Agilent 2100 Bioanalyzer. Sequencing
was performed on the Illumina MiSeq^{TM} v2 platform with 2 x 250 paired end-read chemistry.
573
For shotgun sequencing, the quality of raw reads was evaluated using FastQC (50) (v0.11.9) and
MultiQC (51) (v1.10.1). Raw reads were cleaned using fastp (52) (v0.23.2) with the parameters `-q 20 -p -
3 -5 -M 20 -W 4 -w 15 -c -l 50 --dedup --dup_calc_accuracy 5`. Cleaned reads were used to calculate
relative abundance using both a heuristic alignment approach and an LCA approach. In the heuristic
alignment-based approach, reads were mapped to a custom reference index of the 10 concatenated

microbial genomes using Bowtie2 (37) (v2.4.2) with the following parameters: `-D 20 -R 3 -N 0 -L 20 -i
S,1,0.50 --reorder --no-unal -p 40`. Relative abundance of individual microbes was calculated using
CoverM (53) (v0.6.1) with the following key parameters `coverm genome -m relative abundance`. In the
LCA approach, relative abundance was calculated with Kraken2 (38) (v2.1.2) using trimmed reads (as
described above) and a custom database of the 10 respective community members.
584
For 16S rRNA sequencing, raw reads were quality checked with FastQC (50) (v0.11.9) and then
analysed with QIIME2 (62) (v2023.2) to estimate microbial relative abundance. Reads were processed
with DADA2 (63) (v1.20) to trim primers and truncate sequences with quality score below 30. Denoised
sequences were assigned taxonomic labels using a classifier trained on the SILVA (71) (v138.1) 99%
database.

In silico Illumina reads generation and data analysis

In silico Illumina reads for the SSS community were simulated with InSilicoSeq (54) (v1.5.4) to achieve
known input proportions, using the following parameters: `--mode basic --draft <SSSgenomes> --
coverage <InputCoverage> --n_reads 1.1M`. The genomes of all SSS members were used to generate *in
silico* reads at desired input proportions, determined by coverage. To achieve a wide range of input
coverages for each member, all 5 default `--coverage` values offered by InSilicoSeq (54) were used:
uniform, halfnormal, lognormal, exponential, and zero_inflated_lognormal. Relative abundance from
these *in silico* Illumina shotgun sequencing reads was calculated using both heuristic alignment and LCA
approaches, as described above.

Quantification of Absolute Microbial Counts

Absolute quantification of microbial counts, measured in copies/µL, for both the 10-member GUT and
SSS pools was achieved with droplet digital PCR (ddPCR). All ddPCR reactions were run at Gut4Health
(RRID:SCR_023673). The same microbe-specific primers validated for the GUT and SSS communities
were used in ddPCR (see Table S2). The respective 9-member pools for each primer pair, lacking the
intended target, were run as negative controls. Absolute microbial counts were converted to relative
abundance by dividing the copies/µL of each organism by the sum of copies/µL of all organisms in the
community. Microbial relative abundance measured by ddPCR in the SSS community was calculated
from 9 organisms instead of 10 as we could not design microbe-specific primers for *E. coli* MC4100.
608
Briefly, the 10-member GUT and SSS gDNA pools were diluted in NF-H_2O (Fisher Bioreagents)
to achieve an expected concentration of the target DNA between 0 and 5000 copies/µL. Every ddPCR
reaction was prepared in a semi-skirted ddPCR 96-well plate (Bio-Rad) and contained 11 µL 2X
QX200[TM] ddPCR[TM] EvaGreen Supermix (Bio-Rad), 0.45 µL of forward and reverse primers (final
concentration of 204 nM), 8.1 µL of NF-H_2O, and 2 µL of the diluted gDNA pool. The ddPCR plate was
placed in an Automated Droplet Generator (Bio-Rad), together with DG32 Automated Droplet Generator
Cartridges (Bio-Rad), Pipet Tips for AutoDG[TM] system (Bio-Rad), and Automated Droplet Generation Oil
for EvaGreen (Bio-Rad). Following droplet generation, the ddPCR plate was sealed with pierceable foil
(Bio-Rad) using a PX1[TM] PCR Plate Sealer (Bio-Rad). After sealing, the ddPCR plate was placed in a
C1000 Touch[TM] Thermal Cycler (Bio-Rad) and amplified using the following program: 95°C for 5 min,
40 cycles of (95°C for 30 s, 62°C for 1 min), 4°C for 5 min, 90°C for 5 min and 4°C hold. All changes in
temperature were limited to 2°C/s ramp rate. Droplet reading was performed with a QX200[TM] Droplet
Reader (Bio-Rad) using ddPCR[TM] Droplet Reader Oil (Bio-Rad). Manual selection of the amplitude
threshold and data analysis were performed on the QX Manager v2.0 software (Bio-Rad).

Average Nucleotide Identity Analyses

Average nucleotide identity (ANI) analyses for the GUT and SSS communities were performed using the
ANIb method of pyani (49) (v0.2.9). The ANI outputs were imported into RStudio and heatmaps were
plotted using ggplot2.

M. intestinale **G6 sequencing and assembly**

High molecular weight genomic DNA extraction

M. intestinale G6 was cultured following the conditions outlined in Table S1. High molecular weight
genomic DNA (gDNA) was isolated using a Genomic-tip 100/G Kit (QIAGEN) following the
manufacturer's protocol. Extracted DNA was allowed to relax at 4°C overnight and was then quantified
using a Quant-iT™ 1X dsDNA HS Assay Kit (Invitrogen) and quality controlled using
spectrophotometry. DNA integrity was confirmed on a 1% agarose gel.

Genome sequencing

Library construction on gDNA extracted from pure *M. intestinale* G6 cultures was performed with
NEBNext® Companion Module for Oxford Nanopore Technologies® (ONT) Ligation Sequencing (New
England Biolabs, MA, USA), ONT Ligation Sequencing Kit (SQK-LSK109, Oxford Nanopore
Technologies, UK) and Flongle Sequencing Expansion kit (EXP-FSE001, Oxford Nanopore
Technologies, UK) following the manufacturer's instructions. Sequencing was performed with Flongle
flowcell (FLO-FLG001) on MinION sequencer (ONT).

De novo **genome assembly**

Sequencing raw reads were basecalled with Guppy (64) (v4.5.3) and Bonito (65) (v3.1) model and then
assembled with Flye (66) (v2.7) to generate a draft genome. The draft genome was first polished with
Racon (67) (v1.4.22) using -m 8 -x -6 -g -8 -w 500, then polished with Medaka (68) (v1.4.3) using
Medaka model for Bonito.

The final assembly resulted in a single 3,123,480 bp circular genome, with average coverage of 467x. The
assembly quality was evaluated using CheckM (69) (v1.0.18), which showed 98.4% completeness and
0.26% contamination. The *de novo* genome assembly was annotated using Prokka(70) (v.1.14.5) using
the following settings: I) --addgenes, II) --addmrna III) --gffver 3 IV) --genus Muribaculum, V) --
kingdom Bacteria, VI) --gcode 11 which identified including 2918 CDSs.

Data and Software Availability

Code, including vignettes and tutorials, are available in the PUPpy GitHub page:
https://github.com/Tropini-lab/PUPpy
Data is available at: https://borealisdata.ca/privateurl.xhtml?token=37d3e60b-62ca-40d8-9696-
f77f9d7ccb57

Author contributions:

H.G., Y.M.F., J.C.B., K.M.N., and C.T.: conceptualization, methodology, and software. H.G., Y.M.F., J.C.B., K.M.N., D.M.P., and C.T.: resources. H.G., C.T.: writing. H.G., Y.M.F., D.M.P.: formal analysis, validation, investigation, data curation, and visualization. C.T.: supervision, project administration, and funding acquisition. All authors read and approved the paper prior to submission.

Acknowledgments:

The authors acknowledge that the land we performed this research on is the traditional, ancestral, and unceded territory of the xwməθkwəy̓əm (Musqueam) Nation. We encourage others to learn more about the native lands in which they live and work at https://native-land.ca/

The authors would like to thank Ian Ghezzi, Paula Otto Andrade von Sperling, Bryan Merrill, Giselle McCallum, Jerry He and Thad Hughes for useful discussions, Claudia Stroppa for designing the PUPpy logo, Jerry He for inspiration on the "PUPpy" name, Sam Collins for support with mouse experiments, and Giselle McCallum for reading the manuscript and providing critical feedback. We also thank Angele Arrieta for research support. This work received support from Gut4Health (RRID:SCR_023673) at the BC Children's hospital Research Institute, and the Sequencing + Bioinformatics Consortium (SBC) and Advanced Research Computing (ARC) at the University of British Columbia. The authors acknowledge support from 4-Year Fellowship (to H.G., J.C.B, and D.M.P.), NSERC PGS-D Scholarship (to D.M.P.), Canadian Institute for Advanced Research / Humans and the Microbiome (FL-001253 Appt 3362, to C.T.), Michael Smith Foundation for Health Research Scholar Award (18239, to C.T.), Johnson & Johnson Women in STEM2D Award (015007, to C.T.), and Canada Foundation for Innovation / Infrastructure Operating Fund (38277).

 Fig. legends

Fig. 1
The PUPpy pipeline designs taxon-specific primers in defined microbial communities. (A) Overview of
the PUPpy workflow. Input coding sequence (CDS) files for primer target(s) and non-targets (the
background) are aligned using MMseqs2 (39). Candidate unique or shared genes are selected to design
microbe- and group-specific primers, respectively, with Primer3 (55). (B) The key output of PUPpy is
taxon-specific primers, which include microbe- and group-specific primers. Microbe-specific primers
selectively target individual members of the community, while group-specific primers target user-
determined collections of microbes.

Fig. 2
PUPpy-designed microbe-specific primers selectively detect microbes in a community of 10
phylogenetically distinct bacteria. (A) Pairwise Average Nucleotide Identity (ANI) based on the whole-
genome sequences of the 10 GUT community members (Methods). Color indicates percentage identity,
with darker red indicating greater similarity between sequences. Grey circles indicate percentage identity
lower than 70%. Circle size indicates alignment coverage between sequences, or the percentage of 2
sequences (in this case genomes) that are aligned. (B) PUPpy-generated bar plot showing the number of
unique genes (to the right of the strain name) identified for each member of the input community. The
high number of unique genes found across GUT community members, ranging from 1248 in *B.*
thetaiotaomicron (~26% of the CDSs) to 3269 in *E. coli* BW25113 (~74% of the CDSs), reflects the
genetic diversity observed by ANI (A). (C) Experimental conditions for validation of microbe-specific
primers specificity in the GUT community. The PCR gel image exemplifies the specificity of microbe-
specific primers tested using the conditions above for *F. prausnitzii*. (D) Binary heatmap showing the
targets amplified by each microbe-specific primer pair following the validation in (C). Each primer pair
exclusively amplified the respective intended target (black squares along the diagonal) following primer
optimization. Individual gel imaging data can be found in Fig. S1.

Fig. 3
PUPpy-designed taxon-specific primers selectively detect microbes down to the substrain level. (A)
Pairwise Average Nucleotide Identity (ANI) based on the whole-genome sequences of 10 microbes from
3 increasingly related taxa in the Species-Strain-Substrain (SSS) community. Color indicates percentage
identity, with darker red indicating greater similarity between sequences. Grey circles indicate percentage
identity lower than 70%. Circle size indicates alignment coverage between sequences, or the percentage
of 2 sequences (in this case genomes) that are aligned. ANI analysis, assessing genetic similarity based on
percentage identity and coverage, grouped the *Enterocloster* species, *Bacteroides thetaiotaomicron*
strains, and *Escherichia coli* K-12 sub-strains as 3 distinct clusters consistent with their taxonomic
assignment. *Enterocloster* species displayed lower genetic similarity, with a minimum of ~78% identity
and ~34% coverage, while *E. coli* sub-strains were almost genetically identical, sharing >99.95% identity
and ~99% coverage. (B) PUPpy-generated bar plot. The number of unique genes identified by PUPpy in
SSS community members decreases across increasingly related taxa. Between 19 and 36% of the CDSs in
Enterocloster species are unique within the SSS community, while only 0-1% of CDSs are unique across
E. coli K-12 sub-strains. (C) Experimental conditions for validation of microbe- and group-specific
primers specificity in the SSS community. Group-specific primers were not validated against the 10-
member community (i.e., experimental condition ii). (D) Binary heatmap showing selective amplification
by both microbe- and group-specific primer pair down to the sub-strain level. Microbe-specific primer
pairs could not be designed and validated for *E. coli* K-12 MC4100 because PUPpy could not identify
unique genes for this organism. Individual gel imaging data can be found in Fig. S2. (E) Experimental
conditions for validation of Muribaculaceae-specific primer specificity in a complex microbial
community from conventional mouse fecal samples. (F) Muribaculaceae quantification by 16S rRNA

sequencing of 5 Muribaculaceae-depleted (PEG-treated) (iii) and 5 Muribaculaceae-positive (untreated)
(iv) biological replicates. Relative abundance was calculated using QIIME2 (62) (see Methods section).
741
Fig. 4
PUPpy-designed taxon-specific primers enable more accurate and resolved quantification than 16S and
shotgun sequencing in defined communities. (A) Schematic of the workflow to prepare samples for
quantification (see Methods section for details). (B, C) Stacked bar plots evaluating microbial
quantification with droplet digital PCR (ddPCR), 16S and shotgun sequencing in the GUT (B) and SSS
(C) communities. ddPCR absolute counts were converted to relative abundance for a more direct
comparison to the other methods (see Methods).
749
Fig. 5
Shotgun sequencing quantification of genetically related microbes by heuristic alignment inaccurately
estimates relative abundance due to the random assignment of multimapping reads. (A, B) Scatter plot of
observed vs expected relative abundance from *in silico* Illumina reads of the Species-Strain-Substrain
(SSS) community members, generated with InSilicoSeq (54). Relative abundance was calculated using a
heuristic alignment-based approach with Bowtie2 (37), followed by CoverM (53)(A) and a lowest
common ancestor (LCA) approach with Kraken2 (38) (B) to evaluate different shotgun sequencing
quantification pipelines (see Methods for details). The dotted line represents a 1:1 slope. The blue line and
grey shaded area indicate a linear regression fitted on all data points from each taxon. (C, D) Boxplot of
the ratio of observed and expected relative abundance for each LCA (C) and individual member (D) in the
SSS community, calculated using the Kraken2 (38) pipeline.

Fig. S1
PUPpy-designed microbe-specific primers selectively amplify all targets in the GUT community. Related
to Fig. 2. Experimental validation of all microbe-specific primers by polymerase chain reaction (PCR)
and gel electrophoresis, following the conditions outlined in Fig. 2C. All 10 microbe-specific primers
selectively amplify their respective target, showing no unintended amplification. Faint primer dimers are
visible in *A. muciniphila*, *B. thetaiotaomicron*, *C. sporogenes*, *E. faecalis*, and *E. rectale*.
767
Fig. S2
PUPpy-designed microbe- and group-specific primers selectively amplify the respective targets in the
SSS community. Related to Fig. 3. (A) Experimental validation of all microbe-specific primers by
polymerase chain reaction (PCR) and gel electrophoresis, following the conditions outlined in Fig. 3C.
The initial validation of *E. clostridioformis* specific primers (top right gel) against the (-) pool without the
intended target (condition *iii*) showed unspecific amplification at different sizes than the expected
amplicon. Decreasing the extension time to 30 seconds (instead of 1 minute) prevented unspecific
amplification, as seen in the small gel. The water control was not repeated because condition iii also acts
as a negative control. (B) Experimental validation of group-specific primers by PCR, following the
conditions outlined in (A). The gels were imaged on distinct runs. All group-specific primers selectively
amplified their respective targets without unintended or unspecific amplification.
779
Fig. S3
PUPpy-designed Muribaculaceae-specific primers selectively amplify Muribaculaceae members in a
complex microbial community. Related to Fig. 3E and 3F. Experimental validation of Muribaculaceae-
specific primers by polymerase chain reaction (PCR) and gel electrophoresis, following the conditions
outlined in Fig. 3E. The 2 PCR gel images were taken in distinct instances. The Muribaculaceae-specific
primer mix amplified 10 different Muribaculaceae members, including strains that were not originally
included in the PUPpy input for specificity checks (See Table S1 and S2). *M. intestinale* NM03 was run
twice. *M. intestinale* NM65 was the only member not to be amplified, likely due to low DNA content. In
the fecal samples, the Muribaculaceae-specific primer mix accurately discriminated the absence (*iii*) and

presence (*iv*) of Muribaculaceae members in a complex microbial community, while also enabling strain-
level resolution (*i*). The numbers in brackets for the fecal samples indicate biological replicates.
791
792 **Table legends**
793
794 **Table S1. Metadata of all microbes used for the design and validation of PUPpy-designed taxon-**
795 **specific primers.** Microbes with "NA" noted under the growth conditions columns (last 4) were not
796 cultured but only used for *in silico* primer design. Media ingredients and recipes for culture conditions
797 can be found in Table S3.
798
799 **Table S2. Primers sequences and essential parameters for taxon-specific primers designed and**
800 **validated in this study.** The column 'target organisms' exclusively refers to the specificity within the
801 community for which primers were designed and does not necessarily extend to other microbial
802 communities.
803
804 **Table S3. Media ingredients and recipes used to culture microbes in this study.** Details on culture
805 conditions and media for each microbe can be found in Table S1.
806
807 **References:**
808

1. Cavicchioli R, Ripple WJ, Timmis KN, Azam F, Bakken LR, Baylis M, Behrenfeld MJ, Boetius A,

Boyd PW, Classen AT, Crowther TW, Danovaro R, Foreman CM, Huisman J, Hutchins DA,

Jansson JK, Karl DM, Koskella B, Mark Welch DB, Martiny JBH, Moran MA, Orphan VJ, Reay

DS, Remais JV, Rich VI, Singh BK, Stein LY, Stewart FJ, Sullivan MB, van Oppen MJH, Weaver

SC, Webb EA, Webster NS. 2019. Scientists' warning to humanity: microorganisms and climate

change. 9. Nat Rev Microbiol 17:569–586.

2. Zhu Y-G, Zhu D, Rillig MC, Yang Y, Chu H, Chen Q-L, Penuelas J, Cui H-L, Gillings M. 2023.

Ecosystem Microbiome Science. mLife 2:2–10.

3. Zhu A, Sunagawa S, Mende DR, Bork P. 2015. Inter-individual differences in the gene content of

human gut bacterial species. Genome Biology 16:82.

4. Greenblum S, Carr R, Borenstein E. 2015. Extensive Strain-Level Copy-Number Variation across

Human Gut Microbiome Species. Cell 160:583–594.

5. Engfer MB, Stahl B, Finke B, Sawatzki G, Daniel H. 2000. Human milk oligosaccharides are

resistant to enzymatic hydrolysis in the upper gastrointestinal tract. Am J Clin Nutr 71:1589–1596.

6.   Sela DA, Mills DA. 2010. Nursing our microbiota: molecular linkages between bifidobacteria and

milk oligosaccharides. Trends Microbiol 18:298–307.

7.   Wang M, Osborn LJ, Jain S, Meng X, Weakley A, Yan J, Massey WJ, Varadharajan V, Horak A,

Banerjee R, Allende DS, Chan ER, Hajjar AM, Wang Z, Dimas A, Zhao A, Nagashima K, Cheng

AG, Higginbottom S, Hazen SL, Brown JM, Fischbach MA. 2023. Strain dropouts reveal

interactions that govern the metabolic output of the gut microbiome. Cell 186:2839-2852.e21.

8.   Ma B, Gavzy SJ, Saxena V, Song Y, Piao W, Lwin HW, Lakhan R, Iyyathurai J, Li L, France M,

Paluskievicz C, Shirkey MW, Hittle L, Munawwar A, Mongodin EF, Bromberg JS. 2023. Strain-

specific alterations in gut microbiome and host immune responses elicited by tolerogenic

Bifidobacterium pseudolongum. Sci Rep 13:1023.

9.   Dong H, Rowland I, Yaqoob P. 2012. Comparative effects of six probiotic strains on immune

function in vitro. British Journal of Nutrition 108:459–470.

10.   Webb KA, Olagoke O, Baird T, Neill J, Pham A, Wells TJ, Ramsay KA, Bell SC, Sarovich DS,

Price EP. 2022. Genomic diversity and antimicrobial resistance of Prevotella species isolated from

chronic lung disease airways. Microb Genom 8:000754.

11.   Zlitni S, Bishara A, Moss EL, Tkachenko E, Kang JB, Culver RN, Andermann TM, Weng Z, Wood

C, Handy C, Ji HP, Batzoglou S, Bhatt AS. 2020. Strain-resolved microbiome sequencing reveals

mobile elements that drive bacterial competition on a clinical timescale. Genome Med 12:50.

12.   Sharon I, Morowitz MJ, Thomas BC, Costello EK, Relman DA, Banfield JF. 2013. Time series

community genomics analysis reveals rapid shifts in bacterial species, strains, and phage during

infant gut colonization. Genome Res 23:111–120.

13.   Bi D, Zhu Y, Gao Y, Li H, Zhu X, Wei R, Xie R, Cai C, Wei Q, Qin H. 2022. Profiling

845   Fusobacterium infection at high taxonomic resolution reveals lineage-specific correlations in

846   colorectal cancer. 1. Nat Commun 13:3336.

847   14.   Sonnenborn U. 2016. Escherichia coli strain Nissle 1917-from bench to bedside and back: history of

848   a special Escherichia coli strain with probiotic properties. FEMS Microbiol Lett 363:fnw212.

849   15.   Figler HM, Dudley EG. 2016. The interplay of Escherichia coli O157:H7 and commensal E. coli:

850   the importance of strain-level identification. Expert Review of Gastroenterology & Hepatology

851   10:415–417.

852   16.   Yan Y, Nguyen LH, Franzosa EA, Huttenhower C. 2020. Strain-level epidemiology of microbial

853   communities and the human microbiome. Genome Medicine 12:71.

854   17.   Sandrin TR, Goldstein JE, Schumaker S. 2013. MALDI TOF MS profiling of bacteria at the strain

855   level: a review. Mass Spectrom Rev 32:188–217.

856   18.   Noecker C, McNally CP, Eng A, Borenstein E. 2017. High-Resolution Characterization of the

857   Human Microbiome. Transl Res 179:7–23.

858   19.   Bailén M, Bressa C, Larrosa M, González-Soltero R. 2020. Bioinformatic strategies to address

859   limitations of 16rRNA short-read amplicons from different sequencing platforms. Journal of

860   Microbiological Methods 169:105811.

861   20.   Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. 2017. Shotgun metagenomics, from

862   sampling to analysis. 9. Nat Biotechnol 35:833–844.

863   21.   Gobert G, Cotillard A, Fourmestraux C, Pruvost L, Miguet J, Boyer M. 2018. Droplet digital PCR

864   improves absolute quantification of viable lactic acid bacteria in faecal samples. Journal of

865   Microbiological Methods 148:64–73.

22. Kralik P, Ricchi M. 2017. A Basic Guide to Real Time PCR in Microbial Diagnostics: Definitions,

Parameters, and Everything. Frontiers in Microbiology 8.

23. Dreier M, Berthoud H, Shani N, Wechsler D, Junier P. 2020. SpeciesPrimer: a bioinformatics

pipeline dedicated to the design of qPCR primers for the quantification of bacterial species. PeerJ

8:e8544.

24. Pritchard L, Holden NJ, Bielaszewska M, Karch H, Toth IK. 2012. Alignment-free design of highly

discriminatory diagnostic primer sets for Escherichia coli O104:H4 outbreak strains. PLoS One

7:e34498.

25. Thomsen MCF, Hasman H, Westh H, Kaya H, Lund O. 2017. RUCS: rapid identification of PCR

primers for unique core sequences. Bioinformatics 33:3917–3921.

26. Vijaya Satya R, Kumar K, Zavaljevski N, Reifman J. 2010. A high-throughput pipeline for the

design of real-time PCR signatures. BMC Bioinformatics 11:340.

27. Desai MS, Seekatz AM, Koropatkin NM, Kamada N, Hickey CA, Wolter M, Pudlo NA, Kitamoto

S, Terrapon N, Muller A, Young VB, Henrissat B, Wilmes P, Stappenbeck TS, Núñez G, Martens

EC. 2016. A Dietary Fiber-Deprived Gut Microbiota Degrades the Colonic Mucus Barrier and

Enhances Pathogen Susceptibility. Cell 167:1339-1353.e21.

28. Hsu BB, Gibson TE, Yeliseyev V, Liu Q, Lyon L, Bry L, Silver PA, Gerber GK. 2019. Dynamic

Modulation of the Gut Microbiota and Metabolome by Bacteriophages in a Mouse Model. Cell Host

& Microbe 25:803-814.e5.

29. Elzinga J, van der Oost J, de Vos WM, Smidt H. 2019. The Use of Defined Microbial Communities

To Model Host-Microbe Interactions in the Human Gut. Microbiol Mol Biol Rev 83:e00054-18.

887 30. Ng KM, Pannu S, Liu S, Burckhardt JC, Hughes T, Van Treuren W, Nguyen J, Naqvi K, Nguyen B,

888    Clayton CA, Pepin DM, Collins SR, Tropini C. 2022. Single-strain behavior predicts responses to

889    environmental pH and osmolality in the gut microbiota. bioRxiv 2022.08.31.505752.

890 31. Marcobal A, Kashyap PC, Nelson TA, Aronov PA, Donia MS, Spormann A, Fischbach MA,

891    Sonnenburg JL. 2013. A metabolomic view of how the human gut microbiota impacts the host

892    metabolome using humanized and gnotobiotic mice. 10. ISME J 7:1933–1943.

893 32. Seedorf H, Griffin NW, Ridaura VK, Reyes A, Cheng J, Rey FE, Smith MI, Simon GM, Scheffrahn

894    RH, Woebken D, Spormann AM, Van Treuren W, Ursell LK, Pirrung M, Robbins-Pianka A,

895    Cantarel BL, Lombard V, Henrissat B, Knight R, Gordon JI. 2014. Bacteria from Diverse Habitats

896    Colonize and Compete in the Mouse Gut. Cell 159:253–266.

897 33. Lagkouvardos I, Lesker TR, Hitch TCA, Gálvez EJC, Smit N, Neuhaus K, Wang J, Baines JF, Abt

898    B, Stecher B, Overmann J, Strowig T, Clavel T. 2019. Sequence and cultivation study of

899    Muribaculaceae reveals novel species, host preference, and functional potential of this yet

900    undescribed family. Microbiome 7:28.

901 34. Ormerod KL, Wood DLA, Lachner N, Gellatly SL, Daly JN, Parsons JD, Dal'Molin CGO,

902    Palfreyman RW, Nielsen LK, Cooper MA, Morrison M, Hansbro PM, Hugenholtz P. 2016.

903    Genomic characterization of the uncultured Bacteroidales family S24-7 inhabiting the guts of

904    homeothermic animals. Microbiome 4:36.

905 35. Tropini C, Moss EL, Merrill BD, Ng KM, Higginbottom SK, Casavant EP, Gonzalez CG, Fremin

906    B, Bouley DM, Elias JE, Bhatt AS, Huang KC, Sonnenburg JL. 2018. Transient osmotic

907    perturbation causes long-term alteration to the gut microbiota. Cell 173:1742-1754.e17.

908 36. Zhao C, Shi ZJ, Pollard KS. 2023. Pitfalls of genotyping microbial communities with rapidly

909    growing genome collections. cels 14:160-176.e3.

910   37.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. 4. Nat Methods

911   9:357–359.

912   38.   Wood DE, Lu J, Langmead B. 2019. Improved metagenomic analysis with Kraken 2. Genome

913   Biology 20:257.

914   39.   Steinegger M, Söding J. 2017. MMseqs2 enables sensitive protein sequence searching for the

915   analysis of massive data sets. Nat Biotechnol 35:1026–1028.

916   40.   Callahan BJ, Wong J, Heiner C, Oh S, Theriot CM, Gulati AS, McGill SK, Dougherty MK. 2019.

917   High-throughput amplicon sequencing of the full-length 16S rRNA gene with single-nucleotide

918   resolution. Nucleic Acids Research 47:e103.

919   41.   2023. Method of the Year 2022: long-read sequencing. 1. Nat Methods 20:1–1.

920   42.   Amarasinghe SL, Su S, Dong X, Zappia L, Ritchie ME, Gouil Q. 2020. Opportunities and

921   challenges in long-read sequencing data analysis. Genome Biology 21:30.

922   43.   Hardwick SA, Chen WY, Wong T, Kanakamedala BS, Deveson IW, Ongley SE, Santini NS,

923   Marcellin E, Smith MA, Nielsen LK, Lovelock CE, Neilan BA, Mercer TR. 2018. Synthetic

924   microbe communities provide internal reference standards for metagenome sequencing and analysis.

925   1. Nat Commun 9:3096.

926   44.   Morton JT, Marotz C, Washburne A, Silverman J, Zaramela LS, Edlund A, Zengler K, Knight R.

927   2019. Establishing microbial composition measurement standards with reference frames. 1. Nat

928   Commun 10:2719.

929   45.   McLaren MR, Willis AD, Callahan BJ. 2019. Consistent and correctable bias in metagenomic

930   sequencing experiments. eLife 8:e46923.

46.   Ihaka R, Gentleman R. 1996. R: A Language for Data Analysis and Graphics. Journal of

Computational and Graphical Statistics 5:299–314.

47.   Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis, p. 11–31. *In* Wickham, H (ed.),

ggplot2: Elegant Graphics for Data Analysis. Springer International Publishing, Cham.

48.   Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes A,

Henry L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K, Ooms J, Robinson D,

Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H. 2019. Welcome to the

Tidyverse. Journal of Open Source Software 4:1686.

49.   Pritchard L, Glover RH, Humphris S, Elphinstone JG, Toth IK. 2015. Genomics and taxonomy in

diagnostics for food security: soft-rotting enterobacterial plant pathogens. Anal Methods 8:12–24.

50.   Andrews S, Krueger F, Segonds-Pichon A, Biggins L, Krueger C, Wingett S. 2012. FASTQC: a

quality control tool for high throughput sequence data (0.11.9).

51.   Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize analysis results for

multiple tools and samples in a single report. Bioinformatics 32:3047–3048.

52.   Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics 34:i884–i890.

53.   wwood/CoverM: Read coverage calculator for metagenomics. https://github.com/wwood/CoverM.

Retrieved 10 October 2023.

54.   Gourlé H, Karlsson-Lindsjö O, Hayer J, Bongcam-Rudloff E. 2019. Simulating Illumina

metagenomic data with InSilicoSeq. Bioinformatics 35:521–522.

55.   Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, Rozen SG. 2012. Primer3-

-new capabilities and interfaces. Nucleic Acids Res 40:e115.

56.   McKinney W. 2010. Data Structures for Statistical Computing in Python. Proceedings of the 9th

Python in Science Conference 56–61.

57.   Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck T, Kauff F,

Wilczynski B, de Hoon MJL. 2009. Biopython: freely available Python tools for computational

molecular biology and bioinformatics. Bioinformatics 25:1422–1423.

58.   Rocklin M. 2015. Dask: Parallel Computation with Blocked algorithms and Task Scheduling.

Proceedings of the 14th Python in Science Conference 126–132.

59.   Waskom ML. 2021. seaborn: statistical data visualization. Journal of Open Source Software 6:3021.

60.   Hunter JD. 2007. Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering

9:90–95.

61.   Hartley J. Colorama.

62.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ,

Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT,

Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC,

Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM,

Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H,

Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR,

Keim P, Kelley ST, Knights D, Koester I, Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu Y-

X, Loftfield E, Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV,

Metcalf JL, Morgan SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF, Orchanian SB,

973   Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson MS,

974   Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD, Thompson

975   LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ, Vargas F,

976   Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren J, Weber

977   KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG.

978   2019. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. 8.

979   Nat Biotechnol 37:852–857.

980 63. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2: High-

981   resolution sample inference from Illumina amplicon data. Nat Methods 13:581–583.

982 64. guppy (4.5.3).

983 65. bonito (3.1).

984 66. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using

985   repeat graphs. 5. Nat Biotechnol 37:540–546.

986 67. Vaser R, Sović I, Nagarajan N, Šikić M. 2017. Fast and accurate de novo genome assembly from

987   long uncorrected reads. Genome Res 27:737–746.

988 68. Medaka (1.14.5).

989 69. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the

990   quality of microbial genomes recovered from isolates, single cells, and metagenomes. Genome Res

991   25:1043–1055.

992 70. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069.

993   71.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013. The

994   SILVA ribosomal RNA gene database project: improved data processing and web-based tools.

995   Nucleic Acids Research 41:D590–D596.

996   72.   Herlemann DP, Labrenz M, Jürgens K, Bertilsson S, Waniek JJ, Andersson AF. 2011. Transitions in

997   bacterial communities along the 2000 km salinity gradient of the Baltic Sea. 10. ISME J 5:1571–

998   1579.

999   73.   De Wolfe TJ, Wright ES. 2023. Multi-factorial examination of amplicon sequencing workflows

1000   from sample preparation to bioinformatic analysis. BMC Microbiology 23:107.

1001

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5