1

Impact of the clinically approved BTK inhibitors on the conformation of full-length BTK and analysis of the development of BTK resistance mutations in chronic lymphocytic leukemia.

Raji E. Joseph[#][*][a], Thomas E. Wales[#][b], Sandrine Jayne, Robert G. Britton, D. Bruce Fulton[a],

John R. Engen[b], Martin J. S. Dyer[c] and Amy H. Andreotti[*][a]

[a]Roy J. Carver Department of Biochemistry, Biophysics and Molecular Biology, Iowa State University, Ames, IA 50011, USA.

[b]Department of Chemistry and Chemical Biology, Northeastern University, Boston, MA 02115, USA.

[c]The Ernest and Helen Scott Haematological Research Institute, Leicester Cancer Research Centre, College of Life Sciences, University of Leicester, Leicester LE1 9HN, UK.

[#]These authors contributed equally to this work

[*] Co-corresponding author emails: jraji@iastate.edu, amyand@iastate.edu

ABSTRACT

Inhibition of Bruton's tyrosine kinase (BTK) has proven to be highly effective in the

treatment of B-cell malignancies such as chronic lymphocytic leukemia (CLL), autoimmune

disorders and multiple sclerosis. Since the approval of the first BTK inhibitor (BTKi), Ibrutinib,

several other inhibitors including Acalabrutinib, Zanubrutinib, Tirabrutinib and Pirtobrutinib have

been clinically approved. All are covalent active site inhibitors, with the exception of the reversible

active site inhibitor Pirtobrutinib. The large number of available inhibitors for the BTK target

creates challenges in choosing the most appropriate BTKi for treatment. Side-by-side comparisons

in CLL have shown that different inhibitors may differ in their treatment efficacy. Moreover, the

nature of the resistance mutations that arise in patients appears to depend on the specific BTKi

administered. We have previously shown that Ibrutinib binding to the kinase active site causes

unanticipated long-range effects on the global conformation of BTK (Joseph, R.E., et al., 2020,

https://doi.org/10.7554/eLife.60470). Here we show that binding of each of the five approved BTKi

to the kinase active site brings about distinct allosteric changes that alter the conformational

equilibrium of full-length BTK. Additionally, we provide an explanation for the resistance

mutation bias observed in CLL patients treated with different BTKi and characterize the

mechanism of action of two common resistance mutations: BTK T474I and L528W.

Keywords: BTK, Bruton's Tyrosine Kinase; BTK inhibitors; CLL, chronic lymphocytic leukemia;

Ibrutinib; Acalabrutinib; Zanubrutinib; Tirabrutinib; Pirtobrutinib; Fenebrutinib; drug resistance;

allostery; conformational equilibrium; HDX-MS, hydrogen/deuterium exchange mass

spectrometry and NMR, Nuclear Magnetic Resonance.

Abbreviations: BTKi, BTK inhibitor; LKD, linker-kinase domain; CLL, chronic lymphocytic

leukemia; MCL, Mantle cell lymphoma; MZL, Marginal zone lymphoma; PRR, proline-rich

region; HDX-MS, hydrogen/deuterium exchange-mass spectrometry; NMR, Nuclear Magnetic

Resonance; WT, wild-type.

INTRODUCTION

Ibrutinib (IMBRUVICA®), has revolutionized the treatment of the B cell malignancy, CLL [1]. Ibrutinib is a covalent active site inhibitor of the multi-domain B-cell kinase, BTK (Fig. 1a), that was first approved by the FDA in 2013 [2]. Inhibition of BTK disrupts signaling downstream of the B-cell receptor (BCR), a pathway on which the survival of CLL cells are dependent [3, 4]. The success of Ibrutinib has spurred the development of other BTKi including the now clinically approved BTK inhibitors: Acalabrutinib (CALQUENCE®), Zanubrutinib (BRUKINSA®), Tirabrutinib/ONO-4059 (VELEXBRU®), Pirtobrutinib/LOXO-405 (JAYPIRCA®) and Orelabrutinib (HIBRUKA®) [5-9]. All of the clinically approved BTKi (with the exception of Pirtobrutinib) are covalent active site inhibitors that bind to BTK C481 residue within the kinase active site. Pirtobrutinib is currently the first and only clinically approved non-covalent BTK active site inhibitor [7].

While BTKi are highly effective in the treatment of CLL, they are also being used to treat other B-cell malignancies such as Mantle cell lymphoma (MCL), Waldenström's macroglobulinemia, Marginal zone lymphoma (MZL) and are being evaluated in the treatment of multiple sclerosis and rheumatoid arthritis [5, 10, 11]. The plethora of clinically approved BTKi along with several other promising candidates currently in clinical trials pose a new challenge with respect to patient treatment. We now need a way to assess the suitability of a given BTKi for a given patient or disease state. Indeed, recent clinical trial data comparing one BTKi to another indicate that BTKi differ in their effectiveness in treating various conditions [12-15]. While toxicity, specificity and other criteria are used in these clinical comparisons, a molecular level characterization of the interaction of the inhibitor with full-length BTK is lacking and could be key to understanding these differences. In our previous work, using high resolution structural

biology techniques such as Nuclear Magnetic Resonance (NMR) and Hydrogen Deuterium

Exchange Mass Spectrometry (HDX-MS), we have shown that some inhibitors (Ibrutinib) exert

unanticipated allosteric effects upon binding to BTK while others (Fenebrutinib) do not [16].

Ibrutinib binding to the BTK kinase active site leads to a shift in the conformational ensemble of

full-length BTK towards an open conformation where the regulatory SRC homology (SH3 and

SH2) domains are released from the 'back' surface of the kinase domain, disrupting the closed

autoinhibited conformation of the full-length protein (Fig. 1b, [16]). Allosteric conformational

changes upon drug binding have been shown to alter protein-ligand interactions in other systems

with functional consequences *in vivo* [17-19]. The full impact of the panel of currently approved

BTKi on the conformation of full-length BTK is not known.

Despite the success of BTKi in CLL, a recurring problem is the development of BTKi

resistance. Resistance typically develops in patients ~2 years after the start of BTK inhibitor

treatment [20]. Analysis of CLL patients that stop responding to BTKi has revealed that most

develop mutations within *Btk* or in the substrate of BTK: Phospholipase C gamma 2 (PLCγ2) [21-

23]. Interestingly, the specific resistance mutations that develop in *Btk* seem to be dependent on

the specific BTKi used [23-29]. While data available so far for the BTKi inhibitors Acalabrutinib,

Zanubrutinib, Tirabrutinib and Pirtobrutinib are low in number, there is an emerging trend (Fig.

1c,d). Perhaps not surprisingly, patients treated with the reversible inhibitor Pirtobrutinib do not

develop mutations in C481 (Fig. 1d). Instead, they develop other kinase active site mutations

including BTK T474I and L528W (Fig. 1c,d) [26]. In contrast, 90% or more of CLL patients

treated with Ibrutinib and Acalabrutinib develop mutations in BTK C481 (Fig. 1d, Supp. Table 1

[21, 23]). Substitution of C481 with serine is the most common resistance mutation found in these

patients [30]. While mutation of BTK C481 to residues other than serine have also been reported

(C481F/Y/R), they occur at a much lower frequency [31]. Additionally, other sites within BTK

such as T474 and L528 are rarely (if ever) mutated in Ibrutinib and Acalabrutinib-treated CLL

patients (Fig. 1d, Supp. Table 1, [21-24, 29 , 30, 32-34]). Surprisingly, BTK T474 and L528 are

frequently mutated in CLL patients treated with the covalent inhibitors Zanubrutinib and

Tirabrutinib (Fig. 1d, [24, 27, 28]). BTK T474I and L528W are found at almost equal or higher

frequency compared to C481S in these patients (Fig. 1d, Supp. Table 1, [24, 27, 28]). Moreover,

in Zanubrutinib-treated patients, the L528W mutation was often present together with C481S

mutation (on different alleles), and with the L528W mutation present at a higher allelic frequency

compared to C481S [28]. This mutational bias with respect to covalent BTKi is unexpected given

the shared mode of action across the panel of covalent inhibitors, and the reasons for these

differences are unclear. Additionally, the mechanism of action of the BTK T474I and L528W

mutations is not known. Given the development of T474I and L528W mutations in multiple CLL

patients treated with both covalent (Zanubrutinib and Tirabrutinib) and non-covalent

(Pirtobrutinib) BTK inhibitors, we focused on these BTK mutations and explore their mechanisms

of action.

Here, we probe the impact of four clinically-approved BTKi: Acalabrutinib, Zanubrutinib,

Tirabrutinib and Pirtobrutinib on the conformation of full-length BTK. We find that each of these

BTKi brings about a unique combination of changes in full-length BTK. Acalabrutinib,

Zanubrutinib and Tirabrutinib disrupted the autoinhibited conformation of full-length BTK

similarly to Ibrutinib but did so to varying degrees. Interestingly, Acalabrutinib and Tirabrutinib

altered the dynamics of the kinase G-helix, a region that has been previously characterized as the

PLCγ substrate docking site [35]. Pirtobrutinib on the other hand stabilized the compact

autoinhibited conformation of full-length BTK and is the first BTK inhibitor observed to do so.

Additionally, we probed the mechanism of action of the BTK resistance mutations T474I and

L528W. We show that the catalytically inactive BTK L528W mutant activated the SRC kinase

HCK and that this activation is dependent on the proline-rich region within BTK. The BTK T474I

mutation disrupted binding to Zanubrutinib, Tirabrutinib and Pirtobrutinib and likely evades the

action of these drugs due to reduced binding to these inhibitors. Furthermore, we provide an

explanation for the mutational bias observed in patients treated with different covalent BTKi. The

development of the C481S resistance mutation is dependent on the half-life of the inhibitor which

likely explains the low prevalence of the C481S resistance mutation in patients treated with the

covalent inhibitors Tirabrutinib and Zanubrutinib (which have a long half-life) as compared to

Ibrutinib and Acalabrutinib (which have a short half-life). Characterization of the interaction of

BTKi with full-length BTK allows us to better interpret clinical trial results and will help guide

the choice of BTKi to be used for treatment. Furthermore, understanding the mechanism of action

of resistance mutations allows us to develop treatment strategies that either prevent or delay

development of resistance mutations and ways to treat them when they arise.

RESULTS

Assessing the impact of inhibitor binding on the isolated BTK linker kinase domain by NMR.

To evaluate the impact of inhibitor binding on BTK, we first monitored its effect on the

conformation of the linker kinase domain (LKD) fragment of BTK (Fig. 1a). The catalytic kinase

domain of BTK can adopt an active or inactive conformation in solution and previous studies have

shown that inhibitor binding can stabilize one or more of these conformations [16]. The switch

from an inactive to an active kinase conformation involves changes in key structural elements

within the kinase domain. These changes include the inward movement of the C-helix from an

inactive 'αC out' to the active 'αC in' position, changes in the side chain rotamer conformation of

BTK W395, formation of a conserved BTK K430:E445 salt bridge and unfurling of the collapsed

activation loop, leading to the exposure of the conserved Y551 on the activation loop for

phosphorylation (Fig. 2a, [16]).

Crystal structures of the BTK kinase domain in complex with Acalabrutinib, Zanubrutinib,

Tirabrutinib or Pirtobrutinib are available and show that each of these BTK inhibitors stabilize the

kinase domain in an inactive conformation similar to that of Ibrutinib-bound BTK (Fig. 2b-f, [16,

36-38]). In all the inhibitor bound structures the C-helix is in the 'αC out' position and the

activation loop is collapsed into the kinase active site, burying the conserved BTK Y551 (Fig. 2b-

f). Moreover, superposition of these BTK/inhibitor complexes shows that the structures of the

kinase domain remains largely the same regardless of which inhibitor is bound (Fig. 2g). The

structure of the Tirabrutinib bound BTK is the only structure that shows a difference; the activation

loop in that complex adopts an alternative inactive loop conformation compared to the other

inhibitor bound structures (Fig. 2e,g).

Previous NMR and HDX-MS analysis of the interaction of BTK with a different panel of

active site inhibitors has shown that there are differences between the solution and crystal behavior

[16]. To test whether the BTK inhibitors Acalabrutinib, Zanubrutinib, Tirabrutinib and

Pirtobrutinib can stabilize an inactive BTK kinase conformation in solution as predicted by the

crystal structures, we evaluated the BTK inhibitor bound complexes by NMR. The LKD fragment

of BTK was isotopically labeled with ^{15}N and a ^{1}H-^{15}N TROSY-HSQC spectrum was obtained in

the presence or absence of the inhibitor (Fig. 3). We have previously shown that BTK W395 within

the linker region (L) that precedes the kinase domain provides a useful probe to monitor the

conformational state adopted by the kinase domain in solution [16, 39]. Assignments for the apo

BTK linker kinase domain, which adopts the active (αC in) conformation, show that the W395

indole ^{1}H resonates at 10.21 ppm in the ^{1}H-^{15}N TROSY-HSQC spectrum (Fig. 3, top panel, black

spectrum). An upfield shift in the BTK W395 side chain indole NH resonance is observed upon

Ibrutinib binding and is consistent with the outward movement of the C-helix (αC out), and

stabilization of the inactive kinase domain conformation by Ibrutinib (Fig.3, top panel, cyan

spectrum [16]).

Comparing the tryptophan indole region of the ^{1}H-^{15}N TROSY-HSQC spectra for the BTK

LKD inhibitor bound to Acalabrutinib, Zanubrutinib or Tirabrutinib with that of the apo BTK LKD

protein shows that W395 undergoes an upfield shift in the presence of inhibitor (Fig. 3), suggesting

that, like Ibrutinib [16] and consistent with the crystal structures, all of these inhibitors stabilize

the inactive kinase domain conformation in solution. Interestingly, the Tirabrutinib bound

spectrum shows several peaks corresponding to W395, suggesting that the Tirabrutinib-bound

BTK kinase domain is likely adopting multiple kinase conformations in solution. Additionally, the

magnitude of the upfield shift in W395 resonance is smaller in both the Tirabrutinib and

Zanubrutinib spectra (compared to Ibrutinib and Acalabrutinib), which may reflect a relatively

larger active state population in these samples.

In contrast to the covalent inhibitors, Pirtobrutinib causes a downfield shift in W395, which

might suggest that this inhibitor is stabilizing an active conformation of BTK (Fig. 3, bottom

panel). However, the fluorinated benzene ring in Pirtobrutinib, which is adjacent to the C-helix

and W395 in the Pirtobrutinib bound structure, may cause changes in the local environment of

W395 and thereby give rise to the unusual chemical shift change. Indeed, HDX-MS data (see

below) confirms the stabilization of the inactive kinase conformation by Pirtobrutinib. All W395

assignments in the WT BTK inhibitor bound spectra were confirmed by comparison to the

corresponding ^{1}H-^{15}N TROSY-HSQC spectra of BTK LKD W395A mutant bound to the

inhibitors (Supp. Fig. S1).

Probing the effects of inhibitor binding on full-length BTK by HDX-MS.

We next evaluated the impact of inhibitor binding on the conformation of full-length BTK

by HDX-MS. We have previously shown that HDX-MS can be used to probe conformational

changes in full-length BTK that are brought about by inhibitor binding [16]. The structure of the

full-length BTK protein in the autoinhibited conformation was recently solved [40]. However,

electron density for both the N-terminal PHTH domain and proline rich region is missing in this

structure, suggesting that the N-terminal region of BTK is dynamic. Indeed, CryoEM, SAXS and

solution data all indicate that the PHTH domain is highly dynamic and likely transiently contacts

multiple sites on the core SH3-SH2-kinase domain (Fig. 1b, [40]). Changes in deuterium

incorporation must therefore be mapped on the crystal structure of the SH3-SH2-kinase fragment

of BTK, which adopts a compact inactive autoinhibited conformation (Fig. 1b, [41]). Additionally,

our previous work [16] has shown that for the majority of the inhibitor complexes studied changes

in deuterium incorporation are rarely observed in the PHTH domain suggesting that the PHTH

domain dynamics are not affected by active site occupancy. Inhibitor binding typically leads to

decreased deuterium incorporation in peptides derived from the kinase active site due to

stabilization of this region in the presence of the bound inhibitor [16]. Additionally, inhibitors such

as Ibrutinib that have allosteric effects lead to an increase in deuterium incorporation in peptides

derived from the BTK SH3 and SH2 domains as well as the SH2-kinase linker, indicating a shift

away from the autoinhibited conformation [16].

Full-length BTK was mixed with Acalabrutinib, Zanubrutinib, Tirabrutinib or Pirtobrutinib

and subjected to HDX-MS analysis. Peptides that could be followed in each of the six experimental

conditions (apo, and inhibitor bound) were used for comparison (see Supplemental Datafile). Intact

mass analysis supports a single binding site on BTK for the covalent inhibitors similar to what has

been observed previously with Ibrutinib (Fig. 4a, [16]).

Acalabrutinib, Zanubrutinib, Tirabrutinib and Pirtobrutinib show decreased deuterium

incorporation in peptides derived from the kinase N-lobe and kinase activation loop similar to that

observed previously with Ibrutinib (Fig. 4b,c, Fig. 5a-e, [16]). This is consistent with the binding

of these inhibitors in the kinase active site and stabilization of the inactive kinase conformation

observed by NMR as well as in the crystal structures. Furthermore, we note that each of these

inhibitors induce allosteric effects. Binding of Acalabrutinib, Zanubrutinib and Tirabrutinib to

BTK causes increased deuterium incorporation in peptides derived from the SH3 domain

suggesting that the autoinhibited conformation of full-length BTK is destabilized (Fig. 4b,c, Fig.

5b-d). This is similar to what has been previously observed with Ibrutinib [16]. Additionally,

Acalabrutinib and Tirabrutinib showed increased deuterium incorporation in peptides derived from

the G-helix of the kinase domain, a region that has been previously identified as the PLCγ substrate

docking site (Fig. 4b,c, Fig. 5b-d, [35]). Taken together, Acalabrutinib, Zanubrutinib and

Tirabrutinib binding to BTK leads to hybrid conformations of full-length BTK, where the kinase

domain is stabilized in an inactive conformation and the regulatory domains are disrupted from

their autoinhibitory conformation. In stark contrast, Pirtobrutinib binding to BTK shows decreased

deuterium incorporation in peptides derived from the SH3 and SH2 domains, suggesting that

Pirtobrutinib stabilizes the compact, autoinhibited conformation of full-length BTK (Fig. 4b,c, Fig.

5e). To date, Pirtobrutinib is the only inhibitor that we have tested that uniformly stabilizes both

the kinase domain as well as the regulatory domains in the inactive autoinhibited conformation.

Thus, each BTK inhibitor causes unique changes in the overall conformation of full-length BTK

that is not readily predicted from crystal structures. We next turn our attention to the distinct

resistance mutations that arise upon treatment with the different BTK inhibitors.

Probing the intrinsic effects of the BTK T474I and L528W mutations.

BTK resistance mutations can potentially confer a selective advantage to cells in several

ways: by increasing the activity of the kinase, by changing the conformation/stability of the kinase

which in turn can alter protein-protein interactions, or by disrupting drug binding. To probe the

mechanism by which the BTK T474I and L528W mutations confer resistance to Zanubrutinib,

Tirabrutinib and Pirtobrutinib we first tested the impact of each mutation on the catalytic activity

of the kinase and then investigated the impact on the overall conformation of the protein by HDX-

MS and NMR.

To test the catalytic activity of the BTK mutants we set up *in vitro* kinase assays using

purified full-length WT and mutant BTK proteins. Kinase activity was monitored by following

phosphorylation on the activation loop Y551 by western blotting. The BTK T474I mutant shows

phosphorylation on Y551, however it is lower than that of the WT protein (Fig.6a,b) suggesting

that the T474I mutation reduces the activity of the kinase. In contrast, the BTK L528W mutant is

completely inactive with no detectable phosphorylation on Y551 throughout the time course

(Fig.6c,d). These results are consistent with previous activity reports on both the BTK T474I and

L528W mutants [26, 28, 42, 43]. Given that neither the BTK T474I or the L528W mutant had

increased catalytic activity, augmenting kinase activity is not the mechanism by which these

mutations confer resistance.

Resistance mutations can change the conformation of the protein which in turn can alter

protein-protein interactions. To test whether the BTK T474I and L528W mutations alter the overall

conformation of the protein, we carried out HDX-MS analysis on the apo mutant proteins.

Comparing the BTK T474I mutant to WT BTK, the BTK T474I mutant shows increased deuterium

incorporation within the kinase domain N-lobe β2-β3 strands and the activation loop, but no

changes elsewhere on the protein (Fig.6e,f,h). These increased dynamics within the N-lobe of the

BTK T474I mutant could potentially alter drug binding to the active site. The BTK L528W mutant

on the other hand shows a slight decrease in deuterium incorporation in the kinase domain and the

SH3 domain suggesting that the L528W mutation has a slight stabilizing effect on the full-length

autoinhibited conformation of BTK (Fig.6e,g,h).

Additionally, we tested the BTK T474I and L528W mutants by NMR. An overlay of the

^{1}H-^{15}N TROSY-HSQC spectra of apo BTK LKD T474I with that of apo WT BTK LKD shows

some changes in the mutant spectrum (Fig.6i). The resonance corresponding to BTK A524 located

close (~18 Å) to T474 in the kinase active site shows line broadening in the BTK T474I mutant

spectrum suggesting increased dynamics in the mutant. This is consistent with the increased

deuterium incorporation observed in the kinase domain for this mutant by HDX-MS. The BTK

W395 resonance in the BTK T474I mutant spectrum overlaps with that of the WT suggesting that

the mutant kinase domain adopts an active conformation similar to that of the WT protein. The

BTK L528W mutant on the other hand shows a small upfield shift in the W395 resonance,

suggesting that the BTK L528W mutant kinase domain is shifted slightly towards an inactive

conformation compared to the WT protein. This change in the BTK L528W mutant is consistent

with the HDX-MS changes for this protein which suggest that the mutation has a slight stabilizing

effect on the compact autoinhibited conformation. Additionally, a new peak is observed as

expected in the BTK L528W mutant spectrum consistent with the introduction of an additional

tryptophan due to the mutation. Overall, as the BTK T474I and L528W mutations by themselves

cause only minor conformational changes, such changes are unlikely to constitute the mechanism

by which these mutations confer resistance.

BTK L528W mutant activates the SRC family kinase HCK.

Previous studies have shown that the BTK L528W mutant propagates BCR signaling

despite being catalytically dead [26, 43]. PLCγ phosphorylation and calcium signaling are

maintained in cells carrying this BTK mutation [26, 43]. This suggests that the BTK L528W

mutant might recruit other kinases to compensate for its lack of activity. The mechanism of action

of another catalytically inactive BTK resistant mutation: BTK C481F/Y, which arises in Ibrutinib

treated CLL patients, has been recently reported [44]. In that work the BTK C481F/Y resistance

mutant was shown to bind and activate the SRC family kinase HCK, thereby propagating signals

from the BCR signaling pathway [44]. This activation of HCK by BTK C481F/Y requires

phosphorylation on the BTK kinase activation loop, Y551. The phosphorylated Y551 is suggested

to bind to the HCK SH2 domain, displacing the autoinhibited conformation of HCK, thereby

activating the HCK catalytic function. To determine if a similar mechanism is at work for the

catalytically dead BTK L528W mutant, we tested the ability of the BTK L528W mutant to activate

HCK in a western blot assay by monitoring the phosphorylation levels on two substrates: PLCγ1

(pY783 phosphorylation) and HCK itself (pY levels). In addition, we also included BTK WT that

had been pre-incubated with Zanubrutinib as a control. As shown in Fig.7a,b, phosphorylation

increased on both PLCγ and HCK in the presence of full-length BTK L528W, suggesting that the

catalytically dead BTK mutant is able to activate HCK. Interestingly, Zanubrutinib bound WT

BTK also promoted increased phosphorylation levels on both PLCγ1 and HCK suggesting that

drug-inactivated BTK also activates HCK. This is distinct from what has been previously observed

for the BTK C481Y/F resistant mutation where, compared to WT BTK, the C481Y/F mutant BTK

preferentially engages HCK. Nevertheless, since the BTK L528W mutant activates HCK, we next

designed experiments to probe the requirements for this activation.

We first tested whether the regulatory domains of BTK were required for HCK activation

by the BTK L528W mutant by using the isolated linker kinase domain of BTK (a construct in

which the BTK regulatory domains have been deleted). As shown in Fig. 7c,d, the isolated linker

kinase domain of BTK L528W does not activate HCK. This suggests that the regulatory domain/s

of BTK are required for the activation of HCK and that the isolated linker kinase domain of BTK

alone is insufficient for HCK activation. This again is different from what has been previously

reported; the BTK C481Y/F resistant mutant requires the kinase domain for HCK activation [44].

We therefore turned our focus onto the BTK regulatory domains to isolate the specific region/s

required for HCK activation by the BTK L528W resistance mutation.

Displacement of the SH3 domain from its autoinhibitory conformation upon binding of a

proline-rich ligand is a classic mechanism by which SRC family kinases such as HCK are activated

[45]. BTK contains such a proline-rich ligand sequence within its regulatory region and so we

tested HCK activation by a BTK mutant lacking the proline-rich sequence (BTK L528W/Pro: BTK

L528W/ P189A/P192A/P203A/P206A). As shown in Fig. 7e,f, the BTK L528W proline-depleted

mutant does not activate HCK. This data again emphasizes the difference between the mechanism

of action of the BTK L528W mutant from the previously characterized BTK C481Y/F mutant.

Additionally, this data is consistent with the observation that WT BTK can also activate HCK, as

the proline-rich region is present in both WT and mutant L528W BTK. The ability of both WT

and L528W mutant BTK to activate HCK, however, suggests that there must be additional reasons

for the selection of the BTK L528W resistance mutation in patients.

BTK L528W is more stable than WT BTK.

Changes in protein levels have been shown to cause resistance in several cancers [46].

Cellular protein levels can be altered due to changes in protein expression, stability or degradation,

or changes at the RNA level [46]. An increase in BTK protein levels *in vivo* could drive protein-

protein interactions typically not observed in the WT background. Interestingly, we note that

protein yield after bacterial expression and purification of the full-length BTK L528W mutant is

higher than that of the WT protein, suggesting that the L528W mutant is more stable than the WT

protein. To test whether the BTK L528W mutation has a stabilizing effect on BTK, we measured

the melting temperature (Tm) of the BTK L528W mutant and compared it to that of the WT protein

in a thermal shift assay. As shown in Fig.7g, the BTK L528W mutant is more stable than the WT

protein (a one degree increase in Tm compared to the WT). The increased stability of the L528W

mutant, albeit small, could result in increased protein levels, an increase in the lifetime of the

protein or altered protein trafficking *in vivo*, all of which would drive the interaction of the mutant

protein with HCK unlike the WT protein.

HDX-MS reveals that the BTK T474I and L528W mutants show reduced binding to BTK inhibitors.

Disruption of inhibitor binding due to mutations is a classic mechanism by which resistance

arises [47, 48]. While loss of inhibitor binding is likely irrelevant to the catalytically dead BTK

L528W mutant, we nevertheless evaluated it along with the BTK T474I mutant using HDX-MS

and NMR. HDX-MS data show that Tirabrutinib, Zanubrutinib and Pirtobrutinib induce overall

changes in deuterium incorporation for both WT BTK and the T474I mutant (Fig. 8a,b,c) These

data suggest that the inhibitors are still capable of binding to the T474I mutant. However, in the

presence of Tirabrutinib, Zanubrutinib or Pirtobrutinib, the magnitude of the protection in the

kinase domain (N-lobe β2-β3 strands and activation loop) is consistently reduced in drug bound

T474I mutant as compared to the WT protein (Fig. 8a,b,c,e). The differences in deuterium

exchange for drug binding to WT and mutant BTK suggest that the T474I mutation either causes

a reduction in inhibitor binding or otherwise alters the mode of drug interaction in the active site.

Specifically, the peptides derived from the kinase activation loop in particular do not show as much

protection in the mutant relative to the WT suggesting that the inhibitors do not stabilize the

activation loop in the inactive conformation upon binding to the T474I mutant. Interestingly, the

changes observed in the BTK regulatory domains (SH3 and SH2 domains) upon Zanubrutinib,

Tirabrutinib or Pirtobrutinib binding to WT BTK as well as increased deuterium incorporation in

the G-helix of the kinase domain upon Tirabrutinib binding to WT BTK are maintained or reduced

slightly in the drug bound BTK T474I mutant. In stark contrast to the BTK T474I mutant, the BTK

L528W mutant does not show any change in deuterium incorporation in the presence of

Zanubrutinib, Tirabrutinib or Pirtobrutinib, providing strong evidence that the BTK L528W

mutant does not bind the inhibitors (Fig.8d). Taken together, both the BTK T474I and the BTK

L528W mutation impact inhibitor binding. Indeed, these results are consistent with previous

studies that show that the BTK T474I and the BTK L528W mutations disrupt binding to

Zanubrutinib and Pirtobrutinib [26]. We next tested the impact of the resistance mutations on

inhibitor binding by NMR.

Probing the effects of BTK T474I and L528W mutations on inhibitor binding by NMR.

Comparison of the Zanubrutinib bound WT BTK LKD spectrum to that of BTK T474I

shows that the BTK T474I mutant undergoes minor changes in the presence of Zanubrutinib,

suggesting that the mutant is binding weakly to the inhibitor (Fig. 9a, left and middle panels). The

W395 resonance shows little to no chemical shift change upon addition of Zanubrutinib to the

BTK T474I mutant, suggesting that this drug does not stabilize the inactive conformation of the

mutant kinase (Fig. 9a, middle panel). These results are consistent with HDX-MS changes in the

T474I mutant upon Zanubrutinib binding (Fig. 8a).

Spectral comparison of Tirabrutinib bound to WT and BTK T474I (Fig. 9b, left and middle

panels) shows drug induced chemical shift changes for both complexes, supporting the conclusion

from the HDX-MS data (Fig. 8b) that the drug binds to both WT and mutant BTK. However, the

spectral changes observed in the spectrum of Tirabrutinib bound to BTK T474I are different from

that of the WT/Tirabrutinib bound spectrum (Fig. 9b). Although BTK W395 gives rise to multiple

peaks in the spectra of Tirabrutinib bound to both WT and T474I, the W395 resonances in the

BTK T474I/Tirabrutinib complex are shifted upfield to a lesser extent compared to the same

resonance in the WT/Tirabrutinib spectrum. This suggests that the BTK T474I mutation alters the

binding to Tirabrutinib and the inactive kinase domain conformation (αC out) is less populated in

the drug bound mutant protein compared to WT BTK bound to Tirabrutinib. These results are

consistent with the HDX-MS results that show reduced protection in peptides corresponding to the

kinase activation loop in the mutant compared to WT due to the lack of stabilization of the kinase

inactive conformation upon Tirabrutinib binding.

Spectral overlay of the Pirtobrutinib bound BTK T474I with the apo T474I protein shows

pronounced chemical shift changes in the presence of the inhibitor, providing evidence consistent

with the HDX-MS data (Fig. 8c), that Pirtobrutinib binds the BTK T474I mutant (Fig. 9c, middle).

However, unlike Pirtobrutinib bound to WT BTK, multiple W395 peaks are visible in the spectrum

of Pirtobrutinib bound to BTK T474I. The resonance frequencies of the additional peaks are shifted

toward that of the apo protein suggesting the possibility of fast exchange between inhibitor bound

and unbound states. These observations suggest that the T474I mutation reduces affinity towards

Pirtobrutinib and are consistent with previously published SPR binding data [26].

BTK inhibitors that retain binding to the BTK T474I mutant could serve as potential

alternate treatment option for patients with this mutation. Additionally, the lack of or low

frequency of the BTK T474I mutation in patients treated with Ibrutinib and Acalabrutinib could

be due to the inability of this mutation to disrupt inhibitor binding. We therefore tested the ability

of the BTK T474I mutant to bind Ibrutinib, Acalabrutinib and Fenebrutinib (GDC-0853). The

NMR data suggest that the BTK T474I mutation does not impact Ibrutinib or Fenebrutinib binding

(Fig. 9d,f), but Acalabrutinib binding is reduced (Fig. 9e). Taken together, the T474I mutation

significantly reduces binding to Zanubrutinib, Tirabrutinib and Pirtobrutinib, retains binding to

Ibrutinib and Fenebrutinib, and may have an adverse effect on Acalabrutinib binding. Given that

the BTK T474I mutant retains activity (albeit reduced activity), disruption of drug binding is a

possible mechanism by which this mutation escapes inhibition.

Spectral overlays of the BTK L528W mutant with and without Zanubrutinib show no

chemical shift changes (Fig. 9a, right panel) suggesting that the mutation completely disrupts

inhibitor binding in complete agreement with the HDX-MS data (Fig. 8d). Tirabrutinib does show

chemical shift changes, but the changes are markedly different from that observed in the WT

protein (Fig. 9b, right). In the L528W spectrum in the presence of Tirabrutinib, W395 shifts in the

opposite downfield direction compared to the WT spectrum suggesting that Tirabrutinib could be

binding to the BTK L528W in a different orientation. Alternatively, the distinct chemical shift

change could be due to the mutated L528W residue altering the local chemical environment within

the kinase.

The Pirtobrutinib-bound BTK L528W spectrum (Fig. 9c) shows two resonance positions,

one of which overlaps with the W395 resonance in the apo protein and the other that corresponds

to that of the mutant protein bound to Pirtobrutinib. This data suggests a mixture of inhibitor bound

and unbound BTK kinase domain in solution, likely due to a reduction in Pirtobrutinib affinity

caused by the L528W mutation. Although the L528W mutation alters binding to both Tirabrutinib

and Pirtobrutinib, the NMR data suggests that it retains partial binding unlike the HDX-MS data

that suggests complete disruption of binding. The higher inhibitor concentrations used in the NMR

experiments compared to the HDX-MS experiments likely explain this discrepancy. Interestingly,

we note similarities between the BTK L528W Pirtobrutinib bound spectrum and that of

Tirabrutinib bound spectrum, suggesting that the BTK kinase domain adopts similar

conformations in solution when bound to these different drugs. Additionally, the BTK L528W

mutant retains binding to Fenebrutinib, however Ibrutinib and Acalabrutinib binding are disrupted

(Fig. 9d-f, right). Taken together, the BTK L528W mutation significantly disrupts Zanubrutinib,

Tirabrutinib and Pirtobrutinib binding. However, drugs based on Fenebrutinib could be developed

to treat patients carrying this mutation (see discussion below).

DISCUSSION

Binding of BTK active site inhibitors can have long-range effects on the protein. Here, we build on earlier work to show that three of the clinically approved BTKi (Acalabrutinib, Zanubrutinib and Tirabrutinib) shift the conformational ensemble of full-length BTK, destabilizing the autoinhibited conformation of the SH3 and SH2 domains to varying degrees (Fig.4 and 5). In marked contrast, Pirtobrutinib led to the stabilization of the compact autoinhibited conformation of full-length BTK. The exposure or stabilization of the regulatory domains of BTK by active site BTK inhibitors can alter the interaction of BTK with ligands *in vivo*. Indeed, BTK inhibitors have been shown to differ in their effectiveness at terminating signals downstream of the Fc receptor *versus* the B cell receptor [49, 50]. As more BTKi become available, a molecular-level understanding of the interaction of the inhibitor with full-length BTK will aid the interpretation of the efficacy of different BTK inhibitors in treating disease states driven by different signaling pathways.

The development of specific resistance mutations in patients treated with different BTKi is intriguing. Covalent BTKi rely on BTK C481 within the kinase active site. The C481S mutation prevents the covalent attachment of these inhibitors to BTK and converts the mode of binding of these drugs to that of a reversible inhibitor. Importantly, we and others have shown that the BTK C481S mutation does not prevent binding of Ibrutinib to the BTK kinase domain [16, 26]. In fact, the BTK C481S mutant binds to Ibrutinib just as well as the WT protein *in vitro* under equilibrium conditions where drug concentrations do not vary over time [16]. However, *in vivo*, drug concentrations change with time; inhibitor concentrations peak rapidly after intake followed by a decrease as the drug is cleared (represented by the half-life of the drug). The occupancy of BTK C481S protein by these covalent inhibitors in patients is therefore dictated by the availability of

these covalent inhibitors over time. Interestingly, Ibrutinib and Acalabrutinib have short half-lives

compared to Zanubrutinib and Tirabrutinib (Table 1). The short half-lives of Ibrutinib and

Acalabrutinib suggests that the BTK C481S mutant is likely to be unoccupied when inhibitor

concentrations fall, thereby allowing the BTK C481S mutant to escape from inhibition.

Conversely, the longer half-life of Zanubrutinib and Tirabrutinib in patients suggests that the BTK

C481S mutant is likely to remain occupied (inhibited) for a longer period of time. The C481S

mutation therefore does not offer a selective advantage as a resistance mutation in Zanubrutinib

and Tirabrutinib treated patients and likely explains the lower frequency of occurrence of this

mutation in these patients compared to Ibrutinib and Acalabrutinib treated patients. Taken

together, the BTK C481S mutant can escape covalent inhibition at low drug concentrations *in vivo*

and is therefore predicted to arise in patients treated with BTK covalent inhibitors that have short

half-lives.

The T474I mutation arises frequently in Tirabrutinib and Pirtobrutinib treated CLL patients

(Fig. 1d). Our HDX-MS and NMR binding studies show that the T474I mutation disrupts binding

to both inhibitors. Reduced binding of BTK T474I mutant to Tirabrutinib and Pirtobrutinib, along

with the partial catalytic activity retained by this mutant, would allow BTK T474I mutant to escape

inhibition and maintain BCR signaling. This gives a selective advantage to the cells that carry this

mutation. Additionally, our NMR binding studies show that the BTK T474I mutation does not

disrupt binding of Ibrutinib. This is consistent with the low occurrence of this mutation in

Ibrutinib-treated CLL patients. While the BTK T474I mutation does seem to disrupt binding to

Acalabrutinib, the lower activity of the T474I mutant (compared to the BTK C481S mutant) may

explain the higher prevalence of the BTK C481S mutation (compared to T474I) in Acalabrutinib

treated patients.

Ideas and Speculation

Successful treatment of CLL has relied on the inhibition of BTK catalytic activity to curb BCR signaling. Inhibitor-bound, catalytically-inactive BTK is incapable of propagating BCR signaling. Paradoxically, the catalytically inactive BTK L528W mutant is able to propagate BCR signals [26, 43]. A similar catalytically inactive BTK mutant, the BTK C481Y/F resistance mutation also arises in Ibrutinib and Acalabrutinib treated patients, albeit at a low frequency [44]. Since these BTK mutants are catalytically dead, other kinases must compensate for the absence of BTK activity to account for the intact BCR signaling in these patients. Exogenous kinases do not appear to compensate for WT BTK that is inactivated by BTK inhibitors, and so there must be additional differences between the catalytically dead (BTK L528W and BTK C481Y/F) mutant kinases and inhibitor bound WT BTK that is rendered catalytically inactive by virtue of drug binding. Our HDX-MS and NMR data show that BTK L528W mutation does not drastically change the conformation of BTK. However, this mutation does increase the stability of BTK compared to the WT protein. This increased stability could alter protein levels or change the trafficking/localization of the mutant protein *in vivo* which in turn could alter protein-protein interactions that drive activation of an alternate kinase such as HCK *in vivo*. Comparison of BTK protein levels in patients pre- and post-development of resistance will be needed to test this hypothesis. Alternatively, the BTK L528W mutation may require an additional change/s in the cell that allow for this dead kinase to recruit other kinases such as HCK to propagate BCR signaling. This requirement for additional changes may explain the low frequency of the L528W mutation in Ibrutinib and Acalabrutinib treated patients; the single amino acid change, C481S, combined with the rapid clearance of drug seems sufficient to drive resistance. Finally, the absence of the BTK

L528W resistance mutation in Ibrutinib treated CLL patients could be due to the promiscuity of

Ibrutinib; candidate compensatory kinases may be inactivated by the drug [51]. The increased

specificity of Zanubrutinib, Tirabrutinib and Pirtobrutinib on the other hand could allow for the

utilization of compensatory kinases such as HCK by BTK L528W in order to propagate BCR

signaling.

Understanding the mechanism/s by which resistance mutations evade inhibition allows for

counter strategies to be devised [52-54]. The reversible BTK inhibitor Pirtobrutinib is able to

inhibit the BTK C481S mutant due to its long serum half-life and has been suggested as a treatment

option for patients that develop this mutation [55]. Alternatively, drug occupancy in the context of

the BTK C481S resistance mutant could be increased by administering a twice daily dose of the

inhibitor as opposed to the single daily dose possibly circumventing their short half-lives [56].

This increased exposure time to the covalent inhibitor could cause increased side effects and so

such risks should be considered before altering the dosage regimen. Alternatively, Tirabrutinib,

which exhibits the longest half-life among the available covalent BTK inhibitors, could be used to

slow progression of disease in patients that develop the C481S resistance mutation.

The BTK T474I mutation disrupts binding to Zanubrutinib, Tirabrutinib, Pirtobrutinib and

Acalabrutinib (Fig.8, 9) but retains binding to Ibrutinib and Fenebrutinib creating possible options

for patients that develop the BTK T474I resistance mutation. The BTK L528W mutant is

catalytically inactive, hence treatment with any BTK active site inhibitor is futile. Previous studies

have shown that the similarly inactive BTK C481Y/F mutant can be suppressed by the use of

PROTACS to induce degradation of the full-length protein [44]. Importantly, our data indicate that

although the BTK L528W mutation disrupted binding to most BTK inhibitors it retains binding to

Fenebrutinib (Fig.9), suggesting that PROTACS based on Fenebrutinib or similar backbones could

be developed to treat this resistance mutation [57]. Alternate treatment approaches such as

targeting HCK or other BCR signaling proteins such as BCL-2 [58] could also be used to counter

this resistance mutation. As additional patient data becomes available, the patterns of resistance

mutations for different inhibitors will become clearer and strategies to circumvent resistance using

existing inhibitors should improve. As for many areas of medicine, the treatment landscape should

benefit from the ongoing era of personalized medicine.

MATERIALS AND METHODS

Constructs and reagents: The bacterial expression constructs for murine BTK linker kinase

domain (LKD) and full-length (FL) have been described previously [39]. All BTK constructs carry

the solubilizing Y617P mutation for bacterial expression [39]. All mutations were made using the

site directed mutagenesis kit (Agilent), and the sequences of all constructs were confirmed by

sequencing at the Iowa State University DNA facility. Acalabrutinib, Zanubrutinib, Tirabrutinib

and Pirtobrutinib were purchased from MedChem Express. Ibrutinib was purchased from

Selleckchem. The pCDF-1 Duet HCK SH3-SH2-Kinase domain/YopH construct with a 'YEEI'

tail, a kind gift from Dr. Tom Smithgall, was mutated to convert the 'YEEI' tail residues to that of

the WT protein (YQQQ). The pGEX PLCγ1 cSH2-linker Y771F/Y775F protein substrate

construct has been described previously [59].

Protein expression and purification: Expression and purification of the PLCγ1 cSH2-linker

Y771F/Y775F protein substrate has been described previously [59]. HCK, Full-length BTK and

BTK linker kinase domain (WT and mutants) were produced by co-expressing with YopH in

BL21(DE3) (Millipore Sigma) or BL21-Gold(DE3) cells (Agilent Technologies) as described

previously [39]. Briefly, the culture was grown at 37 °C to an O.D. 600 nm of 0.6 to 0.8. The

temperature of the culture was lowered to 18°C and then induced with either 1.0 mM IPTG for

BTK LKD construct or 0.1 mM IPTG for BTK full-length and HCK. The culture was harvested

24 hours after induction and the pellets were resuspended in lysis buffer (50 mM KH_2PO_4, pH 8.0,

150 mM NaCl, 20 mM imidazole and 0.5 mg/ml lysozyme) and stored at -80°C. Cells were lysed

by thawing and the action of lysozyme, and 3000 U DNAse I (Sigma) and 1 mM PMSF were

added to the lysate, incubated at RT for 20 minutes and then spun at 16,000 rpm for one hour at

4°C. Glycerol was added to the supernatant to a final concentration of 10 % and was then incubated

with Ni-NTA resin (QIAGEN) for two hours, washed with Tris pH 8.0, 75 mM NaCl, 40 mM

imidazole, and eluted in 20 mM Tris pH 8.0, 150 mM NaCl, 250 mM Imidazole, and 10% glycerol.

Eluted protein was flash frozen in liquid nitrogen and stored at -80°C. The proteins were

concentrated and further purified by size exclusion chromatography (Hiload Superdex 26/60 200

pg or Hiload Superdex 26/60 75 pg, GE Healthcare). The fractions containing pure protein were

pooled, concentrated, snap frozen and stored at -80° C. The final buffer consists of 20 mM Tris

pH 8.0, 150 mM Sodium chloride, 0.02 % Sodium azide and 10% glycerol. Initial phosphorylation

levels of all purified BTK and HCK proteins used in this study is below western immuno-detection.

NMR: Uniformly [15]N labeled BTK samples were produced as described earlier by growth in

modified M9 minimal media containing [15]N ammonium chloride (1g/L, Cambridge Isotope

Laboratories, Inc.) as the sole source of nitrogen [39]. The final NMR sample buffer consists of

20 mM Tris, 150 mM Sodium chloride, 10% glycerol, and 0.02 % Sodium azide at pH 8.0. All

NMR spectra were collected at 298 K on a Bruker AVIII HD 800 spectrometer equipped with a 5

mm HCN z-gradient cryoprobe operating at a [1]H frequency of 800.37 NMR samples with

inhibitors consisted of 150 µM [15]N labeled BTK, mixed with 200 µM inhibitor in 2% DMSO. All

data were analyzed using NMRViewJ [60].

HDX-MS: General procedures for HDX-MS of BTK have been described in detail previously [39].

Details specific to experiments conducted here are provided in the Supplemental Datafile in the

format recommended [61] for HDX-MS experimental descriptions. All HDX-MS data have been

deposited to the ProteomeXchange Consortium via the PRIDE [62] partner repository with the

dataset identifier PXD047865. Briefly, prior to continuous labeling HDX experiments, purified

BTK full-length wild-type, T474I or L528W (20 µM) and inhibitor (40 µM), (20 mM Tris pH 8.0,

150mM NaCl, 10% glycerol, 2% DMSO) were allowed to interact at 21 °C for 1 hour. After the

binding reactions, both the free kinase and kinase bound to inhibitor were placed on ice prior to

deuterium labeling. Deuterium labeling proceeded for the times described using labeling buffer,

and labeling was stopped with an equal volume of quench buffer at 0°C (details in Supplemental

Datafile). Quenched samples were immediately analyzed using a Waters nanoACQUITY with

HDX technology using online pepsin digestion with a Waters Enzymate immobilized pepsin

column and UPLC separation of the resulting peptic peptides. Mass spectra were acquired using a

Waters Synapt HDMS[E] mass spectrometer. Peptides generated from online pepsin digestion were

identified with Waters Protein Lynx Global Server 3.0 using separate unlabeled protein that was

prepared in the same manner as protein labeled with deuterium. Deuterium incorporation was

quantified using Waters DynamX 3.0. Deuterium levels for each peptide were calculated by

subtracting the average mass of the undeuterated control sample from that of the deuterium labeled

sample; the data were not corrected for back exchange and are therefore reported as relative [63].

Vertical difference maps in Figures 4, 6 and 8 do not represent a linear sequence of non-

overlapping peptides. All coincident and overlapping peptides for comparisons in each figure are

provided in figure identified tabs of the Supplemental Datafile.

Activity assays: *In vitro* kinase assays were performed as described previously [39]. Briefly, 1 μM

BTK FL, BTK FL T474I or BTK FL L528W proteins were incubated in a kinase assay buffer (50

mM Hepes pH 7.0, 10 mM $MgCl_2$, 1 mM DTT, 5 % glycerol, 1 mM Pefabloc, and 200 μM ATP)

at room temperature for varying time. The reactions were stopped by the addition of SDS-PAGE

loading buffer and the samples were boiled, separated by SDS−PAGE, and Western blotted with

the anti-BTK pY551 antibody (BD Pharmingen[TM]) or anti-His antibody (EMD Millipore) as

described previously [39]. The bands were quantified using the ChemiDoc[TM] (Biorad) gel imaging

system. The phosphorylation levels (Anti-BTK pY551 blot) were normalized to the total protein

level (Anti-His blot). The BTK FL value at 40 minutes was set to 1 and compared to BTK FL

T474I or BTK FL L528W. The HCK activation experiments were performed by preincubating 0.2

μM of BTK WT or L528W proteins with 0.5 μM Zanubrutinib for 15 min at RT. The reaction was

initiated with the addition of 0.2 μM HCK in a kinase assay buffer (50 mM Hepes pH 7.0, 10 mM

$MgCl_2$, 1 mM DTT, 5 % glycerol, 1 mM Pefabloc, 30 μM PLCγ1 cSH2 linker Y771F/Y775F

substrate and 200 μM ATP). Time points were taken at 5, 10 and 15 minutes. The reactions were

stopped by the addition of SDS-PAGE loading buffer and the samples were boiled, separated by

SDS−PAGE, and Western blotted with the anti-PLCγ1-pY783 antibody (Cell Signaling) or anti-

pY antibody (4G10, Millipore Sigma) or anti-His antibody (EMD Millipore) as described

previously [39]. The bands were quantified using the ChemiDoc[TM] (Biorad) gel imaging system.

The HCK alone value was set to 1 and compared to HCK + BTK WT/L528W. Initial

phosphorylation levels of BTK, prior to the start of the activity assay were undetectable by western

immuno-detection.

Thermal shift assays: BTK FL WT or mutant at 5 µM was incubated with 10X final concentration

of SYPRO Orange dye (Thermo Fisher Scientific) in a total volume of 20 µL Tris buffer (20 mM

Tris, 150 mM Sodium chloride, 10% glycerol, pH 8.0). Thermal shift assays were performed on

the StepOnePlus RT-PCR machine in the Iowa State University DNA Facility. The temperature

was raised from 25 °C to 95 °C and measurements recorded with every 0.5 °C increment in

temperature. The derivative plot of the data was analyzed for the melting temperature (Tm). Assays

were performed thrice and the average Tm was calculated.

ACKNOWLEDGEMENTS

This work is supported by a grant from the National Institutes of Health (National Institute

of Allergy and Infectious Diseases, AI43957) to A.H.A., J.R.E and T.E.W., and by funds from the

Scott-Waudby Trust, the Hope Against Cancer charity, Cancer Research UK in conjunction with

the UK Department of Health on an Experimental Cancer Medicine Centre grant

[C10604/A25151] to S.J, R.G.B and M.J.SD. Research at the University of Leicester was carried

out at the National Institute for Health and Care Research (NIHR) Leicester Biomedical Research

Centre (BRC). The authors also thank the Roy J. Carver Charitable Trust, Muscatine, Iowa for

ongoing research support.

REFERENCES

1.   Burger, J.A., et al., *Ibrutinib as Initial Therapy for Patients with Chronic Lymphocytic Leukemia.* N
Engl J Med, 2015. **373**(25): p. 2425-37.
2.   Davids, M.S. and J.R. Brown, *Ibrutinib: a first in class covalent inhibitor of Bruton's tyrosine kinase.*
Future Oncol, 2014. **10**(6): p. 957-67.
3.   Honigberg, L.A., et al., *The Bruton tyrosine kinase inhibitor PCI-32765 blocks B-cell activation and
is efficacious in models of autoimmune disease and B-cell malignancy.* Proc Natl Acad Sci U S A,
2010. **107**(29): p. 13075-80.
4.   Herman, S.E., et al., *Bruton tyrosine kinase represents a promising therapeutic target for
treatment of chronic lymphocytic leukemia and is effectively targeted by PCI-32765.* Blood, 2011.
117(23): p. 6287-96.
5.   Rozkiewicz, D., et al., *Bruton's Tyrosine Kinase Inhibitors (BTKIs): Review of Preclinical Studies and
Evaluation of Clinical Trials.* Molecules, 2023. **28**(5).
6.   Shirley, M., *Bruton Tyrosine Kinase Inhibitors in B-Cell Malignancies: Their Use and Differential
Features.* Target Oncol, 2022. **17**(1): p. 69-84.
7.   Montoya, S. and M.C. Thompson, *Non-Covalent Bruton's Tyrosine Kinase Inhibitors in the
Treatment of Chronic Lymphocytic Leukemia.* Cancers (Basel), 2023. **15**(14).
8.   Kueffer, L.E., R.E. Joseph, and A.H. Andreotti, *Reining in BTK: Interdomain Interactions and Their
Importance in the Regulatory Control of BTK.* Front Cell Dev Biol, 2021. **9**: p. 655489.
9.   Frustaci, A.M., et al., *Next Generation BTK Inhibitors in CLL: Evolving Challenges and New
Opportunities.* Cancers (Basel), 2023. **15**(5).
10.   Mendes-Bastos, P., et al., *Bruton's tyrosine kinase inhibition-An emerging therapeutic strategy in
immune-mediated dermatological conditions.* Allergy, 2022. **77**(8): p. 2355-2366.
11.   Robak, E. and T. Robak, *Bruton's Kinase Inhibitors for the Treatment of Immunological Diseases:
Current Status and Perspectives.* J Clin Med, 2022. **11**(10).
12.   Byrd, J.C., et al., *Acalabrutinib Versus Ibrutinib in Previously Treated Chronic Lymphocytic
Leukemia: Results of the First Randomized Phase III Trial.* J Clin Oncol, 2021. **39**(31): p. 3441-3452.
13.   Brown, J.R., et al., *Zanubrutinib or Ibrutinib in Relapsed or Refractory Chronic Lymphocytic
Leukemia.* N Engl J Med, 2023. **388**(4): p. 319-332.
14.   Lovell, A.R., N. Jammal, and P. Bose, *Selecting the optimal BTK inhibitor therapy in CLL: rationale
and practical considerations.* Ther Adv Hematol, 2022. **13**: p. 20406207221116577.
15.   Buske, C., et al., *Managing Waldenstrom's macroglobulinemia with BTK inhibitors.* Leukemia,
2023. **37**(1): p. 35-46.
16.   Joseph, R.E., et al., *Differential impact of BTK active site inhibitors on the conformational state of
full-length BTK.* Elife, 2020. **9**.
17.   Sonti, R., et al., *ATP Site Ligands Determine the Assembly State of the Abelson Kinase Regulatory
Core via the Activation Loop Conformation.* J Am Chem Soc, 2018. **140**(5): p. 1863-1869.
18.   Tong, M., et al., *Survey of solution dynamics in Src kinase reveals allosteric cross talk between the
ligand binding and regulatory sites.* Nat Commun, 2017. **8**(1): p. 2160.
19.   Skora, L., et al., *NMR reveals the allosteric opening and closing of Abelson tyrosine kinase by ATP-
site and myristoyl pocket inhibitors.* Proc Natl Acad Sci U S A, 2013. **110**(47): p. E4437-45.
20.   Sedlarikova, L., et al., *Resistance-Associated Mutations in Chronic Lymphocytic Leukemia Patients
Treated With Novel Agents.* Front Oncol, 2020. **10**: p. 894.
21.   Maddocks, K.J., et al., *Etiology of Ibrutinib Therapy Discontinuation and Outcomes in Patients With
Chronic Lymphocytic Leukemia.* JAMA Oncol, 2015. **1**(1): p. 80-7.

22.   Ahn, I.E., et al., *Clonal evolution leading to ibrutinib resistance in chronic lymphocytic leukemia.*
Blood, 2017. **129**(11): p. 1469-1479.
23.   Woyach, J.A., et al., *Resistance mechanisms for the Bruton's tyrosine kinase inhibitor ibrutinib.* N
Engl J Med, 2014. **370**(24): p. 2286-94.
24.   Blombery, P., et al., *Enrichment of BTK Leu528Trp mutations in patients with CLL on zanubrutinib:*
potential for pirtobrutinib cross-resistance. Blood Adv, 2022. **6**(20): p. 5589-5592.
25.   Furman, R.R., et al., *Ibrutinib resistance in chronic lymphocytic leukemia.* N Engl J Med, 2014.
370(24): p. 2352-4.
26.   Wang, E., et al., *Mechanisms of Resistance to Noncovalent Bruton's Tyrosine Kinase Inhibitors.* N
Engl J Med, 2022. **386**(8): p. 735-743.
27.   Jackson, R.A., et al., *BTK mutations in patients with chronic lymphocytic leukemia receiving*
tirabrutinib. Blood Adv, 2023. **7**(14): p. 3378-3381.
28.   Handunnetti SM, T.C., Nguyen T, et al., *BTK Leu528Trp - a Potential Secondary Resistance*
Mechanism Specific for Patients with Chronic Lymphocytic Leukemia Treated with the Next
Generation BTK Inhibitor Zanubrutinib. Blood., 2019. **134**: p. 170.
29.   Woyach, J.A., Huang, Y., Rogers, K., Bhat, S.A., et al., *Resistance to Acalabrutinib in CLL Is Mediated*
Primarily By BTK Mutations. Blood, 2019. **134**: p. 504.
30.   Woyach, J.A., et al., *BTK(C481S)-Mediated Resistance to Ibrutinib in Chronic Lymphocytic*
Leukemia. J Clin Oncol, 2017. **35**(13): p. 1437-1443.
31.   Nakhoda, S., A. Vistarop, and Y.L. Wang, *Resistance to Bruton tyrosine kinase inhibition in chronic*
lymphocytic leukaemia and non-Hodgkin lymphoma. Br J Haematol, 2023. **200**(2): p. 137-149.
32.   Kanagal-Shamanna, R., et al., *Targeted multigene deep sequencing of Bruton tyrosine kinase*
inhibitor-resistant chronic lymphocytic leukemia with disease progression and Richter
transformation. Cancer, 2019. **125**(4): p. 559-574.
33.   Gango, A., et al., *Dissection of subclonal evolution by temporal mutation profiling in chronic*
lymphocytic leukemia patients treated with ibrutinib. Int J Cancer, 2020. **146**(1): p. 85-93.
34.   Sharma, S., et al., *Identification of a structurally novel BTK mutation that drives ibrutinib resistance*
in CLL. Oncotarget, 2016. **7**(42): p. 68833-68841.
35.   Xie, Q., et al., *Substrate recognition of PLCgamma1 via a specific docking surface on Itk.* J Mol Biol,
2013. **425**(4): p. 683-96.
36.   Lin, D.Y. and A.H. Andreotti, *Structure of BTK kinase domain with the second-generation inhibitors*
acalabrutinib and tirabrutinib. PLoS One, 2023. **18**(8): p. e0290872.
37.   Gomez, E.B., et al., *Preclinical characterization of pirtobrutinib, a highly selective, noncovalent*
(reversible) BTK inhibitor. Blood, 2023. **142**(1): p. 62-72.
38.   Guo, Y., et al., *Discovery of Zanubrutinib (BGB-3111), a Novel, Potent, and Selective Covalent*
Inhibitor of Bruton's Tyrosine Kinase. J Med Chem, 2019. **62**(17): p. 7923-7940.
39.   Joseph, R.E., et al., *Achieving a Graded Immune Response: BTK Adopts a Range of Active/Inactive*
Conformations Dictated by Multiple Interdomain Contacts. Structure, 2017. **25**(10): p. 1481-1494
e4.
40.   Lin, D.Y., Kueffer, L.E., Juneja, P., Wales, T., et al., *Conformational heterogeneity of the BTK PHTH*
domain drives multiple regulatory states. eLife, 2023.
41.   Wang, Q., et al., *Autoinhibition of Bruton's tyrosine kinase (Btk) and activation by soluble inositol*
hexakisphosphate. Elife, 2015. **4**.
42.   Johnson, A.R., et al., *Battling Btk Mutants With Noncovalent Inhibitors That Overcome Cys481 and*
Thr474 Mutations. ACS Chem Biol, 2016. **11**(10): p. 2897-2907.
43.   Yuan, H., et al., *BTK kinase activity is dispensable for the survival of diffuse large B-cell lymphoma.*
J Biol Chem, 2022. **298**(11): p. 102555.

44. Dhami, K., et al., *Kinase-deficient BTK mutants confer ibrutinib resistance through activation of the*
kinase HCK. Sci Signal, 2022. **15**(736): p. eabg5216.
45. Moarefi, I., et al., *Activation of the Src-family tyrosine kinase Hck by SH3 domain displacement.*
Nature, 1997. **385**(6617): p. 650-3.
46. Pessoa, J., et al., *Editorial: Altered Expression of Proteins in Cancer: Function and Potential*
Therapeutic Targets. Front Oncol, 2022. **12**: p. 949139.
47. Bhullar, K.S., et al., *Kinase-targeted cancer therapies: progress, challenges and future directions.*
Mol Cancer, 2018. **17**(1): p. 48.
48. Cohen, P., D. Cross, and P.A. Janne, *Kinase drug discovery 20 years after imatinib: progress and*
future directions. Nat Rev Drug Discov, 2021. **20**(7): p. 551-569.
49. Bender, A.T., et al., *Ability of Bruton's Tyrosine Kinase Inhibitors to Sequester Y551 and Prevent*
Phosphorylation Determines Potency for Inhibition of Fc Receptor but not B-Cell Receptor
Signaling. Mol Pharmacol, 2017. **91**(3): p. 208-219.
50. Li, W., et al., *Bruton's Tyrosine Kinase Inhibitors With Distinct Binding Modes Reveal Differential*
Functional Impact on B-Cell Receptor Signaling. Mol Cancer Ther, 2023.
51. Yang, G., et al., *HCK is a survival determinant transactivated by mutated MYD88, and a direct*
target of ibrutinib. Blood, 2016. **127**(25): p. 3237-52.
52. Skanland, S.S. and A.R. Mato, *Overcoming resistance to targeted therapies in chronic lymphocytic*
leukemia. Blood Adv, 2021. **5**(1): p. 334-343.
53. Smith, C.I.E. and J.A. Burger, *Resistance Mutations to BTK Inhibitors Originate From the NF-kappaB*
but Not From the PI3K-RAS-MAPK Arm of the B Cell Receptor Signaling Pathway. Front Immunol,
2021. **12**: p. 689472.
54. Alexandra Chirino, S.M., Anita Safronenka and Justin Taylor, *Resisting the Resistance: Navigating*
BTK Mutations in Chronic Lymphocytic Leukemia (CLL). Genes, 2023. **14**: p. 2182.
55. Aslan, B., et al., *Pirtobrutinib inhibits wild-type and mutant Bruton's tyrosine kinase-mediated*
signaling in chronic lymphocytic leukemia. Blood Cancer J, 2022. **12**(5): p. 80.
56. Skanland, S.S. and G.E. Tjonnfjord, *Determining drug dose in the era of targeted therapies: playing*
it (un)safe? Blood Cancer J, 2022. **12**(8): p. 123.
57. Zorba, A., et al., *Delineating the role of cooperativity in the design of potent PROTACs for BTK.* Proc
Natl Acad Sci U S A, 2018. **115**(31): p. E7285-E7292.
58. Molica, S., *Venetoclax: a real game changer in treatment of chronic lymphocytic leukemia.* Int J
Hematol Oncol, 2020. **9**(4): p. IJH31.
59. Joseph, R.E., et al., *A remote substrate docking mechanism for the tec family tyrosine kinases.*
Biochemistry, 2007. **46**(18): p. 5595-603.
60. Johnson, B.A. and R.A. Blevins, *NMR View: A computer program for the visualization and analysis*
of NMR data. J Biomol NMR, 1994. **4**(5): p. 603-14.
61. Masson, G.R., et al., *Recommendations for performing, interpreting and reporting hydrogen*
deuterium exchange mass spectrometry (HDX-MS) experiments. Nat Methods, 2019. **16**(7): p. 595-
602.
62. Vizcaino, J.A., et al., *2016 update of the PRIDE database and its related tools.* Nucleic Acids Res,
2016. **44**(22): p. 11033.
63. Wales, T.E. and J.R. Engen, *Hydrogen exchange mass spectrometry for the analysis of protein*
dynamics. Mass Spectrom Rev, 2006. **25**(1): p. 158-70.
64. Advani, R.H., et al., *Bruton tyrosine kinase inhibitor ibrutinib (PCI-32765) has significant activity in*
patients with relapsed/refractory B-cell malignancies. J Clin Oncol, 2013. **31**(1): p. 88-94.
65. Podoll, T., et al., *Bioavailability, Biotransformation, and Excretion of the Covalent Bruton Tyrosine*
Kinase Inhibitor Acalabrutinib in Rats, Dogs, and Humans. Drug Metab Dispos, 2019. **47**(2): p. 145-
154.

66. Tam, C.S., et al., *Phase 1 study of the selective BTK inhibitor zanubrutinib in B-cell malignancies*
and safety and efficacy evaluation in CLL. Blood, 2019. **134**(11): p. 851-859.
67. Walter, H.S., et al., *A phase 1 clinical trial of the selective BTK inhibitor ONO/GS-4059 in relapsed*
and refractory mature B-cell malignancies. Blood, 2016. **127**(4): p. 411-9.
68. Mato, A.R., et al., *Pirtobrutinib in relapsed or refractory B-cell malignancies (BRUIN): a phase 1/2*
study. Lancet, 2021. **397**(10277): p. 892-901.

 FIGURES

 Figure 1:

Figure 1: (a) Domain organization of full-length (FL) BTK and the BTK linker-kinase domain

(LKD) fragment used in this study: PHTH, Pleckstrin homology-Tec homology domain; PRR,

proline-rich region; SH3, Src homology 3 domain; SH2, Src homology 2 domain, SH2-kinase

linker (L) and the catalytic kinase domain. Key residues are indicated above each domain. (b)

Autoinhibited conformation of FL BTK based on the crystal structure of FL BTK [40]. The PHTH

domain (purple) is dynamic, in transient contact with several regions on the core SH3-SH2-kinase

domain and is not visible in the crystal structure of full-length BTK [40]. Dynamics of the PHTH

domain is represented by the multiple poses of the PHTH domain and the double headed arrow.

(c) Co-crystal structure of BTK LKD (light cyan cartoon) bound to Ibrutinib (PDB ID: 5P9J)

showing the location of C481 (yellow spheres), T474 and L528 (red spheres) within the kinase

active site (broken oval). (c) Pie charts showing the prevalence of the BTK resistance mutations

in CLL patients treated with various BTK inhibitors. The total number of patients with mutations

in BTK are indicated below each chart. See Supp. Table 1 for additional details.

Figure 2:

Figure 2: The BTK kinase domain can interconvert between active and inactive conformations. (a)

Superposition of the structure of BTK linker-kinase domain (LKD) bound to Dasatinib (PDB ID:

3K54) in the active kinase conformation (grey cartoon) with the Ibrutinib bound structure (PDB

ID: 5P9J) in the inactive conformation (red cartoon). The expanded inset shows the inward

movement of the αC-helix, the change in W395 rotamer conformation and the K430/E445 salt

bridge formation that accompanies kinase activation. (b-f) Co-crystal structures of BTK LKD

(light cyan cartoon) bound to Ibrutinib (PDB ID: 5P9J), Acalabrutinib (PDB ID: 8FD9),

Zanubrutinib (PDB ID: 6J6M), Tirabrutinib (PDB ID: 5P9M) and Pirtobrutinib (PDB ID: 8FLL)

in the inactive kinase conformation. The inhibitors are shown as dark blue sticks, the kinase

activation loop is purple and C481, Y551 and W395 residues are shown as sticks with transparent

spheres. Electron density for part of the activation loop is missing in the Ibrutinib co-crystal

structure and is indicated as dotted lines (b). In the Acalabrutinib structure, the activation loop has

several mutations [36] and the SH2-kinase linker (including W395) is absent (c). Electron density

for W395 sidechain is missing in the BTK:Pirtobrutinib co-crystal structure (f). (g) Overlay of the

BTK:Ibrutinib, Acalabrutinib, Zanubrutinib, Tirabrutinib and Pirtobrutinib co-crystal structures.

With the exception of the Tirabrutinib co-crystal structure (e), no major structural variation is

observed in the kinase domains. The activation loop in the Tirabrutinib bound structure adopts a

different conformation compared to the other co-crystal structures.

Figure 3:

Figure 3: BTK inhibitors stabilize the inactive kinase conformation in solution. The tryptophan

side chain region of the ^{1}H-^{15}N TROSY HSQC spectra of ^{15}N-labelled apo BTK linker-kinase

domain (black spectrum) overlaid with that of the inhibitor bound spectrum (cyan spectrum). Here

and in subsequent figures, the broken black and grey lines indicate the position of the BTK W395

resonance in the active (αC-in) and inactive (αC-out) states respectively as shown in Figure 2a.

The shift in the BTK W395 resonance upon inhibitor binding is indicated by an arrow in each

spectrum. The structures of each inhibitor are shown on the right. The BTK W395 indole NH

resonance is in the inactive (αC-out) position in the Ibrutinib (published earlier [16]),

Acalabrutinib, Zanubrutinib and Tirabrutinib bound BTK LKD samples. Multiple peaks

corresponding to W395 are seen in the Tirabrutinib bound spectrum suggesting that the kinase

adopts multiple conformations in solution. The downfield shift observed in W395 in the

Pirtobrutinib bound structure is likely due to local changes in the chemical environment due to the

distinct chemical structure of Pirtobrutinib. W395 assignments in the inhibitor bound spectra were

confirmed by acquiring inhibitor bound spectra with the BTK LKD W395A mutant (see Supp.

Fig. S1).

Figure 4:

Figure 4: Assessing the impact of BTK inhibitors on full-length BTK by HDX-MS. (a) Intact mass

analysis of wild-type FL BTK before (bottom spectrum, black) and after one hour incubation with

a 2-fold molar excess of covalent BTK inhibitors: Ibrutinib (red), Acalabrutinib (purple),

Zanubrutinib (blue) and Tirabrutinib (orange) show a mass increase of one inhibitor molecule. (b)

Clinically approved BTK inhibitors induce allosteric changes in full-length BTK. Relative

deuterium level of peptides in apo BTK was subtracted from the deuterium level of the

corresponding peptide from each drug-bound form of BTK ($D_{WT\ drug-bound} - D_{WT\ apo}$) and the

differences colored according to the scale shown. In this and subsequent figures, peptic peptides are shown from N- to C-terminus, top to bottom, and the amount of time in deuterium is shown left to right. The relative difference data shown here represents a curated set of peptides that are coincident across all 6 states (apo and five drug-bound BTK forms). The identification of these chosen peptides, the relative difference values, and the complete data set for each state can be found in the Supplemental Datafile. The approximate position of the domains of BTK, as described in Figure 1a, is shown at the left. Deuterium incorporation curves of selected peptides (indicated with a gray box in panel b and labelled i-vi) from various regions of the protein are shown below. Data for Ibrutinib has been previously published [16].

Figure 5:

Figure 5: (a-e) Mapping the HDX-MS changes induced by each BTK inhibitor on the structure of

the BTK SH3-SH2-kinase fragment (PDB ID: 4XI2). Major differences greater than 1.0 Da are

shown as dark blue (decrease) or dark green (increase); modest differences between 0.5 Da and

1.0 Da are shown as light blue (decrease) and light green (increase). Localization of the changes

in deuterium incorporation was accomplished using overlapping peptides included in the complete

peptide data set provided in the Supplemental Datafile. The location of peptides i – vi from Figure

4 are indicated in panel a. Data corresponding to Ibrutinib has been previously published [16].

Figure 6:

Figure 6: Probing the impact of the BTK resistance mutations T474I and L528W on BTK. (a-d)

Western blot comparing the kinase activity of full-length (FL) BTK wild-type (WT), T474I and

L528W mutants. BTK autophosphorylation was monitored using the BTK pY551 antibody and

the total protein levels monitored using the Anti-His antibody. (b, d) Histogram quantifying the

western blots shown in (a and c). The blots were quantified and normalized as described in the

Materials and Methods. Data shown are the average of three independent experiments. (e) HDX

difference data for the BTK T474I and L528W mutants ($D_{Mutant\ apo}$ -$D_{WT\ apo}$). Color scale and

peptide/time course arrangement are the same as in Figure 4. See the Supplemental Datafile for

additional information, including all peptide identifications and deuterium values. (f,g) Mapping

the mutational induced HDX-MS changes on the structure of the BTK SH3-SH2-kinase fragment. (h) Deuterium incorporation curves of selected peptides (indicated with a gray box in panel e and labelled i-iv) from various regions of the protein are shown. (i) The tryptophan side chain region of the ^{1}H-^{15}N TROSY HSQC spectra of ^{15}N-labelled apo WT BTK linker-kinase domain (black spectrum) overlaid with that of the apo mutant kinase spectrum (cyan spectrum). The boxed region in the T474I spectral overlay is expanded above.

Figure 7:

Figure 7: The BTK L528W mutant can activate HCK. (a-f) Kinase activity of HCK in the presence

or absence of full-length BTK L528W mutant was compared in a western blot assay by monitoring

PLCγ1 phosphorylation (pY783 antibody) and HCK autophosphorylation (pY antibody). Total

protein levels monitored using the Anti-His antibody. Full-length WT BTK preincubated with

Zanubrutinib was used as a control. (b, d and f) Histogram quantifying the western blots shown in

(a, c and e). The blots were quantified and normalized as described in the Materials and Methods.

Data shown are the average of three independent experiments. (b and c) Kinase activity of HCK

in the presence or absence of the isolated linker kinase domain (LKD) fragment of the BTK L528W

mutant (b) or the full-length proline mutant of BTK L528W (BTK FL L528W/Pro: BTK L528W/

P189A/P192A/P203A/P206A, (c)) was compared as in (a). (g) Thermal stability analysis of BTK

FL WT and BTK FL L528W. Data shown are the average of three independent experiments.

Figure 8:

Figure 8: HDX-MS analysis of BTK inhibitor binding to BTK T474I and L528W mutants. (a-d)

Relative deuterium level of peptides in apo mutant BTK was subtracted from the deuterium level

of the corresponding peptide from each inhibitor-bound form of BTK ($D_{drug-bound}$-D_{apo}) and

compared to the changes in the WT protein. The differences are colored according to the scale

shown. (e) Deuterium incorporation curves of selected peptides (indicated with a gray box in

panels a-c, and labelled i-iv) are shown.

    Figure 9:

Figure 9: NMR analysis of BTK inhibitor binding to the BTK T474I and L528W mutants. The tryptophan side chain region of the ^{1}H-^{15}N TROSY HSQC spectra of ^{15}N-labelled apo BTK linker-kinase domain (black spectrum) overlaid with that of the inhibitor bound spectrum (cyan spectrum). The broken black and grey lines indicate the position of the BTK W395 resonance and have been described earlier in Figure 3. The shift in the BTK W395 indole NH resonance upon

998   inhibitor binding is indicated by an arrow in each spectrum. The red asterisks indicate the presence

999   of unbound kinase domain in the inhibitor bound NMR sample.

1000

1001   TABLES

1002   **Table 1**: Half-life of clinically approved BTK inhibitors: Ibrutinib[64], Acalabrutinib [65],

1003   Zanubrutinib [66], Tirabrutinib [67] and Pirtobrutinib [68].

Inhibitor	Half life (h)
Ibrutinib	2 - 3
Acalabrutinib	0.6 - 2.8
Zanubrutinib	4
Tirabrutinib	6.5 - 8
Pirtobrutinib	20

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014   Supp. Fig. S1

1015

1016

Supp Fig. S1: Assignment of W395 in inhibitor bound spectra of BTK LKD. The tryptophan side

chain region of the ^{1}H-^{15}N TROSY HSQC spectra of ^{15}N-labelled inhibitor bound BTK linker-

kinase domain WT (black spectrum) overlaid with that of the inhibitor bound BTK LKD W395A

spectrum (cyan spectrum). The boxed peak indicates the W395 resonance in each of the WT

inhibitor bound spectrum.

Supplemental Datafile:

Microsoft Excel file providing enhanced experimental details for HDX-MS including minimum

criteria specified by [61], lists of all peptides by residue number, sequence, as well as deuterium

levels measured for each Figure. The value of each deuterium difference for every colored box in

each Figure as well as the complete dataset for each state are also found in this file.

1027

1028 **Supp. Table 1**: BTK mutations detected in CLL patients treated with BTK inhibitors. In addition

1029 to mutations in BTK C481, T474 and L528, other BTK mutations that have been detected include:

1030 R28S, E108K, G164D, V416L, A428D, M437R, R490H, Q516K, V537I and T316A [21-24, 26-

1031 30, 32-34].

BTK inhibitor	Total number of patients with BTK mutations.	Patients with BTK C481 mutated.	Patients with BTK T474 mutated.	Patients with BTK L528 mutated.	Patients with other BTK mutations.
Ibrutinib	106	95 (90%)	1 (1%)	1 (1%)	5 (5%)
Acalabrutinib	11	11 (100%)	1 (9%)	0	0
Zanubrutinib	17	15 (88%)	0	11 (65%)	0
Tirabrutinib	6	4 (67%)	3 (50%)	4 (67%)	1 (17%)
Pirtobrutinib	9	0	2 (22%)	4 (44%)	3 (33%)

1032