1

Impact of the clinically approved BTK inhibitors on the conformation of full-length BTK and analysis of the development of BTK resistance mutations in chronic lymphocytic leukemia.

Raji E. Joseph[#*a], Thomas E. Wales[#b], Sandrine Jayne, Robert G. Britton, D. Bruce Fulton[a],

John R. Engen[b], Martin J. S. Dyer[c] and Amy H. Andreotti[*a]

[a]Roy J. Carver Department of Biochemistry, Biophysics and Molecular Biology, Iowa State University, Ames, IA 50011, USA.

[b]Department of Chemistry and Chemical Biology, Northeastern University, Boston, MA 02115, USA.

[c]The Ernest and Helen Scott Haematological Research Institute, Leicester Cancer Research Centre, College of Life Sciences, University of Leicester, Leicester LE1 9HN, UK.

[#]These authors contributed equally to this work

[*]Co-corresponding author emails: jraji@iastate.edu, amyand@iastate.edu

ABSTRACT

Inhibition of Bruton's tyrosine kinase (BTK) has proven to be highly effective in the treatment of B-cell malignancies such as chronic lymphocytic leukemia (CLL), autoimmune disorders and multiple sclerosis. Since the approval of the first BTK inhibitor (BTKi), Ibrutinib, several other inhibitors including Acalabrutinib, Zanubrutinib, Tirabrutinib and Pirtobrutinib have been clinically approved. All are covalent active site inhibitors, with the exception of the reversible active site inhibitor Pirtobrutinib. The large number of available inhibitors for the BTK target creates challenges in choosing the most appropriate BTKi for treatment. Side-by-side comparisons in CLL have shown that different inhibitors may differ in their treatment efficacy. Moreover, the nature of the resistance mutations that arise in patients appears to depend on the specific BTKi administered. We have previously shown that Ibrutinib binding to the kinase active site causes unanticipated long-range effects on the global conformation of BTK (Joseph, R.E., et al., 2020, https://doi.org/10.7554/eLife.60470). Here we show that binding of each of the five approved BTKi to the kinase active site brings about distinct allosteric changes that alter the conformational equilibrium of full-length BTK. Additionally, we provide an explanation for the resistance mutation bias observed in CLL patients treated with different BTKi and characterize the mechanism of action of two common resistance mutations: BTK T474I and L528W.

Keywords**:** BTK, Bruton's Tyrosine Kinase; BTK inhibitors; CLL, chronic lymphocytic leukemia; Ibrutinib; Acalabrutinib; Zanubrutinib; Tirabrutinib; Pirtobrutinib; Fenebrutinib; drug resistance; allostery; conformational equilibrium; HDX-MS, hydrogen/deuterium exchange mass spectrometry and NMR, Nuclear Magnetic Resonance.

52   Abbreviations: BTKi, BTK inhibitor; LKD, linker-kinase domain; CLL, chronic lymphocytic

53   leukemia; MCL, Mantle cell lymphoma; MZL, Marginal zone lymphoma; PRR, proline-rich

54   region; HDX-MS, hydrogen/deuterium exchange-mass spectrometry; NMR, Nuclear Magnetic

55   Resonance; WT, wild-type.

 INTRODUCTION

Ibrutinib (IMBRUVICA®), has revolutionized the treatment of the B cell malignancy,

CLL [1]. Ibrutinib is a covalent active site inhibitor of the multi-domain B-cell kinase, BTK (Fig.

1a), that was first approved by the FDA in 2013 [2]. Inhibition of BTK disrupts signaling

downstream of the B-cell receptor (BCR), a pathway on which the survival of CLL cells are

dependent [3, 4]. The success of Ibrutinib has spurred the development of other BTKi including

the now clinically approved BTK inhibitors: Acalabrutinib (CALQUENCE®), Zanubrutinib

(BRUKINSA®), Tirabrutinib/ONO-4059 (VELEXBRU®), Pirtobrutinib/LOXO-405

(JAYPIRCA®) and Orelabrutinib (HIBRUKA®) [5-9]. All of the clinically approved BTKi (with

the exception of Pirtobrutinib) are covalent active site inhibitors that bind to BTK C481 residue

within the kinase active site. Pirtobrutinib is currently the first and only clinically approved non-

covalent BTK active site inhibitor [7]. Importantly, all clinically approved BTK inhibitors in this

study, covalent and non-covalent, completely block nucleotide binding.

While BTKi are highly effective in the treatment of CLL, they are also being used to treat

other B-cell malignancies such as Mantle cell lymphoma (MCL), Waldenström's

macroglobulinemia, Marginal zone lymphoma (MZL) and are being evaluated in the treatment of

multiple sclerosis and rheumatoid arthritis [5, 10, 11]. The plethora of clinically approved BTKi

along with several other promising candidates currently in clinical trials pose a new challenge

with respect to patient treatment. We now need a way to assess the suitability of a given BTKi

for a given patient or disease state. Indeed, recent clinical trial data comparing one BTKi to

another indicate that BTKi differ in their effectiveness in treating various conditions [12-15].

While toxicity, specificity and other criteria are used in these clinical comparisons, a molecular

level characterization of the interaction of the inhibitor with full-length BTK is lacking and could

be key to understanding these differences. In our previous work, using high resolution structural

biology techniques such as Nuclear Magnetic Resonance (NMR) and Hydrogen Deuterium

Exchange Mass Spectrometry (HDX-MS), we have shown that some inhibitors (Ibrutinib) exert

unanticipated allosteric effects upon binding to BTK while others (Fenebrutinib) do not [16].

Ibrutinib binding to the BTK kinase active site leads to a shift in the conformational ensemble of

full-length BTK towards an open conformation where the regulatory SRC homology (SH3 and

SH2) domains are released from the 'back' surface of the kinase domain, disrupting the closed

autoinhibited conformation of the full-length protein (Fig. 1b, [16]). Allosteric conformational

changes upon drug binding have been shown to alter protein-ligand interactions in other systems

with functional consequences *in vivo* [17-19]. The full impact of the panel of currently approved

BTKi on the conformation of full-length BTK is not known.

Despite the success of BTKi in CLL, a recurring problem is the development of BTKi

resistance. Resistance typically develops in patients ~2 years after the start of BTK inhibitor

treatment [20]. Analysis of CLL patients that stop responding to BTKi has revealed that most

develop mutations within *Btk* or in the substrate of BTK: Phospholipase C gamma 2 (PLCγ2)

[21-23]. Interestingly, the specific resistance mutations that develop in *Btk* seem to be dependent

on the specific BTKi used [23-29]. While data available so far for the BTKi inhibitors

Acalabrutinib, Zanubrutinib, Tirabrutinib and Pirtobrutinib are low in number, there is an

emerging trend (Fig. 1c,d). Perhaps not surprisingly, patients treated with the reversible inhibitor

Pirtobrutinib do not develop mutations in C481 (Fig. 1d). Instead, they develop other kinase

active site mutations including BTK T474I and L528W (Fig. 1c,d) [26]. In contrast, 90% or

more of CLL patients treated with Ibrutinib and Acalabrutinib develop mutations in BTK C481

(Fig. 1d, Supp. Table 1 [21, 23]). Substitution of C481 with serine is the most common

resistance mutation found in these patients [30]. While mutation of BTK C481 to residues other

than serine have also been reported (C481F/Y/R), they occur at a much lower frequency [31].

Additionally, other sites within BTK such as T474 and L528 are rarely (if ever) mutated in

Ibrutinib and Acalabrutinib-treated CLL patients (Fig. 1d, Supp. Table 1, [21-24, 29 , 30, 32-

34]). Surprisingly, BTK T474 and L528 are frequently mutated in CLL patients treated with the

covalent inhibitors Zanubrutinib and Tirabrutinib (Fig. 1d, [24, 27, 28]). BTK T474I and L528W

are found at almost equal or higher frequency compared to C481S in these patients (Fig. 1d,

Supp. Table 1, [24, 27, 28]). Moreover, in Zanubrutinib-treated patients, the L528W mutation

was often present together with C481S mutation (on different alleles), and with the L528W

mutation present at a higher allelic frequency compared to C481S [28]. This mutational bias with

respect to covalent BTKi is unexpected given the shared mode of action across the panel of

covalent inhibitors, and the reasons for these differences are unclear. Additionally, the

mechanism of action of the BTK T474I and L528W mutations is not known. Given the

development of T474I and L528W mutations in multiple CLL patients treated with both covalent

(Zanubrutinib and Tirabrutinib) and non-covalent (Pirtobrutinib) BTK inhibitors, we focused on

these BTK mutations and explore their mechanisms of action.

Here, we probe the impact of four clinically-approved BTKi: Acalabrutinib,

Zanubrutinib, Tirabrutinib and Pirtobrutinib on the conformation of full-length BTK. We find

that each of these BTKi brings about a unique combination of changes in full-length BTK.

Acalabrutinib, Zanubrutinib and Tirabrutinib disrupted the autoinhibited conformation of full-

length BTK similarly to Ibrutinib but did so to varying degrees. Interestingly, Acalabrutinib and

Tirabrutinib altered the dynamics of the kinase G-helix, a region that has been previously

characterized as the PLCγ substrate docking site [35]. Pirtobrutinib on the other hand stabilized

the compact autoinhibited conformation of full-length BTK and is the first BTK inhibitor

observed to do so. Additionally, we probed the mechanism of action of the BTK resistance

mutations T474I and L528W. We show that the catalytically inactive BTK L528W mutant

activated the SRC kinase HCK and that this activation is dependent on the proline-rich region

within BTK. The BTK T474I mutation disrupted binding to Zanubrutinib, Tirabrutinib and

Pirtobrutinib and likely evades the action of these drugs due to reduced binding to these

inhibitors. Furthermore, we provide an explanation for the mutational bias observed in patients

treated with different covalent BTKi. The development of the C481S resistance mutation is

dependent on the half-life of the inhibitor which likely explains the low prevalence of the C481S

resistance mutation in patients treated with the covalent inhibitors Tirabrutinib and Zanubrutinib

(which have a long half-life) as compared to Ibrutinib and Acalabrutinib (which have a short

half-life). Characterization of the interaction of BTKi with full-length BTK allows us to better

interpret clinical trial results and will help guide the choice of BTKi to be used for treatment.

Furthermore, understanding the mechanism of action of resistance mutations allows us to

develop treatment strategies that either prevent or delay development of resistance mutations and

ways to treat them when they arise.

RESULTS

Assessing the impact of inhibitor binding on the isolated BTK linker kinase domain by NMR.

To evaluate the impact of inhibitor binding on BTK, we first monitored its effect on the

conformation of the linker kinase domain (LKD) fragment of BTK (Fig. 1a). The catalytic kinase

domain of BTK can adopt an active or inactive conformation in solution and previous studies

have shown that inhibitor binding can stabilize one or more of these conformations [16]. The

switch from an inactive to an active kinase conformation involves changes in key structural

elements within the kinase domain. These changes include the inward movement of the C-helix

from an inactive 'αC out' to the active 'αC in' position, changes in the side chain rotamer

conformation of BTK W395, formation of a conserved BTK K430:E445 salt bridge and

unfurling of the collapsed activation loop, leading to the exposure of the conserved Y551 on the

activation loop for phosphorylation (Fig. 2a, [16]).

Crystal structures of the BTK kinase domain in complex with Acalabrutinib,

Zanubrutinib, Tirabrutinib or Pirtobrutinib are available and show that each of these BTK

inhibitors stabilize the kinase domain in an inactive conformation similar to that of Ibrutinib-

bound BTK (Fig. 2b-f, [16, 36-38]). In all the inhibitor bound structures the C-helix is in the 'αC

out' position and the activation loop is collapsed into the kinase active site, burying the

conserved BTK Y551 (Fig. 2b-f). Moreover, superposition of these BTK/inhibitor complexes

shows that the structures of the kinase domain remains largely the same regardless of which

inhibitor is bound (Fig. 2g). The structure of the Tirabrutinib bound BTK is the only structure

that shows a difference; the activation loop in that complex adopts an alternative inactive loop

conformation compared to the other inhibitor bound structures (Fig. 2e,g).

Previous NMR and HDX-MS analysis of the interaction of BTK with a different panel of

active site inhibitors has shown that there are differences between the solution and crystal

behavior [16]. To test whether the BTK inhibitors Acalabrutinib, Zanubrutinib, Tirabrutinib and

Pirtobrutinib can stabilize an inactive BTK kinase conformation in solution as predicted by the

crystal structures, we evaluated the BTK inhibitor bound complexes by NMR. The LKD

fragment of BTK was isotopically labeled with ^{15}N and a ^{1}H-^{15}N TROSY-HSQC spectrum was

obtained in the presence or absence of the inhibitor (Fig. 3). We have previously shown that

BTK W395 within the linker region (L) that precedes the kinase domain provides a useful probe

to monitor the conformational state adopted by the kinase domain in solution [16, 39].

Assignments for the apo BTK linker kinase domain, which adopts the active (αC in)

conformation, show that the W395 indole ^{1}H resonates at 10.21 ppm in the ^{1}H-^{15}N TROSY-

HSQC spectrum (Fig. 3, top panel, black spectrum). An upfield shift in the BTK W395 side

chain indole NH resonance is observed upon Ibrutinib binding and is consistent with the outward

movement of the C-helix (αC out), and stabilization of the inactive kinase domain conformation

by Ibrutinib (Fig.3, top panel, cyan spectrum [16]).

Comparing the tryptophan indole region of the ^{1}H-^{15}N TROSY-HSQC spectra for the

BTK LKD inhibitor bound to Acalabrutinib, Zanubrutinib or Tirabrutinib with that of the apo

BTK LKD protein shows that W395 undergoes an upfield shift in the presence of inhibitor (Fig.

3), suggesting that, like Ibrutinib [16] and consistent with the crystal structures, all of these

inhibitors stabilize the inactive kinase domain conformation in solution. Interestingly, the

Tirabrutinib bound spectrum shows several peaks corresponding to W395, suggesting that the

Tirabrutinib-bound BTK kinase domain is likely adopting multiple kinase conformations in

solution. Additionally, the magnitude of the upfield shift in W395 resonance is smaller in both

the Tirabrutinib and Zanubrutinib spectra (compared to Ibrutinib and Acalabrutinib), which may

reflect a relatively larger active state population in these samples.

In contrast to the covalent inhibitors, Pirtobrutinib causes a downfield shift in W395,

which might suggest that this inhibitor is stabilizing an active conformation of BTK (Fig. 3,

bottom panel). However, the fluorinated benzene ring in Pirtobrutinib, which is adjacent to the

C-helix and W395 in the Pirtobrutinib bound structure, may cause changes in the local

environment of W395 and thereby give rise to the unusual chemical shift change. Indeed, HDX-

MS data (see below) confirms the stabilization of the inactive kinase conformation by

Pirtobrutinib. All W395 assignments in the WT BTK inhibitor bound spectra were confirmed by

comparison to the corresponding ^{1}H-^{15}N TROSY-HSQC spectra of BTK LKD W395A mutant

bound to the inhibitors (Supp. Fig. S1).

Probing the effects of inhibitor binding on full-length BTK by HDX-MS.

We next evaluated the impact of inhibitor binding on the conformation of full-length

BTK by HDX-MS. We have previously shown that HDX-MS can be used to probe

conformational changes in full-length BTK that are brought about by inhibitor binding [16]. The

structure of the full-length BTK protein in the autoinhibited conformation was recently solved

[40]. However, electron density for both the N-terminal PHTH domain and proline rich region is

missing in this structure, suggesting that the N-terminal region of BTK is dynamic. Indeed,

CryoEM, SAXS and solution data all indicate that the PHTH domain is highly dynamic and

likely transiently contacts multiple sites on the core SH3-SH2-kinase domain (Fig. 1b, [40]).

Changes in deuterium incorporation must therefore be mapped on the crystal structure of the

SH3-SH2-kinase fragment of BTK, which adopts a compact inactive autoinhibited conformation

(Fig. 1b, [41]). Additionally, our previous work [16] has shown that for the majority of the

inhibitor complexes studied changes in deuterium incorporation are rarely observed in the PHTH

domain suggesting that the PHTH domain dynamics are not affected by active site occupancy.

Inhibitor binding typically leads to decreased deuterium incorporation in peptides derived from

the kinase active site due to stabilization of this region in the presence of the bound inhibitor

[16]. Additionally, inhibitors such as Ibrutinib that have allosteric effects lead to an increase in

deuterium incorporation in peptides derived from the BTK SH3 and SH2 domains as well as the

SH2-kinase linker, indicating a shift away from the autoinhibited conformation [16].

Importantly, HDX-MS provides information about the conformational average adopted by the

ensemble and in most cases cannot distinguish between individual intermediate conformations.

Full-length BTK was mixed with Acalabrutinib, Zanubrutinib, Tirabrutinib or

Pirtobrutinib and subjected to HDX-MS analysis. Peptides that could be followed in each of the

six experimental conditions (apo, and inhibitor bound) were used for comparison (see

Supplemental Datafile). Intact mass analysis supports a single binding site on BTK for the

covalent inhibitors similar to what has been observed previously with Ibrutinib (Fig. 4a, [16]).

Acalabrutinib, Zanubrutinib, Tirabrutinib and Pirtobrutinib show decreased deuterium

incorporation in peptides derived from the kinase N-lobe and kinase activation loop similar to

that observed previously with Ibrutinib (Fig. 4b,c, Fig. 5a-e, [16]). This is consistent with the

binding of these inhibitors in the kinase active site and stabilization of the inactive kinase

conformation observed by NMR as well as in the crystal structures. Furthermore, we note that

each of these inhibitors induce allosteric effects. Binding of Acalabrutinib, Zanubrutinib and

Tirabrutinib to BTK causes increased deuterium incorporation in peptides derived from the SH3

domain suggesting that the autoinhibited conformation of full-length BTK is destabilized (Fig.

4b,c, Fig. 5b-d). This is similar to what has been previously observed with Ibrutinib [16].

Additionally, Acalabrutinib and Tirabrutinib showed increased deuterium incorporation in

peptides derived from the G-helix of the kinase domain, a region that has been previously

identified as the PLCγ substrate docking site (Fig. 4b,c, Fig. 5b-d, [35]). Taken together,

Acalabrutinib, Zanubrutinib and Tirabrutinib binding to BTK leads to hybrid conformations of

full-length BTK, where the kinase domain is stabilized in an inactive conformation and the

regulatory domains are disrupted from their autoinhibitory conformation. In stark contrast,

Pirtobrutinib binding to BTK shows decreased deuterium incorporation in peptides derived from

the SH3 and SH2 domains, suggesting that Pirtobrutinib stabilizes the compact, autoinhibited

conformation of full-length BTK (Fig. 4b,c, Fig. 5e). To date, Pirtobrutinib is the only inhibitor

that we have tested that uniformly stabilizes both the kinase domain as well as the regulatory

domains in the inactive autoinhibited conformation. Thus, each BTK inhibitor causes unique

changes in the overall conformation of full-length BTK that is not readily predicted from crystal

structures. We next turn our attention to the distinct resistance mutations that arise upon

treatment with the different BTK inhibitors.

Probing the intrinsic effects of the BTK T474I and L528W mutations.

BTK resistance mutations can potentially confer a selective advantage to cells in several

ways: by increasing the activity of the kinase, by changing the conformation/stability of the

kinase which in turn can alter protein-protein interactions, or by disrupting drug binding. To

probe the mechanism by which the BTK T474I and L528W mutations confer resistance to

Zanubrutinib, Tirabrutinib and Pirtobrutinib we first tested the impact of each mutation on the

catalytic activity of the kinase and then investigated the impact on the overall conformation of

the protein by HDX-MS and NMR.

To test the catalytic activity of the BTK mutants we set up *in vitro* kinase assays using

purified full-length WT and mutant BTK proteins. Kinase activity was monitored by following

phosphorylation on the activation loop Y551 by western blotting. The BTK T474I mutant shows

phosphorylation on Y551, however it is lower than that of the WT protein (Fig.6a,b) suggesting

that the T474I mutation reduces the activity of the kinase. In contrast, the BTK L528W mutant is

completely inactive with no detectable phosphorylation on Y551 throughout the time course

(Fig.6c,d). These results are consistent with previous activity reports on both the BTK T474I and

L528W mutants [26, 28, 42, 43]. Given that neither the BTK T474I or the L528W mutant had

increased catalytic activity, augmenting kinase activity is not the mechanism by which these

mutations confer resistance.

Resistance mutations can change the conformation of the protein which in turn can alter

protein-protein interactions. To test whether the BTK T474I and L528W mutations alter the

overall conformation of the protein, we carried out HDX-MS analysis on the apo mutant

proteins. Comparing the BTK T474I mutant to WT BTK, the BTK T474I mutant shows

increased deuterium incorporation within the kinase domain N-lobe β2-β3 strands and the

activation loop, but no changes elsewhere on the protein (Fig.6e,f,h). These increased dynamics

within the N-lobe of the BTK T474I mutant could potentially alter drug binding to the active

site. The BTK L528W mutant on the other hand shows a decrease in deuterium incorporation in

the kinase domain and the SH3 domain suggesting that the L528W mutation has a stabilizing

effect on the full-length autoinhibited conformation of BTK (Fig.6e,g,h).

Additionally, we tested the BTK T474I and L528W mutants by NMR. An overlay of the

^{1}H-^{15}N TROSY-HSQC spectra of apo BTK LKD T474I with that of apo WT BTK LKD shows

some changes in the mutant spectrum (Fig.6i). The resonance corresponding to BTK A524

located close (~18 □) to T474 in the kinase active site shows line broadening in the BTK T474I

mutant spectrum suggesting increased dynamics in the mutant. This is consistent with the

increased deuterium incorporation observed in the kinase domain for this mutant by HDX-MS.

The BTK W395 resonance in the BTK T474I mutant spectrum overlaps with that of the WT

suggesting that the mutant kinase domain adopts an active conformation similar to that of the

WT protein. The BTK L528W mutant on the other hand shows a small upfield shift in the W395

resonance, suggesting that the BTK L528W mutant kinase domain is shifted towards an inactive

conformation compared to the WT protein. This change in the BTK L528W mutant is consistent

with the HDX-MS changes for this protein which suggest that the mutation has a stabilizing

effect on the compact autoinhibited conformation. Additionally, a new peak is observed as

expected in the BTK L528W mutant spectrum consistent with the introduction of an additional

tryptophan due to the mutation. Overall, as the BTK T474I and L528W mutations by themselves

cause only minor conformational changes in the nucleotide free apo form, such changes are

unlikely to constitute the mechanism by which these mutations confer resistance.

BTK L528W mutant activates the SRC family kinase HCK.

Previous studies have shown that the BTK L528W mutant propagates BCR signaling

despite being catalytically dead [26, 43]. PLCγ phosphorylation and calcium signaling are

maintained in cells carrying this BTK mutation [26, 43]. This suggests that the BTK L528W

mutant might recruit other kinases to compensate for its lack of activity. The mechanism of

action of another catalytically inactive BTK resistant mutation: BTK C481F/Y, which arises in

Ibrutinib treated CLL patients, has been recently reported [44]. In that work the BTK C481F/Y

resistance mutant was shown to bind and activate the SRC family kinase HCK, thereby

propagating signals from the BCR signaling pathway [44]. This activation of HCK by BTK

C481F/Y requires phosphorylation on the BTK kinase activation loop, Y551. The

phosphorylated Y551 is suggested to bind to the HCK SH2 domain, displacing the autoinhibited

conformation of HCK, thereby activating the HCK catalytic function. To determine if a similar

mechanism is at work for the catalytically dead BTK L528W mutant, we tested the ability of the

BTK L528W mutant to activate HCK in a western blot assay by monitoring the phosphorylation

levels on two substrates: PLCγ1 (pY783 phosphorylation) and HCK itself (pY levels). In

addition, we also included BTK WT that had been pre-incubated with Zanubrutinib as a control.

As shown in Fig.7a,b, phosphorylation increased on both PLCγ and HCK in the presence of full-

length BTK L528W, suggesting that the catalytically dead BTK mutant is able to activate HCK.

Interestingly, Zanubrutinib bound WT BTK also promoted increased phosphorylation levels on

both PLCγ1 and HCK suggesting that drug-inactivated BTK also activates HCK. This is distinct

from what has been previously observed for the BTK C481Y/F resistant mutation where,

compared to WT BTK, the C481Y/F mutant BTK preferentially engages HCK. Nevertheless,

since the BTK L528W mutant activates HCK, we next designed experiments to probe the

requirements for this activation.

Activation of HCK by BTK C481F/Y requires phosphorylation on the BTK kinase

activation loop, Y551. Therefore, we first tested whether phosphorylation on Y551 is required

for activation of HCK by the BTK L528W mutant. While BTK cannot autophosphorylate itself

on BTK Y551 in this assay (BTK FL WT is inhibited by Zanubrutinib and BTK FL L528W is

catalytically dead), both proteins are able to be phosphorylated by HCK on BTK Y551.

Interestingly, phosphorylation on BTK Y551 is lower in the BTK L528W mutant compared to

the WT BTK. This is likely due to the BTK kinase activation loop Y551 being less accessible to

phosphorylation by HCK in the mutant (which adopts the autoinhibited conformation) compared

to WT BTK (Fig. 7a). Despite the lower pY551 level on BTK FL L528W mutant compared to

BTK FL WT, the BTK FL L528W mutant is able to activate HCK similar to the BTK WT

protein, suggesting that phosphorylation levels on BTK Y551 do not impact HCK activation.

Additionally, we tested whether the BTK L528W/Y551F double mutant is able to activate HCK.

The BTK L528W/Y551F double mutant is able to activate HCK similar to BTK L528W,

suggesting that phosphorylation on Y551 is not required for HCK activation by BTK L528W

(Supp. Fig. S2a). Furthermore, BTK FL L528W increased phosphorylation on HCK and PLCγ1

even in the absence of Zanubrutinib (Fig. S2b) showing that drug interactions are not responsible

for the increased phosphorylation on HCK and PLCγ1.

We next tested whether the regulatory domains of BTK were required for HCK activation

by the BTK L528W mutant by using the isolated linker kinase domain of BTK (a construct in

which the BTK regulatory domains have been deleted). As shown in Fig. 7c,d, the isolated linker

kinase domain of BTK L528W does not activate HCK. This suggests that the regulatory

domain/s of BTK are required for the activation of HCK and that the isolated linker kinase

domain of BTK alone is insufficient for HCK activation. This again is different from what has

been previously reported; the BTK C481Y/F resistant mutant requires the kinase domain for

HCK activation [44]. We therefore turned our focus onto the BTK regulatory domains to isolate

the specific region/s required for HCK activation by the BTK L528W resistance mutation.

Displacement of the SH3 domain from its autoinhibitory conformation upon binding of a

proline-rich ligand is a classic mechanism by which SRC family kinases such as HCK are

activated [45]. BTK contains such a proline-rich ligand sequence within its regulatory region and

so we tested HCK activation by a BTK mutant lacking the proline-rich sequence (BTK

L528W/Pro: BTK L528W/ P189A/P192A/P203A/P206A). As shown in Fig. 7e,f, the BTK

L528W proline-depleted mutant does not activate HCK. This data again emphasizes the

difference between the mechanism of action of the BTK L528W mutant from the previously

characterized BTK C481Y/F mutant. Additionally, this data is consistent with the observation

that WT BTK can also activate HCK, as the proline-rich region is present in both WT and mutant

L528W BTK. The ability of both WT and L528W mutant BTK to activate HCK, however,

suggests that there must be additional reasons for the selection of the BTK L528W resistance

mutation in patients.

BTK L528W is more stable than WT BTK.

Changes in protein levels have been shown to cause resistance in several cancers [46].

Cellular protein levels can be altered due to changes in protein expression, stability or

degradation, or changes at the RNA level [46]. An increase in BTK protein levels *in vivo* could

drive protein-protein interactions typically not observed in the WT background. Interestingly, we

note that protein yield after bacterial expression and purification of the full-length BTK L528W

mutant is higher than that of the WT protein, suggesting that the L528W mutant is more stable

than the WT protein. To test whether the BTK L528W mutation has a stabilizing effect on BTK,

we measured the melting temperature (Tm) of the BTK L528W mutant and compared it to that

of the WT protein in a thermal shift assay. As shown in Fig.7g, the BTK L528W mutant is more

stable than the WT protein (a one degree increase in Tm compared to the WT). The increased

stability of the L528W mutant, albeit small, could result in increased protein levels, an increase

in the lifetime of the protein or altered protein trafficking *in vivo*, all of which would drive the

interaction of the mutant protein with HCK unlike the WT protein.

HDX-MS reveals that the BTK T474I and L528W mutants show reduced binding to BTK

inhibitors.

Disruption of inhibitor binding due to mutations is a classic mechanism by which

resistance arises [47, 48]. While loss of inhibitor binding is likely irrelevant to the catalytically

dead BTK L528W mutant, we nevertheless evaluated it along with the BTK T474I mutant using

HDX-MS and NMR. HDX-MS data show that Tirabrutinib, Zanubrutinib and Pirtobrutinib

induce overall changes in deuterium incorporation for both WT BTK and the T474I mutant (Fig.

8a,b,c) These data suggest that the inhibitors are still capable of binding to the T474I mutant.

However, in the presence of Tirabrutinib, Zanubrutinib or Pirtobrutinib, the magnitude of the

protection in the kinase domain (N-lobe β2-β3 strands and activation loop) is consistently

reduced in drug bound T474I mutant as compared to the WT protein (Fig. 8a,b,c,e). The

differences in deuterium exchange for drug binding to WT and mutant BTK suggest that the

T474I mutation causes a reduction in inhibitor binding. Specifically, the peptides derived from

the kinase activation loop in particular do not show as much protection in the mutant relative to

the WT suggesting that the inhibitors do not stabilize the activation loop in the inactive

conformation upon binding to the T474I mutant. Interestingly, the changes observed in the BTK

regulatory domains (SH3 and SH2 domains) upon Zanubrutinib, Tirabrutinib or Pirtobrutinib

binding to WT BTK as well as increased deuterium incorporation in the G-helix of the kinase

domain upon Tirabrutinib binding to WT BTK are maintained or reduced in the drug bound BTK

T474I mutant. In stark contrast to the BTK T474I mutant, the BTK L528W mutant does not

show any change in deuterium incorporation in the presence of Zanubrutinib, Tirabrutinib or

Pirtobrutinib, providing strong evidence that the BTK L528W mutant does not bind the

inhibitors (Fig.8d). Taken together, both the BTK T474I and the BTK L528W mutation impact

inhibitor binding. Indeed, these results are consistent with previous studies that show that the

BTK T474I and the BTK L528W mutations disrupt binding to Zanubrutinib and Pirtobrutinib

[26]. We next tested the impact of the resistance mutations on inhibitor binding by NMR.

Probing the effects of BTK T474I and L528W mutations on inhibitor binding by NMR.

Comparison of the Zanubrutinib bound WT BTK LKD spectrum to that of BTK T474I

shows that the BTK T474I mutant undergoes minor changes in the presence of Zanubrutinib,

suggesting that the mutant is binding weakly to the inhibitor (Fig. 9a, left and middle panels).

The W395 resonance shows little to no chemical shift change upon addition of Zanubrutinib to

the BTK T474I mutant, suggesting that this drug does not stabilize the inactive conformation of

the mutant kinase (Fig. 9a, middle panel). These results are consistent with HDX-MS changes in

the T474I mutant upon Zanubrutinib binding (Fig. 8a).

Spectral comparison of Tirabrutinib bound to WT and BTK T474I (Fig. 9b, left and

middle panels) shows drug induced chemical shift changes for both complexes, supporting the

conclusion from the HDX-MS data (Fig. 8b) that the drug binds to both WT and mutant BTK.

However, the spectral changes observed in the spectrum of Tirabrutinib bound to BTK T474I are

different from that of the WT/Tirabrutinib bound spectrum (Fig. 9b). Although BTK W395 gives

rise to multiple peaks in the spectra of Tirabrutinib bound to both WT and T474I, the W395

resonances in the BTK T474I/Tirabrutinib complex are shifted upfield to a lesser extent

compared to the same resonance in the WT/Tirabrutinib spectrum. This suggests that the BTK

T474I mutation reduces binding to Tirabrutinib and the inactive kinase domain conformation

(αC out) is less populated in the drug bound mutant protein compared to WT BTK bound to

Tirabrutinib. These results are consistent with the HDX-MS results that show reduced protection

in peptides corresponding to the kinase activation loop in the mutant compared to WT due to the

lack of stabilization of the kinase inactive conformation upon Tirabrutinib binding.

Spectral overlay of the Pirtobrutinib bound BTK T474I with the apo T474I protein shows

pronounced chemical shift changes in the presence of the inhibitor, providing evidence

consistent with the HDX-MS data (Fig. 8c), that Pirtobrutinib binds the BTK T474I mutant (Fig.

9c, middle). However, unlike Pirtobrutinib bound to WT BTK, multiple W395 peaks are visible

in the spectrum of Pirtobrutinib bound to BTK T474I. The resonance frequencies of the

additional peaks are shifted toward that of the apo protein suggesting the possibility of fast

exchange between inhibitor bound and unbound states. These observations suggest that the

T474I mutation reduces affinity towards Pirtobrutinib and are consistent with previously

published SPR binding data [26].

BTK inhibitors that retain binding to the BTK T474I mutant could serve as potential

alternate treatment option for patients with this mutation. Additionally, the lack of or low

frequency of the BTK T474I mutation in patients treated with Ibrutinib and Acalabrutinib could

be due to the inability of this mutation to disrupt inhibitor binding. We therefore tested the ability

of the BTK T474I mutant to bind Ibrutinib, Acalabrutinib and Fenebrutinib (GDC-0853). The

NMR data suggest that the BTK T474I mutation does not impact Ibrutinib or Fenebrutinib

binding (Fig. 9d,f), but Acalabrutinib binding is reduced (Fig. 9e). Taken together, the T474I mutation significantly reduces binding to Zanubrutinib, Tirabrutinib and Pirtobrutinib, retains binding to Ibrutinib and Fenebrutinib, and may have an adverse effect on Acalabrutinib binding. Given that the BTK T474I mutant retains activity (albeit reduced activity), disruption of drug binding is a possible mechanism by which this mutation escapes inhibition.

Spectral overlays of the BTK L528W mutant with and without Zanubrutinib show no chemical shift changes (Fig. 9a, right panel) suggesting that the mutation completely disrupts inhibitor binding in complete agreement with the HDX-MS data (Fig. 8d). Tirabrutinib does show chemical shift changes, but the changes are markedly different from that observed in the WT protein (Fig. 9b, right). In the L528W spectrum in the presence of Tirabrutinib, W395 shifts in the opposite downfield direction compared to the WT spectrum suggesting that Tirabrutinib could be binding to the BTK L528W in a different orientation. Alternatively, the distinct chemical shift change could be due to the mutated L528W residue altering the local chemical environment within the kinase.

The Pirtobrutinib-bound BTK L528W spectrum (Fig. 9c, Supp. Fig. S1c) shows two resonance positions, one of which overlaps with the resonance in the apo protein and the other that corresponds to that of the mutant protein bound to Pirtobrutinib. This data suggests a mixture of inhibitor bound and unbound BTK kinase domain in solution, likely due to a reduction in Pirtobrutinib affinity caused by the L528W mutation. Although the L528W mutation reduces binding to both Tirabrutinib and Pirtobrutinib, the NMR data suggests that it retains partial binding unlike the HDX-MS data that suggests complete disruption of binding. The higher inhibitor concentrations used in the NMR experiments compared to the HDX-MS experiments likely explain this discrepancy. Interestingly, we note similarities between the BTK

L528W Pirtobrutinib bound spectrum and that of Tirabrutinib bound spectrum, suggesting that

the BTK kinase domain adopts similar conformations in solution when bound to these different

drugs. Additionally, the BTK L528W mutant retains binding to Fenebrutinib, however Ibrutinib

and Acalabrutinib binding are disrupted (Fig. 9d-f, right). Taken together, the BTK L528W

mutation significantly disrupts Zanubrutinib, Tirabrutinib and Pirtobrutinib binding. However,

drugs based on Fenebrutinib could be developed to treat patients carrying this mutation (see

discussion below).

DISCUSSION

471   Binding of BTK active site inhibitors can have long-range effects on the protein. Here,

we build on earlier work to show that three of the clinically approved BTKi (Acalabrutinib,

Zanubrutinib and Tirabrutinib) shift the conformational ensemble of full-length BTK,

destabilizing the autoinhibited conformation of the SH3 and SH2 domains to varying degrees

(Fig.4 and 5). In marked contrast, Pirtobrutinib led to the stabilization of the compact

autoinhibited conformation of full-length BTK. The exposure or stabilization of the regulatory

domains of BTK by active site BTK inhibitors can alter the interaction of BTK with ligands *in*

vivo. Indeed, BTK inhibitors have been shown to differ in their effectiveness at terminating

signals downstream of the Fc receptor *versus* the B cell receptor [49, 50]. As more BTKi become

available, a molecular-level understanding of the interaction of the inhibitor with full-length

BTK will aid the interpretation of the efficacy of different BTK inhibitors in treating disease

states driven by different signaling pathways.

483   The development of specific resistance mutations in patients treated with different BTKi

is intriguing. Covalent BTKi rely on BTK C481 within the kinase active site. The C481S

mutation prevents the covalent attachment of these inhibitors to BTK and converts the mode of

binding of these drugs to that of a reversible inhibitor. Importantly, we and others have shown

that the BTK C481S mutation does not prevent binding of Ibrutinib to the BTK kinase domain

[16, 26]. In fact, the BTK C481S mutant binds to Ibrutinib just as well as the WT protein *in vitro*

under equilibrium conditions where drug concentrations do not vary over time [16]. However, in

patients, drug concentrations change with time; inhibitor concentrations peak rapidly after intake

followed by a decrease as the drug is cleared (represented by the half-life of the drug). The

occupancy of BTK C481S protein by these covalent inhibitors in patients is therefore dictated by

the availability of these covalent inhibitors over time. Interestingly, Ibrutinib and Acalabrutinib

have short half-lives compared to Zanubrutinib and Tirabrutinib (Table 1). The short half-lives of

Ibrutinib and Acalabrutinib suggests that the BTK C481S mutant is likely to be unoccupied when

inhibitor concentrations fall, thereby allowing the BTK C481S mutant to escape from inhibition.

Conversely, the longer half-life of Zanubrutinib and Tirabrutinib in patients suggests that the

BTK C481S mutant is likely to remain occupied (inhibited) for a longer period of time. The

C481S mutation therefore does not offer a selective advantage as a resistance mutation in

Zanubrutinib and Tirabrutinib treated patients and likely explains the lower frequency of

occurrence of this mutation in these patients compared to Ibrutinib and Acalabrutinib treated

patients. Taken together, the BTK C481S mutant can escape covalent inhibition at low drug

concentrations *in vivo* and is therefore predicted to arise in patients treated with BTK covalent

inhibitors that have short half-lives.

The T474I mutation arises frequently in Tirabrutinib and Pirtobrutinib treated CLL

patients (Fig. 1d). Our HDX-MS and NMR binding studies show that the T474I mutation

disrupts binding to both inhibitors. Reduced binding of BTK T474I mutant to Tirabrutinib and

Pirtobrutinib, along with the partial catalytic activity retained by this mutant, would allow BTK

T474I mutant to escape inhibition and maintain BCR signaling. This gives a selective advantage

to the cells that carry this mutation. Additionally, our NMR binding studies show that the BTK

T474I mutation does not disrupt binding of Ibrutinib. This is consistent with the low occurrence

of this mutation in Ibrutinib-treated CLL patients. While the BTK T474I mutation does seem to

disrupt binding to Acalabrutinib, the lower activity of the T474I mutant (compared to the BTK

C481S mutant) may explain the higher prevalence of the BTK C481S mutation (compared to

T474I) in Acalabrutinib treated patients.

Analysis of the crystal structures of BTK bound to inhibitors provide clues as to why

some BTK inhibitors are more susceptible to loss of binding due to the BTK T474I and L528W

mutations. All BTK inhibitors evaluated in this study occupy the kinase active site, contact the

kinase hinge region (BTK residues E475-M477) and (with the exception of Fenebrutinib) extend

into the back pocket of the kinase (Fig. 10). BTK T474 is adjacent to the kinase hinge region and

BTK L528 occupies the 'base' of the kinase active site. Given the central location of both these

BTK residues, introduction of the bulky T474I and L528W mutations are likely to impact

multiple inhibitor interactions. Zanubrutinib and Pirtobrutinib in particular, make extensive

contacts with the kinase hinge region with three hydrogen bonds to BTK E475 and M477

backbone atoms. In contrast, Ibrutinib makes only a single hydrogen bond with BTK E475

backbone. This likely explains why the BTK T474I and L528W mutations disrupt Zanubrutinib

and Pirtobrutinib binding compared to Ibrutinib. The BTK inhibitor Fenebrutinib also seems to

be less impacted by the BTK T474I and L528W mutations. The BTK T474I and L528W

mutations are likely accommodated as Fenebrutinib does not occupy the 'back pocket' of the

kinase active site. While these BTK WT inhibitor bound structures provide clues as to why the

efficacy of certain BTK inhibitors are more impacted by the BTK T474I and L528W mutations,

some questions remain. Tirabrutinib for example makes a single hydrogen bond with the hinge

region, however, it is severely impacted by the BTK T474I and L528W mutations. These

mutations likely disrupt additional interactions that Tirabrutinib makes with the kinase such as

the glycine loop of the kinase. Crystal structures of inhibitors bound to the BTK mutants are

needed to conclusively evaluate their effects.

Ideas and Speculation

Successful treatment of CLL has relied on the inhibition of BTK catalytic activity to curb

BCR signaling. Inhibitor-bound, catalytically-inactive BTK is incapable of propagating BCR

signaling. Paradoxically, the catalytically inactive BTK L528W mutant is able to propagate BCR

signals [26, 43]. A similar catalytically inactive BTK mutant, the BTK C481Y/F resistance

mutation also arises in Ibrutinib and Acalabrutinib treated patients, albeit at a low frequency

[44]. Since these BTK mutants are catalytically dead, other kinases must compensate for the

absence of BTK activity to account for the intact BCR signaling in these patients. Exogenous

kinases do not appear to compensate for WT BTK that is inactivated by BTK inhibitors, and so

there must be additional differences between the catalytically dead (BTK L528W and BTK

C481Y/F) mutant kinases and inhibitor bound WT BTK that is rendered catalytically inactive by

virtue of drug binding. Our HDX-MS and NMR data show that BTK L528W mutation does not

drastically change the conformation of BTK. However, this mutation does increase the stability

of BTK compared to the WT protein. This increased stability could alter protein levels or change

the trafficking/localization of the mutant protein *in vivo* which in turn could alter protein-protein

interactions that drive activation of an alternate kinase such as HCK *in vivo*. Comparison of BTK

protein levels in patients pre- and post-development of resistance will be needed to test this

hypothesis. Alternatively, the BTK L528W mutation may require an additional change/s in the

cell that allow for this dead kinase to recruit other kinases such as HCK to propagate BCR

signaling. This requirement for additional changes may explain the low frequency of the L528W

mutation in Ibrutinib and Acalabrutinib treated patients; the single amino acid change, C481S,

combined with the rapid clearance of drug seems sufficient to drive resistance. Finally, the

absence of the BTK L528W resistance mutation in Ibrutinib treated CLL patients could be due to

the promiscuity of Ibrutinib; candidate compensatory kinases may be inactivated by the drug

[51]. The increased specificity of Zanubrutinib, Tirabrutinib and Pirtobrutinib on the other hand

could allow for the utilization of compensatory kinases such as HCK by BTK L528W in order to

propagate BCR signaling.

Understanding the mechanism/s by which resistance mutations evade inhibition allows

for counter strategies to be devised [52-54]. The reversible BTK inhibitor Pirtobrutinib is able to

inhibit the BTK C481S mutant due to its long serum half-life and has been suggested as a

treatment option for patients that develop this mutation [55]. Alternatively, drug occupancy in

the context of the BTK C481S resistance mutant could be increased by administering a twice

daily dose of the inhibitor as opposed to the single daily dose possibly circumventing their short

half-lives [56]. This increased exposure time to the covalent inhibitor could cause increased side

effects and so such risks should be considered before altering the dosage regimen. Alternatively,

Tirabrutinib, which exhibits the longest half-life among the available covalent BTK inhibitors,

could be used to slow progression of disease in patients that develop the C481S resistance

mutation.

The BTK T474I mutation disrupts binding to Zanubrutinib, Tirabrutinib, Pirtobrutinib

and Acalabrutinib (Fig.8, 9) but retains binding to Ibrutinib and Fenebrutinib creating possible

options for patients that develop the BTK T474I resistance mutation. The BTK L528W mutant is

catalytically inactive, hence treatment with any BTK active site inhibitor is futile. Previous

studies have shown that the similarly inactive BTK C481Y/F mutant can be suppressed by the

use of PROTACS to induce degradation of the full-length protein [44]. Importantly, our data

indicate that although the BTK L528W mutation disrupted binding to most BTK inhibitors it

retains binding to Fenebrutinib (Fig.9), suggesting that PROTACS based on Fenebrutinib or

similar backbones could be developed to treat this resistance mutation [57]. Alternate treatment

approaches such as targeting HCK or other BCR signaling proteins such as BCL-2 [58] could

also be used to counter this resistance mutation. As additional patient data becomes available, the

patterns of resistance mutations for different inhibitors will become clearer and strategies to

circumvent resistance using existing inhibitors should improve. As for many areas of medicine,

the treatment landscape should benefit from the ongoing era of personalized medicine.

MATERIALS AND METHODS

Constructs and reagents: The bacterial expression constructs for murine BTK linker kinase

domain (LKD) and full-length (FL) have been described previously [39]. All BTK constructs

carry the solubilizing Y617P mutation for bacterial expression [39]. All mutations were made

using the site directed mutagenesis kit (Agilent), and the sequences of all constructs were

confirmed by sequencing at the Iowa State University DNA facility. Acalabrutinib, Zanubrutinib,

Tirabrutinib and Pirtobrutinib were purchased from MedChem Express. Ibrutinib was purchased

from Selleckchem. The pCDF-1 Duet HCK SH3-SH2-Kinase domain/YopH construct with a

'YEEI' tail, a kind gift from Dr. Tom Smithgall, was mutated to convert the 'YEEI' tail residues

to that of the WT protein (YQQQ). The pGEX PLCγ1 cSH2-linker Y771F/Y775F protein

substrate construct has been described previously [59].

Protein expression and purification: Expression and purification of the PLCγ1 cSH2-linker

Y771F/Y775F protein substrate has been described previously [59]. HCK, Full-length BTK and

BTK linker kinase domain (WT and mutants) were produced by co-expressing with YopH in

BL21(DE3) (Millipore Sigma) or BL21-Gold(DE3) cells (Agilent Technologies) as described

previously [39]. Briefly, the culture was grown at 37 °C to an O.D. 600 nm of 0.6 to 0.8. The

temperature of the culture was lowered to 18°C and then induced with either 1.0 mM IPTG for

BTK LKD construct or 0.1 mM IPTG for BTK full-length and HCK. The culture was harvested

24 hours after induction and the pellets were resuspended in lysis buffer (50 mM KH_2PO_4, pH

8.0, 150 mM NaCl, 20 mM imidazole and 0.5 mg/ml lysozyme) and stored at -80°C. Cells were

lysed by thawing and the action of lysozyme, and 3000 U DNAse I (Sigma) and 1 mM PMSF

were added to the lysate, incubated at RT for 20 minutes and then spun at 16,000 rpm for one

hour at 4°C. Glycerol was added to the supernatant to a final concentration of 10 % and was then

incubated with Ni-NTA resin (QIAGEN) for two hours, washed with Tris pH 8.0, 75 mM NaCl,

40 mM imidazole, and eluted in 20 mM Tris pH 8.0, 150 mM NaCl, 250 mM Imidazole, and

10% glycerol. Eluted protein was flash frozen in liquid nitrogen and stored at -80°C. The

proteins were concentrated and further purified by size exclusion chromatography (Hiload

Superdex 26/60 200 pg or Hiload Superdex 26/60 75 pg, GE Healthcare). The fractions

containing pure protein were pooled, concentrated, snap frozen and stored at -80° C. The final

buffer consists of 20 mM Tris pH 8.0, 150 mM Sodium chloride, 0.02 % Sodium azide and 10%

glycerol. Initial phosphorylation levels of all purified BTK and HCK proteins used in this study

is below western immuno-detection.

NMR: Uniformly [15]N labeled BTK samples were produced as described earlier by growth in

modified M9 minimal media containing [15]N ammonium chloride (1g/L, Cambridge Isotope

Laboratories, Inc.) as the sole source of nitrogen [39]. The final NMR sample buffer consists of

20 mM Tris, 150 mM Sodium chloride, 10% glycerol, and 0.02 % Sodium azide at pH 8.0. All

NMR spectra were collected at 298 K on a Bruker AVIII HD 800 spectrometer equipped with a

5 mm HCN z-gradient cryoprobe operating at a ^{1}H frequency of 800.37 NMR samples with

inhibitors consisted of 150 µM ^{15}N labeled BTK, mixed with 200 µM inhibitor in 2% DMSO.

All data were analyzed using NMRViewJ [60].

HDX-MS: General procedures for HDX-MS of BTK have been described in detail previously

[39]. Details specific to experiments conducted here are provided in the Supplemental Datafile in

the format recommended [61] for HDX-MS experimental descriptions. All HDX-MS data have

been deposited to the ProteomeXchange Consortium via the PRIDE [62] partner repository with

the dataset identifier PXD047865. Briefly, prior to continuous labeling HDX experiments,

purified BTK full-length wild-type, T474I or L528W (20 µM) and inhibitor (40 µM), (20 mM

Tris pH 8.0, 150mM NaCl, 10% glycerol, 2% DMSO) were allowed to interact at 21 °C for 1

hour. After the binding reactions, both the free kinase and kinase bound to inhibitor were placed

on ice prior to deuterium labeling. Deuterium labeling proceeded for the times described using

labeling buffer, and labeling was stopped with an equal volume of quench buffer at 0°C (details

in Supplemental Datafile). Quenched samples were immediately analyzed using a Waters

nanoACQUITY with HDX technology using online pepsin digestion with a Waters Enzymate

immobilized pepsin column and UPLC separation of the resulting peptic peptides. Mass spectra

were acquired using a Waters Synapt HDMS^E mass spectrometer. Peptides generated from

online pepsin digestion were identified with Waters Protein Lynx Global Server 3.0 using

separate unlabeled protein that was prepared in the same manner as protein labeled with

deuterium. Deuterium incorporation was quantified using Waters DynamX 3.0. Deuterium levels

for each peptide were calculated by subtracting the average mass of the undeuterated control

sample from that of the deuterium labeled sample; the data were not corrected for back exchange

and are therefore reported as relative [63]. Vertical difference maps in Figures 4, 6 and 8 do not

represent a linear sequence of non-overlapping peptides. All coincident and overlapping peptides

for comparisons in each figure are provided in figure identified tabs of the Supplemental

Datafile.

Activity assays: *In vitro* kinase assays were performed as described previously [39]. Briefly, 1

μM BTK FL, BTK FL T474I or BTK FL L528W proteins were incubated in a kinase assay

buffer (50 mM Hepes pH 7.0, 10 mM $MgCl_2$, 1 mM DTT, 5 % glycerol, 1 mM Pefabloc, and

200 μM ATP) at room temperature for varying time. The reactions were stopped by the addition

of SDS-PAGE loading buffer and the samples were boiled, separated by SDS−PAGE, and

Western blotted with the anti-BTK pY551 antibody (BD Pharmingen[TM]) or anti-His antibody

(EMD Millipore) as described previously [39]. The bands were quantified using the

ChemiDoc[TM] (Biorad) gel imaging system. The phosphorylation levels (Anti-BTK pY551 blot)

were normalized to the total protein level (Anti-His blot). The BTK FL value at 40 minutes was

set to 1 and compared to BTK FL T474I or BTK FL L528W. The HCK activation experiments

were performed in the absence of Zanubrutinib or by preincubating 0.2 μM of BTK WT, L528W

or L528W/Y551F proteins with 0.5 μM Zanubrutinib for 15 min at RT. The reaction was

initiated with the addition of 0.2 μM HCK in a kinase assay buffer (50 mM Hepes pH 7.0, 10

mM $MgCl_2$, 1 mM DTT, 5 % glycerol, 1 mM Pefabloc, 30 μM PLCγ1 cSH2 linker

Y771F/Y775F substrate and 200 μM ATP). Time points were taken at 5, 10 and 15 minutes. The

reactions were stopped by the addition of SDS-PAGE loading buffer and the samples were

boiled, separated by SDS−PAGE, and Western blotted with the anti-PLCγ1-pY783 antibody

(Cell Signaling) or anti-pY antibody (4G10, Millipore Sigma) or anti-His antibody (EMD

Millipore) as described previously [39]. The bands were quantified using the ChemiDoc^{TM}

(Biorad) gel imaging system. The HCK alone value was set to 1 and compared to HCK + BTK

WT/L528W. Initial phosphorylation levels of BTK, prior to the start of the activity assay were

undetectable by western immuno-detection.

Thermal shift assays: BTK FL WT or mutant at 5 µM was incubated with 10X final

concentration of SYPRO Orange dye (Thermo Fisher Scientific) in a total volume of 20 µL Tris

buffer (20 mM Tris, 150 mM Sodium chloride, 10% glycerol, pH 8.0). Thermal shift assays were

performed on the StepOnePlus RT-PCR machine in the Iowa State University DNA Facility. The

temperature was raised from 25 °C to 95 °C and measurements recorded with every 0.5 °C

increment in temperature. The derivative plot of the data was analyzed for the melting

temperature (Tm). Assays were performed thrice and the average Tm was calculated.

ACKNOWLEDGEMENTS

This work is supported by a grant from the National Institutes of Health (National

Institute of Allergy and Infectious Diseases, AI43957) to A.H.A., J.R.E and T.E.W., and by

funds from the Scott-Waudby Trust, the Hope Against Cancer charity, Cancer Research UK in

conjunction with the UK Department of Health on an Experimental Cancer Medicine Centre

grant [C10604/A25151] to S.J, R.G.B and M.J.SD. Research at the University of Leicester was

carried out at the National Institute for Health and Care Research (NIHR) Leicester Biomedical

Research Centre (BRC). The authors also thank the Roy J. Carver Charitable Trust, Muscatine,

Iowa for ongoing research support.

REFERENCES

1.   Burger, J.A., et al., *Ibrutinib as Initial Therapy for Patients with Chronic Lymphocytic Leukemia.* N
Engl J Med, 2015. **373**(25): p. 2425-37.
2.   Davids, M.S. and J.R. Brown, *Ibrutinib: a first in class covalent inhibitor of Bruton's tyrosine
kinase.* Future Oncol, 2014. **10**(6): p. 957-67.
3.   Honigberg, L.A., et al., *The Bruton tyrosine kinase inhibitor PCI-32765 blocks B-cell activation and
is efficacious in models of autoimmune disease and B-cell malignancy.* Proc Natl Acad Sci U S A,
2010. **107**(29): p. 13075-80.
4.   Herman, S.E., et al., *Bruton tyrosine kinase represents a promising therapeutic target for
treatment of chronic lymphocytic leukemia and is effectively targeted by PCI-32765.* Blood, 2011.
117(23): p. 6287-96.
5.   Rozkiewicz, D., et al., *Bruton's Tyrosine Kinase Inhibitors (BTKIs): Review of Preclinical Studies
and Evaluation of Clinical Trials.* Molecules, 2023. **28**(5).
6.   Shirley, M., *Bruton Tyrosine Kinase Inhibitors in B-Cell Malignancies: Their Use and Differential
Features.* Target Oncol, 2022. **17**(1): p. 69-84.
7.   Montoya, S. and M.C. Thompson, *Non-Covalent Bruton's Tyrosine Kinase Inhibitors in the
Treatment of Chronic Lymphocytic Leukemia.* Cancers (Basel), 2023. **15**(14).
8.   Kueffer, L.E., R.E. Joseph, and A.H. Andreotti, *Reining in BTK: Interdomain Interactions and Their
Importance in the Regulatory Control of BTK.* Front Cell Dev Biol, 2021. **9**: p. 655489.
9.   Frustaci, A.M., et al., *Next Generation BTK Inhibitors in CLL: Evolving Challenges and New
Opportunities.* Cancers (Basel), 2023. **15**(5).
10.   Mendes-Bastos, P., et al., *Bruton's tyrosine kinase inhibition-An emerging therapeutic strategy in
immune-mediated dermatological conditions.* Allergy, 2022. **77**(8): p. 2355-2366.
11.   Robak, E. and T. Robak, *Bruton's Kinase Inhibitors for the Treatment of Immunological Diseases:
Current Status and Perspectives.* J Clin Med, 2022. **11**(10).
12.   Byrd, J.C., et al., *Acalabrutinib Versus Ibrutinib in Previously Treated Chronic Lymphocytic
Leukemia: Results of the First Randomized Phase III Trial.* J Clin Oncol, 2021. **39**(31): p. 3441-
3452.
13.   Brown, J.R., et al., *Zanubrutinib or Ibrutinib in Relapsed or Refractory Chronic Lymphocytic
Leukemia.* N Engl J Med, 2023. **388**(4): p. 319-332.
14.   Lovell, A.R., N. Jammal, and P. Bose, *Selecting the optimal BTK inhibitor therapy in CLL: rationale
and practical considerations.* Ther Adv Hematol, 2022. **13**: p. 20406207221116577.
15.   Buske, C., et al., *Managing Waldenstrom's macroglobulinemia with BTK inhibitors.* Leukemia,
2023. **37**(1): p. 35-46.
16.   Joseph, R.E., et al., *Differential impact of BTK active site inhibitors on the conformational state of
full-length BTK.* Elife, 2020. **9**.
17.   Sonti, R., et al., *ATP Site Ligands Determine the Assembly State of the Abelson Kinase Regulatory
Core via the Activation Loop Conformation.* J Am Chem Soc, 2018. **140**(5): p. 1863-1869.
18.   Tong, M., et al., *Survey of solution dynamics in Src kinase reveals allosteric cross talk between
the ligand binding and regulatory sites.* Nat Commun, 2017. **8**(1): p. 2160.
19.   Skora, L., et al., *NMR reveals the allosteric opening and closing of Abelson tyrosine kinase by
ATP-site and myristoyl pocket inhibitors.* Proc Natl Acad Sci U S A, 2013. **110**(47): p. E4437-45.
20.   Sedlarikova, L., et al., *Resistance-Associated Mutations in Chronic Lymphocytic Leukemia
Patients Treated With Novel Agents.* Front Oncol, 2020. **10**: p. 894.
21.   Maddocks, K.J., et al., *Etiology of Ibrutinib Therapy Discontinuation and Outcomes in Patients
With Chronic Lymphocytic Leukemia.* JAMA Oncol, 2015. **1**(1): p. 80-7.

22. Ahn, I.E., et al., *Clonal evolution leading to ibrutinib resistance in chronic lymphocytic leukemia.*
Blood, 2017. **129**(11): p. 1469-1479.

23. Woyach, J.A., et al., *Resistance mechanisms for the Bruton's tyrosine kinase inhibitor ibrutinib.* N
Engl J Med, 2014. **370**(24): p. 2286-94.

24. Blombery, P., et al., *Enrichment of BTK Leu528Trp mutations in patients with CLL on*
zanubrutinib: potential for pirtobrutinib cross-resistance. Blood Adv, 2022. **6**(20): p. 5589-5592.

25. Furman, R.R., et al., *Ibrutinib resistance in chronic lymphocytic leukemia.* N Engl J Med, 2014.
370(24): p. 2352-4.

26. Wang, E., et al., *Mechanisms of Resistance to Noncovalent Bruton's Tyrosine Kinase Inhibitors.* N
Engl J Med, 2022. **386**(8): p. 735-743.

27. Jackson, R.A., et al., *BTK mutations in patients with chronic lymphocytic leukemia receiving*
tirabrutinib. Blood Adv, 2023. **7**(14): p. 3378-3381.

28. Handunnetti SM, T.C., Nguyen T, et al., *BTK Leu528Trp - a Potential Secondary Resistance*
Mechanism Specific for Patients with Chronic Lymphocytic Leukemia Treated with the Next
Generation BTK Inhibitor Zanubrutinib. Blood., 2019. **134**: p. 170.

29. Woyach, J.A., Huang, Y., Rogers, K., Bhat, S.A., et al., *Resistance to Acalabrutinib in CLL Is*
Mediated Primarily By BTK Mutations. Blood, 2019. **134**: p. 504.

30. Woyach, J.A., et al., *BTK(C481S)-Mediated Resistance to Ibrutinib in Chronic Lymphocytic*
Leukemia. J Clin Oncol, 2017. **35**(13): p. 1437-1443.

31. Nakhoda, S., A. Vistarop, and Y.L. Wang, *Resistance to Bruton tyrosine kinase inhibition in*
chronic lymphocytic leukaemia and non-Hodgkin lymphoma. Br J Haematol, 2023. **200**(2): p. 137-
149.

32. Kanagal-Shamanna, R., et al., *Targeted multigene deep sequencing of Bruton tyrosine kinase*
inhibitor-resistant chronic lymphocytic leukemia with disease progression and Richter
transformation. Cancer, 2019. **125**(4): p. 559-574.

33. Gango, A., et al., *Dissection of subclonal evolution by temporal mutation profiling in chronic*
lymphocytic leukemia patients treated with ibrutinib. Int J Cancer, 2020. **146**(1): p. 85-93.

34. Sharma, S., et al., *Identification of a structurally novel BTK mutation that drives ibrutinib*
resistance in CLL. Oncotarget, 2016. **7**(42): p. 68833-68841.

35. Xie, Q., et al., *Substrate recognition of PLCgamma1 via a specific docking surface on Itk.* J Mol
Biol, 2013. **425**(4): p. 683-96.

36. Lin, D.Y. and A.H. Andreotti, *Structure of BTK kinase domain with the second-generation*
inhibitors acalabrutinib and tirabrutinib. PLoS One, 2023. **18**(8): p. e0290872.

37. Gomez, E.B., et al., *Preclinical characterization of pirtobrutinib, a highly selective, noncovalent*
(reversible) BTK inhibitor. Blood, 2023. **142**(1): p. 62-72.

38. Guo, Y., et al., *Discovery of Zanubrutinib (BGB-3111), a Novel, Potent, and Selective Covalent*
Inhibitor of Bruton's Tyrosine Kinase. J Med Chem, 2019. **62**(17): p. 7923-7940.

39. Joseph, R.E., et al., *Achieving a Graded Immune Response: BTK Adopts a Range of Active/Inactive*
Conformations Dictated by Multiple Interdomain Contacts. Structure, 2017. **25**(10): p. 1481-1494
e4.

40. Lin, D.Y., Kueffer, L.E., Juneja, P., Wales, T., et al., *Conformational heterogeneity of the BTK PHTH*
domain drives multiple regulatory states. eLife, 2023.

41. Wang, Q., et al., *Autoinhibition of Bruton's tyrosine kinase (Btk) and activation by soluble inositol*
hexakisphosphate. Elife, 2015. **4**.

42. Johnson, A.R., et al., *Battling Btk Mutants With Noncovalent Inhibitors That Overcome Cys481*
and Thr474 Mutations. ACS Chem Biol, 2016. **11**(10): p. 2897-2907.

43. Yuan, H., et al., *BTK kinase activity is dispensable for the survival of diffuse large B-cell*
lymphoma. J Biol Chem, 2022. **298**(11): p. 102555.

44. Dhami, K., et al., *Kinase-deficient BTK mutants confer ibrutinib resistance through activation of*
the kinase HCK. Sci Signal, 2022. **15**(736): p. eabg5216.
45. Moarefi, I., et al., *Activation of the Src-family tyrosine kinase Hck by SH3 domain displacement.*
Nature, 1997. **385**(6617): p. 650-3.
46. Pessoa, J., et al., *Editorial: Altered Expression of Proteins in Cancer: Function and Potential*
Therapeutic Targets. Front Oncol, 2022. **12**: p. 949139.
47. Bhullar, K.S., et al., *Kinase-targeted cancer therapies: progress, challenges and future directions.*
Mol Cancer, 2018. **17**(1): p. 48.
48. Cohen, P., D. Cross, and P.A. Janne, *Kinase drug discovery 20 years after imatinib: progress and*
future directions. Nat Rev Drug Discov, 2021. **20**(7): p. 551-569.
49. Bender, A.T., et al., *Ability of Bruton's Tyrosine Kinase Inhibitors to Sequester Y551 and Prevent*
Phosphorylation Determines Potency for Inhibition of Fc Receptor but not B-Cell Receptor
Signaling. Mol Pharmacol, 2017. **91**(3): p. 208-219.
50. Li, W., et al., *Bruton's Tyrosine Kinase Inhibitors With Distinct Binding Modes Reveal Differential*
Functional Impact on B-Cell Receptor Signaling. Mol Cancer Ther, 2023.
51. Yang, G., et al., *HCK is a survival determinant transactivated by mutated MYD88, and a direct*
target of ibrutinib. Blood, 2016. **127**(25): p. 3237-52.
52. Skanland, S.S. and A.R. Mato, *Overcoming resistance to targeted therapies in chronic*
lymphocytic leukemia. Blood Adv, 2021. **5**(1): p. 334-343.
53. Smith, C.I.E. and J.A. Burger, *Resistance Mutations to BTK Inhibitors Originate From the NF-*
kappaB but Not From the PI3K-RAS-MAPK Arm of the B Cell Receptor Signaling Pathway. Front
Immunol, 2021. **12**: p. 689472.
54. Alexandra Chirino, S.M., Anita Safronenka and Justin Taylor, *Resisting the Resistance: Navigating*
BTK Mutations in Chronic Lymphocytic Leukemia (CLL). Genes, 2023. **14**: p. 2182.
55. Aslan, B., et al., *Pirtobrutinib inhibits wild-type and mutant Bruton's tyrosine kinase-mediated*
signaling in chronic lymphocytic leukemia. Blood Cancer J, 2022. **12**(5): p. 80.
56. Skanland, S.S. and G.E. Tjonnfjord, *Determining drug dose in the era of targeted therapies:*
playing it (un)safe? Blood Cancer J, 2022. **12**(8): p. 123.
57. Zorba, A., et al., *Delineating the role of cooperativity in the design of potent PROTACs for BTK.*
Proc Natl Acad Sci U S A, 2018. **115**(31): p. E7285-E7292.
58. Molica, S., *Venetoclax: a real game changer in treatment of chronic lymphocytic leukemia.* Int J
Hematol Oncol, 2020. **9**(4): p. IJH31.
59. Joseph, R.E., et al., *A remote substrate docking mechanism for the tec family tyrosine kinases.*
Biochemistry, 2007. **46**(18): p. 5595-603.
60. Johnson, B.A. and R.A. Blevins, *NMR View: A computer program for the visualization and*
analysis of NMR data. J Biomol NMR, 1994. **4**(5): p. 603-14.
61. Masson, G.R., et al., *Recommendations for performing, interpreting and reporting hydrogen*
deuterium exchange mass spectrometry (HDX-MS) experiments. Nat Methods, 2019. **16**(7): p.
595-602.
62. Vizcaino, J.A., et al., *2016 update of the PRIDE database and its related tools.* Nucleic Acids Res,
2016. **44**(22): p. 11033.
63. Wales, T.E. and J.R. Engen, *Hydrogen exchange mass spectrometry for the analysis of protein*
dynamics. Mass Spectrom Rev, 2006. **25**(1): p. 158-70.
64. Advani, R.H., et al., *Bruton tyrosine kinase inhibitor ibrutinib (PCI-32765) has significant activity*
in patients with relapsed/refractory B-cell malignancies. J Clin Oncol, 2013. **31**(1): p. 88-94.
65. Podoll, T., et al., *Bioavailability, Biotransformation, and Excretion of the Covalent Bruton*
Tyrosine Kinase Inhibitor Acalabrutinib in Rats, Dogs, and Humans. Drug Metab Dispos, 2019.
47(2): p. 145-154.

66.   Tam, C.S., et al., *Phase 1 study of the selective BTK inhibitor zanubrutinib in B-cell malignancies*
and safety and efficacy evaluation in CLL. Blood, 2019. **134**(11): p. 851-859.
67.   Walter, H.S., et al., *A phase 1 clinical trial of the selective BTK inhibitor ONO/GS-4059 in relapsed*
and refractory mature B-cell malignancies. Blood, 2016. **127**(4): p. 411-9.
68.   Mato, A.R., et al., *Pirtobrutinib in relapsed or refractory B-cell malignancies (BRUIN): a phase 1/2*
study. Lancet, 2021. **397**(10277): p. 892-901.

FIGURES

Figure 1:

Figure 1: (a) Domain organization of full-length (FL) BTK and the BTK linker-kinase domain

(LKD) fragment used in this study: PHTH, Pleckstrin homology-Tec homology domain; PRR,

proline-rich region; SH3, Src homology 3 domain; SH2, Src homology 2 domain, SH2-kinase

linker (L) and the catalytic kinase domain. Key residues are indicated above each domain. (b)

Autoinhibited conformation of FL BTK based on the crystal structure of FL BTK [40]. The

PHTH domain (purple) is dynamic, in transient contact with several regions on the core SH3-

SH2-kinase domain and is not visible in the crystal structure of full-length BTK [40]. Dynamics

of the PHTH domain is represented by the multiple poses of the PHTH domain and the double

headed arrow. (c) Co-crystal structure of BTK LKD (light cyan cartoon) bound to Ibrutinib

(PDB ID: 5P9J) showing the location of C481 (yellow spheres), T474 and L528 (red spheres)

within the kinase active site (broken oval). (c) Pie charts showing the prevalence of the BTK

resistance mutations in CLL patients treated with various BTK inhibitors. The total number of

patients with mutations in BTK are indicated below each chart. See Supp. Table 1 for additional

details.

Figure 2:

Figure 2: The BTK kinase domain can interconvert between active and inactive conformations.

(a) Superposition of the structure of BTK linker-kinase domain (LKD) bound to Dasatinib (PDB

ID: 3K54) in the active kinase conformation (grey cartoon) with the Ibrutinib bound structure

(PDB ID: 5P9J) in the inactive conformation (red cartoon). The expanded inset shows the inward

movement of the αC-helix, the change in W395 rotamer conformation and the K430/E445 salt

bridge formation that accompanies kinase activation. (b-f) Co-crystal structures of BTK LKD

(light cyan cartoon) bound to Ibrutinib (PDB ID: 5P9J), Acalabrutinib (PDB ID: 8FD9),

Zanubrutinib (PDB ID: 6J6M), Tirabrutinib (PDB ID: 5P9M) and Pirtobrutinib (PDB ID: 8FLL)

in the inactive kinase conformation. The inhibitors are shown as dark blue sticks, the kinase

activation loop is purple and C481, Y551 and W395 residues are shown as sticks with

transparent spheres. Electron density for part of the activation loop is missing in the Ibrutinib co-

crystal structure and is indicated as dotted lines (b). In the Acalabrutinib structure, the activation

loop has several mutations [36] and the SH2-kinase linker (including W395) is absent (c).

Electron density for W395 sidechain is missing in the BTK:Pirtobrutinib co-crystal structure (f).

(g) Overlay of the BTK:Ibrutinib, Acalabrutinib, Zanubrutinib, Tirabrutinib and Pirtobrutinib co-

crystal structures. With the exception of the Tirabrutinib co-crystal structure (e), no major

structural variation is observed in the kinase domains. The activation loop in the Tirabrutinib

bound structure adopts a different conformation compared to the other co-crystal structures.

Figure 3:

Figure 3: BTK inhibitors stabilize the inactive kinase conformation in solution. The tryptophan

side chain region of the ^{1}H-^{15}N TROSY HSQC spectra of ^{15}N-labelled apo BTK linker-kinase

domain (black spectrum) overlaid with that of the inhibitor bound spectrum (cyan spectrum).

Here and in subsequent figures, the broken black and grey lines indicate the position of the BTK

W395 resonance in the active (αC-in) and inactive (αC-out) states respectively as shown in

Figure 2a. The shift in the BTK W395 resonance upon inhibitor binding is indicated by an arrow

in each spectrum. The structures of each inhibitor are shown on the right. The BTK W395 indole

NH resonance is in the inactive (αC-out) position in the Ibrutinib (published earlier [16]),

Acalabrutinib, Zanubrutinib and Tirabrutinib bound BTK LKD samples. Multiple peaks

corresponding to W395 are seen in the Tirabrutinib bound spectrum suggesting that the kinase

adopts multiple conformations in solution. The downfield shift observed in W395 in the

Pirtobrutinib bound structure is likely due to local changes in the chemical environment due to

the distinct chemical structure of Pirtobrutinib. W395 assignments in the inhibitor bound spectra

were confirmed by acquiring inhibitor bound spectra with the BTK LKD W395A mutant (see

Supp. Fig. S1).

937 Figure 4:

940 Figure 4: Assessing the impact of BTK inhibitors on full-length BTK by HDX-MS. (a) Intact

941 mass analysis of wild-type FL BTK before (bottom spectrum, black) and after one hour

942 incubation with a 2-fold molar excess of covalent BTK inhibitors: Ibrutinib (red), Acalabrutinib

943 (purple), Zanubrutinib (blue) and Tirabrutinib (orange) show a mass increase of one inhibitor

molecule. (b) Clinically approved BTK inhibitors induce allosteric changes in full-length BTK. Relative deuterium level of peptides in apo BTK was subtracted from the deuterium level of the corresponding peptide from each drug-bound form of BTK ($D_{WT\ drug\text{-}bound}$-$D_{WT\ apo}$) and the differences colored according to the scale shown. In this and subsequent figures, peptic peptides are shown from N- to C-terminus, top to bottom, and the amount of time in deuterium is shown left to right. The relative difference data shown here represents a curated set of peptides that are coincident across all 6 states (apo and five drug-bound BTK forms). The identification of these chosen peptides, the relative difference values, and the complete data set for each state can be found in the Supplemental Datafile. The approximate position of the domains of BTK, as described in Figure 1a, is shown at the left. Deuterium incorporation curves of selected peptides (indicated with a gray box in panel b and labelled i-vi) from various regions of the protein are shown below. Data for Ibrutinib has been previously published [16].

Figure 5:

Figure 5: (a-e) Mapping the HDX-MS changes induced by each BTK inhibitor on the structure of the BTK SH3-SH2-kinase fragment (PDB ID: 4XI2). Changes mapped onto the structure represent the maximal HDX-MS change that occurred at any time point. Major differences greater than 1.0 Da are shown as dark blue (decrease) or dark green (increase); modest differences between 0.5 Da and 1.0 Da are shown as light blue (decrease) and light green (increase). Localization of the changes in deuterium incorporation was accomplished using overlapping peptides included in the complete peptide data set provided in the Supplemental

Datafile. The location of peptides i – vi from Figure 4 are indicated in panel a. Data

corresponding to Ibrutinib has been previously published [16].

Figure 6:

Figure 6: Probing the impact of the BTK resistance mutations T474I and L528W on BTK. (a-d)

Western blot comparing the kinase activity of full-length (FL) BTK wild-type (WT), T474I and

L528W mutants. BTK autophosphorylation was monitored using the BTK pY551 antibody and

the total protein levels monitored using the Anti-His antibody. (b, d) Histogram quantifying the

western blots shown in (a and c). The blots were quantified and normalized as described in the

Materials and Methods. Data shown are the average of three independent experiments. (e) HDX

difference data for the BTK T474I and L528W mutants ($D_{\text{Mutant apo}}$ -$D_{\text{WT apo}}$). Color scale and

peptide/time course arrangement are the same as in Figure 4. See the Supplemental Datafile for

additional information, including all peptide identifications and deuterium values. (f,g) Mapping

the mutational induced HDX-MS changes on the structure of the BTK SH3-SH2-kinase

fragment. Changes mapped onto the structure represent the maximal HDX-MS change that

occurred at any time point. (h) Deuterium incorporation curves of selected peptides (indicated

with a gray box in panel e and labelled i-iv) from various regions of the protein are shown. (i)

The tryptophan side chain region of the ^{1}H-^{15}N TROSY HSQC spectra of ^{15}N-labelled apo WT

BTK linker-kinase domain (black spectrum) overlaid with that of the apo mutant kinase

spectrum (cyan spectrum). The boxed region in the T474I spectral overlay is expanded above.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011   Figure 7:

1012

Figure 7: The BTK L528W mutant can activate HCK. (a-f) Kinase activity of HCK in the presence or absence of full-length BTK L528W mutant was compared in a western blot assay by monitoring PLCγ1 phosphorylation (pY783 antibody) and HCK autophosphorylation (pY antibody). BTK phosphorylation on Y551 was monitored using the Anti-pY551 antibody. Total

protein levels monitored using the Anti-His antibody. Full-length WT BTK preincubated with Zanubrutinib was used as a control. (b, d and f) Histogram quantifying the western blots shown in (a, c and e). The blots were quantified and normalized as described in the Materials and Methods. Data shown are the average of three independent experiments. (b and c) Kinase activity of HCK in the presence or absence of the isolated linker kinase domain (LKD) fragment of the BTK L528W mutant (b) or the full-length proline mutant of BTK L528W (BTK FL L528W/Pro: BTK L528W/ P189A/P192A/P203A/P206A, (c)) was compared as in (a). (g) Thermal stability analysis of BTK FL WT and BTK FL L528W. Data shown are the average of three independent experiments.

Figure 8:

Figure 8: HDX-MS analysis of BTK inhibitor binding to BTK T474I and L528W mutants. (a-d) Relative deuterium level of peptides in apo mutant BTK was subtracted from the deuterium level of the corresponding peptide from each inhibitor-bound form of BTK ($D_{drug-bound}$-D_{apo}) and compared to the changes in the WT protein. The differences are colored according to the scale shown. (e) Deuterium incorporation curves of selected peptides (indicated with a gray box in panels a-c, and labelled i-iv) are shown.

1046   Figure 9:

1047

1048

Figure 9: NMR analysis of BTK inhibitor binding to the BTK T474I and L528W mutants. The

tryptophan side chain region of the ^{1}H-^{15}N TROSY HSQC spectra of ^{15}N-labelled apo BTK

linker-kinase domain (black spectrum) overlaid with that of the inhibitor bound spectrum (cyan

spectrum). The broken black and grey lines indicate the position of the BTK W395 resonance

and have been described earlier in Figure 3. The shift in the BTK W395 indole NH resonance

upon inhibitor binding is indicated by an arrow in each spectrum. The red asterisks indicate the

presence of unbound kinase domain in the inhibitor bound NMR sample.

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073   Figure 10:

1074

1075

Figure 10: Close up of BTK inhibitors bound to the BTK kinase active site. The covalent BTK

inhibitors (Zanubrutinib, Tirabrutinib, Acalabrutinib and Ibrutinib, represented as sticks) are

bound to BTK C481. All BTK inhibitors, covalent and non-covalent (Pirtobrutinib and

Fenebrutinib) interact with the hinge region of the kinase and with the exception of Fenebrutinib,

extend into the back pocket of the kinase. Hydrogen bonds between the inhibitor and hinge

region are shown as dotted lines. The binding of these BTK inhibitors are incompatible with

nucleotide binding. The location of BTK T474 and L528, residues that are frequently mutated in

CLL patients are shown as sticks and spheres. BTK T474 is adjacent to the hinge region and

BTK L528 is at the 'base' of the kinase active site.

1085

1086   TABLES

1087   **Table 1**: Half-life of clinically approved BTK inhibitors: Ibrutinib[64], Acalabrutinib [65],

1088   Zanubrutinib [66], Tirabrutinib [67] and Pirtobrutinib [68].

Inhibitor	Half life (h)
Ibrutinib	2 - 3
Acalabrutinib	0.6 - 2.8
Zanubrutinib	4
Tirabrutinib	6.5 - 8
Pirtobrutinib	20

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101   Supp. Fig. S1

1102

1103

Supp Fig. S1: NMR analysis of inhibitor binding. (a) Cartoon of the BTK linker kinase domain

(LKD) fragment used for NMR analysis. (b) Assignment of W395 in inhibitor bound spectra of

BTK LKD. The tryptophan side chain region of the ^{1}H-^{15}N TROSY HSQC spectra of ^{15}N-

labelled inhibitor bound BTK LKD WT (black spectrum) overlaid with that of the inhibitor

bound BTK LKD W395A spectrum (cyan spectrum). The boxed peak indicates the W395

resonance in each of the WT inhibitor bound spectrum. (c) Overlay of the ^{1}H-^{15}N TROSY HSQC

spectra of ^{15}N-labelled apo BTK LKD L528W (black spectrum) with that of the Pirtobrutinib

bound BTK LKD L528W spectrum (cyan spectrum). The boxed resonances show resonance

doubling in the Pirtobrutinib bound spectrum, one of which overlaps with the apo BTK LKD

L528W spectrum, indicating a mixture of bound and unbound protein.

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127   Supp. Fig. S2:

1128

1129

Supp Fig. S2: BTK FL L528W is able to activate HCK. (a) Phosphorylation on BTK Y551 is not

required for HCK activation by BTK FL L528W. Kinase activity of HCK in the presence or

absence of full-length BTK L528W single or L528W/Y551F double mutant was compared in a

western blot assay by monitoring PLCγ1 phosphorylation (pY783 antibody) and HCK

autophosphorylation (pY antibody). Total protein levels monitored using the Anti-His antibody.

(b) Zanubrutinib is not required activation of HCK by BTK FL L528W. Kinase activity of HCK

in the presence or absence of full-length BTK L528W mutant, without Zanubrutinib was

compared in a western blot assay by monitoring PLCγ1 phosphorylation and HCK

autophosphorylation (pY antibody). Total protein levels monitored using the Anti-His antibody.

1139

1140

1141

1142

1143   Supplemental Datafile:

1144   Microsoft Excel file providing enhanced experimental details for HDX-MS including minimum

1145   criteria specified by [61], lists of all peptides by residue number, sequence, as well as deuterium

1146   levels measured for each Figure. The value of each deuterium difference for every colored box in

1147   each Figure as well as the complete dataset for each state are also found in this file.

1148

1149   **Supp. Table 1**: BTK mutations detected in CLL patients treated with BTK inhibitors. In addition

1150   to mutations in BTK C481, T474 and L528, other BTK mutations that have been detected

1151   include: R28S, E108K, G164D, V416L, A428D, M437R, R490H, Q516K, V537I and T316A

1152   [21-24, 26-30, 32-34].

BTK inhibitor	Total number of patients with BTK mutations.	Patients with BTK C481 mutated.	Patients with BTK T474 mutated.	Patients with BTK L528 mutated.	Patients with other BTK mutations.
Ibrutinib	106	95 (90%)	1 (1%)	1 (1%)	5 (5%)
Acalabrutinib	11	11 (100%)	1 (9%)	0	0
Zanubrutinib	17	15 (88%)	0	11 (65%)	0
Tirabrutinib	6	4 (67%)	3 (50%)	4 (67%)	1 (17%)
Pirtobrutinib	9	0	2 (22%)	4 (44%)	3 (33%)

1153