New Results

Detailed microbiome analysis of sticker-stripped surface materials of acne lesions revealed acne-related *Cutibacterium acnes* subtypes: a pilot study

Yutaka Shimokawa[1], Osamu Funatsu[1], Ohata Kazuma[1], Fukashi Inoue[2], Kota Tachibana[2], Itaru Dekio[3]*

1). KINS LAB, Tokyo, Japan

2). MySkin Corporation, Tokyo, Japan

3). Department of Dermatology, The Jikei University School of Medicine, Tokyo, Japan

*Correspondence: Itaru Dekio, M.D., Ph.D.

Department of Dermatology, School of Medicine, The Jikei University School of Medicine

3-25-8 Nishi-shinbashi, Minato-ku, Tokyo 105-8461, Japan

Tel: +81-3-3433-1111 ext 3341, Fax: +81-3-5401-0125.

E-mail: d-12@umin.org

Word count: 3614, References: 42, Figures: 7, Tables: 2, Supplementary figures: 2.

Abstract

Cutibacterium acnes (C. acnes) is known to play a central role in pathogenesis of acne vulgaris. It has been understood that multiple phylotypes of *C. acnes* exist, with certain types being more prevalent in patient with acne vulgaris and others more common in healthy individuals. In this context, we conducted a preliminary study using self-collected samples via an adhesive sticker (MySkin®) to analyze the skin microbiome of Japanese women. The study aimed to determine the role of *C. acnes* and its specific phylotypes in the development of acne vulgaris.

Participants in this study were Japanese females aged between their 20s and 40s. Dermatologists evaluate the data from web-based questionnaires and smartphone image submissions to classify subjects into either the Acne or Non-acne group. Quality assessment of DNA extracted from the sticker was conducted, followed by amplification of the *16S rRNA* region using PCR. Subsequent microbial community analysis was performed using next-generation sequencing techniques. Genetic classification of *C. acnes* was accomplished through single locus sequence typing.

Results indicated a bacterial community composition on the facial skin surface predominantly consisting of *C. acnes* clusters, with over half of these clusters constituted by *C. acnes*. Notably, the Acne group exhibited a significantly higher proportion of *C. acnes* relative to total bacterial presence compared to the Non-acne group. Analysis of *C. acnes*

phylotypes revealed a markedly lower presence of type III in the Acne group (vs. Non-acne

group, p<0.05). No significant differences were observed in the prevalence of Types IA_1, IA_2,

II, and IB between the two groups. The predominant phylotypes of *C. acnes* identified were

IA2_2_F0 (23.9%), IA1_4_A0 (20.6%), and II_2_K0 (18.6%). Within the Acne group, an

increase in IA2_1_F1 and a decrease in III_1_L0 were observed (vs. Non-acne group,

p<0.05).

This study underscores the feasibility of using self-collected and mailed-in samples

for qPCR and microbiome analysis, maintaining diagnostic quality comparable to in-person

assessments. Furthermore, the variation in the expression of *C. acnes* phylotypes across

skin surfaces between acne-afflicted and healthy individuals could suggest that shifts in

phylotype expression patterns may be indicative of skin susceptibilities to acne

development.

Introduction

Human skin serves as a barrier that separates the external and internal environments. It prevents the invasion of harmful substances into the body and blocks the loss of moisture and beneficial substances from the body. Its surface is home to many microorganisms, including bacteria, fungi, and viruses, which form colonies. These microorganisms interact with each other and with their host, creating a stable and complex ecosystem through cross-talk. This skin microbiome changes with factors such as age, gender, temperature, pH, ultraviolet exposure, moisture and humidity, and sebum content[1,2]. In recent years, comprehensive analyses of the skin microbiome have been conducted using next-generation sequencing techniques, specifically the analysis of bacterial *16S rRNA* gene. Reports have been emerging about its involvement in inflammatory skin diseases such as acne vulgaris, atopic dermatitis, and psoriasis[3–6].

Among the skin microbiome, the most common commensal bacterial species is *Cutibacterium acnes*[7,8]. *C. acnes* is a ubiquitous Gram-positive anaerobic bacterium belonging to the phylum *Actinobacteria* and is primarily found deep within the sebaceous hair follicles where it comes into contact with keratinocytes[9]. Furthermore, *C. acnes* is known to metabolize free fatty acids from sebum, which reduces the surface pH. This acidic environment potentially inhibits pathogenic bacteria such as *Staphylococcus aureus* and promotes the growth of other healthy commensal bacteria, suggesting its contribution to

barrier homeostasis[10–12].

Based on genotyping, it has become clear that *C. acnes* has multiple subtypes,

ranging from those involved in the onset of acne vulgaris to those existing on healthy

skin[13–15]. Moreover, the subspecies of *C. acnes* are classified as *C. acnes* subspecies

acnes (*C. acnes* type I), subspecies *defendens* (*C. acnes* type II), and subspecies

elongatum (*C. acnes* type III)[16,17]. Further, based on single locus sequence typing (SLST)

and whole genome sequencing (WGS) methods, the genetic sequence analysis of this

species has been phylogenetically refined and categorized into subtypes IA_1, IA_2, IB, IC, II,

and III[17]. Among these, subtype IA_1 of *C. acnes* has been isolated from acne vulgaris and is

believed to have strong pathogenicity[13,18,19]. On the other hand, subtypes IB[15] and II[14] are

expressed in healthy skin and are not believed to be associated with the onset of acne

vulgaris[19]. Thus, while there are multiple subtypes of *C. acnes* that seem to have different

roles, many aspects of their detailed ecology remain unknown.

At present, various methods such as cotton swabs, scraping, cyanoacrylate gel

biopsy, and needle biopsy are used to collect skin bacteria for analysis of the skin

microbiome. However, reports suggest that the detection rate of *C. acnes* varies depending

on the sampling method[20–22]. The most common sampling method is the swabbing method

using a cotton swab[23,24], but it is difficult to control the sampling efficiency depending on the

collection site and technique, and it has the disadvantage of poor preservation[20]. In any

case, it has not been envisaged that subjects would collect their own samples, and until

now, it has been difficult to collect samples by mail and analyze the skin microbiome.

However, with recent technological advancements, a sticker stripping kit for self-sampling

called MySkin®, which possesses features like preservation and quantification and can

withstand outdoor temperatures for postal services, has been developed. Therefore, this

study conducted a preliminary investigation with the aim to elucidate the relationship

between the skin resident bacterium *C. acnes* and acne vulgaris using samples collected by

self-sampling with the sticker stripping kit MySkin® among Japanese women.

Materials and Methods

Samples

From the female Japanese customers in their 20s to 40s who had purchased

products from KINS Co., Ltd., 296 individuals who agreed to the content of this study were

selected as the subjects. This research was conducted based on the Declaration of Helsinki

and the ethical guidelines for medical research, with the approval by the SOUKEN Ethics

Committee in April 2021 (Approval number: 146853_rn-30080). Individuals who had used

antibiotics within a month prior to sample collection were excluded from the study. Subjects

were not given any lifestyle restrictions such as the use of skincare products, except just

before sampling. The condition of the facial skin was determined from a web questionnaire

and a smartphone photo image of the face. The web questionnaire acquired responses to

the question, "Do you currently have visible pimples or acne on your face? Yes/No," and

concurrently, subjects sent photos taken with their smartphones. Dermatologists evaluated

these photos and categorized the patients into Acne group/Non-acne group. Subjects, after

waking up and before washing their faces, applied the dedicated adhesive sticker MySkin®

(an oval-shaped sticker with the area of 6.4 mm^2) to the specified locations on their facial

skin for 1 min to collect samples. After collection, the samples were promptly mailed to the

analytical institution. The flow from sample collection to data analysis is shown in Fig.1. The

dedicated skin sticker used in this study, designed for collecting skin microbes, has superior

microbial preservation and has been demonstrated to monitor relative bacterial count

fluctuations[25].

Skin microbiome analysis

Bacterial community analysis was performed by MySkin Co., Ltd. The DNA was

extracted from the skin sticker fragments and after performing a Quality Check (QC), the

obtained DNA was used as a template to amplify the bacterial *16S rRNA* by PCR.

Subsequently, indexes for next-generation sequencer migration were added to the PCR.

The DNA concentration of each sample was measured, and all samples were adjusted to

be of the same concentration. The adjusted DNA was purified and then sequenced using a

next-generation sequencer (MiSeq, Illumina Inc.) to produce the library. The library was

analyzed using the next-generation sequencer, targeting the V1-V2 regions of the *16S*

rRNA gene for bacterial community analysis. The microbiome analysis tool QIIME2[52] was

used, and for bacterial species prediction, the *16S rRNA* gene database Greengenes[26] was

employed. For the single gene-based subtype structure analysis of *C. acnes*, a *C.*

acnes-specific genomic region that can classify into six subtypes (IA$_1$, IA$_2$, IB, IC, II, III) was

utilized. A unique database regarding *C. acnes* was used to determine the subtypes.

Bacterial species in each sample were summarized by taxonomic rank, and from the

number of 16S rRNA reads, the relative abundance of each bacterium was calculated, and

the bacterial composition and percentages were determined. Based on these data,

hierarchical clustering using the Ward's method was performed to group the microbial

communities. Alpha diversity was evaluated using the Shannon index. Moreover, for the *C.*

acnes subtype analysis, in conjunction with the *16S rRNA* gene-based microbiome analysis,

a single locus sequence typing (SLST) analysis of *C. acnes* was conducted.

Polymerase chain reaction (PCR) was conducted using ExTaq (TaKaRa) to amplify

the SLST locus. Following amplification, the PCR products were purified using magnetic

beads (Sera-Mag™ Select, Cytiva). These purified samples were then analyzed using the

MiSeq sequencing platform. The resulting sequences were trimmed and classified into

SLST classes using the self-built SLST database .

 The primer sequences used for the SLST analysis are shown below.

   Fw-R3B:TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGacaccaggggggtcaacttgg

Rv-R3AB:GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGatgttacccatgtaatgggcagg

The uppercase represents the adapter sequences for MiSeq sequencing, while the

lowercase indicates the sequences used for acne bacteria typing.

Copy number analysis by qPCR

      For *C. acnes*, the copy numbers in the materials were determined by calculating

bacterial cell numbers per fixed area of the sticker from qPCR quantification. First, the skin

adhesive sticker fragments were placed in a tube and immersed in 1 ml of buffer, followed

by a spin-down and then left at 100℃ for 15 minutes. For the residual liquid after removing

the sticker, nucleic acid extraction was performed using the phenol/chloroform method. A

standard solution was adjusted and used by stepwise dilution of the standard DNA with a

known number of copies. The sequence of the PCR primers is confidential, but it is possible

to provide information upon request. PowerUp™ SYBR™ was used for detection, and

nucleic acid amplification was performed using the qPCR device (QuantStudio, Applied

Biosystems).

Statistical Analysis

The measurement results are shown as mean ± standard error, and if the *p*-value is less than 0.05, it was judged as statistically significant. For comparisons between two groups, Welch's *t*-test was used, and for comparisons among three or more groups, one-way analysis of variance (ANOVA) and *post-hoc* testing using the Bonferroni method were appropriately applied. Statistical processing was performed using HAD[27].

Results

Sample Quality

Quality control of the DNA extracted from the mailed samples was conducted. After PCR (both the first and second times), agarose gel electrophoresis was performed to confirm that the PCR product was of the expected size (Data not shown). In addition, quantification was carried out using qPCR, utilizing PCR primers corresponding to a portion of the P5 and P7 sequences to verify that adapter sequences had been added to both ends of the library. Based on these verifications, it was determined that the extracted DNA was suitable for use in the *16S rRNA* gene-based microbiome analysis.

Skin microbiome analysis

Out of 296 participants aged between 20 to 40, a comprehensive analysis of the resident bacteria on the facial skin was initially conducted on 76 randomly selected Japanese women (56 in the Acne group and 20 in the Non-acne group) (Table 1). As a result of hierarchical clustering analysis, the bacterial flora was classified into three groups: *C. acnes*-dominant, *Neisseriaceae*-dominant, and others (Fig2-A). The most prevalent was the cluster where *C. acnes* was dominant, in which 50-90% or more was composed of *C. acnes*. Subsequently, there were many clusters dominant with *Neisseriaceae*, which are resident bacteria of the oral cavity. On the other hand, regarding the fungal flora, it was divided into five major groups, consisting of clusters dominated by *Malassezia restricta* and *Malassezia globosa*, which are also suggested to be involved in acne vulgaris, and other clusters (Fig2-B). Interestingly, in the other clusters, there were participants who were dominated by *Rothia mucilaginosa*[28], which is a resident bacterium of the oral cavity, and *Wickerhamomyces anomalus*[29], known as a pathogenic yeast.

Alpha diversity

The *α*-diversity (species richness) on the facial skin was compared between the bacterial and fungal communities. The Shannon index for bacterial communities showed a median value of 0.382 in the Acne group and 0.702 in the Non-Acne group, with the Acne

group being significantly lower ($p<0.05$, Fig. 3-A). On the other hand, there was no

noticeable difference in the α-diversity of the fungal communities between the two groups

(Fig. 3-B). Therefore, subsequent analyses focused solely on bacteria.

Relationship between Acne vulgaris and C. acnes

Among the 296 participants, 219 were categorized into the Acne group, identified as

having acne vulgaris, while 76 were categorized into the Non-acne group, determined as

not having acne. Using the DNA samples extracted from facial skin, the proportion of *C.*

acnes within the entire bacterial community was investigated. As a result, the proportion of

C. acnes was significantly higher in the Acne group (51.7%) compared to the non-acne

participants (44.3%) ($p<0.001$, Fig.4-A). The logarithmic count of C. acnes per single sticker

strip was significantly reduced in the Acne group (4.12) compared to the Non-acne group

(3.72) ($p<0.001$, Fig.4-B). Furthermore, the proportion of C. acnes within all bacteria

decreased with age, and the proportion in individuals in their 40s was significantly lower

compared to those in their 20s and 30s ($p<0.05$, Fig.5).

C. acnes phylotypes

Next, we investigated the proportion of *C. acnes* subtypes in all 296 participants.

Subtype IA_1(42.8%) was the most common, followed by the subtype IA_2 (24.8%), Type II

(21.4%), subtype IB (10.0%), and type III (1.1%), as shown in Fig.6. When comparing the

proportions of C. acnes subtypes between the Acne group and the Non-acne group, there

was no difference observed in the proportions of IA_1, IA_2, Type II, and IB types (Data not

shown). On the other hand, for Type III, while the Acne group had a proportion of 0.6%, the

Non-acne group showed a higher proportion of 2.5%. The Acne group had a significantly

($p<0.005$) lower proportion of type III as shown in Fig.7.

Clonal complex of C. acnes

Upon further investigation of the detailed sequence types (ST) of *C. acnes*, the

constituent ratios, starting from the highest, were as follows: IA2_2_F0 23.9%, IA1_4_A0

20.6%, II_2_K0 18.6%, IA1_5_A2 11.1%, IB_1_H0 10.5%, IA1_6_D0 3.0%, IA1_5_A1

3.0%, II_3_K1 2.5%, III_1_L0 2.0%, IA2_1_F1 1.7% (Table 2). The result was sorted by

presence or absence of acne vulgaris. As a result, for ST IA1_1_E0, the Acne group had

2.4% while the non-acne group had 0.2%, showing a significant difference (p<0.05, without

correction). For ST IA2_1_F1, the Acne group had 1.5%, while the non-acne group had

3.5%, displaying a significant difference (p<0.05, without correction) (Table 3). Upon

comparison by age group, the biggest differences between ages 20-40 were found in

IA1_4_A0 (11.1%), IA1_5_A2 (6.6%), IA1_6_D0 (-6.0%), and IA2_2_F0 (-6.3%). However,

no statistically significant differences were observed.

Discussion

There is a growing demand for skin microbiome analysis from the perspective of skin disease treatment and beauty. Consequently, a technique that can sample skin bacteria easily and accurately is highly sought after. Various sampling methods, such as cotton swabs, scrapings, cyanoacrylate gel biopsy, and needle biopsy, have been devised and used in research for collecting the skin microbiome (flora)[20]. Among them, the swabbing method using a moistened cotton swab has been reported to be able to collect a bacterial flora equivalent to that of a biopsy, and it has been relatively commonly used[23,24]. However, the swab method had drawbacks, such as difficulty controlling efficiency depending on the sampling site and technique, and poor preservation.

In this study, we used a patch adhesive sticker with high preservation and adhesiveness (MySkin®), allowing subjects to sample microbes from their own facial surface. We evaluated samples sent by post and confirmed that there was no issue using them for microbiome analysis. Previously, Ogai *et al.* collected samples using either the cotton swab method or the sticker-stripping method and investigated the microbial composition using both next-generation sequencing and culture methods. They reported that the results of both methods were equivalent[24]. MySkin® is designed for even more convenient collection, such as having a constant seal area, and also possesses excellent

bacterial preservation properties, making it possible for subjects to collect samples

themselves. Indeed, this study demonstrated that samples could be easily collected and

sent by mail.

C. acnes is the most abundant resident bacterium of facial skin and is found in both

acne patients and healthy subjects[30,31]. Since it is anaerobic, its main niche is believed to

be the deeper area of the sebaceous hair follicles[32,33]. Thus, to study the relationship

between acne vulgaris and *C. acnes*, methods involving follicle or skin biopsy have been

primarily used[20]. However, follicle tissue sampling such as skin biopsy is more challenging.

One study that have employed swab sampling, *C. acnes* has been reported to constitute

more than 30% of the facial microbial community in acne patients[21]. On the other hand,

another report that sampled 55 facial acne patients using the swab method and found that

even in acne-affected areas, the proportion of propionic acid-producing bacteria such as*C.*

acnes was much lower (less than 2%)[22].

In this study, we obtained web questionnaires and image analysis from photographs

taken with smartphones from subjects to differentiate between the Acne group and the

Non-Acne group. It is widely known that the proliferation of *C. acnes* is a factor in the onset

of acne vulgaris[5,13,34]. In this study, we also examined the relative abundance of *C. acnes* in

the samples. As a result, the rate of *C. acnes* was significantly higher in the Acne group.

From this, it was believed that even with data from web questionnaires and images taken

with smartphones[35,36], one can grasp the skin condition of the subjects, and it is possible to

differentiate between the Acne group and the Non-Acne group.

Results of hierarchical clustering analysis on skin resident bacteria showed a group

where the microbial community was dominated by *C. acnes*, and the proportion of *C. acnes*

was high. When comparing the Acne group and the Non-Acne group, both the proportion

and copy number of *C. acnes* were significantly higher in the Acne group. However,

considering that many subjects were diagnosed with acne vulgaris even in groups

dominated by *Neisseriaceae* and others, it is unlikely that *C. acnes* alone has a pronounced

influence on the disease. Our new finding is that there might be various factors other than *C.*

acnes in the pathogenesis of acne vulgaris. Moreover, the alpha diversity (species

richness) of the facial skin microbial community was significantly lower in the Acne group

(Fig. 6-A). This result is consistent with the observation that the incidence of acne vulgaris

decreases with age, and α-diversity increases[37,38].

C. acnes is generally composed of several phylogenetic subtypes, commonly

referred to as IA_1, IA_2, IB, IC, II, III[17]. At the genome level, even finer classifications are

possible[17,39]. In this study, a genetic analysis was conducted to determine the phylotype

ratio of *C. acnes* in all subjects. IA_1 was found to be the most prevalent type, consistent with

a previous report[40]. It has been reported that the IA_1 type, isolated from common acne,

possesses strong pathogenicity[13,18,19]. In the hair follicles of acne patients, the dominant

strain of *C. acnes* is IA_1. On the surface of the face, not only IA_1 but also IA_2, IB, and II are

present[40,41]. When we examined the phylotype of *C. acnes* on the facial skin surface, we

found a cocktail of IA_1, IA_2, IB, II, and III, with no significant difference between the Acne

and Non-Acne groups. This result may be due to the wider area of sampling rather than

pinpoint acne lesions. Meanwhile, type III, which showed a significant difference between

the groups, has been found in normal skin but not in common acne areas[14,19]. In our

research, type III was confirmed in both Acne and Non-acne groups. It's suggested that the

reason that type III, which is rarely reported in the face, was found in the Acne group might

be due to the inability to strictly separate common acne lesion from surrounding areas

during sticker sampling. Further detailed verification is needed in the future, but for narrow

lesion sampling, biopsies or swabs might be suitable. Type III is reported to lack toxic

factors and has a different morphology and metabolic pathway than types I and II[42]. In our

results, type III in the Acne group showed significantly lower values compared to the

Non-acne group. Further analysis is needed to understand how the decreased expression

of type III in normal skin areas relates to the onset of common acne.

When examining the detailed sequence types of *C. acnes*, the strains with the

highest proportion in all subjects were IA2_2_F0 (23.9%), IA1_4_A0 (20.6%), and II_2_K0

(18.6%). IA2_1_F1 ($p < 0.05$ vs non-acne group) higher in acne patients, while III_1_L0 was

lower ($p < 0.05$ vs non-acne group) (Table3, Supplemental figure 2). Also, our study revealed

that various types of *C. acnes* phylotypes were expressed on the skin surface of the face.

Our method of using specialized sticker to collect samples from specific facial areas

captures both acne and healthy tissues. This may indicate skin conditions prone to

developing acne. Dagnelie *et al*. reported that the detailed phylotype expression patterns of

C. acnes differ with the onset of acne[40]. In the future, increasing the number of subjects and

conducting larger-scale analyses may clarify statistical differences between groups.

In this study, we could successfully conduct postally collected skin microbiome

samples and made analysis by online basis from customers of a company.While

preliminary, our results suggest the potential for online consultations to provide diagnostic

quality comparable to in-person assessments in testing and diagnosing skin-related

diseases. Especially in the post-COVID era, the advancement of such research could

significantly contribute to the diversification of diagnostic methods and the development of

dermatology.

References

1.   Skowron, K. *et al.* Human Skin Microbiome: Impact of Intrinsic and Extrinsic Factors on Skin Microbiota. *Microorganisms* **9**, (2021).

2.   Smythe, P. & Wilkinson, H. N. The Skin Microbiome: Current Landscape and Future Opportunities. *Int. J. Mol. Sci.* **24**, (2023).

3.   Chilicka, K., Dzieńdziora-Urbińska, I., Szyguła, R., Asanova, B. & Nowicka, D. Microbiome and Probiotics in Acne Vulgaris-A Narrative Review. *Life* **12**, (2022).

4.   Chen, L. *et al.* Skin and Gut Microbiome in Psoriasis: Gaining Insight Into the Pathophysiology of It and Finding Novel Therapeutic Strategies. *Front. Microbiol.* **11**, 589726 (2020).

5.   Huang, C. *et al.* The updates and implications of cutaneous microbiota in acne. *Cell Biosci.* **13**, 113 (2023).

6.   Fyhrquist, N. *et al.* Microbe-host interplay in atopic dermatitis and psoriasis. *Nat. Commun.* **10**, 4703 (2019).

7.   Christensen, G. J. M. & Brüggemann, H. Bacterial skin commensals and their role as host guardians. *Benef. Microbes* **5**, 201–215 (2014).

8.   Allhorn, M., Arve, S., Brüggemann, H. & Lood, R. A novel enzyme with antioxidant capacity produced by the ubiquitous skin colonizer Propionibacterium acnes. *Sci. Rep.* **6**, 36412 (2016).

9.   Elston, M. J., Dupaix, J. P., Opanova, M. I. & Atkinson, R. E. Cutibacterium acnes (formerly Proprionibacterium acnes) and Shoulder Surgery. *Hawaii J Health Soc Welf* **78**, 3–5 (2019).

10.   Yang, M., Zhou, M. & Song, L. A review of fatty acids influencing skin condition. *J. Cosmet. Dermatol.* **19**, 3199–3204 (2020).

11.   O'Neill, A. M. & Gallo, R. L. Host-microbiome interactions and recent progress into understanding the biology of acne vulgaris. *Microbiome* **6**, 177 (2018).

12.   Ahle, C. M., Feidenhansl, C. & Brüggemann, H. Cutibacterium acnes. *Trends Microbiol.* **31**, 419–420 (2023).

13.   Fitz-Gibbon, S. *et al.* Propionibacterium acnes strain populations in the human skin microbiome associated with acne. *J. Invest. Dermatol.* **133**, 2152–2160 (2013).

14.   McDowell, A. *et al.* An expanded multilocus sequence typing scheme for propionibacterium acnes: investigation of 'pathogenic', 'commensal' and antibiotic resistant strains. *PLoS One* **7**, e41480 (2012).

15.   Lomholt, H. B. & Kilian, M. Population genetic analysis of Propionibacterium acnes identifies a subpopulation and epidemic clones associated with acne. *PLoS One* **5**, e12277 (2010).

16.   Dekio, I., Asahina, A. & Shah, H. N. Unravelling the eco-specificity and pathophysiological properties of Cutibacterium species in the light of recent taxonomic

changes. *Anaerobe* **71**, 102411 (2021).

17.   Mayslich, C., Grange, P. A. & Dupin, N. Cutibacterium acnes as an Opportunistic

Pathogen: An Update of Its Virulence-Associated Factors. *Microorganisms* **9**, (2021).

18.   McLaughlin, J. *et al.* Propionibacterium acnes and Acne Vulgaris: New Insights

from the Integration of Population Genetic, Multi-Omic, Biochemical and Host-Microbe

Studies. *Microorganisms* **7**, (2019).

19.   McDowell, A., Nagy, I., Magyari, M., Barnard, E. & Patrick, S. The opportunistic

pathogen Propionibacterium acnes: insights into typing, human disease, clonal

diversification and CAMP factor evolution. *PLoS One* **8**, e70897 (2013).

20.   Platsidaki, E. & Dessinioti, C. Recent advances in understanding Propionibacterium

acnes (Cutibacterium acnes) in acne. *F1000Res.* **7**, (2018).

21.   Grice, E. A. *et al.* Topographical and temporal diversity of the human skin

microbiome. *Science* **324**, 1190–1192 (2009).

22.   Dreno, B. *et al.* Skin microbiome and acne vulgaris: Staphylococcus, a new actor in

acne. *Exp. Dermatol.* **26**, 798–803 (2017).

23.   Bjerre, R. D. *et al.* Effects of sampling strategy and DNA extraction on human skin

microbiome investigations. *Sci. Rep.* **9**, 17287 (2019).

24.   Dekio, I. *et al.* Detection of potentially novel bacterial components of the human

skin microbiota using culture-independent molecular profiling. *J Med Microbiol.* **54**,

1231-1238 (2005).

25.   Inoue, F., Tachibana, K. & Sutani, T. Adhesive tape for collecting skin indigenous

microorganisms, method of evaluating physical condition of subject, method of presenting

information to subject and method of screening for substance capable of improving or

preventing physical condition, and adhesive tape for collecting skin metabolites. *US Patent*

Publication number 20220338795 (2022).

26.   Shannon, C. E. A mathematical theory of communication. *The Bell System*

Technical Journal **27**, 379–423 (1948).

27.   Shimizu, H. (2016). An Introduction to the Statistical Free Software HAD

Suggestions to Improve Teaching, Learning and Practice Data Analysis. Journal of Media,

Information and Communication, 1, 59-73. - references - scientific research publishing.

https://www.scirp.org/(S(351jmbntvnsjt1aadkozje))/reference/ReferencesPapers.aspx?Ref

erenceID=1990046.

28.   Fukukawa, Y., Okazaki, M. & Ohkusu, K. A case of Rothia mucilaginosa

(Stomatococcus mucilaginosus) bacteremia in a patient with acute myelocytic leukemia and

evaluation of the ability to identify for this organism used identification kits. *The journal of*

the Japanese Society for Clinical Microbiology **20**, 50–55 (2010).

29.   Mehta, V. *et al. Wickerhamomyces anomalous*: A Rare Cause of Fungemia

Causing Febrile Neutropenia in Acute Lymphoblastic Leukemia. *Case Rep. Infect. Dis.*

2020, 8847853 (2020).

30.   Rozas, M. *et al.* From Dysbiosis to Healthy Skin: Major Contributions of

Cutibacterium acnes to Skin Homeostasis. *Microorganisms* **9**, (2021).

31.   Miura, Y. *et al.* Quantitative PCR of Propionibacterium acnes DNA in samples

aspirated from sebaceous follicles on the normal skin of subjects with or without acne. *J.*

Med. Dent. Sci. **57**, 65–74 (2010).

32.   Gehse, M., Höffler, U., Gloor, M. & Pulverer, G. Propionibacteria in patients with

acne vulgaris and in healthy persons. *Arch. Dermatol. Res.* **275**, 100–104 (1983).

33.   Bek-Thomsen, M., Lomholt, H. B., Scavenius, C., Enghild, J. J. & Brüggemann, H.

Proteome analysis of human sebaceous follicle infundibula extracted from healthy and

acne-affected skin. *PLoS One* **9**, e107908 (2014).

34.   Dréno, B., Dagnelie, M. A., Khammari, A. & Corvec, S. The Skin Microbiome: A

New Actor in Inflammatory Acne. *Am. J. Clin. Dermatol.* **21**, 18–24 (2020).

35.   Lee, S. *et al.* Sample-to-answer platform for the clinical evaluation of COVID-19

using a deep learning-assisted smartphone-based assay. *Nat. Commun.* **14**, 2361 (2023).

36.   Mikołajczyk, M., Patrzyk, S., Nieniewski, M. & Woźniacka, A. Evaluation of a

smartphone application for diagnosis of skin diseases. *Postepy Dermatol Alergol* **38**,

761–766 (2021).

37.   Jugé, R. *et al.* Shift in skin microbiota of Western European women across aging. *J.*

Appl. Microbiol. **125**, 907–916 (2018).

38.   Heng, A. H. S. & Chew, F. T. Systematic review of the epidemiology of acne vulgaris. *Sci. Rep.* **10**, 5754 (2020).

39.   Both, A. *et al.* Genomics of Invasive Cutibacterium acnes Isolates from Deep-Seated Infections. *Microbiol Spectr* **11**, e0474022 (2023).

40.   Dagnelie, M.-A. *et al.* Decrease in Diversity of Propionibacterium acnes Phylotypes in Patients with Severe Acne on the Back. *Acta Derm. Venereol.* **98**, 262–267 (2018).

41.   Lomholt, H. B., Scholz, C. F. P., Brüggemann, H., Tettelin, H. & Kilian, M. A comparative study of Cutibacterium (Propionibacterium) acnes clones from acne patients and healthy controls. *Anaerobe* **47**, 57–63 (2017).

42.   McDowell, A., Perry, A. L., Lambert, P. A. & Patrick, S. A new phylogenetic group of Propionibacterium acnes. *J. Med. Microbiol.* **57**, 218–224 (2008).

Figure legends

Figure 1

A simplified schema summarizing the skin microbiome analysis. Skin sampling was

conducted using the sticker-stripping method (for 1 minute, before face wash in the

morning).

Figure 2

Taxonomic composition of bacteria (A) and fungi (B) based on *16S rRNA* and ITS

sequences. Acneic subjects are highlighted in red bar at the bottom of figure.

Figure 3

Alpha diversity (Shannon index) of facial skin bacteria (A) or fungi (B) microbiome

samples from acne (n=56) or non-acne (n=20) subjects. The lines inside the rectangles

represent the median, and the whiskers indicate the maximum and minimum values.

Statistical differences were calculated using Welch's *t*-test with Holm's correction

Figure 4

The percentage (A) and log copy number (B) of *C. acnes* among total bacteria in

facial skin samples from acne (n=219) or non-acne (n=77) subjects. Statistical differences

were calculated using Welch's *t*-test with Holm's correction.

Figure 5

Changes in the proportion of *C. acnes* in total bacteria in facial skin samples were

observed among the following age groups: 20s (n=43), 30s (n=181) and 40s (n=70).

Statistical differences were calculated using one-way ANOVA followed by the Holm's *post*

hoc test.

Figure 6

C. acnes phylotype composition of all subjects.

Figure 7

The percentage comparison among *C. acnes* phylotype in facial skin samples from

acne (n=56) or non-acne (n=20) subjects. Statistical differences were calculated using

Welch's *t*-test with Holm's correction.

Tables

Table 1. The composition of *C. acnes* strain level phylotypes

Level4	Acne(%)	Non-acne(%)	Defference(%)
II_2_K0	17.5	21.8	4.3
IA1_1_E0	0.1	2.5	2.4
III_1_L0	1.5	3.5	2.0
IA1_6_D0	2.6	4.4	1.8
IA1_5_A1	3.3	2.3	-1.0
II_3_K1	2.8	1.6	-1.2
IA1_5_A2	11.5	10.0	-1.5
IA2_1_F1	2.3	0.2	-2.1
IA1_4_A0	21.2	18.9	-2.3

*$p<0.05$ vs. Non-acne

Table 2 The composition of *C. acnes* strain level phylotypes

Strain types	Acne (%)	Non-acne (%)	Defference (%)
II_2_K0	17.5	21.8	4.3
IA1_1_E0	0.1	2.5	2.4
III_1_L0	1.5	3.5	2.0
IA1_6_D0	2.6	4.4	1.8
IA1_5_A1	3.3	2.3	-1.0
II_3_K1	2.8	1.6	-1.2
IA1_5_A2	11.5	10.0	-1.5
IA2_1_F1	2.3	0.2	-2.1
IA1_4_A0	21.2	18.9	-2.3

*$p<0.05$ vs. Non-acne

Fig. 1

A Bacteria

C. acnes *Neisseriaceae* *Streptococcus*

B Fungi

Malassezia_restricta *Malassezia_globosa* *Wickerhamomyces_anomalus* *Rhodotorula_mucilaginosa*

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7