Associations between music and dance relationships, rhythmic proficiency, and spatiotemporal movement modulation ability in adults with and without mild cognitive impairment

Alexandra Slusarenko[a,†], Michael C. Rosenberg[b,†], Meghan E. Kazanski[c], J. Lucas McKay[d,e], Laura Emmery[f], Trisha M. Kesar[g], Madeleine E. Hackney[c,h-j]*

[a]College of Arts and Sciences, Emory University, Atlanta, GA, USA
[b]Neuromechanics Laboratory, Department of Biomedical Engineering, Emory University & Georgia Institute of Technology, Atlanta, GA, USA
[c]Department of Medicine, Division of Geriatrics and Gerontology, Emory University School of Medicine, Atlanta, GA, USA
[d]Department of Neurology, Emory University School of Medicine, Atlanta, GA, USA
[e]Department of Biomedical Informatics, Emory University School of Medicine, Atlanta, GA, USA
[f]Department of Music, Emory University College of Arts and Sciences, Atlanta, GA, USA
[g]Department of Rehabilitation Medicine, Emory University School of Medicine, Atlanta, GA, USA
[h]Emory University School of Nursing, Atlanta, GA, USA
[i]Atlanta VA Center for Visual & Neurocognitive Rehabilitation, Atlanta, GA, USA
[j]Birmingham/Atlanta VA Geriatric Research Education and Clinical Center, Atlanta, GA, USA

Running title: Dance, music, and movement ability in MCI

†Alexandra Slusarenko and Michael C. Rosenberg contributed equally to this work.

*Correspondence:
Madeleine E. Hackney
mehackn@emory.edu
314-412-4852

Keywords: Alzheimer's disease, mild cognitive impairment, rehabilitation, therapy, gait analysis, dance, music, rhythm

Abstract

Background: Personalized dance-based movement therapies may improve cognitive and motor function in individuals with mild cognitive impairment (MCI), a precursor to Alzheimer's disease. While age- and MCI-related deficits reduce individuals' abilities to perform dance-like rhythmic movement sequences (RMS)—spatial and temporal modifications to movement—it remains unclear how individuals' relationships to dance and music affect their ability to perform RMS.

Objective: Characterize associations between RMS performance and music or dance relationships, as well as the ability to perceive rhythm and meter (rhythmic proficiency) in adults with and without MCI.

Methods: We used wearable inertial sensors to evaluate the ability of 12 young adults (YA; age=23.9±4.2 yrs; 9F), 26 older adults without MCI (OA; age=68.1±8.5 yrs; 16F), and 18 adults with MCI (MCI; age=70.8±6.2 yrs; 10F) to accurately perform spatial, temporal, and spatiotemporal RMS. To quantify self-reported music and dance relationships and rhythmic proficiency, we developed Music (MRQ) and Dance Relationship Questionnaires (DRQ), and a rhythm assessment (RA), respectively. We correlated MRQ, DRQ, and RA scores against RMS performance for each group separately.

Results: The OA and YA groups exhibited better MRQ and RA scores than the MCI group ($p < 0.006$). Better MRQ and RA scores were associated with better temporal RMS performance for only the YA and OA groups ($r^2 = 0.18\text{-}0.41$; $p < 0.045$). DRQ scores were not associated with RMS performance in any group.

54   **Conclusions:** Cognitive deficits in adults with MCI likely limit the extent to which music relationships or

55   rhythmic proficiency improve the ability to perform temporal aspects of movements performed during

56   dance-based therapies.

Introduction

Dance-based movement therapy is a cognitively engaging physical activity that helps mitigate

neurodegeneration and improve cognitive and motor function in individuals with mild cognitive

impairment (MCI), a precursor to Alzheimer's disease and dementia [1-5]. The level of motor and

cognitive challenge of dance-based therapies may be customized by selecting different therapy

parameters (e.g., prescribed dance movements or musical elements). Selecting therapy parameters that

challenge each individual, without being discouragingly difficult, may enhance therapeutic efficacy [3, 6].

However, we currently lack objective approaches to personalize dance-based therapy parameters [1, 7].

The ability to perform dance-based therapies likely depends on multiple factors, including aspects of

motor and cognitive function, as well as each individual's relationships to music and dance (defined as

individuals' histories, experiences, and attitudes towards music and dance) and rhythmic proficiency

(defined as individuals' abilities to perceive and replicate rhythms) [1, 3, 5, 7]. While we previously

showed that age-related declines in motor and cognitive function reduce the ability to perform dance-

like movements, it remains unclear how relationships with dance and music, or rhythmic proficiency,

impact this ability [8].

Assaying an individual's ability to perform dance-like movements is critical for determining challenging,

engaging, and individual-specific dance therapy protocols. We recently developed a library of Rhythmic

Movement Sequences (RMS) that isolate spatial (e.g., modified lower-extremity joint range of motion

and coordination) and temporal (e.g., modified timing of stepping patterns, synchronized to music)

features of forward movement (*i.e.*, walking) [8, 9]. Note that "RMS" used throughout the manuscript

refers to the movement sequences, not "root-mean-square" error. Similar to RMS, dance therapies

often involve modifications to spatial and temporal aspects of movement, but lack principles to

determine modification difficulty based on participants' abilities [3, 5, 10]. By requiring participants to

walk while modifying either spatial or temporal aspects of movement, or both simultaneously, the RMS

experimental paradigm provides insights regarding participant's abilities to modify spatial and temporal

aspects of movement that may be performed during therapy.

RMS performance can be quantified as the ability to achieve prescribed spatial and temporal targets [8].

Spatial RMS consist of modifications to the joint kinematics of typical walking to achieve prescribed

spatial targets (e.g., 90 degrees of peak hip flexion during the swing phase of gait), with no prescribed

changes in step timing [11]. Deviations from spatial modification targets may reflect an inability to

perceive, comprehend, or recall spatial patterns and execute appropriate motor commands to

accurately modulate spatial aspects of movement. Temporal RMS involve modifying step timing to

achieve prescribed patterns of quick and slow steps, synchronized to perceived concurrent rhythmic

cues in music, with no prescribed spatial modifications [9]. Our approach to temporal RMS is grounded

in principles of music theory that suggest that auditory cues can influence the temporal progression of

movement [12-14]. Deviations from the prescribed tempo or step pattern reflect an inability to perceive

musical cues and recall or execute stepping patterns. Spatiotemporal RMS assays the additional

challenge of simultaneously performing spatial and temporal RMS. The performance of these three

types of RMS may be differentially influenced by multiple factors including motor function, cognitive

function, and experience perceiving rhythms and executing motor commands.

We previously showed detrimental effects of age and MCI on individuals' abilities to perform RMS,

though these factors did not comprehensively explain individual differences in RMS performance [8].

We found worse performance on spatial, but not temporal RMS in older adults without MCI, compared

to young adults, suggesting that age-related motor deficits are primarily related to a reduced ability to

accurately modulate spatial features of movement. Conversely, we found worse performance on some

spatial and some temporal RMS in older adults with MCI, compared to older adults without MCI,

suggesting that cognitive deficits in adults with MCI are related to a reduced ability to accurately modify

both spatial and temporal features of movement. However, we found that RMS performance was not

fully explained by differences in age or cognitive status [8]. Individual differences in the ability to

perceive meter and rhythm and synchronize motor commands to music may help explain variability in

the temporal aspects of RMS performance in individuals of similar age and cognitive status [14, 15].

Similarly, individual differences in motor experiences stemming from music or dance relationships may

explain variability in the spatial aspects of RMS performance with similar age-related motor function

levels [11, 15].

Relationships to music and dance reflect individuals' diverse histories, perceptions, and attitudes

towards music and dance, while rhythmic proficiency reflects the ability to entrain motor commands to

perceived rhythms. Merely by existing within a cultural context and through past experiences,

individuals develop such relationships with these ubiquitous and often interconnected art forms [14,

16]. Lifelong exposure to music and dance may alter the neural and biomechanical processes underlying

the ability to perceive auditory cues and produce corresponding rhythmic motor commands, both of

which are required during dance-based therapies [3, 11, 15, 17, 18]. Stronger relationships with music

may enhance the ability to perceive and predict the rhythmic patterns, musical groupings, and meter

needed to follow temporal movement cues in music [15, 19, 20]. Stronger relationships with dance may

enhance the ability to sense and accurately modify spatial aspects of movement (e.g., joint kinematics)

and entrain movements to musical cues [11, 15]. Conversely, stronger rhythmic proficiency reflects a

better ability to perceive rhythm and meter, anticipate beats in music, and entrain motor commands to

prescribed rhythmic cues [14]. Rhythmic proficiency, therefore, reflects both perceptual acuity and

motor skill pertinent to RMS performance, making it distinct from music and dance relationships [21].

Here, we investigated how individuals' relationships with dance and music, and their rhythmic

proficiency impacted their performance on spatial, temporal, and spatiotemporal RMS. The central

hypothesis of this work is that stronger relationships and past experiences with music and dance, and

rhythmic proficiency, contribute to an improved ability to accurately modulate spatial and temporal

aspects of movement, as captured during RMS. To test this hypothesis, we developed Music

Relationship (MRQ) and Dance Relationship Questionnaires (DRQ) that survey individuals' relationships

to music and dance, respectively. We also developed a Rhythm Assessment (RA) that assesses rhythmic

proficiency, defined by the ability to recognize meter in music and reproduce auditory or visual rhythms.

First, we characterized the relationships between the MRQ, DRQ, and RA. We then compared these

novel assessment scores to RMS performance in younger adults (YA group) and older adults without (OA

group) and with MCI (MCI group). We predicted that, within each group, stronger relationships to music

(higher MRQ scores) would be associated with more accurate performance of temporal and

spatiotemporal RMS. We similarly predicted that, within groups, stronger relationships to dance (higher

DRQ scores) would be associated with more accurate performance of spatial and spatiotemporal RMS.

Finally, we predicted that better rhythmic proficiency (higher RA scores) would be associated with more

accurate performance of temporal and spatiotemporal RMS. Furthermore, to better understand how

music-related perceptual acuity and motor skill impact RMS performance, we conducted an exploratory

correlational analysis regressing two RA subcomponents—*auditory rhythm reproduction* and *auditory*

meter recognition—against RMS performance.

150   Materials and Methods

This study was approved by the Emory University Institutional Review Board (STUDY00003507). All

participants provided written, informed consent before participation.

Participants

An observational cross-sectional study with 56 participants was performed in non-disabled younger

adults (YA; N=12), older adults without MCI (OA; N = 26), and older adults with MCI (N = 18; Table 1).

The inclusion criteria for all participants were the ability to walk 20m without an assistive device, 6 years

of education (e.g., kindergarten through fifth grade), proficiency in the English language, and no

hospitalizations in the last 60 days. YA participants included in the study were 18-35 years of age and OA

and MCI participants were 55 years and older. For participants with MCI, additional inclusion criteria

included amnestic MCI, as defined using the Alzheimer's Disease Neuroimaging Initiative criteria and

standard clinical assessments showing reduced executive function, working memory, and spatial

cognition [22]. Assessments characterizing cognitive function included the Montreal Cognitive

Assessment (MoCA), Reverse Corsi Blocks, Body Position Spatial Task, and Trail Making Test (Table 1)

[23-26]. Table 1 shows the average (±1 SD) participant demographic information and clinical assessment

scores for each group, along with p-values reflecting the probability of statistical differences across the

groups (omnibus ANOVA or Chi-Squared tests, where appropriate; α = 0.05). When we identified

significant differences across groups, we performed post-hoc independent-samples t-tests to identify

differences between groups the YA and OA groups (daggers in Table 1; α = 0.05) and the OA and MCI

groups (double-daggers in Table 1; α = 0.05).

Table 1: Demographic characteristics for each participant group.

	YA	OA	MCI	p-value*
N	12	26	18	
Age*,†				< 0.01
Mean (SD)	23.9 (4.2)	68.1 (8.5)	70.8 (6.2)	
Range	18.0 – 30.0	53.0 – 85.0	57.0 – 79.0	
Height (m)				0.14
Mean (SD)	1.65 (0.09)	1.73 (0.13)	1.69 (0.09)	
Range	1.57 – 1.91	1.56 – 2.07	1.52 – 1.83	

	YA	OA	MCI	
Mass (kg)*,†				0.03
Mean (SD)	62.5 (12.7)	73.1 (15.1)	76.3 (13.6)	
Range	49.9 – 90.7	51.7 – 100.0	52.6 – 108.9	
Sex				0.55
F	9 (75%)	16 (62%)	10 (56%)	
M	3 (25%)	10 (38%)	8 (44%)	
Years since MCI[a]				-
	-	-	2.7 (1.9)	
	-	-	0.0-7.0	
Montreal Cognitive Assessment (MoCA)[b],*,‡				< 0.01
Mean (SD)	28.4 (2.2)	27.4 (2.3)	24.8 (3.6)	
Range	22.0 – 30.0	23.0 – 30.0	18.0 – 30.0	
Reverse Corsi Blocks[c],*,†,‡				< 0.01
Mean (SD)	59.6 (21.2)	37.3 (14.7)	27.4 (14.2)	
Range	15.0 – 98.0	20.0 – 77.0	12.0 – 54.0	
Body Position Spatial Task[d],*,†				< 0.01
Mean (SD)	31.2 (15.5)	17.4 (7.9)	14.0 (6.0)	
Range	12.0 – 70.0	6.0 – 35.0	4.0 – 25.0	
Trail Making Test (B-A)[e],*,‡				0.03
Mean (SD)	24.3 (13.9)	29.7 (21.6)	44.9 (26.0)	
Range	7.9-50.4	-28.6 – 68.4	-19.7 – 92.5	

Abbreviations: N, number of participants; YA, younger adults; OA, older adults without mild cognitive impairment;
MCI, older adults with mild cognitive impairment; m, meters; kg, kilograms; SD, standard deviation; F, female; M,
male.
*Significant between-group differences (omnibus ANOVA or Chi-Squared tests, where appropriate; α = 0.05).
†Significant differences between the YA and OA groups (post-hoc independent-samples t-tests; α = 0.05).
‡Significant differences between the OA and MCI groups (post-hoc independent-samples t-tests; α = 0.05).
Superscript letters denote the number of missing participants for each analysis: [a]N = 13 (MCI group only); [b]N = 55;
[c]N = 54, [d]N = 52, [e]N = 53.
We did not test for differences between the YA and MCI groups.

Rhythmic Movement Sequences (RMS)

Participants performed an overground assessment battery of 9 *spatial*, 9 *temporal*, and 4

spatiotemporal RMS, using the same protocol as described by Rosenberg and colleagues (2023) [8]. Each

RMS consisted of spatial and/or temporal modifications to movement: Spatial RMS involved modifying

leg joint angles during the stance or swing phases during forward movement (i.e., walking), or during

both phases of gait. While dance does not always prescribe strict joint angles, and can focus instead on

movement fluidity or interpersonal interactions, control of one's kinematic configuration remains an

essential aspect of some dance styles and therapies [5, 7, 17, 27]. Each spatial RMS consisted of two

different spatial *gait modifications* to leg joint kinematics, with one modification per leg or both

modifications in the same leg. Spatial modifications did not prescribe step timing or rhythms. The spatial

modifications used in this study have corollaries in ballet, alter the typical leg joint flexion-extension

patterns of walking, and were designed to be feasible for OA and individuals with MCI [28]. We defined

three sub-classes of spatial RMS: *swing*, *stance*, and *swing-stance* (Figure 1). RMS sub-class names were

defined by whether participants were instructed to modify joint kinematics during swing, stance, or both

phases of gait, respectively. Each sub-class contained three different RMS. For *swing* and *stance* RMS,

different gait modifications were performed by each leg. For example, the *Attitude-Developpé* RMS

involved performing the *Attitude* and *Developpé* modifications during the swing phases of the left and

right legs, respectively (Figure 1; bullets denote target joint angles). Similarly, for *swing-stance* RMS, two

different modifications were performed by the left leg, while the right leg could move freely. For

example, the *Attitude-Relevé* modification involved performing the *Attitude* and *Relevé* modifications

both with the left leg during the swing and stance phases of gait, respectively.

Swing		Stance	
Spatial target	Gait modification	Spatial target	Gait modification
▪ 45° knee flexion ▪ Max. ankle plantarflexion ▪ Toe at stance leg ankle	Coupé	▪ 0° knee flexion ▪ Max. ankle plantarflexion throughout stance	Piqué
▪ 90° knee flexion ▪ Max. knee flexion ▪ Toe at stance leg knee	Passé	▪ 30° knee flexion ▪ 30° ankle dorsiflexion	Tombé
▪ 90° hip flexion ▪ 90° knee flexion ▪ Max. ankle plantarflexion	Attitude	▪ 0° knee flexion ▪ Max. ankle dorsiflexion	Heel strike
▪ 90° hip flexion ▪ Max. early swing knee flexion ▪ 0° late-swing knee flexion ▪ Max. ankle plantarflexion	Developpé	▪ 0° knee flexion ▪ Max. ankle plantarflexion in mid-stance	Relevé
▪ 90° hip flexion ▪ 0° knee flexion ▪ Max. ankle plantarflexion	Battement		

Figure 1: **Spatial modifications and corresponding targets used in spatial and spatiotemporal rhythmic movement sequences (RMS)**. A) Spatial modifications. The left two columns correspond to modifications to swing-phase kinematics during movement, while the right two columns correspond to modifications to stance-phase kinematics. The bullets describe joint kinematics defining each spatial target variable for the corresponding modification. Deviations from these target values quantified RMS performance. The colored lines denote the hip (purple), knee (orange), and ankle (red) target values. *Adapted, with permission, from Rosenberg et al., 2023 [8].*

Temporal RMS involved walking while performing repeating sequences of 2-6 steps of varying duration,

with footfalls synchronized to musical pieces that served as external rhythmic cues [8]. Each temporal

RMS consisted of only one temporal gait modification. The spatial aspects of movement were not

constrained; participants could walk with any gait pattern and any speed that let them achieve the

desired step sequence. Steps were performed at different rates relative to prescribed music: half steps

spanned half of a beat (*&* in Figure 2), quick (q) steps spanned one beat (solid note in Figure 2), and slow

(S) steps spanned two beats (open note in Figure 2). We defined three meter-based subclasses of

temporal RMS from ballroom dance: *simple duple* (2-count), *complex duple* (2-count), and *waltz* (3-

count) metric modifications, shown in Figure 2. Each RMS subclass contained three different step

sequences, performed to the same rhythm and meter. For example, the *simple duple – qqSS* RMS

involved taking two quick steps at a cadence of one step per beat, followed by two slow steps at a

cadence of one step every two beats. Each modification class was expected to challenge different

aspects of Western music listeners' experiences [8, 14]. Participants performed duple and waltz

modifications synchronously to modified versions of Libertango (by Astor Piazzolla, 1974) and Waltz No.

2 (by Dmitri Shostakovich, 1938), respectively, with superfluous accents and cues removed [8, 29]. All

participants performed the duple modifications at 100 beats per minute (bpm). For the waltz

modifications, the YA groups performed modifications to music at 80 bpm, while the music was slowed

to 60 bpm for the OA and MCI groups to ensure that they could perform the modifications. Empirical

tapping studies and theoretical models of motor-auditory synchronization suggest that the OA and MCI

groups could be expected to exhibit better synchronization to the waltz 60bpm tempo than would the

YA group to the 80bpm tempo [30-33]. However, we previously found that any differences in motor-

auditory synchronization at 60bpm were small compared to within-group variability in RMS

performance: the YA group did not perform waltz RMS less accurately than the OA group [8].

A) Simple Duple (2-count)

B) Complex Duple (2-count)

C) Waltz (3-count)

Figure 2: **Rhythmic stepping sequences used in temporal and spatiotemporal rhythmic movement sequences (RMS)**. Each sequence consisted of 2-6 steps, synchronized to either a duple (2-count) or waltz (3-count) meter. Sequences were comprised of very-quick (*&* = half-beat per step), quick (*q* = one beat per step), and slow (*S* = two beats per step) steps. The numbers above each musical note reflect the beat count. A) Simple duple sequences were two-count rhythms with the strong beat on the downbeat and spanned 1-2 measures. B) Complex duple sequences were also two-count rhythms with a weak beat on the downbeat and spanned 2 measures. C) Waltz sequences were three-count rhythms spanning 1-2 measures. *Re-used, with permission, from Rosenberg et al., 2023 [8].*

Each spatiotemporal RMS involved performing spatial and temporal modifications. Participants

performed four spatiotemporal RMS consisting of different combinations of two spatial modifications

and one temporal modification (Figure 1 & 2).

The RMS trial order was block-randomized for each participant. Spatial or temporal RMS were randomly

selected to be performed first and second, followed by spatiotemporal RMS. Within each RMS class

(spatial, temporal, spatiotemporal), the order of RMS sub-classes was randomized, as was the order of

the three RMS within each subclass.

RMS protocol

Participants performed each of the 22 RMS once. For each RMS, YA performed RMS while walking

overground for four lengths of an 11-meter walkway. To mitigate the effects of fatigue on RMS

performance, participants in the OA and MCI groups performed RMS for at least 11 meters (one

walkway length) [8]. Participants who walked shorter distances also took more strides per walkway

length. All participants performed each RMS for at least 15 strides. Participants could take as much time

as needed to safely complete each trial. Across participants, trials typically lasted less than 180s, though

longer trials took up to 360s. However, longer trials typically occurred once per participant, after which

we reduced the number of walkway lengths to mitigate fatigue.

During RMS assessments, sagittal-plane hip, knee, and ankle kinematic timeseries were recorded at 128

Hz using Opal V2R inertial measurement units (APDM, Inc., Portland, USA). Fifteen sensors were

attached to the forehead, sternum, lumbar region and bilaterally to the hands, wrists, upper arms,

thighs, shanks, and feet in a standard configuration [34]. For each trial, joint kinematics were estimated

using validated proprietary software (APDM Moveo Explorer) [35]. Computation of RMS performance is

described in the next section.

Before performing each spatial RMS, participants watched a tutorial video of the sequence (two spatial

modifications) being performed by an expert and received instructions on how to achieve the spatial

targets of the movement (*i.e.*, discrete joint angle targets; Figure 1) [8]. Before performing each

temporal RMS, participants watched a tutorial video that guided progressive entrainment of the trial's

rhythmic pattern in five steps of increasing complexity: clapping, tapping with one foot, shifting weight

between the feet, marching in place, and walking to the rhythmic pattern. Our prior work found this

approach to be an effective tool to teach new rhythms to dance-based therapy participants [17]. We

found the technique useful for participants in the present study who had little music and dance

experience. Before performing each spatiotemporal RMS, participants were reminded of the spatial

targets (spatial) and step sequences (temporal) and could review the tutorial videos if needed.

Participants practiced each RMS with assessor feedback until the assessor determined that the

participant understood the RMS. Practice typically spanned less than one walkway length. At the

beginning of each walkway length during temporal trials, the assessor would clap the rhythm for two

sequences to help participants identify the rhythm.

RMS performance targets and quantification

Performance of each RMS was quantified as each participant's ability to achieve pre-defined spatial

targets (spatial and spatiotemporal RMS) and temporal targets (temporal and spatiotemporal RMS).

Targets were specific to each modification and are described in Figure 1 & 2. Spatial targets were

defined by sagittal-plane joint angles of the hip, knee, and/or ankle in the stance or swing phases for

each RMS (Figure 1 & 3A). For example, the *Attitude* modification involved joint angle targets of 90-

degree hip flexion, 90-degree knee flexion, and maximal ankle plantarflexion (Figure 3A, middle; stars

denote target values).

Figure 3: Calculation of error on spatial and temporal modifications comprising RMS. The left column shows timeseries data. The middle column depicts how percent errors computed within spatial and/ or temporal modifications comprising a given RMS. The right column depicts how percent errors within multiple modifications combine to produce error for each RMS. Dashed lines denote spatial and temporal targets. Yellow dots denote the portion of the stride (e.g., swing vs. stance) where the joint angles were compared to target values. A) One example spatial modification (*Attitude*). Percent errors of peak joint angles relative to target values were computed for each joint separately (left), then averaged across strides and target variables to compute and overall modification error (middle). The middle column shows stride-averaged data from one participant from each of the YA (orange), OA (gray), and MCI (purple) groups. B) One example temporal modification (bottom; *simple duple–quick-quick-Slow-Slow; qqSS*). The target rhythm is shown as musical notation below the temporal timeseries. Temporal targets were defined by step sequences, shown by the q-q-S-S notation. Deviations in the timing of kinematic peaks relative to the prescribed temporal pattern constituted error. The middle column shows a histogram of percent errors, which were averaged across strides (dashed line). The bottom middle plot depicts a Fast Fourier Transform of the timeseries data, in which the dominant frequencies (peaks) are compared to target frequencies (dashed lines). Temporal modification performance was defined as the average of these percent errors. Finally, to compute RMS performance error, all spatial and/or temporal modifications constituting an RMS were averaged (w*ithin-RMS performance*).

Temporal targets for each modification were defined by the modification's pattern of quick (q) and slow

(S) steps and the prescribed tempos (Figure 2). For example, the *simple duple – qqSS* modification

(Figure 2A) involved performing two quick steps, followed by two slow steps at cadences of 100 steps

per minute for the quick steps and 50 steps per minute for the slow steps.

For each RMS, performance was defined as the percent error relative to the spatial and temporal

targets, averaged over all gait cycles, targets, and modifications comprising the RMS (Figure 3). For

spatial modifications, percent errors were computed for executed joint angles relative to the prescribed

joint targets evaluated at key instances of the gait cycle (stance or swing; Figure 3A, yellow dots). The

percent errors were averaged across all strides and target joints for the modification (targets are

specified in Figure 1; multiple strides are shown in Figure 3A, left). For temporal modifications, percent

errors were computed for the temporal sequence of hip flexion angles relative to the prescribed step

sequence (Figure 3B, timeseries peaks deviate from target sequence timing), and the dominant

frequencies of the kinematics relative to target frequencies that would be expected by the pattern of

quick and slow steps (Figure 3B, middle). This approach, rather than using step times, was necessary

because foot contact events could not be reliably detected for all gait modifications using the inertial

sensors. We identified dominant frequencies at the maxima of a Fast Fourier Transform of the joint

kinematics (Figure 3B, middle). If a temporal modification consisted of quick and slow steps, we

identified two dominant frequencies. If half, quick, and slow steps were included, we identified three

dominant frequencies. For example, the *qqSS* simple duple sequence at 100 bpm would have target

dominant frequencies of 100 (q) and 50 (S) bpm, or 1.67 and 0.83 steps/s.

To compute an overall percent error for each RMS, errors computed for each target and gait cycle in a

modification were first computed for each variable of the modification's target variables (e.g., yellow

dots in Figure 3A and temporal error in Figure 3B). The errors within each modification were then

averaged (Figure 3, middle). The RMS performance was then defined as the average percent error

across modifications comprising that RMS (Figure 3, right). For example, RMS performance error on the

spatiotemporal *coupé*-passé – *simple duple qqSS* was defined as the average of percent errors for the

coupé, *passé*, and *simple duple qqSS* modifications. Finally, for RMS classes (spatial, temporal,

spatiotemporal), percent errors for each RMS within the corresponding class were averaged. Lower

spatial or temporal percent error implied better RMS performance.

Music and dance relationships and proficiency assessments

To quantify individuals' relationships to music, including their prior experiences and how they interact

with music in daily life, we developed a Music Relationship Questionnaire (MRQ; Table 2) [36]. The MRQ

was administered via REDCap and consisted of ten introspective questions on a Likert scale with seven

response categories. A composite MRQ score was defined as the average score of the ten Likert scale

questions, producing a maximum possible score of 7, with larger scores reflecting stronger relationships

to music. The MRQ also included an open-ended question about active engagement in music and a

multiple-choice selection of musical genre preferences (see *Supplemental – S3* for more information).

Table 2: List of questions asked for the Music Relationships Questionnaire (MRQ) and Dance
Relationship Questionnaire (DRQ).

Item	MRQ	DRQ	Scoring Scale**
1	Music is important in my life.	Dance is important in my life.	Strongly disagree = 1 Strongly agree = 7
2	I listen to music in my typical day.	I dance in my typical day.*	Strongly disagree = 1 Strongly agree = 7

3	I focus on what I am hearing while listening to music.	While listening to music, I often physically respond (ex: tapping, moving, clapping, nodding, snapping).	Strongly disagree = 1 Strongly agree = 7
4	I focus on other things while music is playing.	I do not often move while music is playing.	Strongly agree = 1 Strongly disagree = 7
5	I actively choose the music I listen to.	I actively choose to move when music is playing.	Strongly disagree = 1 Strongly agree = 7
6	I usually listen to whatever music happens to be playing.	I usually dance to whatever music happens to be playing.	Strongly disagree = 1 Strongly agree = 7
7	I play an instrument/sing.	I dance regularly.	Strongly disagree = 1 Strongly agree = 7
8	I have played an instrument/sung for most of my life.	I have danced for most of my life.	Strongly disagree = 1 Strongly agree = 7
9	I played an instrument/sang only as a child.	I danced only as a child.	Strongly disagree = 1 Strongly agree = 7
10	In a typical day, I play an instrument/sing.	In a typical day, I dance.*	Strongly disagree = 1 Strongly agree = 7

*Two items that were worded similarly and highly correlated (Pearson's r = 0.89) were averaged before computing the composite score for the DRQ.

**The Scoring Scale was identical for the MRQ and DRQ. Note that the Scoring Scale on Item 4 is reversed relative to all other items.

Similarly, to quantify individuals' relationships to dance, including their prior experience and how they

interact with dance in daily life, we developed a Dance Relationship Questionnaire (DRQ; Table 2) [36].

The DRQ was administered via REDCap and consisted of ten introspective questions on a Likert scale

with seven response categories. Like the MRQ, a composite DRQ score was initially defined as the

average score of the 10 Likert scale questions, producing a maximum possible score of 7, with larger

scores reflecting stronger relationships to dance. However, because questions 2 and 10 were similarly

worded and responses were strongly correlated (Pearson's r = 0.89), we took the average of these

questions before computing the composite score across all questions. The DRQ also included an open-

ended question about active engagement in dance and a multiple-choice selection of dance style

preferences (see *Supplemental – S3* for more information).

To quantify individuals' proficiency in perceiving rhythms and executing motor commands in-sync with

those rhythms, we developed an objective rhythm assessment (RA; Table 3) [36]. The RA was conducted

in-person or virtually by trained personnel and took 10-15 minutes to complete. The assessment

consisted of three parts: 1) To assess participants' abilities to perceive, comprehend, and reproduce

rhythmic patterns from *auditory* stimuli, participants listened to four recordings of rhythmic patterns of

quick and slow claps (listened three times per pattern), then attempted to accurately clap each rhythm

twice (Table 3; items 1-4). These tasks are denoted *auditory rhythm reproduction* tasks. 2) To assess

participants' abilities to perceive, comprehend, and replicate rhythmic patterns from *visual* stimuli,

participants read two measures of Western music notation in 4/4 time and attempted to accurately

clapback the rhythm (Table 3; item 5). This task is denoted *visual rhythm reproduction* tasks. Participants

could practice the rhythms for up to one minute before their clapping was scored and were instructed

to attempt to clap the rhythm even if they could not read musical notation. Anecdotally, participants

who verbally indicated they could not read musical notation typically received a score of zero on this

item. The audio of all clapped rhythms was recorded for scoring. 3) To assess participants' abilities to

recognize different meters in music, participants listened to five music passages, then identified whether

that passage consisted of either "two's" (*duple*), "three's" (*triple*), or "other" meters (Table 3; items 6-

10). These tasks are denoted *auditory meter recognition* tasks. All passages were less than 60 seconds in

length and played up to three times for the participant. The RA composite score was defined as the sum

of scores on the 10 items, with a maximum score of 10 points (Table 3). Correct answers received one

point, partially correct answers (e.g., correct on one of two clapbacks on the *auditory* or *visual rhythm*

production tasks) received one-half point, and incorrect answered received zero points. RA instructions

are provided in *Supplemental – S2*.

365 **Table 3**: Description of assessment items in the Rhythm Assessment (RA).

Item	Item Description	Scoring Scale – Response (Points)		
1	Auditory rhythm reproduction A			
2	Auditory rhythm reproduction B			
3	Auditory rhythm reproduction C	Correct (1)	Partial (0.5)	Incorrect (0)
4	Auditory rhythm reproduction D			
5	Visual rhythm reproduction A			
6	Auditory meter recognition A (Duple)	Duple (1)	Waltz (0)	Other (0)
7	Auditory meter recognition B (Waltz)	Duple (0)	Waltz (1)	Other (0)
8	Auditory meter recognition C (Duple)	Duple (1)	Waltz (0)	Other (0)
9	Auditory meter recognition D (Waltz)	Duple (0)	Waltz (1)	Other (0)
10	Auditory meter recognition E (Other)	Duple (0)	Waltz (0)	Other (1)

"Partially correct" indicated that the participant clapped the rhythm correctly once and incorrectly once.
Auditory items involved listening to a rhythm, then clapping it back.
The *Visual* item involved reading a sequence of 10 musical notes (eighth, quarter, half beats) then clapping the
notes.
Meter recognition items involved listening to a rhythm, then identifying whether the meter was "*twos" (two-*
count), "*threes" (three-count),* or "*other.*"

The RA documentation and audio recordings, along with the data used in this manuscript and a data

dictionary, are freely available in the supplemental materials (see *Supplemental – S1* for a description of

these files).

Statistical analysis

To determine if groups differed in their experiences with dance and music, as well as their rhythmic

proficiency, we tested for differences in MRQ, DRQ, and RA scores between all pairs of groups (YA, OA,

and MCI) using independent-sample t-tests ($\alpha = 0.05$). Because our sample was larger than that in [8]

and [36], we replicated the study's group-wise comparisons of spatial, temporal, and spatiotemporal

RMS performance using independent-sample t-tests ($\alpha = 0.05$). Because we were previously interested

in how age and cognitive status impacted RMS performance, we only compared RMS performance

between YA and OA (age effect) and between OA and MCI (cognitive status effect) [8].

To characterize relationships between the MRQ, DRQ, and RA, we performed univariate linear

regression of the composite scores between pairs of the MRQ, DRQ, and RA. Similarly, to determine if

the MRQ, DRQ, and RA were associated with measures of cognitive and motor-cognitive function, we

performed univariate linear regression between these composite scores and scores on the MoCA, Trail

Making Task (B-A), Reverse Corsi Blocks test, and the Body Position Spatial Task. Regressions were

performed separately for each assessment pair and separately for the YA, OA, and MCI groups.

To determine if, within each participant group, stronger relationships to music (higher MRQ composite

scores) were associated with better temporal modulation ability (lower RMS performance error), we

performed linear regression between MRQ composite scores and temporal RMS performance separately

for each group. Analyzing groups separately was based on our prior finding that RMS performance

differed between groups [8]. Similarly, to determine if, within groups, stronger relationships to music

were associated with better spatiotemporal modulation ability, we performed linear regression between

MRQ composite scores and spatiotemporal RMS performance.

To determine if, within groups, stronger relationships to dance (higher DRQ composite scores) were

associated with better spatial modulation ability, we performed linear regression between DRQ

composite scores and spatial RMS performance. Similarly, to determine if, within groups, stronger

relationships to dance were associated with better spatiotemporal RMS performance, we performed

linear regression between DRQ scores and spatiotemporal RMS performance.

To determine if stronger rhythmic proficiency (higher RA composite scores) was associated with better

temporal and spatiotemporal modulation ability, we performed linear regression between RA composite

scores and temporal and spatiotemporal RMS performance, respectively, for each group separately.

Finally, because we identified associations between RA composite scores and RMS performance, we

sought to determine if the ability to reproduce auditory rhythms and identify meter in musical pieces

differentially impacted RMS performance. We conducted an exploratory regression analysis of the

auditory rhythm reproduction and *auditory meter recognition* sub-components of the RA against

temporal and spatiotemporal RMS separately for each group.

For all analyses, we report regression accuracy as coefficients of determination (r^2), regression slopes,

and significance levels according to Wald tests ($\alpha = 0.05$). Group-level comparisons of demographics and

clinical characteristics were performed using the software package *R*. Regression analyses were

performed using MATLAB 2021b (Mathworks, Natick, USA).

Results:

Clinical assessments suggest worse cognitive function in adults with MCI, compared to older adults

Clinical assessments revealed group-level differences in motor-cognitive and cognitive function (Table

1). Compared to the OA group, the YA group exhibited better performance on assessments of working

memory (Reverse Corsi Blocks; $p < 0.001$) and motor-cognitive integration (Body Position Spatial Task; p

< 0.001). Similarly, the MCI group exhibited worse performance than the OA group on assessments of

global cognitive function (MoCA; $p = 0.005$), working memory (Reverse Corsi Blocks; $p = 0.035$), and set-

shifting (Trail Making Test; $p = 0.050$).

Music relationships, rhythmic proficiency, and RMS performance differed between groups

Relationships to music and rhythmic proficiency differed between groups (Figure 4A). The YA and OA

groups exhibited stronger relationships to music and greater rhythmic proficiency than the MCI group.

On the MRQ, the YA group scored an average of 1.0 points (out of 7) higher (p = 0.001) and the OA

group scored 0.7 points higher (p = 0.006) than the MCI group. The YA and OA groups exhibited similar

relationships to music, as indicated by similar MRQ scores (p = 0.228). On the RA, the YA group scored

an average of 3.2 points (out of 10) higher (p < 0.001) and the OA group scored 2.4 points higher (p <

0.001) compared to the MCI group. YA and OA groups exhibited similar levels of rhythmic proficiency, as

indicated by similar RA scores (p = 0.223). No groups exhibited differences in the strength of their dance

relationships, as indicated by similar DRQ scores (p > 0.575).

Figure 4: Boxplots representing distributions of Music Relationship Questionnaire (MRQ), Dance Relationship Questionnaire (DRQ), and Rhythm Assessment (RA) scores. A) MRQ, DRQ, and RA scores for the three participant groups (YA, OA, MCI). Higher MRQ and DRQ scores (max = 7) reflect stronger music and dance relationships, respectively. Higher RA scores (max = 10) reflect greater rhythmic proficiency. B) RMS performance error on each of spatial, temporal, and spatiotemporal RMS for the three participant groups. For both plots, dots represent individual participants. Higher composite scores indicate better performance on the MRQ, DRQ, and RA (upper green arrow). Lower RMS error indicates better performance on RMS (lower green

arrow). Each spatial and temporal RMS, error was averaged across 9 RMS from their respective domains and spatiotemporal error was averaged across 4 RMS. For all boxplots, p-values denote significant differences according to independent-samples t-tests ($\alpha = 0.05$).

Compared to OA, the YA group exhibited better RMS performance, as indicated by lower percent error

on spatial ($p < 0.001$), temporal ($p = 0.025$), and spatiotemporal ($p < 0.001$) RMS (Figure 4B). The OA

group did not perform RMS with different accuracy than the MCI group (all $p > 0.257$).

MRQ, DRQ, and RA composite scores were not strongly correlated across groups

Greater MRQ composite scores were associated with greater DRQ and RA composite scores only in the

OA group ($r^2 < 0.27$; $p < 0.024$; Table 4). Conversely, greater DRQ composite scores were associated with

greater RA scores only in the YA group ($r^2 < 0.43$; $p < 0.021$).

Table 4: Univariate linear regression results testing for covariation between MRQ, DRQ, RA composite
scores.

Questionnaire	Questionnaire / assessment	YA			OA			MCI		
		r^2	slope	p	r^2	slope	p	r^2	slope	p
MRQ	DRQ	0.28	0.8	0.080	0.20	0.8	**0.024**	0.04	0.4	0.399
	RA	0.26	0.3	0.087	0.27	0.2	**0.010**	0.22	0.2	0.064
DRQ	RA	0.43	0.5	**0.021**	0.00	0.0	0.848	0.17	0.2	0.109

Abbreviations: YA, younger adults; OA, older adults without mild cognitive impairment; MCI, older adults with mild
cognitive impairment; MRQ, Music Relationship Questionnaire; DRQ, Dance Relationship Questionnaire; RA,
Rhythm Assessment

For each group, regression results are shown using the coefficient of determination (r^2), regression slope, and p-
value (p) denoting a slope that is significantly different from zero ($\alpha = 0.05$). Bold p-values denote $p < 0.05$.

Clinical measures of cognitive and motor-cognitive function were not associated with the MRQ, DRQ, or

RA scores

Across four clinical measures of cognitive and motor-cognitive function, we identified only one

significant association between the Body Position Spatial Task and MRQ in the OA group ($r^2 = 0.37$; slope

= 0.06; p = 0.001; Table 5). Scores on the MoCA, Trail Making Task (B-A), and Reverse Corsi Blocks test

were not associated with the MRQ, DRQ, or RA in any group.

Table 5: Univariate linear regression results between clinical assessments of cognitive and motor-
cognitive function and the MRQ, DRQ, and RA composite scores.

Questionnaire / assessment	Assessment	YA	OA	MCI
MRQ	MoCA	0.00	0.00	0.12
	Trails B-A	0.02	0.00	0.02
	Reverse Corsi blocks	0.06	0.05	0.04
	Body position spatial task	0.00	**0.37**	0.06
DRQ	MoCA	0.01	0.01	0.00
	Trails B-A	0.00	0.00	0.09
	Reverse Corsi blocks	0.00	0.00	0.01
	Body position spatial task	0.14	0.01	0.22
RA	MoCA	0.01	0.01	0.00
	Trails B-A	0.04	0.01	0.04
	Reverse Corsi blocks	0.13	0.03	0.19
	Body position spatial task	0.14	0.11	0.03

Abbreviations: YA, younger adults; OA, older adults without mild cognitive impairment; MCI, older adults with mild
cognitive impairment; RMS, Rhythmic Movement Sequence; MRQ, Music Relationship Questionnaire; DRQ, Dance
Relationship Questionnaire; RA, Rhythm Assessment; MoCA, Montreal Cognitive Assessment; Trails B-A, Trail
Making Task items B-A.
For each group, regression results are shown using the coefficient of determination (r^2), boldfaced values denote a
correlation with slope that was significantly different from zero (α = 0.05) without correction for multiple
comparisons.

Higher MRQ scores were associated with reduced temporal RMS errors in young and older adults without

MCI

In participants without MCI, individuals with higher MRQ scores (stronger music relationships) exhibited

lower error on temporal RMS (i.e., better temporal RMS performance). MRQ scores were associated

with temporal RMS errors in the YA (r^2 = 0.41; slope = -3.2; p = 0.026; Table 6; orange in Figure 5A, top)

and OA groups (r^2 = 0.18; slope = -3.5; p = 0.030; grey in Figure 5A, top), explaining a small-to-moderate

amount of the variance in temporal RMS errors. Conversely, the MCI group did not exhibit significant

associations between MRQ scores and temporal RMS performance (r^2 = 0.06; p = 0.314; purple in Figure

5A, top). No group exhibited significant associations between MRQ scores and spatiotemporal RMS

performance (all r^2 < 0.15; p > 0.114; Figure 5A, bottom).

Table 6: Univariate linear regression results between MRQ, DRQ, and RA composite scores and RMS
performance.

Questionnaire / assessment	RMS class	YA			OA			MCI		
		r^2	slope	p	r^2	slope	p	r^2	slope	p
MRQ	Temporal	0.41	-3.21	**0.026**	0.18	-3.49	**0.030**	0.06	-2.62	0.317
	Spatiotemporal	0.15	-1.63	0.215	0.10	-1.23	0.114	0.02	-0.62	0.551
DRQ	Spatial	0.03	-0.60	0.610	0.01	-0.30	0.575	0.01	-0.29	0.732
	Spatiotemporal	0.03	-0.47	0.612	0.00	-0.03	0.942	0.00	-0.12	0.841
RA	Temporal	0.40	-1.67	**0.026**	0.17	-1.65	**0.045**	0.08	-0.91	0.299
	Spatiotemporal	0.05	-0.48	0.498	0.18	-0.75	**0.043**	0.00	0.00	0.994
RA rhythm reproduction	Temporal	0.21	-2.6	0.135	0.05	-2.0	0.296	0.01	-0.5	0.722
	Spatiotemporal	0.02	-0.7	0.667	0.00	0.2	0.791	0.02	0.3	0.623
RA meter recognition	Temporal	0.26	-2.6	0.090	0.18	-2.7	**0.037**	0.09	-1.9	0.265
	Spatiotemporal	0.01	-0.4	0.743	0.34	-1.5	**0.003**	0.03	-0.4	0.534

492   Abbreviations: YA, younger adults; OA, older adults without mild cognitive impairment; MCI, older adults with mild
493   cognitive impairment; RMS, Rhythmic Movement Sequence; MRQ, Music Relationship Questionnaire; DRQ, Dance
494   Relationship Questionnaire; RA, Rhythm Assessment
495   For each group, regression results are shown using the coefficient of determination (r^2), regression slope, and p-
496   value (p) denoting a slope that is significantly different from zero (α = 0.05). Bold p-values denote p < 0.05.
497

Figure 5: Linear regression testing for within-group associations between RMS performance and each of the MRQ, DRQ, and RA. Each dot represents a single participant. Lines indicate within-group linear fits. Colors denote the groups (YA: orange, OA: gray, MCI: purple). Regression R-squared values, slopes, and p-values are shown on the corresponding plots for fits that were significantly different from zero (Wald tests; α = 0.05), with colors corresponding to groups. Each of the spatial and temporal RMS percent errors were averaged across the 9 respective RMS and spatiotemporal errors were averaged across 4 RMS. Lower RMS error indicates better performance (green arrow), on average, across RMS within the corresponding RMS class. A) MRQ vs. temporal (top) and spatiotemporal (bottom) RMS performance errors. Higher MRQ scores represent stronger music relationships. B) DRQ vs. spatial (top) and spatiotemporal (bottom) RMS performance errors. Higher DRQ scores represent stronger dance relationships. C) Comparisons of RA with percent errors on temporal (top) and spatiotemporal (bottom) RMS. Higher RA scores imply greater rhythmic proficiency. D) Comparisons of RA subcomponents—*auditory rhythm reproduction* (max score = 4) and *auditory meter recognition* (max score = 5)—to temporal (top) and spatiotemporal (bottom) RMS.

DRQ scores were not associated with spatial or spatiotemporal RMS performance

No group exhibited significant associations between DRQ scores and spatial or spatiotemporal RMS

performance (all $r^2 < 0.03$; $p > 0.575$; Table 6; Figure 5B).

Higher RA scores were associated with reduced temporal RMS errors in young and older adults without

MCI

In participants without MCI, individuals with higher RA scores tended to exhibit better temporal RMS

performance (i.e., lower error). Negative associations between RA scores and temporal RMS error were

observed in the YA (r^2 = 0.40; slope = -1.7; p = 0.026; Table 6; orange in Figure 5C, top) and OA groups (r^2

= 0.22; slope = -1.7; p = 0.021; grey in Figure 5C, top), explaining a small-to-moderate amount of

variance in temporal RMS errors. Conversely, the MCI group did not exhibit significant associations

between RA scores and temporal RMS performance, in part due to heterogeneity in RMS performance

across individuals with low RA scores (r^2 = 0.11; p = 0.210; purple in Figure 5C, top). Only the OA group

exhibited an association between better RA scores and better spatiotemporal RMS performance (all r^2 =

0.18; slope = -0.75; p > 0.043; Figure 5C, bottom).

Auditory rhythm reproduction and auditory meter recognition scores were only associated with RMS

performance in older adults

When analyzing the RA sub-components, *auditory rhythm reproduction* (4 items in the RA) alone was

not associated with temporal or spatiotemporal RMS performance in any group (r^2 < 0.21; p > 0.135;

Table 6; Figure 5D). For the second RA sub-component, better *auditory meter recognition* (5 items in the

RA), was associated with better temporal (r^2 > 0.18; slope = -2.7; p = 0.037) and spatiotemporal (r^2 >

0.34; slope = -1.5; p = 0.003) RMS only in the OA group (Figure 5D).

Discussion

This study shows that in young and older adults without MCI, stronger relationships to music and better

rhythmic proficiency are associated with a better ability to accurately modulate temporal aspects of

movement during RMS. Conversely, these relationships are not apparent in adults with MCI, suggesting

that cognitive deficits associated with MCI hinder the ability to transform relationships to music or

rhythmic proficiency into an ability to modulate temporal aspects of movement during RMS. Our central

hypothesis was, therefore, only supported for temporal RMS in younger and older adults without MCI.

Stronger relationships to dance do not appear to impact the ability to perform RMS involving spatial

modifications to movement in any group. These findings suggest that only in the absence of MCI-related

cognitive deficits will individuals' relationships to music or rhythmic proficiency be indicative of the

complexity of musical rhythms and step sequences that they can successfully perform during dance-

based therapies [3, 5, 6].

Stronger relationships to music and rhythmic proficiency indicate a better ability to accurately modulate

temporal features of movement during RMS in adults without MCI, but not with MCI. In adults without

MCI, these associations suggest that constructs that are developed by prior relationships to music and

support rhythmic proficiency also improve the ability to perceive rhythm and meter and entrain

movement to these rhythms. For example, individuals with stronger music relationships may exhibit

more precise mapping of temporal stimuli between auditory processing and motor planning regions in

the brain [14, 32]. Such mapping may enable more accurate entrainment of step timing to music-based

auditory cues during temporal RMS [37]. Given the likelihood of similar perception-action pathways

invoked by the RA and temporal and spatiotemporal RMS [31, 32], it is reasonable to expect that within-

group differences in RA and RMS performance are driven by similar constructs, which merits future

investigation. For example, the ability to identify meter in musical pieces appears to be moderately

predictive of temporal RMS performance in the absence of MCI. Our findings suggest that this skill may

be developed through prior experiences with either music or dance.

Cognitive deficits in adults with MCI may have masked associations between music relationships or

rhythmic proficiency and temporal RMS performance. Individual-specific working memory deficits in the

MCI group may explain the more variable temporal RMS performance and may be exacerbated for

longer temporal sequences. For example, we previously found that adults with MCI performed longer

temporal RMS sequences (6-step duple sequences) less accurately than older adults and that lower

temporal RMS performance was associated with worse working memory (i.e., lower scores on the

Reverse Corsi Blocks test) [8, 24]. Note that the group differences in RMS performance presented in the

present study are consistent with those previously reported using a subset of participants included here

[8]. Similarly, lower rhythmic proficiency in adults with MCI, compared to those without MCI, suggests

that attentional or motor-cognitive deficits accompanying MCI diagnosis may manifest as a reduced

ability to perceive musical cues and entrain motor commands to music [14]. Despite adults with MCI

reporting weaker relationships to music than older adults without MCI, our findings suggest individual

differences in music and dance relationships are not driven by declines in cognitive or motor-cognitive

function. Further research is required to determine the extent to which MCI-related impairments

contribute to declines in rhythmic proficiency, which may impact individuals' abilities to synchronize

movements to auditory cues during dance-based therapies [12-14].

Unlike music relationships, stronger relationships to dance are not indicative of a better ability to

accurately perform spatial RMS. Rather, our prior study found that differences in spatial RMS

performance are better explained by differences in age-related motor and cognitive function [8].

Therefore, stronger dance relationships do not likely enhance constructs that are beneficial to spatial

RMS performance consistently across individuals. For example, the experiences assessed by the DRQ

neither require nor develop dynamic balance or the large joint ranges of motion required by spatial RMS

[38]. Prior experience in non-dance activities like gymnastics may, therefore, be a better indicator of an

individual's ability to perform spatial RMS [39, 40]. Identifying specific subcomponents of dance

relationships or other dance-related experiences that predict the ability to modulate spatial aspects of

movement is an interesting future research direction. For example, only three participants reported a

preference for ballet, which was the basis for the spatial modifications during RMS (*Supplemental – S3*).

Adjusting the modifications comprising RMS to be derived from dance styles that more-strongly align

with participants' preferences may yield stronger relationships between DRQ scores and spatial RMS

performance.

Better rhythmic proficiency, but not relationships to music or dance, may be indicative of better

spatiotemporal RMS performance. However, we observed this trend only in older adults without MCI. As

discussed in the prior paragraph, constructs not evaluated by the MRQ, DRQ, or RA likely impact spatial

RMS performance, which constitutes half of the spatiotemporal RMS error. The relationship between

better rhythmic proficiency and temporal RMS performance likely also drives better spatiotemporal

RMS performance. The lack of association in young adults is not necessarily surprising; participants may

choose to prioritize spatial, rather than temporal aspects of movement, potentially decoupling rhythmic

proficiency from spatiotemporal RMS performance. However, our prior findings suggest that differences

in motor function (e.g., balance) or cognitive function (e.g., set shifting), appear to be stronger

predictors of the ability to perform spatiotemporal RMS [8, 23, 24, 41].

Several factors limit the generalizability and interpretation of our results. Limitations related to our

experimental protocol and RMS performance quantification are discussed in [8]. While the OA and MCI

group sample sizes in this study were larger than in [8] and [36], still larger sample sizes would improve

confidence in our regression results for groups with highly variable RMS performance. Further, our

analyses did not adjust for within-group effects of motor or cognitive function, or their interactions with

music or dance relationships, on RMS performance [8]. Complex interactions between motor and

cognitive decline, music or dance relationships, and rhythmic proficiency in individuals with MCI, may

contribute to RMS performance. Future larger-sample studies could use multivariate regression

techniques to extend our understanding of how individual differences in motor and cognitive function

moderate the impacts of music and dance relationships or rhythmic proficiency on RMS performance.

Kinematic estimates from the APDM Mobility Lab software may also induce errors, particularly at large

joint excursions for which the software and inertial sensors have not been validated [34, 35, 42, 43].

Prior studies of large joint ranges of motion suggest that IMU-based kinematic errors can reach 5-8

degrees, which may introduce non-trivial errors in RMS performance [44, 45]. However, these errors

would likely reduce our ability to detect inter-individual differences in RMS performance: extreme joint

angles would likely increase RMS error for the best performers. Further, experiment administrators

reported, anecdotally, that participants with worse RMS percent errors struggled more to perform RMS.

Additionally, the MRQ, DRQ, and RA represent a preliminary characterization of music relationships,

dance relationships, and rhythmic proficiency, respectively. The questionnaire/assessment items

encompassed multiple constructs that may impact RMS performance. Computing MRQ, DRQ, and RA

composite scores by averaging the scores for all items may mask more nuanced associations between

music and dance relationships and RMS performance. Further, moderate covariation between the MRQ,

DRQ, and RA scores suggests that these assessments encode some overlapping constructs. Analyzing

individual questionnaire items may reveal subcomponents of music and dance relationships that are

more strongly associated with RMS performance. Further, established questionnaires, such as the

Goldsmith's Music and Dance Sophistication Indices and the Barcelona Music Reward Questionnaire test

similar perceptual and motor constructs (e.g., active listening, singing ability, sensory-motor) as those

assayed in the MRQ, DRQ, and RA [46-48]. These previously established questionnaires include a

broader range of questions, from proficiency to emotional engagement. The MRQ and DRQ were

developed to assess broad relationships with music and dance, and did not include more nuanced

emotional or social concepts. Unlike the MRQ, the DRQ assessed active, but not passive aspects of

dance relationships. As dance observation is known to impact neural activity and dance performance,

recording information about dance relationships will be valuable for future studies [27, 47, 49]. The

inclusion of the additional constructs described above could provide a more detailed understanding of

how prior experience with music or dance impacts RMS performance. Including a broader set of

questions from the MRQ, DRQ, and RA, as well as established questionnaires, may be useful to identify

specific aspects of music and dance relationships that impact the ability to perform dance-like

movements.

Further, we did not control for individual preferences for different musical genres or dance styles, active

(e.g., singing, playing an instrument, dancing) versus passive (e.g., listening to music, watching dance)

engagement in music or dance, and extent of experience, all of which may affect the relationships

between music or dance relationships and RMS performance. All participants in the present study

reported listening to one or more Western musical genres. However, non-Western music listeners may

struggle to identify rhythm and meter in Western musical pieces, possibly reducing RMS performance

[14]. Most participants in the present study reported listening to classical music, and classical music

listeners tended to perform better on temporal RMS, which used classical music pieces (see

Supplemental – S3 for more information). However, this trend may not be specific to classical music.

Similarly, we did not find that participants who reported active music or dance engagement performed

better on RMS than those who did not (see *Supplemental – S3* for more information). Finally, we did not

directly measure the extent (e.g., number of years) of experience in music or dance, which may impact

RMS performance. Characterizing how group or individual differences in RMS performance generalize

across preference, engagement style, and extent of experience in music and dance would improve our

understanding of the clinical utility of RMS.

Finally, future studies should characterize the extent to which RMS performance is associated with

participants' capabilities within the many other dance styles and musical genres encountered during

dance-based therapies, as well as the participant-reported difficulty, efficacy, and engagement in such

therapies. Determining the extent to which RMS, which incorporate ballet fundamentals with classical

musical selections, generalize to other music or dance styles would support the utility of RMS as a tool

to quantify and predict therapeutic performance. Spatial and temporal modification outcomes may

need to be refined when applying RMS to other dance styles, such as partner-based dances or styles

that place less emphasis on kinematic configuration [50, 51]. Quantifying how RMS maps to participant-

reported difficulty and engagement could enable the selection of therapy parameters to optimize the

level of challenge during therapy [3, 6, 52]. Future work should relate RMS performance to therapy

efficacy, participant-reported difficulty and engagement, and therapy impacts of motor and/or cognitive

function [6]. Ultimately, to establish RMS as a clinical tool to help clinicians optimize dance-based

therapies, we envision a unifying multivariate model capable of predicting RMS performance across

participants of different ages, cognitive function levels, prior relationships to music or dance, and

rhythmic proficiency.

Conclusions

We investigated the relationships between rhythmic movement sequence (RMS) performance and novel

assessments of individuals' relationships to music and dance, as well as rhythmic proficiency. The

associations of only music relationships and rhythmic proficiency with temporal RMS in young and older

adults without, but not with, MCI suggest that cognitive deficits in adults with MCI likely hinder the

ability of music relationships or rhythmic proficiency to improve performance on dance-like RMS. These

findings contribute to a growing understanding of the factors influencing the ability to accurately

perform dance-like movements, which may inform the personalization of dance-based therapies.

Author contributions

Alexandra Slusarenko (Conceptualization; Formal Analysis; Investigation; Visualization; Writing – original draft; Writing – review & editing)

Michael C. Rosenberg (Conceptualization; Data Curation; Funding Acquisition; Investigation; Methodology; Project administration; Software; Validation; Visualization; Writing – original draft; Writing – review & editing)

Meghan E. Kazanski (Conceptualization; Investigation; Methodology; Project administration; Validation; Writing – review & editing)

J. Lucas McKay (Conceptualization; Data Curation; Funding Acquisition; Methodology; Software; Supervision; Writing – review & editing)

Laura Emmery (Conceptualization; Funding Acquisition; Methodology; Supervision; Writing – review & editing)

Trisha M. Kesar (Conceptualization; Funding Acquisition; Methodology; Supervision; Writing – review & editing)

Madeleine E. Hackney (Conceptualization; Data Curation; Funding Acquisition; Methodology; Project administration; Resources; Supervision; Writing – review & editing)

Acknowledgments

We thank K. Cao, T. Prusin, and C. Carroll-Sauer for their assistance in data collection and participant recruitment.

Funding

Research reported in this manuscript was supported by the National Institute of Child Health and Human Development and the National Institute on Aging of the National Institutes of Health under award numbers F32HD108927 and R01AG062691, respectively. This research was supported by Emory University through a Goizueta Alzheimer's Disease Research Center CEP Innovation Accelerator Seed Grant and an award from the Office of the Emory University Senior Vice President of Research Intersection Fund.

Conflict of Interest

Madeleine E. Hackney is an Editorial Board Member of this journal, but was not involved in the peer-review process nor had access to any information regarding its peer-review.

Data availability

The data supporting the findings of this study are openly available as Supplemental Material. These data were derived from the following resources available in the public domain:

https://www.frontiersin.org/articles/10.3389/fnhum.2023.1040930/full.

References

[1]   Zhu Y, Zhong Q, Ji J, Ma J, Wu H, Gao Y, Ali N, Wang T (2020) Effects of aerobic dance on cognition in older adults with mild cognitive impairment: a systematic review and meta-analysis. *J Alzheimers Dis* **74**, 679-690.

[2]   Gauthier S, Reisberg B, Zaudig M, Petersen RC, Ritchie K, Broich K, Belleville S, Brodaty H, Bennett D, Chertkow H, Cummings JL, de Leon M, Feldman H, Ganguli M, Hampel H, Scheltens P, Tierney MC, Whitehouse P, Winblad B (2006) Mild cognitive impairment. *The Lancet* **367**, 1262-1270.

[3]   Lazarou I, Parastatidis T, Tsolaki A, Gkioka M, Karakostas A, Douka S, Tsolaki M (2017) International ballroom dancing against neurodegeneration: a randomized controlled trial in Greek community-dwelling elders with mild cognitive impairment. *Am J Alzheimers Dis Other Demen* **32**, 489-499.

[4]   Zhu Y, Wu H, Qi M, Wang S, Zhang Q, Zhou L, Wang S, Wang W, Wu T, Xiao M (2018) Effects of a specially designed aerobic dance routine on mild cognitive impairment. *Clin Interv Aging* **13**, 1691.

[5]   McKee KE, Hackney ME (2013) The effects of adapted tango on spatial cognition and disease severity in Parkinson's disease. *J Mot Behav* **45**, 519-529.

[6]   Guadagnoli MA, Lee TD (2004) Challenge point: a framework for conceptualizing the effects of various practice conditions in motor learning. *J Motor Behav* **36**, 212-224.

[7]   Hackney ME, Earhart GM (2009) Effects of dance on movement control in Parkinson's disease: a comparison of Argentine tango and American ballroom. *J Rehabil Med* **41**, 475-481.

[8]   Rosenberg MC, Slusarenko A, Cao K, Lucas McKay J, Emmery L, Kesar TM, Hackney ME (2023) Motor and cognitive deficits limit the ability to flexibly modulate spatiotemporal gait features in older adults with mild cognitive impairment. *Front Hum Neurosci* **17**.

[9]   Rallis I, Doulamis N, Doulamis A, Voulodimos A, Vescoukis V (2018) Spatio-temporal summarization of dance choreographies. *Computers & Graphics* **73**, 88-101.

[10]   Hackney ME, Kantorovich S, Levin R, Earhart GM (2007) Effects of tango on functional mobility in Parkinson's disease: a preliminary study. *J Neurol Phys Ther* **31**, 173-179.

[11]   Wilson M, Kwon Y-H (2008) The role of biomechanics in understanding dance movement: a review. *J Dance Med Sci* **12**, 109-116.

[12]   Thaut MH, Abiru M (2010) Rhythmic auditory stimulation in rehabilitation of movement disorders: a review of current research. *Music Percept* **27**, 263-269.

[13]   Thaut M, Kenyon G, Schauer M, McIntosh G (1999) The connection between rhythmicity and brain function. *IEEE Eng Med Biol Mag* **18**, 101-108.

[14]   Emmery L, Hackney ME, Kesar T, McKay JL, Rosenberg MC (2023) An integrated review of music cognition and rhythmic stimuli in sensorimotor neurocognition and neurorehabilitation. *Ann N Y Acad Sci* **1530**, 74-86.

[15]   Zatorre RJ, Chen JL, Penhune VB (2007) When the brain plays music: auditory–motor interactions in music perception and production. *Nat Rev Neurosci* **8**, 547-558.

[16]   Akombo D (2016) *The unity of music and dance in world cultures*, McFarland & Company, Inc., Jefferson, North Carolina.

[17]   Hackney ME, Earhart GM (2010) Recommendations for implementing tango classes for persons with Parkinson disease. *Am J Dance Ther* **32**, 41-52.

[18]   Chen JL, Penhune VB, Zatorre RJ (2008) Moving on time: brain network for auditory-motor synchronization is modulated by rhythm complexity and musical training. *J Cogn Neurosci* **20**, 226-239.

757   [19]   Lerdahl F, Jackendoff R (1983) An overview of hierarchical structure in music. *Music Percept*,
758   229-252.
759   [20]   Chen JL, Penhune VB, Zatorre RJ (2008) Listening to musical rhythms recruits motor regions of
760   the brain. *Cereb Cortex* **18**, 2844-2854.
761   [21]   London J (2012) *Hearing in time: Psychological aspects of musical meter*, Oxford University
762   Press.
763   [22]   Mueller SG, Weiner MW, Thal LJ, Petersen RC, Jack C, Jagust W, Trojanowski JQ, Toga AW,
764   Beckett L (2005) The Alzheimer's disease neuroimaging initiative. *Neuroimag Clin* **15**, 869-877.
765   [23]   Bowie CR, Harvey PD (2006) Administration and interpretation of the Trail Making Test. *Nat*
766   *Protoc* **1**, 2277-2281.
767   [24]   Vandierendonck A, Kemps E, Fastame MC, Szmalec A (2004) Working memory components of
768   the Corsi blocks task. *Br J Psychol* **95**, 57-79.
769   [25]   Hackney ME, Hall CD, Echt KV, Wolf SL (2013) Dancing for balance: feasibility and efficacy in
770   oldest-old adults with visual impairment. *Nurs Res* **62**, 138-143.
771   [26]   Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL,
772   Chertkow H (2005) The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild
773   cognitive impairment. *J Am Geriatr Soc* **53**, 695-699.
774   [27]   Bläsing B, Schack T (2012) Mental representation of spatial movement parameters in dance.
775   *Spat Cogn Comput* **12**, 111-132.
776   [28]   Ivanenko YP, Cappellini G, Dominici N, Poppele RE, Lacquaniti F (2005) Coordination of
777   locomotion with voluntary movements in humans. *J Neurosci* **25**, 7238-7253.
778   [29]   Styns F, van Noorden L, Moelants D, Leman M (2007) Walking on music. *Hum Movment Sci* **26**,
779   769-785.
780   [30]   Will U, Berg E (2007) Brain wave synchronization and entrainment to periodic acoustic stimuli.
781   *Neurosci Lett* **424**, 55-60.
782   [31]   Van Noorden L, Moelants D (1999) Resonance in the perception of musical pulse. *J New Music*
783   *Res* **28**, 43-66.
784   [32]   Drake C, Jones MR, Baruch C (2000) The development of rhythmic attending in auditory
785   sequences: attunement, referent period, focal attending. *Cognition* **77**, 251-288.
786   [33]   Lakatos P, Karmos G, Mehta AD, Ulbert I, Schroeder CE (2008) Entrainment of neuronal
787   oscillations as a mechanism of attentional selection. *Science* **320**, 110-113.
788   [34]   Mancini M, Horak FB (2016) Potential of APDM mobility lab for the monitoring of the
789   progression of Parkinson's disease. *Expert Rev Med Devices* **13**, 455-462.
790   [35]   Washabaugh EP, Kalyanaraman T, Adamczyk PG, Claflin ES, Krishnan C (2017) Validity and
791   repeatability of inertial measurement units for measuring gait parameters. *Gait Posture* **55**, 87-
792   93.
793   [36]   Slusarenko A (2022) in *Neuroscience and Behavioral Biology* Emory University.
794   [37]   Bengtsson SL, Ullen F, Ehrsson HH, Hashimoto T, Kito T, Naito E, Forssberg H, Sadato N (2009)
795   Listening to rhythms activates motor and premotor cortices. *Cortex* **45**, 62-71.
796   [38]   Hackney ME, Nocera J, Creel T, Riebesell MD, Kesar T (2017) Exercise and balance in older adults
797   with movement disorders In *Locomotion and Posture in Older Adults* Springer, pp. 323-346.
798   [39]   Hrysomallis C (2011) Balance ability and athletic performance. *Sports Med* **41**, 221-232.
799   [40]   Busquets A, Ferrer-Uris B, Angulo-Barroso R, Federolf P (2021) Gymnastics experience enhances
800   the development of bipedal-stance multi-segmental coordination and control during
801   proprioceptive reweighting. *Front Psychol* **12**, 661312.
802   [41]   Reimann H, Ramadan R, Fettrow T, Hafer JF, Geyer H, Jeka JJ (2020) Interactions between
803   different age-related factors affecting balance control in walking. *Front Spor Act Living* **2**, 94.

[42] Mancini M, King L, Salarian A, Holmstrom L, McNames J, Horak FB (2011) Mobility lab to assess
balance and gait with synchronized body-worn sensors. *J Bioeng Biomed Sci*, 007.
[43] Spain R, George RS, Salarian A, Mancini M, Wagner J, Horak F, Bourdette D (2012) Body-worn
motion sensors detect balance and gait deficits in people with multiple sclerosis who have
normal walking speed. *Gait Posture* **35**, 573-578.
[44] Morrow MM, Lowndes B, Fortune E, Kaufman KR, Hallbeck MS (2017) Validation of inertial
measurement units for upper body kinematics. *J Appl Biomech* **33**, 227-232.
[45] Tulipani L, Boocock MG, Lomond KV, El-Gohary M, Reid DA, Henry SM (2018) Validation of an
inertial sensor system for physical therapists to quantify movement coordination during
functional tasks. *J Appl Biomech* **34**, 23-30.
[46] Müllensiefen D, Gingras B, Musil J, Stewart L (2014) The musicality of non-musicians: An index
for assessing musical sophistication in the general population. *PLoS One* **9**, e89642.
[47] Rose D, Müllensiefen D, Lovatt P, Orgs G (2022) The Goldsmiths Dance Sophistication Index
(Gold-DSI): A psychometric tool to assess individual differences in dance experience. *Psychol
Aesthet Crea* **16**, 733-745.
[48] Mas-Herrero E, Marco-Pallares J, Lorenzo-Seva U, Zatorre RJ, Rodriguez-Fornells A (2012)
Individual differences in music reward experiences. *Music Percept* **31**, 118-138.
[49] Karpati FJ, Giacosa C, Foster NE, Penhune VB, Hyde KL (2015) Dance and the brain: a review. *Ann
N Y Acad Sci* **1337**, 140-146.
[50] Hackney ME, Earhart GM (2010) Effects of dance on gait and balance in Parkinson's disease: a
comparison of partnered and nonpartnered dance movement. *Neurorehabil Neural Repair* **24**,
384-392.
[51] Hackney ME, Kantorovich S, Earhart GM (2007) A study on the effects of Argentine tango as a
form of partnered dance for those with Parkinson disease and the healthy elderly. *Am J Dance
Ther* **29**, 109-127.
[52] Chan JS, Wu J, Deng K, Yan JH (2020) The effectiveness of dance interventions on cognition in
patients with mild cognitive impairment: a meta-analysis of randomized controlled trials.
Neurosci Biobehav Rev **118**, 80-88.