A robust pipeline for the systematic validation of sRNA effectors suggests cross-kingdom communication in the symbiosis of *Serendipita indica* with *Arabidopsis thaliana*

Sabrine Nasfi, Saba Shahbazi, Katharina Bitterlich, Ena Šečić, Karl-Heinz Kogel & Jens Steinbrenner

Institute of Phytopathology, Research Centre for BioSystems, Land Use and Nutrition, Justus-Liebig-University Giessen, Heinrich-Buff-Ring 26, 35392 Giessen, Germany.

*Correspondence:

karl-heinz.kogel@agrar.uni-giessen.de

All E-mail addresses:

Sabrine.Nasfi@agrar.uni-giessen.de (ORCID no.: 0000-0001-5916-0824)

Karl-Heinz.Kogel@agrar.uni-giessen.de (ORCID no.: 0000-0003-1226-003X)

Jens.Steinbrenner@gmail.com (ORCID no.: 0000-0003-1008-2268)

Ena.Secic@agrar.uni-giessen.de (ORCID no.: 0000-0002-6469-4895)

Shahbazi.saba@yahoo.com (ORCID no.: 0009-0005-9222-3715)

katha.bitterlich@gmail.com (ORCID no.: 0009-0003-5976-1322)

Abstract

Cross-kingdom RNA interference (ckRNAi), the bidirectional communication between microbes and their host counterparts, is a key element in the outcome of host colonization. Whether mutualistic fungi of the Serendipitaceae family with their broad host range use small RNA effectors (sRNAs) to colonize plant roots is still under debate. To investigate if ckRNAi is a factor in the symbiosis of *Serendipita indica* (*Si*) with *Arabidopsis thaliana* (*Ath*), we established a pipeline to validate expression, translocation and post-transcriptional gene silencing of host genes by *Si*-derived sRNAs (*Si*sRNAs). First, we confirmed the expression of *Si*sRNAs both in axenic fungal culture and during *Ath* root colonization using stem-loop PCR. Then, to verify the translocation of putative *Si*sRNA effectors, an ARGONAUTE 1 immuno-purification assay (*At*AGO1-IP) was employed, detecting fungal *Si*sRNAs being loaded into the plant RNAi machinery in *Si*-colonised roots. Subsequently, *Si*sRNAs and artificial sRNAs (amiRNAs), were transiently expressed in *Ath* protoplasts to test their gene silencing activity. Stem-loop PCR confirmed expression of sRNA effectors and qPCR validated post-transcriptional gene silencing of their predicted target genes involved in cell wall organization, hormonal signalling regulation, plant immunity and gene expression. Moreover, 5'-RLM-RACE analysis revealed amiRNA-mediated canonical cleavage in *Arabidopsis* targets. In conclusion, this study provides a blueprint for rapid selection and analysis of sRNA effectors in plant-microbe interactions in general and suggests cross-kingdom communication in the Sebacinoid symbiosis.

Keywords

Arabidopsis, AGO1-IP, cross-kingdom communication, *Serendipita indica,* sRNA effectors, symbiosis, 5'-RLM-RACE, RNA interference

Introduction

RNA interference (RNAi) is a regulatory process in most eukaryotes in which gene expression is silenced at the transcriptional or post-transcriptional level by the action of small (s) RNA and is associated with the control of genome stability, developmental processes, and responses to biotic and abiotic stresses (Zhan and Meyers, 2023). Two types of sRNAs predominate in plants: Micro (mi) RNAs are 20-22 nucleotides (nt) in length processed from primary miRNAs (pri-miRNAs) containing a stem-loop structure encoded by microRNA (*MIR)* genes (Meyers and Axtell, 2019; Reinhart *et al.,* 2002). In contrast, small-interfering (si) RNAs are RNA molecules of 20-24 nt in length produced from longer endogenous RNA templates or from exogenous RNA sequences (viruses or transgenes). miRNA and siRNA precursors are processed by RNase III-like endonucleases named DICER-like (DCL), then bound by ARGONAUTE (AGO) proteins, which are incorporated into the RNA-Induced Silencing Complex (RISC), where they target complementary mRNA transcripts to induce post-transcriptional gene silencing (PTGS) (Iwakawa & Tomari, 2022). In addition to the involvement in intracellular RNAi, plant sRNAs can move intercellularly across plasmodesmata and systemically across the phloem and thus act cell non-autonomous and can also move between interacting species. In the latter case, sRNA effectors are transferred bidirectionally between plant hosts and pathogenic microbes to modulate defence and virulence, a phenomenon called cross-kingdom RNAi (ckRNAi) (Göhre and Weiberg, 2023; Mahanty *et al*., 2023; Wang *et al*., 2017). ckRNAi in plant-pathogen interactions was discovered in 2013 when the necrotrophic fungus *Botrytis cinerea* was shown to induce silencing of host defence genes in *Solanum lycopersicum* and *Arabidopsis thaliana* (*Ath*) (Weiberg *et al.,* 2013). Since then, several reports supported the existence of ckRNAi (Derbyshire *et al.,* 2019; Wang *et al.,* 2016) and showed that sRNA effectors exchange can be bidirectional (Zhang *et al.,* 2016).

Already in 2006, Huang and colleagues reported reduced nematode infectivity when root-knot nematodes (RKN) fed on roots of Arabidopsis plants expressing double-stranded (ds) RNA targeting the *16D10* gene, which encodes a conserved

secretory RKN root growth-stimulating peptide known to be involved in parasitism (Huang *et al.,* 2006). Later, it was shown that sRNA can indeed move from a plant to an attacking pathogen in the interaction of barley and wheat with the powdery mildew fungus (*Blumeria graminis*) (Nowara *et al.,* 2010). Transient transformation with a plasmid overexpressing dsRNA directed against the fungal *AVR10* gene increased the susceptibility of leaf epidermal cells to the corresponding *Blumeria graminis* races, a technique termed host-induced-gene-silencing (HIGS). Since then, numerous reports have shown that HIGS is an efficient strategy for pests and disease control in crops (Cai et al., 2018; Koch et al., 2013; Liu et al., 2020; Sang & Kim, 2020; Zand Karimi & Innes, 2022).

Although ckRNAi and related artificial processes such as HIGS were discovered in plants more than 10 years ago, the molecular mechanisms are still not fully understood. For instance, the mode of bidirectional transfer of RNAs in extracellular vesicles (EVs) is still a matter of debate (He, Cai, *et al.,* 2021; He, Hamby, *et al.,* 2021; Nasfi & Kogel, 2022; Rutter & Innes, 2018), and the occurrence in mutualistic symbioses is little researched but an increasing number of studies are shedding the light on ckRNAi in plant-symbiont interactions (Qiao *et al.,* 2023).

ckRNAi in mycorrhizal symbioses seems to include miRNA and transfer RNA fragments (tRFs). For example, Pmic_miR-8, a miRNA from ectomycorrhizal fungus *Pisolithus microcarpus* enhanced the colonization of its host *Eucalyptus grandis and* thus plays a role in the beneficial ectomycorrhizal symbiosis (Wong-Bajracharya *et al.,* 2022). In addition, recent *in silico* analysis of the sRNA population from *Rhizophagus irregularis*, supported by degradome data from *Medicago truncatula*, has shown that some sRNAs of Rhizophagus modulate root colonization (Silvestri *et al.,* 2019). Moreover, tRFs identified in root nodule symbioses between soybean and the bacterial symbiont *Bradyrhizobium japonicum* hijack *Glycine max* AGO1 and are positive regulators of rhizobial infection and nodule formation (Ren *et al.,* 2019), further emphasising the existence of ckRNAi in plant-mutual interactions.

Key to understanding ckRNAi is the identification and functional validation of sRNA effectors, and tools are needed to study the mode of action of numerous sRNAs derived from extensive sRNA sequencing data. Here, we report on a pipeline for sRNA effectors validation by studying the interaction between the mutualistic basidiomycete fungus *Serendipita indica* (*Si*) and the dicot model plant *Ath*. *Si* was originally isolated from the roots of the shrubs *Prosopis juliflora* and *Ziziphus nummularia* in the Indian Thar desert (Verma *et al.,* 1998). The beneficial activity of the fungus includes plant growth promotion, plant resistance priming, enhanced nitrate and phosphate delivery, promotion of adventitious root and root hair formation, early flowering, support of higher seed yield, alteration in secondary metabolites, and hardening of tissue cultured plants (Glaeser *et al.,* 2016; Qiang *et al.,* 2012; N. Verma *et al.,* 2022; Weiß *et al.,* 2016; Xu *et al.,* 2018; Zuccaro *et al.,* 2011). *Si* transfers protein effectors to host cells to exploit the host´s metabolism and promote microbial colonization (Akum *et al.,* 2015; Osborne *et al.,* 2023). To further explore the potential of *Si* we recently demonstrated the global change in sRNA profiles in the interaction of *Si* and the grass model *Brachypodium distachyon* (*Bd*) (Šečić et al., 2021). Among *Bd*- and *Si*-generated sRNAs with putative functions in the interacting organism, we found proteins involved in circadian clock, flowering regulation and immunity as potential targets of putative fungal sRNA effectors.

In the present work, we analysed the function of *Si*-derived sRNAs (*Si*sRNAs) in more detail by investigating the interaction between *Si* and *Ath*. By performing an *At*AGO1 immunoprecipitation assay, we confirmed the loading of *Si*sRNA into the RNAi machinery of *Ath*. We developed a functional protoplast assay to validate putative fungal sRNA effectors by testing their gene-silencing activity. Furthermore, we confirmed sRNA transformation and expression in plant protoplasts with stem-loop PCR and recorded downregulation of the corresponding predicted host target genes by qPCR. Finally, we confirmed sRNA effector candidate-mediated degradation of host mRNA by 5'-RLM-RACE. The data

support naturally occurring ckRNAi between the mutualistic fungus *Serendipita indica* and *Arabidopsis thaliana*.

Materials and methods

Plant material and growth conditions

For protoplast isolation, *Arabidopsis thaliana* plants of ecotype Columbia-0 (Col-0) were cultivated in a 4:1 ratio of T-type soil (F.-E. Typ Nullerde, Hawita) and crystal quartz sand mixture for 4 to 5 weeks in a growth chamber in a 19℃ day/18℃ night (9 h of light, 150 µmol photons $m^{-2}\,s^{-1}$) and a constant relative humidity of 60%. For interaction studies of *Ath* roots and *Serendipita indica* (IPAZ-11827, Institute of Phytopathology, Giessen, Germany), plants were grown on vertical square Petri dishes on an ATS medium (Lincoln *et al.,* 1990) without sucrose and supplemented with 4.5 g/L Gelrite (Duchefa #G1101) in a 22°C day/18°C night cycle (8 h of light). Roots of 14-day-old plants were inoculated with 1 mL of a suspension of 500,000 chlamydospores/mL in $H_2O_{0.02\%\ Tween\ 20}$ per Petri dish.

Plasmid construct

AtMIR390a distal stem-loop was digested from the pMDC32B-AtMIR390a-B/c vector (Plasmid #51776, https://www.addgene.org, Carbonell *et al.,* 2014) using Fast digest enzymes EcoRI and HindIII (Thermo Fisher Scientific, FD0274 & FD0505). The pUC18-AtMIR390a-B/c construct was generated by ligating the AtMIR390a precursor into the pUC18 vector (plasmid #50004) backbone using T4 DNA Ligase (Thermo Fischer Scientific, EL0011). The *RFP* insert was digested from the pBeaconRFP_GR vector (Bargmann & Birnbaum, 2009) using the restriction enzyme NdeI (NEB, R0111S) and further ligated into the NdeI site of the pUC18-AtMIR390a-B/c vector. 75mer oligonucleotides comprising the amiRNAs and *Si*sRNAs sequences were cloned into the BsaI-sites by Golden Gate strategy using BsaI-HF®v2 (NEB, R3733) and T4 DNA Ligase (Thermo Fischer Scientific, EL0011) in a single reaction, replacing the toxic ccdB gene which allows the selection of 75-

mer-positive clones. Before cloning 75-mer oligonucleotides, a third BsaI-site in
the pUC18 backbone which would interfere with the Golden-Gate cloning was
removed by site-directed mutagenesis using Phusion High-Fidelity DNA
polymerase (Thermo Fisher Scientific, F553S) and the Fast digest Dpn I
endonuclease (Thermo Fisher Scientific, FD1703) as well as custom-designed site-
directed mutagenesis primers pUC18-Mut-Fwd and pUC18-Mut-Rev.

The artificially designed amiRNAs and *Si*sRNAs and their corresponding predicted
targets are listed in Supplemental Tables 1 & 2. 75-mer oligonucleotides for
amiRNAs and *Si*sRNAs were designed following the P-SAMS algorithm developed
by the James Carrington lab (Fahlgren *et al.,* 2016) available at http://p-
sams.carringtonlab.org. Oligonucleotides are listed in Supplemental Table 3. An
in-silico design of the cloning was performed using Snapgene software
(www.snapgene.com) and the final pUC18-AtMIR390a-RFP-sRNA vector map is
shown in Supplement Figure 1.

Protoplast transformation

The "Tape-Arabidopsis Sandwich" method was used for protoplast isolation (Wu
et al., 2009). The transformation was performed following the transient expression
of recombinant genes using the Arabidopsis mesophyll protoplast (TEAMP)
approach (Yoo *et al.,* 2007) with minor modifications. Since the transformation of
protoplasts was key to all subsequent experiments, we attempted to optimize its
efficiency by varying *i.* protoplast concentration (10x 10^4 and 5x 10^4
protoplasts/mL), *ii.* plasmid concentration (20 µg, 30 µg, and 40 µg) pUC18-
AtMIR390a-RFP-sRNA and *iii*. incubation time (24 and 48 h). High transformation
efficiency was reported when using 5x 10^4 protoplasts/ml with a plasmid
concentration of 30 µg and an incubation time of 24 h, which was comparable to
the reported transformation efficiencies of *Ath* leaf protoplasts (Yoo *et al.,* 2007).
Total cell numbers were counted using a Fuchs-Rosenthal counting chamber
under a light microscope. A microscopic check of protoplast transformation was
done using an epifluorescence microscope (TCS SP2 Leica). Images were acquired

with the image software Leica Application Suite (LAS) and all counting was

performed in triplicates. Transformed protoplasts were stored at -80°C for further

studies.

RNA extraction and quality analysis

Total RNA was extracted from protoplasts using QuickRNA™ Miniprep Kit (Zymo

Research, R1050) with an on-column DNase I treatment. Total RNA from *Si-Ath*

interaction growing on ATS plates was extracted using Direct-zol RNA Miniprep

(Zymo Research, R2051) with an on-column DNase I treatment. The RNA

concentration was determined with the NanoDrop ND-1000 Spectrophotometer

(Thermo Fisher Scientific, USA) and the purity was determined by measuring

A260/280 and A260/230 ratios. The quality of the RNA extracted from transformed

protoplasts was checked using the Agilent 2100 Bioanalyzer Nano Chip (Agilent,

Germany).

Stem-Loop-RT-End-PCR

*Si*sRNAs and the designed amiRNAs were used as templates to design specific

stem-loop (SL) primers matching the corresponding sRNA over six nt at the 3´ end.

Hairpin primers (cDNAhp) and forward primers were designed using the tool

published in Adhikari et al (2013) based on Varkonyi-Gasic et al., (2007) (Adhikari

et al., 2013; Varkonyi-Gasic & Hellens, 2011). Stem-loop PCR was performed as

described by Werner et al., (2021) (Werner *et al.,* 2021). End-point PCR was used to

assess the expression of *Si*sRNAs and amiRNAs. The PCR program was optimized

to 95 °C for 5 min; followed by 40 cycles of 95 C for 30 s, 60°C for 30 s, 72°C for 30

s; and 72°C for 5 min and then held at 4 °C. PCR products were separated and

visualized in 2 % TBE -agarose gels. Stem-loop PCR primers are listed in

Supplemental Table 4.

AGO-IP

To perform AGO1 immunoprecipitation, 5 g of *Si* mycelium or *Ath* roots inoculated

with *Si* were ground to a fine powder using a precooled mortar and pestle. Co-

immunoprecipitation, sRNA recovery, and western blot analysis were performed

as described in the publication of Dunker et *al.,* 2021. Anti-AGO1 polyclonal

antibody (Agrisera, AS09 527) and IRDye® 800CW Goat anti-Rabbit IgG Secondary

Antibody (LI-COR, 926-32211) were used.

PTGS detection by qPCR

The standard curve method was used to test the efficiency of the qPCR transcript

primers using a serial dilution of *Ath* cDNA library along with 3 different primer

concentrations (0.4 µM, 0.2 µM and 0.1 µM for each primer) and 5 µL of SybrGreen

(Sigma-Aldrich). The total volume of 10 µL and three technical replicates are

considered for each reaction. Before master mix preparation, 2 µL ROX (CRX

reference dye, Promega, C5411) were added to 1 mL SybrGreen as a passive

reference dye that allows fluorescent normalization for qPCR data. For the

preparation of cDNA first strands, 500 ng of RNA samples from transformed *Ath*

protoplasts and control protoplasts were used. qPCR was performed using the

QuantStudio 5 Real-Time PCR system (Applied Biosystems) in the 384-well plate as

follows: 50°C for 2 min, 95°C for 10 min, followed by 40 cycles of 95°C for 15 s, and

60°C for 1 min, with an additional melt curve stage consisting of 15 s at 95°C, 1

min at 60°C, and 15 s at 95°C.

To evaluate the relative quantification of *Ath* transcripts by qPCR, the level of gene

expression was standardized to the control protoplasts. Fold change of *ΔΔ*ct values

and standard errors were calculated for all mean values. All qPCR primers are

listed in Supplemental Table 5.

5´-RLM-RACE

RNA ligase-mediated rapid amplification of cDNA ends (5´-RLM-RACE) was

performed using FirstChoice® RLM-RACE kit (Thermo Fisher Scientific) following

the manufacturer protocol and omitting the dephosphorylation and decapping

steps. One µg of RNA isolated from *At*h protoplasts and control protoplasts were

used as a template and ligated to the 5´-RACE adapter using T4 RNA Ligase [10

U/µL] (Thermo Fisher Scientific). The ligation reaction was used entirely to

generate the first cDNA strand. Two rounds of nested hot-start touch-down PCR

were performed using outer (first) and inner (second) 5´-RLM-RACE universal

primers in combination consecutively with gene outer specific and gene inner

specific primers. PCR products were evaluated in a 1.5% agarose gel and bands of

the expected size were excised. Products were cleaned with the Wizard® SV Gel

and PCR Clean-Up System (Promega) and cloned with the pGEM®-T Vector Systems

(Promega). For each band, six clones were selected for sequencing at LGC

genomics. The oligonucleotides used are listed in the Supplemental Table 6.

T-vector cloning and sequencing

Stem-loop PCR and 5´-RLM-RACE PCR amplification products were purified using

The Wizard® SV Gel and PCR Clean-Up System (Promega). Cloning of the different

PCR products was performed as described in the pGEM®-T Vector system

according to the instructor´s instructions (Promega). Sequencing was performed

at LGC Genomics (Berlin, Germany) and analysed using the Snapgene tool (GSL

Biotech, available at snapgene.com). All PCR primers are presented in

Supplemental Table 7.

Statistics Analysis

To assess the downregulation of *Ath* genes via qPCR, the 2−ΔΔCt method was used

for the quantification of *Ath* target transcript. The results are expressed as values

of the fold change of the transcript expression level in comparison to the control

and compared using a two-sided paired student t-test. Data showed a normal

distribution in the Shapiro-Wilk test or D´Agostino & Pearson test.

Results

We developed a pipeline for the systematic validation of sRNA effectors in the

mutualistic interaction of *Serendipita indica* and *Arabidopsis thaliana*, comprising i.

detection of fungal sRNAs (*Si*sRNAs) in axenic culture and *Si*-colonized *Ath* roots, ii.

detection of the association of *Si*sRNAs with the *At*AGO1 protein, and iii. validation

of the gene silencing activity of *Si*sRNAs in *Ath* protoplasts (Figure 1).

Figure 1. Schematic flowchart showing the molecular methods and outputs in this study. The turquoise colour indicates the different input material and purple indicates the wet-lab steps. Green represents the validated outputs of the study. For details also see Material and Methods. *Si*, *Serendipita indica*; *Ath*, *Arabidopsis thaliana*; PTGS, post-transcriptional gene silencing; amiRNA, artificial miRNA.

_Si_sRNAs are present in axenic cultures of _Serendipita indica_

In a previous RNAseq study, we identified potential fungal sRNA effectors in axenic *Si* cultures and *Si*-colonised *Brachypodium distachyon* roots (Šečić *et al.,* 2021). Here we further investigated a set of 15 putative *Si*sRNA effector candidates by focusing on 21 nt sRNAs encompassing all four 5'-terminal nucleotides: adenine (A), uracil (U), cytosine (C) and guanine (G) (Supplementary Table 8). We used multiplexed stem-loop PCR as a specific and sensitive method to amplify sRNAs followed by gel electrophoresis and showed that all 15 *Si*sRNAs could be amplified from 4-week-old axenic *Si* culture (Figure 2). Next, we cloned a subset of *Si*sRNAs (*Si*sRNA21, *Si*sRNA23, *Si*sRNA24, *Si*sRNA28, *Si*sRNA154 and *Si*sRNA296) and subjected them to Sanger sequencing, which further confirmed the expected *Si*sRNA sequences.

Figure 2: Stem-loop PCR analysis of *Si*sRNA detected in axenic cultures of

Serendipita indica. Multiplexed stem-loop PCR confirmed the presence of *Si*sRNAs

in a 4-week-old *Si* axenic culture. A PCR product of the expected size of 62 bp (sum

of the 21 bp of the sRNA sequence, loop region, plus the forward primer and the

reverse primer length) is visualized in a 2% agarose gel. The plant AtmiR159a which

served as a negative control was not detected.

_Si_sRNAs are present in colonized _Arabidopsis thaliana_ roots

Next, we analysed the presence of nine *Si*sRNAs (*Si*sRNA21, *Si*sRNA23, *Si*sRNA24,

*Si*sRNA25, *Si*sRNA26, *Si*sRNA28, *Si*sRNA37, *Si*sRNA47, and *Si*sRNA62) (Supplemental

Table 8) in *Si*-colonized *Ath* roots. Roots of 14-day-old seedlings were inoculated

with *Si* chlamydospores and harvested at 3-, 7-, and 14 days post-inoculation (dpi).

Fungal colonization of roots was confirmed after 7 days both by confocal laser

scanning microscopy (CLSM) using the chitin-specific dye Wheat Germ Agglutinin

with Alexa Fluor 488 (WGA-AF-488) (Figure 3 A) and by PCR using both specific *ITS*

gene primers and *Si*-specific *Ubiquitin* primers (Supplemental Figure 2).

Multiplexed stem-loop PCR of total RNA extracted from *Si*-colonized *Ath* roots at

different time points detected eight out of the nine selected *Si*sRNAs (Figure 3 B-

D). Bands for *Si*sRNA21 and *Si*sRNA24 did not consistently show up at all three-

time points, likely because of the limitations associated with multiplexing the

hairpin cDNA primers (Colombo *et al.,* 2013; Kramer, 2011; Varkonyi-Gasic and

Hellens, 2011). Consistent with this notion, we detected the expression of

*Si*sRNA21 and *Si*sRNA24 at 7 dpi in a stem-loop PCR with less multiplexing

(Supplemental Figure 3). A subset of three *Si*sRNAs (*Si*sRNA21, *Si*sRNA24,

*Si*sRNA28) were cloned and submitted for Sanger sequencing, confirming the

expected *Si*sRNA sequences in *Si*-colonized *Ath* roots.

Figure 3: Arabidopsis roots colonized by *Serendipita indica* and gel analysis of

***Si*sRNAs in the *Si-Ath* interaction**. **(A)** Root colonization pattern at 7 dpi.

Fluorescence microscopy shows green chitin-specific WGA-AF488 staining of *Si*

hyphal walls (λexc494 nm, λem515) **(B)** Detection of *Si*sRNAs in roots at 3-, 7- and

14 dpi using multiplexed stem-loop PCR. Agarose gel analysis of *Si*sRNA detected

bands of the expected size of 62 bp. Plant AtmiR159a was used as a positive

control for a plant-expressed sRNA.

SisRNAs are loaded onto Arabidopsis AGO1

To further substantiate the hypothesis that ckRNAi occurs in the interaction

between *Si* and *Ath* and that *Si*sRNAs are likely to be loaded into the Arabidopsis

RISC complex, we conducted *At*AGO1/sRNA co-immunopurification (Co-IP) using

samples of *Si*-colonised roots, followed by stem-loop PCR analysis (Carbonell,

2017b; Dunker *et al.,* 2021). To exclude the possibility that the *Ath*-specific anti-

AGO1 antibody would cross-react with fungal AGOs, we first performed an

*At*AGO1/sRNA Co-IP using a 4-week-old *Si* axenic culture, followed by Western-Blot

(WB) analysis. No binding of the *At*AGO1-specific antibody was detected in samples

from *Si* axenic cultures confirming the specificity of the plant anti-AGO1 antibody

(Supplementary Figure 4A). In line with this, no *Si*sRNAs could be detected by stem-

loop PCR using *Si*sRNA-specific stem-loop primers (Supplementary Figure 4B).

Next, we performed *At*AGO1/sRNA Co-IP on samples from *Si*-colonized roots (*Si-*

Ath Co-IP). *Si*sRNA21, *Si*sRNA23, *Si*sRNA24, *Si*sRNA28 and the *At*AGO1-bound

AtmiR159a were detected by stem-loop PCR (Figure 4), whereas AtmiR393a*,

which is known to preferably bind *At*AGO2 (Mi *et al.,* 2008), was undetectable.

Stem-loop PCR products from the *Si-Ath* Co-IP fraction were cloned and

sequenced. The sequencing results confirmed the presence of plant AtmiR159a as

well as *Si*sRNA2*4* and *Si*sRNA28 with the exact same sequences detected from *Si*

axenic cultures and *Si*-colonized *Ath* roots. In addition, two *Si*sRNAs with almost

identical sequences to *Si*sRNA21 and *Si*sRNA23 were also co-immunoprecipitated

(Supplementary Table 9). These results show that detected *Si*sRNAs are bound to

*At*AGO1, supporting the notion that *Si*sRNAs are transferred into *Ath* root cells

during root colonisation and loaded into the plant RNAi machinery.

Figure 4: *Si*sRNAs co-immunoprecipitated with *At*AGO1 in *Si*-colonized

Arabidopsis roots. Stem-loop PCR analysed in agarose gel confirms the presence

of *Si*sRNA21, *Si*sRNA23 *Si*sRNA24, *Si*sRNA28, as well as AGO1 bound AtmiR159a but

not AGO2 bound AtmiR393a* in AGO1-co-IP samples from *Si*-colonized *Ath* roots.

***Si*sRNAs mediate PTGS in transformed Arabidopsis protoplasts**

Next, we transiently transformed protoplasts with *Si*sRNAs to demonstrate their

gene-silencing activity. *Ath* protoplasts were transformed with a *35S::AtMIR390a*

expression construct containing *Si*sRNA21, *Si*sRNA24, *Si*sRNA28 and *Si*sRNA154

(confirmed in *Si* axenic culture). Information on these *Si*sRNAs, including their

predicted *Ath* target genes, their target predicted scores (TPS), indicating the

degree of mismatches with the predicted *Ath* target sequence, their 5'-terminal

nucleotides and an alignment to their corresponding target fragment are shown

in Supplemental Table 1. We achieved a transformation efficiency of 65% to 90%,

calculated as the ratio between the red fluorescing protoplasts and the total

number of viable protoplasts (Supplemental Figure 5). Duplexed stem-loop PCR

confirmed the expression of the four *Si*sRNAs. All four *Si*sRNAs were successfully

amplified (Figure 5). The amplification products were cloned into the pGEM-T®

cloning vector and sequencing confirmed their identity.

Figure 5. Duplexed stem-loop PCR confirming expression of *Si*sRNAs in Arabidopsis protoplasts. Agarose gel analysis of stem-loop PCR confirmed the detection of all four expressed *Si*sRNA in transformed protoplasts 24 hours post-transformation (hpt), but not in control protoplasts. Plant AtmiR159a was used as a positive control.

Next, we investigated the ability of the expressed *Si*sRNAs to induce PTGS of predicted target genes in protoplasts. *Si*sRNA24 was predicted to target five *Ath* genes: *AT1G57590 (PECTIN ACETYESTERASE 2)* involved in cell wall organization, *AT1G63180 (UDP-D-GALACTOSE 4-EPIMERASE 3)* involved in pollen development, *AT1G65090 (SEED LIPID DROPLET PROTEIN1*) involved in lipid droplet-plasma membrane tethering, *AT4G15765 (FAD/NAD(P)-binding oxidoreductase family protein)* involved in jasmonic acid-mediated signalling, and *AT5G16680* (PAIPP2, PHD2) involved in regulation of gene expression (see Supplemental Table 1).

Expression of *Si*sRNA24, which has U at the 5'-terminus, and thus a preference for AGO1 (Mi *et al.*, 2008; Montgomery *et al.*, 2008; Takeda *et al.*, 2008), significantly reduced the mRNA abundance of *AT1G63180*, *AT1G65090*, *AT4G15765* and *AT5G16680* in protoplasts 24 h after transformation by 63%, 56%, 45%, and 73%, respectively, (Figure 6 B). Similarly, *Si*sRNA154, also with 5´ U, is predicted to target *AT2G47600 (MAGNESIUM/PROTON EXCHANGER)* involved in magnesium, iron and zinc ion transport and *AT4G32160 (Phox (PX) domain-containing protein)* involved in signal transduction. Expression of *Si*sRN154 significantly downregulated the accumulation of *AT2G47600* mRNA by 37% but not of *AT4G32160*. (Figure 6 D). *Si*sRNA28 has 5' Adenine (A), therefore a loading preference for AGO2 and AGO4 (Mi et al., 2008; Montgomery et al., 2008; Takeda et al., 2008). Consistent with this, expression of *Si*sRNA28 did not result in downregulation of any of the three predicted target genes: *AT1G05180 (AUXIN RESISTANT 1)* involved in auxin-activated signalling pathway and response to cytokinin, *AT5G55930 (ARABIDOPSIS THALIANA OLIGOPEPTIDE TRANSPORTER 1)* an oligopeptide transporter, and *AT3G06670* (*PLATINUM SENSITIVE 2 LIKE*) with a regulatory function in ncRNA processing (Figure 6 C). Finally, expression of *Si*sRNA21 with 5' Cytosine (C) and thus a preference for AGO5 (Mi et al., 2008; Montgomery et al., 2008; Takeda et al., 2008) resulted in downregulation of predicted target transcripts *AT5G25350* (*EIN3-BINDING F BOX PROTEIN 2*), a negative regulator of the ethylene-activated signalling pathway, and *AT5G37600 (ARABIDOPSIS GLUTAMINE SYNTHASE 1)* involved in nitrate assimilation, by 51%, and 16%, respectively (Figure 6 A).

Figure 6. qPCR analysis of target gene silencing in Arabidopsis protoplasts 24 h post transformation with *Si*sRNAs vs. control (protoplast without *Si*sRNA construct). The mRNA transcript levels in transformed protoplasts (+) expressing **(A)** *Si*sRNA21, **(B)** *Si*sRNA24, **(C)** *Si*sRNA28 and **(D)** *Si*sRNA154 normalized against housekeeping gene *Ubiquitin* (*AT3G62250*) are displayed as fold change relative to control protoplasts (-). Data are the average of two to three biological replicates ± standard deviation. Asterisks indicate difference at p ≤ 0.05(*), p ≤ 0.01 (**) and p ≤ 0.001 (***) according to Student's t-test. ns= not significant.

5'-RLM-RACE reveals non-canonical cleavage of Arabidopsis target genes in protoplasts expressing *Si*sRNAs with preference for AGO1

We selected *Si*sRNA24 to study the cleavage pattern of its significantly down-regulated target transcripts *AT1G65090* and *AT4G15765* using RNA-ligase-mediated

Rapid Amplification of cDNA Ends (5'-RLM-RACE) (Llave *et al.,* 2002; Ueno *et al.,*

2022). For the predicted target transcript *AT1G65090,* no band corresponding to

an expected canonical cleavage product was detected (Supplemental Figure 6A)

(Carbonell, 2017c; Carthew & Sontheimer, 2009). For target gene *AT4G15765*, we

detected a single band of 300 bp in protoplasts expressing *Si*sRNA24*,* which was

shorter than the expected size (443 bp), while double bands were observed for the

control protoplasts. PCR amplicon from both transformed and control protoplasts

were cloned into the pGEM-T® cloning vector and sequenced. Sequence analysis

confirmed the alignment of the 300 bp RLM-RACE product with *AT4G15765*.

However, it did not confirm the 5´-end position, between nt 10 and 11, predicted

for *Si*sRNA24-guided canonical cleavage (Supplemental Figure 6B). However, the

absence of cleavage and/or a canonical cleavage product does not exclude the

potential of *Si*sRNA24 in mediating predicted *Ath* gene silencing as many previous

studies have highlighted the challenge of identifying sRNAs cleavage sites in plants

(Brousse et al., 2014; Werner *et al.,* 2021).

amiRNAs mediate PTGS in transformed Arabidopsis protoplasts

Working with the naturally occurring *Si*sRNAs can be challenging due to the varying

degree of mismatches with their predicted target genes. The position and the

significance of these mismatches are crucial factors for PTGS and for the

mediation of a canonical target cleavage. To overcome these challenges and to

have a proof of concept for the sRNA effector validation pipeline we used

artificially designed microRNA (amiRNAs) that, unlike *Si*sRNAs, match their

predicted target genes at a higher degree and exhibit a U at their 5´-end. To this

end, four amiRNAs were designed: amir21, amir24, amir154, and amir296, cloned

into pUC18-AtMIR390a-RFP vector and transformed into *Ath* protoplasts.

Subsequently, their expression was confirmed as described for *Si*sRNAs using

duplexed stem-loop PCR. Supplemental Table 2 provides information on these

amiRNAs, including their predicted *Ath* target genes, their target predicted scores

(TPS) indicating the degree of mismatches between the amiRNAs and the *Ath*

target sequence, their 5'-terminal nucleotides, and an alignment to their

corresponding target fragment. The four amiRNAs were successfully expressed in

protoplast (Figure 7), amplification products were cloned into the pGEM-T® cloning

vector, and sequencing confirmed amiRNAs sequences.

Figure 7. Duplexed stem-loop PCR confirming expression of amiRNAs in

Arabidopsis protoplasts. Agarose gel analysis of all four expressed amiRNAs in

transformed protoplasts 24 hours post-transformation (hpt), but not in control

protoplasts. Plant AtmiR159a was used as a positive control.

The abundance of the four *Ath* transcripts *AT5G25350* (previously investigated for

PTGS in protoplasts transformed with *Si*sRNA21), *AT2G39020 (GCN5-RELATED N-*

ACETYLTRANSFERASE 8) involved in RNA metabolic process, *AT4G32160* (previously

investigated for PTGS in protoplasts transformed with *Si*sRNA154), and *AT2G45240*

(METHIONINE AMINOPEPTIDASE 1A) involved in removing the N-terminal methionine

from proteins, predicted to be targeted by amir21, amir24, amir154, and amir296,

respectively. Target gene expression was quantified 24 h after the transformation

of amiRNAs into protoplasts by qPCR. Downregulation of mRNA transcript level

was confirmed for *AT5G25350*, *AT2G39020*, *AT4G32160*, and *AT2G4524*0 by 68%,

43%, 50%, and 65%, respectively, compared to control (Figure 8).

Figure 8. qPCR analysis of target gene silencing in Arabidopsis protoplasts 24 h post transformation with amiRNAs. The mRNA transcript levels in transformed protoplasts (+) expressing amir21, amir24, amir154 and amir296 were normalized against housekeeping gene *Ubiquitin* (*AT3G62250*) and displayed as fold change relative to control protoplasts (-) (mock-treated without amiRNA construct). Data are the average of two to three biological replicates ± standard deviation. Asterisks indicate difference at $p \leq 0.05$(*), $p \leq 0.01$ (**) and $p \leq 0.001$ (***) according to Student's t-test. ns= not significant.

5'-RLM-RACE reveals canonical cleavage in protoplasts transformed with amiRNA

We selected amir296 and its predicted target gene *AT2G45240* to investigate the canonical PTGS cleavage site using 5-RLM-RACE. Samples were obtained from transformed protoplasts at 24 hpt, and from control samples. A distinct band of approximately 383 bp was amplified from protoplasts expressing amir296, consistent with the expected size of the canonical cleavage product predicted to be targeted by amir296. Unexpectedly, weaker DNA fragments of a similar size were detected in both control and non-treated protoplasts (Figure 9A). Cloning and subsequent sequencing of the amplicons confirmed that all sequences aligned with the *AT2G45240* sequence, and 50% of the obtained sequences from protoplasts transformed with pUC18-AtMIR390a-RFP-amir296 (three of six clones) exhibited the canonical cleavage site between nucleotide 10 and 11 of amir296. In contrast, PCR products from control and natural protoplasts contained an *AT2G45240* fragment but did not correspond to a canonical cleavage site, suggesting a possible cleavage by an endogenous miRNA near the amir296 binding site or RNA degradation (Figure 9B).

A

B

Figure 9. 5'-RLM-RACE *AT2G45240* target genes of amir296 after expression in Arabidopsis protoplasts (A) PCR products of the second nested 5´-RLM-RACE-PCR visualized in an EtBr-agarose gel. RNA was extracted from transformed

protoplasts with amir296 (T) and from the corresponding controls (control protoplasts (C) and natural protoplasts (NP). **(B)** Mapping of *AT2G45240* target cleavage products by 5´-RLM-RACE. The predicted base pairing between amir296 and *AT2G45240* is shown in blue. The red arrow indicates the detected canonical cleavage sites. The proportion of cloned 5´-RLM-RACE products at the different cleavage sites is shown in brackets. For amir296 transformed protoplasts, three colonies have the 5´-end at the expected position, opposite to nucleotides 10-11 of amir296, which is not the case for control and non-treated protoplasts.

Discussion

In this work, we developed an experimental pipeline for the analysis of putative fungal sRNA effectors and investigated their role in host target gene silencing. As we have previously reported on the identification and analysis of fungal sRNAs (*Si*sRNAs) in the interaction of *Si* and *Brachypodium distachyon* (Šečić *et al.,* 2021), we aimed to bridge the gap between the large amount of bioinformatic data available from this previous study, by establishing here a rapid, simple, and reproducible roadmap for the analysis of *Si*sRNA effectors and their target genes based on an Arabidopsis protoplast expression system. We validated the effectiveness of our pipeline, by investigating sRNAs detected in the mutualistic interaction between the root endophyte *Serendipita indica* and the model plant *Arabidopsis thaliana* and we confirmed the accumulation of a set of *Si*sRNAs in *Si*-colonized roots and their transfer to the plant cells. By immunopurification pull-down assay, we showed that *Si*sRNAs are loaded into the plant RNAi machinery, confirming the translocation of *Si*sRNA effectors into host cells. Using *At*AGO/sRNA co-IP we detected two *Si*sRNAs, *Si*sRNA24 and *Si*sRNA28 supporting the ckRNAi hypothesis. Additional *Si*sRNAs of 21 nt were identified in the *At*AGO1 Co-IP assay that had some mismatches when compared to the *Si*sRNA detected in *Si* axenic culture (Supplemental Table 9).

To gain more information about ckRNAi between *Si* and *Ath,* we generated a plant expression construct under the 35S promotor for sRNA expression in protoplasts based on pUC18 backbone vector and AtMIR390a distal stem-loop which can generate 21 nt amiRNAs or *Si*sRNAs. The AtMIR390a precursor belongs to the MIR390 family, which is highly conserved and precisely produced and processed in several plant species (Axtell *et al.,* 2006; Cuperus *et al.,* 2011). Carbonell et al (2014) tested the functionality of the AtMIR390a-based amiRNAs vector in repressing target transcript accumulation in Arabidopsis plants using the plant vector pMDC32BAtMIR390a-B/c, which enables the directed cloning of amiRNAs and their stable expression in dicots plants (Carbonell *et al.,* 2014). In our approach, we combined the advantages of plant expression vectors and protoplast transient expression systems, both considered as valuable tools, to study gene silencing and the molecular mechanisms of RNAi in plants (Tyurin *et al.,* 2020).

Arabidopsis protoplasts were transformed with the putative *Si*sRNA24, and its expression was validated by stem-loop PCR. Four of five predicted targets of *Si*sRNA24 exhibited significantly reduced abundance in the transformed protoplasts. Further analysis of the molecular cleavage pattern of *Si*sRNA24 by 5´-RLM-RACE identified noncanonical cut sites in the target gene *AT4G15765.* This finding may be explained by the fact that cleavage products might be present at a low abundance, falling below the detection threshold of the RLM-RACE method and making the detection of a precise canonical cleavage a challenge. Alternatively, it might be also possible that the cleavage site within the target mRNA could be located at a different position.

Another putative *Si*sRNA, *Si*sRNA154, which also exhibits a 5´ U and contains some mismatches to two potential mRNA targets, showed a significant downregulation only for the target *AT2G47600*. One major factor to consider when studying PTGS via sRNA effectors is a high target predicted score (TPS) between the sRNA and

their predicted targets. Previous studies have demonstrated that the number and position of mismatches between the sRNA and the corresponding target mRNA can significantly impact their binding affinity, eventually weakening the interaction and resulting in either non-significant or no downregulation (Ossowski *et al.,* 2008; Rhoades *et al.,* 2002). Studies have shown that mismatches within the sRNA seed region spanning nt 2 to 12/13 are possible, but they may reduce the sRNA activity, whereas mismatches at position 1 or between nt 14 and 21 are more acceptable (Carbonell, 2017a; Mallory *et al.*, 2004). Moreover, mismatches can also lead to a different mode of action for the sRNA, and instead of an mRNA degradation pathway, mismatches can trigger a deadenylation pathway or a translational repression pathway, thus causing a decrease in protein synthesis instead of an mRNA target degradation (Baulcombe, 2004; Carthew & Sontheimer, 2009). We also hypothesize that a high abundance level of a target mRNA may lead to less detectable downregulation, as its accessibility for the sRNA may have an impact on the silencing process and the detection of a canonical cleavage. It is also conceivable that the presence of other endogenous miRNAs, which may bind near or at the same mRNA binding site might disrupt sRNA/mRNA binding, modulate the regulatory outcome and affect the gene expression pattern.

To overcome the drawbacks of low target specificity of *Si*sRNAs, we designed additional experiments based on amiRNAs with almost a perfect match to *Ath's* predicted target transcript. Four amiRNAs, amir21, amir24, amir154 and amir296 (Supplemental Table 2), each with a 5´U were expressed in *Ath* protoplasts. All amiRNAs induced a significant reduction in mRNA abundance of their corresponding host-target transcripts, confirming their effective silencing activity in *Ath* protoplasts.

Although MIR390 associates preferably with AGO7, the association of AtMIR390a-derived amiRNAs that have a 5´ U is redirected to AGO1 (Carbonell, 2017b; Carbonell *et al.,* 2014; Cuperus *et al.,* 2010; Montgomery *et al.,* 2008).

Conventionally, an sRNA-AGO1-mediated cleavage results in a canonical cut

between nucleotides 10 and 11 of the sRNA at the corresponding mRNA site. This

cleavage is a confirmation of sRNA loading into AGO1, formation of the RISC

complex and degradation of the mRNA target (Addo-Quaye *et al.*, 2009; Huntzinger

& Izaurralde, 2011). amiRNAs used in our proof of concept approach are

algorithmically designed to exhibit a 5' U, thereby enabling their loading into the

AGO1 protein (Carbonell, 2017c). Consequently, they can enter the RISC complex

and induce a downregulation of the target gene. Using 5'-RLM-RACE, we showed

that amir296 can induce a potential *At*AGO1-mediated degradation of *AT2G45240*

(MAP1A, METHIONINE AMINOPEPTIDASE 1A) mRNA as we detected a canonical

cleavage site between positions 10 and 11 of amir296 in the selected transcript.

Expression of all four miRNAs resulted in a significant downregulation of their

corresponding mRNA targets. AT5G25350 is predicted to be targeted by both

amir21 and *Si*sRNA21. When expressing amir21 (with 5´ U and a TPS of 0), the

mRNA abundance of *AT5G25350* showed a downregulation of 68%. However, when

expressing *Si*sRNA21 (with a 5´C and a TPS of 5) we observed a downregulation of

51% of the same target gene. Another example is the predicted target gene

AT4G32160 targeted by amir154 and *Si*sRNA154 both with a 5´ U. The mRNA

abundance of *AT4G32160* in protoplasts expressing amir154 (with a TPS of 0)

resulted in a reduction of 50% but protoplasts expressing *Si*sRNA154 (with a TPS

of 4.5) did not show a reduction of *AT4G32160* transcript (Supplemental Figure 7).

These results raise the possibility that amiRNAs are more prone to mediate target

gene silencing but further studies with additional amiRNAs and putative sRNAs are

required to confirm these observations.

We also investigated whether amir296 with a 5´ U can mediate a canonical

cleavage of the predicted target gene *AT2G45240,* and indeed detected a cleavage

site of *AT2G45240* mRNA transcript between position 10 and 11 of amir296. 5´-

RLM-RACE is a challenging method to detect cleavage sites. High throughput

sequencing of samples from the *Si-Ath* interaction including transcriptional analysis and degradome sequencing are needed to better understand *Si*sRNA-mediated target cleavage. Moreover, consideration for sRNA non-canonical cleavage site may also be further investigated as different studies speculate that PTGS can still be induced through sRNA or phasiRNA and generate a non-canonical cleavage site (Brousse *et al*., 2014; Ivanova *et al*., 2022).

In conclusion, we have developed a transient protoplast expression system to investigate the potential role of putative *Si*sRNA effectors in regulating target gene expression in *Arabidopsis thaliana* and to explore whether *Si* uses sRNAs to establish a successful beneficial interaction with its host plant. This powerful validation tool will not only help to understand the underlying molecular mechanism of *Si*sRNA-mediated PTGS but will also allow us to examine ckRNAi in mutualistic plant-fungal association.

By studying *Si*sRNA-mediated PTGS, we have gained valuable insights into the possibility that the endophytic fungus *Si* exploits sRNA effectors for the establishment of its mutualistic symbiosis with plants. More targeted studies will show whether *Si* utilizes its sRNAs to potentially modulate the plant immune system and identify more genes that can be silenced or upregulated during the *Si-Ath* interaction. With the great potential of *Si* to colonize a wide host range of plants, we expect that findings from the study of the *Si-Ath* interaction can be translated to crop plants like barley, wheat, maize and rice. Moreover, amiRNAs have been shown to be effective in fine-tuning gene expression in plants with a high silencing level (Carbonell et al., 2014; Cisneros & Carbonell, 2020). Therefore, they can be viewed as a potential tool for RNAi-mediated crop protection strategies. Computational tools, along with *in vivo* and *in vitro* experimental setups, can expand our knowledge of ckRNAi and reveal the unknown about mutualistic

and pathogenic interaction for cost-effective basic research and agronomic
application.

Acknowledgements

The authors gratefully thank R. Eichmann for critically reading and correcting the
manuscript. We thank M. Ladera-Carmona and B.T. Werner for their comments
and suggestions. We thank the German Research Foundation (DFG) for supporting
our research on RNA in plant-pathogen interactions by granting the DFG research
unit RU5116 on exRNA to KHK. SN is supported by the Dr. Ernst-Leopold Klipstein-
Stiftung, Paderborn, Giessen.

Conflict of interest

The authors declare that they have no competing interests.

Author contributions

S.N. and J.S. conceived the study; S.N. and K-H.K. wrote the paper; J.S. and E.S.
generated Arabidopsis target predicted genes data. S.N., K.B., and S.S. conducted
the functional experiments. All authors read and approved the final manuscript.

References

References

Addo-Quaye, C., Snyder, J. A., Park, Y. B., Li, Y.-F., Sunkar, R., & Axtell, M. J. (2009).

Sliced microRNA targets and precise loop-first processing of MIR319

hairpins revealed by analysis of the Physcomitrella patens degradome.

RNA, *15*(12), 2112–2121. https://doi.org/10.1261/rna.1774909

Adhikari, S., Turner, M., & Subramanian, S. (2013). Hairpin Priming Is Better

Suited than In Vitro Polyadenylation to Generate cDNA for Plant miRNA

qPCR. *Molecular Plant*, *6*(1), 229–231. https://doi.org/10.1093/mp/sss106

Akum, F. N., Steinbrenner, J., Biedenkopf, D., Imani, J., & Kogel, K.-H. (2015). The

Piriformospora indica effector PIIN_08944 promotes the mutualistic

Sebacinalean symbiosis. *Frontiers in Plant Science*, *6*, 906.

https://doi.org/10.3389/fpls.2015.00906

Axtell, M. J., Jan, C., Rajagopalan, R., & Bartel, D. P. (2006). A Two-Hit Trigger for

siRNA Biogenesis in Plants. *Cell*, *127*(3), 565–577.

https://doi.org/10.1016/j.cell.2006.09.032

Bargmann, B. O. R., & Birnbaum, K. D. (2009). Positive Fluorescent Selection

Permits Precise, Rapid, and In-Depth Overexpression Analysis in Plant

Protoplasts. *Plant Physiology*, *149*(3), 1231–1239.

https://doi.org/10.1104/pp.108.133975

Baulcombe, D. (2004). RNA silencing in plants. *Nature*, *431*(7006), Article 7006.

https://doi.org/10.1038/nature02874

Brousse, C., Liu, Q., Beauclair, L., Deremetz, A., Axtell, M. J., & Bouché, N. (2014). A

non-canonical plant microRNA target site. *Nucleic Acids Research*, *42*(8),

5270–5279. https://doi.org/10.1093/nar/gku157

Cai, Q., Qiao, L., Wang, M., He, B., Lin, F.-M., Palmquist, J., Huang, S.-D., & Jin, H.

(2018). Plants send small RNAs in extracellular vesicles to fungal pathogen

to silence virulence genes. *Science (New York, N.Y.)*, *360*(6393), 1126–1129.

https://doi.org/10.1126/science.aar4142

Carbonell, A. (2017a). Artificial small RNA-based strategies for effective and

specific gene silencing in plants. *Plant Gene Silencing: Mechanisms and*

Applications, 110–127. https://doi.org/10.1079/9781780647678.0110

Carbonell, A. (2017b). Immunoprecipitation and High-Throughput Sequencing of

ARGONAUTE-Bound Target RNAs from Plants. In A. Carbonell (Ed.), *Plant*

Argonaute Proteins: Methods and Protocols (pp. 93–112). Springer.

https://doi.org/10.1007/978-1-4939-7165-7_6

Carbonell, A. (2017c). Plant ARGONAUTEs: Features, Functions, and Unknowns. In

A. Carbonell (Ed.), *Plant Argonaute Proteins: Methods and Protocols* (pp. 1–

21). Springer. https://doi.org/10.1007/978-1-4939-7165-7_1

Carbonell, A., Takeda, A., Fahlgren, N., Johnson, S. C., Cuperus, J. T., & Carrington,

J. C. (2014). New generation of artificial MicroRNA and synthetic trans-

acting small interfering RNA vectors for efficient gene silencing in

Arabidopsis. *Plant Physiology*, *165*(1), 15–29.

https://doi.org/10.1104/pp.113.234989

Carthew, R. W., & Sontheimer, E. J. (2009). Origins and Mechanisms of miRNAs

and siRNAs. *Cell*, *136*(4), 642–655. https://doi.org/10.1016/j.cell.2009.01.035

Cisneros, A. E., & Carbonell, A. (2020). Artificial Small RNA-Based Silencing Tools

for Antiviral Resistance in Plants. *Plants*, *9*(6), Article 6.

https://doi.org/10.3390/plants9060669

Colombo, S., Nielsen, H. M., & Foged, C. (2013). Evaluation of carrier-mediated

siRNA delivery: Lessons for the design of a stem-loop qPCR-based

approach for quantification of intracellular full-length siRNA. *Journal of*

Controlled Release, *166*(3), 220–226.

https://doi.org/10.1016/j.jconrel.2013.01.006

Cuperus, J. T., Carbonell, A., Fahlgren, N., Garcia-Ruiz, H., Burke, R. T., Takeda, A.,

Sullivan, C. M., Gilbert, S. D., Montgomery, T. A., & Carrington, J. C. (2010).

Unique Functionality of 22 nt miRNAs in Triggering RDR6-Dependent siRNA

Biogenesis from Target Transcripts in Arabidopsis. *Nature Structural &*

Molecular Biology, *17*(8), 997–1003. https://doi.org/10.1038/nsmb.1866

Cuperus, J. T., Fahlgren, N., & Carrington, J. C. (2011). Evolution and Functional

Diversification of MIRNA Genes. *The Plant Cell*, *23*(2), 431–442.

https://doi.org/10.1105/tpc.110.082784

Derbyshire, M., Mbengue, M., Barascud, M., Navaud, O., & Raffaele, S. (2019).

Small RNAs from the plant pathogenic fungus Sclerotinia sclerotiorum

highlight host candidate genes associated with quantitative disease

resistance. *Molecular Plant Pathology*, *20*(9), 1279–1297.

https://doi.org/10.1111/mpp.12841

Dunker, F., Lederer, B., & Weiberg, A. (2021). Plant ARGONAUTE Protein

Immunopurification for Pathogen Cross Kingdom Small RNA Analysis. *Bio-

Protocol*, *11*(3), e3911. https://doi.org/10.21769/BioProtoc.3911

Fahlgren, N., Hill, S. T., Carrington, J. C., & Carbonell, A. (2016). P-SAMS: A web site

for plant artificial microRNA and synthetic trans-acting small interfering

RNA design. *Bioinformatics*, *32*(1), 157–158.

https://doi.org/10.1093/bioinformatics/btv534

Glaeser, S. P., Imani, J., Alabid, I., Guo, H., Kumar, N., Kämpfer, P., Hardt, M., Blom,

J., Goesmann, A., Rothballer, M., Hartmann, A., & Kogel, K.-H. (2016). Non-

pathogenic Rhizobium radiobacter F4 deploys plant beneficial activity

independent of its host Piriformospora indica. *The ISME Journal*, *10*(4), 871–

884. https://doi.org/10.1038/ismej.2015.163

Göhre, V., & Weiberg, A. (2023). RNA Dialogues in Fungal–Plant Relationships. In

B. Scott & C. Mesarich (Eds.), *Plant Relationships: Fungal-Plant Interactions*

(pp. 31–51). Springer International Publishing. https://doi.org/10.1007/978-

3-031-16503-0_2

He, B., Cai, Q., Qiao, L., Huang, C.-Y., Wang, S., Miao, W., Ha, T., Wang, Y., & Jin, H.

(2021). RNA-binding proteins contribute to small RNA loading in plant

extracellular vesicles. *Nature Plants*, *7*(3), 342–352.

https://doi.org/10.1038/s41477-021-00863-8

He, B., Hamby, R., & Jin, H. (2021). Plant extracellular vesicles: Trojan horses of

cross-kingdom warfare. *FASEB BioAdvances*, *3*(9), 657–664.

https://doi.org/10.1096/fba.2021-00040

Huang, G., Allen, R., Davis, E. L., Baum, T. J., & Hussey, R. S. (2006). Engineering

broad root-knot resistance in transgenic plants by RNAi silencing of a

conserved and essential root-knot nematode parasitism gene. *Proceedings*

of the National Academy of Sciences, *103*(39), 14302–14306.

https://doi.org/10.1073/pnas.0604698103

Huntzinger, E., & Izaurralde, E. (2011). Gene silencing by microRNAs:

Contributions of translational repression and mRNA decay. *Nature Reviews*

Genetics, *12*(2), Article 2. https://doi.org/10.1038/nrg2936

Ivanova, Z., Minkov, G., Gisel, A., Yahubyan, G., Minkov, I., Toneva, V., & Baev, V.

(2022). The Multiverse of Plant Small RNAs: How Can We Explore It?

International Journal of Molecular Sciences, *23*(7), 3979. PubMed.

https://doi.org/10.3390/ijms23073979

Iwakawa, H., & Tomari, Y. (2022). Life of RISC: Formation, action, and degradation

of RNA-induced silencing complex. *Molecular Cell*, *82*(1), 30–43.

https://doi.org/10.1016/j.molcel.2021.11.026

Koch, A., Kumar, N., Weber, L., Keller, H., Imani, J., & Kogel, K.-H. (2013). Host-

induced gene silencing of cytochrome P450 lanosterol C14α-demethylase–

encoding genes confers strong resistance to Fusarium species. *Proceedings*

of the National Academy of Sciences, *110*(48), 19324–19329.

https://doi.org/10.1073/pnas.1306373110

Kramer, M. F. (2011). STEM-LOOP RT-qPCR for miRNAS. *Current Protocols in*

Molecular Biology / Edited by Frederick M. Ausubel ... [et Al.], CHAPTER,

Unit15.10. https://doi.org/10.1002/0471142727.mb1510s95

Lincoln, C., Britton, J. H., & Estelle, M. (1990). Growth and development of the

axr1 mutants of Arabidopsis. *The Plant Cell*, *2*(11), 1071–1080.

Liu, S., Jaouannet, M., Dempsey, D. A., Imani, J., Coustau, C., & Kogel, K.-H. (2020).

RNA-based technologies for insect control in plant production.

Biotechnology Advances, *39*, 107463.

https://doi.org/10.1016/j.biotechadv.2019.107463

Llave, C., Xie, Z., Kasschau, K. D., & Carrington, J. C. (2002). Cleavage of Scarecrow-

like mRNA Targets Directed by a Class of Arabidopsis miRNA. *Science*,

297(5589), 2053–2056. https://doi.org/10.1126/science.1076311

Mahanty, B., Mishra, R., & Joshi, R. K. (2023). Cross-kingdom small RNA

communication between plants and fungal phytopathogens-recent

updates and prospects for future agriculture. *RNA Biology*, *20*(1), 109–119.

https://doi.org/10.1080/15476286.2023.2195731

Mallory, A. C., Reinhart, B. J., Jones-Rhoades, M. W., Tang, G., Zamore, P. D.,

Barton, M. K., & Bartel, D. P. (2004). MicroRNA control of PHABULOSA in

leaf development: Importance of pairing to the microRNA 5' region. *The*

EMBO Journal, *23*(16), 3356–3364.

https://doi.org/10.1038/sj.emboj.7600340

Meyers, B. C., & Axtell, M. J. (2019). MicroRNAs in Plants: Key Findings from the

Early Years. *The Plant Cell*, *31*(6), 1206–1207.

https://doi.org/10.1105/tpc.19.00310

Mi, S., Cai, T., Hu, Y., Chen, Y., Hodges, E., Ni, F., Wu, L., Li, S., Zhou, H., Long, C.,

Chen, S., Hannon, G. J., & Qi, Y. (2008). Sorting of Small RNAs into

Arabidopsis Argonaute Complexes Is Directed by the 5' Terminal

Nucleotide. *Cell*, *133*(1), 116–127. https://doi.org/10.1016/j.cell.2008.02.034

Montgomery, T. A., Howell, M. D., Cuperus, J. T., Li, D., Hansen, J. E., Alexander, A.

L., Chapman, E. J., Fahlgren, N., Allen, E., & Carrington, J. C. (2008).

Specificity of ARGONAUTE7-miR390 interaction and dual functionality in

TAS3 trans-acting siRNA formation. *Cell*, *133*(1), 128–141.

https://doi.org/10.1016/j.cell.2008.02.033

Nasfi, S., & Kogel, K.-H. (2022). Packaged or unpackaged: Appearance and

transport of extracellular noncoding RNAs in the plant apoplast. *ExRNA; Vol*

4 (May 31, 2022): ExRNA. https://exrna.amegroups.com/article/view/64855

Nowara, D., Gay, A., Lacomme, C., Shaw, J., Ridout, C., Douchkov, D., Hensel, G.,

Kumlehn, J., & Schweizer, P. (2010). HIGS: Host-Induced Gene Silencing in

the Obligate Biotrophic Fungal Pathogen *Blumeria graminis*. *The Plant Cell*,

22(9), 3130–3141. https://doi.org/10.1105/tpc.110.077040

Osborne, R., Rehneke, L., Lehmann, S., Roberts, J., Altmann, M., Altmann, S.,

Zhang, Y., Köpff, E., Dominguez-Ferreras, A., Okechukwu, E., Sergaki, C.,

Rich-Griffin, C., Ntoukakis, V., Eichmann, R., Shan, W., Falter-Braun, P., &

Schäfer, P. (2023). Symbiont-host interactome mapping reveals effector-

targeted modulation of hormone networks and activation of growth

promotion. *Nature Communications*, *14*(1), Article 1.

https://doi.org/10.1038/s41467-023-39885-5

Ossowski, S., Schwab, R., & Weigel, D. (2008). Gene silencing in plants using

artificial microRNAs and other small RNAs. *The Plant Journal*, *53*(4), 674–

690. https://doi.org/10.1111/j.1365-313X.2007.03328.x

Qiang, X., Weiss, M., Kogel, K.-H., & Schäfer, P. (2012). Piriformospora indica—A

mutualistic basidiomycete with an exceptionally large plant host range.

Molecular Plant Pathology, *13*(5), 508–518. https://doi.org/10.1111/j.1364-

3703.2011.00764.x

Qiao, S. A., Gao, Z., & Roth, R. (2023). A perspective on cross-kingdom RNA

interference in mutualistic symbioses. *New Phytologist*, *240*(1), 68–79.

https://doi.org/10.1111/nph.19122

Reinhart, B. J., Weinstein, E. G., Rhoades, M. W., Bartel, B., & Bartel, D. P. (2002).

MicroRNAs in plants. *Genes & Development*, *16*(13), 1616–1626.

https://doi.org/10.1101/gad.1004402

Ren, B., Wang, X., Duan, J., & Ma, J. (2019). Rhizobial tRNA-derived small RNAs are

signal molecules regulating plant nodulation. *Science (New York, N.Y.)*,

365(6456), 919–922. https://doi.org/10.1126/science.aav8907

Rhoades, M. W., Reinhart, B. J., Lim, L. P., Burge, C. B., Bartel, B., & Bartel, D. P.

(2002). Prediction of Plant MicroRNA Targets. *Cell*, *110*(4), 513–520.

https://doi.org/10.1016/S0092-8674(02)00863-2

Rutter, B. D., & Innes, R. W. (2018). Extracellular vesicles as key mediators of

plant–microbe interactions. *Current Opinion in Plant Biology*, *44*, 16–22.

https://doi.org/10.1016/j.pbi.2018.01.008

Sang, H., & Kim, J.-I. (2020). Advanced strategies to control plant pathogenic fungi

by host-induced gene silencing (HIGS) and spray-induced gene silencing

(SIGS). *Plant Biotechnology Reports*, *14*(1), 1–8.

https://doi.org/10.1007/s11816-019-00588-3

Šečić, E., Zanini, S., Wibberg, D., Jelonek, L., Busche, T., Kalinowski, J., Nasfi, S.,

Thielmann, J., Imani, J., Steinbrenner, J., & Kogel, K.-H. (2021). A novel plant-

fungal association reveals fundamental sRNA and gene expression

reprogramming at the onset of symbiosis. *BMC Biology*, *19*(1), 171.

https://doi.org/10.1186/s12915-021-01104-2

Silvestri, A., Fiorilli, V., Miozzi, L., Accotto, G. P., Turina, M., & Lanfranco, L. (2019).

In silico analysis of fungal small RNA accumulation reveals putative plant

mRNA targets in the symbiosis between an arbuscular mycorrhizal fungus

and its host plant. *BMC Genomics*, *20*, 169. https://doi.org/10.1186/s12864-019-5561-0

Takeda, A., Iwasaki, S., Watanabe, T., Utsumi, M., & Watanabe, Y. (2008). The Mechanism Selecting the Guide Strand from Small RNA Duplexes is Different Among Argonaute Proteins. *Plant and Cell Physiology*, *49*(4), 493–500. https://doi.org/10.1093/pcp/pcn043

Tyurin, A. A., Suhorukova, A. V., Kabardaeva, K. V., & Goldenkova-Pavlova, I. V. (2020). Transient Gene Expression is an Effective Experimental Tool for the Research into the Fine Mechanisms of Plant Gene Function: Advantages, Limitations, and Solutions. *Plants*, *9*(9), 1187. https://doi.org/10.3390/plants9091187

Ueno, D., Yamasaki, S., & Kato, K. (2022). Methods for detecting RNA degradation intermediates in plants. *Plant Science*, *318*, 111241. https://doi.org/10.1016/j.plantsci.2022.111241

Varkonyi-Gasic, E., & Hellens, R. P. (2011). Quantitative stem-loop RT-PCR for detection of microRNAs. *Methods in Molecular Biology (Clifton, N.J.)*, *744*, 145–157. https://doi.org/10.1007/978-1-61779-123-9_10

Verma, N., Narayan, O. P., Prasad, D., Jogawat, A., Panwar, S. L., Dua, M., & Johri, A. K. (2022). Functional characterization of a high-affinity iron transporter (PiFTR) from the endophytic fungus Piriformospora indica and its role in plant growth and development. *Environmental Microbiology*, *24*(2), 689–706. https://doi.org/10.1111/1462-2920.15659

Verma, S., Varma, A., Rexer, K.-H., Hassel, A., Kost, G., Sarbhoy, A., Bisen, P.,

Bütehorn, B., & Franken, P. (1998). Piriformospora indica, gen. Et sp. Nov.,

a new root-colonizing fungus. *Mycologia*, *90*(5), 896–903.

https://doi.org/10.1080/00275514.1998.12026983

Wang, M., Thomas, N., & Jin, H. (2017). Cross-kingdom RNA trafficking and

environmental RNAi for powerful innovative pre- and post-harvest plant

protection. *Current Opinion in Plant Biology*, *38*, 133–141.

https://doi.org/10.1016/j.pbi.2017.05.003

Wang, M., Weiberg, A., Lin, F.-M., Thomma, B. P. H. J., Huang, H.-D., & Jin, H.

(2016). Bidirectional cross-kingdom RNAi and fungal uptake of external

RNAs confer plant protection. *Nature Plants*, *2*(10), Article 10.

https://doi.org/10.1038/nplants.2016.151

Weiberg, A., Wang, M., Lin, F.-M., Zhao, H., Zhang, Z., Kaloshian, I., Huang, H.-D., &

Jin, H. (2013). Fungal small RNAs suppress plant immunity by hijacking host

RNA interference pathways. *Science (New York, N.Y.)*, *342*(6154), 118–123.

https://doi.org/10.1126/science.1239705

Weiß, M., Waller, F., Zuccaro, A., & Selosse, M.-A. (2016). Sebacinales – one

thousand and one interactions with land plants. *New Phytologist*, *211*(1),

20–40. https://doi.org/10.1111/nph.13977

Werner, B. T., Koch, A., Šečić, E., Engelhardt, J., Jelonek, L., Steinbrenner, J., &

Kogel, K.-H. (2021). Fusarium graminearum DICER-like-dependent sRNAs

are required for the suppression of host immune genes and full virulence.

PLOS ONE, *16*(8), e0252365. https://doi.org/10.1371/journal.pone.0252365

Wong-Bajracharya, J., Singan, V. R., Monti, R., Plett, K. L., Ng, V., Grigoriev, I. V.,

Martin, F. M., Anderson, I. C., & Plett, J. M. (2022). The ectomycorrhizal

fungus Pisolithus microcarpus encodes a microRNA involved in cross-

kingdom gene silencing during symbiosis. *Proceedings of the National*

Academy of Sciences of the United States of America, *119*(3), e2103527119.

https://doi.org/10.1073/pnas.2103527119

Wu, F.-H., Shen, S.-C., Lee, L.-Y., Lee, S.-H., Chan, M.-T., & Lin, C.-S. (2009). Tape-

Arabidopsis Sandwich—A simpler Arabidopsis protoplast isolation

method. *Plant Methods*, *5*(1), 16. https://doi.org/10.1186/1746-4811-5-16

Xu, L., Wu, C., Oelmüller, R., & Zhang, W. (2018). Role of Phytohormones in

Piriformospora indica-Induced Growth Promotion and Stress Tolerance in

Plants: More Questions Than Answers. *Frontiers in Microbiology*, *9*.

https://www.frontiersin.org/articles/10.3389/fmicb.2018.01646

Yoo, S.-D., Cho, Y.-H., & Sheen, J. (2007). Arabidopsis mesophyll protoplasts: A

versatile cell system for transient gene expression analysis. *Nature*

Protocols, *2*(7), 1565–1572. https://doi.org/10.1038/nprot.2007.199

Zand Karimi, H., & Innes, R. W. (2022). Molecular mechanisms underlying host-

induced gene silencing. *The Plant Cell*, *34*(9), 3183–3199.

https://doi.org/10.1093/plcell/koac165

Zhan, J., & Meyers, B. C. (2023). Plant Small RNAs: Their Biogenesis, Regulatory

Roles, and Functions. *Annual Review of Plant Biology*, *74*(1), 21–51.

https://doi.org/10.1146/annurev-arplant-070122-035226

Zhang, T., Zhao, Y.-L., Zhao, J.-H., Wang, S., Jin, Y., Chen, Z.-Q., Fang, Y.-Y., Hua, C.-

L., Ding, S.-W., & Guo, H.-S. (2016). Cotton plants export microRNAs to

inhibit virulence gene expression in a fungal pathogen. *Nature Plants*, *2*(10),

16153. https://doi.org/10.1038/nplants.2016.153

Zuccaro, A., Lahrmann, U., Güldener, U., Langen, G., Pfiffi, S., Biedenkopf, D.,

Wong, P., Samans, B., Grimm, C., Basiewicz, M., Murat, C., Martin, F., &

Kogel, K.-H. (2011). Endophytic Life Strategies Decoded by Genome and

Transcriptome Analyses of the Mutualistic Root Symbiont Piriformospora

indica. *PLOS Pathogens*, *7*(10), e1002290.

https://doi.org/10.1371/journal.ppat.1002290