Short title: Black-veined white extirpation

The last days of *Aporia crataegi* (L.) in Britain: evaluating genomic erosion in

an extirpated butterfly

Rebecca Whitla[1], Korneel Hens[1,2], James Hogan[3], Geoff Martin[4], Casper Breuker[1,2], Timothy G.

Shreeve[1], Saad Arif[1,2,*]

[1] Department of Biological and Medical Sciences, Oxford Brookes University, Oxford, OX3 0BP, UK

[2] Centre for Functional Genomics, Oxford Brookes University Oxford, OX3 0BP, UK

[3] Oxford University Museum of Natural History, Parks Road, Oxford, OX1 3PW, UK

[4] Insects Division, Natural History Museum, London, SW7 5BD, UK

* Author for correspondence

Keywords: *Aporia crataegi*, inbreeding, genetic load, historical DNA, genomic erosion, genetic diversity

Correspondence:

Saad Arif

Centre for Functional Genomics, Oxford Brookes University, OX3 0BP Oxford, United Kingdom

Department of Biological and Medical Sciences, Oxford Brookes University, OX3 0BP, United Kingdom

Tel: +44 (0)1865 483437

Email: sarif@brookes.ac.uk

Abstract

Current rates of habitat degradation and climate change are causing unprecedented declines in global biodiversity. Studies on vertebrates highlight how conservation genomics can be an effective tool to identify and manage threatened populations, but it is unclear how vertebrate derived metrics of genomic erosion translate to invertebrate species, with their markedly different population sizes and life histories. The Black-veined White butterfly (*Aporia crataegi)* was extirpated from Britain in the 1920s. Here, we sequenced historical DNA from 17 museum specimens collected between 1854-1924 to reconstruct demography and compare levels of genomic erosion between extirpated British and extant European mainland populations. We contrast these results using modern samples of the Common Blue butterfly (*Polyommatus icarus*); a species with relatively stable demographic trends in Great Britain. We provide evidence for bottlenecks in both these species around the period of post-glacial colonisation of the British Isles. Our results reveal different demographic histories and N_e for both species, consistent with their fates in Britain, likely driven by differences in life history, ecology, genome lengths and body size. Despite a difference, by an order of magnitude, in historical effective population sizes (N_e), reduction in genome-wide heterozygosity in *A. crataegi* was comparable to that in *P. icarus*. Symptomatic of *A. crataegi*'s disappearance were marked increases in runs-of-homozygosity (RoH), potentially indicative of recent inbreeding, and accumulation of putatively mildly and weakly deleterious variants. Our results support the idea that metrics of inbreeding and accumulation of deleterious mutations could be more informative markers of population or species in decline.

Introduction

Global extinction rates have increased over the last 500 years (Barnosky *et al.*, 2011, Ceballos *et al.*, 2015, Cowie *et al.*, 2022), accelerating over the last 100 years with *circa* 25% of species recently assessed by the IUCN are now threatened with extinction (IPBES, 2019). In Great Britain, *c.* 16.1% of assessed flora and fauna species (10,008) are classified as threatened, and 146 species have gone extinct (with 52 species 'likely extinct') since 1500 (Burns *et al.*, 2023). Invertebrates count for 97% of all animals globally and are a major component of terrestrial biodiversity with important roles in ecosystem functioning. Assessments for many invertebrates are data deficient, precluding accurate determination of their status (Karam-Gemael *et al.*, 2020). Reasons for insufficient data include underfunding for less charismatic and economically important groups (Cardoso and Leather, 2019) and difficulties in sampling and identification (Wagner *et al.*, 2021).

Among insects, butterflies are one of the best-studied groups (for example, Warren, 1992; Thomas, 2005; Dincă *et al.*, 2015; Middleton-Welling *et al.*, 2020; Warren *et al.*, 2021), can be used as bioindicators of environmental and climate change (Dennis *et al.*, 2003; Hill *et al.*, 2021) and serve as an important invertebrate model for communicating science to society (Preston *et al.*, 2021). Since the 1970s, standardised and systematic monitoring of butterflies across Europe has provided a comprehensive and continuous record of changes in biodiversity. Compared to most of continental Europe, a larger proportion of UK butterflies are now either extirpated (4 of 62 species) or considered as critically endangered or vulnerable (19/62) (Fox *et al.*, 2022).

By quantifying genomic erosion, conservation genomics can be a valuable tool in conservation, helping identify vulnerable populations and informing translocations and/or breeding programmes (Díez-del-Molino *et al.,* 2018; Bosse and van Loon, 2022). Metrics for genomic erosion, associated with reductions of fitness (Bosse and van Loon, 2022), aim to estimate genetic diversity, effective population size (N_e), levels of inbreeding and genetic load. All have a strong theoretical basis to be correlated with and potentially exacerbate extinction risk in small and vulnerable populations (Gomulkiewicz and Holt, 1995; Lynch *et al.*, 1995a; Charlesworth and Willis, 2009). Negative impact of inbreeding depression in small populations is well established (Saccheri *et al.*, 1998; Mattila *et al.*,

2012; Neaves *et al.*, 2015; Rowe *et al.,* 2017) but the utility of such metrics of, particularly those

relying solely on genome-wide, putatively neutral genetic markers, in predicting extinction risk is

currently under debate (Kardos *et al.*, 2016; Teixeira and Huber, 2021; Dussex *et al.*, 2023). A better

understanding is emerging from the accumulation of genomic data of small and threatened

populations using whole genomes to estimate multiple metrics of genomic erosion simultaneously

(DeWoody *et al.*, 2021; Bosse and van Loon, 2022; Dussex *et al.*, 2023), but is largely limited to

examples from terrestrial vertebrates and our understanding of genomic erosion in insects is far more

limited (Webster *et al.*, 2023; but see de-Dios *et al.*, 2023; Bortoluzzi *et al.*, 2023). Insects can have

markedly higher population sizes and different life histories, compared to vertebrates, and it is unclear

how this would impact readouts of genetic diversity, effective population size (N_e), levels of inbreeding

and genetic load in threatened insect populations. Comparative analyses of genomic erosion between

endangered and stable or expanding insect populations, could prove extremely useful for detecting

insect declines or identifying vulnerable species or populations from small or limited samples and for

determining effective metrics to identify these in insects.

Here, we examine trends of genomic erosion in *Aporia crataegi* (L.) (Black-veined White butterfly),

which was extirpated from the British Isles *c.* 1925 (Pratt, 1983), contrasting these with populations in

continental Europe. We also provide a comparison between British and continental European

populations of *Polyommatus icarus* (Rott.) (Common Blue butterfly); a species that has exhibited little

change in long term population trends (https://www.gov.uk/government/statistics/butterflies-in-the-

wider-countryside-uk). *A. crataegi* is widespread throughout much of Europe, Asia and north Africa. It

has recently been lost from the Netherlands, Czechia (van Swaay *et al.*, 2010), and South Korea (Kim

et al., 2015), and is declining in other European countries (van Swaay *et al.*, 2010). Prior to extinction

in Britain, it was restricted to southern and central England and Wales and there is evidence that

populations were in decline from the mid to late 19[th] century (Dale, 1887; South 1906; Pratt, 1983)

and was primarily only found in southeast England by the end of the 19[th] Century. Its decline in

Europe is typically attributed to habitat loss, but this does not seem to be the case for its extirpation in

Britain (MacLachlan, 1893) with pesticide use, parasites, microbial or fungal pathogens and predation

by small birds all being suggested as potential reasons instead (reviewed in Pratt, 1983). Numerous

"accidental" releases, recorded as early as the 19[th] century (Pratt, 1983), from butterfly enthusiasts to

reintroduce the species from continental European stock have been made, but none have been

successful for establishment of the species.

To examine whether *A. crataegi* in Britain exhibited any symptoms of genomic erosion, we sequenced

historical specimens of *A. crataegi* from Britain and continental Europe dating from 1854-1924. We

specifically addressed whether the British *A. crataegi* were distinct from continental Europe based on

genetic structuring and demographic history prior to 10 Kya. We examine individual heterozygosity,

runs-of-homozygosity ((RoH), a genomic measure of inbreeding), and genetic load to explain the

vulnerability of the British Isles populations prior to extinction and contextualise these finding by

comparison with corresponding analyses of the relatively stable British and continental European *P.*

icarus.

Materials and Methods

Historical **A. crataegi** *Specimen Selection*

Seventeen *A. crataegi* specimens from Britain (GB), Belgium, and France, (Fig. 1A, Table S1) were

sampled from collections at Oxford University Museum of Natural History (OUMNH), Oxford, England

and the Natural History Museum (NHM), London, England. Specimen dates ranged from 1854-1924.

For comparison, we only used museum samples of continental populations to avoid any potential bias

stemming from comparing degraded historical DNA with modern DNA. Up to two legs per specimen

were taken from pinned samples for DNA extraction, although in some cases only a single leg was

available.

Extraction, Library Preparation and Sequencing of Historical DNA

DNA extractions were undertaken on a laboratory bench used only for historical DNA (hDNA)

extractions to prevent contamination from modern DNA. Extraction blanks were prepared alongside

samples to monitor potential contamination. Legs of specimens were fully macerated using a

Tissuelyser II bead mill (Qiagen) and extracted using a QIAamp DNA Micro kit (Qiagen) using the

standard protocol with a final elution volume of 20μL. Total DNA yield was measured using a Qubit 4

Fluorometer (Life Technologies) with a Qubit dsDNA HS Assay Kit. Eluted DNA was stored in

Eppendorf DNA LoBind Tubes at -20°C until library preparation.

Genomic library preparation of hDNA was performed using the NEBNext Ultra II DNA Library Prep Kit

for Illumina. A CUT&RUN (Liu, 2019) protocol was used, modified for use with low yield samples and

samples with short fragment sizes. NEBNext® Dual Index Multiplex Oligos for Illumina (Set 1) Primers

were used to construct multiplexed libraries. SPRIselect beads (Beckman Coulter) were used for size

exclusion post primer ligation at 1.75X or 2.2X concentration.

The number of PCR cycles used for amplification was dependent on the input DNA quantity; 12

cycles for samples < 5 ng and 10 cycles for samples ε 5 ng. Libraries were then purified with 1.2X

SPRIselect beads to remove adapter dimer. Library enrichment was performed using either a Q5U®

Hot Start High-Fidelity DNA Polymerase (New England Biolabs) or the NEBNext Ultra II Q5DNA

polymerase (Table S1). We did not attempt to repair the hDNA template prior to library preparation.

However, the NEBNext Ultra II DNA Library Prep Kit makes use of USER enzyme during the adapter

ligation step. This excises Uracil bases from the ligated adapters and is required when using

NEBNext adapters, but in case of hDNA template with post-mortem deamination damage, the

enzyme will also excise Uracil bases on the 5' end of the hDNA template. With subsequent paired-

end Illumina sequencing this results in partial repair with 5' ends exhibiting little to no signs of

historical damage whereas the damage profile on 3' ends reflects typical post-mortem DNA damage.

Quality control for the libraries was performed using a Qubit 4 Fluorometer (Life Technologies) and an

Agilent Bioanalyser, using the Agilent High Sensitivity DNA analysis Chip Kit. Sample libraries were

pooled equimolarly, with extraction blanks pooled at a 1:10 molar ratio in proportion to sample

libraries. Sequencing was performed using the NovaSeq Illumina platform using a 150 base pair (bp)

paired-end (150PE) strategy at Novogene (Cambridge).

***Sampling Modern specimens of* P. icarus**

Four individuals of *P. icarus* were collected from across the British Isles and an additional three from

Averyon, France between 2017-2018 (Table S2, Fig. 1A) and DNA extraction and further details of

these samples is as described in Arif *et al.* (2021). DNA was normalised to 10 ng/ul and library

preparation and sequencing, to a theoretical coverage of 15-50X (Table S2), was performed using the

NovaSeq Illumina platform with a 150PE strategy at Novogene (Cambridge).

Data Processing

The GenErode pipeline v0.4.2 (Kutschera *et al.*, 2022), suitable for processing both modern and

historical short-read data, was used to trim and process all modern and historical sequence data

along with highly contiguous reference genomes for *A. crataegi* (Ebdon *et al.*, 2022) or *P. icarus*

(Lohse *et al.*, 2023). Both these reference genome assemblies are high quality, produced from single

male specimens, with an assembled Z chromosome (Lepidopteran males are commonly the

homogametic sex). The *A. crataegi* assembly is ~230 Megabases (Mb) long and is scaffolded into 26

pseudochromosomes and a Z chromosome. The *P. icarus* assembly is ~512 Mb scaffolded into 23

pseudochromosomes and a Z.

FastQC v0.11.9 (Andrews *et al.*, 2012) and fastp v0.20.1 (Chen *et al.*, 2018) were used to trim

adapters and merge paired sequences (for historical samples). Trimmed reads were aligned to

reference genomes using BWA aln for hDNA and BWA mem for modern data (Li and Durbin, 2009)

with default settings in the GenErode pipeline. Post-mapping, any samples sequenced across multiple

lanes were merged and PCR duplicates removed. MapDamage v2.0 (Jónsson *et al.*, 2013) was used

to estimate damage in historical samples and rescale base quality scores. Next, GATK

IndelRealigner, (Van der Auwera and O'Connor, 2020) was used to remap any reads around indels to

improve mapping accuracy. For the modern, *P. icarus* data, all resulting BAM files were subsampled

to 10X to mitigate the influence of varying coverage on heterozygous calls. These BAM files were

processed further to call variants and reconstruct historical demography. For the latter, we used full

coverage data for the modern *P. icarus* samples.

Variants calls were made on a per-sample basis using Bcftools mpileup (Danecek *et al*., 2021) with a

minimum depth threshold of 1/3 of average coverage (with an absolute minimum of 3X) and a

maximum of ten times the average coverage while minimum mapping quality and minimum base

quality were both set to 30 (-Q 30 -q 30). Additional filters as a part of the GenErode pipeline, included

an allelic imbalance filter for heterozygous sites, removal of variants in repetitive regions of the

genome, and removal of variants in CpG regions. A final VCF was generated that consisted of only

biallelic single nucleotide polymorphisms (SNPS) and genotypes with no more than 10% missing data

across all samples (historical *A. crataegi* samples, containing 305,394 SNPs) or no missing data

(modern *P .icarus* data, containing 5,656,704 SNPs). SNPs from these VCFs were used for

downstream analyses, unless otherwise stated. Our configuration files for the GenErode pipeline are

available at https://github.com/rmwhitla/BVWpaper.

Population Structure

Population structure of both the historical *A. crataegi* and modern *P. icarus* samples were compared

using principal component analysis (PCA) based on filtered genotypes in VCF format. Before PCA,

any SNPs on the Z chromosome were filtered out. Furthermore, we pruned the remaining SNP sets

for linkage disequilibrium over windows of 10 kilobases with a step size of 10 bases and an r^2 of 0.1

using Plink 1.9 (Purcell *et al.,* 2007) resulting in a total of 14,930 (*A. crataegi*) or 197,182 (*P. icarus*)

SNPs. PCA was performed using Plink v1.9 with a minor allele frequency (MAF) threshold of 10%

(historical *A. crataegi)* or 15% (modern *P. icarus*).

Given the low and variable coverage (2-13X) of the historical *A. crataegi* samples, we also performed

PCA based on genotype likelihoods, which accounts for statistical uncertainty in called genotypes

arising from low coverage data (Korneliussen *et al*., 2014). Genotype likelihoods were calculated

using ANGSD (Korneliussen *et al*., 2014) from BAM files produced as part of the GenErode pipeline

with a minimum mapping quality at 30 and minimum base quality at 20 (-minMapQ 30 -minQ 20)

filtering only for the same sites used with the PCA for hard genotype calls. Pcangsd (Meisner and

Albrechtsen, 2018) was used for the PCA of genotype likelihoods with a minimum allele frequency of

10% (--minMaf 0.1).

Historical Demography Reconstruction

We used the Pairwise Sequentially Markovian Coalescent model as implemented in PSMC v0.6.5 (Li

and Durbin, 2011) to determine if British populations of *A. crataegi* and *P. icarus* exhibit similar trends

of historical change in effective population size (N_e) following permanent colonization of the British

Isles after the Younger Dryas *c.*11.5 Kya and permanent flooding of the Channel *c.* 9 Kya (Dennis,

1992). As input for PSMC, we generated a consensus fastq file from BAM files, using only samples

with average coverage greater than 5X. Consensus fastq files were generated with Bcftools pileup

and vcfutils.pl, filtering positions with minimum depth less than 5X or maximum depth greater than

100X in *A. crataegi* and with minimum depth less than 5X or maximum depth greater than 500X in *P.*

icarus. Additionally, any positions in repeat masked regions or CpG regions (historical samples only)

or Z chromosomes were also removed. PSMC was run with: number of iterations -N = 25, maximum

coalescent time -t = 8 for *A. crataegi* or t = 9 fpr *P. icarus* and the atomic time interval (-p) at 66 (for *A.*

crataegi: 4+25*2+3+9) and 66 (for *P. icarus*: 27*2+4+8). The atomic time interval and maximum

coalescent time was chosen to avoid overfitting by ensuring that at least 10 recombination events

occurred over each interval after the 20[th] iteration of PSMC (Li and Durbin, 2011). To account for low

coverage and stochastic loss of heterozygosity, the initial theta/rho ratio parameter for each sample

was calculated based on false negative rates (FNR) corresponding to the coverage of that sample.

FNR for different coverages were determined based on the recommendation of Li and Durbin (2011)

(https://github.com/lh3/psmc), and as implemented by Sarabia *et al.* (2020). To obtain FNR estimated

for *A. crataegi*, we used a subsampled version of a publicly available high coverage (23X) modern *A.*

crataegi (accession: SRR7948941) specimen from Japan that was sequentially subsampled to 5X,

8X, 10X, 15X and 20X. For FNR correction for low coverage *P. icarus* samples, we used the same

subsampling strategy as for *A. crataegi* using one of our high coverage samples (DGCM100,

coverage: >50X). The initial theta/rho (-r) parameter for PSMC was then set to 5/(1-FNR). To visualise

the variability in our PSMC estimates, we ran 100 bootstraps for each sample over 5 Mb segments

using the same initial parameters as the original run for each sample.

To scale population size and times resulting from the PSMC analysis, we used a direct estimate of

spontaneous mutation rate of 2.9 x10^{-9} from *Heliconius melpomene* (Keightley *et al.,* 2015). For *A.*

crataegi, which is univoltine in the geographic range our samples came from (Solonkin *et al.,* 2021),

we used 1 generation per year. For *P. icarus*, we used 2 generations per year, although the number

of broods are variable in this species (one in Scotland but two or three in Southern England and parts

of Europe). We tested the sensitivity of the PSMC curves for *P. icarus* by changing the generation

time to 1 and 3 broods per year.

Genetic Diversity

Genome-wide heterozygosity was estimated using ANGSD, which utilises a genotype likelihood

approach that is more suitable for low coverage data (Korneliussen *et al.*, 2014). We calculated

genotype likelihoods as described above (PCA using genotype likelihoods), except in this case no LD

pruning was performed. To estimate global heterozygosity, we calculated the unfolded site frequency

spectrum (SFS) using the realSFS function in ANGSD.

Inbreeding and Contemporary N_e

We estimated individual inbreeding coefficients based on the detection of the number and size of

Runs of Homozygosity (RoH). RoH were calculated using Plink v1.9 (Purcell *et al.,* 2007). We used a

sliding window size of 100 SNPs (homozyg-window-snp 100). The maximum number of

heterozygotes per window was 2 (homozyg-window-het 2). Minimum SNP count for each RoH was 25

(homozyg-snp 25) and the minimum size was set to 50kb (homozyg-kb 50). We allowed 10 missing

sites per window (intermediate filtering - homozyg-window-missing 10). For the maximum number of

heterozygote sites within a RoH, we allowed 3 (homozyg-het3). Using these results, F_{RoH}, the

inbreeding coefficient, was determined using the overall proportion of the genome in RoH. Results

from Plink were summarised in R v.4.3.0 (R Core Team, 2023) using the package DetectRuns

(Biscarini *et al.,* 2018). F_{ROH} was estimated for RoH $\geq$ 100 Kilobases (Kb), RoH $\geq$ 500 Kb or RoH $\geq$ 1

Megabase (Mb) for samples with average coverage $\geq$ 5X. RoH $\geq$ 100 Kb and less than 1 Mb may

result from background inbreeding due to drift whereas RoH ϵ 1 Mb result from recent

consanguineous mating (Ceballos *et al.*, 2018).

Inference of N_e from PSMC is limited over the last 10 Kya but the length and distributions of RoHs can

be informative of more recent N_e. To compare our historical estimates of N_e (~10 Kya) estimated from

PSMC with those observed from RoH we calculated the probability of observing at least one RoH = 1

Mb, in a genome of size 230 Mb (*A. crataegi*) or 512 Mb (*P. icarus)*. Assuming a neutrally evolving

Wright-Fisher population, the probability of a randomly chosen region of length 10^6 bases being a

ROH can be computed as the probability that pairwise coalescence happens before either sequence

recombines (Hudson, 1990):

$$P(RoH=1Mb) \approx \frac{\frac{1}{2N_e}}{\left(\frac{1}{2N_e} + 2*10^6 r\right)}$$

Where N_e was based on the PSMC estimate at ~10 Kya and *r* represents the recombination rate and

was set to 20 times 2.9 x10^{-9} (the spontaneous mutation rate). This choice of recombination rate is

empirically plausible for butterflies which can be 20-fold higher than the mutation rate. (e.g. Jiggins *et*

al., 2005; Palahí i Torres *et al.,* 2023)

Genetic Load Estimation

To estimate the number of deleterious mutations or the genetic load we used snpEff (Cingolani *et al.,*

2012) to annotate genetic variants of high impact (likely highly disruptive to protein, e.g., loss of

function (LoF) variants), moderate impact (e.g. missense variants) and low impact (synonymous, likely

harmless but may be weakly deleterious under strong codon bias), as well as modifier (non-coding)

variants. We focused on annotated variants within coding regions (LoF, missense or synonymous).

snpEFF only predicts the potential phenotypic impact of the variants, and we lack any prior knowledge

of the selection or dominance coefficients for these variants.

For the analysis of genetic load, we only used samples with ϵ 5X average depth and no missing data

across samples. This resulted in 179,135 SNPs for historical *A. crataegi* and 5,656,704 SNPs for *P.*

icarus. Gene annotations (as GTF files) for both species were retrieved from Ensembl's Darwin Tree

of Life data portal (https://projects.ensembl.org/darwin-tree-of-life/). Both annotations were generated

by the Ensembl Gene Annotation System (Aken *et al.*, 2017) optimised for non-vertebrates. The

annotation for *A. crataegi* (Ebdon *et al.,* 2022) is based on a single male from Catalunya, Spain and

the annotation for *P. icarus* (Lohse, 2023) is based on a single male from Scotland.

We also determined the relative excess in the frequency of shared derived alleles (ignoring the impact

of *de novo* mutations) across the LoF, missense and synonymous categories in both species. We

used an approach previously described by Xue *et al.* (2015) as implemented by van der Valk *et al.*

(2019). For each variant category we calculate a ratio R_{xy} that is a measure of shared derived alleles

with a higher frequency in population X (British) compared to population Y (European). An $R_{xy} > 1$

implies a relative increase in frequency in population and X compared to Y, whereas < 1 implies a

relative decrease in X relative to Y. To estimate confidence intervals for our R_{xy} ratios we used a

jackknife approach dropping a single chromosome at a time.

Statistical Analysis

For all statistical analysis between historical specimens of *A. crataegi* we used a linear model with

average coverage as a covariate to account for variation in coverage between specimens. Marginal

means of all estimates for each group were estimated using the R package emmeans (Lenth, 2023).

For *P. icarus*, statistical comparisons were made using Welch's two sample t-tests. All statistical

comparisons were performed in R v.4.3.0 (R Core Team, 2023).

Results

Resequencing of hDNA

The final sequencing depth of the 17 historical *A. crataegi* specimens varied between 2X-13X, with

higher mean coverage of European samples (10.13X) compared to British specimens (6.03X) and

sequencing depth varied from 2X-13X (Table S1). The average proportion of the genome sequenced

to a depth of 5X or greater was 50.77% (Table S1).

Post-mortem degradation was apparent in the 3' ends of our historical reads (Fig. S1) as an increase

in G>A transversions. The 5' ends of our fragments show little to no sign of historical damage due to

the application of the USER enzyme during library preparation. Mean fragment length of sequencing

reads, after trimming and merging reads, was 49bp with average fragment length per sample ranging

from 44-65bp (Table S1). There was a significant positive correlation between mean fragment length

and age of sample (Fig. S2) with fragment lengths increasing closer to more recent times (Mullin *et*

al., 2023).

In our extraction blanks (BL1 and BL2), 0.49% and 0.43% of reads mapped to the *A. crataegi* genome

respectively. In the blanks, 0.2% of bases had 1X coverage, and 0% of bases had higher coverage.

The average read lengths in the two blanks were 91 and 97. Overall, analysis of the mapping profile

of the blanks suggests no evidence for cross contamination during DNA extraction and library

preparation of the historical samples.

Spatial Genetic Structuring between Great British and European Mainland Butterflies

A PCA using LD-pruned SNPs of all historical *A. crataegi* specimens identifies spatial genetic

structuring across historical samples regardless of collection date (Fig. 1B). A PCA based on

genotype likelihoods (Fig. S3), undertaken because of low to moderate coverage of the historic

specimens, yielded visually similar results to the one based on hard genotype calls (Fig. 1B).

PC1 explained 10.8% of the variation, while PC2 explained 8.6% of variation. Spatial genetic

structuring is apparent along PC1 with European samples tightly clustering on one end with negative

values for PC1, whereas points with greater values of PC1 represent two specimens from range

margins (Wales; see Fig. S3 for a labelled PCA- but note the axes are reversed) but also include a

sample with unknown origins (K05L) and one specimen from Kent (NHM026). In contrast to the

geographical cline apparent along PC1, there is no clear cline present along PC2 but the points with

the most extreme values represent the most recent specimens from Kent (NHM928 and NHM019). *P.*

icarus exhibit similar structuring along PC1 with the southern England sample (Fig. 1C) sorting in

between the European and marginal Scottish samples. Overall, sorting of specimens along PC1 for

both species reveals a tight clustering among European butterflies and more geographic structuring

within GB samples.

Three historical *A. crataegi* specimens (NHM898, NHM027 and NHM918; Fig. 1B) labelled as GB

clustered closer to French and Belgium samples rather than the GB samples. Rather than a case of

mislabelling, these specimens may be potential European migrants or European specimens released

in Britain. The latter was not an uncommon practise from the late 19[th] to early 20[th] century (Pratt,

1983). For subsequent analyses, these three samples are grouped with the other mainland European

samples.

Demographic reconstruction

Historical demographic reconstruction of *A. crataegi* using PSMC shows similar trajectories of

effective population size (N_e; Fig. 2A, Fig. S4) for both British and European samples up to the last

glacial maximum (LGM, *c.* 22 to 17 Kya). British *A. crataegi,* show a marked decrease in N_e around

the LGM and Younger Dryas (12.8–11.5 Kya). A similar pattern of trajectories is also observed in *P.*

icarus (Fig. 2B, Fig. S5) but, the divergence in N_e appears to happen earlier, just before the LGM, but

this trend is sensitive to the number of generations per year (Fig. S6). Both reconstructions indicate

British *A. crataegi* and *P. icarus* show reduced N_e before or around 12 Kya.

Approximation of N_e at the time of the Younger Dryas (~10 Kya) are 3.87 x 10^4 for *A. crataegi,* and

2.14 x 10^5 for *P. icarus* (Fig. 2). We note that although we used higher coverage samples to estimate

the FNR for low coverage samples, for samples below 8X (*A. crataegi*) or 14X (*P. icarus*) we could not

recover N_e estimated for the high coverage data before 50 Kya (Fig. S7 and S8). However, this

discrepancy was limited to the magnitude of N_e and not the trajectories themselves.

Estimates of heterozygosity

Individual genome-wide heterozygosity (Fig. 3A-B) rates were lower for British samples for both *A. crataegi* and *P. icarus*. Heterozygosity between British and European *A. crataegi* was significantly different ($F_{1,14}$=25.21, *p*=0.000187, Fig. 3A), after accounting for variability in sample coverage. British *A. crataegi* exhibited an average reduction of 17.9% in heterozygosity relative to European samples. Heterozygosity was also significantly different between the British and French *P. icarus* (*t*=9.26, *degrees of freedom* (*df*) = 3.52, *p*= 0.001356; Fig. 3B) samples, with an average reduction of 14.5%.

Inbreeding and concordance of historical and recent N_e

For detecting RoHs we used three thresholds: ≥ 100 kb, ≥ 500 kb or ≥ 1 Mb. British *A. crataegi* showed a much higher proportion of the genome in RoH (7.6% in ≥ 100 Kb, 3.2% in ≥ 500 kb and 1.6% in ≥ 1 Mb; Fig. 3C) than European *A. crataegi* (0.6% in ≥ 100 Kb, 0.2% in ≥ 500 kb, and 0.1% in ≥ 1 Mb; Fig. 3C). Additionally, the difference in all RoH categories were significantly different (≥ 100 Kb: $F_{1,10}$=43.84, *p*=0.00005919; ≥ 500 kb: $F_{1,10}$=28.47 *p*=0.00033; ≥ 1MB: $F_{1,10}$=16.69, *p*=0.002918) with all British *A. crataegi* harbouring at least one RoH ≥ 1 Mb (Fig. S9), whereas only one of seven amongst European specimens harboured an RoH ≥ 1 Mb. In contrast, British *P. icarus* had a relatively smaller proportion (1.9%) of its genome in RoHs ≥ 100 Kb, 0.2% in ≥ 500 Kb, and none in regions ≥ 1 Mb (Fig. 3D), while French *P. icarus* had negligible RoHs in ≥ 100 Kb (0.03%) and none in ≥ 500 kb or ≥ 1 Mb. The difference in in RoH segments ≥ 100 Kb between British and French *P. icarus* was marginally significant (*t*=-3.15, *df* = 3, *p*= 0.0507; Fig. 3D).

Given a historical effective population size of 3.87×10^4 for *A. crataegi*, the probability of observing at least one RoH ≥ 1 Mb in a genome of 230 Mb is 0.051. Given that all British *A. crataegi* harbour at least one RoH ≥ 1 Mb (Fig. S9) this could indicate a lower recent N_e consistent with more recent inbreeding. With a historical of N_e of 2.14×10^5, the same probability for *P. icarus* was 0.021 respectively, consistent with this historical N_e, no RoH ≥ 1 Mb were observed.

Estimation of genetic load

In *A. crataegi* (Fig. 4A), the mean number of LoF variants was not significantly different ($F_{1,10}$=2.8,

p=0.125) between the British (17.5) and European specimens (20.2). However, the total numbers of

missense and synonymous were significantly lower in British *A. crataegi* (Fig. 4B & C*).* In *P. icarus,* all

variant classes were significantly lower in British samples (Fig. 4D-C). These are indicative of either

no change (LoF) or reduction (missense and synonymous) in genetic load in British specimens.

Realised genetic load (deleterious variants exposed in homozygosity, assuming most deleterious

variants are partially recessive) was determined by comparing the predicted variant classes in

homozygous (Fig. S10A-C) state. For *A. crataegi* there was no significant increase homozygosity for

LoF and missense classes, but there was a statistically significant increase in synonymous variants

($F_{1,10}$ = 6.37, *p*= 0.03) between the British specimens (893) and European (864) specimens. British *P.*

icarus show no significant change in homozygosity across all three variant classes (Fig S11).

Genetic load in terms of derived alleles for LoF variants was not significantly different between

British *A. crataegi* and mainland specimens albeit with a trend of decrease in the British specimens

(Fig. 5A). However, there was a statistically significant increase in frequencies of missense and

synonymous variants in British specimens relative to mainland specimens. Shared derived alleles in

British *P. icarus* (Fig. 5B) exhibit a significant decrease in frequencies of missense and synonymous

variants, but no statistically significant change in LoF frequency.

Discussion

Our extraction of historic DNA was contaminant free and we obtained short fragment lengths which

decreased in length with specimen age, a degradation pattern consistent with other studies using

historical DNA (e.g. Mullin *et al.*, 2023). We used these historical genomes of *A. crataegi* to provide a

genomic perspective on its extirpation from the British Isles in the early twentieth century and their

post-glacial colonization history in Britain. We complemented our analysis with a similar approach

using *P. icarus*, a butterfly with long-term stable population size within the British Isles. Our results

also provide genomic insights into the extirpation of *A. crataegi* and the utility of genomic erosion

metrics to identify vulnerable insect populations. This approach should be applicable to other insect

species using both museum and contemporary specimens. Comparative data on genomic erosion in

threatened, stable and expanding species, but also those with different life histories could improve the

predictive capacity for using metrics of genome erosion for identifying and managing threatened

populations and species (Bortoluzzi *et al.*, 2023; Kyriazis *et al.*, 2023; Webster *et al.*, 2023).

Different demographic histories of* A. crataegi *and* P. icarus *in Britain

Our population structure analysis (Fig. 1B-C) and PSMC-based demographic reconstruction (Fig. 2)

suggest that British populations of both *A. crataegi* and *P. icarus* are genetically distinct from their

continental European counterparts. However, there was evidence of strong structuring within the

British populations (Fig. 1B-C). Structuring apparent within the British specimens could result from

recent gene flow with European mainland populations or from sequential founder events and isolation

by distance following colonisation and subsequent founder events or a combination of these two

hypotheses. Gene flow and/or sequential founder events are a likely explanation for the structuring

observed in *P. icarus* as has been suggested by Arif *et al.*, (2021). However, for *A. crataegi*, the

consistency of demographic reconstructions (Fig. 2 and Fig S4), genome-wide heterozygosity (Fig.

2A), RoHs (Fig. 2C) and genetic load (Fig. 3A-C) amongst the British specimens, do not favour a

hypothesis of gene flow or admixture from the continent.

The post-glacial colonisation of the British Isles by *A. crataegi* and *P. icarus* are likely to have been

different (Fig. 2), based on their differing resource requirements (Middleton-Welling *et al.*, 2020) and

climate tolerances (Dennis 1992). *A. crataegi* uses wood edge shrub and tree species as hostplants

and is primarily a species associated with grassland edge/scrub matrices and requires relatively warm

summer temperatures whilst *P. icarus* is associated with pioneer or persistent short grassland

locations in which its hostplants occur (Howe *et al.*, 2007; De Keyser, *et al.*, 2012), and is tolerant of

lower temperatures than *A. crataegi* (Dennis, 1992). Conditions in Britain and the adjacent European

mainland during the Younger Dryas would not have permitted the occurrence of *A. crataegi* and it is

most likely to have colonized Britain in the short period from the late Boreal to the elimination of

channel land c.11.5 – 8.5 Kya (Dennis, 1992), probably arriving from mid-European latitudes. By

contrast, *P. icarus* may have persisted in sheltered southern valleys further north than *A. crataegi*

during the Younger Dryas, possibly in what is currently southern Britain, northern France and possibly

channel land, having originally reached southern Britain as early as the late Glacial period (c.14.5 –

13.5 Kya) (Dennis 1992). Thus, gene flow and admixture during the colonization period is more likely

for *P. icarus* than for *A. crataegi*. For *P. icarus* it is also possible that there were at least two

colonization periods, with good evidence for mixing of different lineages from distinct glacial refugia

(Arif *et al.,* 2021). Furthermore, admixture and gene flow are likely to be higher in *P. icarus* because of

its multivoltine life-history, compared to the univoltine strategy of *A. crataegi* which reduces the

number of generations over which these can occur over any time period.

Our PSMC-based demographic reconstructions and species life histories explain changes of effective

population sizes and the spatial structuring of populations. Both species underwent decreases of N_e

(Fig. 2 and Fig. S4) around the passing of the Eemian interglacial period (~125 Kya) until around the

last glacial maximum (LGM, ~20 Kya). After this time (~15 Kya) British *A. crataegi* exhibit a marked

reduction in N_e, consistent with a genetic bottleneck associated with a founder event and earlier

northward spread from southern glacial refugia. For British *A. crataegi* this N_e ($\approx 10^4$) is a magnitude

smaller than the commonly observed sizes of 10^5 or greater reported in butterflies (Mackintosh *et al.*,

2019; Ebdon *et al.*, 2021), but is consistent with PSMC-based estimates of several other British

Lepidoptera (Bortoluzzi *et al.*, 2023). At the same time, European *A. crataegi* show a drastic increase

in N_e presumably because of mixing of distinct populations from previously isolated glacial refugia.

(Hewitt, 1999; Hinojosa *et al.,* 2019). It is likely that colonization occurred over a relatively short time-

period, and possibly by individuals from few source populations. Our results also indicate a reduction

of N_e and a bottleneck for *P. icarus* with divergence from mainland European populations just before

the LGM but also note that inference of the precise timing of the *P. icarus* bottleneck is dependent on

the number of generations per year which may vary from 1 to 3 (Fig. S6). A bottleneck around the

time of colonization of Britain at the end of the last glacial would be consistent with the interpretation

of allozyme data by De Keyser *et al.* (2012), however it is also possible that the lineages of *P. icarus*

that colonised Britain never recovered from pre-LGM reductions in N_e, which may be the case for

rapidly colonising species. The discrepancy in the timings of our demographic reconstructions (Fig. 2)

and the exact timing of the colonisation of the British Isles by these species may also result from bias

in the PSMC method. Simulations by Bortoluzzi *et al.* (2023) suggest, that PSMC reconstructions for

species with small genomes and a high ratio of recombination to mutation rate, may reflect linked

selection rather than true demographic history. We cannot rule this out for either species, and this

bias could potentially explain differences in the demographic timing (along with uncertainty in

generation time for *P. icarus*). We note that our PSMC-based trajectories recapitulate changes in N_e

that are largely consistent with major climatic events of the last 10-125 Kya. Finally, we note that

absolute N_e of British *A. crataegi* ($\approx 10^4$) and *P. icarus* ($\approx 10^5$) are a function of their species-specific

N_e, which could be considerably smaller in *A. crataegi*, given that genetic diversity correlates

negatively with body size and positively with genome size in Lepidoptera (Mackintosh *et al.*, 2019).

Genomic symptoms of the demise of* A. crataegi *in Britain

The last recorded sighting of *A. crataegi* in Britain dates from *c.* 1925 (Pratt, 1983). However, by this

time, the species was thought to be limited to the south-eastern regions of Kent and had already

disappeared from other parts of its range as early as 1880 (Pratt, 1983 and references therein).

Consistent with these observations, three of 13 of our British samples, samples collected in 1888,

1908 and 1924 , appear to be of European origin in terms of population structure, demographic

history, and genomic erosion. These specimens may have been captive-bred or wild-caught

European (potentially from imported German larvae (Pratt, 1983)) stock released into British habitats

or potential migrants.

There was a reduction in genome-wide heterozygosity in British specimens of both species, but

reductions (compared to European populations) in both the extirpated *A. crataegi* and the extant and

stable *P. icarus* were roughly comparable (17.9 and 14.5%, respectively). As a comparison, the

observed reduction in genome-wide heterozygosity in the extinct *Glaucopsyche xerces* (Xerces Blue

butterfly; de-Dios *et al.*, 2023) was 22%, hence greater than what we observed for British *A. crataegi*.

However, the former was calculated in comparison to a sister species (*Glaucopsyche lygdamus*) and

hence it is unclear whether these estimates of reductions in genome-wide heterozygosity are strictly

comparable.

We saw significant increases in homozygous regions of small (≥ 100 Kb), intermediate (≥ 500 Kb) and

large (≥ 1 Mb) variety (Fig. 3C, Fig. S9) in British *A. crataegi*.. In contrast, no category of F_{RoH} was

significantly different between British and continental European *P. icarus* suggesting this population

likely does not suffer from inbreeding. The probability of observing large (≥ 1 Mb) RoHs, given the

PSMC-estimated historical N_e for both *A. crategi* and *P. icarus* were of the same magnitude (0.051

and 0.021, respectively), yet such RoH were more prevalent in *A. crategi*. The presence of the large

RoHs raise the possibility of more recent inbreeding, consistent with a spasmodic rather than instant

decline of the species in the 19[th] century. It is important to note that the proportions of genome in RoH

detected in *A. crataegi* genomes are likely an underestimate due to limited coverage in the museum

specimens (Table S1), as the fraction of detectable RoHs is a function of genome coverage

(Meyermans *et al*., 2020).

Accumulation of mildly or weakly deleterious variants, solely due to drift after a bottleneck, is expected

to lead to decrease in the total number of such variants but increase in homozygosity across all

variant classes (Dussex *et al.,* 2023). However, purging of deleterious variants through purifying

selection should lead to a decrease in strongly deleterious variants (LoF) in both homozygous and

heterozygous state, whereas homozygosity of mildly or weakly deleterious variants can still increase

and inflate the genetic load. The accumulation of genetic load in small populations has been observed

in many vertebrate populations (see references in Dussex *et al*. 2023) and in the Xerces blue butterfly

(de-Dios *et al*., 2023). However, there has also been evidence documentation of purging of the most

deleterious variants and accumulation of less deleterious variants in the same system in Alpine Ibex

(Grossen *et al*., 2020) and Indian tigers (Khan *et al*., 2021) and is supported by simulations (Dussex

et al., 2023). Although we observed no clear evidence of purging of LoF in British *A. crataegi* (Fig. 4A,

Fig. S10A and D, Fig. 5A), we did see an increase in shared derived allele frequency in the missense

and synonymous categories which may have contributed to extirpation by reducing fitness and

population viability. Additionally, we also observed a significant increase in realised load but this was

limited to only synonymous variants. Although synonymous variants are normally classified as neutral,

they can be weakly deleterious under codon bias and Lepidoptera do exhibit an A/T codon usage bias

(Näsvall *et al.*, 2023). In contrast, *P. icarus* displayed a decrease in realised load (Fig. S11) and a

decrease in frequencies of shared derived alleles of the missense and synonymous categories (Fig.

5B), suggesting these populations may have been less influenced by genetic drift. These results

raise possibility that the accumulation of mildly and weakly deleterious mutations contributed to the

decline of *A. crataegi* in Britain by triggering a mutational meltdown (Lynch *et al.*, 1995b). However, it

should be noted that the actual fitness effects including selection and dominance coefficients of the

annotated variants remains unknown. Moreover, given our limited sample sizes and lack of statistical

significance in some cases, our results on genetic load should be interpreted with caution. Finally,

given that the annotation for *P. icarus* (Lohse *et al.*, 2023) is based on a single male from Scotland,

our predictions of phenotypic impact of variants for this species may suffer from annotation bias, for

example, by reducing the number of predicted LoF and missense variants.

Overall, we provide compelling evidence that *A. crataegi* suffered a severe bottleneck during the

colonisation of the British Isles. Based on our analysis it is likely that N_e likely remained low following

the bottleneck and further reduced by more recent inbreeding, leading to the accumulation of weak or

mild deleterious mutations and potentially an increase in realised load, which may have further

exacerbated population persistence in the short term. Interestingly, two other butterfly species,

Lycaena dispar (Large copper butterfly) and *Cyaniris semiargus* (Mazarine blue butterfly) were also

extirpated in Britain between the mid to late 19[th] century. Like *A. crataegi*, both species expressed

single broods in Great Britain, but unlike *A. crataegi* both species are smaller with potentially larger

genomes and hence are expected to have higher ancestral N_e (Mackintosh *et al.*, 2019), however as

yet nothing is known about their demographic history in Britain. Hence, it would be instructive to

examine whether genomes of these extirpated populations exhibit similar symptoms of genomic

erosion observed in *A. crataegi* and *G. xerces.*

Metrics of genomic erosion for identifying vulnerable insect populations

Conservation genomics could be an extremely helpful tool in identifying populations or species at risk

of extinction or extirpation. This could be particularly relevant for invertebrates, whereas opposed to

vertebrates, reductions in census population size may not be easily monitored (Cardoso *et al.*, 2011;

Cardoso and Leather, 2019; Didham *et al.,* 2020). Our results suggest that a consistently small N_e on

the order of $\approx 10^4$, inbreeding in the recent past and accumulation of deleterious variants could be

viable indicators of population decline in species with generally large N_e. Indeed, similar signals of

genomic erosion have also been detected in the genomes of the extinct *G. xerces* (de-Dios *et al.*,

2023).

To evaluate how genomic erosion translates into population persistence or extinction we require

further empirical studies of not only declining but stable and expanding populations as well. Isolated

populations on islands, including the British Isles, offer an excellent system to study the genomics of

populations with contrasting or unknown fates (e.g. Bortoluzzi *et al.*, 2023; Kyriazis *et al.*, 2023).

Museum collections can provide a substantial and unequivocal source of threatened populations (e.g.,

samples from extirpated or extinct populations). Recent advances in resequencing of museum

samples (Korlević *et al.*, 2021, Mullin *et al.*, 2023), bioinformatic tools (e.g., Korneliussen *et al.*, 2014,

Kutschera *et al.,* 2022) and the availability of high-quality genomes with annotations (Rhie *et al.*,

2021; Formenti *et al.,* 2022; DToL, 2022) are providing much needed resources for such work.

Although challenges remain, for example, evaluating and interpreting fitness effects of deleterious

mutations (de Valles-Ibáñez *et al.*, 2016; Robinson *et al.*, 2023), we expect museomics in combination

with contemporary samples will expand our understanding of the genomics of extinction but also

improve feasibility to monitor and manage threatened insect populations (Díez-del-Molino *et al.,* 2018;

Jensen *et al.*, 2022; Bortoluzzi *et al.*, 2023).

Acknowledgements

This work was supported by a Nigel Groome PhD Studentship (Department of Biological and Medical

Sciences, Oxford Brookes University) to RW and a pilot funding grant from the Centre for Functional

Genomics, Oxford Brookes University to SA. We would like to thank the Oxford University Natural

History Museum (OUMNH) and the Natural History Museum, London (NHMUK) for permission to obtain

the Black-veined white museum samples. We would also like to thank Joe Middleton-Welling for

obtaining the samples for a pilot study from the OUMNH and the Darwin Tree of Life for the Black-

veined White genome assembly. We would also like to thank Konrad Lohse for help in calculating coalescent probabilities. We thank Laura Lukens for helpful comments on improving the manuscript.

Author Contributions

TGS, SA and RW designed the study. JH and GM helped procure museum specimens. TGS, SA and RW collected data. RW completed all molecular work with supervision and support from KH, CB and SA. RW analysed all the data with the help of SA. SA and RW wrote the paper with input from all authors.

Data Availability Statement

Raw sequence data is available under project accessing PRJEB70473 (for *A. crataegi*) and PRJEB70726 (for *P. icarus*) on the European Nucleotide Archive (ENA). Configuration files for GenErode and additional script files are available at https://github.com/rmwhitla/BVWpaper. Protocol for library preparation of historical samples is available at https://www.protocols.io/view/library-prep-for-cut-run-with-nebnext-ultra-ii-dna-bwgjpbun

References

Aken, B.L., Achuthan, P., Akanni, W., Amode, M.R., Bernsdorff, F., Bhai, J., Billis, K., Carvalho-Silva, D., Cummins, C., Clapham, P., Gil, L., Girón, C.G., Gordon, L., Hourlier, T., Hunt, S.E., Janacek, S.H., Juettemann, T., Keenan, S., Laird, M.R., Lavidas, I., Maurel, T., McLaren, W., Moore, B., Murphy, D.N., Nag, R., Newman, V., Nuhn, M., Ong, C.K., Parker, A., Patricio, M., Riat, H.S., Sheppard, D., Sparrow, H., Taylor, K., Thormann, A., Vullo, A., Walts, B., Wilder, S.P., Zadissa, A., Kostadima, M., Martin, F.J., Muffato, M., Perry, E., Ruffier, M., Staines, D.M., Trevanion, S.J., Cunningham, F., Yates, A., Zerbino, D.R., Flicek, P. (2017). Ensembl 2017. *Nucleic Acids Research*, **45**, D635–D642. https://doi.org/10.1093/nar/gkw1104

Andrews, S., (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Arif, S., Gerth, M., Hone-Millard, W.G., Nunes, M.D.S., Dapporto, L., Shreeve, T.G. (2021). Evidence for multiple colonisations and *Wolbachia* infections shaping the genetic structure of the widespread butterfly *Polyommatus icarus* in the British Isles. *Molecular Ecology*, **30**, 5196–5213. https://doi.org/10.1111/mec.16126

Barnosky, A.D., Matzke, N., Tomiya, S., Wogan, G.O.U., Swartz, B., Quental, T.B., Marshall, C., McGuire, J.L., Lindsey, E.L., Maguire, K.C., Mersey, B., Ferrer, E.A. (2011). Has the Earth's sixth mass extinction already arrived? *Nature*, **471**, 51–57. https://doi.org/10.1038/nature09678

Biscarini, F., Cozzi, P., Gaspa, G., Marras, G. (2018). detectRUNS: Detect runs of homozygosity and runs of heterozygosity in diploid genomes [WWW Document]. URL https://cran.r-project.org/web/packages/detectRUNS/index.html

Bortoluzzi, C., Wright, C. J., Lee, S., Cousins, T., Genez, T. A. L., Thybert, D., Martin, F. J., Haggerty, L., The Darwin Tree of Life Project Consortium, Blaxter, M., Durbin, R. (2023). Lepidoptera genomics based on 88 chromosomal reference sequences informs population

genetic parameters for conservation. *bioRxiv* 2023.04.14.536868. [PREPRINT]

https://doi.org/10.1101/2023.04.14.536868

Bosse, M., van Loon, S. (2022). Challenges in quantifying genome erosion for conservation.

Frontiers in Genetics, **13**, 960958. https://doi.org/10.3389/fgene.2022.960958

Burns, F., Mordue, S., al Fulaij, N., Boersch-Supan, P. H., Boswell, J., Boyd, R. J., Bradfer-

Lawrence, T., de Ornellas, P., de Palma, A., de Zylva, P., Dennis, E. B., Foster, S., Gilbert, G.,

Halliwell, L., Hawkins, K., Kaysom, K. A., Holland, M. M., Hughes, J., Jackson, A. C., Mancini,

F., Mathews, F., McQuatters-Gollop, A., Noble, D. G., O'Brien, D., Pescott, O. L., Purvis, A.,

Simkin, J., Smith, A., Stanbury, A. J., Villemot, J., Walker, K. J., Walton, P., Webb, T. J.,

Williams, K., Wilson, R., Gregory, R. D. (2023). State of Nature 2023, the State of Nature

Partnership. Available at: www.stateofnature.org.uk

Cardoso, P., Erwin, T.L., Borges, P.A.V., New, T.R. (2011). The seven impediments in invertebrate

conservation and how to overcome them. *Biological Conservation*, **144**, 2647–2655.

https://doi.org/10.1016/j.biocon.2011.07.024

Cardoso, P., Leather, S.R. (2019). Predicting a global insect apocalypse. *Insect Conservation and*

Diversity, **12**, 263–267. https://doi.org/10.1111/icad.12367

Ceballos, G., Ehrlich, P.R., Barnosky, A.D., García, A., Pringle, R.M., Palmer, T.M. (2015).

Accelerated modern human-induced species losses: Entering the sixth mass extinction.

Science Advances, **1**, e1400253. https://doi.org/10.1126/sciadv.1400253

Ceballos, F.C., Joshi, P.K., Clark, D.W., Ramsay, M., Wilson, J.F. (2018). Runs of homozygosity:

windows into population history and trait architecture. *Nature Reviews Genetics*, **19**, 220–234.

https://doi.org/10.1038/nrg.2017.109

Charlesworth, D., Willis, J.H. (2009). The genetics of inbreeding depression. *Nature Reviews*

Genetics, **10**, 783–796. https://doi.org/10.1038/nrg2664

Chen, S., Zhou, Y., Chen, Y., Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics, **34**, i884–i890. https://doi.org/10.1093/bioinformatics/bty560

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X., Ruden, D.M.

(2012). A program for annotating and predicting the effects of single nucleotide polymorphisms,

SnpEff. *Fly (Austin)*, **6**, 80–92. https://doi.org/10.4161/fly.19695

Cowie, R.H., Bouchet, P., Fontaine, B. (2022). The Sixth Mass Extinction: fact, fiction or speculation?

Biological Reviews, **97**, 640–663. https://doi.org/10.1111/brv.12816

Dale, C.W. (1887). Query regarding *Aporia crataegi*. *The Entomologist's Monthly Magazine*, **23**, 214.

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane,

T., McCarthy, S.A., Davies, R.M., Li, H. (2021). Twelve years of SAMtools and BCFtools.

Gigascience, **10**, giab008. https://doi.org/10.1093/gigascience/giab008

Darwin Tree of Life Project Consortium. (2022). Sequence locally, think globally: The Darwin Tree of

Life Project. *Proceedings of the National Academy of Sciences of the United States of America*,

119, e2115642118. https://doi.org/10.1073/pnas.2115642118

de-Dios, T., Fontsere, C., Renom, P., Stiller, J., Llovera, L., Uliano-Silva, M., Sánchez-Gracia, A.,

Wright, C., Lizano, E., Caballero, B., Navarro, A., Civit, S., Robbins, R.K., Blaxter, M., Marquès-

Bonet, T., Vila, R., Lalueza-Fox, C. (2023). Whole-genomes from the extinct Xerces Blue

butterfly can help identify declining insect species. *eLife*, **12**. https://doi.org/10.7554/eLife.87928

De Keyser, R., Shreeve, T.G., Breuker, C.J., Hails, R.S., Schmitt, T. (2012). *Polyommatus icarus*

butterflies in the British Isles: evidence for a bottleneck. *Biological Journal of the Linnean*

Society, **107**, 123–136. https://doi.org/10.1111/j.1095-8312.2012.01925.x

de Valles-Ibáñez, G., Hernandez-Rodriguez, J., Prado-Martinez, J., Luisi, P., Marquès-Bonet, T.,

Casals, F. (2016). Genetic Load of Loss-of-Function Polymorphic Variants in Great Apes.

Genome Biology and Evolution, **8**, 871–877. https://doi.org/10.1093/gbe/evw040

Dennis, R.L.H. (1992). An evolutionary history of British butterflies. In Dennis, R.L.H. (Ed.) *The*

Ecology of Butterflies in Britain. Oxford, Oxford University Press, pp. 217-245.

Dennis, R.L.H., Shreeve, T.G., Van Dyck, H. (2003). Towards a Functional Resource-Based Concept

for Habitat: A Butterfly Biology Viewpoint. *Oikos*, **102**, 417–426. https://doi.org/10.1034/j.1600-

0579.2003.12492.x

DeWoody, J.A., Harder, A.M., Mathur, S., Willoughby, J.R. (2021). The long-standing significance of

genetic diversity in conservation. *Molecular Ecology*, **30**, 4147–4154.

https://doi.org/10.1111/mec.16051

Didham, R.K., Basset, Y., Collins, C.M., Leather, S.R., Littlewood, N.A., Menz, M.H.M., Müller, J.,

Packer, L., Saunders, M.E., Schönrogge, K., Stewart, A.J.A., Yanoviak, S.P., Hassall, C.

(2020). Interpreting insect declines: seven challenges and a way forward. *Insect Conservation*

and Diversity, **13**, 103–114. https://doi.org/10.1111/icad.12408

Díez-del-Molino, D., Sánchez-Barreiro, F., Barnes, I., Gilbert, M.T.P., Dalén, L. (2018). Quantifying

Temporal Genomic Erosion in Endangered Species. *Trends in Ecology & Evolution*, **33**, 176–

185. https://doi.org/10.1016/j.tree.2017.12.002

Dincă, V., Montagud, S., Talavera, G., Hernández-Roldán, J., Munguira, M.L., García-Barros, E.,

Hebert, P.D.N., Vila, R. (2015). DNA barcode reference library for Iberian butterflies enables a

continental-scale preview of potential cryptic diversity. *Scientific Reports*, **5**, 12395.

https://doi.org/10.1038/srep12395

Dussex, N., van der Valk, T., Morales, H.E., Wheat, C.W., Díez-del-Molino, D., von Seth, J., Foster,

Y., Kutschera, V.E., Guschanski, K., Rhie, A., Phillippy, A.M., Korlach, J., Howe, K., Chow, W.,

Pelan, S., Mendes Damas, J.D., Lewin, H.A., Hastie, A.R., Formenti, G., Fedrigo, O., Guhlin, J.,

Harrop, T.W.R., Le Lec, M.F., Dearden, P.K., Haggerty, L., Martin, F.J., Kodali, V., Thibaud-

Nissen, F., Iorns, D., Knapp, M., Gemmell, N.J., Robertson, F., Moorhouse, R., Digby, A.,

Eason, D., Vercoe, D., Howard, J., Jarvis, E.D., Robertson, B.C., Dalén, L. (2021). Population

genomics of the critically endangered kākāpō. *Cell Genomics*, **1**, 100002.

https://doi.org/10.1016/j.xgen.2021.100002

Dussex, N., Morales, H.E., Grossen, C., Dalén, L., Oosterhout, C. van. (2023). Purging and

accumulation of genetic load in conservation. *Trends in Ecology & Evolution*, **38**, 961–969.

https://doi.org/10.1016/j.tree.2023.05.008

Ebdon, S., Laetsch, D. R., Dapporto, L., Hayward, A., Ritchie, M. G., Dincă, V., Vila, R.,

& Lohse, K. (2021). The Pleistocene species pump past its prime: Evidence from European

butterfly sister species. *Molecular Ecology*, **30**, 3575–3589. https://doi.org/10.1111/mec.15981

Ebdon, S., Mackintosh, A., Lohse, K., Hayward, A., Arif, S., Whitla, R. (2022). The genome sequence

of the black-veined white butterfly, *Aporia crataegi* (Linnaeus, 1758) [version 1; peer review: 1

approved, 1 not approved]. *Wellcome Open Research* **7**:81.

https://doi.org/10.12688/wellcomeopenres.17709.1

Formenti, G., Theissinger, K., Fernandes, C., Bista, I., Bombarely, A., Bleidorn, C., Ciofi, C., Crottini,

A., Godoy, J.A., Höglund, J., Malukiewicz, J., Mouton, A., Oomen, R.A., Paez, S., Palsbøll,

P.J., Pampoulie, C., Ruiz-López, María J., Svardal, H., Theofanopoulou, C., de Vries, J.,

Waldvogel, A.-M., Zhang, Guojie, Mazzoni, C.J., Jarvis, E.D., Bálint, M., Formenti, G.,

Theissinger, K., Fernandes, C., Bista, I., Bombarely, A., Bleidorn, C., Čiampor, F., Ciofi, C.,

Crottini, A., Godoy, J.A., Hoglund, J., Malukiewicz, J., Mouton, A., Oomen, R.A., Paez, S.,

Palsbøll, P., Pampoulie, C., Ruiz-López, María José, Svardal, H., Theofanopoulou, C., de

Vries, J., Waldvogel, A.-M., Zhang, Goujie, Mazzoni, C.J., Jarvis, E., Bálint, M., Aghayan, S.A.,

Alioto, T.S., Almudi, I., Alvarez, N., Alves, P.C., Amorim, I.R., Antunes, A., Arribas, P., Baldrian,

P., Berg, P.R., Bertorelle, G., Böhne, A., Bonisoli-Alquati, A., Boštjančić, L.L., Boussau, B.,

Breton, C.M., Buzan, E., Campos, P.F., Carreras, C., Castro, L.Fi., Chueca, L.J., Conti, E.,

Cook-Deegan, R., Croll, D., Cunha, M.V., Delsuc, F., Dennis, A.B., Dimitrov, D., Faria, R.,

Favre, A., Fedrigo, O.D., Fernández, R., Ficetola, G.F., Flot, J.-F., Gabaldón, T., Galea Agius,

D.R., Gallo, G.R., Giani, A.M., Gilbert, M.T.P., Grebenc, T., Guschanski, K., Guyot, R.,

Hausdorf, B., Hawlitschek, O., Heintzman, P.D., Heinze, B., Hiller, M., Husemann, M., Iannucci,

A., Irisarri, I., Jakobsen, K.S., Jentoft, S., Klinga, P., Kloch, A., Kratochwil, C.F., Kusche, H.,

Layton, K.K.S., Leonard, J.A., Lerat, E., Liti, G., Manousaki, T., Marques-Bonet, T., Matos-

Maraví, P., Matschiner, M., Maumus, F., Mc Cartney, A.M., Meiri, S., Melo-Ferreira, J.,

Mengual, X., Monaghan, M.T., Montagna, M., Mysłajek, R.W., Neiber, M.T., Nicolas, V., Novo,

M., Ozretić, P., Palero, F., Pârvulescu, L., Pascual, M., Paulo, O.S., Pavlek, M., Pegueroles, C.,

Pellissier, L., Pesole, G., Primmer, C.R., Riesgo, A., Rüber, L., Rubolini, D., Salvi, D.,

Seehausen, O., Seidel, M., Secomandi, S., Studer, B., Theodoridis, S., Thines, M., Urban, L.,

Vasemägi, A., Vella, A., Vella, N., Vernes, S.C., Vernesi, C., Vieites, D.R., Waterhouse, R.M.,

Wheat, C.W., Wörheide, G., Wurm, Y., Zammit, G. (2022). The era of reference genomes in

conservation genomics. *Trends in Ecology & Evolution*, **37**, 197–202.

https://doi.org/10.1016/j.tree.2021.11.008

Fox, R., Dennis, E.B., Brown, A.F., Curson, J. (2022). A revised Red List of British butterflies. *Insect

Conservation and Diversity*, **15**, 485–495. https://doi.org/10.1111/icad.12582

Gomulkiewicz, R., Holt, R.D. (1995). When does Evolution by Natural Selection Prevent Extinction?

Evolution, **49**, 201–207. https://doi.org/10.2307/2410305

Grossen, C., Guillaume, F., Keller, L.F., Croll, D. (2020). Purging of highly deleterious mutations

through severe bottlenecks in Alpine ibex. *Nature Communications*, **11**, 1001.

https://doi.org/10.1038/s41467-020-14803-1

Hewitt, G.M. (1999). Post-glacial re-colonization of European biota. *Biological Journal of the Linnean*

Society, **68**, 87–112. https://doi.org/10.1111/j.1095-8312.1999.tb01160.x

Hill, G.M., Kawahara, A.Y., Daniels, J.C., Bateman, C.C., Scheffers, B.R. (2021). Climate change

effects on animal ecology: butterflies and moths as a case study. *Biological Reviews of the*

Cambridge Philosophical Society, **96**, 2113–2126. https://doi.org/10.1111/brv.12746

Hinojosa, J.C., Koubínová, D., Szenteczki, M.A., Pitteloud, C., Dincă, V., Alvarez, N., Vila, R. (2019).

A mirage of cryptic species: Genomics uncover striking mitonuclear discordance in the butterfly

Thymelicus sylvestris. *Molecular Ecology*, **28**, 3857–3868. https://doi.org/10.1111/mec.15153

Howe, P.D., Bryant, S.R., Shreeve, T.G. (2007). Predicting body temperature and activity of adult

Polyommatus icarus using neural network models under current and projected climate

scenarios. *Oecologia,* **153**(4), 857-69. https://doi.org/10.1007/s00442-007-0782-3

Hudson, R. R. (1990). Gene Genealogies and the coalescent process. In D. Futuyma and J.

Antonovics (Eds*.), Oxford Surveys in Evolutionary Biology, Vol. 7*. Oxford University Press,

New York, pp. 1-44.

IPBES, 2019. Summary for policymakers of the global assessment report on biodiversity and

ecosystem services. IPBES Plenary at its seventh session (IPBES 7, Paris, 2019) 29 April - 4

May 2019. https://doi.org/10.5281/zenodo.3553579

Jensen, E.L., Díez-del-Molino, D., Gilbert, M.T.P., Bertola, L.D., Borges, F., Cubric-Curik, V., de

Navascués, M., Frandsen, P., Heuertz, M., Hvilsom, C., Jiménez-Mena, B., Miettinen, A.,

Moest, M., Pečnerová, P., Barnes, I., Vernesi, C. (2022). Ancient and historical DNA in

conservation policy. *Trends in Ecology & Evolution*, **37**, 420–429.

https://doi.org/10.1016/j.tree.2021.12.010

Jiggins, C.D., Mavarez, J., Beltrán, M., McMillan, W.O., Johnston, J.S., Bermingham, E. (2005) A

Genetic Linkage Map of the Mimetic Butterfly *Heliconius melpomene*, *Genetics*, **171**(2), 557–

570, https://doi.org/10.1534/genetics.104.034686

Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P.L.F., Orlando, L. (2013). mapDamage2.0: fast

approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics*, **29**,

1682–1684. https://doi.org/10.1093/bioinformatics/btt193

Karam-Gemael, M., Decker, P., Stoev, P., Marques, M.I., Jr, A.C. (2020). Conservation of terrestrial

invertebrates: a review of IUCN and regional Red Lists for Myriapoda. *ZooKeys*, **930**, 221–229.

https://doi.org/10.3897/zookeys.930.48943

Kardos, M., Taylor, H.R., Ellegren, H., Luikart, G., Allendorf, F.W. (2016). Genomics advances the

study of inbreeding depression in the wild. *Evolutionary Applications*, **9**, 1205–1218.

https://doi.org/10.1111/eva.12414

Keightley, P.D., Pinharanda, A., Ness, R.W., Simpson, F., Dasmahapatra, K.K., Mallet, J., Davey,

J.W., Jiggins, C.D. (2015). Estimation of the Spontaneous Mutation Rate in *Heliconius*

melpomene. *Molecular Biology and Evolution*, **32**, 239–243.

https://doi.org/10.1093/molbev/msu302

Khan, A., Patel, K., Shukla, H., Viswanathan, A., van der Valk, T., Borthakur, U., Nigam, P.,

Zachariah, A., Jhala, Y.V., Kardos, M., Ramakrishnan, U. (2021). Genomic evidence for

inbreeding depression and purging of deleterious genetic variation in Indian tigers. *Proceedings*

of the National Academy of Sciences, **118**, e2023018118.

https://doi.org/10.1073/pnas.2023018118

Kim, T., Han, Y.-G., Kwon, O., Cho, Y. (2015). Changes in *Aporia crataegi*'s potential habitats in

accordance with climate changes in the northeast Asia. *Journal of Ecology and Environment*,

38, 15–23. https://doi.org/10.5141/ecoenv.2015.002

Korlević, P., McAlister, E., Mayho, M., Makunin, A., Flicek, P., Lawniczak, M.K.N. (2021). A Minimally

Morphologically Destructive Approach for DNA Retrieval and Whole-Genome Shotgun

Sequencing of Pinned Historic Dipteran Vector Species. *Genome Biology and Evolution*, **13**,

evab226. https://doi.org/10.1093/gbe/evab226

Korneliussen, T.S., Albrechtsen, A., Nielsen, R. (2014). ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics*, **15**, 356. https://doi.org/10.1186/s12859-014-0356-4

Kutschera, V.E., Kierczak, M., van der Valk, T., von Seth, J., Dussex, N., Lord, E., Dehasque, M.,

Stanton, D.W.G., Khoonsari, P.E., Nystedt, B., Dalén, L., Díez-del-Molino, D. (2022).

GenErode: a bioinformatics pipeline to investigate genome erosion in endangered and extinct

species. *BMC Bioinformatics*, **23**, 228. https://doi.org/10.1186/s12859-022-04757-0

Kyriazis, C.C., Beichman, A.C., Brzeski, K.E., Hoy, S.R., Peterson, R.O., Vucetich, J.A., Vucetich,

L.M., Lohmueller, K.E., Wayne, R.K. (2023). Genomic Underpinnings of Population Persistence

in Isle Royale Moose. *Molecular Biology and Evolution*, **40**, msad021.

https://doi.org/10.1093/molbev/msad021

Lenth, R. (2023). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.8.7. https://github.com/rvlenth/emmeans

Li, H., Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics, **25**, 1754–1760. https://doi.org/10.1093/bioinformatics/btp324

Li, H., Durbin, R. (2011). Inference of human population history from individual whole-genome

sequences. *Nature*, **475**, 493–496. https://doi.org/10.1038/nature10231

Liu, N. (2019). Library Prep for CUT&RUN with NEBNext® Ultra™ II DNA Library Prep Kit for

Illumina® (E7645). protocols.io https://dx.doi.org/10.17504/protocols.io.wvgfe3w

Lohse, K. (2023). The genome sequence of the Common Blue, *Polyommatus icarus* (Rottemburg,

1775). [version 1; peer review: 1 approved]. *Wellcome Open Research* **8**:72.

https://doi.org/10.12688/wellcomeopenres.18772.1

Lynch, M., Conery, J., Bürger, R. (1995a). Mutational meltdowns in sexual populations. *Evolution*,

49, 1067–1080. https://doi.org/10.1111/j.1558-5646.1995.tb04434.x

Lynch, M., Conery, J., Burger, R. (1995b). Mutation Accumulation and the Extinction of Small

Populations. *The American Naturalist*, **146**, 489–518. https://doi.org/10.1086/285812

Mackintosh, A., Laetsch, D. R., Hayward, A., Charlesworth, B., Waterfall, M., Vila, R., & Lohse,

K. (2019). The determinants of genetic diversity in butterflies. *Nature Communications*, **10**,

3466. https://doi.org/10.1038/s41467-019-11308-4

Mackintosh, A., Vila, R., Martin, S.H., Setter, D., Lohse, K. (2023). Do chromosome rearrangements

fix by genetic drift or natural selection? Insights from *Brenthis* butterflies. *Molecular Ecology*,

00, 1–15. https://doi.org/10.1111/mec.17146

MacLachlan, R. (1893). The Decadence of British Butterflies. *The Entomologist's Monthly Magazine*,

29, 132–138.

Mattila, A.L.K., Duplouy, A., Kirjokangas, M., Lehtonen, R., Rastas, P., Hanski, I. (2012). High

genetic load in an old isolated butterfly population. *Proceedings of the National Academy of*

Sciences, **109**, E2496–E2505. https://doi.org/10.1073/pnas.1205789109

Meisner, J., Albrechtsen, A. (2018). Inferring Population Structure and Admixture Proportions in Low-

Depth NGS Data. *Genetics*, **210**, 719–731. https://doi.org/10.1534/genetics.118.301336

Meyermans R, Gorssen W, Buys N, Janssens S. (2020). How to study runs of homozygosity using

PLINK? A guide for analyzing medium density SNP data in livestock and pet species. *BMC*

Genomics, **29**;21(1):94. https://doi: 10.1186/s12864-020-6463-x.

Middleton-Welling, J., Dapporto, L., García-Barros, E., Wiemers, M., Nowicki, P., Plazio, E., Bonelli,

S., Zaccagno, M., Šašić, M., Liparova, J., Schweiger, O., Harpke, A., Musche, M., Settele, J.,

Schmucki, R., Shreeve, T. (2020). A new comprehensive trait database of European and

Maghreb butterflies, Papilionoidea. *Scientific Data*, **7**, 351. https://doi.org/10.1038/s41597-020-

00697-7

Mullin, V.E., Stephen, W., Arce, A.N., Nash, W., Raine, C., Notton, D.G., Whiffin, A., Blagderov, V.,

Gharbi, K., Hogan, J., Hunter, T., Irish, N., Jackson, S., Judd, S., Watkins, C., Haerty, W.,

Ollerton, J., Brace, S., Gill, R.J., Barnes, I. (2023). First large-scale quantification study of DNA

preservation in insects from natural history collections using genome-wide sequencing.

Methods in Ecology and Evolution, **14**, 360–371. https://doi.org/10.1111/2041-210X.13945

Näsvall, K., Boman, J., Talla, V., Backström, N. (2023). Base Composition, Codon Usage, and

Patterns of Gene Sequence Evolution in Butterflies. *Genome Biology and Evolution*, **15**,

evad150. https://doi.org/10.1093/gbe/evad150

Neaves, L.E., Eales, J., Whitlock, R., Hollingsworth, P.M., Burke, T., Pullin, A.S. (2015). The fitness

consequences of inbreeding in natural populations and their implications for species

conservation – a systematic map. *Environmental Evidence*, **4**:5. https://doi.org/10.1186/s13750-

015-0031-x

Neukamm, J., Peltzer, A., Nieselt, K. (2021). DamageProfiler: fast damage pattern calculation for

ancient DNA. *Bioinformatics*, **37**, 3652–3653. https://doi.org/10.1093/bioinformatics/btab190

1000

1001 Palahí I Torres, A., Höök, L., Näsvall, K., Shipilina, D., Wiklund, C., Vila, R., Pruisscher, P.,

1002   Backström, N. (2023) The fine-scale recombination rate variation and associations with

1003   genomic features in a butterfly. *Genome Research*. **33**(5), 810-823.

1004   https://doi.org/10.1101/gr.277414.122

1005

1006 Pratt, C. (1983). A modern review of the demise of *Aporia crataegi L.*: the black veined white. *The*

1007   *entomologist's record and journal of variation*, **95**, 161--166.

1008

1009 Preston, S.D., Liao, J.D., Toombs, T.P., Romero-Canyas, R., Speiser, J., Seifert, C.M. (2021). A

1010   case study of a conservation flagship species: the monarch butterfly. *Biodiversity and*

1011   *Conservation*, **30**, 2057–2077. https://doi.org/10.1007/s10531-021-02183-x

1012

1013 Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J., Sklar, P.,

1014   de Bakker, P.I.W., Daly, M.J., Sham, P.C. (2007). PLINK: A Tool Set for Whole-Genome

1015   Association and Population-Based Linkage Analyses. *American Journal of Human Genetics*,

1016   **81**, 559–575. https://doi.org/10.1086/519795

1017

1018 R Core Team. (2023). R: A language and environment for statistical computing. R Foundation for

1019   Statistical Computing, Vienna, Austria.

1020

1021 Rhie, A., McCarthy, S.A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow,

1022   W., Fungtammasan, A., Kim, J., Lee, C., Ko, B.J., Chaisson, M., Gedman, G.L., Cantin, L.J.,

1023   Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., Haase, B., Mountcastle, J., Winkler, S.,

1024   Paez, S., Howard, J., Vernes, S.C., Lama, T.M., Grutzner, F., Warren, W.C., Balakrishnan,

1025   C.N., Burt, D., George, J.M., Biegler, M.T., Iorns, D., Digby, A., Eason, D., Robertson, B.,

1026   Edwards, T., Wilkinson, M., Turner, G., Meyer, A., Kautt, A.F., Franchini, P., Detrich, H.W.,

1027   Svardal, H., Wagner, M., Naylor, G.J.P., Pippel, M., Malinsky, M., Mooney, M., Simbirsky, M.,

1028   Hannigan, B.T., Pesout, T., Houck, M., Misuraca, A., Kingan, S.B., Hall, R., Kronenberg, Z.,

1029   Sović, I., Dunn, C., Ning, Z., Hastie, A., Lee, J., Selvaraj, S., Green, R.E., Putnam, N.H., Gut, I.,

Ghurye, J., Garrison, E., Sims, Y., Collins, J., Pelan, S., Torrance, J., Tracey, A., Wood, J.,

Dagnew, R.E., Guan, D., London, S.E., Clayton, D.F., Mello, C.V., Friedrich, S.R., Lovell, P.V.,

Osipova, E., Al-Ajli, F.O., Secomandi, S., Kim, H., Theofanopoulou, C., Hiller, M., Zhou, Y.,

Harris, R.S., Makova, K.D., Medvedev, P., Hoffman, J., Masterson, P., Clark, K., Martin, F.,

Howe, Kevin, Flicek, P., Walenz, B.P., Kwak, W., Clawson, H., Diekhans, M., Nassar, L., Paten,

B., Kraus, R.H.S., Crawford, A.J., Gilbert, M.T.P., Zhang, G., Venkatesh, B., Murphy, R.W.,

Koepfli, K.-P., Shapiro, B., Johnson, W.E., Di Palma, F., Marques-Bonet, T., Teeling, E.C.,

Warnow, T., Graves, J.M., Ryder, O.A., Haussler, D., O'Brien, S.J., Korlach, J., Lewin, H.A.,

Howe, Kerstin, Myers, E.W., Durbin, R., Phillippy, A.M., Jarvis, E.D. (2021). Towards complete

and error-free genome assemblies of all vertebrate species. *Nature*, **592**, 737–746.

https://doi.org/10.1038/s41586-021-03451-0

1041

1042 Robinson, J., Kyriazis, C.C., Yuan, S.C., Lohmueller, K.E. (2023). Deleterious Variation in Natural

1043 Populations and Implications for Conservation Genetics. *Annual Review of Animal Biosciences*,

1044 **11**, 93–114. https://doi.org/10.1146/annurev-animal-080522-093311

1045

1046 Rowe, G., Sweet, M., Beebee, T. (2017). *An Introduction to Molecular Ecology*, Third Edition. Oxford,

1047 Oxford University Press.

1048

1049 Saccheri, I., Kuussaari, M., Kankare, M., Vikman, P., Fortelius, W., Hanski, I. (1998). Inbreeding and

1050 extinction in a butterfly metapopulation. *Nature*, **392**, 491–494. https://doi.org/10.1038/33136

1051

1052 Sarabia, C., vonHoldt, B., Larrasoaña, J.C., Uríos, V., Leonard, J.A. (2021). Pleistocene climate

1053 fluctuations drove demographic history of African golden wolves (*Canis lupaster*). *Molecular*

1054 *Ecology*, **30**, 6101–6120. https://doi.org/10.1111/mec.15784

1055

1056 Solonkin, I.A., Shkurikhin, A.O., Oslina, T.S., Zakharova, E.Y. (2021). Changes in the body size of

1057 black-veined white, *Aporia crataegi* (Lepidoptera: Pieridae), recorded in a natural population in

1058 response to different spring weather conditions and at different phases of an outbreak.

1059 *European Journal of Entomology*, **118**, 214–224. https://doi.org/10.14411/eje.2021.023

1060

1061   South, R. (1906). *The Butterflies of the British Isles.* London, Frederick Warne.

1062

1063   Teixeira, J.C., Huber, C.D. (2021). The inflated significance of neutral genetic diversity in

1064   conservation genetics. *Proceedings of the National Academy of Sciences of the United States*

1065   *of America*, **118**, e2015096118. https://doi.org/10.1073/pnas.2015096118

1066

1067   Thomas, J. A. (2005). Monitoring change in the abundance and distribution of insects using

1068   butterflies and other indicator groups*. Philosophical Transactions of the Royal Society B:*

1069   *Biological Sciences*, **360**, 339–357. https://doi.org/10.1098/rstb.2004.1585

1070

1071   van der Auwera, G., O'Connor, B.D. (2020). Genomics in the Cloud: Using Docker, GATK, and WDL

1072   in Terra. *O'Reilly Media*, *Incorporated*.

1073

1074   van der Valk, T., Díez-Del-Molino, D., Marques-Bonet, T., Guschanski, K., Dalén, L. (2019).

1075   Historical Genomes Reveal the Genomic Consequences of Recent Population Decline in

1076   Eastern Gorillas. *Current Biology*, **29**, 165-170.e6. https://doi.org/10.1016/j.cub.2018.11.055

1077

1078   van Swaay, C., Wynhoff, I., Verovnik, R., Wiemers, M., López Munguira, M., Maes, D., Sasic, M.,

1079   Verstrael, T., Warren, M. & Settele, J. (2010)*. Aporia crataegi* (Europe assessment). The IUCN

1080   Red List of Threatened Species 2010: e.T159814A5339278. Accessed on 17 January 2024.

1081

1082   Wagner, D.L., Grames, E.M., Forister, M.L., Berenbaum, M.R., Stopak, D. (2021). Insect decline in

1083   the Anthropocene: Death by a thousand cuts. *Proceedings of the National Academy of*

1084   *Sciences*, **118**, e2023989118. https://doi.org/10.1073/pnas.2023989118

1085

1086   Warren, M.S. (1992). The Conservation of British Butterflies, in Dennis, R.L.H. *The Ecology of*

1087   *Butterflies in Britain.* Oxford, Oxford University Press, pp. 246–274.

1088

1089 Warren, M.S., Maes, D., van Swaay, C.A.M., Goffart, P., Van Dyck, H., Bourn, N.A.D., Wynhoff, I.,

1090   Hoare, D., Ellis, S. (2021). The decline of butterflies in Europe: Problems, significance, and

1091   possible solutions. *Proceedings of the National Academy of Sciences of the United States of*

1092   *America*, **118**, e2002551117. https://doi.org/10.1073/pnas.2002551117

1093

1094 Webster, M.T., Beaurepaire, A., Neumann, P., Stolle, E. (2023). Population Genomics for Insect

1095   Conservation. *Annual Review of Animal Biosciences*, **11**, 115–140.

1096   https://doi.org/10.1146/annurev-animal-122221-075025

1097

Xue, Y., Prado-Martinez, J., Sudmant, P.H., Narasimhan, V., Ayub, Q., Szpak, M., Frandsen, P.,

Chen, Y., Yngvadottir, B., Cooper, D.N., de Manuel, M., Hernandez-Rodriguez, J., Lobon, I.,

Siegismund, H.R., Pagani, L., Quail, M.A., Hvilsom, C., Mudakikwa, A., Eichler, E.E., Cranfield,

M.R., Marques-Bonet, T., Tyler-Smith, C., Scally, A. (2015). Mountain gorilla genomes reveal

the impact of long-term population decline and inbreeding. *Science*, **348**, 242–245.

https://doi.org/10.1126/science.aaa3952

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

Figure Legends

Figure 1 (A): Collection locations (based on museum records) of historical A. crataegi *specimens (circles) and of modern* P. icarus *(triangles). NHM prefix refers to specimens obtained from Natural History Museum, London and OX for those obtained from Oxford University Natural History Museum, Oxford. (B): Principal component analysis (PCA) for* A. crataegi *SNPs. The PCA is based on 14,930 LD-pruned SNPs based on hard genotype calls. A PCA using genotype likelihoods shows a similar pattern (Fig S3). Asterisks indicate UK-collected specimens that have clustered with European specimens. (C): PCA for* P. icarus *specimens based on 197,182 LD-pruned SNPs showing similar differentiation between British and European specimens. Inset images: A pinned* A. crataegi *specimen and live* P. icarus *in their respective plots.*

Figure 2: (A): PSMC plot for A. crataegi *specimens, showing changing N_e over time and population divergence between a GB specimen (blue curve) and a European specimen (yellow curve). All A. crataegi specimens PSMC curves are shown in S4. (B): PSMC for modern* P. icarus *specimens, with a GB specimen (blue curve) and a European specimen (yellow curve). Divergence timing is shown as before the LGM due to interval settings during the PSMC run (-p). YD = Younger Dryas, LGM= Last Glacial Maximum, EEM= Eemian Interglacial Period*

Figure 3: (A): Individual heterozygosity rates in A. crataegi *specimens from Europe and GB, calculated using ANGSD. Dashed lines show heterozygosity difference between Europe and GB. Significance codes: n.s. = not significant, * = p <0.05, ** = p <0.01, *** = p < 0.001. (B): Individual heterozygosity rates in* P.icarus *specimens from Europe and GB (C): RoH in* A. crataegi *specimens from Europe and GB. Open bars show the total proportion of the genome in RoH ≥ 100 KB, light grey bars show RoH ≥ 500kb and dark grey bars show proportions in RoH ≥ 1 MB. (D): RoH in* P.icarus *European and GB specimens. Dashed lines in C and D show significance of difference between European and GB specimens.*

1143   *Figure. 4: (A-C): Total LoF, Missense and Synonymous variants found using snpEff in historic* A.

1144   crataegi *(A.c.) specimens. (D-F): Total LoF, Missense and Synonymous variants found in* P. icarus.

1145   *Dashed lines show difference between Europe and GB. Significance codes: n.s. = not significant, * =*

1146   *p <0.05, ** = p <0.01, *** = p < 0.001.* A.c*: linear model with average coverage as a covariate*, P.i.

1147   *Welch's two sample t-test.*

1148

1149   *Figure 5 (A): R_{xy} ratio of shared derived alleles classified Synonymous, Missense and LoF variants in*

1150   A. crataegi. *Red whiskers represent 95% confidence intervals based on jackknife estimates over*

1151   *chromosomes. A ratio of > 1 represents an increase in frequency of alleles in population X (GB)*

1152   *relative to population Y (Europe), 1 reflects no change and < 1 a decrease. (B) R_{xy} ratio of shared*

1153   *derived alleles of each class in* P. icarus.