# Effect of eye globe and optic nerve morphologies on gaze-induced optic nerve head deformations

Tingting Liu[1], Pham Tan Hung[2], Xiaofei Wang[1, 3*], Michaël J. A. Girard[2, 4, 5, 6*]

1.  Key Laboratory for Biomechanics and Mechanobiology of Ministry of Education, Beijing Advanced Innovation Center for Biomedical Engineering, School of Biological Science and Medical Engineering, Beihang University, Beijing, China

2.  Singapore Eye Research Institute, Singapore National Eye Centre, Singapore

3.  School of Ophthalmology and Optometry and School of Biomedical Engineering, Wenzhou Medical University, Wenzhou, China

4.  Duke-NUS Medical School, Singapore

5.  Institute for Molecular and Clinical Ophthalmology, Basel, Switzerland

6.  Department of Biomedical Engineering, College of Design and Engineering, National University of Singapore, Singapore

*These authors contributed equally to this work and are both corresponding authors

**Keywords:**            Optic Nerve Head, Eye Movements, Finite Element
                         Analysis, Glaucoma, Biomechanics

**Word count:**          4599 (Manuscript Text)
                         230 (Abstract)

**Tables:**              3

**Figures:**             6

**Corresponding Author:**    Xiaofei Wang
                             School of Biological Science and Medical Engineering
                             Beihang University
                             Room 424, Building 5
                             37 Xueyuan Road
                             Beijing 10083, China
                             xiaofei.wang@buaa.edu.cn

                             Michaël J.A. Girard
                             Ophthalmic Engineering & Innovation Laboratory (OEIL)
                             Singapore Eye Research Institute (SERI)
                             The Academia, 20 College Road
                             Discovery Tower Level 6,
                             Singapore 169856
                             mgirard@duke-nus.edu.sg

**Submission to *IOVS*:** 19/12/2023

## ABSTRACT

**Purpose**: To investigate the effect of globe and optic nerve (ON) morphologies and tissue stiffnesses on gaze-induced optic nerve head deformations using parametric finite element modeling and a Design of Experiment (DOE) approach.

**Methods**: A custom software was developed to generate finite element models of the eye using 10 morphological parameters: dural radius, scleral, choroidal, retinal, pial, peripapillary border tissue thicknesses, prelaminar tissue depth, lamina cribrosa (LC) depth, ON radius, and ON tortuosity. A 10-factor 2-level full-factorial analysis (1,024 models) was used to predict the effects of each morphological factor and their interactions on LC strains induced by 13º adduction. Subsequently, a further DOE analysis (1,024 models) was conducted to study the effects and potential interactions between the top 5 morphological parameters identified from the initial DOE study and 5 critical tissue stiffnesses.

**Results**: In the DOE analysis of 10 morphological parameters, the five most significant factors were ON tortuosity, dural radius, ON radius, scleral thickness and LC depth. Further DOE analysis incorporating biomechanical parameters highlighted the importance of dural and LC stiffness. A larger dural radius and stiffer dura increased LC strains but the other main factors had the opposite effects. Notably, a significant interaction was found between dural radius and dural stiffness.

**Conclusions**: This study highlights the significant impact of morphological factors on LC deformations during eye movements, with key morphological effects being more pronounced than tissue stiffnesses.

## INTRODUCTION

Glaucoma is one of the most common causes of blindness worldwide.[1] The biomechanical theory of glaucoma suggests that the deformations of the optic nerve (ON) head (ONH) tissues, especially the lamina cribrosa (LC), may lead to the apoptosis of retinal ganglion cells and visual field defects, either directly or indirectly.[2] Intraocular pressure (IOP) and cerebrospinal fluid pressure (CSFP) are the two main mechanical loads acting on the ONH that have been shown to be correlated to glaucoma pathogenesis[3–5]. Recent studies using finite element (FE) modeling,[6–8] optical coherence tomography [9–13] and magnetic resonance imaging (MRI)[14–16] have highlighted that ON traction during eye movements can yield large ONH deformations, which may be as large as or significantly larger than those caused by a substantial IOP elevation to 40 or 50 mmHg.

In vivo studies have shown that gaze-induced ONH deformations vary widely across individuals[17,18]. The differences are likely due to variations in the biomechanical properties and morphologies of the eye globe, ON and ONH. For instance, ON tortuosity varies across individuals, as shown in **Figure 1**. A less tortuous ON has been hypothesized to generate a larger traction force, and thus potentially larger ONH deformations.[15] Therefore, to identify those who are vulnerable to ON traction during eye movements, it would be critical to identify the biomechanical and morphological factors (and their interactions) that significantly affect ONH deformations. Using FE modeling, we have previously investigated the effects of biomechanical properties on gaze-induced ONH deformations and predicted that a stiffer dura would generate a larger ONH deformation.[6] However, the effects of eye-globe and ON

morphologies on gaze-induced ONH deformations and their potential

interactions with biomechanical properties remain unexplored.

The aim of this study was to explore the effects of eye-globe and ON

morphologies on gaze-induced ONH deformations, and to examine any

potential interactions between morphological parameters and biomechanical

parameters of tissues, using parametric FE modeling and design of experiment

(DOE).

## METHODS

In this study, we developed a methodology to automatically generate

thousands of 3D eye models to study the effects of eye-globe and ON

morphologies, as well as tissue biomechanical properties, on gaze-induced

ONH deformations. Specifically, a custom-written software (C++) was designed

to automatically generate FE models of the eye, each with a set of pre-

determined morphological and material parameters. These models were then

fed into the FE solver FEBio (Musculoskeletal Research Laboratories,

University of Utah, UT, US) to predict gaze-induced ONH deformations.

Configurations of all key factors were generated by a DOE approach. The initial

DOE analysis evaluated 10 morphological parameters to determine the top 5,

which were then combined with the 5 key tissue stiffnesses from previous

studies. This resulted in a refined set of 10 parameters, covering both

morphology and tissue stiffness, for a follow-up DOE study. Since adduction is

known to induce significant ONH deformations compared to abduction,[6,7,9] an

adduction of 13° was chosen for each model, as employed in our previous work.

The response of each model was characterized by the magnitude of the

effective strain within the LC. Below is a detailed description of the methodology.

### *Geometry and Biomechanical Properties of the Baseline FE Model*

A whole-eye FE model was established, including the sclera, choroid, prelaminar neural tissue, LC, ON, pia mater, dura mater, orbital fat-muscle complex (OFM) and orbital bone. The baseline geometric parameters of eye global tissues were set to averaged values reported in the literature, shown in **Table 1**. To maintain simplicity, we opted to combine and simulate the extraocular muscles and the orbital fat as a unified entity referred to as OFM. Only half of the eye was reconstructed because the FE model was assumed to be symmetric about a transverse plane passing through the center of the eye globe (**Figure 2**).

The baseline biomechanical properties were the same as those used in our previous studies.[6,7] Briefly, both the sclera and LC were modeled as soft tissues reinforced with collagen fibers. Those fibers can exhibit stretch-induced stiffening and they are typically distributed within a 2D plane (following a von-Mises probability distribution). The collagen fibers in the peripapillary sclera surrounding the disc were organized into a ring, while those in the peripheral sclera were organized randomly (as specified by the kf parameter in **Table 2**) and parallel to the anterior scleral surface. The collagen fibers in the LC exhibited lower anisotropy than that in the peripapillary sclera and were aligned radially, extending from the central vessel trunk to the LC insertion sites.[20] All other tissues were considered either hyperelastic or linear elastic as shown in **Table 2**. Among them, peripapillary border tissue (PBT) is the border tissue of the choroid and sclera [21]. The peripapillary choroid is separated from the prelaminar neural tissue by a collagenous layer, which constitutes the border

153  tissue of the choroid. Likewise, the scleral flange is separated from the LC by

154  the border tissue of the sclera. Since the biomechanical behavior of the PBT

155  has not yet been reported, we assumed that the PBT shared the same

156  biomechanical properties as those of the pia[22]. All soft tissues were assumed

157  to be incompressible. The orbital wall was considered a rigid body.

### *Parameterization of Morphological and Biomechanical Properties*

The morphology of the FE model was parameterized using 10 factors

representing the geometry of the eye globe, ONH, and that of the ON. These

factors were: dural radius, scleral thickness, choroidal thickness, retinal

thickness, ON radius (excluding the pia and dura), pial thickness, the thickness

of PBT,[23] prelaminar tissue depth, central LC depth, and ON tortuosity (**Figure**

**3**). Specifically, the thicknesses of the eye globe tissues (sclera, choroid, and

retina) were modified by adjusting the distance between each tissue's

boundaries and the fixed sclera-choroid interface, while maintaining the

thickness of other tissue unchanged. For the ON tissues (pia and PBT), a

similar approach was used where the inner surface of each specific tissue was

fixed and the outer surface were altered to vary its thickness. The radii of the

dura and ON were adjusted by changing their distance from the central axis of

the ON and ONH. ON tortuosity was altered by adjusting the positions of three

control points along its central path. Refer to **Supplementary Material A-1** for

more details on how the morphological parameters of the eye globe and ONH

were varied.

The biomechanical properties of each tissue were directly modified in

the input file of the FE model.

### Contact Definitions, Boundary and Loading Conditions

Contacts between tissues were the same as those defined in our previous study.[6,7] Briefly, the OFM and the dura were tied together; the OFM was able to slide over the bony orbital margin with a friction factor of 0.5; the cornea-scleral shell and the OFM had a sliding contact with no friction to mimic the Tenon's capsule enveloping the eyeball. Rigid contact was assigned between the horizontal rectus muscle insertions and a 'non-physiological' rigid body. This latter had a center of mass at the center of the eyeball, which was constrained with a prescribed rotation to simulate an adduction of 13°. For boundary conditions, the OFM and ON were fixed at the orbital apex to mimic the fibrous adhesion of those tissues to the optic canal. The orbital bone was also held in place by fixing its outer margin. In addition to an adduction of 13°, an IOP of 15 mm Hg was applied to the surface of the retinal and prelaminar tissues and a CSFP of 12.9 mm Hg was applied within the subarachnoid space of the ON. Loading was applied in two steps: first, IOP and CSFP were applied and maintained, then an adduction of 13° was applied. All contact patterns, boundaries and loading conditions are illustrated in **Figure 2**.

### FE Simulations, Post-processing and Output Measurement

All FE models were consistently meshed with 73,922 nodes and 62,821 elements, including 62,521 8-node hexahedra and 300 6-node pentahedra elements. All tissues were bonded by shared nodes at tissue boundaries (**Figure 2**). The mesh density was numerically validated through a convergence test as described in our previous study.[6] A single model required about 20 minutes to solve on a desktop workstation (Intel Xeon Silver 4114 CPU @ 2.20GHz, 32GB of memory).

The preprocessing Matlab script executed FEBio to solve the generated

models and outputted the Lagrange strain tensors for each step and the

volumes of all LC elements into a text file. Effective strains were calculated from

the principal components of the Green Lagrange strain tensor. To isolate the

specific effect of eye movement, effective strains of each LC element after the

first load step (with only IOP and CSFP) and those after the second load step

(following eye movement) were extracted. The differential, termed "delta

effective strain", was then calculated for each element. This delta effective

strain for each LC element was multiplied by its volume, and these values were

summed and divided by the total LC volume, yielding the volume-weighted

mean LC delta effective strain. For simplicity, this will be referred to as the LC

effective strain in the subsequent manuscript.

***DOE of Morphological Factors***

For the 10 morphological parameters (**Figure 3**), a two-level full factorial

design was used, resulting in a total of 1,024 models. This comprehensive

design allowed us to assess the effects of the main factors and any potential

interactions among them. The low and high levels of all morphological

parameters (excluding ON tortuosity) were set by varying them by 20% around

their baseline values (**Table 1**). For simplicity, we varied ON tortuosity within a

narrower range, from 1.013 (low level) to 1.1 (high level), as these values have

typically been observed in MRI studies. Detailed information on the

morphological parameters for each model is available in the **Supplementary**

**Material B-Sheet1.**

***DOE of Both Morphological and Biomechanical Factors***

We conducted another DOE analysis to examine the potential

interactions between morphological parameters and tissue stiffnesses. This

analysis included five key morphological parameters identified from the initial

DOE analysis and the stiffnesses of five tissues (LC, ON, sclera, pia and dura)

informed by our previous study.[6,7]

In this DOE analysis, high and low levels of the biomechanical properties

of LC, ON, sclera, pia and dura were set by varying the material constants by

20% around their baseline values (see **Table 3** for the exact values). The

variation in morphological parameters was consistent with the initial DOE

analysis. A two-level full factorial design was employed, resulting in 1,024

models. Detailed information on the morphological parameters and tissue

stiffnesses for each model is available in the **Supplementary Material B-**

**Sheet2.**

### *Statistical Analysis*

All statistical analyses were carried out in Minitab (release 20, Minitab,

LLC, Pennsylvania, USA). A P-value of less than 0.05 was considered

statistically significant.

For each model, we reported gaze-induced LC effective strain as

responses. Main effects and interaction effects were reported, and the

significance of factors was tested and ranked as detailed below. The main effect

indicates the average change in the response when a factor's level shifts from

low (512 models) to high (512 models), in which other factors vary between

both levels. Interaction effects refer to how the influence of one factor on the

response changes depending on the level of another factor. Essentially, it

examines whether the combined impact of two factors differs from the sum of their individual effects.

Analysis of Variance (ANOVA) was used to assess the significance of individual factors or interactions when they vary from low to high levels. It is important to note that in DOE, the use of ANOVA is not limited by the number of levels a factor has. Even when a factor is set at just two levels, as in this study, ANOVA remains well-suited for assessing the factor's significance on the response variable and continues to be the standard method in Minitab. This is because ANOVA focuses on partitioning the total variability in the data into components attributable to different sources, including the main effects of factors and their interactions. As ANOVA was performed individually for all factors and their possible interactions, we applied the Bonferroni correction to our p-values to account for the increased risk of Type I errors due to multiple comparisons.

In the DOE analysis, a linear relationship between the factors (and their interactions, if included) and the response is assumed and liner models were fitted with these factors as independent variables. In this context, the $R^2$ value indicates how well the linear model fits the experimental data, essentially assessing how effectively the factors explain the variation in the response. Statistically significant factors were further ranked based on the magnitude of change they induced in the response variable (equals to the absolute value of the regression coefficients), facilitating the identification of the most influential factors and interactions.

# RESULTS

## *The Effects of 10 Morphological Parameters*

The average LC effective strain (i.e., delta strain after removing the

effects of IOP and CSFP) across all models was 0.031. In the DOE analysis,

we examined 55 factors, comprising 10 main factors and 45 pairwise

interactions. Out of these, 25 factors were statistically significant ($p < 0.05$).

Among the significant factors, only five factors contribute to more than 1% of

the total effects: ON tortuosity, dural radius, ON radius, scleral thickness and

LC depth. A linear regression model showed that these five factors accounted

for 96.69% of the total effects in the responses. Details on the statistically

significant factors with less than 1% impact on the total effects are available in

**Supplementary Material B-Sheet3**.

Larger ON tortuosity, scleral thickness, ON radius and LC depth

decreased LC strains following eye movements, while a lager dural radius

increased LC strains. **Figure 4** illustrates the magnitude and trend of the effects.

**Figure 5** shows the morphology of the undeformed and deformed FE

models and color-coded strains (ONH and LC) in models with low and high

levels of these five factors. In the baseline model, the mean LC effective strain

caused by 13° adduction was 0.042. The mean LC effective strains for models

with larger ON tortuosity, dural radius, ON radius, scleral thickness and LC

depth were 0.026, 0.054, 0.026, 0.035, and 0.037, respectively.

***The Effects of 5 Morphological Parameters and 5 Tissue***

***Stiffnesses***

A total of 55 factors, including main factors and their interactions, were

examined; 39 of these were statistically significant ($p<0.05$). The factors that

individually contributed to more than 1% of the total effects, in descending order,

are ON tortuosity, dural radius, ON radius, dural stiffness, scleral thickness, LC

stiffness, LC depth, the interaction between the dural radius and the dural stiffness, together accounting for 95.52% of the total effects in the responses. For detailed information on factors that were statistically significant but had an effect of less than 1% of the total effects, refer to **Supplementary Material B-Sheet4**.

**Figure 6** illustrates the magnitude and trend of the effects. The trend for the 5 morphological parameters are consistent with the results from the above morphological DOE analysis. For biomechanical properties, a stiffer dura increased LC strains, whereas a stiffer LC reduced strains. The most pronounced interaction occurred between dural radius and dural stiffness. Specifically, an increase in LC strain associated with a larger dural radius was found to be amplified when combined with a stiffer dura.

# DISCUSSION

In this study, we developed a parametric FE model and studied the effects of eye-globe and ON morphologies, as well as tissue stiffnesses on ONH deformations during eye movements. Our models demonstrated that ON tortuosity, dural radius, ON radius, scleral thickness and LC depth were the five main morphological factors that significantly affect gaze-induced ONH deformations. These parameters retained their significance in a combined analysis with tissue stiffnesses. We also observed a significant interaction between the dural radius and dural stiffness, proving to be a considerable factor in the ONH's response to eye movement.

***A Larger Dural Radius Increased LC Strains During Eye Movements***

Our study found that a larger dural radius (i.e., the optic nerve sheath inner radius) leads to higher LC strains, similar to the effect of a stiffer dura.

During eye movements, peripapillary tissues are sheared in the transverse

plane by the optic nerve sheath, resulting in significant deformations.[7] An

increased dural radius would tend to restrict eye movements by exerting a

larger pulling force onto the ONH, as evidenced by the calculated traction force

from the FE models (**Supplementary Material A-2**). The potential for such

forces to cause axonal death in glaucoma requires further investigation.

In this study, it is important to acknowledge that variations in the dural

radius were accompanied by changes in the length of the scleral flange, as the

insertion point of the dura into the sclera was altered. Consequently, an

increase in dural radius resulted in an enlargement of the scleral flange in the

model. Although this relationship aligns with anatomical observations, where

the dural radius and scleral flange size are positively correlated,[4,24] this

confounding factor complicates the interpretation of the effect of a larger dura

radius. Specifically, a larger scleral flange could potentially indicate a weaker

ONH, which is more susceptible to deformation. To dissect the potential

confounding impacts of an increased scleral flange and dural radius, we

conducted additional simulations (**Supplementary Material A-3**). In these

models, we modified the dural and ON radius without varying the scleral flange

size. Keeping the scleral flange size constant, we observed that an increase in

dural radius (along with a concurrent increase in ON radius, which can actually

reduce LC strain) led to an increase in LC strains. This approach allowed us to

confirm that an increased dural radius indeed contributes to higher LC strains.

Computed tomography (CT) studies have shown that the optic nerve

sheath diameter (ONSD) is significantly larger in patients with normal tension

glaucoma (NTG) compared to healthy controls[25,26], suggesting that NTG eyes

may exhibited more ONH deformation due to eye movement. However, other

studies have found no significant difference in ONSD between NTG and healthy

controls,[27] and that NTG subjects may have a smaller ONSD owing to a lower

CSFP.[28,29] These conflicting findings could be due to ethnic differences or

underlying differences in the pathogenesis of various NTG subtypes. Our

previous studies demonstrated a negative correlation between IOP-induced

ONH strain and retinal sensitivity in high tension glaucoma subjects[17], whereas

NTG subjects showed a stronger correlation with gaze-induced ONH

deformation[30]. Given these results, it is quite possible that NTG itself may have

different subtypes, with some subjects being sensitive to IOP and others

sensitive to gaze. The role of morphological differences in the dura in these

variations remains unexplored, and further studies are warranted.

***A Large ON Tortuosity Decreased LC Strains During Eye Movements***

Our study suggested that increased ON tortuosity may lead to lower LC

strains during eye movements. A relatively taut ON has the potential to exhibit

rapid straightening during eye movements and thus exert more force on the

ONH tissue, a phenomenon observed in our FE models (**Supplementary**

**Material A-2**). In a previous study involving a small Chinese population, we

found that ONs in glaucoma subjects (mean IOP: 26.4 ± 4.6 mmHg) were tauter

than in normal controls (mean IOP: 15.3 ± 3.6 mmHg). This smaller ON

tortuosity in glaucoma subjects may exert more force on the ONH tissues during

eye movements, indicating a potential risk factor for glaucoma.[15] However,

another study[31] reported that ON path redundancy was greater in NTG than in

normal controls in primary gaze and abduction. The discrepancy between those

two studies may be attributed to: 1) differences in subjects (high-tension

glaucoma subjects of Chinese ethnicity versus NTG subjects of unknown

ethnicity); 2) small sample sizes; or simply; 3) differences in methods to assess

ON tortuosity.

A recent study reported that ON tortuosity in highly myopic subjects was

significantly larger than that in emmetropic controls.[16] It would seem that, in

high myopia, the 'slack' ON (i.e., increased ON tortuosity) might act as a

protective mechanism against ONH deformations. However, it is crucial to

acknowledge that high myopic subjects with more tortuous ONs might still be

susceptible to greater ONH deformations during eye movements due to other

influencing factors. For instance, in high myopic eyes, the ON-globe junction

must travel a longer distance for the same amount of eye movement compared

to normal eyes. This is due to the extreme elongation of high myopic eyes,

which can exhaust the redundancy in ON tortuosity. Additionally, the weakened

structural stiffness of the sclera and other ONH structures in high myopia could

make them more susceptible to ONH deformations.

***A Thicker Sclera Decreased LC Strains During Eye Movements***

Our study showed that scleral thickness significantly affects LC strains,

with a thicker sclera associated with lower LC strains during eye movements.

This finding aligns with other studies examining the effects of factors on IOP-

[32,33] and CSFP-[5] induced LC strains. Since other ONH tissues are relatively

compliant compared to the sclera, scleral deformation induced by eye

movement can be directly transmitted to surrounding tissues, suggesting that

eyes with a thinner sclera may be more sensitive to eye movements. In case of

high myopia, scleral thickness decreases significantly with increasing axial

length.[34] In severe cases, scleral thickness can be as low as 31% of that in

normal subjects[35], potentially leading to the development of staphylomas. The

reduced scleral thickness in high myopia could result in large LC deformations

and increasing susceptibility to ONH damage during eye movements. However,

as discussed above, ONH deformations in high myopic eyes are also affected

by axial length and ON tortuosity. Further studies are warranted to investigate

the interactions of these factors in high myopia.

***Other Factors Affecting LC Strains During Eye Movements***

Our study showed that a larger ON radius (i.e., the ON parenchyma,

excluding the dural and pial sheaths) had a protective effect, resulting in smaller

LC strains. There is a typical shear deformation due to ON traction, with clear

temporal pulling from the dura in adduction. We speculated that a larger ON

radius might possibly provide more mechanical support to the LC during ON

traction, which could potentially lead to smaller LC strains. ON radius is

associated with disc size. Previous studies have reported conflicting results

regarding the relationship between disc size and glaucoma. Some studies[2,36,37]

demonstrated that a larger optic disc size is associated with higher glaucoma

susceptibility, while other studies suggested that smaller discs with less space

for nerve fibers to travel through increase glaucoma susceptibility[38,39]. There

are also studies[40,41] found no significant correlation between the degree of ON

atrophy in glaucoma and disc size. Note that disc size measured in these

studies were not equal to the ON size posterior to LC. Direct measurement of

ON size with MRI imaging[42,43] have shown that the ON radius of glaucoma

subjects was smaller than that of normal controls, which was significantly

correlated with retinal nerve fiber layer thickness thinning and perimetric loss.

A recent study suggested that myopes also tended to have smaller ONs.[44] It is

worth noting that histologic studies showed that optic atrophy led to a smaller

retrobulbar ON[45], suggesting that ON diameter may correlate with the extent of

optic atrophy and a smaller ON diameter in glaucoma subjects might be the

consequence of retinal ganglion cells (RGC) apoptosis[31]. The link between ON

size and glaucoma needs further exploration.

Our study also revealed that a larger LC depth resulted in smaller gaze-

induced LC strains. In this study, a greater LC depth corresponds to a more

curved LC. Previous studies[46–49] showed that the LC depth and LC curvature

were significantly larger in POAG eyes than in healthy eyes. These differences

could be the consequences of glaucoma.[50] However, it remains unclear

whether an initial larger LC depth/curvature is protective or detrimental in the

development of glaucoma. Since LC morphology varies with race, sex, age and

axial length[51–54], the relationship between the LC morphology and gaze-

induced LC deformations needs to be further studied.

Lastly, our study reconfirmed the significance of tissue stiffness on gaze-

induced LC deformations. A stiffer dura increases LC strains, which is

consistent with our previous study.[6] In addition, LC stiffness also had a strong

influence on gazed-induced LC strains, where a stiffer LC resulted in a

reduction of LC strains. This is straightforward as a stiffer material will deform

less under the same loading condition. This observation is consistent with other

studies investigating LC strains induced by IOP[32,33] and CSFP[55]. However, the

effect of tissue stiffness on gazed-induced LC strains is smaller than that of the

three main morphologies (ON tortuosity, dural radius, and ON radius; **Figure 6**

and **Supplementary Material B-Sheet4**).

***The Interactions Affecting LC Strains During Eye Movements***

This study revealed significant interactions among various factors, with

the most notable being between dural radius and dural stiffness. A larger dural

radius tends to increase LC strains, an effect that is amplified with stiffer dura

and diminished when the dura is more compliant. These findings underscore

the importance of considering individual-specific characteristics such as eye

globe and ON morphologies, as well as their biomechanical properties, in

assessing the susceptibility of LC deformation during eye movements. Given

the complex and multifaceted nature of morphological and biomechanical

properties of the ONH, our parametric FE models provided an ideal platform for

studying and quantifying the main factors and their interactions in a systematic

manner to inform future experimental study design and analysis.

***Limitations.*** In this study, several limitations warrant further discussion.

First, our models only predicted acute ONH deformations during eye

movements and could not account for the long-term growth & remodeling

processes that are known to take place in ocular tissues.

Second, there are some inherent limitations in a two-level full factorial

design. As each factor has only two levels, it cannot account for the nonlinear

effect between the factor and the response. Additionally, in a full factorial design,

all possible combinations of the factors are tested, resulting in a high number

of experiments. This would increase the time and cost to conduct the study.

Considering more advanced experimental designs that account for nonlinear

effects and minimize the number of experiments can enhance the efficiency

and precision of future studies in this field.

Third, the morphological factors in our study included dural radius, but

not dural thickness. Previous computational studies and this study have shown

that a stiffer dura could significantly increase LC strains during eye movements.[6] To enhance our understandings, future studies should examine the effect of an increased dural thickness on LC deformation during eye movements.

Fourth, to rank the effects of all morphological factors, we varied these parameters by 20% from their baseline values, as proper physiologic ranges for each parameter are not known. As a result, morphological size variations from larger tissues were higher. A more precise understanding of the physiological ranges for these parameters would be valuable for more accurate assessments in future studies.

Finally, the simplified morphological properties of our models provided a reasonable approximation, allowing us to improve our understanding of ONH biomechanics during eye movements. It will be necessary to update this work as more biomechanical information on eye and orbital tissues becomes available.

***Conclusion.*** Our parametric finite element models demonstrated that ON tortuosity, dural radius, ON radius, scleral thickness and LC depth were the five most important morphological factors influencing gaze-induced ONH deformations. Additionally, the stiffnesses of dura and LC were the most important biomechanical factors influencing gaze-induced ONH deformations, and the interactions between dural radius and the dural stiffness was significant. Our study provides an ideal platform for studying and quantifying the main factors and interactions between factors to inform experimental design and analysis. Further experimental and clinical studies are needed to explore the

role of effect of individual-specific characteristics on gaze-induced ONH

deformations in ocular diseases, such as myopia and glaucoma.

## ACKNOWLEDGMENTS

Acknowledgement is made to (1) the National Natural Science

Foundation of China (12272030, 12002025), (2) the donors of the National

Glaucoma Research, a program of the BrightFocus Foundation, for support of

this research (G2021010S [MG]), (3) NMRC-LCG grant 'TAckling & Reducing

Glaucoma Blindness with Emerging Technologies (TARGET)', award ID: MOH-

OFLCG21jun-0003 [MG], (4) the "Retinal Analytics through Machine learning

aiding Physics (RAMP)" project that is supported by the National Research

Foundation, Prime Minister's Office, Singapore under its Intra-Create Thematic

Grant "Intersection Of Engineering And Health" - NRF2019-THE002-0006

awarded to the Singapore MIT Alliance for Research and Technology (SMART)

Centre [MG].

# REFERENCES

1. Kapetanakis VV, Chan MPY, Foster PJ, Cook DG, Owen CG, Rudnicka AR. Global variations and time trends in the prevalence of primary open angle glaucoma (POAG): a systematic review and meta-analysis. *Br J Ophthalmol*. 2016;100(1):86-93. doi:10.1136/bjophthalmol-2015-307223

2. Burgoyne CF, Crawford Downs J, Bellezza AJ, Francis Suh JK, Hart RT. The optic nerve head as a biomechanical structure: a new paradigm for understanding the role of IOP-related stress and strain in the pathophysiology of glaucomatous optic nerve head damage. *Progress in Retinal and Eye Research*. 2005;24(1):39-73. doi:10.1016/j.preteyeres.2004.06.001

3. Tun TA, Atalay E, Baskaran M, et al. Association of Functional Loss With the Biomechanical Response of the Optic Nerve Head to Acute Transient Intraocular Pressure Elevations. *JAMA Ophthalmol*. 2018;136(2):184. doi:10.1001/jamaophthalmol.2017.6111

4. Wang N, Xie X, Yang D, et al. Orbital Cerebrospinal Fluid Space in Glaucoma: The Beijing Intracranial and Intraocular Pressure (iCOP) Study. *Ophthalmology*. 2012;119(10):2065-2073.e1. doi:10.1016/j.ophtha.2012.03.054

5. Hua Y, Voorhees AP, Sigal IA. Cerebrospinal Fluid Pressure: Revisiting Factors Influencing Optic Nerve Head Biomechanics. *Invest Ophthalmol Vis Sci*. 2018;59(1):154. doi:10.1167/iovs.17-22488

6. Wang X, Rumpel H, Lim WEH, et al. Finite Element Analysis Predicts Large Optic Nerve Head Strains During Horizontal Eye Movements. *Invest Ophthalmol Vis Sci*. 2016;57(6):2452. doi:10.1167/iovs.15-18986

7. Wang X, Fisher LK, Milea D, Jonas JB, Girard MJA. Predictions of Optic

Nerve Traction Forces and Peripapillary Tissue Stresses Following

Horizontal Eye Movements. *Invest Ophthalmol Vis Sci*. 2017;58(4):2044.

doi:10.1167/iovs.16-21319

8. Shin A, Yoo L, Park J, Demer JL. Finite Element Biomechanics of Optic

Nerve Sheath Traction in Adduction. *Journal of Biomechanical Engineering*.

2017;139(10):101010. doi:10.1115/1.4037562

9. Wang X, Beotra MR, Tun TA, et al. In Vivo 3-Dimensional Strain Mapping

Confirms Large Optic Nerve Head Deformations Following Horizontal Eye

Movements. *Invest Ophthalmol Vis Sci*. 2016;57(13):5825.

doi:10.1167/iovs.16-20560

10. Chang MY, Shin A, Park J, et al. Deformation of Optic Nerve Head and

Peripapillary Tissues by Horizontal Duction. *American Journal of

Ophthalmology*. 2017;174:85-94. doi:10.1016/j.ajo.2016.10.001

11. Suh SY, Le A, Shin A, Park J, Demer JL. Progressive Deformation of the

Optic Nerve Head and Peripapillary Structures by Graded Horizontal

Duction. :7.

12. Sibony PA. Gaze Evoked Deformations of the Peripapillary Retina in

Papilledema and Ischemic Optic Neuropathy. *Invest Ophthalmol Vis Sci*.

2016;57(11):4979. doi:10.1167/iovs.16-19931

13. Lee WJ, Kim YJ, Kim JH, Hwang S, Shin SH, Lim HW. Changes in the

optic nerve head induced by horizontal eye movements. Hamann S, ed.

*PLoS ONE*. 2018;13(9):e0204069. doi:10.1371/journal.pone.0204069

14. Demer JL, Clark RA, Suh SY, et al. Magnetic Resonance Imaging of Optic

Nerve Traction During Adduction in Primary Open-Angle Glaucoma With

Normal Intraocular Pressure. *Invest Ophthalmol Vis Sci.* 2017;58(10):4114.

doi:10.1167/iovs.17-22093

15. Wang X, Rumpel H, Baskaran M, et al. Optic Nerve Tortuosity and Globe

Proptosis in Normal and Glaucoma Subjects: *Journal of Glaucoma*.

2019;28(8):691-696. doi:10.1097/IJG.0000000000001270

16. Wang X, Chang S, Grinband J, et al. Optic nerve tortuosity and

displacements during horizontal eye movements in healthy and highly

myopic subjects. *Br J Ophthalmol*. Published online May 26,

2021:bjophthalmol-2021-318968. doi:10.1136/bjophthalmol-2021-318968

17. Chuangsuwanich T, Tun TA, Braeu FA, et al. Differing Associations

between Optic Nerve Head Strains and Visual Field Loss in Patients with

Normal- and High-Tension Glaucoma. *Ophthalmology*. 2023;130(1):99-

110. doi:10.1016/j.ophtha.2022.08.007

18. Thanadet Chuangsuwanich, Tin A Tun, Fabian A Braeu, et al. Adduction

induces large optic nerve head deformations in subjects with normal-

tension glaucoma. *Br J Ophthalmol*. Published online April 3, 2023:bjo-

2022-322461. doi:10.1136/bjo-2022-322461

19. Le A, Chen J, Lesgart M, Gawargious BA, Suh SY, Demer JL. Age-

dependent Deformation of the Optic Nerve Head and Peripapillary Retina

by Horizontal Duction. *American Journal of Ophthalmology*. 2020;209:107-

116. doi:10.1016/j.ajo.2019.08.017

20. Zhang L, Albon J, Jones H, et al. Collagen Microstructural Factors

Influencing Optic Nerve Head Biomechanics. *Investigative Ophthalmology*

*& Visual Science*. 2015;56(3):2031-2042. doi:10.1167/iovs.14-15734

21. Wang YX, Panda-Jonas S, Jonas JB. Optic nerve head anatomy in myopia and glaucoma, including parapapillary zones alpha, beta, gamma and delta: Histology and clinical features. *Progress in Retinal and Eye Research*. 2021;83:100933. doi:10.1016/j.preteyeres.2020.100933

22. Sigal IA, Ethier CR. Biomechanics of the optic nerve head. *Experimental Eye Research*. 2009;88(4):799-807. doi:10.1016/j.exer.2009.02.003

23. Jonas JB, Holbach L, Panda-Jonas S. Peripapillary ring: histology and correlations. *Acta Ophthalmologica*. 2014;92(4):e273-e279. doi:10.1111/aos.12324

24. Liu H, Yang D, Ma T, et al. Measurement and Associations of the Optic Nerve Subarachnoid Space in Normal Tension and Primary Open-Angle Glaucoma. *American Journal of Ophthalmology*. 2018;186:128-137. doi:10.1016/j.ajo.2017.11.024

25. Jaggi GP, Miller NR, Flammer J, Weinreb RN, Remonda L, Killer HE. Optic nerve sheath diameter in normal-tension glaucoma patients. *Br J Ophthalmol*. 2012;96(1):53-56. doi:10.1136/bjo.2010.199224

26. Pircher A, Montali M, Berberat J, Remonda L, Killer HE. Relationship between the optic nerve sheath diameter and lumbar cerebrospinal fluid pressure in patients with normal tension glaucoma. *Eye*. 2017;31(9):1365-1372. doi:10.1038/eye.2017.70

27. Abegão Pinto L, Vandewalle E, Pronk A, Stalmans I. Intraocular pressure correlates with optic nerve sheath diameter in patients with normal tension glaucoma. *Graefes Arch Clin Exp Ophthalmol*. 2012;250(7):1075-1080. doi:10.1007/s00417-011-1878-3

28. Wang N, Xie X, Yang D, et al. Orbital Cerebrospinal Fluid Space in Glaucoma: The Beijing Intracranial and Intraocular Pressure (iCOP) Study. *Ophthalmology*. 2012;119(10):2065-2073.e1. doi:10.1016/j.ophtha.2012.03.054

29. Liu H, Yang D, Ma T, et al. Measurement and Associations of the Optic Nerve Subarachnoid Space in Normal Tension and Primary Open-Angle Glaucoma. *American Journal of Ophthalmology*. 2018;186:128-137. doi:10.1016/j.ajo.2017.11.024

30. Chuangsuwanich T, Tun TA, Braeu FA, et al. *Adduction Induces Large Optic Nerve Head Deformations in Subjects with Normal Tension Glaucoma*. Bioengineering; 2021. doi:10.1101/2021.08.25.457300

31. Demer JL, Clark RA, Suh SY, et al. Optic Nerve Traction During Adduction in Open Angle Glaucoma with Normal versus Elevated Intraocular Pressure. *Current Eye Research*. 2020;45(2):199-210. doi:10.1080/02713683.2019.1660371

32. Sigal IA, Yang H, Roberts MD, Burgoyne CF, Downs JC. IOP-Induced Lamina Cribrosa Displacement and Scleral Canal Expansion: An Analysis of Factor Interactions Using Parameterized Eye-Specific Models. *Invest Ophthalmol Vis Sci*. 2011;52(3):1896. doi:10.1167/iovs.10-5500

33. Sigal IA. Interactions between Geometry and Mechanical Properties on the Optic Nerve Head. *Invest Ophthalmol Vis Sci*. 2009;50(6):2785. doi:10.1167/iovs.08-3095

34. Shen L, You QS, Xu X, et al. Scleral Thickness in Chinese Eyes. *Invest Ophthalmol Vis Sci*. 2015;56(4):2720. doi:10.1167/iovs.14-15631

35. Cheng H, Singh O, Kwong K, Xiong J, Woods B, Brady T. Shape of the
myopic eye as seen with high-resolution magnetic resonance imaging.
*Optom Vis Sci.* 1992;69(9):698-701. doi:10.1097/00006324-199209000-
00005

36. Bellezza AJ, Rintalan CJ, Thompson HW, Downs JC, Hart RT, Burgoyne
CF. Deformation of the Lamina Cribrosa and Anterior Scleral Canal Wall
in Early Experimental Glaucoma. *Invest Ophthalmol Vis Sci.*
2003;44(2):623. doi:10.1167/iovs.01-1282

37. Wang X, Tun TA, Nongpiur ME, et al. Peripapillary sclera exhibits a v-
shaped configuration that is more pronounced in glaucoma eyes. *Br J*
*Ophthalmol.* Published online December 16, 2020:bjophthalmol-2020-
317900. doi:10.1136/bjophthalmol-2020-317900

38. Jonas JB, Budde WM, Panda-Jonas S. Ophthalmoscopic Evaluation of the
Optic Nerve Head. *Survey of Ophthalmology.* 1999;43(4):293-320.
doi:10.1016/S0039-6257(98)00049-6

39. Teng Y, Yu X, Teng Y, et al. Evaluation of crowded optic nerve head and
small scleral canal in intrapapillary hemorrhage with adjacent peripapillary
subretinal hemorrhage. *Graefes Arch Clin Exp Ophthalmol.*
2014;252(2):241-248. doi:10.1007/s00417-013-2459-4

40. Jonas JB, Fernández MC, Naumann GOH. Correlation of the Optic Disc
Size to Glaucoma Susceptibility. *Ophthalmology.* 1991;98(5):675-680.
doi:10.1016/S0161-6420(91)32234-6

41. Jonas JB, Sturmer J, Papastathopoulos KI, Meier-Gibbons F, Dichtl A.
Optic disc size and optic nerve damage in normal pressure glaucoma.

*British Journal of Ophthalmology*. 1995;79(12):1102-1105.

doi:10.1136/bjo.79.12.1102

42. Zhang YQ, Li J, Xu L, et al. Anterior visual pathway assessment by

magnetic resonance imaging in normal-pressure glaucoma. *Acta*

*Ophthalmologica*. 2012;90(4):e295-e302. doi:10.1111/j.1755-

3768.2011.02346.x

43. Lagre`ze WA, Gaggl M, Weigel M, et al. Retrobulbar Optic Nerve Diameter

Measured by High-Speed Magnetic Resonance Imaging as a Biomarker

for Axonal Loss in Glaucomatous Optic Atrophy. *Invest Ophthalmol Vis Sci*.

2009;50(9):4223. doi:10.1167/iovs.08-2683

44. Nguyen BN, Cleary JO, Glarin R, et al. Ultra-High Field Magnetic

Resonance Imaging of the Retrobulbar Optic Nerve, Subarachnoid Space,

and Optic Nerve Sheath in Emmetropic and Myopic Eyes. *Trans Vis Sci*

*Tech*. 2021;10(2):8. doi:10.1167/tvst.10.2.8

45. Jonas JB, Schmidt AM, Müller-Bergh JA, Naumann GOH. Optic nerve fiber

count and diameter of the retrobulbar optic nerve in normal and

glaucomatous eyes. *Graefe's Arch Clin Exp Ophthalmol*. 1995;233(7):421-

424. doi:10.1007/BF00180945

46. Lee KM, Kim TW, Lee EJ, Girard MJA, Mari JM, Weinreb RN. Association

of Corneal Hysteresis With Lamina Cribrosa Curvature in Primary Open

Angle Glaucoma.

47. Lee SH, Kim TW, Lee EJ, Girard MJA, Mari JM. Diagnostic Power of

Lamina Cribrosa Depth and Curvature in Glaucoma. *Invest Ophthalmol Vis*

*Sci*. 2017;58(2):755. doi:10.1167/iovs.16-20802

48. Kim YW, Jeoung JW, Kim DW, et al. Clinical Assessment of Lamina
Cribrosa Curvature in Eyes with Primary Open-Angle Glaucoma. Wedrich
A, ed. *PLoS ONE*. 2016;11(3):e0150260.
doi:10.1371/journal.pone.0150260

49. Ha A, Kim TJ, Girard MJA, et al. Baseline Lamina Cribrosa Curvature and
Subsequent Visual Field Progression Rate in Primary Open-Angle
Glaucoma. *Ophthalmology*. 2018;125(12):1898-1906.
doi:10.1016/j.ophtha.2018.05.017

50. Park SC, Brumm J, Furlanetto RL, et al. Lamina Cribrosa Depth in Different
Stages of Glaucoma. *Invest Ophthalmol Vis Sci*. 2015;56(3):2059.
doi:10.1167/iovs.14-15540

51. Tun TA, Wang X, Baskaran M, et al. Determinants of lamina cribrosa depth
in healthy Asian eyes: the Singapore Epidemiology Eye Study. *Br J
Ophthalmol*. Published online May 20, 2020:bjophthalmol-2020-315840.
doi:10.1136/bjophthalmol-2020-315840

52. Luo H, Yang H, Gardiner SK, et al. Factors Influencing Central Lamina
Cribrosa Depth: A Multicenter Study. *Invest Ophthalmol Vis Sci*.
2018;59(6):2357. doi:10.1167/iovs.17-23456

53. Vianna JR, Lanoe VR, Quach J, et al. Serial Changes in Lamina Cribrosa
Depth and Neuroretinal Parameters in Glaucoma. *Ophthalmology*.
2017;124(9):1392-1402. doi:10.1016/j.ophtha.2017.03.048

54. Ren R, Yang H, Gardiner SK, et al. Anterior Lamina Cribrosa Surface
Depth, Age, and Visual Field Sensitivity in the Portland Progression Project.
*Invest Ophthalmol Vis Sci*. 2014;55(3):1531. doi:10.1167/iovs.13-13382

55. Feola AJ, Myers JG, Raykin J, et al. Finite Element Modeling of Factors

Influencing Optic Nerve Head Deformation Due to Intracranial Pressure.

*Invest Ophthalmol Vis Sci*. 2016;57(4):1901. doi:10.1167/iovs.15-17573

56. Vaiman M, Abuita R, Bekerman I. Optic nerve sheath diameters in healthy

adults measured by computer tomography. *Int J Ophthalmol*.

2015;8(6):1240-1244. doi:10.3980/j.issn.2222-3959.2015.06.30

57. Norman RE, Flanagan JG, Rausch SMK, et al. Dimensions of the human

sclera: Thickness measurement and regional changes with axial length.

*Experimental Eye Research*. 2010;90(2):277-284.

doi:10.1016/j.exer.2009.11.001

58. Jiang R, Wang YX, Wei WB, Xu L, Jonas JB. Peripapillary Choroidal

Thickness in Adult Chinese: The Beijing Eye Study. *Invest Ophthalmol Vis*

*Sci*. 2015;56(6):4045. doi:10.1167/iovs.15-16521

59. Alamouti B. Retinal thickness decreases with age: an OCT study. *British*

*Journal of Ophthalmology*. 2003;87(7):899-901. doi:10.1136/bjo.87.7.899

60. Sigal IA, Flanagan JG, Tertinegg I, Ethier CR. Finite Element Modeling of

Optic Nerve Head Biomechanics. *Invest Ophthalmol Vis Sci*.

2004;45(12):4378. doi:10.1167/iovs.04-0133

61. Bowd C, Weinreb RN, Lee B, Emdadi A, Zangwill LM. Optic disk

topography after medical treatment to reduce intraocular pressure.

*American Journal of Ophthalmology*. 2000;130(3):280-286.

doi:10.1016/S0002-9394(00)00488-8

62. Jonas RA, Holbach L. Peripapillary border tissue of the choroid and

peripapillary scleral flange in human eyes. *Acta Ophthalmol*. 2020;98(1).

doi:10.1111/aos.14206

63. Girard MJA, Suh JKF, Bottlang M, Burgoyne CF, Downs JC. Scleral

Biomechanics in the Aging Monkey Eye. *Invest Ophthalmol Vis Sci*.

2009;50(11):5226. doi:10.1167/iovs.08-3363

64. Friberg TR, Lace JW. A comparison of the elastic properties of human

choroid and sclera. *Experimental Eye Research*. 1988;47(3):429-436.

doi:10.1016/0014-4835(88)90053-X

65. Miller K. Constitutive model of brain tissue suitable for finite element

analysis of surgical procedures. *Journal of Biomechanics*. 1999;32(5):531-

537. doi:10.1016/S0021-9290(99)00010-X

66. Schoemaker I, Hoefnagel PPW, Mastenbroek TJ, et al. Elasticity, Viscosity,

and Deformation of Orbital Fat. *Invest Ophthalmol Vis Sci*.

2006;47(11):4819. doi:10.1167/iovs.05-1497

# FIGURES

**Figure 1.** MRI images of the orbital region demonstrate the morphological diversity of the optic nerve (ON). These six figures show examples of ONs displaying varying degrees of curvature, ranging from straight to highly tortuous. The white lines represent the ON middle curve.

**Figure 2.** Left panel shows the reconstructed geometry and FE mesh of the eye movement model with boundary conditions and tissue connections. Right panel shows an enlarged view of the detailed ONH region (sclera, scleral fiber ring, the peripapillary border tissue, choroid, Bruch's membrane, lamina cribrosa, neural tissues, pia and dura) illustrating the IOP and CSFP applied to each model in the primary gaze position.

**Figure 3.** Input factors defining the parametric FE model geometry (only the ONH region of the entire eye is shown). See **Table 1** for the ranges of input factors. The blue dashed line represents the symmetry axis of ONH.

**Figure 4.** Ranking of the effects of morphological factors (only the five most significant factors contributing to more than 1% of the total effects were shown) on the mean effective strain of LC. A longer bar indicates a more significant effect when varying parameters from a low to a high level. Blue bars indicate positive effects (strain reduction) and red bars indicate negative effects (strain increase).

**Figure 5.** ONH deformations induced by an adduction of 13° with the five main factors (ON tortuosity, dural radius, ON radius, scleral thickness and LC depth) at their low and high levels, respectively. The enlarged views of the ONH and LC show the color-coded effective strain. In the enlarged views, "Strain" represents the total LC effective strain induced by IOP, CSFP and eye movement, and "EM effect" represents the mean LC delta effective strains after removing the effects of IOP and CSFP. Note that the LC deformations were exaggerated 5 times for illustration purposes.

**Figure 6.** (A) Ranking of the effects of morphological factors, tissue stiffnesses and their interactions (factors contributing to more than 1% of the total effects were shown) on the mean effective strain of LC. A longer bar indicates a more significant effect when varying parameter from a low to a high level. Blue bars indicate positive effects (strain reduction) and red bars indicate negative effects (strain increase). (B) The interactions between dural radius and dural stiffness. When dural stiffness is at the low level, LC effective strain increased by 0.0093 (from 0.0210 to 0.0303) with an increase in dural radius. At the high level of dural stiffness, LC effective strain increased by 0.0140 (from 0.0259 to 0.0399) with the increase of dural radius. 0, low level; 1, high level.

# TABLES

**Table 1.** Morphological factors and their ranges

| Morphological factors | Baseline | Low | High | References |
|---|---|---|---|---|
| Dural radius, mm | 2.17 | 1.74 | 2.61 | Vaiman et al.[56] |
| Scleral thickness, mm | 0.996 | 0.7968 | 1.19 | Norman et al.[57] |
| Choroidal thickness, mm | 0.134 | 0.1072 | 0.1608 | Jiang et al.[58] |
| Retinal thickness, mm | 0.249 | 0.1992 | 0.29 | Alamouti [59] |
| ON radius, mm | 1.333 | 1.0664 | 1.5996 | Sigal et al.[60] |
| Pial thickness, mm | 0.06 | 0.048 | 0.072 | Sigal et al. [60] |
| Prelaminar tissue depth, mm | 0.33 | 0.264 | 0.396 | Bowd et al.[61] |
| LC depth, mm | 0.3 | 0.24 | 0.36 | Tun et al. [51] |
| PBT thickness, mm | 0.083 | 0.0664 | 0.0996 | Jonas et al.[62] |
| ON tortuosity | 1.013 | 1.013 | 1.1 | Wang et al.[15] |

**Table 2.** Tissue biomechanical properties

| Tissue | Constitutive Model | Biomechanical Properties | References |
|---|---|---|---|
| Sclera | Mooney-Rivlin Von Mises Distributed Fibers | $c_1$ = 0.805 MPa<br>$c_3$ = 0.0127 MPa<br>$c_4$ = 1102.25<br>$k_f$ = 2 (scleral ring)<br>$k_f$ = 0 (other region of sclera)<br>$\theta_p$: preferred fiber orientations* | Girard et al.[63] |
| PBT | Yeoh model | $c_1$ = 0.1707 MPa<br>$c_2$ = 4.2109 MPa<br>$c_3$ = -4.9742 MPa | Wang et al. [6] |
| Choroid | Isotropic Elastic | Elastic modulus = 0.6 MPa<br>Poisson's ratio = 0.49 | Friberg et al.[64] |
| Retina | Isotropic Elastic | Elastic modulus = 0.03MPa<br>Poisson's ratio = 0.49 | Miller [65] |
| Lamina Cribrosa | Mooney-Rivlin Von Mises Distributed Fibers | $c_1$ = 0.05 MPa<br>$c_3$ = 0.0025 MPa<br>$c_4$ = 100<br>$k_f$ = 1<br>$\theta_p$: preferred fiber orientation§ | Zhang et al. [20] |
| Optic nerve | Isotropic Elastic | Elastic modulus = 0.03MPa<br>Poisson's ratio = 0.49 | Miller [65] |
| Pia | Yeoh model | $c_1$ = 0.1707 MPa<br>$c_2$ = 4.2109 MPa<br>$c_3$ = -4.9742 MPa | Wang et al. [6] |
| Dura | Yeoh model | $c_1$ = 0.1707 MPa<br>$c_2$ = 4.2109 MPa<br>$c_3$ = -4.9742 MPa | Wang et al.[6] |
| Fat | Isotropic Elastic | Elastic modulus = 0.027MPa<br>Poisson's ratio = 0.49 | Schoemaker et al. [66] |
| Orbit | Isotropic Elastic | Elastic modulus = 300 MPa<br>Poisson's ratio = 0.49 | Schoemaker et al. [66] |

*Collagen fibers orientations in the scleral fiber ring were aligned circumferentially around the scleral canal. Fibers in other parts of the sclera were organized randomly. Fiber orientations were assigned to these elements using a custom-written MATLAB code.

§ Collagen fibers orientations in the LC were along the radial direction, extending from the central vessel trunk to the scleral canal.

**Table 3.** Biomechanical properties of five tissues and their ranges

| Tissue | Constitutive Model | Material Constants (Baseline) | Material Constants (Low level) | Material Constants (High level) |
|---|---|---|---|---|
| Sclera | Mooney-Rivlin Von Mises Distributed Fibers | c1 = 0.805 MPa<br>c3 = 0.0127 MPa<br>c4 = 1102.25 | c1 = 0.644 MPa<br>c3 = 0.01016 MPa<br>c4 = 881.8 | c1 = 0.966 MPa<br>c3 = 0.0152 MPa<br>c4 = 1322.7 |
| Dura | Yeoh model | c1 = 0.1707 MPa<br>c2 = 4.2109 MPa<br>c3 = -4.9742 MPa | c1 = 0.13656 MPa<br>c2 = 3.36872 MPa<br>c3 = -5.96904 MPa | c1 = 0.20484 MPa<br>c2 = 5.05308 MPa<br>c3 = -3.97936 MPa |
| Pia | Yeoh model | c1 = 0.1707 MPa<br>c2 = 4.2109 MPa<br>c3 = -4.9742 MPa | c1 = 0.13656 MPa<br>c2 = 3.36872 MPa<br>c3 = -5.96904 MPa | c1 = 0.20484 MPa<br>c2 = 5.05308 MPa<br>c3 = -3.97936 MPa |
| ON | Isotropic Elastic | 0.03 MPa | 0.024 MPa | 0.036 MPa |
| Lamina Cribrosa | Mooney-Rivlin Von Mises Distributed Fibers | c1 = 0.05 MPa<br>c3 = 0.0025 MPa<br>c4 = 100 | c1 = 0.04 MPa<br>c3 = 0.002 MPa<br>c4 = 80 | c1 = 0.06 MPa<br>c3 = 0.003 MPa<br>c4 = 120 |