Regional response to light illuminance across the human hypothalamus

Islay Campbell[1#], Roya Sharifpour[1#], Jose Fermin Balda Aizpurua[1], Elise Beckers[1,2], Ilenia Paparella[1], Alexandre Berger[1,3,4], Ekaterina Koshmanova[1], Nasrin Mortazavi[1], John Read[1], Mikhail Zubkov[1], Puneet Talwar[1], Fabienne Collette[1], Siya Sherif[1], Christophe Phillips[1], Laurent Lamalle[1], Gilles Vandewalle[1]

[1]GIGA-CRC Human Imaging, University of Liège, Liège, 4000 Belgium

[2]Faculty of Health, Medicine and Life Sciences, School for Mental Health and Neuroscience, Alzheimer Centre Limburg, Maastricht University, Maastricht, 6229 ER The Netherlands.

[3]Institute of Neuroscience (IoNS), Department of Clinical Neuroscience,Université Catholique de Louvain (UCLouvain), Woluwe-Saint-Lambert, 1200 Belgium.

[4]Synergia Medical SA, Mont-Saint-Guibert, 1435 Belgium.

[#] shared first authorship

*Corresponding author: Gilles Vandewalle, GIGA-Cyclotron Research Centre-In Vivo Imaging, Bâtiment B30, 8 Allée du Six Août, University of Liège-Sart Tilman, 4000 Liège, Belgium.

Email: gilles.vandewalle@uliege.be

Author Contributions: I.C., R. S. and G.V. designed the research. I.C., R.S., J.F.B.A., E.B. and I.P. acquired the data. A.B., E.K., N.M., J.R., M.Z., P.T., F.C., S.S., C.P., and L.L. provided valuable insights while acquiring, interpreting, and discussing the data. I.C. and R.S. analysed the data supervised by G.V. I.C. and G.V. wrote the paper. All authors edited and approved the final version of the manuscript.

Competing Interest Statement: The authors declare no conflict of interest.

Abstract

Light exerts multiple non-image-forming biological effects on physiology including the stimulation of alertness and cognition. However, the subcortical circuitry underlying the stimulating impact of light is not established in humans. We used 7 Tesla functional magnetic resonance imaging to assess the impact of variations in light illuminance on the regional activity of the hypothalamus while healthy young adults (N=26; 16 women; 24.3 ± 2.9y) were completing two auditory cognitive tasks. We find that, during both the executive and emotional tasks, higher illuminance triggered an activity increase over the posterior part of the hypothalamus, which includes part of the tuberomamillary nucleus and the posterior part of the lateral hypothalamus. In contrast, increasing illuminance evoked a decrease in activity over the anterior and ventral parts of the hypothalamus, encompassing notably the suprachiasmatic nucleus and another part of the tuberomammillary nucleus. Critically, the performance of the executive task was improved under higher illuminance and was negatively correlated with the activity of the posterior hypothalamus area. These findings reveal the distinct local dynamics of different hypothalamus regions that underlie the impact of light on cognition.

Introduction

Light exerts multiple non-image-forming (NIF) biological effects that influence the quality of sleep and wakefulness, and higher illuminance is known to stimulate alertness and cognition (Campbell, Sharifpour, et al., 2023). The biological effects of light primarily rely on a subclass of retinal ganglion cells that are intrinsically photosensitive (ipRGCs) because they express the photopigment melanopsin, which is maximally sensitive to photons with wavelength ~480nm. IpRGCs combine the light signalling of rods and cones to their intrinsic photosensitivity and, collectively, the biological effects of light present a maximal sensitivity to the shorter blue wavelength of visible light (Tri & Do, 2019). IpRGCs project to multiple subcortical brain areas and their denser projections are found within the hypothalamus, particularly in nuclei involved in sleep and wakefulness regulation (Scammell et al., 2017; Tri & Do, 2019). The suprachiasmatic nucleus (SCN), which is the site of the principal circadian clock, receives the strongest inputs from ipRGC inputs, over the anterior part of the hypothalamus (Hattar et al., 2006). Other nuclei also receive ipRGC projections: the subparaventricular zone, one of the main output routes of the SCN, the ventrolateral preoptic nucleus (VLPO) and the preoptic nucleus (PON) involved in sleep initiation and also found in the anterior part of the hypothalamus; the lateral hypothalamus (LH), site of the orexinergic wake-promoting neurons and melanin-concentrating hormone sleep-promoting neurons, found in contrast over the lateral and posterior parts of the hypothalamus (Scammell et al., 2017; Tri & Do, 2019).

The brain circuitry underlying the biological effects of light has mostly been uncovered in nocturnal rodent models (Campbell, Sharifpour, et al., 2023; Tri & Do, 2019). Translation to diurnal human beings, where the later maturation of the cortex allows for complex cognitive processing (Braak & Del Tredici, 2015), remains scarce. In particular, whether hypothalamus nuclei contribute to the stimulating impact of light on cognition in humans is not established.

We addressed this question using ultra-high-field (UHF) 7 Tesla (7T) functional magnetic resonance imaging (fMRI) in healthy young adults exposed to light of various illuminance while engaged in two different auditory cognitive tasks. We find that higher illuminance increased the activity of the posterior part of the hypothalamus encompassing the mamillary bodies (MB) and parts of the LH and tuberomammillary nucleus (TMN). In contrast, higher illuminance decreased the activity over the anterior and ventral parts of the hypothalamus encapsulating notably the SCN and another part of the TMN. Critically, the pattern of modulation was consistent across the two cognitive tasks. Importantly, the performance of the complex cognitive task was improved under higher illuminance while the activity of the posterior part of the hypothalamus was correlated to task performance. The findings reveal the distinct local dynamics of different hypothalamus areas in response to changing illuminance that may contribute to light's impact on cognition.

Results

Twenty-six healthy young adults (16 women; 24.3 ± 2.9 y; **Suppl. Table 1**) completed two auditory cognitive tasks encompassing, respectively, the executive (Collette et al., 2005) and emotional (Grandjean et al., 2005) domains, while alternatively maintained in darkness or exposed to short periods (< 1 min) of light of four different illuminances (0.16, 37, 92, 190 melanopic equivalent daylight illuminance - mel EDI- lux; **Suppl. Table 2**) [**Fig.1**]. The hypothalamus of each participant was segmented into 5 subparts – inferior-anterior, superior-anterior, inferior-tubular, superior-tubular, and posterior [**Fig.2 A**] – so we could consistently extract the regional effect of illuminance change on fMRI blood-oxygen-level-dependent (BOLD) signal over most of the hypothalamus volume.

The impact of illuminance variations on the activity of the hypothalamus is not uniform

The main analyses aimed at isolating differences in the overall impact of illuminance changes

among the 5 hypothalamus subparts. For each subpart, we extracted an index of the illuminance

impact as their average regression coefficients between their responses to the tasks and the

illuminance levels. These analyses showed significant differences between the hypothalamus

subparts for the executive (generalized linear mixed models (GLMM); main effect of the

subparts; $p = 0.002$) and emotional (GLMM; main effect of the subparts; $p < 0.0001$) tasks,

revealing that, during both tasks, the variations in illuminance affected the activity of the 5

hypothalamus subparts differently [**Fig.2 B, C**; **Table 1**]. A nominal main effect of the task was

detected for the emotional task [$p = 0.049$; Table 1] but not for the n-back task. For both tasks,

there was no significant main effect for any of the other covariates and post hoc analyses showed

that the index of the illuminance impact was consistently different in the posterior hypothalamus

subpart compared to the other subparts [$p_{corrected} \leq 0.05$, **Table 1**]. Importantly, whole-brain

analyses confirmed that increasing illuminance resulted in a local increase and decrease of

activity that could be detected, respectively, over the posterior and inferior subparts of the

hypothalamus [**Fig.2 D-G**]. This shows that our results do not come from a relatively unspecific

and widespread increase in BOLD signal surrounding the hypothalamus subparts and that the

effect of light was most prominent over the posterior and inferior-anterior subparts.

Opposite dynamics between the posterior and inferior/anterior hypothalamus at higher

illuminance

This prompted us to assess the activity of the hypothalamus subparts under each illuminance to

detail the different regional activity dynamics across the hypothalamus. The statistical analyses

confirmed that the activity dynamics across illuminance levels differed between the 5 subparts

during the executive and the emotional tasks (GLMM; subparts-by-illuminance interaction; $p =$

0.041) tasks [**Fig.2 H-K**; **Table 2**]. Post hoc contrasts first considered the impact of the changes in

illuminance within each subpart (**Suppl. Tables 3, 4**). The activity of the posterior hypothalamus

subpart significantly (p <0.05) increased under the highest illuminance (190 mel EDI) compared with darkness for both tasks and with the lower illuminances (37 and 92 mel EDI lux) for the emotional task. In contrast, for both tasks, the activity in the inferior-anterior and inferior-tubular hypothalamus subparts significantly (p < 0.05) decreased under the highest illuminance compared with darkness, and with lower illuminances for the emotional task. Finally, the activity of the superior anterior hypothalamus subpart decreased under higher illuminance during the emotional but not the executive task, while the activity of the fifth hypothalamus subpart, the superior tubular subpart, was not significantly affected by illuminance changes in either task.

Post hoc analyses also yielded several significant differences between hypothalamus subparts (p < 0.05) (**Table 2**; **Suppl. Tables 5, 6**). For both tasks, the activity of the posterior hypothalamus subpart was consistently significantly higher than the activity inferior-tubular subpart under the highest illuminances (92 and 190 mel EDI lux). For the executive task, the activity of the posterior hypothalamus subpart was also significantly higher than the superior-anterior subpart under the highest illuminances (92 and 190 mel EDI lux). For the emotional task, the activity of the posterior hypothalamus subpart was also significantly higher than the superior-anterior subpart under the highest illuminances (190 mel EDI lux), while the activity superior-tubular hypothalamus subpart was significantly higher than the activity of the inferior-tubular, inferior-anterior and superior-anterior hypothalamus subparts (92 and/or190 mel EDI lux). The overall picture arising from these comparisons is that higher illuminance increased the activity of the posterior and superior hypothalamus subparts while it decreased the activity of the inferior and anterior hypothalamus subparts.

Performance to the executive task is improved by light and related to the activity of the posterior hypothalamus

Following these analyses, we explored whether the changes in activity across illuminances were related to cognitive performance. We first considered the more difficult (2-back) subtask of the

executive task as it requires higher cognitive functions (see method for a full rationale) (Collette
et al., 2005). The analysis revealed that accuracy to the executive task was high in all participants,
but accuracy to the more difficult subtask (2-back) improved with increasing illuminance (GLMM;
main effect of illuminance; F = 2.72; p = 0.034; Partial R2 215 = 0.1; **Fig.3 A**), controlling for age,
sex and BMI. Critically, the analysis also showed that performance under each illuminance was
significantly related to the activity of the posterior hypothalamus subpart (GLMM; main effect
of posterior subpart activity; F = 9.43; p = 0.0027; Partial R2 219 = 0.09). Surprisingly, the
association was negative (**Fig.3 B**), suggesting that the part of variance explained by the
hypothalamus subpart is distinct from the impact of light on performance. In contrast, no
significant association was found when considering the activity of the other four subparts
(GLMM; main effect of posterior subpart activity; F < 0.62; p > 0.4; **Fig.3 C,D**; **Suppl. Table 7**). We
went on and found that the accuracy to the simpler control subtask of the executive tasks (0-
back, see method) was not associated with the activity of the posterior hypothalamus subpart
(GLMM controlling for age, sex and BMI; main effect of subpart activity; F = 0.57; p = 0.45; **Fig.3
E**), suggesting that the association with performance is specific to the 2-back subtask.

In the last step, we explored the reaction times during the emotional task (accuracy to
the lure task is not meaningful). We found that reaction times to the emotional stimuli were not
significantly affected by illuminance (GLMM controlling for age, sex and BMI; main effect of
illuminance; F = 1.01; p = 0.41; **Fig.3 F**) and yet, they were significantly associated with the activity
of the posterior hypothalamus subpart across each illuminance (GLMM; main effect of subpart
activity; F = 4.34; p = 0.04; **Fig.3 G**). The association was positive meaning that reaction times
were longer if activity estimates were higher, which could indicate a reinforcement of the
emotional response characterized by longer reaction times (Grandjean et al., 2005). No such
significant association was detected when considering reaction times to the neutral items of the
task (GLMM controlling for age, sex and BMI; main effect of illuminance; F = 1.5; p = 0.21; main
effect of subpart activity; F = 0.28; p = 0.6; **Fig.3 H**).

<u>Discussion</u>

Animal research has established that the biological impact of light illuminance impinges on many

subcortical structures, many of which regulate sleep and wakefulness (Campbell, Sharifpour, et

al., 2023; Hattar et al., 2006; Scammell et al., 2017; Tri & Do, 2019). How these findings translate to

human beings is not established. Here, we took advantage of the relatively high resolution and

signal-to-noise ratio of UHF 7T fMRI to determine how illuminance affects the activity of the

hypothalamus as, based on animal research, it receives the densest projections from ipRGCs

(Hattar et al., 2006; Tri & Do, 2019). We find that the activity of the posterior part of the

hypothalamus increases with increasing illuminance while, in contrast, the inferior and anterior

hypothalamus show a seemingly opposite pattern and see their activity decrease under higher

illuminance. The pattern of light-induced changes was consistent across an executive and an

emotional task which consisted of block and an event-related fMRI design, respectively. This

suggests that a robust anterior-posterior gradient of activity modulation by illuminance is

present in hypothalamus across cognitive domains. Importantly, performance to the complex

cognitive task was improved under higher illuminance and was correlated to the activity of the

posterior part of the hypothalamus under the different illuminance, though negatively. The

results demonstrate that the human hypothalamus does not respond uniformly to variations in

illuminance while engaged in a cognitive challenge and suggest that the posterior part of the

hypothalamus may be key in mediating the stimulating impact of light on cognition.

The different nuclei of the hypothalamus do not have clear contrast boundaries based

on MRI signals (Billot et al., 2020). As a result, achieving nucleus resolution over the human

hypothalamus even using UFH MRI remains out of reach (Sharifpour et al., 2022). Therefore, we

cannot assign the effects we report to a specific nucleus. We can only speculate and present a

selection of plausible scenarios that would need to be tested. The posterior part of the

hypothalamus - delineated in each participant based on an automatic reproducible procedure

(Billot et al., 2020) - encompasses the MB as well as parts of the LH and the TMN. All these nuclei

could participate to the increased BOLD signal we detect under higher illuminance. The LH and

TMN, respectively, produce orexin and histamine, which are both known to promote

wakefulness, while animal histology reports direct projection of the ipRGCs to the LH (Hattar et

al., 2006; Scammell et al., 2017; Tri & Do, 2019). Orexin is a good candidate to constitute the

circadian signal that promotes wakefulness and to counter the progressive increase in sleep

needs with prolonged wakefulness (Zeitzer, 2013). Our data may therefore be compatible with

an increase in orexin release by the LH with increasing illuminance. In line with this assumption,

chemoactivation of ipRGCs lead to increase c-fos production, a marker of cellular activation, over

several nuclei of the hypothalamus, including the lateral hypothalamus (Milosavljevic et al.,

2016). If this initial effect of light we observe over the posterior part of the hypothalamus was

maintained over a longer period of exposure, this would stimulate cognition and maintain or

increase alertness (Campbell, Sharifpour, et al., 2023) and may also be part of the mechanisms

through which daytime light increases the amplitude in circadian variations of several

physiological features (Bano-Otalora et al., 2021; Dijk et al., 2012). It could then also be part of

the mechanisms through which evening light may disturb subsequent sleep (Chellappa et al.,

2013) when illuminance is higher than the recommended maximum of 10 mel EDI lux for evening

light (Brown et al., 2022). If the TMN was the hypothalamus nucleus underlying the regional

increase in the BOLD signal we report, it could confer a role to histamine in mediating the

stimulating impact of light. Of interest, the TMN receives orexin signal from the LH (Scammell et

al., 2019). Alternatively, our findings may suggest a role for the MBs in mediating the impact of

light on ongoing cognition, potentially influenced through its innervation by the TMN (Vann,

2010).

Previous research indicated that increasing illuminance reduced the activity of the

anterior part of the hypothalamus encompassing the SCN, either following the exposure to light

(Perrin et al., 2004) or during the exposure (Schoonderwoerd et al., 2022). We extend this finding

by showing that the significant decrease in activity extends beyond the inferior anterior

hypothalamus and therefore much beyond the SCN. Chemoactivation of ipRGCs in rodents led

to an increase activity of the SCN, over the inferior anterior hypothalamus, but had no impact on

the activity of the VLPO, over the superior anterior hypothalamus (Milosavljevic et al., 2016).

How our findings fit with these fine-grained observations and whether there are species-specific

differences in the responses to light over the different part of the hypothalamus remains to be

established. The inferior-tubular and inferior-anterior subparts of the hypothalamus – that we

isolated also based on an automatic reproducible procedure (Billot et al., 2020) - encompass

several nuclei such as notably the SCN, SON, ventromedial nucleus of the hypothalamus, arcuate

nucleus and part of the TMN. Again, all these nuclei may be involved in the reduction in BOLD

signal we observe at higher illuminance. In terms of chemical communication, these changes in

activity could be the results of an inhibitory signal from a subclass of ipRGCs, potentially through

the release aminobutyric acid (GABA), as a rodent study found that a subset of ipRGCs release

GABA at brain targets including the SCN (and intergeniculate leaflet and ventral lateral geniculate

nucleus), leading to a reduction in the ability of light to affect pupil size and circadian

photoentrainment (Sonoda et al., 2020). Whatever the signalling of ipRGC, our finding over the

anterior hypothalamus could correspond to a modification of GABA signalling of the SCN which

has been reported to have excitatory properties, such that the BOLD signal changes we report

may correspond to a reduction in excitation arising in part from the SCN (Albers et al., 2017).

Likewise, the SCN is also producing other neuropeptides that could affect its downstream

targets. As the inferior-tubular subpart of the hypothalamus also includes part of the TMN it may

be the TMN and its GABA production that is decreased by higher illuminance (Scammell et al.,

2019). The decrease in BOLD signal with increasing illuminance we report could therefore

arguably reflect a decreased inhibitory signal arising from the anterior and inferior nuclei of the

hypothalamus.

Importantly, none of the scenarios we elaborated on are mutually exclusive and we may

have overlooked the potential implication of several nuclei as well as the cellular diversity of the

nuclei of the hypothalamus (Adamantidis et al., 2019; Scammell et al., 2017). We further note

that the anterior-superior hypothalamus subpart of the hypothalamus encompassing the VLPO

and PON sees its activity decreasing under higher illuminance during the emotional task, similarly

to the anterior-inferior and inferior-tubular areas. Likewise, similar to the posterior subpart, the

activity of the superior-tubular hypothalamus subpart may be increased under higher

illuminance during the emotional task. Whether this represents a task-specific effect arising, for

instance, from differences in the salience of the auditory stimulus, remains to be determined.

Subcortical structures, and particularly those receiving direct retinal projections,

including those of the hypothalamus, are likely to receive light illuminance signal first before

passing on the light modulation to the cortical regions involved in the ongoing cognitive process

(Campbell, Sharifpour, et al., 2023). A critical aspect of our results is that the performance to the

2-back (executive) subtask was significantly increased when exposed to higher illuminance light.

The extent of this increase was limited, likely because performance was overall high at all

illuminances, but was not detected for the simpler detection letter subtask (0-back). The result

contrasts with many previous 3T MRI investigations on the biological effects of light on human

brain function which did not report behavioural changes induced by repeated short exposures

to light (e.g. (Vandewalle, Gais, et al., 2007; Vandewalle, Schmidt, et al., 2007; Vandewalle,

Archer, et al., 2011) but see (Daneault et al., 2018; Vandewalle et al., 2006)). Our 7T MRI study,

which includes a sample size larger than many of these previous studies, supports that BOLD

fMRI is sensitive in detecting subtle impacts of light on the brain and that these detected changes

can arguably contribute to the behavioural changes others reported using longer light exposure

and other approaches (e.g.(Cajochen et al., 2011; Lockley et al., 2006) but see (Smolders et al.,

2018)).

Importantly, we find that activity of the posterior hypothalamus subpart is negatively related to the performance to the executive task, making it unlikely that it mediates directly the positive impact of light on performance. The activity of the posterior hypothalamus was, however, associated with an increased behavioural response to emotional stimuli. The association between behaviour and the posterior hypothalamus is therefore likely to be complex and may depend on the context, with for instance different nuclei or neuronal populations contributing in some instances but not in others (Adamantidis et al., 2019; Scammell et al., 2019). It is likely also to work jointly with the decreased activity of the anterior/inferior hypothalamus we detected as well as with other non-hypothalamus subcortical structures regulating wakefulness to influence behaviour, which intrinsically primarily depends on cortical activity.

Future research may consider data-driven analyses of hypothalamus voxels time series as an alternative to the parcellation approach we adopted here. This may refine the delineation of the subparts of the hypothalamus undergoing decreased or increased activity with increasing illuminance. Future research should also assess the impact of light on other subcortical structures and on the entire subcortical network to determine how illuminance modifies their crosstalk as well as their interaction with the cortex, to eventually lead to behavioural impacts. These analyses could for instance address whether the regional changes in activity of the hypothalamus we find are upstream of the repeatedly reported impact of light illuminance on the activity of the pulvinar in the thalamus (Paparella et al., 2023; Vandewalle et al., 2006). Although it does not receive direct dense input from ipRGCs, it is likely to indirectly mediate the biological impact of light on ongoing cognitive activity (Paparella et al., 2023).

We based our rationale and part of our interpretations on ipRGC projections, which have been demonstrated in rodents to channel the NIF biological impact of light and incorporate the inputs from rods and cones with their intrinsic photosensitivity into a light signal that can impact the brain (Güler et al., 2008; Tri & Do, 2019). Given the polychromatic nature of the light we used, classical photoreceptors and their projections to visual brain areas are, however, very likely to

have directly or indirectly contributed to the modulation by light of the regional activity of the

hypothalamus. Furthermore, we cannot exclude that colour and/or spectral differences between

the orange and 3 blue-enriched light conditions may have contributed to our findings. Research

in rodent models demonstrated that variation in the spectral composition of light was perceived

by the suprachiasmatic nucleus to set circadian timing (Walmsley et al., 2015). No such

demonstration has, however, been reported yet for the acute impact of light on alertness,

attention, cognition or affective state. Future human studies could isolate the contribution of each

photoreceptor class to the impact of light on cognitive brain functions by manipulating prior light

history (Chellappa et al., 2014) or through the use of silent substitutions between metameric light

exposures (Viénot et al., 2012).

All these knowledge gaps are important to address because acting on light stands as a

promising means to reduce high sleepiness and improve cognitive deficits during wakefulness as

well as to facilitate sleep in the few hours preceding bedtime (Brown et al., 2022; Didikoglu et

al., 2023; Gooley et al., 2011; Munch et al., 2016; Riemersma-van der Lek et al., 2008;

Scheuermaier et al., 2018; Wirz-Justice et al., 2020). Light therapy is also a validated means to

improve mood and treat mood disorders (Glickman et al., 2006; Lam et al., 2016). Light

administration can also be considered a simple means to disturb the brain circuitry regulating

sleep and wakefulness such that it can provide insights about novel means to improve their

quality. For instance, if orexin and histamine were part of the mechanism through which natural

light affects brain functions, their administration may be the most ecological and/or natural

means to affect alertness and cognition. Likewise, as both orexin and histamine are targets for

the treatment of brain disorders, our findings could suggest that light may constitute a non-

pharmacological complementary intervention to compounds that are being developed to treat

arousal, sleep, or cognitive dysfunction in brain disorders (Ma et al., 2018). It remains, however,

premature in our view to base recommendations on the therapeutic use of light based on the

MRI findings gathered to date. Targeted lighting for interventions or for precise interference of

subcortical circuits will require a full understanding of how light affects the brain, particularly at

the subcortical level. Our findings represent an important step towards this goal, at the level of

the hypothalamus.

Materials and Methods

The data used in this paper arise from a large study that is leading to several publications and

part of the methods have been published previously (Beckers et al., 2023; Campbell, Beckers, et

al., 2023; Paparella et al., 2023). The protocol was approved by the Ethics Committee of the

Faculty of Medicine at the University of Liège. Participants gave their written informed consent

to take part in the study and received monetary compensation for their participation.

Participants. Thirty healthy young adults (19 women; 24.3 ± 2.9 y; **Suppl. Table S1**) were included

in the analyses. Exclusion criteria were assessed through questionnaires and a semi-structured

interview: history of psychiatric and neurological disorders, sleep disorders, use of psychoactive

drugs or addiction; history of ophthalmic disorders or auditory impairments; colour blindness;

night shift work during the last year or recent trans-meridian travel during the last 2 months;

excessive caffeine (>4 caffeine units/day) or alcohol consumption (>14 alcohol units/week);

medication affecting the central nervous system; smoking; pregnancy or breast-feeding

(women); counter indication for MRI-scanning. All participants had to score < 18 on the 21-item

Beck Anxiety Inventory (up to mild anxiety) (Beck, Epstein, et al., 1988), and <14 on the Beck

Depression Inventory-II (up to mild depression) (Beck, Steer, et al., 1988), < 12 on the Epworth

Sleepiness Scale (Johns, 1993), and < 8 on the Pittsburgh Sleep Quality Index (Buysse et al., 1989).

Questionnaires further assessed chronotype with the Horne-Östberg questionnaire (Horne &

Ostberg, 1976) and seasonality with the Seasonal Pattern Assessment Questionnaire (Rosenthal

& Bradt, 1984), but the latter two questionnaires were not used for the inclusion of the

participants.

For each task, 4 datasets were missing or had corrupt data such that 26 participants were included in the analyses of each task (23 participants had valid datasets for both tasks). For the emotional task, two participants' data failed the MRI quality control (QC) check, and the other two participants were excluded as they did not complete the entire task. For the executive task, four of the participants' data failed the MRI QC check. **Suppl. Table S2** summarises participants' characteristics respective to each task.

Overall Protocol. Participants completed an MRI session at least one week before the experiment during which structural images of the brain were acquired and which served as habituation to the experimental conditions. Participants maintained a loose sleep-wake schedule (± 1h from the habitual sleep/wake-up time) during the 7 days preceding the fMRI experiment to warrant similar circadian entrainment across participants and avoid excessive sleep loss while maintaining realistic real-life conditions (verified using sleep diaries and wrist actigraphy - AX3 accelerometer, Axivity, United Kingdom). Volunteers were requested to refrain from all caffeine and alcohol-containing beverages, and extreme physical activity for 3 days before participating in the fMRI acquisitions. Data acquisitions took place in Liège, Belgium, between December 2020 and May 2023.

Participants arrived at the laboratory 1.5 to 2h after habitual wake time for the fMRI scan. They were first exposed for 5 min to a bright polychromatic white light (1000 lux) and then maintained in dim light (< 10 lux) for 45 min to standardise the participant's recent light history. During this period participants were given instructions about the fMRI cognitive tasks and completed practice tasks on a luminance-controlled laptop (< 10 lux). The fMRI session consisted of participants completing three auditory cognitive tasks while alternatively maintained in darkness or exposed to light: an executive task (25 min), an emotional task (20 min) and an attentional task (15 min) [**Fig.1 A**]. The executive task was always completed first, as it was the most demanding task. The order of the following two tasks was counterbalanced. Because it included only 3 light conditions (see below) instead of 5 for the other two tasks, the attentional

task was not included in the present analyses. An eye-tracking system (EyeLink 1000Plus, SR

Research, Ottawa, Canada) was monitored for proper eye opening during all data acquisitions.

Light exposure. An 8-m long MRI-compatible optic fibre (1-inch diameter, Setra Systems, MA,

USA) transmitted light from a light box (SugarCUBE, Ushio America, CA, USA) to the dual end of

the fibre which was attached to a stand fitted at the back of the MRI coil that allowed

reproducible fixation and orientation of the optic fibre ends. The dual branches illuminated the

inner walls of the head coil to ensure relatively uniform and indirect illumination of participants'

eyes. A filter wheel (Spectral Products, AB300, NM, USA) and optical fibre filters (monochromatic

narrowband orange filter - 589mn; full width at half maximum: 10 nm - or a UV highpass filter -

433–1650nm) were used to create the light conditions needed for the experiment (see **Fig.1 B**

and Suppl. Table 2 for in-detail light characteristics).

Illuminance and spectra could not be directly measured within the MRI scanner due to

the ferromagnetic nature of measurement systems. The coil of the MRI and the light stand,

together with the lighting system were therefore placed outside of the MR room to reproduce

the experimental conditions of the in a completely dark room. A sensor was placed 2 cm away

from the mirror of the coil that is mounted at eye level, i.e. where the eye of the first author of

the paper would be positioned, to measure illuminance and spectra. The procedure was

repeated 4 times for illuminance and twice for spectra and measurements were averaged. This

procedure does not take into account inter-individual variation in head size and orbit shape such

that the reported illuminance levels may have varied slightly across subjects. The relative

differences between illuminance are, however, very unlikely to vary substantially across

participants such that statistics consisting of tests for the impact of relative differences in

illuminance were not affected. The detailed values reported in Supplementary Table 2 were

computed combining spectra and illuminance using the excel calculator associated with a

published work (Lucas et al., 2014).

Blue-enriched light illuminances were set according to the technical characteristics of

the light source and to keep the overall photon flux similar to prior 3T MRI studies of our team

(between ~10^{12} and 10^{14} ph/cm²/s) (Vandewalle et al., 2010; Vandewalle, Hébert, et al., 2011).

The orange light was introduced as a control visual stimulation for potential secondary whole-

brain analyses. For the present region of interest analyses, we discarded colour differences

between the light conditions and only considered illuminance as indexed by mel EDI lux. This

constitutes a limitation of our study as it does not allow attributing the findings to a particular

photoreceptor class.

For the executive and emotional task, the light conditions consisted of three different

illuminance of a white, blue-enriched polychromatic LED light (37, 92, 190 mel EDI lux; 6500K)

and one illuminance level of monochromatic orange light (.16 mel EDI lux; 590nm full width at

half maximum - FWHM: 10 nm). For the present analyses, we discarded colour differences

between the light conditions and only considered illuminance as indexed by mel EDI lux,

constituting a limitation of our study. In the executive task, participants were exposed to 30s to

70s (median 30s) of light blocks separated by 10s of darkness (< 0.1 lux) and the light blocks were

repeated 11 times for each light condition. For the emotional task, participants were exposed to

30 to 40s (median 35s) light blocks separated by 20s of darkness (< 0.1 lux) and the light blocks

were repeated five times for each light condition.

The attentional task only included a single illuminance level of the blue-enriched

polychromatic LED light (92 mel EDI lux) and one illuminance level of the monochromatic orange

light (.16 mel EDI lux), otherwise, the task would have been too long (> 30 min). Participants were

exposed to 30s of light blocks separated by 10s of darkness (<0.1 lux). The light blocks were

repeated 7 times for each light condition. As mentioned above it is not considered for the present

analyses.

Cognitive Tasks

Prior work of our team showed that the n-back task and emotional task included in the present

protocol were successful probes to demonstrate that light illuminance modulates cognitive

activity, including within subcortical structures (though resolution did not allow precise isolation

of nuclei or subparts) (e.g. (Vandewalle et al., 2010; Vandewalle, Schmidt, et al., 2007)). When

taking the step of ultra-high-field imaging, we therefore opted for these tasks as our goal was to

show that illuminance affects brain activity across cognitive domains while not testing for task-

specific aspects of these domains.

Auditory cognitive tasks. The tasks were programmed with Opensesame (3.2.8) (Mathôt

et al., 2012). Participants heard the auditory stimuli through MR-compatible earbuds

(Sensimetrics, Malden, MA). Before starting the tasks, to ensure optimal auditory perception of

task stimuli, participants set the volume through a volume check procedure. Participants used

an MRI-compatible keypad to respond to task items (Current Designs, Philadelphia, PA), which

was placed in the participant's dominant hand. The tasks were separated by about 5 minutes in

near darkness, to recalibrate the eye tracking system and to clarify instructions about the next

task to the participant.

Executive task. The task consisted of an auditory variant of the n-back task (Collette et al., 2005)

with a working memory 2-back task and a control letter detection 0-back task. Participants were

either asked to detect whether the current item was identical to the letter presented 2-items

earlier (2-back) or whether the current item consisted of the letter "K" (0-back) or using the

keypad (one button for "yes", one button for "no"). A block design was used for this task in which

each block included 15 items and lasted 30s. Task blocks were separated by 10-20s rest periods

and were preceded by an auditory instruction (500 ms) indicating the type of task to be

completed. Task levels were pseudo-randomised across the 4 light conditions with 3 blocks of 0-

back and 4 blocks of 2-back per light condition (see **Fig.1 C**).

Emotional Task. The task consisted of gender discrimination of auditory vocalizations that were

either pronounced with emotional or neutral prosody (Grandjean et al., 2005). Participants were

asked to use the keypad to indicate what they believed the gender of the person pronouncing

each token was. The gender classification was a lure task ensuring participants paid attention to

the auditory stimulation. The purpose of the task was to trigger an emotional response as

participants were not told that part of the stimuli was pronounced with angry prosody. The 240

auditory stimuli were pronounced by professional actors (50% women) and consisted of three

meaningless words ("goster", "niuvenci", "figotleich"). The stimuli were expressed in either an

angry or neutral prosody, which has been validated by behavioural assessments (Banse & Scherer,

1996) and in previous experiments (Grandjean et al., 2005; Sander et al., 2005; Vandewalle et al.,

2010). The stimuli were also matched for the duration (750 ms) and mean acoustic energy to

avoid loudness effects. During each 30 to 40-s light block, four angry prosody stimuli and four

neutral prosody stimuli were presented in a pseudorandom order and delivered every 3 to 5

seconds. A total of 160 distinct voice stimuli (50% angry; 50% neutral) were distributed across

the four light conditions. The darkness period separating each light block contained two angry

and two neutral stimuli. A total of 80 distinct voice stimuli (50% angry; 50% neutral) were

distributed across the darkness periods [see **Fig.1 D**].

Data acquisition

The MRI data were acquired in a 7T MAGNETOM Terra MR scanner (Siemens Healthineers,

Erlangen, Germany) with a 32-channel receive and 1-channel transmit head coil (Nova Medical,

Wilmington, MA, USA). Dielectric pads (Multiwave Imaging, Marseille, France) were placed

between the subject's head and receiver coil to homogenize the magnetic field of Radio

Frequency (RF) pulses.

Multislice T2*-weighted fMRI images were obtained with a multi-band Gradient-

Recalled Echo - Echo-Planar Imaging (GRE-EPI) sequence using axial slice orientation (TR = 2340

ms, TE = 24 ms, FA = 90°, no interslice gap, in-plane FoV = 224 mm × 224 mm, matrix size = 160

× 160 × 86, voxel size = $1.4 \times 1.4 \times 1.4$ mm^3). To avoid saturation effects, the first three scans

were discarded. To correct for physiological noise in the fMRI data the participants' pulse and

respiration movements were recorded using a pulse oximeter and a breathing belt (Siemens

Healthineers, Erlangen, Germany). Following the fMRI acquisition a 2D GRE field mapping

sequence to assess B0 magnetic field inhomogeneity with the following parameters: TR = 5.2 ms,

TEs = 2.26 ms and 3.28 ms, FA = 15°, bandwidth = 737 Hz/pixel, matrix size = 96 × 128, 96 axial

slices, voxel size = (2x2x2) mm^3, acquisition time = 1:38 min, was applied.

For the anatomical image, a high-resolution T1-weighted image was acquired using a

Magnetization-Prepared with 2 RApid Gradient Echoes (MP2RAGE) sequence: TR = 4300 ms, TE

= 1.98 ms, FA = 5°/6°, TI = 940ms/2830 ms, bandwidth = 240 Hz, matrix size = 256x256, 224 axial

slices, acceleration factor = 3, voxel size = (0.75x0.75x0.75) mm^3.

Data Processing

For the MP2RAGE images, the background noise was removed using an extension (extension:

https://github.com/benoitberanger/mp2rage) of Statistical Parametric Mapping 12 (SPM12;

https://www.fil.ion.ucl.ac.uk/spm/software/spm12/) under Matlab R2019 (MathWorks, Natick,

Massachusetts) (O'Brien et al., 2014). Then the images were reoriented using the

'spm_auto_reorient'   function

(https://github.com/CyclotronResearchCentre/spm_auto_reorient) and corrected for intensity

non-uniformity using the bias correction method implemented in the SPM12 "unified

segmentation" tool (Ashburner & Friston, 2005). To ensure optimal co-registration, brain

extraction   was   done   using   SynthStrip   (Hoopes   et   al.,   2022)   in   Freesurfer

(http://surfer.nmr.mgh.harvard.edu/). The brain-extracted T1-images were used to create a T1-

weighted   group   template   using   Advanced   Normalization   Tools   (ANTs,

http://stnava.github.io/ANTs/) prior to normalization to the Montreal Neurological Institute

(MNI) space using ANTs (1mm³ voxel; MNI 152 template). The hypothalamus of each participant

was segmented within 1mm³ MNI 152 template into 5 subparts - inferior anterior, superior

anterior, inferior tubular, superior tubular, posterior [cf. **Fig.2 A**] using an automatic

computational approach (Billot et al., 2020).

For the EPI images, auto reorientation was applied on the images first. Then, voxel-

displacement maps were computed from the phase and magnitude images associated with B0

map acquisition (taken right after the task), using the SPM fieldmap toolbox. To correct for head

motion and static and dynamic susceptibility-induced variance, the "Realign & Unwarp" of

SPM12 was then applied to the EPI images. The realigned and distortion-corrected EPI images

then underwent brain extraction using the SynthStrip and then the final images were smoothed

with a Gaussian kernel characterized by a FWHM = 3mm. The first level analyses were performed

in the native space to prevent any possible error that may be caused by co-registration.

Statistical analyses

The whole-brain univariate analyses consisted of a general linear model (GLM) computed with

SPM12. For the executive task, task blocks and light blocks were modelled as block functions. For

the emotional task, the auditory stimuli were modelled as stick functions. For both tasks, a high-

pass filter with a 256s cut-off was applied to remove low-frequency drifts. For both tasks, stick

or block functions were convolved with the canonical hemodynamic response function.

Movement and physiological parameters (cardiac, and respiration), which were computed with

the PhysIO Toolbox (Translational Neuromodeling Unit, ETH Zurich, Switzerland), were included

as covariates of no interest (Kasper et al., 2017).

Two separate analyses were completed. In the main analyses, we sought to test whether

brain responses during the tasks were modulated by overall changes in illuminance level. The

regressors of task blocks or events were accompanied by a single parametric modulation

regressor corresponding to the light melanopic illuminance level (.16, 37, 92, 190 mel EDI). The

contrasts of interest consisted of the main effects of the parametric modulation. In the

subsequent post hoc analysis, we estimated the responses to the stimuli under each light

condition. Separate regressors modelled each task's block or event type under each light

condition (0, 0.16, 37, 92, 190 mel EDI). The contrasts of interest consisted of the main effects of

each regressor.

The output masks of the segmentation procedure we used to extract regression betas associated with each of the hypothalamus subparts using the REX Toolbox (https://web.mit.edu/swg/software.htm) (Duff et al., 2007). Betas were averaged (mean) within each subpart and then across the homologous subparts of each hemisphere. In the main analyses this yielded 1 activity estimate per stimulus type and per hypothalamus subpart (i.e. 10 per individual), while in the subsequent analyses, we obtained 5 activity estimates per stimulus type and per subpart (50 per individual).

For visualization of whole-brain results over the entire sample, all statistical maps obtained from the first level analysis were first transferred to the group template space and then the MNI space ($1x1x1mm^3$ image resolution). All the registration steps were performed with ANTs. The visualisation was focused on the hypothalamus regions to assess whether increasing illuminance resulted in a local increase and/ or decrease of beta estimates within the hypothalamus or whether beta estimates were mainly influenced by a relatively unspecific and widespread increase in BOLD signal surrounding the hypothalamus.

Statistical analyses of the activity of the hypothalamus subparts were performed in SAS 9.4 (SAS Institute, NC, USA). Analyses consisted of Generalised Linear Mixed Models (GLMM) with the subject as a random factor (intercept and slope) and were adjusted for the dependent variable distribution. As the main statistical analysis was completed for each task, the significance threshold was corrected for multiple comparisons and was set at $p < 0.025$. Direct post hoc of the main analyses were corrected for multiple comparisons using a Tukey adjustment. The subsequent more detailed analyses were considered as post hoc that were not corrected for multiple comparisons ($p < 0.05$). To detect outlier values within the data sets, Cook's distance > 1 was used for exclusion. No outliers were detected for activity estimates of both tasks, while four outlier values were removed from the analyses of the 2-back and 0-back performance.

The main analyses included the activity estimates modulated by light illuminance as a
dependent variable and the hypothalamus subpart and stimulus type (2-back/0-back -
neutral/emotional) as repeated measures (autoregressive (1) correlation), together with age, sex
and BMI as covariates. The second set of post hoc GLMM analyses included the activity estimates
of the hypothalamus subparts as the dependent variable and hypothalamus subpart, stimulus
type and illuminance (0, 0.16, 37, 92, 190 mel EDI lux) as the repeated measures (autoregressive
(1) correlation), together with age, sex, and BMI as covariates and interaction term between
illuminance and hypothalamus subpart. The final set of analyses included performance metrics
as dependent variables (accuracy to the 2-back or 0-back task - as percentage of correct
responses; reaction time – ms - to emotional or neutral stimuli during the emotional task) and
included the same repeated measures and covariates ads in the preceding set as well as activity
of the relevant hypothalamus subpart.

Optimal sensitivity and power analyses in GLMMs remain under investigation (e.g. (Kain
et al., 2015)). We nevertheless computed a prior sensitivity analysis to get an indication of the
minimum detectable effect size in our main analyses given our sample size. According to
G*Power 3 (version 3.1.9.4) (Erdfelder et al., 2009), taking into account a power of 0.8, an error
rate α of 0.025 (correcting for 2 tasks), and a sample of 26 allowed us to detect large effect sizes
r > 0.54 (two-sided; absolute values; CI: 0.19–0.77; R^2 > 0.29, R^2 CI: 0.04–0.59) within a multiple
linear regression framework including one tested predictor (illuminance effect) and three
covariates (age, sex and BMI).

Data and code availability statement
The processed data and analysis scripts supporting the results included in this manuscript are
publicly   available   via   the   following   open   repository:
https://gitlab.uliege.be/CyclotronResearchCentre/Public/xxxx (the repository will be created
with the version of records).

Acknowledgements

The study was conducted at the GIGA-In Vivo Imaging technological platform of ULiège, Belgium. The authors thank Christine Bastin, Annick Claes, Christian Degueldre, Catherine Hagelstein, Gregory Hammad, Brigitte Herbillon, Patrick Hawotte, Sophie Laloux, Erik Lambot, Benjamin Lauricella, André Luxen, Pierre Maquet, and Eric Salmon for their help over the different steps of the study.

It was supported by the Belgian Fonds National de la Recherche Scientifique (FNRS; CDR J.0222.20), the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 860613, the Fondation Léon Frédéricq, ULiège - U. Maastricht Imaging Valley, ULiège-Valeo Innovation Chair "Health and Well-Being in Transport" and Sanfran (LIGHT-CABIN), the European Regional Development Fund (Biomed-Hub), and Siemens. None of these funding sources had any impact on the design of the study nor the interpretation of the findings. AB is supported by Synergia Medial SA and the Walloon Region (Industrial Doctorate Program, convention n°8193). EB is supported by the Maastricht University - Liège University Imaging Valley. RS and FB are supported by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 860613. IC, EK, IP, NM, and GV are supported by the FRS-FNRS. SS was supported by ULiège-Valeo Innovation Chair and Siemens Healthineers. PT and LL are supported by the EU Joint Programme Neurodegenerative Disease Research (JPND) IRONSLEEP and SCAIFIELD projects, respectively – FNRS references: PINT-MULTI R.8011.21 & 8006.20).

References

Adamantidis, A. R., Gutierrez Herrera, C., & Gent, T. C. (2019). Oscillating circuitries in the sleeping brain. In *Nature Reviews Neuroscience* (Vol. 20, Issue 12, pp. 746–762). Nature Publishing Group. https://doi.org/10.1038/s41583-019-0223-4

Albers, H. E., Walton, J. C., Gamble, K. L., McNeill, J. K., & Hummer, D. L. (2017). The dynamics of GABA signaling: Revelations from the circadian pacemaker in the suprachiasmatic nucleus. *Frontiers in Neuroendocrinology*, *44*, 35–82. https://doi.org/10.1016/J.YFRNE.2016.11.003

Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *NeuroImage*, *26*(3), 839–851. https://doi.org/10.1016/J.NEUROIMAGE.2005.02.018

Bano-Otalora, B., Martial, F., Harding, C., Bechtold, D. A., Allen, A. E., Brown, T. M., Belle, M. D. C., & Lucas, R. J. (2021). Bright daytime light enhances circadian amplitude in a diurnal mammal. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(22), e2100094118. https://doi.org/10.1073/PNAS.2100094118/SUPPL_FILE/PNAS.2100094118.SAPP.PDF

Banse, R., & Scherer, K. R. (1996). Acoustic profiles in vocal emotion expression. *Journal of Personality and Social Psychology*, *70*(3), 614–636. https://doi.org/10.1037//0022-3514.70.3.614

Beck, A. T., Epstein, N., Brown, G., & Steer, R. A. (1988). An inventory for measuring clinical anxiety: Psychometric properties. *Journal of Consulting and Clinical Psychology*, *56*(6), 893–897. https://doi.org/10.1037/0022-006X.56.6.893

Beck, A. T., Steer, R. A., & Carbin, M. G. (1988). Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clinical Psychology Review*, *8*(1), 77–100. https://doi.org/10.1016/0272-7358(88)90050-5

Beckers, E., Campbell, I., Sharifpour, R., Paparella, I., Berger, A., Aizpurua, J. F. B., Koshmanova, E., Mortazavi, N., Talwar, P., Sherif, S., Jacobs, H. I. L., & Vandewalle, G. (2023). Impact of repeated short light exposures on sustained pupil responses in an fMRI environment. *Journal of Sleep Research*, e14085. https://doi.org/10.1111/JSR.14085

Billot, B., Bocchetta, M., Todd, E., Dalca, A. V., Rohrer, J. D., & Iglesias, J. E. (2020). Automated segmentation of the hypothalamus and associated subunits in brain MRI. *NeuroImage*, *223*, 117287. https://doi.org/10.1016/J.NEUROIMAGE.2020.117287

Braak, H., & Del Tredici, K. (2015). The preclinical phase of the pathological process underlying sporadic Alzheimer's disease. *Brain : A Journal of Neurology*, *138*(Pt 10), 2814–2833. https://doi.org/10.1093/BRAIN/AWV236

Brown, T. M., Brainard, G. C., Cajochen, C., Czeisler, C. A., Hanifin, J. P., Lockley, S. W., Lucas, R. J., Münch, M., O'Hagan, J. B., Peirson, S. N., Price, L. L. A., Roenneberg, T., Schlangen, L. J. M., Skene, D. J., Spitschan, M., Vetter, C., Zee, P. C., & Wright, K. P. (2022). Recommendations for daytime, evening, and nighttime indoor light exposure to best support physiology, sleep, and wakefulness in healthy adults. *PLoS Biology*, *20*(3), e3001571. https://doi.org/10.1371/JOURNAL.PBIO.3001571

Buysse, D. J., Reynolds, C. F. 3rd, Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Research*, *28*(2), 193–213. https://doi.org/10.1016/0165-1781(89)90047-4

Cajochen, C., Frey, S., Anders, D., Späti, J., Bues, M., Pross, A., Mager, R., Wirz-Justice, A., & Stefani, O. (2011). Evening exposure to a light-emitting diodes (LED)-backlit computer screen affects circadian physiology and cognitive performance. *Journal of Applied Physiology (Bethesda, Md. : 1985)*, *110*(5), 1432–1438. https://doi.org/10.1152/JAPPLPHYSIOL.00165.2011

Campbell, I., Beckers, E., Sharifpour, R., Berger, A., Paparella, I., Aizpurua, J. F. B., Koshmanova, E., Mortazavi, N., Sherif, S., & Vandewalle, G. (2023). Impact of light on task-evoked pupil responses during cognitive tasks. *Journal of Sleep Research*, e14101. https://doi.org/10.1111/JSR.14101

Campbell, I., Sharifpour, R., & Vandewalle, G. (2023). Light as a Modulator of Non-Image-Forming Brain Functions—Positive and Negative Impacts of Increasing Light Availability. *Clocks & Sleep*, *5*(1), 116. https://doi.org/10.3390/CLOCKSSLEEP5010012

Chellappa, S. L., Ly, J. Q. M., Meyer, C., Balteau, E., Degueldre, C., Luxen, A., Phillips, C., Cooper, H. M., & Vandewalle, G. (2014). Photic memory for executive brain responses. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(16), 6087–6091. https://doi.org/10.1073/pnas.1320005111

Chellappa, S. L., Steiner, R., Oelhafen, P., Lang, D., Götz, T., Krebs, J., & Cajochen, C. (2013). Acute exposure to evening blue-enriched light impacts on human sleep. *Journal of Sleep Research*, *22*(5), 573–580. https://doi.org/10.1111/jsr.12050

Collette, F., Van Der Linden, M., Laureys, S., Delfiore, G., Degueldre, C., Luxen, A., & Salmon, E. (2005). Exploring the unity and diversity of the neural substrates of executive functioning. *Human Brain Mapping*, *25*(4), 409–423. https://doi.org/10.1002/hbm.20118

Daneault, V., Dumont, M., Massé, É., Forcier, P., Boré, A., Lina, J. M., Doyon, J., Vandewalle, G., & Carrier, J. (2018). Plasticity in the sensitivity to light in aging: Decreased non-visual impact of light on cognitive brain activity in older individuals but no impact of lens replacement. *Frontiers in Physiology, 9*(NOV). https://doi.org/10.3389/fphys.2018.01557

Didikoglu, A., Mohammadian, N., Johnson, S., van Tongeren, M., Wright, P., Casson, A. J., Brown, T. M., & Lucas, R. J. (2023). Associations between light exposure and sleep timing and sleepiness while awake in a sample of UK adults in everyday life. *Proceedings of the National Academy of Sciences of the United States of America*, *120*(42). https://doi.org/10.1073/PNAS.2301608120

Dijk, D. J., Duffy, J. F., Silva, E. J., Shanahan, T. L., Boivin, D. B., & Czeisler, C. A. (2012). Amplitude reduction and phase shifts of melatonin, cortisol and other circadian rhythms after a gradual advance of sleep and light exposure in humans. *PloS One*, *7*(2). https://doi.org/10.1371/JOURNAL.PONE.0030037

Duff, E. P., Cunnington, R., & Egan, G. F. (2007). REX: Response exploration for neuroimaging datasets. *Neuroinformatics*, *5*(4), 223–234. https://doi.org/10.1007/S12021-007-9001-Y/METRICS

Erdfelder, E., FAul, F., Buchner, A., & Lang, A. G. (2009). Statistical power analyses using G*Power 3.1: tests for correlation and regression analyses. *Behavior Research Methods*, *41*(4), 1149–1160. https://doi.org/10.3758/BRM.41.4.1149

Glickman, G., Byrne, B., Pineda, C., Hauck, W. W., & Brainard, G. C. (2006). Light therapy for Seasonal Affective Disorder with blue narrow-band light-emitting diodes (LEDs). *Biological Psychiatry*, *59*(6), 502–507. https://doi.org/10.1016/j.biopsych.2005.07.006

Gooley, J. J., Chamberlain, K., Smith, K. A., Khalsa, S. B. S., Rajaratnam, S. M. W., Van Reen, E., Zeitzer, J. M., Czeisler, C. A., & Lockley, S. W. (2011). Exposure to room light before bedtime suppresses melatonin onset and shortens melatonin duration in humans. *The Journal of Clinical Endocrinology and Metabolism*, *96*(3), E463-72. https://doi.org/10.1210/jc.2010-2098

Grandjean, D., Sander, D., Pourtois, G., Schwartz, S., Seghier, M. L., Scherer, K. R., & Vuilleumier, P. (2005). The voices of wrath: Brain responses to angry prosody in meaningless speech. *Nature Neuroscience*, *8*(2), 145–146. https://doi.org/10.1038/nn1392

Güler, A. D., Ecker, J. L., Lall, G. S., Haq, S., Altimus, C. M., Liao, H. W., Barnard, A. R., Cahill, H., Badea, T. C., Zhao, H., Hankins, M. W., Berson, D. M., Lucas, R. J., Yau, K. W., & Hattar, S. (2008). Melanopsin cells are the principal conduits for rod-cone input to non-image-forming vision. *Nature*, *453*(7191), 102–105. https://doi.org/10.1038/nature06829

Hattar, S., Kumar, M., Park, A., Tong, P., Tung, J., Yau, K. W., & Berson, D. M. (2006). Central projections of melanopsin-expressing retinal ganglion cells in the mouse. *Journal of Comparative Neurology*, *497*(3), 326–349. https://doi.org/10.1002/cne.20970

Hoopes, A., Mora, J. S., Dalca, A. V., Fischl, B., & Hoffmann, M. (2022). SynthStrip: skull-stripping for any brain image. *NeuroImage*, *260*, 119474. https://doi.org/10.1016/J.NEUROIMAGE.2022.119474

Horne, J. A., & Ostberg, O. (1976). A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *International Journal of Chronobiology*, *4*(2), 97–110.

Johns, M. W. (1993). Daytime sleepiness, snoring, and obstructive sleep apnea; The Epworth Sleepiness Scale. *Chest*, *103*(1), 30–36. https://doi.org/10.1378/chest.103.1.30

Kain, M. P., Bolker, B. M., & McCoy, M. W. (2015). A practical guide and power analysis for GLMMs: Detecting among treatment variation in randomeffects. *PeerJ*, *2015*(9), e1226. https://doi.org/10.7717/PEERJ.1226/SUPP-1

Kasper, L., Bollmann, S., Diaconescu, A. O., Hutton, C., Heinzle, J., Iglesias, S., Hauser, T. U., Sebold, M., Manjaly, Z. M., Pruessmann, K. P., & Stephan, K. E. (2017). The PhysIO Toolbox for Modeling Physiological Noise in fMRI Data. *Journal of Neuroscience Methods*, *276*, 56–72. https://doi.org/10.1016/J.JNEUMETH.2016.10.019

Lam, R. W., Levitt, A. J., Levitan, R. D., Michalak, E. E., Cheung, A. H., Morehouse, R., Ramasubbu, R., Yatham, L. N., & Tam, E. M. (2016). Efficacy of Bright Light Treatment, Fluoxetine, and the Combination in Patients With Nonseasonal Major Depressive Disorder: A Randomized Clinical Trial. *JAMA Psychiatry*, *73*(1), 56–63. https://doi.org/10.1001/JAMAPSYCHIATRY.2015.2235

Lockley, S. W., Evans, E. E., Scheer, F. A. J. L., Brainard, G. C., Czeisler, C. A., & Aeschbach, D. (2006). Short-wavelength sensitivity for the direct effects of light on alertness, vigilance, and the waking electroencephalogram in humans. *Sleep*, *29*(2), 161–168. https://doi.org/10.1093/sleep/29.2.161

Lucas, R. J., Peirson, S. N., Berson, D. M., Brown, T. M., Cooper, H. M., Czeisler, C. A., Figueiro, M. G., Gamlin, P. D., Lockley, S. W., O'Hagan, J. B., Price, L. L. A., Provencio, I., Skene, D. J., & Brainard, G. C. (2014). Measuring and using light in the melanopsin age. *Trends in Neurosciences*, *37*(1), 1–9. https://doi.org/10.1016/j.tins.2013.10.004

Ma, S., Hangya, B., Leonard, C. S., Wisden, W., & Gundlach, A. L. (2018). Dual-transmitter systems regulating arousal, attention, learning and memory. *Neuroscience and Biobehavioral Reviews*, *85*, 21–33. https://doi.org/10.1016/J.NEUBIOREV.2017.07.009

Mathôt, S., Schreij, D., & Theeuwes, J. (2012). OpenSesame: an open-source, graphical experiment builder for the social sciences. *Behavior Research Methods*, *44*(2), 314–324. https://doi.org/10.3758/S13428-011-0168-7

Milosavljevic, N., Cehajic-Kapetanovic, J., Procyk, C. A., & Lucas, R. J. (2016). Chemogenetic Activation of Melanopsin Retinal Ganglion Cells Induces Signatures of Arousal and/or Anxiety in Mice. *Current Biology*, *26*(17), 2358–2363. https://doi.org/10.1016/j.cub.2016.06.057

Munch, M., Nowozin, C., Regente, J., Bes, F., De Zeeuw, J., Hadel, S., Wahnschaffe, A., & Kunz, D. (2016). Blue-Enriched Morning Light as a Countermeasure to Light at the Wrong Time: Effects on Cognition, Sleepiness, Sleep, and Circadian Phase. *Neuropsychobiology*, *74*(4), 207–218. https://doi.org/10.1159/000477093

O'Brien, K. R., Kober, T., Hagmann, P., Maeder, P., Marques, J., Lazeyras, F., Krueger, G., & Roche, A. (2014). Robust T1-Weighted Structural Brain Imaging and Morphometry at 7T Using MP2RAGE. *PLOS ONE*, *9*(6), e99676. https://doi.org/10.1371/JOURNAL.PONE.0099676

Paparella, I., Campbell, I., Sharifpour, R., Beckers, E., Berger, A., Aizpurua, J. F. B., Koshmanova, E., Mortazavi, N., Talwar, P., Degueldre, C., Lamalle, L., Sherif, S., Phillips, C., Maquet, P., & Vandewalle, G. (2023). Light modulates task-dependent thalamo-cortical connectivity during an auditory attentional task. *Communications Biology 2023 6:1*, *6*(1), 1–10. https://doi.org/10.1038/s42003-023-05337-5

Perrin, F., Peigneux, P., Fuchs, S., Verhaeghe, S., Laureys, S., Middleton, B., Degueldre, C., Del Fiore, G., Vandewalle, G., Balteau, E., Poirrier, R., Moreau, V., Luxen, A., Maquet, P., & Dijk, D. J. (2004). Nonvisual responses to light exposure in the human brain during the Circadian night. *Current Biology*, *14*(20), 1842–1846. https://doi.org/10.1016/j.cub.2004.09.082

Riemersma-van der Lek, R. F., Swaab, D. F., Twisk, J., Hol, E. M., Hoogendijk, W. J., & Van Someren, E. J. (2008). Effect of Bright Light and Melatonin on Cognitive and Noncognitive Function in Elderly Residents of Group Care Facilities. *Jama*, *299*(22), 2642. https://doi.org/10.1001/jama.299.22.2642

Rosenthal, N. E., & Bradt, G. W. T. (1984). Seasonal pattern assessment questionnaire (SPAQ). *National Institute of Mental Health, Bethesda, MD, USA*.

Sander, D., Grandjean, D., Pourtois, G., Schwartz, S., Seghier, M. L., Scherer, K. R., & Vuilleumier, P. (2005). Emotion and attention interactions in social cognition: Brain regions involved in processing anger prosody. *NeuroImage*, *28*(4), 848–858. https://doi.org/10.1016/J.NEUROIMAGE.2005.06.023

Scammell, T. E., Arrigoni, E., & Lipton, J. O. (2017). Neural Circuitry of Wakefulness and Sleep. In *Neuron* (Vol. 93, Issue 4, pp. 747–765). Cell Press. https://doi.org/10.1016/j.neuron.2017.01.014

Scammell, T. E., Jackson, A. C., Franks, N. P., Wisden, W., & Dauvilliers, Y. (2019). Histamine: neural circuits and new medications. *Sleep*, *42*(1), 1–8. https://doi.org/10.1093/SLEEP/ZSY183

Scheuermaier, K., Münch, M., Ronda, J. M., & Duffy, J. F. (2018). Improved cognitive morning performance in healthy older adults following blue-enriched light exposure on the previous evening. *Behavioural Brain Research*, *348*, 267–275. https://doi.org/10.1016/J.BBR.2018.04.021

Schoonderwoerd, R. A., de Rover, M., M Janse, J. A., Hirschler, L., Willemse, C. R., Scholten, L., Klop, I., van
Berloo, S., P van Osch, M. J., Swaab, D. F., Meijer, J. H., & by Joseph Takahashi, E. (2022). *The*
photobiology of the human circadian clock. https://doi.org/10.1073/pnas

Sharifpour, R., Campbell, I., Beckers, E., Balda, F., Mortazavi, N., Koshmanova, E., Paparella, I., Sherif, S.,
Phillips, C., & Vandewalle, G. (2022). Pitfalls in Recording Bold Signal Responses to Light in Small
Hypothalamic Nuclei Using Ultra-High-Field 7 Tesla MRI. *SSRN Electronic Journal*.
https://doi.org/10.2139/SSRN.4174993

Smolders, K. C. H. J., Peeters, S. T., Vogels, I. M. L. C., & de Kort, Y. A. W. (2018). Investigation of Dose-
Response Relationships for Effects of White Light Exposure on Correlates of Alertness and Executive
Control during Regular Daytime Working Hours. *Journal of Biological Rhythms*, *33*(6), 649–661.
https://doi.org/10.1177/0748730418796438

Sonoda, T., Li, J. Y., Hayes, N. W., Chan, J. C., Okabe, Y., Belin, S., Nawabi, H., & Schmidt, T. M. (2020). A
noncanonical inhibitory circuit dampens behavioral sensitivity to light. *Science (New York, N.Y.)*,
368(6490), 527–531. https://doi.org/10.1126/SCIENCE.AAY3152

Tri, M., & Do, H. (2019). Melanopsin and the Intrinsically Photosensitive Retinal Ganglion Cells: Biophysics
to Behavior. *Neuron*, *104*, 205–226. https://doi.org/10.1016/j.neuron.2019.07.016

Vandewalle, G., Archer, S. N., Wuillaume, C., Balteau, E., Degueldre, C., Luxen, A., Dijk, D. J., & Maquet, P.
(2011). Effects of light on cognitive brain responses depend on circadian phase and sleep
homeostasis. *Journal of Biological Rhythms*, *26*(3), 249–259.
https://doi.org/10.1177/0748730411401736

Vandewalle, G., Balteau, E., Phillips, C., Degueldre, C., Moreau, V., Sterpenich, V., Albouy, G., Darsaud, A.,
Desseilles, M., Dang-Vu, T. T., Peigneux, P., Luxen, A., Dijk, D. J., & Maquet, P. (2006). Daytime Light
Exposure Dynamically Enhances Brain Responses. *Current Biology*, *16*(16), 1616–1621.
https://doi.org/10.1016/j.cub.2006.06.031

Vandewalle, G., Gais, S., Schabus, M., Balteau, E., Carrier, J., Darsaud, A., Sterpenich, V., Albouy, G., Dijk,
D. J., & Maquet, P. (2007). Wavelength-dependent modulation of brain responses to a working
memory task by daytime light exposure. *Cerebral Cortex (New York, N.Y. : 1991)*, *17*(12), 2788–
2795. https://doi.org/10.1093/cercor/bhm007

Vandewalle, G., Hébert, M., Beaulieu, C., Richard, L., Daneault, V., Garon, M. Lou, Leblanc, J., Grandjean,
D., Maquet, P., Schwartz, S., Dumont, M., Doyon, J., & Carrier, J. (2011). Abnormal hypothalamic
response to light in seasonal affective disorder. *Biological Psychiatry*, *70*(10), 954–961.
https://doi.org/10.1016/j.biopsych.2011.06.022

Vandewalle, G., Schmidt, C., Albouy, G., Sterpenich, V., Darsaud, A., Rauchs, G., Berken, P. Y., Balteau, E.,
Dagueldre, C., Luxen, A., Maquet, P., & Dijk, D. J. (2007). Brain responses to violet, blue, and green
monochromatic light exposures in humans: Prominent role of blue light and the brainstem. *PLoS*
ONE, *2*(11), e1247. https://doi.org/10.1371/journal.pone.0001247

Vandewalle, G., Schwartz, S., Grandjean, D., Wuillaume, C., Balteau, E., Degueldre, C., Schabus, M.,
Phillips, C., Luxen, A., Dijk, D. J., & Maquet, P. (2010). Spectral quality of light modulates emotional
brain responses in humans. *Proceedings of the National Academy of Sciences of the United States of*
America, *107*(45), 19549–19554. https://doi.org/10.1073/pnas.1010180107

Vann, S. D. (2010). Re-evaluating the role of the mammillary bodies in memory. *Neuropsychologia*, *48*(8),
2316–2327. https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2009.10.019

Viénot, F., Brettel, H., Dang, T.-V., & Le Rohellec, J. (2012). Domain of metamers exciting intrinsically
photosensitive retinal ganglion cells (ipRGCs) and rods. *Journal of the Optical Society of America A*,
29(2), A366. https://doi.org/10.1364/josaa.29.00a366

Walmsley, L., Hanna, L., Mouland, J., Martial, F., West, A., Smedley, A. R., Bechtold, D. A., Webb, A. R.,
Lucas, R. J., & Brown, T. M. (2015). Colour As a Signal for Entraining the Mammalian Circadian Clock.
PLOS Biology, *13*(4), e1002127. https://doi.org/10.1371/journal.pbio.1002127

Wirz-Justice, A., Skene, D. J., & Münch, M. (2020). The relevance of daylight for humans. In *Biochemical
Pharmacology* (p. 114304). Elsevier Inc. https://doi.org/10.1016/j.bcp.2020.114304

Zeitzer, J. M. (2013). Control of sleep and wakefulness in health and disease. *Progress in Molecular
Biology and Translational Science*, *119*, 137–154. https://doi.org/10.1016/B978-0-12-396971-
2.00006-3

Figures and Tables

Figure 1. Experimental protocol.

(A) Overall timeline. After prior light history standardisation, participants performed executive (always
first), emotional and attentional tasks (pseudo-randomly 2nd or 3rd, blue arrow). As the attentional task
included fewer light conditions, it is not considered in the present manuscript (see methods for more
details).

(B) Spectral power distribution of light exposures. Monochromatic orange: 0.16 mel EDI lux;
Polychromatic, blue-enriched light (6500K); LOW, MID, HIGH: 37, 92, 190 mel EDI lux. For the present
analyses, we discarded colour differences between the light conditions and only considered illuminance
as indexed by mel EDI lux, constituting a limitation of our study. See **Suppl. Table 2** for full details).

(C-D) Tasks procedures. Time is reported in seconds relative to session onset; participants were pseudo-
randomly exposed to the 4 light conditions. **(C)** Executive task: alternation of letter detection blocks (0-
back) and working memory blocks (2-back). **(D)** Emotional task: lure gender discrimination of vocalizations
(50% angry (red), 50% neutral (white).

Figure 2. Illuminance impact on the hypothalamus subparts.

(A) Segmentation of the hypothalamus in five subparts in a representative participant. The nuclei encompassed by the different subparts are indicated in the right inset – according to (Billot et al., 2020). ARC: arcuate nucleus; DMH; dorsomedial nucleus; LH lateral hypothalamus; LTN: lateral tubular nucleus; MB: mamillary body; POA: preoptic area; PVN: paraventricular nucleus; PNH: posterior nucleus of the hypothalamus; SCN: suprachiasmatic nucleus; SON: supraoptic nucleus; TMN: tuberomammillary nucleus; VMN: ventromedial nucleus.

(B-C) Estimates (beta; arbitrary unit – a.u.) of the collective impact of illuminance variation on the activity of each hypothalamus subpart (Refer to Table 1 full statistics). **(B)** Executive task: significant main effect of hypothalamus subparts (p=0.002), no significant main of task type (p=0.4) or subpart-by-task-type interaction (p=0.61). **(C)** Emotional task: significant main effect of hypothalamus subparts (p<.0001), and of stimulus type (p=0.048) or subpart-by-stimulus-type interaction (p=0.74).

(D-G) Whole brain analyses of the collective impact of the variations in illuminance over the hypothalamus area - for illustration. A local positive peak (red; $p_{uncorrected}$<0.001) was detected over the posterior hypothalamus subpart (light blue) in executive **(E)** and emotional **(G)**. A local negative peak (red; $p_{uncorrected}$<0.001) was detected over the inferior-tubular hypothalamus subparts (light orange) during the executive task **(D)**, while local negative peak (red; $p_{uncorrected}$<0.001) was detected over the inferior-anterior (yellow) and superior-anterior (blue) hypothalamus subparts during the emotional task **(F)** – insets correspond to enlargements over the hypothalamus area. Arrows from panels B and C arise from and are colour coded according to the hypothalamus subpart that is displayed in panels D to G. These results indicate that our finding does not arise from a nearby "leaking" activation/deactivation.

(H-K) Estimates of the impact of each illuminance on the activity of the hypothalamus subparts. (Refer for Table 2 and Suppl. Tables 3/4/5/6 for full statistics) Activity dynamics across illuminance for each subpart (colour code as in A). Results are displayed per task or stimulus type although no interactions with task or stimulus type were detected. Significant illuminance-by-hypothalamus-subpart interactions were detected for **(H-I)** the executive task (p=0.041) and **(J-K)** the emotional task (p=0.041). Small letter indicate significant difference (p < 0.05) between the following subparts at illuminance: a. 92 mel EDI lux: posterior vs. superior-anterior & inferior-tubular; b. 190 mel EDI lux: posterior vs. inferior-anterior, superior-anterior & inferior-tubular; c. 0 mel EDI lux: posterior vs. superior-tubular; d. 92 mel EDI lux: posterior vs. superior-anterior; superior-anterior vs. superior-tubular; e. 190 mel EDI lux: posterior vs. inferior-anterior, superior-anterior & inferior-tubular; superior-tubular vs. superior-anterior , inferior-tubular & inferior-anterior.

Figure 3. Impact of illuminance on performance and relationships with the activity of the posterior hypothalamus subpart.

(A) Accuracy (percentage of correct responses) to the 2-back increased with increasing illuminance (p = 0.034).

(B) Accuracy to the 2-back task is negatively correlated to the activity of the posterior hypothalamus subpart (p = 0.0027).

(C-D) Accuracy to the 2-back task is not correlated to the activity of the inferior-anterior **(C)** and inferior-tubular **(D)** hypothalamus subparts (p > 0.4). Association between superior-anterior and superior-tubular subparts are not displayed but were not significant (p > 0.6). See Suppl Table 7 for full details.

(E) Accuracy to the 0-back task is not correlated to the activity of the posterior hypothalamus subpart (p = 0.45).

(F) Reaction times to the emotional stimuli did not significantly change with increasing illuminance (p =
0.41).

(G) Reaction times to the emotional stimuli are correlated to the activity of the posterior hypothalamus
subpart (p = 0.04) with higher activity associated to slower reaction times. **(H)** Reaction times to the neutral
stimuli are not correlated to the activity of the posterior hypothalamus subpart (p = 0.6). Solid and dashed
lines correspond to the significant and not significant linear regression lines, respectively.

Table 1. Differences between hypothalamus subparts in the collective impact of the variation in
illuminance on their activity.

Executive task							
Main GLMM				Pairwise comparisons			
Effect	F value (df)	P value*	Partial R²	Contrast#	t-value	P$_{uncorrected}$	P$_{corrected}$
Hypothalamus subparts	4.36 (4,225)	**0.002**	**0.08**	1 vs. 2	-0.30	0.76	0.99
				1 vs. 3	-3.48	**0.0006**	**0.0056**
				1 vs. 4	< 0.01	0.99	1
Task	0.74 (1,25)	0.4		1 vs. 5	-0.57	0.57	0.98
Hypothalamus subparts x task type	0.68 (4,225)	0.61		2 vs. 3	-3.17	**0.0017**	**0.015**
				2 vs. 4	0.31	0.76	0.99
				2 vs. 5	-0.27	0.79	0.99
Age	0.33 (1,22)	0.57		3 vs. 4	3.48	**0.0006**	**0.0055**
BMI	0.59 (1,22)	0.45		3 vs. 5	2.54	**0.0041**	**0.033**
Sex	0.01 (1,22)	0.91		4 vs. 5	-0.5	0.57	0.98
Emotional task							
Main GLMM				Pairwise comparisons			
Effect	F Value	P value*	Partial R²	Contrast#	t-value	P$_{uncorrected}$	P$_{corrected}$
Hypothalamus subparts	9.38 (4,194)	**<.0001**	**0.22**	1 vs. 2	0.67	0.67	0.99
				1 vs. 3	-4.76	**< 0.0001**	**< 0.0001**
				1 vs. 4	0.00	0.99	1
Task	4.33 (1,25)	**0.048**	**0.15**	1 vs. 5	-1.88	0.06	0.33
Hypothalamus subparts x stimulus type	0.5 (4,194)	0.74		2 vs. 3	-5.43	**< 0.0001**	**0.0001**
				2 vs. 4	-0.66	0.51	0.96
				2 vs. 5	-2.54	**0.012**	0.086
Age	0.43 (1,22)	0.52		3 vs. 4	4.76	**< 0.0001**	**< 0.0001**
BMI	0.05 (1,22)	0.83		3 vs. 5	2.85	**0.0048**	**0.038**
Sex	1.47 (1,22)	0.24		4 vs. 5	-1.88	0.061	0.32

Outputs of the generalized linear mixed model (GLMM) with subject as the random factor (intercept and slope), and task and subpart as repeated measures (ar(1) autocorrelation).

* The corrected p-value for multiple comparisons over 2 tests is p < 0.025.

Refer to Fig.2A for correspondence of subpart numbers

Table 2. Statistical outputs of GLMM testing for differences between the activity of each subpart of the hypothalamus under each illuminance

Executive task

Main GLMM				Comparisons between subparts per illuminance #			
Effect	F-value (df)	P value	Partial R²	Illuminance*	contrast	t-value	p-value
Subpart	1.4 (4,228)	0.23		92	2 vs. 3	-2.25	**0.025**
Illuminance	2.15 (4,1017)	0.073		92	3 vs. 4	2.58	**0.01**
				190	1 vs. 3	-2.80	**0.0053**
Task	3.24 (1,228)	0.073		190	2 vs. 3	-2.24	**0.025**
Subpart x Illuminance	1.7 (16,1017)	**0.041**	**0.09**	190	3 vs. 4	3.15	**0.0017**
Age	1.19 (1,22)	0.29					
BMI	0.01 (1,22)	0.9					
Sex	0.38 (1,22)	0.54					

Emotional task

Main GLMM				Comparisons between subparts per illuminance #			
Effect	F-value (df)	P value	Partial R²	Illuminance*	contrast	t-value	p-value
Subpart	4.29 (4,229)	**0.0023**	**0.07**	0	3 vs. 5	-2.05	**0.04**
				92	2 vs. 3	-2.53	**0.012**
				92	2 vs. 5	-2.96	**0.0032**
Illuminance	9.41 (4,1020)	**< 0.0001**	**0.035**	190	1 vs. 3	-3.31	**0.001**
				190	2 vs. 3	-4.75	**< 0.0001**
Task	0.13 (1,229)	0.72		190	1 vs. 5	-2.5	**0.013**
				190	2 vs. 5	-4.04	**<0.0001**
				190	3 vs. 4	3.13	**0.0018**
Subpart x Illuminance	1.7 (16,1020)	**0.041**	**0.026**	190	4 vs. 5	-2.32	**0.021**
Age	0.59 (1,22)	0.45					
BMI	1.54 (1,22)	0.23					
Sex	0.05 (1,22)	0.83					

* illuminance in mel EDI lux

Only significant comparisons are reported in the main text. For the full table, including post hocs comparing light levels within a subpart, refer to **Suppl. Table 3/4/5/6/7**