A new GUV-based assay to reconstitute membrane tethering *in vitro* demonstrates the vesicle tethering ability of the selective autophagy scaffold Atg11*.*

Devika Andhare and Michael J Ragusa

Department of Chemistry, Dartmouth College, Hanover, NH 03755, USA

*Correspondence: michael.J.ragusa@dartmouth.edu

ABSTRACT

Membrane tethering is essential for the generation of organelle contact sites and to anchor incoming vesicles to their target membranes before vesicle fusion. During autophagy in yeast, tethering of 30 nm vesicles to cargo is one of the first steps in the generation of the isolation membrane that engulfs the cargo to be degraded. While membrane tethering is critical for cellular function, many of the current biochemical techniques to assay for membrane tethering rely on indirect readouts and are limited in their ability to monitor protein localization at sites of tethering. As such, we developed a fluorescence-microscopy-based GUV liposome tethering assay (GLT) to directly visualize membrane tethering and monitor protein localization at tethering sites simultaneously. We initially used GLT with engineered membrane tethers to demonstrate the ease of use,

versatility and sensitivity of the assay. We also demonstrated that the selective autophagy

scaffolding protein Atg11 can bind, and tether negatively charged membranes but is

unable to bind GUVs mimicking the charge of the outer mitochondrial membrane. Atg11

instead requires the selective autophagy receptor Atg32 to be recruited to mitochondrial

membranes. Lastly, we demonstrate that Atg11 bound to Atg32 on GUVs can tether

negatively charged vesicles to GUVs. Collectively, our results reconstitute one of the first

steps during the initiation of mitochondrial autophagy and highlight the versatility of GLT

to study a range of membrane tethering events biochemically.

INTRODUCTION

Over the last two decades, an increasing number of studies have highlighted the importance of membrane tethering events in cells[1]. Membrane tethering occurs when individual proteins or multiprotein complexes bridge two different membranes holding them within a defined distance, frequently less than 100 nm. These tethering events can be homotypic, where both membrane compositions are identical, or more commonly heterotypic, where two distinct membranes such as the plasma membrane and the endoplasmic reticulum (ER) are tethered[2]. The main functions of membrane tethering in cells are to generate organelle contact sites and to recruit vesicles to their target organelles prior to fusion[3,4].

Due to the diversity of membrane tethering events in cells, the proteins required for tethering vary dramatically in their size, domain architecture and mechanism by which they tether membranes. One of the most direct membrane tethers is the extended synaptotagmin (E-synt) family of proteins which functions to tether the plasma membrane (PM) to the ER[5]. E-synt proteins interact directly with both the ER and PM allowing for a single protein to generate an organelle contact site[6]. However, most organelle contact sites require at least two proteins[2]. In these cases, the contact site is mediated by separate proteins that are either integral membrane proteins, soluble proteins anchored to the membrane by posttranslational modifications (PTMs), or peripheral membrane binding proteins[1]. In some contact sites, additional proteins are required to bridge the proteins that interact with membranes.

In vesicle fusion, large multisubunit tethering complexes, like the HOPS and CORVET

complexes, bind to RAB GTPases, that are anchored to the membranes by PTMs, and

SNARE proteins, embedded in membranes, to facilitate tethering[4,7,8]. These multisubunit

tethering complexes are soluble protein complexes that do not bind membranes directly

and instead require RABs and SNAREs for their recruitment to membranes[9]. Coiled-coil

tethers are another important family of proteins required for vesicle fusion in cells[10]. These

tethers contain long coiled-coil domains that bind to membrane-anchored RAB GTPases

on one side and either RAB GTPases or membrane directly on the other.

In addition to vesicle trafficking and contact site generation, membrane tethering also

plays a role in organizing vesicles for the catabolic process of autophagy. During

autophagy, cytosolic material, including organelles, are captured in double-membrane

vesicles, termed autophagosomes, and targeted to lysosomes to facilitate their

degradation[11]. As a part of this process, an autophagosomal membrane sheet is built

around the outside of the cargo to be degraded. Membrane tethering plays a role at

several stages in autophagy. In organelle autophagy, selective autophagy receptors

(SARs) embedded in organelle membranes recruit the autophagy initiation machinery to

the organelle to be degraded[12]. This results in tethering between the cargo membrane

and autophagosomal membrane which requires the selective autophagy scaffolding

protein, Atg11 in yeast or FIP200 in mammals[13-15]. After initiation, the SAR interacts with

a family of ubiquitin-like proteins, Atg8 in yeast and LC3 family members in mammals,

which are conjugated to phosphatidylethanolamine (PE) in the autophagosomal

membrane[16]. This interaction maintains tethering between the cargo and

autophagosomal membrane independent of Atg11 or FIP200[17]. In addition, we recently

demonstrated that the dimeric peripheral membrane binding protein Atg23 can tether

small autophagy precursor vesicles and that these tethering events play a role in

membrane recruitment during autophagy initiation[18].

Given the diversity of tethering events and proteins that facilitate these events, studying

membrane tethering has been challenging. This is particularly true in cells due to the

complexity of contact sites, the transient nature of tethering in vesicular trafficking and the

indirect nature of the available assays[19]. Fluorescently labeled proteins and bimolecular

fluorescence complementation have been used to determine if specific proteins or protein

complexes are located at potential tethering sites[19,20]. However, it's unclear if these

proteins are actively participating in tethering or if they are localized to these sites for

other purposes. Disrupting potential tethers in cells may not lead to a loss in the contact

site due to redundancy in tethers at the site[21]. In addition, a loss of contact sites or fusion

events in response to protein depletion or knockout could be due to indirect effects and

not necessarily due to direct loss in tethering ability. Therefore, biochemical approaches

have been critical to investigate if a protein or a protein complex is capable of mediating

membrane tethering. For example, biochemical studies have demonstrated the

importance of RABs and SNAREs for HOPS-mediated homotypic tethering events[22,23].

While these assays have been essential to shed light on the mechanisms of membrane

tethering, many commonly used biochemical assays rely on indirect measures of

tethering. This includes light scattering, turbidity assays and microscopy of liposome

clustering, which all monitor the increasing size of vesicle clusters as they are tethered

rather than two differentially labeled membranes being held together[24].

To investigate membrane tethering in vitro, we aimed to create an assay capable of

reconstituting and examining the diverse spectrum of membrane tethering events that

occur in cells. This encompasses both homotypic and heterotypic tethering events.

Additionally, we sought to establish an assay providing a more direct readout, enabling

independent monitoring of two distinct membranes while simultaneously tracking protein

recruitment to these membranes. Our objective was also to design an assay where one

of the membranes could be immobilized, facilitating real-time manipulation of tethering

events and allowing for the determination of tethering kinetics. This led to the

development of the GUV and Liposome Tethering (GLT) assay, building upon a method

recently employed in the study of membrane tethering by the autophagy protein Atg23[18].

We anticipate that GLT will find broad applicability in the biochemical investigation of

membrane tethering events.

RESULTS

Fluorescence-microscopy-based GLT assay

To establish an assay to study membrane tethering *in vitro* we added extruded liposomes

to giant unilamellar vesicles (GUVs) labeled with different fluorophores in the presence of

a tethering protein (Figure 1). We reasoned that the large size of GUVs (typically 1 to 100

μm in diameter) would allow us to directly visualize liposomes tethered on the GUV

surface, thus eliminating ambiguities arising due to aggregation of liposomes and/or

protein. Moreover, GUVs would allow us to observe if a fluorescently tagged protein also

localizes at tethering sites. To initially demonstrate the feasibility of this assay we

engineered an artificial membrane tether by fusing the monomeric phosphatidylinositol-

3-phosphate(PI3P)-binding EEA1 $FYVE_{1337-1411}$ with GST (GST-FYVE)[25]. Since GST

dimerizes, this generates a stable dimer of EEA1 $FYVE_{1337-1411}$ with two PI3P binding sites

in a single protein complex. Similarly engineered tethers generated by linking membrane

binding domains to GST have been shown to facilitate tethering[26]. GUVs containing PI3P

were first incubated with GST-FYVE, followed by incubation with 400nm liposomes

containing an identical lipid composition to the GUVs except for a different fluorophore for

detection. GUVs mixed with PI3P-liposomes alone or with the addition of purified GST

showed no co-localization between GUVs and liposomes (Figure 2A and B). In contrast,

addition of PI3P-liposomes to GUVs pre-incubated with 1μM GST-FYVE showed robust

recruitment of liposomes to GUVs as illustrated by fluorescence intensity traces along the

surface of a GUV in the liposome fluorescence channel (Figure 2C). This demonstrates

that GST-FYVE mediates homotypic tethering. Comparison of liposome fluorescence per

micron calculated along the surface of GUVs further shows that tethering of liposomes to

GUVs only occurred in the presence of GST-FYVE thereby validating GLT for the study

of homotypic membrane tethering (Figure 2D). To determine if protein localization could

be observed at tethering sites, we inserted a GFP tag between the GST and FYVE

domains in our construct generating GST-GFP-FYVE. Repeating the assay with GST-

GFP-FYVE, we observed that the protein coats GUVs and tethers liposomes (Figure 2E).

In addition, fluorescence intensity traces revealed that areas with tethered liposomes

showed increased intensity of GST-GFP-FYVE demonstrating that protein is localized to

tethering sites highlighting the applicability of this technique to monitor protein

colocalization with tethering events *in vitro* (Figure 2F).

The sensitivity of GLT is comparable to commonly used tethering assays

Having demonstrated that GLT can be used to monitor homotypic membrane tethering

we next sought to compare the sensitivity of this assay with some commonly used

techniques to measure membrane tethering. In most membrane tethering assays

liposomes are mixed with protein tethers which results in the clustering of vesicles. The

resulting increase in the size of liposome clusters, as measured by dynamic light

scattering (DLS), or an increase in the fluorescence intensity of liposomes as monitored

by microscopy have been frequently used to assay for membrane tethering[27-29].

Therefore, we compared tethering at increasing concentrations of GST-FYVE using DLS

and fluorescence microscopy-based liposome clustering with GLT. Extruded 400nm-

liposomes containing PI3P were mixed with GST-FYVE at concentrations ranging from

0.5μM to 4μM. Fluorescence microscopy (Figure 3A and B) and DLS (Figure 3C) revealed

that larger clusters of liposomes could be observed with the addition of 0.5μM of GST-

FYVE. These clusters grew progressively larger in response to increasing GST-FYVE

concentration and didn't show a significant increase in size beyond 2μM GST-FYVE

(Figure 3A and B). Similarly, GLT showed the recruitment of liposomes to GUVs with the

addition of 0.5μM of GST-FYVE and showed increasing tethering with increasing protein

concentration (Figure 3D and E). While all of these techniques demonstrated tethering,

non-significant differences in the amount of tethering were observed for all of the

techniques over some concentration ranges. This indicates that while these techniques

all demonstrate that membrane tethering occurred and was concentration-dependent,

they have differential sensitivity across small changes in protein concentration. These

results demonstrate that the sensitivity of GLT to detect tethering is comparable with DLS

and fluorescence microscopy-based methods to measure membrane tethering.

GLT can distinguish between membrane binding and tethering

One challenge in studying membrane tethering is differentiating between membrane

binding and tethering. This often requires two assays, one to examine membrane binding

and a second to monitor membrane tethering. However, we reasoned that GLT could

distinguish between these two possibilities in a single experiment which would make it a

powerful tool to combine with mutagenesis studies aimed to disrupt membrane binding or

tethering. To test this, we purified GFP-EEA1 FYVE$_{1337-1411}$ which lacks the dimerizing

GST and is therefore a monomer that should not be able to tether membranes. However,

without the dimerizing GST tag this protein showed minimal membrane binding and was

therefore not useful to distinguish between membrane binding and tethering

(Supplemental Figure 1). This result is consistent with previous studies that showed the

monomeric construct (amino acids 1337-1411) of EEA1 FYVE has a very low affinity for

PI3P[25]. Instead, we used the monomeric pleckstrin homology (PH) domain of

phospholipase C-delta 1 (PLC-delta) which has a high affinity for $PI(4,5)P_2$[30]. We purified

GFP-tagged PH domain with (GST-GFP-PH) and without (GFP-PH) the dimerizing GST

tag and tested their abilities to tether membranes using GLT. Both constructs of the PH

domain bound GUVs containing $PI(4,5)P_2$, although GST-GFP-PH showed higher binding

than GFP-PH which is consistent with an increased avidity for the membrane due to the

dimerization of the two PH domains (Figure 4A). Liposomes were recruited to GUVs in

the presence of the dimeric GST-GFP-PH demonstrating that this construct can efficiently

tether membranes. Fluorescence intensity traces along GUVs showed the peaks in

liposome fluorescence (cyan) were coincident with peaks in GST-GFP-PH fluorescence

further validating the use of GLT to monitor the presence of protein at membrane tethering

events (Figure 4B). If GFP-PH is capable of tethering liposomes to GUVs, albeit with less

efficiency due to the reduced membrane binding of GFP-PH, then normalizing the

integrated liposome fluorescence intensity to the GFP fluorescence intensity should give

similar ratios between both GST-GFP-PH and GFP-PH. However, GST-GFP-PH had

significantly higher liposome-to-protein fluorescence ratios as compared to GFP-PH

(Figure 4D). We therefore concluded that while GFP-PH retains membrane binding it is

incapable of tethering membranes. As such, GLT can be used to distinguish between

membrane binding and tethering by proteins using a single assay.

GLT demonstrates heterotypic tethering by the selective autophagy

Atg32-Atg11 complex

Next, we turned our attention to adapting GLT to reconstitute heterotypic tethering, which

involves protein-mediated bridging of membranes with distinct lipid compositions. For this,

we focused on tethering events that occur during mitochondrial selective autophagy,

termed mitophagy. Mitophagy begins with the selective autophagy receptor (SAR) Atg32,

embedded in the outer mitochondrial membrane, binding to the soluble scaffolding protein

Atg11[31,32]. Atg11 in turn recruits 30 nm precursor vesicles to mitochondria through its

interaction with the integral membrane protein Atg9 residing in these vesicles[13,33]. The

mammalian homolog of Atg11, FIP200, has been reported to bind membranes directly

once it binds to a SAR[15]. Based on this, we hypothesized that the binding of Atg32 to

Atg11 could initiate membrane binding by Atg11. If so, this suggests that we may be able

to reconstitute heterotypic membrane tethering by Atg32-Atg11 complex. To begin we

tested to see if Atg11 alone could bind, and tether vesicles made from yeast polar lipids

(YPL). To our surprise, GLT demonstrated that Atg11 alone was not only capable of

binding to YPL GUVs but it also was able to tether YPL GUVs and liposomes (Figure 5).

This demonstrates that Atg11 can bind membranes containing a high percentage of

negatively charged lipids (~42%) and suggests that Atg11 may serve as a homotypic

tether for autophagic precursor vesicles in cells through its interaction with both Atg9 and

membranes.

We next asked if Atg11 could be recruited to GUVs containing a lipid composition with a

similar charge (~20% negative charged lipids) to the outer mitochondrial membrane

(OMM) that are displaying the cytosolic domain of Atg32[34]. For this we fused the cytosolic

domain of Atg32(1-381), which binds to Atg11, with an N-terminal GFP and a C-terminal

10xHistidine tag (Supplemental Figure 2A). GUVs included 5 mol% of DGSNi^{2+}NTA to

allow for the recruitment of GFP-Atg32$_{1-381}$ to GUVs. To monitor Atg11 we labeled it with

Alexa Flour 594. GFP-Atg32$_{1-381}$ bound to GUVs containing DGSNi^{2+}NTA was able to

recruit Atg11. In contrast, Atg11 was not recruited to GUVs in the absence of Atg32

demonstrating that while Atg11 can bind YPL membranes it cannot bind GUVs containing

a similar charge to the OMM and instead requires Atg32 for its recruitment to autophagic

cargos (Supplemental Figure 2B).

To test if the Atg32-Atg11 complex could tether heterotypic membranes, we added

extruded YPL liposomes to GFP-Atg32$_{1-381}$ bound GUVs in the presence and absence of

Atg11 (Figure 6). YPL liposomes were tethered on GUVs only in the presence of Atg11

and the tethered liposomes colocalized with regions of higher Atg32 fluorescence (Figure

6C-E). These results demonstrate the ability of GLT to reconstitute and directly visualize

heterotypic tethering while also monitoring protein localization at these sites. From a

biological standpoint, while Atg9 is required for tethering *in vivo* this assay suggests that

mitochondrial-recruited Atg11 may also participate in these tethering events by binding

membranes directly to further enhance their recruitment and/or stabilize these tethering

events. Moreover, these results underscore the dual nature of some autophagy proteins,

wherein they are recruited to the cargo membrane through an interaction with SARs and

simultaneously engage the isolation membrane due to the presence of a high

concentration of negatively charged lipids in the autophagosomal membrane.

GLT allows real-time visualization and kinetic analysis of membrane tethering

To test if GLT can be used to monitor membrane tethering in real-time, we attempted to follow Atg11-mediated tethering of YPL liposomes on GFP-Atg32$_{1-381}$ bound GUVs. We found that after 10 minutes of addition, GUVs settled and were non-specifically adhered to LabTek chamber wells. We added GFP-Atg32$_{1-381}$ bound GUVs mixed with Atg11 on LabTek wells and allowed them to settle. Next, we started acquiring a time-lapse imaging sequence; added YPL liposomes and continued to monitor the fluorescence channel for the liposomes (Supplemental Movie 1 and 2). On GUVs displaying GFP-Atg32$_{1-381}$ with Atg11, liposomes started tethering on the GUV surface at 20 seconds after the liposomes appeared in the background (Figure 7A). Liposomes remained bound to GUVs and liposome fluorescence on GUVs increased over time. In contrast, on GUVs displaying GFP-Atg32$_{1-381}$ alone, liposomes did not bind GUVs and could be seen diffusing around GUVs throughout the length of the movie (Figure 7B). Analysis of the build-up of liposome fluorescence on a line drawn along the GUV confirmed that liposome fluorescence increased only in the presence of Atg11 (Figure 7C). The increase in liposome fluorescence in the presence of Atg11 fit to a plateau followed by a one-phase exponential rise equation and showed a t$_{half}$ of 155 seconds, suggesting that tethering occurred in a physiologically relevant timescale.

DISCUSSION

We demonstrated the broad usefulness of GLT to study membrane tethering *in vitro*. This

assay can be easily applied to study homotypic or heterotypic tethering events making it

ideal to reconstitute and examine tethering occurring at organelle contact sites and within

vesicle trafficking. We have shown that GLT has some significant advantages over

standard techniques for monitoring membrane tethering. First, it can be used to monitor

the localization of protein components at sites of tethering through three-color imaging.

Second, GLT can be used to distinguish membrane binding from tethering. In this way,

mutations made to individual proteins can be tested using GLT to determine if the

mutations lead to a loss of membrane binding, tethering or both. Lastly, GLT can be used

to monitor membrane tethering in real-time and the kinetics of membrane tethering can

be determined. We hope that real-time GLT can be employed in the future to monitor not

only tethering but also detethering providing new tools to study the regulation and

dynamics of tethering events.

In all of our GLT assays RhPE and DiD were used to label GUVs and liposomes,

respectively. It's important to note that exchange of these fluorophores was never

observed when membranes were tethered. This suggests that even though the

membranes are tethered in close proximity they are not brought close enough to trigger

fusion of the two membranes which would allow for the exchange of the fluorophores.

This suggests that GLT could be further adapted to monitor the exchange of the

fluorophores due to fusion events making it an even more powerful technique to study

membrane tethering.

To monitor tethering events, GLT quantifies the localization of liposomes on the surface

of GUVs. Consequently, GLT is limited in its ability to investigate tethering when both

membranes exhibit high curvature. GUVs are ideal for mimicking relatively flat membrane

surfaces in cells such as the PM, ER and mitochondria; while liposomes can be extruded

to mimic the curvature of vesicles, such as COP-II or clathrin-coated vesicles which are

between 40 and 100 nm in diameter[35,36].This makes GLT optimal for reconstituting

heterotypic tethering events where small vesicles are tethered to flat membranes prior to

vesicle fusion. For example, GLT may be a useful assay to reconstitute exocyst-mediated

heterotypic tethering of secretory vesicles to the PM[37].

Our work for the first time demonstrates that Atg11 can directly interact with and tether

membranes made from YPL. This result is in contrast to previous studies which

demonstrated that Atg11 was unable to bind membranes[13]. However, two key differences

could account for the observed variations in the biochemical abilities of Atg11. First, we

recombinantly expressed and purified Atg11 from HEK cell culture where it is a stable

dimer as opposed to recombinant expression and purification from *E. coli* where Atg11

behaves as a monomer. Second, we used YPL membranes that contain approximately

42% negatively charged lipids. In contrast, Atg11 was previously shown to not bind

liposomes composed of 60% POPC, 20% Cholesterol, 10% POPE and 10% POPS which

contain only 10% negatively charged lipids[13]. Therefore, it seems possible that Atg11

membrane binding requires a high negative charge which is consistent with the

composition of the autophagosomal membrane that contains 37% phosphatidylinositol

and low concentrations of phosphatidylserine and phosphatidic acid[38]. Atg11 membrane

binding may play important roles early in selective autophagy initiation. It may help in the

recruitment of Atg9 vesicles to selective autophagy initiation sites or it could help stabilize

tethering between cargo and autophagosomal membranes. In previous studies, we have

demonstrated that several autophagy proteins, including Atg13, bind to liposomes made

from YPL but not to membranes with a low concentration of negatively charged lipids[18,39].

This observation may suggest a common characteristic among several autophagy

proteins, wherein these proteins bind to the highly charged autophagosomal membrane

but are unable to bind to the less charged membrane of organelle cargos.

Liposome

Tethering protein

GUV

326
327
328 **FIGURE 1. Schematic showing the fluorescence microscopy-based GLT assay.**
329 GUVs and liposomes containing different fluorophores are mixed with a candidate
330 tethering protein or proteins capable of binding and bridging both membranes. Tethering
331 is visualized by the co-localization of liposomes on GUVs in the presence of the protein.
332

FIGURE 2. GLT using an engineered tether. 400 nm-extruded liposomes containing 74.9 mol% DOPC, 20 mol% DOPS, 5 mol% PI3P and 0.1 mol% of the fluorescent lipophilic dye DiD were incubated with GUVs containing 74.9 mol% DOPC, 20 mol% DOPS, 5 mol% PI3P and 0.1 mol% rhodamine PE (RhPE) in the presence or absence of different proteins. Representative images for liposomes mixed with GUVs (A), GUVs and

GST (B), or GUVs and GST-FYVE (C) are shown. Fluorescence intensity profiles for the
GUV and liposome channels along the circumference of the GUV are shown to the right
of each representative image. D) Scatter plots showing the percent of liposome
fluorescence per length of GUV normalized to the largest signal from A to C. Data from 3
independent repeats (N=3) were pooled and analyzed using non-parametric Mann-
Whitney's test. **** indicates $p<0.0001$. n indicates total number of GUVs analyzed for
each condition. E) GUVs containing PI3P were mixed with GST-GFP-FYVE followed by
incubation with PI3P-containing liposomes. Representative fluorescence images are
shown. F) Fluorescence intensity profiles for the GUV, liposome and protein fluorescence
channels around the circumference of the GUV shown in E.

FIGURE 3. Comparison of GLT with two commonly used methods to study tethering. 400nm-extruded liposomes containing 74.9 mol% DOPC, 20 mol% DOPS, 5 mol% PI3P and 0.1 mol% of the fluorescent lipophilic dye DiD were incubated with increasing concentrations of GST-FYVE. Tethering was measured using fluorescence

microscopy (A and B) or DLS (C). A) Representative fluorescence images showing clustering of PI3P-liposomes with increasing concentration of GST-FYVE. Scale bar=10μm B) Quantitation of A. Data from 3 independent repeats (N=3) were pooled and analyzed using Mann-Whitney's test. C) Quantitation of the increase in PI3P-liposome size in response to increasing GST-FYVE concentration as measured by DLS. Data from 4 independent repeats (N=4) were analyzed using Mann-Whitney's test. ** indicates p=0.0025, *** indicates p=0.0006. D) Representative fluorescence images from GLT using liposomes as in A and GUVs containing 74.9 mol% DOPC, 20 mol% DOPS, 5 mol% PI3P and 0.1 mol% RhPE in response to the addition of GST-FYVE at increasing concentration. E) Quantitation of D. Data from 3 independent repeats (N=3) were pooled and analyzed using Mann-Whitney's test. *** indicates p<0.0001. n indicates the total number of GUVs analyzed per condition.

FIGURE 4: GLT of dimeric and monomeric PH domains with PI(4,5)P$_2$ membranes.
GUVs containing PI(4,5)P$_2$ (magenta) were incubated with either GST-GFP-PH or GFP-

PH and subsequently mixed with PI(4,5)P$_2$-containing liposomes (cyan). A)
Representative fluorescence images of GLT where GUVs were mixed with liposomes and
either GST-GFP-PH (top panel) or GFP-PH (bottom panel). B) Fluorescence intensity
profiles from a line drawn along the GUV in A (top panel) showing liposome and GST-
GFP-PH fluorescence along the length of the GUV. C) Fluorescence intensity profiles
from a line drawn along the GUV in A (bottom panel) showing liposome and GFP-PH
fluorescence along the length of the GUV. D) Quantitation of A showing mean liposome
fluorescence intensity from a segmented line drawn along the GUV surface normalized
to mean protein fluorescence intensity. Data from 3 independent repeats (N=3) were
pooled and analyzed using non-parametric Mann-Whitney's test. **** indicates $p<0.0001$.
n indicated total number of GUVs analyzed per condition. GUVs were made with DOPC
(74.9mol%), DOPS(20mol%), DOPI(4,5)P$_2$(5mol%), and RhPE(0.1mol%). Liposomes
were made with identical composition except RhPE which was replaced with DiD.
386

Figure 5. GLT of Atg11 with GUVs and liposomes made from YPL. GUVs (magenta)
made from YPL alone or premixed with GFP-Atg11 (green) were incubated with
liposomes (cyan) made from YPL. For visualization GUVs were made with 0.1% RhPE
and liposomes were made with 0.1% DiD. A) Representative fluorescence images with
GFP-Atg11 (top panel) and without any protein (bottom panel). B) Fluorescence intensity
profiles from a line drawn along the GUV in A (top panel). C) Quantitation of A showing
fluorescence intensity of liposomes normalized to the length of GUV in the presence and
absence of GFP-Atg11. n represents the number of GUVs analyzed. **** indicates
$p < 0.0001$ analyzed using Mann-Whitney's test.

FIGURE 6. GLT testing the heterotypic tethering activity of the Atg32-Atg11 complex. A) Schematic showing heterotypic tethering reconstituted using GLT. B) Representative fluorescence images of GLT with GUVs (magenta) containing 5% mol DGS-Ni^{+2}NTA displaying His-GFP-Atg32$_{1-381}$ (green) which were sequentially incubated

with Atg11 and YPL liposomes (cyan) (top panel) or YPL liposomes (cyan) alone (bottom
panel). C) Fluorescence intensity profiles from a line drawn along the GUV in A (top panel)
showing liposome and GFP-Atg32$_{1-381}$ fluorescence intensities along the GUV. D)
Fluorescence intensity profiles from a line drawn along the GUV in A (bottom panel)
showing liposome and GFP-Atg32$_{1-381}$ fluorescence intensities along the GUV. E)
Quantification of the colocalization of liposomes and GFP-Atg32$_{1-381}$ fluorescence from
B. Data from 3 independent repeats (N=3) were pooled and analyzed using non-
parametric Mann-Whitney's test. **** indicates $p < 0.0001$. n indicates the total number of
GUVs analyzed per condition. GUVs were made with DOPC (74.9mol%), DOPS
(20mol%), DGS-Ni^{+2}NTA (5mol%), and RhPE (0.1mol%). Liposomes were made with YPL
(99.9mol%) and DiD (0.1mol%).
414

FIGURE 7. Real-time visualization and kinetic analysis of tethering using GLT. GUVs
containing 5 mol% of DGS-Ni^{+2}NTA incubated with GFP-Atg32$_{1\text{-}381}$ and Atg11 or GFP-
Atg32$_{1\text{-}381}$ alone were allowed to settle in a LabTek well. A time-lapse series was recorded

during the addition of YPL liposomes. A) A montage showing the real-time addition and tethering of liposomes (cyan) to a GUV (magenta) bound to GFP-Atg32$_{1-381}$(green) and Atg11. B) Montage showing the real-time addition of liposomes (cyan) to a GUV (magenta) bound to GFP-Atg32$_{1-381}$(green). C) Representative trace showing the buildup of liposome fluorescence on a line drawn along the GFP-Atg32$_{1-381}$ bound GUV surface in the presence (filled circles) and absence (empty circles) of Atg11. The red line shows the fit to a plateau followed by a one-phase exponential rise equation. GUVs were made with DOPC (74.9mol%), DOPS(20mol%), DGS-Ni^{+2}NTA (5mol%), and RhPE(0.1mol%). Liposomes were made with YPL (99.9mol%) and DiD (0.1mol%)

MATERIALS AND METHODS

Recombinant protein expression and purification

All constructs used for protein purification in this paper are listed in Supplemental Table 1. The PH domain from GFP-C1-PLCdelta-PH was subcloned into pET His6 GST TEV LIC cloning vector (1G) (Addgene plasmid # 29655 which was a gift from Scott Gradia) to generate 12XHis-GST-GFP-PH. GFP-C1-PLCdelta-PH was a gift from Tobias Meyer (Addgene plasmid # 21179)[40]. 6XHis-GST, 12XHis-GST-EEA1 FYVE, 12XHis-GST-GFP-EEA1 FYVE, 12XHis-GST-GFP-PH and StrepII-GFP-Atg32(1-381)-10XHis were recombinantly expressed in *E. coli* Star cells (Invitrogen C601003). Cultures were grown in LB broth (Fisher Scientific, BP1426) to an OD_{600} of 0.6 to 0.8 at 37°C while shaking at 210 rpm. Cultures were cooled at 4°C for 20 min. Protein expression was induced by adding 1 mM isopropylthio-β-D-galactopyranoside (IPTG; IBI Scientific, IB02125) and the cultures were grown for an additional 18 h at 18°C. Cells were harvested and cell pellets were stored at −80°C.

For purification of 6XHis-GST, 12XHis-GST-EEA1 FYVE, 12XHis-GST-GFP-EEA1 FYVE and StrepII-GFP-Atg32(1-381)-10XHis, cell pellets were thawed and resuspended in 50 mM Tris pH 8, 500 mM NaCl, 2.5mM imidazole, 5mM $MgCl_2$, 1 mM phenylmethanesulfonyl fluoride (PMSF; VWR, 0754) and cOmplete Mini EDTA-free protease inhibitor tablets (Roche, 11836170001). Cells were lysed using 3 strokes of a microfluidizer at 18000 psi. Triton X-100 (VWR, AAA16046-AE), was subsequently added to lysates at 1% vol/vol. Lysates were spun at 25,000g for 30 minutes. The supernatant was then applied to pre-equilibrated Talon resin (Clontech, 635504). The

resin was washed with 50 mM Tris, pH 8, 500 mM NaCl, 10 mM imidazole. Bound

proteins were eluted with 50mM Tris pH8, 500mM NaCl, 250mM imidazole. For

purification of 6XHis-GST, 12Xhis-GST-EEA1 FYVE, 12Xhis-GST-GFP-EEA1, peak

fractions containing protein were pooled and dialyzed against 20mM Tris pH 8, 300mM

NaCl and 0.2 mM TCEP. Proteins were aliquoted, flash-frozen in liquid nitrogen and

stored at −80°C. For purification of StrepII-GFP-Atg32(1-381)-10XHis, elutions from Talon

resin were pooled and incubated with pre-equilibrated Strep resin (IBA). The resin was

washed with 20 mM Tris pH8 and 300 mM NaCl. Protein was eluted with 20 mM Tris pH8,

300 mM NaCl and 50 mM biotin. Peak fractions containing protein were dialyzed against

20 mM Tris pH 8, 300 mM NaCl and 0.2 mM TCEP. Dialyzed protein was aliquoted, flash-

frozen in liquid nitrogen and stored at −80°C until use.

For purification of 12XHis-GST-GFP-PH constructs, cell pellets were resuspended and

thawed in 20 mM Tris pH 8.0, 500 mM NaCl, 2.5 mM imidazole, 1mM DTT and a cOmplete

Mini protease inhibitor tablet (Roche, 11836170001). Cells were lysed using 3 strokes of

a microfluidizer at 18000 psi. Triton X-100 (VWR, AAA16046-AE) was subsequently

added to lysates at 1% vol/vol. Lysates were spun at 25,000g for 30 minutes. Supernatant

was added to pre-equilibrated GST resin (add cat no) and incubated on a rocker for 1.5

hr at 4°C. Resin was washed with 20 mM Tris pH 8.0, 500 mM NaCl. The GST resin was

divided into two parts, one part was used for elution of GST-GFP-PH using 20 mM Tris

pH 8.0, 300 mM NaCl and 40 mM reduced glutathione. Eluted protein was dialyzed

against 20mM Tris pH 8.0, 300mM NaCl and 0.2mM TCEP. The other half of the GST

resin was incubated with TEV overnight at 4°C. Flowthrough from the GST resin

containing GFP-PH was subsequently incubated with Talon resin to remove TEV.

Flowthrough from the Talon resin was subjected to size exclusion chromatography using

a Superdex75 column. Peak fractions containing proteins were aliquoted, flash frozen in

liquid nitrogen and stored at −80°C until use.

Atg11 and GFP-Atg11, both with an N-terminal 2XStrep II tag, were recombinantly

expressed in Freestyle cells for 3 days post transfection. Cells were harvested by spinning

by 1000g for 15 mins. Cell pellets were flash frozen in liquid nitrogen and stored at −80°C.

Cell pellets were resuspended and thawed in 50mM Tris pH 8.0, 500mM NaCl, 1mM DTT,

1mM PMSF and a cOmplete Mini protease inhibitor tablet (Roche, 11836170001). Cells

were lysed using 3 strokes of a microfluidizer at 18000 psi. Triton X-100 (VWR,

AAA16046-AE), was subsequently added to lysates at 1% vol/vol. Lysates were spun at

25,000g for 30 minutes. Supernatant was applied to Strep resin (IBA) at 1mL/min using

a peristaltic pump. Resin was washed with 20mM Tris pH 8, 300 mM NaCl and 1 mM

DTT. Bound protein was eluted with 20mM Tris pH 8.0, 300mM NaCl, 50mM biotin and

1mM DTT. Fractions containing protein were pooled and concentrated to $2\mu M$ and flash-

frozen in liquid nitrogen. For GFP-Atg32 binding experiments, Atg11 was fluorescently

labeled with a three-fold molar excess of Alexa fluor 594 for 1hr at room temperature.

Labelling was quenched using 1 mM DTT. Unreacted dye was removed by dialyzing the

labelled protein against 20 mM Tris pH 8.0, 300 mM NaCl, 0.2 mM TCEP and 1 mM DTT

for 48 hrs. Fresh buffer was added after 24 hrs of dialysis.

Liposome preparation

All lipids were purchased from Avanti Polar lipids. Chloroform stocks of lipid mixtures at a

desired ratio containing 0.1mol% of the fluorescent lipophilic dye DiD were aliquoted in a

clean glass tube and dried under a gentle nitrogen stream until all the chloroform had

evaporated. Residual chloroform was removed under vacuum for 18 hrs. Dried lipids were

resuspended in 20 mM Tris pH 8, 150 mM NaCl to a final concentration of 1 mM. For

experiments with Atg11, YPL liposomes were resuspended in 20 mM Tris pH 8, 100 mM

NaCl to a final concentration on 1 mM. Liposomes were extruded using an Avanti Mini

Extruder using 0.4µm Nuclepore Track-Etched membrane (Sigma Aldrich,

WHA10417104).

Giant unilamellar vesicle preparation

GUVs were made using the gel-assisted formation method[41]. Briefly, chloroform stocks

containing lipids mixed at desired ratios containing the fluorescent lipid RhPE were

aliquoted in a glass tube. Chloroform was dried under a stream of nitrogen followed by

drying under vacuum (1 hr). The dried lipids were resuspended in fresh chloroform to

make a 1 mM lipid mixture. Polyvinyl alcohol (PVA, Mw = 14500; Sigma, 814894) was

dissolved in boiling water to make a 5% stock and degassed in a vacuum chamber. PVA

solution (10 µL) was spread into a thin film on a glass coverslip kept on a heating block

set to 55°C. Chloroform-dissolved lipid mix (10 µL) was then spread on the dried PVA

film. The dried PVA-lipid films were peeled from the glass coverslips and transferred to a

1.5-mL microcentrifuge tube. Films were hydrated in 300 µL assay buffer (20 mM Tris, pH

7.4, 100 mM NaCl) for 20 mins, and GUVs were released by gentle tapping. The PVA

films were removed using a pipette.

GUV liposome tethering (GLT)

For GLT experiments, GUVs were first mixed with tethering protein or control protein in a

1.5 mL Eppendorf tube for 10 min at room temperature. GUV mixtures were tapped to

resuspend settled GUVs and subsequently incubated with 10 µL of 400-nm extruded

liposomes for 5mins at room temperature. Liposomes were used within 3 days of

extrusion. The GLT mixtures (300µL total volume) were transferred to a BSA-passivated

LabTek chamber for imaging. GUVs were allowed to settle for 5 min prior to imaging.

LabTek chambers were treated with 3 M NaOH for 10 min followed by extensive washes

with milliQ water. NaOH-treated chambers were then passivated by incubation with BSA

(3mg/mL). BSA was removed by three washes with the assay buffer.

For real-time monitoring of GLT, GFP-Atg32$_{1-381}$ bound GUVs were incubated with Atg11.

GUVs displaying the GFP-Atg32$_{1-381}$-Atg11 complex were added to the BSA-passivated

LabTek chamber and allowed to settle for 5min. A field containing settled GUVs was

chosen, and real-time acquisition was started at 5 sec intervals. Liposomes (10 uL) were

added near the field under observation taking care not to disturb the settled GUVs.

Dynamic light scattering assay and microscopy-based liposomes clustering assay

To monitor liposome tethering using DLS, GST-FYVE was spun at 100,000g for 20 min

to remove large aggregates. 50μL of PI3P-containing 400nm extruded liposomes were

mixed with increasing concentrations of GST-FYVE in a total volume of 80uL. The

liposome and GST-FYVE mixture was placed on a rocker and incubated for 30 minutes

at room temperature. Liposome size was measured using DynaPro NanoStar.

To measure liposome clustering, PI3P-containing liposomes (with 0.1 mol% of DiD) mixed

with GST-FYVE prepared as described above were added to a BSA-passivated LabTek

chamber. Liposomes were allowed to settle for 5 mins and ten fields were randomly

imaged for each GST-FYVE concentration.

Fluorescence microscopy

549 GLT and liposome clustering were imaged using a Nikon TiE microscope fitted with a

550 Yokogawa CSU-W1 spinning disk system using a 100 X, 1.45 NA oil immersion objective.

551 Images were acquired with a Photometrics Prime BSI sCMOS camera using the NIS

552 elements software.

Image analysis

Fluorescence microscopy images were analyzed using Fiji[42]. and statistical analysis was

performed using GraphPad Prism (version 5.0a).

All experiments were performed in at least triplicates. Data from three independent

repeats (N = 3) were pooled and plotted as scatterplots. Error bars represent standard

deviation.

Percent liposome fluorescence per length of GUV was calculated by normalizing the

liposome fluorescence, collected from a two-pixel wide segmented line drawn along a

GUV, to the length of the GUV. For calculating the liposome/protein fluorescence ratio,

fluorescence intensities in liposome and protein channels from a two-pixel wide

segmented line drawn along a GUV were collected. Mean liposome fluorescence was

divided by mean protein fluorescence to yield liposome/protein fluorescence ratios.

To analyze tethering using the microscopy-based liposome clustering assay, images were

first background corrected using the mode intensity value. Images were auto-thresholded.

Liposome fluorescence per micron was calculated using the integrated density values of

all particles with an area more than 0.2 μm^2.

571   AUTHOR CONTRIBUTIONS

572   D.A. and M.J.R. conceived and planned the experiments. D.A. carried out the

573   experiments. D.A. and M.J.R. drafted the manuscript.

575   ACKNOWLEDGMENTS

This work was supported by the National Institute of General Medical Sciences (NIGMS)

grant R35GM128663 to M.J.R. Expression of Atg11 was performed in the BioMT service

center and protein purification was performed in part performed at the BioMT Molecular

Tools Core which are both supported by NIGMS grant P20GM113132. Sequencing of

plasmids was performed by the Molecular Biology Shared Resource Center which is

supported by NCI Cancer Center Support Grant P30CA023108. We would also like to

thank Ann Lavanway for her work and support at the Dartmouth Imaging Facility.

REFERENCES

1.   Szentgyörgyi V, Spang A. Membrane tethers at a glance. J Cell Sci 2023; 136. doi:ARTN jcs260471
10.1242/jcs.260471.
2.   Scorrano L, De Matteis MA, Emr S, Giordano F, Hajnoczky G, Kornmann B, et al. Coming together to define membrane contact sites. Nat Commun 2019; 10:1287. doi:10.1038/s41467-019-09253-3.
3.   Phillips MJ, Voeltz GK. Structure and function of ER membrane contact sites with other organelles. Nat Rev Mol Cell Biol 2016; 17:69-82. doi:10.1038/nrm.2015.8.
4.   Ungermann C, Kümmel D. Structure of membrane tethers and their role in fusion. Traffic 2019; 20:479-90. doi:10.1111/tra.12655.
5.   Saheki Y, De Camilli P. The Extended-Synaptotagmins. Biochim Biophys Acta Mol Cell Res 2017; 1864:1490-3. doi:10.1016/j.bbamcr.2017.03.013.
6.   Bian X, Saheki Y, De Camilli P. Ca(2+) releases E-Syt1 autoinhibition to couple ER-plasma membrane tethering with lipid transport. Embo J 2018; 37:219-34. doi:10.15252/embj.201797359.
7.   Spang A. Membrane Tethering Complexes in the Endosomal System. Front Cell Dev Biol 2016; 4:35. doi:10.3389/fcell.2016.00035.
8.   Jahn R, Scheller RH. SNAREs--engines for membrane fusion. Nat Rev Mol Cell Biol 2006; 7:631-43. doi:10.1038/nrm2002.
9.   Lurick A, Kummel D, Ungermann C. Multisubunit tethers in membrane fusion. Curr Biol 2018; 28:R417-R20. doi:10.1016/j.cub.2017.12.012.
10.   Cheung PY, Pfeffer SR. Transport Vesicle Tethering at the Trans Golgi Network: Coiled Coil Proteins in Action. Front Cell Dev Biol 2016; 4:18. doi:10.3389/fcell.2016.00018.
11.   Parzych KR, Klionsky DJ. An overview of autophagy: morphology, mechanism, and regulation. Antioxid Redox Signal 2014; 20:460-73. doi:10.1089/ars.2013.5371.
12.   Vargas JNS, Hamasaki M, Kawabata T, Youle RJ, Yoshimori T. The mechanisms and roles of selective autophagy in mammals. Nat Rev Mol Cell Biol 2023; 24:167-85. doi:10.1038/s41580-022-00542-2.
13.   Matscheko N, Mayrhofer P, Rao Y, Beier V, Wollert T. Atg11 tethers Atg9 vesicles to initiate selective autophagy. PLoS Biol 2019; 17:e3000377. doi:10.1371/journal.pbio.3000377.
14.   Sawa-Makarska J, Baumann V, Coudevylle N, von Bülow S, Nogellova V, Abert C, et al. Reconstitution of autophagosome nucleation defines Atg9 vesicles as seeds for membrane formation. Science 2020; 369:1206-+. doi:ARTN eaaz7714
10.1126/science.aaz7714.
15.   Shi X, Chang C, Yokom AL, Jensen LE, Hurley JH. The autophagy adaptor NDP52 and the FIP200 coiled-coil allosterically activate ULK1 complex membrane recruitment. Elife 2020; 9. doi:10.7554/eLife.59099.
16.   Johansen T, Lamark T. Selective Autophagy: ATG8 Family Proteins, LIR Motifs and Cargo Receptors. J Mol Biol 2020; 432:80-103. doi:10.1016/j.jmb.2019.07.016.
17.   Kirkin V, Rogov VV. A Diversity of Selective Autophagy Receptors Determines the Specificity of the Autophagy Pathway. Mol Cell 2019; 76:268-85. doi:10.1016/j.molcel.2019.09.005.

18.   Hawkins WD, Leary KA, Andhare D, Popelka H, Klionsky DJ, Ragusa MJ.
Dimerization-dependent membrane tethering by Atg23 is essential for yeast autophagy.
Cell Rep 2022; 39. doi:ARTN 110702
10.1016/j.celrep.2022.110702.
19.   Huang X, Jiang C, Yu L, Yang A. Current and Emerging Approaches for Studying
Inter-Organelle Membrane Contact Sites. Front Cell Dev Biol 2020; 8:195.
doi:10.3389/fcell.2020.00195.
20.   Jing J, Liu G, Huang Y, Zhou Y. A molecular toolbox for interrogation of
membrane contact sites. J Physiol 2020; 598:1725-39. doi:10.1113/JP277761.
21.   Eisenberg-Bord M, Shai N, Schuldiner M, Bohnert M. A Tether is a Tether is a
Tether: Tethering at Membrane Contact Sites. Developmental Cell 2016; 39:395-409.
doi:10.1016/j.devcel.2016.10.022.
22.   Shvarev D, Schoppe J, Konig C, Perz A, Fullbrunn N, Kiontke S, et al. Structure
of the HOPS tethering complex, a lysosomal membrane fusion machinery. Elife 2022;
11. doi:10.7554/eLife.80901.
23.   Stroupe C, Hickey CM, Mima J, Burfeind AS, Wickner W. Minimal membrane
docking requirements revealed by reconstitution of Rab GTPase-dependent membrane
fusion from purified components. Proc Natl Acad Sci U S A 2009; 106:17626-33.
doi:10.1073/pnas.0903801106.
24.   Ueda S, Tamura N, Mima J. Membrane Tethering Potency of Rab-Family Small
GTPases Is Defined by the C-Terminal Hypervariable Regions. Frontiers in Cell and
Developmental Biology 2020; 8. doi:ARTN 577342
10.3389/fcell.2020.577342.
25.   Dumas JJ, Merithew E, Sudharshan E, Rajamani D, Hayes S, Lawe D, et al.
Multivalent endosome targeting by homodimeric EEA1. Mol Cell 2001; 8:947-58.
doi:10.1016/s1097-2765(01)00385-9.
26.   Song H, Wickner W. Tethering guides fusion-competent trans-SNARE assembly.
Proc Natl Acad Sci U S A 2019; 116:13952-7. doi:10.1073/pnas.1907640116.
27.   Ho R, Stroupe C. The HOPS/Class C Vps Complex Tethers High-Curvature
Membranes via a Direct Protein-Membrane Interaction. Traffic 2016; 17:1078-90.
doi:10.1111/tra.12421.
28.   Nakatogawa H, Ichimura Y, Ohsumi Y. Atg8, a ubiquitin-like protein required for
autophagosome formation, mediates membrane tethering and hemifusion. Cell 2007;
130:165-78. doi:10.1016/j.cell.2007.05.021.
29.   Segawa K, Tamura N, Mima J. Homotypic and heterotypic trans-assembly of
human Rab-family small GTPases in reconstituted membrane tethering. J Biol Chem
2019; 294:7722-39. doi:10.1074/jbc.RA119.007947.
30.   Garcia P, Gupta R, Shah S, Morris AJ, Rudge SA, Scarlata S, et al. The
pleckstrin homology domain of phospholipase C-delta 1 binds with high affinity to
phosphatidylinositol 4,5-bisphosphate in bilayer membranes. Biochemistry 1995;
34:16228-34. doi:10.1021/bi00049a039.
31.   Kanki T, Wang K, Cao Y, Baba M, Klionsky DJ. Atg32 Is a Mitochondrial Protein
that Confers Selectivity during Mitophagy. Developmental Cell 2009; 17:98-109.
doi:10.1016/j.devcel.2009.06.014.

32.   Okamoto K, Kondo-Okamoto N, Ohsumi Y. Mitochondria-anchored receptor
Atg32 mediates degradation of mitochondria via selective autophagy. Dev Cell 2009;
17:87-97. doi:10.1016/j.devcel.2009.06.013.
33.   Coudevylle N, Banas B, Baumann V, Schuschnig M, Zawadzka-Kazimierczuk A,
Kozminski W, et al. Mechanism of Atg9 recruitment by Atg11 in the cytoplasm-to-
vacuole targeting pathway. J Biol Chem 2022; 298:101573.
doi:10.1016/j.jbc.2022.101573.
34.   Sperkagottlieb CDM, Hermetter A, Paltauf F, Daum G. Lipid Topology and
Physical-Properties of the Outer Mitochondrial-Membrane of the Yeast,
Saccharomyces-Cerevisiae. Biochim Biophys Acta 1988; 946:227-34. doi:Doi
10.1016/0005-2736(88)90397-5.
35.   Miller EA, Schekman R. COPII - a flexible vesicle formation system. Curr Opin
Cell Biol 2013; 25:420-7. doi:10.1016/j.ceb.2013.04.005.
36.   Kirchhausen T, Owen D, Harrison SC. Molecular structure, function, and
dynamics of clathrin-mediated membrane traffic. Cold Spring Harb Perspect Biol 2014;
6:a016725. doi:10.1101/cshperspect.a016725.
37.   Lepore DM, Martinez-Nunez L, Munson M. Exposing the Elusive Exocyst
Structure. Trends Biochem Sci 2018; 43:714-25. doi:10.1016/j.tibs.2018.06.012.
38.   Schütter M, Giavalisco P, Brodesser S, Graef M. Local Fatty Acid Channeling into
Phospholipid Synthesis Drives Phagophore Expansion during Autophagy. Cell 2020;
180:135-+. doi:10.1016/j.cell.2019.12.005.
39.   Gatica D, Damasio A, Pascual C, Klionsky DJ, Ragusa MJ, Popelka H. The
carboxy terminus of yeast Atg13 binds phospholipid membrane via motifs that overlap
with the Vac8-interacting domain. Autophagy 2020; 16:1007-20.
doi:10.1080/15548627.2019.1648117.
40.   Stauffer TP, Ahn S, Meyer T. Receptor-induced transient reduction in plasma
membrane PtdIns(4,5)P2 concentration monitored in living cells. Curr Biol 1998; 8:343-
6. doi:10.1016/s0960-9822(98)70135-6.
41.   Weinberger A, Tsai FC, Koenderink GH, Schmidt TF, Itri R, Meier W, et al. Gel-
assisted formation of giant unilamellar vesicles. Biophys J 2013; 105:154-64.
doi:10.1016/j.bpj.2013.05.024.
42.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al.
Fiji: an open-source platform for biological-image analysis. Nat Methods 2012; 9:676-
82. doi:10.1038/nmeth.2019.
708