Reconstitution of autophagosomal membrane tethering reveals that Atg11 can bind and cluster vesicles on cargo mimetics

Devika Andhare[1], Sarah Katzenell[1], Sarah I Najera[2], Katherine M Bauer[2] and Michael J Ragusa[*1,2]

[1]Department of Chemistry, Dartmouth College, Hanover, NH 03755, USA

[2]Department of Biochemistry and Cell Biology, Geisel School of Medicine, Dartmouth College, Hanover, New Hampshire 03755, United States.

[*]Correspondence: michael.j.ragusa@dartmouth.edu

Keywords

biochemistry; mitophagy; selective autophagy; reconstitution; yeast

ABSTRACT

Autophagy is essential for the degradation of mitochondria from yeast to humans. Mitochondrial autophagy in yeast is initiated when the selective autophagy scaffolding protein Atg11 is recruited to mitochondria through its interaction with the selective autophagy receptor Atg32. This also results in the recruitment of small 30 nm vesicles that fuse to generate the initial autophagosomal membrane. We demonstrate that Atg11 can bind to autophagosomal-like membranes in vitro in a curvature dependent manner via a predicted amphipathic helix. Deletion of the amphipathic helix from Atg11 results in a delay in the formation of mitophagy initiation sites in yeast. Furthermore, using a novel biochemical approach we demonstrate that the interaction between Atg11 and Atg32 results in the tethering of autophagosomal-like vesicles in clusters to giant unilamellar vesicles containing a lipid composition designed to mimic the outer mitochondrial membrane. Taken together our results demonstrate an important role for autophagosomal membrane binding by Atg11 in the initiation of mitochondrial autophagy.

ABBREVIATIONS

Atg11: autophagy related 11; Atg32: autophagy related 32; DLS: dynamic light scattering; GLT: GUV liposome tethering assay; GUV: giant unilamellar vesicle; PE: phosphatidyl ethanolamine; PI3P: phosphatidyl inositol 3-phosphate; SAR: selective autophagy receptor; YPL: yeast polar lipids

INTRODUCTION

Macroautophagy, hereafter autophagy, is the cellular process in which double-membrane vesicles, termed autophagosomes, are generated around cytosolic material leading to their capture within these vesicles[1]. Completed autophagosomes fuse with the vacuole, in yeast, or lysosomes, in higher eukaryotes, leading to the degradation of the captured contents. Autophagy can capture and degrade a wide range of cytosolic material including protein aggregates, intracellular pathogens and membrane-bound organelles. As such, autophagy is essential to maintain cellular homeostasis and also serves as an integral component of innate immunity[2,3].

Cargo selection during autophagy can occur via a non-selective or selective mechanism[4]. In selective autophagy, cargos are marked as targets for autophagy by selective autophagy receptors (SARs)[5]. SARs are either soluble cytosolic proteins that commonly recognize ubiquitin moieties on the cargo to be degraded or, in the case of membrane-bound organelles, they can also be integral membrane proteins on the organelle to be degraded. To initiate selective autophagy, SARs recruit the selective autophagy scaffolding protein Atg11, in yeast, or FIP200, in higher eukaryotes, to the cargo to be degraded[6]. In yeast, Atg11 also binds to Atg9, an integral membrane protein embedded in 30 nm vesicles that serve as the initial membrane source for autophagy[7-9]. Once recruited to the cargo, Atg9 vesicles fuse to generate an initial membrane sheet, termed the phagophore[10]. The phagophore expands around the cargo through lipid transport from Atg2 in addition to other mechanisms[11-13]. Once the autophagosomal membrane has fully surrounded the cargo, it is closed leading to the capture of the marked cargo in a double membrane autophagosome[14].

One of first critical steps in selective autophagy is the recruitment of Atg9 vesicles by Atg11 to

autophagic cargo. Deletion of Atg11 in yeast was initially shown to block the recruitment of Atg9

to autophagy cargos highlighting the important role of Atg11 in recruiting membrane to cargo

during selective autophagy initiation[15]. Atg11 was later shown to be required for the recruitment

of Atg9 vesicles to cargo mimetic beads in vitro using biochemical reconstitution[9]. Following on

this work, Atg11 was shown to directly interact with the flexible N-terminus of Atg9 via two PLF

motifs[8]. More recently, truncations or mutations within the N-terminal domain of Atg11 have been

shown to lead to a loss in the interaction between Atg11 and Atg9 in yeast[16]. These studies all

point towards an essential role for Atg11 in the recruitment of membrane to cargos, specifically

through the interaction between Atg11 and Atg9. However, while mutation of both PLF motifs in

Atg9 led to a near complete loss in the interaction between Atg11 and Atg9, it did not fully prevent

the recruitment of Atg9 to selective autophagy cargos[8]. Furthermore, the mammalian analog of

Atg11, FIP200, can bind membranes directly, although the exact function of membrane binding

by FIP200 is unknown[17]. Given these observations, we wanted to test if Atg11 is capable of binding

membranes and whether this plays a role in selective autophagy.

RESULTS

Atg11 binds autophagosomal membranes in a clustered manner

To test if Atg11 can bind membranes we first purified Atg11 and GFP-Atg11 using Freestyle 293

cells as a recombinant expression system (**Figure S1**). Size exclusion chromatography of Atg11

in buffer containing 300mM NaCl had a similar elution profile and estimated molecular weight to

previously published purifications of Atg11 from SF9 insect cells where it was shown to be a dimer

in solution (**Figure S1B**)[18]. Next, we tested the ability of Atg11 to bind membranes. Both

autophagosomal membranes and Atg9 vesicles are known to be composed of an unusually high

concentration of negatively charged lipids, including between 37% and 44% of the negatively

charged phosphatidylinositol (PI)[9,19]. Previously we have shown that several autophagy proteins

including Atg23 and Atg16 bind membranes generated from yeast polar lipids (YPL), which also

contain a high concentration of negatively charged lipids[20,21]. Therefore, we first tested if Atg11

can bind giant unilamellar vesicles (GUVs) made from YPL or a lipid mixture designed to mimic

the autophagosomal membrane composition (43.9% phosphatidylcholine (PC), 19%

phosphatidylethanolamine (PE), 32% phosphatidylinositol (PI), 5% phosphatidylinositol-3-

phosphate (PI3P), and 0.1% rhodamine PE (RhPE) (automix+PI3P))[19]. 5% PI3P was added at the

expense of PI as autophagosomal membranes contain PI3P but the exact concentration is unknown.

GFP-Atg11 showed robust binding to YPL GUVs but was unable to bind autophagosomal

membrane GUVs (**Figure S2A and B**). Given that Atg9 vesicles are highly curved while GUVs

are essentially flat relative to the size of Atg11, we next investigated if curvature impacts the ability

of Atg11 to bind membranes. For this we employed supported membrane templates (SMrT) that

consist of an array of highly curved membrane tubes ranging from 20-60 nm in diameter and

supported lipid bilayers (SLB) resting on a PEGylated glass coverslip[22,23]. The size of the SMrT

nanotubes match the size of Atg9 vesicles that are reported to be 30-60 nm in diameter while SLBs

are primarily flat membranes[10,24]. We made SMrT tubes and SLBs containing 43.9% PC, 19% PE

and either 37% PI alone (automix) or 32% PI with 5% PI3P (automix+PI3P). Interestingly, GFP-

Atg11 bound autophagosomal-like membranes as discreet clusters (**Figure 1A**). The presence of

PI3P in membranes enhanced GFP-Atg11 recruitment. Importantly, the quantification of the ratio

of GFP to membrane fluorescence showed that Atg11 binding was 5-fold higher on tubes (mean

= 0.90 ± 0.4) than on SLB (mean = 0.16 ± 0.07) suggesting that Atg11 membrane binding is

curvature sensitive (**Figure 1B**). As an independent validation of membrane binding, we also

performed a floatation assay with automix and automix+PI3P liposomes. Atg11 migrated to the

top layer of the sucrose density gradient in the presence of 100 nm diameter but not 1 µm diameter

automix+PI3P liposomes further supporting the curvature sensitivity of Atg11 membrane binding

(**Figure S3A**). DLS analysis also demonstrated that Atg11 binding to 100 nm liposomes increased

the diameter of these vesicles from 107 nm to 260 nm suggesting that Atg11 may also be able to

tether automix+PI3P liposomes (**Figure S3B**). These results suggest that Atg11 can bind

autophagosomal-like membranes in a curvature dependent manner and that this binding is

enhanced by the presence of PI3P.

The N-terminal (amino acids 1-646) and C-terminal (amino acids 700-1178) regions of Atg11 can

be purified using *E. coli* as a recombinant expression system[18]. Structure prediction of the N-

terminal region of Atg11 (Atg11-Nterm) using AlphaFold 3 demonstrates that it forms a helical

dimer with a structure that is similar to the N-terminal domain of FIP200 except that it is predicted

to form a Z shaped dimer instead of a C shaped dimer (**Figure S4A**)[25,26]. The C-terminal region of

Atg11 (Atg11-Cterm) contains a long coiled-coil domain, connected by linker regions to the Claw

domain that binds SARs[27,28]. We purified GFP-tagged Atg11-Nterm and -Cterm and analyzed

their membrane binding abilities on SMrT tubes and SLBs (**Figure 1C**). Quantitative fluorescence

microscopy showed that Atg11-Nterm had significantly higher membrane binding than Atg11-

Cterm suggesting that Atg11-Nterm contains the primary membrane binding region (**Figure 1D**).

Moreover, unlike the full-length protein, Atg11-Nterm did not cluster on membranes and instead

bound both membrane tubes and SLBs in a diffuse manner suggesting that membrane binding and

clustering requires distinct regions of Atg11.

Atg11 contains an amphipathic helix required for membrane binding

Sequence and structural analysis of the Atg11-Nterm revealed that it contains a predicted

amphipathic helix spanning residues 612-646, with the highest hydrophobic moment (μH=0.546)

occurring between residues 626 and 643 (**Figure 2A**). Amphipathic helices are known to mediate

curvature- and charge-dependent membrane binding[29,30]. Importantly, this region in the predicted

Atg11-Nterm structure has low and very low pLDDT scores, indicating that it is likely not in the

correct conformation in the structure prediction (**Figure S4B**). This suggests that this helix may

instead be available for membrane binding. Deletion of amino acids 612-646 from the Atg11-

Nterm sequence and rerunning the AlphaFold3 prediction did not alter the overall dimer structure

further suggesting that the predicted amphipathic helix is not required for formation of the Atg11-

Nterm dimer structure (**Figure S4C**). We therefore deleted the predicted amphipathic helix in

Atg11-Nterm (Atg11-Nterm$^{\Delta612\text{-}646}$) and purified it using *E. coli* as a recombinant expression

system. We compared the elution profile of Atg11-Nterm$^{\Delta612\text{-}646}$ from size exclusion

chromatography with that of Atg11-Nterm to verify that deletion of this helix did not disrupt the

overall shape of the protein (**Figure S4D**). The ability of GFP-Atg11-Nterm$^{\Delta612\text{-}646}$ to bind

membrane tubes was tested using SMrT tubes (**Figure 2B**). Comparison of GFP to membrane

fluorescence ratios revealed that Atg11-Nterm$^{\Delta612\text{-}646}$ had 2.6-fold lower binding to membrane

tubes than Atg11-Nterm (**Figure 2C**). Together these results suggest that Atg11 binds

autophagosomal-like membranes primarily via an amphipathic helix between amino acids 612-

646.

To determine if the amphipathic helix of Atg11 is required for selective autophagy in cells we

monitored the selective autophagy of mitochondria (mitophagy) using the established OM45-GFP

assay[31]. In this assay, the outer mitochondrial membrane (OMM) protein OM45 is tagged with a

C-terminal GFP. Once OM45-GFP is delivered to the vacuole during mitophagy, OM45 is rapidly

degraded resulting in the production of free GFP. We performed the OM45-GFP processing assay

to assess the effect of the putative amphipathic helix deletion on mitophagy. While we observed a

slight drop in the level of free GFP produced at 2 hours, this change was not significant suggesting

that deletion of the amphipathic helix from Atg11 does not impair mitochondrial degradation by

autophagy (**Figure 2D** and **E**). Furthermore, immunoblotting against 2xGFP-Atg11 and 2xGFP-

Atg11$^{\Delta612-646}$, expressed from their endogenous promoters in *atg11Δ* cells, showed that the deletion

of amino acids 612-646 did not alter the expression of Atg11, suggesting that deletion of this helix

did not destabilize Atg11 (**Figure 2F**).

Atg32, an integral membrane protein of the OMM, serves as the selective autophagy receptor

(SAR) for mitophagy in yeast[32,33]. Under mitophagy-inducing conditions, we and others have

shown that Atg32 clusters on the mitochondrial surface, forming discrete punctum[27,34]. These

Atg32 puncta initially colocalize with Atg11 and subsequently with the autophagosomal marker

Atg8, before being transported to the vacuole[32,35]. Thus, these puncta are thought to represent sites

of mitophagy initiation. In contrast to OM45-GFP, which shows only a faint vacuolar GFP signal

after 2 hours of starvation, we and others have observed that Atg32 clustering into mitophagy

initiation sites occurs more rapidly, with GFP-Atg32 puncta forming within 30 minutes of

starvation[27,35]. This rapid clustering suggests that monitoring Atg32 puncta formation may provide

a more effective approach for studying the early stages of mitophagy initiation. We therefore tested

to see if Atg11$^{\Delta612-646}$ altered the formation of Atg32 puncta. To test if Atg11$^{\Delta612-646}$ altered the

kinetics of mitophagy initiation, we monitored the formation of GFP-Atg32 puncta, in response to

nitrogen-starvation, in real-time using the temperature- and flow rate- controlled FCS2 chamber

(**Figure 2G**). We found that the rate of GFP-Atg32 puncta appearance was significantly lowered

in cells expressing Atg11$^{\Delta612-646}$ than in cells expressing WT Atg11 (**Figure 2H, Supplemental**

Movie 1). At 28 minutes post starvation, 97% of cells expressing WT Atg11 showed at least one

GFP-Atg32 punctum, whereas only 43% of cells expressing Atg11$^{\Delta612-646}$ displayed at least one

GFP-Atg32 punctum at the same time point (**Supplemental Movie 2**). Moreover, 2xGFP-

Atg11$^{\Delta612-646}$ was still recruited to the RFP-Atg32 puncta, similar to 2xGFP-Atg11$^{\Delta612-646}$,

suggesting that deletion of the amphipathic helix did not affect recruitment of Atg11 during

mitophagy (**Figure S5**). Together, our OM45-GFP processing and GFP-Atg32 clustering results

suggest that the deletion of the predicted amphipathic helix in Atg11 causes a delay in the kinetics

of mitophagy initiation but does not alter the bulk degradation of mitochondria.

Atg11 forms clusters in solution

Formation of autophagy protein condensates have been proposed to enhance weak protein

interactions, direct the formation of protein-complexes, and play a role in the overall organization

of autophagy initiation sites[36,37]. We observed that GFP-Atg11 clustered on membrane tubes and

SLBs which prompted us to investigate if Atg11 may cluster in solution or if the clustering of

Atg11 is induced by membrane binding. GFP-Atg11 was purified in the presence of 300 mM NaCl

as this was previously reported to allow for soluble full-length Atg11(**Figure S1A**)[9,18]. We

monitored 1 µM GFP-Atg11 by confocal microscopy to determine if it formed puncta, which

would indicate clustering of Atg11 (**Figure 3A**). We found that while GFP-Atg11 appeared largely

diffuse at 300 mM NaCl, it formed puncta when the NaCl concentration was reduced to 150 mM

or less (**Figure 3A and B**). We next asked if the formation of these puncta was dependent on the

concentration of Atg11. We diluted GFP-Atg11 to concentrations from 0.05 µM to 5.0 µM into

buffer with a final concentration of 150 mM NaCl (**Figure 3C**). We observed that increasing the

concentration of Atg11 increased the number of Atg11 puncta indicating that Atg11 clustering is

both salt and concentration dependent (**Figure 3D**). The puncta we observed were not liquid-like

condensates as they are not entirely circular, nor did we observe the fusion of individual puncta. It

is possible that diluting Atg11 into 150 mM NaCl resulted in an irreversible aggregation event.

Therefore, to test if the formation of Atg11 clustering was reversible, we first generated Atg11

puncta in 150 mM NaCl, and then increased the NaCl concentration to 300 mM by mixing the 1

µM Atg11 in 150 mM NaCl with 1 µM of Atg11 in 450 mM NaCl (**Figure 3E**). After the samples

were mixed, we saw a near complete loss in the formation of Atg11 puncta suggesting that these

Atg11 clustering events are reversible in solution (**Figure 3F**). We next wanted to determine

whether Atg11-Nterm or Atg11-Cterm could undergo similar clustering events as full-length

Atg11 in solution. Therefore, we diluted GFP-Atg11, GFP-Atg11-Nterm and GFP-Atg11-Cterm

into buffer containing 150 mM NaCl. While GFP-Atg11 formed puncta at 1 µM in 150 mM NaCl,

neither the Nterm nor Cterm regions of Atg11 formed puncta at 1 µM in 150 mM NaCl, suggesting

that the ability of Atg11 to cluster is a property of the full-length protein (**Figure 3G and H**). This

data demonstrates that while full-length Atg11 can form reversible clusters in solution that are salt

and concentration dependent, the Nterm and Cterm regions of Atg11 alone are unable to form

clusters. This is consistent with our membrane binding observations in **Figure 1** where full-length

Atg11 bound to membranes in a clustered fashion while the Nterm and Cterm regions of Atg11

bound membranes in a diffuse manner.

The Atg32-Atg11 complex tethers autophagosomal-like liposomes to mitochondria-like GUVs

We have demonstrated that Atg11 can bind membranes in a clustered manner via a predicted

amphipathic helix and that a loss of this helix leads to a delay in the formation of Atg32 puncta

and the onset of mitophagy in yeast. Recruitment of the highly-curved Atg9 vesicles by the Atg32-

Atg11 complex is a critical step in initiating autophagosomal membrane formation during

mitophagy[7]. To investigate whether the ability of Atg11 to bind membranes enables the tethering

of liposomes (mimicking the size and composition of Atg9 vesicles) to mitochondrial membranes,

we utilized the <u>G</u>UV and <u>L</u>iposome <u>T</u>ethering (GLT) assay, which we recently developed[20]. GLT

monitors tethering by measuring the co-localization of liposomes with GUVs in the presence and

absence of potential tethering proteins. We made GUVs containing a lipid composition similar to

the OMM (OMM-GUVs)[38-40]. GFP-Atg11 alone was unable to bind these OMM-GUVs further

highlighting that Atg11 is not a broad membrane binding protein (**Figure S6A**). Next, we

displayed the cytosolic domain of the mitophagy SAR Atg32 (1-381), which contains the Atg11

binding region, fused to an N-terminal GFP and a C-terminal 10xHistidine tag, on OMM-GUVs

(**Figure S6B**)[41]. GUVs included 5 mol% of DGSNi^{2+}NTA to recruit the cytosolic domain of GFP-

Atg32$_{1-381}$. To monitor if Atg11 can bind to these Atg32-displaying OMM-GUVs, we labeled

purified Atg11 with Alexa Flour 594. Atg11 was recruited to GFP-Atg32$_{1-381}$ bound GUVs. In

contrast, no Atg11 recruitment occurred to GUVs displaying 6xHistidine tagged GFP (**Figure**

S6C), demonstrating that Atg11 requires Atg32 for its recruitment to the surface of OMM-GUVs

(**Figure S6D**). To test if the Atg32-Atg11 complex could tether autophagosomal-like liposomes,

we added sonicated automix+PI3P liposomes, mimicking the lipid composition and size of Atg9

vesicles (30 nm diameter), to GFP-Atg32$_{1-381}$ OMM-GUVs in the presence and absence of Atg11

(**Figure 4A**). Automix+PI3P liposomes were recruited to Atg32$_{1-381}$ bound GUVs only in the

presence of Atg11 (**Figure 4B-E**). Fluorescence intensity traces along the GUVs further confirm

that peaks of liposome fluorescence correspond with peaks of GFP-Atg32$_{1-381}$ fluorescence and

were only seen in presence of Atg11 (**Figure 4C** and **D**). Interestingly, tethered automix+PI3P

liposomes appeared punctate on GUVs and these overlap with peaks of Atg32 fluorescence,

suggesting that Atg11 may be able to cluster liposomes on the surface of cargo (**Figure 4C**). While

Atg11 robustly tethered 30 nm automix+PI3P liposomes to Atg32 OMM-GUVs, the ability of

Atg11 to recruit larger automix+PI3P liposomes (1 µm diameter) was significantly reduced

(**Figure 4F** and **G**). This is in agreement with our results on SMrT tubes showing that

automix+PI3P membrane binding by Atg11 occurred in a curvature-dependent manner.

We hypothesized that using YPL could override the need for curvature as Atg11 was able to bind

GUVs containing YPL but not automix+PI3P GUVs (**Figure S2**). This may be due to the

heterogeneity of lipids present in YPL which could lead to packing defects similar to what would

be expected for more highly curved vesicles[48]. To test if Atg11 was also able to tether larger YPL

liposomes to GUVs, we performed the GLT assay with 400 nm YPL liposomes and GFP-Atg32$_{1-}$

$_{381}$ bound GUVs in the presence and absence of Atg11 (**Figure S7A**). As expected, YPL liposomes

were tethered on GUVs only in the presence of Atg11 (**Figure S7D**) and the tethered liposomes

colocalized with regions of higher Atg32 fluorescence (**Figure S7B** and **C**).

The N-terminal region of Atg11 is critical for membrane tethering

Our results till now show that Atg11 binds membranes via a putative amphipathic helix located in

its N-terminus and that it can self-associate to form clusters. Additionally, the interaction between

Atg11 and Atg32 leads to membrane tethering in vitro. Therefore, to test if the Atg11-Nterm

contributes to tethering of autophagosomal-like liposomes, we carried out GLT using GFP-Atg32$_{1-}$

$_{381}$ bound OMM-GUVs and automix+PI3P liposomes with Atg11 (1µM) alone and Atg11(1µM)

mixed with 20-fold molar excess of Atg11-Nterm (20µM) (**Figure 5A**). We reasoned that excess

Atg11-Nterm would compete with the full-length Atg11 and inhibit tethering of automix+PI3P

liposomes to GUVs. As a control we also carried out the GLT assay using Atg11 (1µM) mixed

with 20-fold molar excess of GST (20µM). Quantification of the fluorescence intensity of

liposomes normalized to that of GFP-Atg32$_{1-381}$ showed that the addition of excess Atg11-Nterm

led to a near complete block in automix+PI3P liposome tethering (**Figure 5B**). Atg11 mixed with

an excess of GST had only a modest reduction in the tethering of automix+PI3P liposomes. This

lends further support to the idea that autophagosomal-like liposome tethering by Atg11 occurs via

membrane binding by the N-terminal region of Atg11.

Next, we compared the liposome tethering abilities of full-length Atg11 with that of the Atg11-

Nterm. Full-length Atg11 was mixed with GFP-Atg32$_{1-381}$ bound OMM-GUVs and subsequently

incubated with automix+PI3P liposomes (**Figure 5C,** top panel). Because the Atg32 binding

interface lies at the C-terminus of Atg11, we directly recruited Atg11-Nterm to NTA-containing

OMM-GUVs and subsequently added automix+PI3P liposomes. GUV-bound Atg11-Nterm was

also able to tether liposomes further supporting the role of the Atg11-Nterm in membrane binding

and tethering (**Figure 5C,** bottom panel). Intriguingly, while liposomes tethered via full-length

Atg11 showed a punctate distribution, liposomes tethered via Atg11-Nterm uniformly coated the

GUVs (**Figure 5C** zoom images). Fluorescence intensity traces along the GUVs revealed that

automix+PI3P liposomes tethered by full-length Atg11 showed discreet peaks. In comparison,

liposomes tethered via Atg11-Nterm had a lower and more uniformly distributed intensity (**Figure**

5C). To quantitate the distribution of liposomes, we compared the coefficient of variance (COV)

of liposome fluorescence, calculated as standard deviation by mean, which serves a measure for

the extent of liposomes clustering. We found that the COV for Atg11-Nterm is significantly lower

than the COV for full-length Atg11 (**Figure 5D)**, suggesting that while OMM-GUV bound Atg11-

Nterm alone can tether autophagosomal-like liposomes, it fails to cluster these liposomes.

Importantly, the liposome fluorescence per micron calculated along the surface of GUVs was

significantly higher for full-length Atg11 than Atg11-Nterm, suggesting that the clustering by

Atg11 enhances the tethering of autophagosomal-like liposomes (**Figure 5E**).

299   DISCUSSION

Our results demonstrate, for the first time, that Atg11 can bind highly curved membranes

containing an autophagosomal-like lipid composition. Intriguingly, we observed that Atg11

membrane binding occurred in a punctate manner and that this clustering of Atg11 could be

recapitulated in the absence of membrane. We also identified a putative amphipathic helix in the

N-terminal region of Atg11 as the primary membrane binding interface. We show that the deletion

of this helix results in a delay in the formation of mitophagy initiation sites in yeast by monitoring

the formation of Atg32 puncta in real-time. To further explore the role of Atg11 membrane

binding, we used a method we recently developed—GLT—to reconstitute membrane tethering in

vitro[20].   Using GLT, we demonstrated that Atg11, in complex with Atg32, recruits

autophagosomal-like liposomes to OMM-GUVs. Comparing the tethering properties of Atg11

full-length and Atg11-Nterm, we show that the ability of Atg11 to cluster enhances liposome

tethering and causes clustering of tethered liposomes on OMM-GUVs. Taken together our work

suggests a possible role for Atg11 in the recruitment or organization of Atg9 vesicles during

selective autophagy.

The interaction between Atg9 and Atg11 was previously shown to be only partially required for

the recruitment of Atg9 vesicles to cargo in yeast[8]. Our results showing that Atg11 can bind

autophagosomal-like membranes directly and tether them to OMM-GUVs, could provide a

mechanistic explanation to this observation. Given these results it is possible that mitochondrial-

recruited Atg11 may tether Atg9 vesicles to mitochondria through dual interactions—binding both

Atg9 and autophagosomal membranes. While the deletion of the membrane binding helix of Atg11

results in a delay in the formation of Atg32 puncta, this deletion did not significantly hamper

delivery to the mitochondrial protein OM45 to the vacuole at 2 hours post mitophagy induction.

Therefore, we propose that Atg11 membrane binding via its amphipathic helix is critical for the

initial recruitment of Atg9 vesicles during the early stages of mitophagy (**Figure 6**). Once the low-

affinity protein interactions at mitophagy initiation sites are stabilized via avidity, membrane

binding via the amphipathic helix may become less significant. During this initial phase, Atg11

membrane association via the amphipathic helix may: 1) work synergistically with Atg11-Atg9

interaction to enhance or organize Atg9 vesicle recruitment, or 2) act as co-incident targeting

mechanism ensuring selective binding of Atg9 vesicles specifically at autophagy initiation sites.

However, which of these possibilities is correct will need to be further explored in subsequent

work.

We also demonstrate that GLT can be used for reconstituting the earliest steps of organelle

selective autophagy. We propose that GLT is a useful alternative to other reconstitution methods

that rely on selective autophagy receptors immobilized on Sepharose or agarose beads which may

more accurately mimic protein-based cargos such as aggregated proteins[9,42]. In contrast to

Sepharose or agarose-based reconstitution methods, selective autophagy receptors immobilized on

GUVs can diffuse freely along the bilayer and provide a more biologically relevant system for

studying organelle selective autophagy initiation where integral membrane selective autophagy

receptors such as Atg32 can diffuse along the cargo membrane.

Our work demonstrating that Atg11 can directly interact with membranes contrasts a previous

study reporting that Atg11 does not bind membranes[7]. However, two key differences could account

for the observed discrepancy in the membrane binding abilities of Atg11. First, we purified Atg11

using Freestyle 293 cells as a recombinant expression system where Atg11 appears to be a dimer

similarly to SF9 insect cells, while the Matscheko et al. study used monomeric Atg11 purified

from *E. coli* (**Figure S1B**). Second, we used liposomes containing YPL and autophagosomal-like

lipid compositions. Both mixtures contain approximately 42% negatively charged lipids. In

contrast, monomeric Atg11 was previously shown to be unable to bind liposomes composed of

60% POPC, 20% Cholesterol, 10% POPE and 10% POPS which contain only 10% negatively

charged lipids[7]. We have previously demonstrated that Atg13, a member of the autophagy

initiation machinery, binds liposomes containing at least 42.5% negatively charged lipids[20,43].

These findings may also suggest a common mechanism among autophagy proteins, where

membrane association occurs preferentially with the highly negatively charged autophagosomal

membranes, rather than the less charged membrane of the cargo. This differential lipid composition

may facilitate the spatial organization of autophagy proteins during autophagosome expansion.

Our data shows that the deletion of the amphipathic helix in Atg11 reduces but does not

completely abolish membrane binding. Interestingly, the mammalian homolog of Atg11, FIP200,

has been shown to interact with membranes via its large coiled-coil (CC) domain[17]. Our

experiments with SMrT tubes show a weak membrane binding for the Atg11 C-terminus, raising

the possibility that Atg11 has additional weaker membrane binding regions located in its C-

terminus. Given the enhanced recruitment and clustering of liposomes observed in the presence of

full-length Atg11 but not Atg11-Nterm it is possible that secondary membrane binding sites might

further strengthen Atg11 membrane binding and clustering.

While this work was being completed, a recent study demonstrated that the diffusion of selective

autophagy receptors on the cargo surface promotes the recruitment and phase separation of Atg11

into liquid-like droplets. These phase-separated assemblies, stabilized by dynamic weak

interactions, are proposed to stabilize the initiation machinery through avidity while providing

spatio-temporal regulation to phagophore initiation[44]. Our findings on Atg11 clustering on

autophagosomal membranes, and the enhancement of autophagosomal-like liposome tethering to

OMM-GUVs by Atg11 clustering, are in close agreement with this model. Intriguingly, the

properties of Atg11 clustering that we observed appear to be similar to those reported in Licheva

et al. in that Atg11 began to form significant clusters at 0.5 µM in 150 mM NaCl (**Figure 3**).

However, Atg11 clusters we observed are much smaller, less circular and do not appear to be

liquid-like as was described in Licheva et al. This may be due to differences in the expression

systems used or the imaging conditions. We purified Atg11 using the Freestyle 293 expression

system, whereas Licheva et al. purified Atg11 from SF9 insect cells. Additionally, our imaging

was performed on glass coverslips, while Licheva et al. used a six-channel chamber. Despite these

differences both our study and the Licheva et al. study point towards the importance of Atg11

higher order assemblies (either clustering or phase separation) in the initiation of selection

autophagy.

There is now a significant body of work that highlights the clustering of selective autophagy

receptors into high order assemblies to facilitate selective autophagy initiation[5]. In addition,

several studies also demonstrate a critical role for higher order assembly, primarily phase

separation, for the assembly of the autophagy initiation machinery[37,44]. Our work extends this

understanding by demonstrating that Atg11 can form clustered scaffolds on membranes. The

clustering of Atg11 also enhances the recruitment of and organization of small autophagosomal-

like vesicles on membrane-based cargo mimetics. Therefore, it is plausible that weak multivalent

interactions of Atg11 with the membrane, with itself and other autophagy proteins including

selective autophagy receptors may all contribute to the recruitment and organization of membrane

during selective autophagy initiation.

396
Figure 1. Atg11 binds curved autophagosomal-like membranes. (**A**) Representative images
showing binding of GFP-Atg11 on an array of SMrT tubes and flat supported lipid bilayers (SLB).
Membranes (magenta) are made of autophagosomal-like lipid composition with PI3P
(Automix+PI3P, top panel) and without PI3P (Automix, bottom panel). Scale bar=5μm. (**B**)
Quantification of GFP-Atg11 fluorescence normalized to the membrane fluorescence from SMrT
tubes and SLB made with automix+PI3P and automix. A total 153 tubes or 2 to 4um^2 individual
patches of SLB (n=153) were analyzed across three independent repeats (N=3). Data represent
mean ± SD. **** indicates p<0.0001 calculated using one-way ANOVA with Šídák's multiple
comparisons test. (**C**) Representative fluorescence images showing the binding of GFP-Atg11-
Nterm and GFP-Atg11-Cterm on an array of tubes and SLB made of automix+PI3P. Scale
bar=5μm. (**D**) Quantification of GFP-Atg11-Nterm or -Cterm normalized to membrane

fluorescence for SMrT tubes and SLB made with automix+PI3P. A total of 138 to 150 tubes or 2

to 4um^2 individual patches of SLB (n≥138) were analyzed across three independent repeats (N=3).

Data represent mean ± SD. **** indicates p<0.0001 and ** indicates p=0.0086 calculated using

one-way ANOVA with Šídák's multiple comparisons test. Autophagosomal-like membranes with

PI3P (automix+PI3P) were made of 43.9mol% PC, 19mol% PE, 32mol% PI, 5mol% PI3P with

0.1mol% RhPE. Autophagosomal-like membranes without PI3P (automix) were made similarly

except for the omission of PI3P and increase of PI to 37mol%.

Figure 2. A putative amphipathic helix mediates autophagosomal membrane binding by Atg11 and contributes to mitophagy. (**A**) Helical wheel representation of the predicted amphipathic helix in Atg11-Nterm. (**B**) Representative images showing GFP-Atg11-Nterm or GFP-Atg11-Nterm$^{\Delta612-646}$ binding on SMrT tubes made with 43.9% PC, 19% PE, 32% PI, 5% PI3P with 0.1mol% RhPE (automix+PI3P). Scale bar=5μm. **C**) Quantification of GFP-Atg11-Nterm or GFP-Atg11-Nterm$^{\Delta612-646}$ fluorescence normalized by membrane fluorescence from **B**. Data

represent mean ± SD of 150 tubes (n=150) analyzed across three independent repeats (N=3). ****
indicates p<0.0001 calculated using the non-parametric Mann-Whitney's test. (**D**) *atg11Δ* cells
transformed with Atg11 or Atg11$^{Δ612-646}$ were grown in media containing lactate and shifted to
nitrogen starvation media (SD-N) to induce mitophagy. Western blot showing OM45-GFP at 0, 2,
4 and 6 hours after nitrogen starvation. Actin was used as a loading control. (**E**) Quantification of
the background subtracted free GFP band normalized to total OM45-GFP at 0, 2, 4 and 6 hours
after nitrogen starvation. Data represent mean ± SD from three independent repeats (N=3). ns
indicates non-significant analyzed using one-way ANOVA with Šídák's multiple comparisons test.
(**F**) Western blot showing the expression of 2xGFP-Atg11 and 2xGFP-Atg11$^{Δ612-646}$. (**G**)
Timelapse fluorescence images showing *atg11Δ* cells expressing GFP-Atg32 with Atg11 (top
panel) or Atg11$^{Δ612-646}$ (bottom panel). Cells grown in nutrient rich (SMD) media were attached to
a concanavalin A-treated coverslip, assembled in a flow cell maintained at 30°C. Nitrogen-
starvation medium (SD-N) was introduced under controlled laminar flow, and time-lapse images
were acquired three minutes after the flow initiation. Images were captured at 4 minutes interval.
Yellow arrowheads indicate the onset of GFP-Atg32 puncta formation. Scale bar=5μm. (**H**)
Quantification of cells subjected to SD-N showing percent of cells with at least one GFP-Atg32
punctum. Data represents mean ± SD of a total of 270 cells for Atg11 and 290 cells for Atg11$^{Δ612-}$
^{646} (n≥270) from three independent repeats (N=3). * indicates p=0.0151 to p=0.0487, ** indicates
p=0.0089, ns indicates non-significant calculated using multiple t tests.

Figure 3. Atg11 self-associates in a salt and concentration dependent manner. Purified GFP-Atg11 was diluted into different buffers and imaged by confocal microscopy. (**A**) GFP-Atg11 puncta formation at 1uM protein concentration in different NaCl concentrations (as indicated). Scale bar=5μm. (**B**) Quantification of A. (**C**) GFP-Atg11 puncta formation in 150 mM NaCl at different protein concentrations (as indicated). Scale bar=5μm. (**D**) Quantification of C. (**E**) GFP-Atg11 was diluted to 1 μM in 150 mM or 450 mM NaCl, then imaged. Samples were mixed to produce 300 mM NaCl at 1 μM protein concentration. (**F**) Quantification of E. (**G**) Representative images of clustering observed when GFP-Atg11 constructs (FL, Atg11-Nterm or Atg11-Cterm) were diluted to 1 μM in 150 mM NaCl and imaged as before. (**H**) Quantification of clustering observed when GFP-Atg11 constructs (FL, Atg11-Nterm or Atg11-Cterm) were diluted to 1 μM in 150 mM NaCl and imaged as before. Each experiment was conducted in 3 independent repeats, with 7 images acquired at each repeat, amounting to a total area of 30492 $μm^2$ per repeat. Puncta analyzed varied from ~30 to ~19,000, depending on the group. The number of puncta were

459 normalized to the area imaged, yielding particles per mm^2. The data are presented as means, with
460 error bars indicating standard deviation. Groups were compared using a one-way ANOVA test
461 with Tukey's correction for multiple comparisons. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, ****
462 $p < 0.0001$, ns is not significant.

Figure 4. Atg11 and Atg32 coordinate to tether curved autophagosomal membranes to cargo mimetics. (**A**) Schematic showing tethering of autophagosomal membranes to mitochondrial mimetic GUV reconstituted using the GLT assay. OMM-GUVs containing mitochondrial lipids and DGS-Ni^{+2}NTA (5 mol%) displaying the cytosolic domain of Atg32 (His-GFP-Atg32$_{1-381}$) are used as a mitochondria mimic to recruit Atg11. 35nm-automix+PI3P liposomes are then added to reconstitute heterotypic tethering by the Atg32-Atg11 complex. (**B**) Representative fluorescence images of GLT with GUVs displaying His-GFP-Atg32$_{1-381}$ which were sequentially incubated with

Atg11 and automix+PI3P liposomes (top panel) or automix+PI3P liposomes alone (bottom panel).
Scale bar=10μm. Yellow box marks the region used for tracing line profiles in C and D. (**C**)
Fluorescence intensity profiles from a line drawn along the GUV in B (top panel) showing
liposome and GFP-Atg32$_{1\text{-}381}$ fluorescence intensities along the GUV. (**D**) Fluorescence intensity
profiles from a line drawn along the GUV in B (bottom panel) showing liposome and GFP-Atg32$_{1\text{-}}$
$_{381}$ fluorescence intensities along the GUV. (**E**) Quantification of the ratio of liposomes to GFP-
Atg32$_{1\text{-}381}$ fluorescence from B. Data are presented as means ± SD from 3 independent repeats
(N=3). ****p<0.0001 analyzed using Mann-Whitney's test. n indicates total number of GUVs
analyzed per condition. (**F**) Representative image of a GUV displaying His-GFP-Atg32$_{1\text{-}381}$
sequentially incubated with Atg11 and 1μm-automix+PI3P liposomes. (**G**) Quantification of the
ratio of liposomes to GFP-Atg32$_{1\text{-}381}$ fluorescence from F. Merge images show an overlay of GFP-
Atg32$_{1\text{-}381}$ and liposomes fluorescence channels. Data are presented as means ± SD from three
independent repeats (N=3). n indicates the total number of GUVs analyzed per condition. ****
indicates p<0.0001 analyzed using non-parametric Mann-Whitney's test. OMM-GUVs were made
with 69.9% DOPC, 10% DOPE, 10% DOPS, 5% cardiolipin, 5% DGS-Ni^{+2}NTA, and 0.1% RhPE.
Automix+PI3P were made of 43.9% PC, 19% PE, 32% PI, 5% PI3P with 0.1% DiD. Liposomes
were either sonicated to obtain 30-50nm curved liposomes or extruded through a 1μm-filter.

Figure 5. Membrane binding by Atg11-Nterm and self-association promotes autophagosomal vesicle tethering. OMM-GUVs displaying GFP-Atg32$_{1-381}$ were incubated with Atg11 alone, Atg11 with 20-fold molar excess of Atg11-Nterm or GST and subsequently mixed with autophagosomal liposomes. (**A**) Representative images of autophagosomal liposome recruitment to OMM-GUVs with Atg11 alone, Atg11 with excess of Atg11-Nterm or Atg11 with excess of GST. Scale bar=10μm. (**B**) Quantification of the ratio of liposomes to GFP-Atg32$_{1-381}$ fluorescence from A. Data from three independent experiments (N=3) are presented as mean ± SD. **** indicates p<0.0001 calculated using one-way ANOVA with Šídák's multiple comparisons test. n indicates the total number of GUVs analyzed per condition. (**C**) Representative images of autophagosomal liposome recruitment to OMM-GUVs displaying GFP-Atg32$_{1-381}$ with

Atg11 or GFP-Atg11-Nterm alone (recruited via 6X-His tag). Scale bar=10μm. Fluorescence
intensity profiles for the GUV and liposome channels along the circumference of GUVs from the
zoomed inset are shown to the right of each representative image. (**D**) Quantification of the
coefficient of variance (COV) for liposome fluorescence along the circumference of the GUV from
C. Data are presented as mean ± SD from three independent repeats (N=3). n indicates the total
number of GUVs analyzed per condition. **** indicates p<0.0001 analyzed using the non-
parametric Mann-Whitney's test (**E**) Quantification of the ratio of liposomes to GFP-Atg32$_{1-381}$
fluorescence from C. Data are presented as mean ± SD from three independent repeats (N=3). n
indicates the total number of GUVs analyzed per condition. **** indicates p<0.0001 analyzed
using non-parametric Mann-Whitney's test. OMM-GUVs were made with 69.9% DOPC, 10%
DOPE, 10% DOPS, 5% cardiolipin, 5% DGS-Ni^{+2}NTA, and 0.1% RhPE. Sonicated
Automix+PI3P were made of 43.9% PC, 19% PE, 32% PI, 5% PI3P with 0.1% DiD.

Figure 6. Proposed role of Atg11 membrane binding in mitophagy initiation. In response to nitrogen starvation, Atg32 clusters and recruits Atg11 on mitochondria. An amphipathic helix in Atg11 engages with negatively charged lipids in Atg9 vesicles. This membrane binding may serve to enhance, stabilize or organize the recruitment of Atg9 vesicles helping in rapid onset of mitophagy.

MATERIALS AND METHODS

Recombinant protein expression and purification

All constructs used for protein purification in this paper are listed in **Table S1**. StrepII-GFP-Atg32(1-381)-10XHis were recombinantly expressed in *E. coli* Star cells (Invitrogen C601003). Cultures were grown in LB broth (Fisher Scientific, BP1426) to an OD600 of 0.6 to 0.8 at 37°C while shaking at 210 rpm. Cultures were cooled at 4°C for 20 min. Protein expression was induced by adding 1 mM isopropylthio-β-D-galactopyranoside (IBI Scientific, IB02125) and the cultures were grown for an additional 18 h at 18°C. Cells were harvested and cell pellets were stored at −80°C. For purification of StrepII-GFP-Atg32(1-381)-10XHis, cell pellets were thawed and resuspended in 50 mM Tris pH 8.0, 500 mM NaCl, 2.5mM imidazole, 5 mM $MgCl_2$, 1 mM phenylmethanesulfonyl fluoride (PMSF; VWR, 0754) and cOmplete Mini EDTA-free protease inhibitor tablets (Roche, 11836170001). Cells were lysed using 3 strokes of a microfluidizer at 18, 000 psi. Triton X- 100 (VWR, AAA16046-AE), was subsequently added to lysates at 1% vol/vol. Lysates were spun at 25,000 g for 30 minutes. The supernatant was then applied to pre-equilibrated Talon resin (Clontech, 635504). The resin was washed with 50 mM Tris pH 8.0, 500 mM NaCl, 10 mM imidazole. Bound proteins were eluted with 50 mM Tris pH8.0, 500 mM NaCl, 250 mM imidazole. Elutions from Talon resin were pooled and incubated with pre- equilibrated Strep resin (IBA, 2-5030-025). The resin was washed with 20 mM Tris pH 8.0 and 300 mM NaCl. Protein was eluted with 20 mM Tris pH 8.0, 300 mM NaCl and 50 mM biotin. Peak fractions containing protein were dialyzed against 20 mM Tris pH 8.0, 300 mM NaCl and 0.2 mM TCEP. Dialyzed protein was aliquoted, flash-frozen in liquid nitrogen and stored at −80°C until use.

Atg11 and GFP-Atg11, both with an N-terminal 2XStrep II tag, were recombinantly expressed in

Freestyle cells for 3 days post transfection. Cells were harvested by spinning by 1000 g for 15 min.

Cell pellets were flash frozen in liquid nitrogen and stored at −80°C.

Cell pellets were resuspended and thawed in 50 mM Tris pH 8.0, 500 mM NaCl, 1 mM DTT, 1

mM PMSF and a cOmplete Mini protease inhibitor tablet (Roche, 11836170001). Cells were lysed

using 3 strokes of a microfluidizer at 10, 000 psi. Triton X-100 (VWR, AAA16046-AE), was

subsequently added to lysates at 1% vol/vol. Lysates were spun at 25,000 g for 30 min. Supernatant

was applied to Strep resin (IBA, 2- 5030-025) at 1 mL/min using a peristaltic pump. Resin was

washed with 20 mM Tris pH 8.0, 300 mM NaCl and 1 mM DTT. Bound protein was eluted with

20 mM Tris pH 8.0, 300 mM NaCl, 50 mM biotin and 1 mM DTT. Fractions containing protein

were pooled and concentrated to 2 μM and flash-frozen in liquid nitrogen. For GFP-Atg32 binding

experiments, Atg11 was fluorescently labeled with a three-fold molar excess of Alexa Fluor™ 594

C5 Maleimide (Invitrogen, A10256) for 1 h at room temperature. Labelling was quenched using 1

mM DTT. Unreacted dye was removed by dialyzing the labelled protein against 20 mM Tris pH

8.0, 300 mM NaCl, 0.2 mM TCEP and 1 mM DTT for 48 h. Fresh buffer was added after 24 h of

dialysis.

Yeast strains, plasmids and culture conditions

Δ*atg11*cell line was obtained from the *S. cerevisiae* knockout collection (Invitrogen). TKYM22

(SEY6210 OM45-GFP::TRP1) was a gift from Daniel Klionsky[45]. TKYM22 atg11Δ::NAT

(XXY003) was generated by replacing atg11 with the NAT cassette from pAG25 in TKYM22[27].

GFP-Atg32 and RFP-Atg32 (pRS415-MET25) plasmids were kind gifts from Dr. Benedikt

Westermann[46]. Plasmid expressing 2XGFP-Atg11 (yCPLAC33) and Atg11 (yCPLAC33) were

reported earlier[27]. Atg11$^{\Delta612-646}$ was generated from Atg11 (yCPLAC33) by site directed

mutagenesis. Cells were grown in SMD (0.67% yeast nitrogen base, 2% glucose and supplemented

with the appropriate amino acid and vitamin mixture).

Live imaging of GFP-Atg32 puncta formation

For GFP-Atg32 puncta formation experiments, cells expressing GFP-Atg32 under the MET25

promoter were grown to mid-log phase in SMD supplemented with 10 mg/mL methionine. GFP-

Atg32 expression was then induced by shifting cells to SMD without methionine for 1hr.

Subsequently, cells were transferred onto a concavalinA-coated glass coverslip assembled in a

temperature- and flow-controlled FCS2 chamber (Bioptechs). Chamber temperature was

maintained at 30 °C. To induce mitophagy, SD-N (0.17% yeast nitrogen base without amino acids

and ammonium sulfate; 2% glucose) prewarmed to 30 °C was flowed into the chamber at a

constant flowrate. Chamber was equilibrated with SD-N for 3 mins. Subsequently, timelapse

images were acquired at every 4 mins interval for a total of 30-40 mins.

OM45-GFP processing assay

The OM45-GFP processing assay was performed as described previously[31]. XXY003 cells were

transformed with Atg11 and Atg11$^{\Delta612-646}$. Cells were grown at 30 °C in SMD to an OD600 of 0.8–

1.0. Cells were pelleted at 1000g for 10 min, washed and resuspended in SML (0.67% yeast

nitrogen base, 2% lactic acid and supplemented with the appropriate amino acid and vitamin

mixture) and grown for 16 h. Subsequently, cells were pelleted and resuspended in SD-N for 6 h.

Finally, cells were harvested, lysed by bead beating and GFP was detected by western blot.

Liposome preparation

All lipids were purchased from Avanti Polar lipids. Chloroform stocks of lipid mixtures at a desired

ratio containing 0.1 mol% of the fluorescent lipophilic dye DiD were aliquoted in a clean glass

tube and dried under a gentle nitrogen stream until all the chloroform had evaporated. Residual

chloroform was removed under vacuum for 18 h. Dried lipids were resuspended in 20 mM Tris

pH 8.0, 100 mM NaCl to a final concentration on 1 mM. Automix+PI3P liposomes were sonicated

using a probe sonicator at 10% amplitude for 5 mins with 1 sec on and 2 sec off. Sonication was

carried out in an ice-water bath to prevent heating. 1-μm liposomes were prepared using an Avanti

Mini Extruder using 1-μm Nuclepore Track-Etched membrane filters (Sigma Aldrich,

WHA10417104).

Giant unilamellar vesicle preparation

GUVs were made using the gel-assisted formation method[51]. Briefly, chloroform stocks containing

lipids mixed at desired ratios containing the fluorescent lipid RhPE were aliquoted in a glass tube.

Chloroform was dried under a stream of nitrogen followed by drying under vacuum (1 h). The

dried lipids were resuspended in fresh chloroform to make a 1 mM lipid mixture. Polyvinyl alcohol

(PVA, Mw = 14500; Sigma, 814894) was dissolved in boiling water to make a 5% stock and

degassed in a vacuum chamber. PVA solution (10 μL) was spread into a thin film on a glass

coverslip kept on a heating block set to 55°C. Chloroform-dissolved lipid mix (10 μL) was then

spread on the dried PVA film. The dried PVA-lipid films were peeled from the glass coverslips

and transferred to a 1.5-mL microcentrifuge tube. Films were hydrated in 300 μL assay buffer (20

mM Tris pH 8, 100 mM NaCl) for 20 min, and GUVs were released by gentle tapping. The PVA

films were removed using a pipette.

GUV liposome tethering (GLT)

For GLT experiments, GUVs (120 to150 μL) were first mixed with tethering protein or control

protein in a 1.5 mL microcentrifuge tube for 10 min at room temperature. GUV mixtures were first

tapped to resuspend settled GUVs and subsequently incubated with 10 μL of liposomes (1mM)

for 5mins at room temperature. Liposomes were used within 3 days of extrusion or sonication. The

GLT mixtures (300 μL total volume) were transferred to a BSA-passivated LabTek chamber

(Nunc, Z734853) for imaging. GUVs were allowed to settle for 5 min prior to imaging. LabTek

chambers were treated with 3 M NaOH for 10 min followed by extensive washes with milliQ

water. NaOH-treated chambers were then passivated by incubation with BSA (3 mg/mL). BSA

was removed by three washes with the assay buffer. GLT experiments with Atg11 were performed

in 20 mM Tris pH 8, 100 mM NaCl. All proteins were spun at 100,000 g prior to use to remove

any aggregates.

Atg11 binding on SMrT tubes

SMrTs were prepared as described earlier[22,23]. Briefly, 3 µL of chloroform-dissolved automix and

automix+PI3P stocks (1mM) were spread on a PEGylated coverslip. Chloroform was dried using

a gentle stream of nitrogen. A ~35 µL flow cell (Bioptechs) was assembled by placing a 0.1 mm

silicone spacer between the PEGylated coverslip and an ITO-coated slide. Rehydration of the dried

lipid film with 20mM Tris pH 8.0 and 100mM NaCl yielded supported lipid bilayers and vesicles.

Lipid nanotubes were extruded from the vesicles using buffer flow. Subsequently, 300 µL of GFP-

Atg11 (0.5 µM), GFP-Atg11-Nterm (2 µM) or GFP-Atg11-Cterm (2 µM) were incubated with

SMrTs for 10 mins. Excess protein was washed with 3x300 µL of buffer (20mM Tris pH 8 and

100mM NaCl) and SMrTs were imaged. All proteins were spun at 100,000 g prior to use to remove

any aggregates.

Atg11 Clustering Assay

All buffers used for these assays contained 20 mM Tris pH 8.0 at 4°C and the indicated NaCl

concentrations. Purified GFP-Atg11 was thawed on ice and centrifuged at 20,000 x g for 15

minutes at 4°C. GFP-Atg11 was diluted to a concentration of 1 µM in buffer containing 50, 100,

150, 300 or 450 mM NaCl, or to a concentration of 0.05, 0.1, 0.5, 1 or 5 µM in buffer containing

150 mM NaCl. After dilution 40 µl of sample was spotted onto a glass coverslip, mounted on a 63

X oil immersion objective on a Zeiss LSM 880 microscope. Samples were left on the microscope

for 5 minutes before imaging. All images were acquired within 10 minutes after the sample was

applied to the coverslip to prevent evaporation and thus concentration of the sample. For

reversibility assays, GFP-Atg11 was diluted to 1µM in buffer containing 150 mM NaCl or 450

mM NaCl and imaged as described above. The remaining samples of 1µM GFP-Atg11 in 150 mM

NaCl and 450 mM NaCl were mixed in a 1:1 ratio to generate 1µM GFP-Atg11 in 300 mM NaCl.

For the comparison of GFP-Atg11, GFP-Atg11-Nterm and GFP-Atg11-Cterm protein was diluted

to 1 µM in buffer containing 150 mM NaCl and imaged as described above.

Fluorescence microscopy

Atg11 clustering was imaged on a Zeiss Airyscan LSM 880 microscope using 63 X oil immersion

objective using Zeiss Immersol 518 F immersion oil at room temperature. GFP was excited using

a 488 argon laser. Emission was detected using a Zeiss GaAsP-PMT detector.

All other images were acquired using a Nikon TiE microscope fitted with a Yokogawa CSU-W1

spinning disk system using a 100 X, 1.45 NA oil immersion objective. Images were acquired with

a Photometrics Prime BSI sCMOS camera using the NIS elements software.

Image analysis

Fluorescence microscopy images were analyzed using Fiji[52].

To analyze tethering in GLT, images were first background corrected by subtracting the mode

intensity value. Percent liposome fluorescence per length of GUV was calculated by normalizing

the liposome fluorescence, collected from a two-pixel wide segmented line drawn along a GUV,

to the length of the GUV. For calculating the liposome/protein fluorescence ratio, fluorescence

intensities in liposome and protein channels from a two-pixel wide segmented line drawn along a

GUV were collected. Mean liposome fluorescence was divided by mean protein fluorescence to

yield liposome/protein fluorescence ratios.

For analysis of protein bound on SMrTs, images were background corrected by subtracting the

mode value. Integrated density of protein and membrane fluorescence was measured from a 20x10

pixel rectangle placed centrally on the tubes or from a 2 to 4 μm^2 square box placed on SLBs.

GFP-Atg32 puncta were analyzed by placing a segmented line centered on the punctum. Only

those GFP-Atg32 puncta that were over 2.5-fold intensity of the background mitochondrial

fluorescence intensity were counted.

For Atg11 clustering analysis images were analyzed in Fiji. A threshold was set for each image

based on the average background intensity of the image. The analyze particle command was used

with parameters of 0.2 µm to infinity for the particle size and 0.8-1.0 for the circularity.

Statistical analysis

Statistical analysis was performed using GraphPad Prism (version 5.0a).

All experiments were performed in independent triplicates or duplicates. Liposomes and GUVs

were prepared independently for each GLT experiment using freshly aliquoted lipids. Data from

all repeats were pooled and plotted as scatterplots. Error bars represent standard deviation.

Significance was calculated using the non-parametric Mann-Whitney's test, one-way ANOVA

with Tukey's correction for multiple comparisons or Šídák's multiple comparisons test, or t test as

appropriate.

Author Contributions

D.A., S.K., K.M.B., and M.J.R. conceived and planned the experiments. D.A., S.K., S.I.N. and

K.M.B. carried out the experiments. D.A. and M.J.R. drafted the manuscript. All authors edited

the manuscript.

Acknowledgements

This work was supported by the National Institute of General Medical Sciences (NIGMS) grant

R35GM128663 to M.J.R. Expression of Atg11 was performed in the BioMT service center and

protein purification was performed in part performed at the BioMT Molecular Tools Core which

are both supported by NIGMS grant P20GM113132. Sequencing of plasmids was performed by

the Molecular Biology Shared Resource Center which is supported by NCI Cancer Center Support

Grant P30CA023108. We would also like to thank Ann Lavanway for her work and support at the

Dartmouth Imaging Facility. We thank Dr. Benedikt Westermann for the kind gift of GFP-Atg32

plasmid.

Disclosure Statement

707   The authors have no conflicts of interest to declare.

REFERENCES

1. Wen X, Klionsky DJ. An overview of macroautophagy in yeast. J Mol Biol 2016; 428:1681-99. doi:10.1016/j.jmb.2016.02.021.
2. Germic N, Frangez Z, Yousefi S, Simon HU. Regulation of the innate immune system by autophagy: monocytes, macrophages, dendritic cells and antigen presentation. Cell Death Differ 2019; 26:715-27. doi:10.1038/s41418-019-0297-6.
3. Dikic I, Elazar Z. Mechanism and medical implications of mammalian autophagy. Nat Rev Mol Cell Biol 2018; 19:349-64. doi:10.1038/s41580-018-0003-4.
4. Lamark T, Johansen T. Mechanisms of Selective Autophagy. Annu Rev Cell Dev Biol 2021; 37:143-69. doi:10.1146/annurev-cellbio-120219-035530.
5. Adriaenssens E, Ferrari L, Martens S. Orchestration of selective autophagy by cargo receptors. Current Biology 2022; 32:R1357-R71. doi:10.1016/j.cub.2022.11.002.
6. Yorimitsu T, Klionsky DJ. Atg11 links cargo to the vesicle-forming machinery in the cytoplasm to vacuole targeting pathway. Mol Biol Cell 2005; 16:1593-605. doi:10.1091/mbc.e04-11-1035.
7. Matscheko N, Mayrhofer P, Rao Y, Beier V, Wollert T. Atg11 tethers Atg9 vesicles to initiate selective autophagy. PLoS Biol 2019; 17:e3000377. doi:10.1371/journal.pbio.3000377.
8. Coudevylle N, Banas B, Baumann V, Schuschnig M, Zawadzka-Kazimierczuk A, Kozminski W, et al. Mechanism of Atg9 recruitment by Atg11 in the cytoplasm-to-vacuole targeting pathway. J Biol Chem 2022; 298:101573. doi:10.1016/j.jbc.2022.101573.
9. Sawa-Makarska J, Baumann V, Coudevylle N, von Bülow S, Nogellova V, Abert C, et al. Reconstitution of autophagosome nucleation defines Atg9 vesicles as seeds for membrane formation. Science 2020; 369:1206-+. doi:ARTN eaaz7714 10.1126/science.aaz7714.
10. Mari M, Griffith J, Rieter E, Krishnappa L, Klionsky DJ, Reggiori F. An Atg9-containing compartment that functions in the early steps of autophagosome biogenesis. J Cell Biol 2010; 190:1005-22. doi:10.1083/jcb.200912089.
11. Melia TJ, Lystad AH, Simonsen A. Autophagosome biogenesis: From membrane growth to closure. J Cell Biol 2020; 219. doi:10.1083/jcb.202002085.
12. Maeda S, Otomo C, Otomo T. The autophagic membrane tether ATG2A transfers lipids between membranes. Elife 2019; 8. doi:ARTN e45777 10.7554/eLife.45777.
13. Valverde DP, Yu S, Boggavarapu V, Kumar N, Lees JA, Walz T, et al. ATG2 transports lipids to promote autophagosome biogenesis. J Cell Biol 2019; 218:1787-98. doi:10.1083/jcb.201811139.
14. Jiang W, Chen X, Ji C, Zhang W, Song J, Li J, et al. Key Regulators of Autophagosome Closure. Cells 2021; 10. doi:10.3390/cells10112814.
15. He C, Song H, Yorimitsu T, Monastyrska I, Yen WL, Legakis JE, et al. Recruitment of Atg9 to the preautophagosomal structure by Atg11 is essential for selective autophagy in budding yeast. J Cell Biol 2006; 175:925-35. doi:10.1083/jcb.200606084.
16. Meyer MD, Winzeler J, Taylor SM, Kilgore A, Edicha K, Chitwood C, et al. Mapping Critical Residues in ATG11's Coiled-Coil 2 Domain that Block Multiple Interactions and Disrupt Selective Autophagy. Front Cell Dev Biol 2021; 9:775364. doi:10.3389/fcell.2021.775364.
17. Shi X, Chang C, Yokom AL, Jensen LE, Hurley JH. The autophagy adaptor NDP52 and the FIP200 coiled-coil allosterically activate ULK1 complex membrane recruitment. Elife 2020; 9. doi:10.7554/eLife.59099.
18. Suzuki H, Noda NN. Biophysical characterization of Atg11, a scaffold protein essential for selective autophagy in yeast. FEBS Open Bio 2018; 8:110-6. doi:10.1002/2211-5463.12355.
19. Schutter M, Giavalisco P, Brodesser S, Graef M. Local Fatty Acid Channeling into Phospholipid Synthesis Drives Phagophore Expansion during Autophagy. Cell 2020; 180:135-49 e14. doi:10.1016/j.cell.2019.12.005.

20.   Hawkins WD, Leary KA, Andhare D, Popelka H, Klionsky DJ, Ragusa MJ. Dimerization-dependent
membrane tethering by Atg23 is essential for yeast autophagy. Cell Rep 2022; 39:110702.
doi:10.1016/j.celrep.2022.110702.
21.   Popelka H, Reinhart EF, Metur SP, Leary KA, Ragusa MJ, Klionsky DJ. Membrane Binding and
Homodimerization of Atg16 Via Two Distinct Protein Regions is Essential for Autophagy in Yeast.
J Mol Biol 2021; 433:166809. doi:10.1016/j.jmb.2021.166809.
22.   Dar S, Kamerkar SC, Pucadyil TJ. A high-throughput platform for real-time analysis of membrane
fission reactions reveals dynamin function. Nat Cell Biol 2015; 17:1588-96. doi:10.1038/ncb3254.
23.   Dar S, Kamerkar SC, Pucadyil TJ. Use of the supported membrane tube assay system for real-time
analysis of membrane fission reactions. Nat Protoc 2017; 12:390-400. doi:10.1038/nprot.2016.173.
24.   Yamamoto H, Kakuta S, Watanabe TM, Kitamura A, Sekito T, Kondo-Kakuta C, et al. Atg9 vesicles
are an important membrane source during early steps of autophagosome formation. J Cell Biol 2012;
198:219-33. doi:10.1083/jcb.201202061.
25.   Shi X, Yokom AL, Wang C, Young LN, Youle RJ, Hurley JH. ULK complex organization in
autophagy by a C-shaped FIP200 N-terminal domain dimer. J Cell Biol 2020; 219.
doi:10.1083/jcb.201911047.
26.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein
structure prediction with AlphaFold. Nature 2021; 596:583-9. doi:10.1038/s41586-021-03819-2.
27.   Margolis HK, Katzenell S, Leary KA, Ragusa MJ. The Third Coiled Coil Domain of Atg11 Is
Required for Shaping Mitophagy Initiation Sites. J Mol Biol 2020; 432:5752-64.
doi:10.1016/j.jmb.2020.08.025.
28.   Turco E, Witt M, Abert C, Bock-Bierbaum T, Su MY, Trapannone R, et al. FIP200 Claw Domain
Binding to p62 Promotes Autophagosome Formation at Ubiquitin Condensates. Mol Cell 2019;
74:330-46 e11. doi:10.1016/j.molcel.2019.01.035.
29.   Gallop JL, Jao CC, Kent HM, Butler PJ, Evans PR, Langen R, et al. Mechanism of endophilin N-
BAR domain-mediated membrane curvature. EMBO J 2006; 25:2898-910.
doi:10.1038/sj.emboj.7601174.
30.   Nishimura T, Lazzeri G, Mizushima N, Covino R, Tooze SA. Unique amphipathic helix drives
membrane insertion and enzymatic activity of ATG3. Sci Adv 2023; 9. doi:ARTN
eadh128110.1126/sciadv.adh1281.
31.   Kanki T, Kang D, Klionsky DJ. Monitoring mitophagy in yeast: the Om45-GFP processing assay.
Autophagy 2009; 5:1186-9. doi:10.4161/auto.5.8.9854.
32.   Kanki T, Wang K, Cao Y, Baba M, Klionsky DJ. Atg32 is a mitochondrial protein that confers
selectivity during mitophagy. Dev Cell 2009; 17:98-109. doi:10.1016/j.devcel.2009.06.014.
33.   Okamoto K, Kondo-Okamoto N, Ohsumi Y. Mitochondria-anchored receptor Atg32 mediates
degradation of mitochondria via selective autophagy. Dev Cell 2009; 17:87-97.
doi:10.1016/j.devcel.2009.06.013.
34.   Furukawa K, Fukuda T, Yamashita SI, Saigusa T, Kurihara Y, Yoshida Y, et al. The PP2A-like
Protein Phosphatase Ppg1 and the Far Complex Cooperatively Counteract CK2-Mediated
Phosphorylation of Atg32 to Inhibit Mitophagy. Cell Rep 2018; 23:3579-90.
doi:10.1016/j.celrep.2018.05.064.
35.   Mao K, Wang K, Liu X, Klionsky DJ. The scaffold protein Atg11 recruits fission machinery to drive
selective mitochondria degradation by autophagy. Dev Cell 2013; 26:9-18.
doi:10.1016/j.devcel.2013.05.024.
36.   Hollenstein DM, Licheva M, Konradi N, Schweida D, Mancilla H, Mari M, et al. Spatial control of
avidity regulates initiation and progression of selective autophagy. Nat Commun 2021; 12:7194.
doi:10.1038/s41467-021-27420-3.
37.   Fujioka Y, Alam JM, Noshiro D, Mouri K, Ando T, Okada Y, et al. Phase separation organizes the
site of autophagosome formation. Nature 2020; 578:301-5. doi:10.1038/s41586-020-1977-6.

38. Sperka-Gottlieb CD, Hermetter A, Paltauf F, Daum G. Lipid topology and physical properties of the
outer mitochondrial membrane of the yeast, Saccharomyces cerevisiae. Biochim Biophys Acta 1988;
946:227-34. doi:10.1016/0005-2736(88)90397-5.
39. van Meer G, Voelker DR, Feigenson GW. Membrane lipids: where they are and how they behave.
Nat Rev Mol Cell Bio 2008; 9:112-24. doi:10.1038/nrm2330.
40. Decker ST, Funai K. Mitochondrial membrane lipids in the regulation of bioenergetic flux. Cell
Metab 2024; 36:1963-78. doi:10.1016/j.cmet.2024.07.024.
41. Aoki Y, Kanki T, Hirota Y, Kurihara Y, Saigusa T, Uchiumi T, et al. Phosphorylation of Serine 114
on Atg32 mediates mitophagy. Mol Biol Cell 2011; 22:3206-17. doi:10.1091/mbc.E11-02-0145.
42. Adriaenssens E, Schaar S, Cook ASI, Stuke JFM, Sawa-Makarska J, Nguyen TN, et al.
Reconstitution of BNIP3/NIX-mediated autophagy reveals two pathways and hierarchical flexibility
of the initiation machinery. bioRxiv 2024. doi:10.1101/2024.08.28.609967.
43. Gatica D, Damasio A, Pascual C, Klionsky DJ, Ragusa MJ, Popelka H. The carboxy terminus of
yeast Atg13 binds phospholipid membrane via motifs that overlap with the Vac8-interacting domain.
Autophagy 2020; 16:1007-20. doi:10.1080/15548627.2019.1648117.
44. Licheva M, Pflaum J, Babic R, Mancilla H, Elsasser J, Boyle E, et al. Phase separation of initiation
hubs on cargo is a trigger switch for selective autophagy. Nat Cell Biol 2025. doi:10.1038/s41556-
024-01572-y.
45. Kanki T, Klionsky DJ. Mitophagy in Yeast Occurs through a Selective Mechanism. Journal of
Biological Chemistry 2008; 283:32386-93. doi:10.1074/jbc.M802403200.
46. Bockler S, Westermann B. Mitochondrial ER contacts are crucial for mitophagy in yeast. Dev Cell
2014; 28:450-8. doi:10.1016/j.devcel.2014.01.012.
828
829