c-fos induction in the choroid plexus, tanycytes and pars tuberalis is an early indicator of spontaneous arousal from torpor in a deep hibernator

Fredrik A.F Markussen, Fernando Cázarez-Marquez, Vebjørn J. Melum, David Hazlerigg, Shona

Wood[*]

Arctic seasonal timekeeping initiative (ASTI), Arctic Chronobiology & Physiology, Arctic & Marine

Biology, BFE, UiT - Arctic University of Norway, Tromsø, Norway

* Correspondence: shona.wood@uit.no

Keywords: Hibernation, Thermoregulation, _c-fos,_ Golden hamster, choroid plexus, tanycytes,

dorsomedial hypothalamus, pars tuberalis

Summary statement

Non-neuronal sites linked to control of daily and seasonal metabolic physiology, peak in _c-fos_

expression during arousal from torpor, suggesting metabolic feedback-based initiation of the

spontaneous arousal process.

Abstract

Hibernation is an extreme state of seasonal energy conservation, reducing metabolic rate to as little as

1% of the active state. During the hibernation season, many species of hibernating mammals cycle

repeatedly between the active (aroused) and hibernating (torpid) states (T-A cycling), using brown

adipose tissue (BAT) to drive cyclical rewarming. The regulatory mechanisms controlling this

process remain undefined but are presumed to involve thermoregulatory centres in the hypothalamus.

Here, we use the golden hamster (*Mesocricetus auratus*), and high-resolution monitoring of BAT,

core body temperature (T_b), and ventilation rate, to sample at precisely defined phases of the T-A

cycle. Using *c-fos* as a marker of cellular activity we show that although the dorso-medial

hypothalamus (DMH) is active during torpor entry, neither it nor the pre-optic area (POA) show any

significant changes during the earliest stages of spontaneous arousal. Contrastingly, in 3 non-

neuronal sites previously linked to control of metabolic physiology over seasonal and daily

timescales, the choroid plexus (CP), pars tuberalis (PT) and third ventricle tanycytes, peak *c-fos*

expression is seen at arousal initiation. We suggest that through their sensitivity to factors in the

blood or cerebrospinal fluid (CSF), these sites may mediate metabolic feedback-based initiation of

the spontaneous arousal process.

Introduction

Hibernation is a seasonally regulated process of energy conservation achieved through extreme shut down of metabolic energy expenditure, sometimes to as little as 1% of that in the euthermic state, allowing species to survive extended winter periods of low energy availability (Kenagy et al., 1989; Mckee and Andrews, 1992; Ruf and Geiser, 2015). Hibernation is characterised by multi-day bouts of torpor during which metabolic rate is suppressed and core body temperature (T_b) falls, in some species to within a few degrees of ambient temperature (T_a), while breathing and heart rate slow to an extent not expected to support life in endothermic organisms (Lyman et al., 1982).

In many hibernating species, torpor bouts are separated by relatively short arousal episodes during which T_b rises back to euthermic levels, so that the hibernation season as a whole comprises a series of torpor – arousal (T-A) cycles. This presents a paradox, why would an organism undergo a costly arousal if the ultimate goal of hibernation is to save energy? Since homeostatic processes consume energy to maintain healthy cellular environments, one scenario is that arousal episodes allow restoration of healthy cellular conditions by reactivating aspects of homeostasis that are compromised or suspended during torpor.

While this narrative provides plausible ultimate reasoning for the evolution of periodic arousal, it does not address the proximate mechanisms by which T-A cycling is controlled. Here, two basic models can be considered: a clock-based mechanism and a metabolic feedback-based mechanism. In its most explicit formulation, the former is conceived as an extension of circadian function (Körtner and Geiser, 2000; Malan, 2010). The hypothalamic circadian pacemaker in the suprachiasmatic nucleus (SCN) controls daily T_b cycles in euthermic mammals and the expression of daily torpor episodes in Siberian hamsters (Körtner and Geiser, 2000; Ruby and Zucker, 1992; Ruf, T., Steinlechner, S., and Heldmaier, 1989). A role for circadian control mechanisms in multiday T-A

cycling is not favoured, however. Except for the first torpor bout of the season, T-A cycles show no

circadian organisation (Hut et al., 2002; Williams et al., 2012; Williams et al., 2017), and SCN

molecular clock gene oscillations flatten out (Ikeno et al., 2017; Revel et al., 2007). Malan proposed

a 'poorly temperature compensated circadian mechanism' in which slow running of the circadian

oscillation at low temperatures determines timing of arousals (Malan, 2010; Malan et al., 2018), but

this conjecture fails to account for changes in T-A cycle periodicity, particularly at the beginning and

end of the hibernation season, or for the finding that the period of T-A cycling is the same in wild-

type Syrian hamsters and in *tau* mutant hamsters in which the circadian clock runs fast (Oklejewicz

et al., 2001). Hence current models focus on the second possibility, namely that T-A cycle regulation

stems from accumulation (or depletion) during the torpid state of metabolite(s) which act as signals

to stimulate the arousal process, and that levels of these metabolic signals are then reset during

arousal permitting torpor re-entry (reviewed in:(van Breukelen and Martin, 2015)). Since arousal is

initiated by combined increases in pulmonary-cardio-vascular activity and brown adipose tissue

(BAT) activation, this framing places an emphasis on metabolic signals that access brain mechanisms

controlling sympathetic stimulation of these tissues.

A study in the Siberian chipmunk identified a "hibernation factor", produced in the liver and

transported to the brain via the choroid plexus (Kondo et al., 2006), indicating a link between

peripheral communication to the brain via the circulation and cerebrospinal fluid (CSF) in the

regulation of hibernation. However, this study did not investigate changes in hibernation factor levels

during T-A cycling, instead emphasizing a role as a putative seasonally permissive factor (Kondo et

al., 2006). The circuits controlling BAT thermogenic activity in euthermic rodents include the

temperature-sensitive pre-optic area (POA), which acts via the dorsomedial hypothalamus

(DMH)(Cannon and Nedergaard, 2004; Morrison, 2016). In mice, chemo/opto-genetic manipulation

of the activity of neurons within the POA can suppress T_b and metabolic rate (Hrvatin et al., 2020;

Takahashi et al., 2020; Upton et al., 2021) to produce a torpor-like state, suggesting that metabolic

feedback to these sites could be involved in natural torpor regulation.

To identify key signals controlling spontaneous T-A cycling, it is necessary to overcome the

challenge of repeatable sampling immediately prior to, and in the earliest stages of spontaneous

arousal from torpor. Because this is difficult, "forced" arousal by handling or changing room

temperature has been employed (Regan et al., 2019). Unfortunately this necessarily alters the

dynamics of the T-A cycle and causes stress (Tähti and Antti, 1977; Utz and van Breukelen, 2013),

potentially confounding attempts to characterize brain mechanisms involved in spontaneous T-A

cycling. Bratincsak *et al*. combined T_b measurement and expression of the immediate early gene *c-*

fos (Bratincsák et al., 2007) to explore brain changes during spontaneous T-A cycles in thirteen lined

ground squirrels. While this revealed *c-fos* changes in many brain sites, the use of T_b as the sole

indicator of arousal status inevitably misses the early stages of the arousal process, which may

precede core T_b by almost one hour (Markussen et al., 2020).

Here, we have developed an enhanced telemetric approach based on BAT temperature combined

with T_b and ventilation rate to characterize T-A cycling at high temporal and temperature resolution

in hibernating golden hamsters, a species showing highly reproducible T-A cycles (Chayama et al.,

2016; Lyman, 1948; Lyman et al., 1982). Using *c-fos* RNA expression, we sought to capture the

early dynamics of brain activation during spontaneous T-A cycling. Our data reveal changes in the

choroid plexus, pars tuberalis and tanycytes lining the 3^{rd} ventricle of the hypothalamus, as the

earliest brain indicators of initiation of spontaneous arousal.

Materials and Methods

Animals & Experimental set up

All experimental procedures were carried out at the animal facility of Arctic Chronobiology and Physiology at UiT – the Arctic University of Norway. All procedures and experiments were approved under national legislation by the Norwegian Food Safety Authority under the laboratory animal administration's supervision and application system ID: 24904. For 4 weeks, 5 months old male golden hamsters (n=30, *Mesocricetus auratus,* RJHan:AURA*)* purchased from Janvier Labs (Le Genest-Saint-Isle, France) were kept in sibling groups with *ad libitum* access to food and water, housed under long photoperiod (LP: 16L:8D) and 21°C. Prior to induction of hibernation, Anipill® temperature loggers (Animals-monitoring, Hérouville Saint-Clair, France) and IPTT-300 (BMDS/Plexx BV,Netherlands) tags were implanted under surgical anesthesia in the abdominal cavity and interscapular brown fat deposit, respectively. The animals were then single housed and allowed 2 weeks to recover in LP. To induce hibernation, single housed hamsters were switched to short photoperiod (SP: 8L:16D) for 4 weeks at 21°C (SP-warm) before lowering the ambient temperature to 6.5 °C (SP-cold).

Physiological and behavioural monitoring

To reduce stress and disturbance to the animals during the SP-cold phase, a select group of trained personnel conducted daily monitoring, thereby maintaining a stable and controlled environment conducive to animal welfare.

Real-time monitoring of interscapular BAT temperature (T_{iBAT}) was done using a mix of proprietary hardware and readily available electronics components using open-source practices, together with an open-source software stack. Iterative reads of T_{iBAT} were done using BMDS DAS-8027 (BMDS/Plexx BV, Netherlands) reader, and fixed in position near the scapular region of the

hibernating hamster. The reader was configured to operate in automatic scanning mode, recording the

IPTT-300 every 3-5 seconds. Data acquisition from the reader was wirelessly transmitted through

BMDS communications module to a computer in a separate room. The data was written into a text

document using Atom text editor (https://github.com/atom/atom) and automatically saved on change.

A program written using NodeRED v.19.9.0 (OpenJS Foundation) monitors this file and

automatically extracts the last row of information as soon as data is written, formats it, and pushes

the data into an InfluxDB v.1.8.1 (InfluxData Inc, San Francisco, CA, USA) database housed in the

local computer (JavaScript for replication of the NodeRED program is available here:

https://github.com/ShonaWood/cFOSGHam).   Ambient temperature (T_a) was also continuously

monitored using RuuviTag Bluetooth sensors (Ruuvi Innovations Ltd, Riihimäki, Finland), data were

pushed to Influx DB using NodeRed. InfluxDB data was accessed and displayed using Grafana

v.10.0.1 (GrafanaLabs, New York, NY, USA), this allowed real-time monitoring from any computer

or smart phone. We established a torpor baseline T_{iBAT} temperature by averaging T_{iBAT} values below

10°C and subtracting the T_a to calculate the thermal distance (TD). Subsequently we calculated the

change from this baseline (ΔT), therefore as the animal remains torpid the value remains at zero, even

with ambient temperature fluctuations. Mathematically this is simply expressed as:

$\quad \Delta T = T_{iBAT} - (T_a + TD) \mid TD = mean(T_{iBAT} < 10) - T_a).$

Departures from ΔT were then used to detect animals spontaneously arousing.

In parallel an infrared sensitive camera (Raspberry Pi Camera Module 2 NoIR, Raspberry Pi

Foundation, UK) connected via Raspberry Pi (model 3B+, Raspberry Pi Foundation, UK) was used

to stream a video feed to monitor and record animal activity and respiration rate using software

developed by (Singh et al., 2019).

We then analyzed twelve T-A cycles from 6 individuals to select physiological sampling criteria with

the goal of reducing inter-individual variation and identify the earliest possible reliable predictors of

arousal. Our criteria were;

Animals entering torpor (ENT<25°C) must demonstrate a minimum of a 12-hour inter bout

euthermic (IBE) duration, followed by a steady decrease in observed anipill T_b to below 25°C.

Additionally the IPTT read must be below 25°C and the animal must be in curled up torpid posture.

Torpid animals (T-40h) were sampled 40 hours after ENT<25°C, if Tb and T_{iBAT} were within 2°C of

T_a, and ventilation frequency (VF) was less than 2 per minute. Arousal 1 (AΔ0.5°C) was defined

only if an animal had entered torpor and remained torpid for longer than 40 hours, and subsequently

showed a VF >10 per minute and a 0.5°C change in T_{iBAT} from torpor baseline (ΔT). Arousal 2

(AΔ3°C) followed the same criteria as arousal 1 but was sampled at ΔT=3°C. IBE was defined as an

animal that had entered torpor, remained in torpor for longer than 40 hours, subsequently aroused and

then reached a T_b of 30°C. Animals were sampled 1 hour (IBE-1h) and 12 hours (IBE-12h) after

euthermia was reached.

Prior to sampling all animals had shown at least 3 full T-A cycles, these data were used to verify the

sampling criteria prior to sampling in the 4[th] T-A cycle. On reaching the above criteria the animals

were euthanized by cervical dislocation. The brain was rapidly dissected and frozen in isopentane

before storage at -80°C until sectioning.

***In-Situ* hybridization**

Regions of interest, preoptic area (POA, Bregma ca. 0.8 to 0.95) and mediobasal hypothalamus

(MBH, Bregma ca. -2.5 to -2.8), were collected in 14 µm sections onto Fisherbrand Superfrost Plus

slides using a Leica CM3050S cryostat and stored at -80°C until further processing.

On the day of hybridization, sections were dried for 15 min at 50°C before fixation with 4%

formaldehyde solution for 20 min. Sections were acetylated and delipidated using 1%

(volume/volume (v/v)) triethylamine (Sigma, 90335) with 0.25% (v/v) acetic anhydride (Sigma,

539996) and 0.1% (v/v) Triton-X-100 (Sigma, X100) in nuclease-free water, respectively. The

sections were hybridized with a 200 ng/mL *c-fos* antisense probe (A gift from Valerie Simonneaux

and Paul Klosen, University of Strasbourg, 793-1193 of Genbank accession: XM_005086369.4)

overnight at 58°C in hybridization buffer containing 5x Denhardt's solution (Sigma, D2532) , 0.25

mg/mL yeast tRNA (Roche, 10109525001), 0.2 mg/mL fish sperm ssDNA (Roche, 11467140001),

5x saline-sodium citrate (SSC, Sigma, 8310-OP) , and 50% (v/v) formamide (Sigma F9037) in

nuclease-free water (Sigma, F9037). Following hybridization, washes were made in decreasing

concentrations of SSC (5x and 2x) at 55°C, 50% formamide in 0.2x SSC at 55°C, and 0.2x SSC at

room temperature. After blocking using 1% Roche blocking solution (Roche, 11096176001), sections

were incubated for 3 hours with Anti-Digoxigenin-AP Fab fragments (dilution 1:3000) (Roche,

11093274910) at room temperature, followed by washes in A-dig and alkaline phosphate buffer. The

signal was then visualized by incubating the slides for 48 hours in a solution containing 0.33 mg/mL

NBT (Nitro-blue tetrazolium chloride, Roche, 11383213001) and 0.165 mg/mL BCIP (5-Bromo-4-

chloro-3-indolyl phosphate, Roche, 11383221001) in alkaline phosphatase buffer. Once the signal

had developed properly, the slides were prepared for immunohistochemical processing to visualize

vimentin and cell nuclei. After three five-minute washes in PBS pH 7.6 containing 0.05% (v/v)

Tween-20 (PBS-Tw, Sigma, P1379), the slides were incubated in blocking buffer containing PBS-

Tw, 0.1% (weight/v) cold water fish skin gelatin (Sigma, G7041), 0.1% (w/v) bovine serum albumin

(Sigma, A2153) , and 3 mM glycine (Sigma, G7126) for one hour at room temperature, followed by

overnight incubation at 4°C in blocking buffer containing mouse anti-vimentin antibody (Sigma,

MAB3400) diluted 1:2000. After four 10-minute washes in PBS-Tw at room temperature, the

secondary antibody was applied by incubating the slides for 1 hour at room temperature with

blocking buffer containing Alexa Fluor 647-conjugated goat anti-mouse antibody (1:1000, Sigma,

SAB4600354). The slides were then briefly rinsed in PBS-Tw before a 10-minute incubation in PBS-

Tw containing 1:2000 SYTOX Orange (Thermo Fisher, S11368), followed by four 10-minute washes

in PBS-Tw and a final rinse in PBS. The slides were then cover slipped and sealed using nail polish

and 1.5H coverslips over antifade solution containing 90% (v/v) glycerol (Sigma, G7757), 10% (v/v)

PBS, and 2.5% (w/v) diazabicyclo 2.2.2 octane (DABCO, Sigma, 290734)

The slides were scanned using an Olympus VS120 slide scanning microscope at the core microscopy

facility at UiT. Overview images of the sections were obtained using a Cy3 filter set, then regions of

interest were marked up in the OlyVIA software and, by using the x40 objective, high-resolution

images were obtained. The images were analyzed using QuPath v0.4.3 by annotating regions of

interest (ROIs). The number of cells within each ROI was detected and quantified using the SytoxO

channel in the standard cell detection plugin in QuPath. To quantify the amount of *c-fos* within each

ROI, a threshold was used to select the signal, and the *c-fos*-positive area within the ROI was

quantified by threshold. Cells are then classified as either positive or negative depending on the

intensity above threshold within cell boundaries. The percentage of *c-fos* positive cells within each

ROI was calculated. Figures were made by exporting ROIs to ImageJ v1-54f where they were

assembled using FigureJ package.

Statistical analysis

The physiological and cell counts data were analyzed in R version 4.3.1. The scripts for the analysis

and the data are available on github: https://github.com/ShonaWood/cFOSGHam

Time of day preference was assessed using Rayleigh test of circular uniformity using R. Significant

differences between numbers of *c-fos* positive cells was assessed by one-way ANOVA and post-hoc

testing by Dunnett's multiple comparisons test using R.

Results

Reproducible induction of hibernation and reliable prediction of torpor entry and arousal

We switched animals from a long photoperiod (16L:8D) at 21°C (LP) to a short photoperiod (8L:16D) at 21°C (SP-warm) for 4 weeks, then we lowered the temperature to 6.5°C (SP-cold) and monitored core body temperature (T_b) continuously (Figure 1A). Within 8 weeks of exposure to SP-cold conditions, 80% of our animals had commenced multi-day T-A cycling, this increased to 93% by 12 weeks (Supplementary Figure 1A). Similar to previous observations, we observed a gradual resetting of mean T_b from 36.4°C (Standard deviation (SD):0.83°C) to 33.4°C (SD:0.98°C) after 4 weeks in SP-cold, indicating a seasonal preparatory phase prior to the expression of hibernation (Chayama et al., 2016)(Figure 1A & Supplementary Figure 1B). We also observed so-called ´test-drops´ (Figure 1A, B)(Sheriff et al., 2012) in 86% of our animals prior to the initiation of T-A cycling. During test drops the mean T_b is 21.4°C (SD: 7.35) and the mean test-drop duration was only 5.3 hours. Both entry to, and, arousal from test drops showed circadian organisation (Supplementary Figure 1C). Not all animals showed test-drops prior to multiday T-A cycling, indicating that they are not a pre-requisite for hibernation.

Once initiated, T-A cycles were reproducible within and between animals, with T_b falling to within an average of 0.75°C of T_a. Torpor bout length increased from 56.5 hours (SD: 9.9) at the first bout to a maximum of 75.5 hours (SD: 8.9) by the sixth bout, while the interbout euthermic (IBE) interval decreased (from 61.2 hrs (SD:38.3) to 22.9 hrs (SD:7.86), Supplementary Figure 1D).

The T_b dynamics of the entry phase to torpor follows an inverse sigmoidal trajectory (Figure 1C). This is somewhat variable in between 33 – 26°C, with some animals showing a "step-wise" entry pattern; thereafter Tb declines approximately linearly at a max cooling rate of -3.24°C per hour, until about 15°C when the approach to nadir values slows (Figure 1C). Initiation of torpor is more likely to

occur in the dark phase (approx.17 to 22 hours after lights on (p<0.01 by Rayleigh test of circular

uniformity, Figure 1D).

Consistent with previous studies, arousal from torpor showed no circadian organisation (Figure 1D)

(Hut et al., 2002; Oklejewicz et al., 2001; Williams et al., 2012; Williams et al., 2017). Since T_b is

not a good early indicator of arousal (Markussen et al., 2020), we used IPTT-300 tags to measure

interscapular BAT temperature (T_{iBAT}) every 3-5 seconds, and monitored ventilation frequency (VF)

by video-streaming (Figure 1E). During spontaneous arousal, increases in T_{iBAT} precede T_b by up to

40 minutes (Figure 1F). We noted some variability in the arousal process when T_{iBAT} was between

4°C and 10°C from the torpor baseline, after which ventilation rates rapidly increased towards peak

values together with the max rewarming speed of T_{iBAT} and T_b (+18.3 °C / h). Despite variability in

its initiation, arousal was completed within 2.5-3 h in all animals (Figure 1G). To determine the

earliest possible time at which we could reliably predict an arousal event and clearly differentiate it

from animals thermoregulating during torpor, we monitored 12 separate arousals. We found that an

increase in T_{iBAT} of 0.5°C above torpor baseline coupled with a VF > 10 was the earliest reliable

predictor of a forthcoming arousal event, approximately 40 minutes before a detectable change in T_b

(Figure 1F, G).

The DMH expresses *c-fos* during entry to torpor

We used RNA expression of the immediate early gene *c-fos* by in-situ hybridization and validated

our probes using a light pulse paradigm and staining of the suprachiasmatic nucleus (Supplementary

Figure 2A)(Earnest et al., 1990). Then we sampled animals that had undergone at least 3 full T-A

cycles at the following phases (for details of calculation see Methods): Entering torpor (ENT<25°C),

40 hours torpid (T-40h), the earliest spontaneous arousal (AΔ0.5°C), early spontaneous arousal prior

to phase of high inter-animal variation (AΔ3°C), and interbout euthermic animals after 1 hour (IBE-

1h) and 12 hours (IBE-12h) (Figure 2A).

Surveying the thermoregulatory circuits and comparing *c-fos* expression over these timepoints

revealed that the POA (Figure 2B) showed an approximately 0.5% increase in the proportion of *c-fos*

positive cells after 40 hours of torpor compared to IBE-12h (Figure 2C & D , $p < 0.01$, one-way

ANOVA, Dunnett´s multiple comparison test). Contrastingly, we saw significant changes in the

number of *c-fos* positive cells in the DMH during the entry phase of the T-A cycle (2.25% positive

cells, Figure 2E & F, $p < 0.005$, one-way ANOVA, Dunnett's multiple comparison test). In the DMH

during arousal *c-fos* positive cells were detected but these changes were variable and therefore not

significant. We also surveyed other neuronal centres in the hypothalamus (Figure 2B; ARC and

VMH), less than 1% of cells showed *c-fos* expression during torpor and almost no cells were positive

in the transition phases of the T-A cycle (Supplementary Figure 2B & C).

***c-fos* induction in choroid plexus, tanycytes and pars tuberalis and during early arousal**

We noted high expression of *c-fos* in the choroid plexus (CP) (Figure 3A, B & C), with an average of

65% of CP cells showing positive staining in torpid animals, rising to 93% in the earliest arousing

animals (AΔ0.5°C), compared to 0.3% of positive cells in IBE-12h (Figure 3C, IBE-12h versus

AΔ0.5°C p=0.0005, IBE-12h versus T-40h p=0.0063, one-way ANOVA, post-hoc testing with

Dunnett's multiple comparisons). The circumventricular organs, the vascular organ of lamina

terminalis (OVLT) and median eminence (ME) showed approximately 5% of cells as positive during

torpor and arousal but there was high inter-individual variation within the groups (Supplementary

Figure 3A-D). Whilst surveying the ME, we noted that the pars tuberalis (PT) of the adjacent

pituitary stalk (Figure 3A) showed very high *c-fos* expression (22.6% of cells) at AΔ0.5°C (Figure

3D & E, IBE12h versus AΔ0.5°C p <0.0001, one-way ANOVA, post-hoc testing with Dunnett's

multiple comparisons). Similarly, alpha tanycytic cells lining the 3[rd] ventricle of the mediobasal

hypothalamus (αTC, Figure 3A), showed increased *c-fos* expression in AΔ0.5°C (7.3%, Figure 3F &

G, IBE-12h versus AΔ0.5°C p=0.0064, one-way ANOVA, post-hoc testing with Dunnett's multiple

comparisons). Beta tanycytes lining the 3[rd] ventricle but in closer proximity to the ME also showed

an increase in *c-fos* positive cells but to a lesser degree (Supplementary Figure 3E & F). Tanycytes

have long projections into the hypothalamus and down to the pars tuberalis (Guerra et al., 2010;

Güldner and Wolff, 1973; Rodríguez et al., 2005), we used vimentin to stain these projections,

showing infiltration to the DMH, VMH, ARC and PT (Figure 3H & I).

Discussion

We have monitored spontaneous T-A cycles in the golden hamster at high temporal and temperature

resolution to capture the initiation of spontaneous arousal from torpor. In two hypothalamic neuronal

centres linked to thermoregulation, the POA and the DMH (Cannon and Nedergaard, 2004;

Morrison, 2016), no significant change in *c-fos* expression was seen during spontaneous arousal, and

the only significant change observed was increased expression during torpor entry in the DMH,

potentially reflecting the tightly controlled cooling process (Florant and Heller, 1977). This contrasts

with the pattern seen in 3 non-neuronal tissues, the CP, PT and tanycytes, in which we detected the

highest level of *c-fos* RNA expression when BAT temperature had risen only 0.5°C above torpor

baseline. This indicates that these regions are sensitive to physiological changes at an early stage in

the arousal process. This extends similar findings in the thirteen lined ground squirrel (Bratincsák et

al., 2007) and is consistent with involvement of these sites in metabolic feedback regulation of T-A

cycling.

The expression of *c-fos* in non-neuronal cells in the choroid plexus, the mediobasal hypothalamus

and the PT may reflect ionic changes that are not so clearly linked to "activation" as is the case for

neuronal *c-fos* expression (Lara Aparicio et al., 2022), but are nonetheless interesting given the

highly time- and tissue-restricted nature of this response. The primary function of the choroid plexus

is to produce and control the composition of CSF, maintaining a blood-CSF barrier that gates the

exchange of metabolites and factors into the brain (Lun et al., 2015). In the mouse, the CP shows

pronounced daily changes in its production of major transport proteins, notably the thyroid hormone

carrier protein transthyretin (Duarte et al., 2020; Fame et al., 2023), while accumulation of

hibernation factor in the CP of Siberian chipmunks shows clear circannual variation (Kondo et al.,

2006). These observations suggest that CP function changes over time, correlated with daily and

seasonal changes in brain energy demand.

This framing of CP function has clear parallels with the function of tanycytes in the mediobasal

hypothalamus: These modified glial cells line the walls of the 3^{rd} ventricle, and serve a gating

function modulating the influence of nutrients, hormones and metabolites as feedback signals on

neuronal populations involved in appetite, energy balance and reproduction (Balland et al., 2014;

Bolborea et al., 2020; Frayling et al., 2011; Langlet, 2014; Langlet et al., 2013; Lhomme et al.,

2021). Tanycytes have been linked to circadian control of glucose homeostasis (Rodríguez-Cortés et

al., 2022) and have emerged as key cellular substrate for seasonal modulation of hypothalamic

function (Dardente et al., 2014; Dardente et al., 2019; Hazlerigg and Loudon, 2008). This latter

aspect depends on photoperiod-dependent production of thyrotropin by neighbouring PT cells which

in turn acts via tanycytic TSH-receptor expression (Hanon et al., 2008). This then modulates

tanycytic uptake of thyroid hormone by monocarboxylate transporter 8 (SLC16A2) (Friesema et al.,

2003; Petri et al., 2016) and tanycytic conversion of thyroid hormone by deiodinases (dio2 / dio3)

(Dardente et al., 2014; Dardente et al., 2019; Hazlerigg and Loudon, 2008). The relevance of this

pathway for expression of torpor is suggested by the blockade of daily torpor expression in Siberian

hamsters by exogenous thyroid hormone delivery to the DMH (Murphy et al., 2012), a site to which

tanycyte cell processes project (Figure 3H) (Guerra et al., 2010; Güldner and Wolff, 1973; Rodríguez

et al., 2005). Furthermore, in the Arctic ground squirrel, circannual termination of hibernation

correlates with increased deiodinase 2 gene expression in the mediobasal hypothalamus (Chmura et

al., 2022).

Hence it can be seen that involvement in temporal changes in metabolic energy demand and thyroid

metabolism constitutes a unifying functional framework for the 3 non-neuronal tissues in which we

see peak *c-fos* expression at arousal initiation. This leads us to hypothesise that *c-fos* induction occurs

in response to metabolic changes developing during torpor, reflecting the direct sensitivity of these

tissues to changing concentrations of blood- or CSF-borne metabolites. Further experiments are

required to test this hypothesis and to explore how changes in tanycyte or CP activity might couple to

the neural circuits driving thermogenesis at arousal.

Acknowledgements

The authors would like to thank our animal technicians Hans Lian, Hans-Arne Solvang and Renate

Thorvaldsen. The authors also thank Valerie Simonneaux and Paul Klosen for the gift of the *c-fos*

plasmid. We acknowledge the UiT core microscopy facility for the use of the confocal microscope.

Funding

The work was supported by grants from the Tromsø forskningsstiftelse (TFS) starter grant

TFS2016SW and the TFS infrastructure grant (IS3_17_SW) awarded to S.H.W. It was also co-

funded by the European Union (ERC, HiTime, 101086671). Views and opinions expressed are

however those of the author(s) only and do not necessarily reflect those of the European Union or the

European Research Council. Neither the European Union nor the granting authority can be held

responsible for them. The Arctic seasonal timekeeping initiative (ASTI) grant and UiT strategic

funds support D.G.H., S.H.W., F.A.F.M., and F.C.M.

Data availability

The analyzed data and code for figures for this study can be found in GitHub repository

https://github.com/ShonaWood/cFOSGHam

References

Balland, E., Dam, J., Langlet, F., Caron, E., Steculorum, S., Messina, A., Rasika, S., Falluel-

Morel, A., Anouar, Y., Dehouck, B., et al. (2014). Hypothalamic Tanycytes Are an ERK-

Gated Conduit for Leptin into the Brain. *Cell Metab.* **19**, 293–301.

Bolborea, M., Pollatzek, E., Benford, H., Sotelo-Hitschfeld, T. and Dale, N. (2020).

Hypothalamic tanycytes generate acute hyperphagia through activation of the arcuate neuronal

network. *Proc. Natl. Acad. Sci.* **117**, 14473–14481.

Bratincsák, A., McMullen, D., Miyake, S., Tóth, Z. E., Hallenbeck, J. M. and Palkovits, M.

(2007). Spatial and temporal activation of brain regions in hibernation:c-fos expression during

the hibernation bout in thirteen-lined ground squirrel. *J. Comp. Neurol.* **505**, 443–458.

Cannon, B. and Nedergaard, J. (2004). Brown Adipose Tissue: Function and Physiological

Significance. *Physiol. Rev.* **84**, 277–359.

Chayama, Y., Ando, L., Tamura, Y., Miura, M. and Yamaguchi, Y. (2016). Decreases in body

temperature and body mass constitute pre-hibernation remodelling in the Syrian golden hamster,

a facultative mammalian hibernator. *R. Soc. Open Sci.* **3**, 160002.

Chmura, H. E., Duncan, C., Saer, B., Moore, J. T., Barnes, B. M., Loren Buck, C., Christian, H.

C., Loudon, A. S. I. and Williams, C. T. (2022). Hypothalamic remodeling of thyroid hormone

signaling during hibernation in the arctic ground squirrel. *Commun. Biol.* **5**, 492.

Dardente, H., Hazlerigg, D. G. and Ebling, F. J. P. (2014). Thyroid hormone and seasonal

rhythmicity. *Front. Endocrinol. (Lausanne).* **5**, 19.

Dardente, H., Wood, S., Ebling, F. and Sáenz de Miera, C. (2019). An integrative view of

mammalian seasonal neuroendocrinology. *J. Neuroendocrinol.* **31**, e12729.

Duarte, A. C., Furtado, A., Hrynchak, M. V., Costa, A. R., Talhada, D., Gonçalves, I., Lemos,

M. C., Quintela, T. and Santos, C. R. A. (2020). Age, Sex Hormones, and Circadian Rhythm

Regulate the Expression of Amyloid-Beta Scavengers at the Choroid Plexus. *Int. J. Mol. Sci.* **21**,

6813.

Earnest, D. J., Iadarola, M., Yeh, H. H. and Olschowka, J. A. (1990). Photic regulation of c-fos

expression in neural components governing the entrainment of circadian rhythms. *Exp. Neurol.*

109, 353–361.

Fame, R. M., Kalugin, P. N., Petrova, B., Xu, H., Soden, P. A., Shipley, F. B., Dani, N., Grant,

B., Pragana, A., Head, J. P., et al. (2023). Defining diurnal fluctuations in mouse choroid

plexus and CSF at high molecular, spatial, and temporal resolution. *Nat. Commun.* **14**, 3720.

Florant, G. L. and Heller, H. C. (1977). CNS regulation of body temperature in euthermic and

hibernating marmots (Marmota flaviventris). *Am. J. Physiol. Integr. Comp. Physiol.* **232**, R203–

R208.

Frayling, C., Britton, R. and Dale, N. (2011). ATP-mediated glucosensing by hypothalamic

tanycytes. *J. Physiol.* **589**, 2275–2286.

Friesema, E. C. H., Ganguly, S., Abdalla, A., Manning Fox, J. E., Halestrap, A. P. and Visser,

T. J. (2003). Identification of monocarboxylate transporter 8 as a specific thyroid hormone

transporter. *J. Biol. Chem.* **278**, 40128–40135.

Guerra, M., Blázquez, J. L., Peruzzo, B., Peláez, B., Rodríguez, S., Toranzo, D., Pastor, F. and

Rodríguez, E. M. (2010). Cell organization of the rat pars tuberalis. Evidence for open

communication between pars tuberalis cells, cerebrospinal fluid and tanycytes. *Cell Tissue Res.*

339, 359–81.

Güldner, F. H. and Wolff, J. R. (1973). Neurono-glial synaptoid contacts in the median eminence

of the rat: ultrastructure, staining properties and distribution on tanycytes. *Brain Res.*

Hanon, E. A., Lincoln, G. A., Fustin, J.-M., Dardente, H., Masson-Pévet, M., Morgan, P. J. and

Hazlerigg, D. G. (2008). Ancestral TSH mechanism signals summer in a photoperiodic

mammal. *Curr. Biol.* **18**, 1147–52.

Hazlerigg, D. and Loudon, A. (2008). New insights into ancient seasonal life timers. *Curr. Biol.* **18**,

R795–R804.

Hrvatin, S., Sun, S., Wilcox, O. F., Yao, H., Lavin-Peter, A. J., Cicconet, M., Assad, E. G.,

Palmer, M. E., Aronson, S., Banks, A. S., et al. (2020). Neurons that regulate mouse torpor.

Nature **583**, 115–121.

Hut, R., Barnes, B. and Daan, S. (2002). Body temperature patterns before, during, and after semi-

natural hibernation in the European ground squirrel. *J. Comp. Physiol. B Biochem. Syst.*

Environ. Physiol. **172**, 47–58.

Ikeno, T., Williams, C. T., Buck, C. L., Barnes, B. M. and Yan, L. (2017). Clock Gene Expression

in the Suprachiasmatic Nucleus of Hibernating Arctic Ground Squirrels. *J. Biol. Rhythms* **32**,

246–256.

Kenagy, G. J., Sharbaugh, S. M. and Nagy, K. A. (1989). Annual cycle of energy and time

expenditure in a golden-mantled ground squirrel population. *Oecologia* **78**, 269–282.

Kondo, N., Sekijima, T., Kondo, J., Takamatsu, N., Tohya, K. and Ohtsu, T. (2006). Circannual

Control of Hibernation by HP Complex in the Brain. *Cell* **125**, 161–172.

Körtner, G. and Geiser, F. (2000). The temporal organization of daily torpor and hibernation:

circadian and circannual rhythms. *Chronobiol. Int.* **17**, 103–128.

Langlet, F. (2014). Tanycytes: A Gateway to the Metabolic Hypothalamus. *J. Neuroendocrinol.* **26**,

753–760.

Langlet, F., Mullier, A., Bouret, S. G., Prevot, V. and Dehouck, B. (2013). Tanycyte-like cells

form a blood-cerebrospinal fluid barrier in the circumventricular organs of the mouse brain. *J.*

Comp. Neurol. **521**, 3389–3405.

Lara Aparicio, S. Y., Laureani Fierro, Á. de J., Aranda Abreu, G. E., Toledo Cárdenas, R.,

García Hernández, L. I., Coria Ávila, G. A., Rojas Durán, F., Aguilar, M. E. H., Manzo

Denes, J., Chi-Castañeda, L. D., et al. (2022). Current Opinion on the Use of c-Fos in

Neuroscience. *NeuroSci* **3**, 687–702.

Lhomme, T., Clasadonte, J., Imbernon, M., Fernandois, D., Sauve, F., Caron, E., da Silva

Lima, N., Heras, V., Martinez-Corral, I., Mueller-Fielitz, H., et al. (2021). Tanycytic

networks mediate energy balance by feeding lactate to glucose-insensitive POMC neurons. *J.*

Clin. Invest. **131**,.

Lun, M. P., Monuki, E. S. and Lehtinen, M. K. (2015). Development and functions of the choroid

plexus–cerebrospinal fluid system. *Nat. Rev. Neurosci.* **16**, 445–457.

Lyman, C. P. (1948). The oxygen consumption and temperature regulation of hibernating hamsters.

J. Exp. Zool. **109**, 55–78.

Lyman, C. P., Willis, J. ., Malan, A. and Wang, L. C. . (1982). *Hibernation and torpor in*

mammals and birds. (ed. Lyman, C. P.) New York: Academic Press.

Malan, A. (2010). Is the Torpor-Arousal Cycle of Hibernation Controlled by a Non-Temperature-

Compensated Circadian Clock? *J. Biol. Rhythms* **25**, 166–175.

Malan, A., Ciocca, D., Challet, E. and Pévet, P. (2018). Implicating a Temperature-Dependent

Clock in the Regulation of Torpor Bout Duration in Classic Hibernation. *J. Biol. Rhythms* **33**,

626–636.

Markussen, F. A. S., Melum, V. J., Bothorel, B., Hazlerigg, D. G., Simonneaux, V. and Wood, S.

H. (2020). A refined method to monitor arousal from hibernation in the European hamster.

Mckee, G. and Andrews, J. (1992). The costs and savings of hibernation in the golden hamster,

Proceedings of the Physiological Society, 20☐21 September 1991, Dublin Meeting:

Communications. In *The Journal of Physiology*, p. 451.

Morrison, S. (2016). Central neural control of thermoregulation and brown adipose tissue. *Auton.*

Neurosci. **196**, 14–24.

Murphy, M., Jethwa, P. H., Warner, A., Barrett, P., Nilaweera, K. N., Brameld, J. M. and

Ebling, F. J. P. (2012). Effects of Manipulating Hypothalamic Triiodothyronine Concentrations

on Seasonal Body Weight and Torpor Cycles in Siberian Hamsters. *Endocrinology* **153**, 101–

112.

Oklejewicz, M., Daan, S. and Strijkstra, A. (2001). Temporal organisation of hibernation in wild-

type and tau mutant Syrian hamsters. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **171**,

431–439.

Petri, I., Diedrich, V., Wilson, D., Fernandez-Calleja, J., Herwig, A., Steinlechner, S., Barrett,

P., Fernández-Calleja, J., Herwig, A., Steinlechner, S., et al. (2016). Orchestration of gene

expression across the seasons: Hypothalamic gene expression in natural photoperiod throughout

the year in the Siberian hamster. *Sci. Rep.* **6**, 29689.

Regan, M. D., Chiang, E., Martin, S. L., Porter, W. P., Assadi-Porter, F. M. and Carey, H. V.

(2019). Shifts in metabolic fuel use coincide with maximal rates of ventilation and body surface

rewarming in an arousing hibernator. *Am. J. Physiol. Integr. Comp. Physiol.* **316**, R764–R775.

Revel, F. G., Herwig, A., Garidou, M.-L., Dardente, H., Menet, J. S., Masson-Pévet, M.,

Simonneaux, V., Saboureau, M. and Pévet, P. (2007). The circadian clock stops ticking

during deep hibernation in the European hamster. *Proc. Natl. Acad. Sci.* **104**, 13816–13820.

Rodríguez-Cortés, B., Hurtado-Alvarado, G., Martínez-Gómez, R., León-Mercado, L. A.,

Prager-Khoutorsky, M. and Buijs, R. M. (2022). Suprachiasmatic nucleus-mediated glucose

entry into the arcuate nucleus determines the daily rhythm in blood glycemia. *Curr. Biol.*

Rodríguez, E. M., Blázquez, J. L., Pastor, F. E., Peláez, B., Peña, P., Peruzzo, B. and Amat, P.

(2005). Hypothalamic Tanycytes: A Key Component of Brain–Endocrine Interaction. *Int. Rev.*

Cytol. **247**, 89–164.

Ruby, N. F. and Zucker, I. (1992). Daily torpor in the absence of the suprachiasmatic nucleus in

Siberian hamsters. *Am. J. Physiol. Integr. Comp. Physiol.* **263**, R353–R362.

Ruf, T., Steinlechner, S., and Heldmaier, G. (1989). Rhythmicity of body temperature and torpor

in the Djungarian hamster, Phodopus sungorus. In *Proceedings of the Living in the Cold: 2nd*

International Symposium (ed. Malan, A.) and Canguilhem, B.), pp. 53–61. Le Hohwald:

Colloques INSERM/John Libbey Eurotext Ldt.

Ruf, T. and Geiser, F. (2015). Daily torpor and hibernation in birds and mammals. *Biol. Rev.* **90**,

891–926.

Sheriff, M. J., Williams, C. T., Kenagy, G. J., Buck, C. L. and Barnes, B. M. (2012).

Thermoregulatory changes anticipate hibernation onset by 45 days: Data from free-living arctic

ground squirrels. *J. Comp. Physiol. B Biochem. Syst. Environ. Physiol.* **182**, 841–847.

Singh, S., Bermudez-Contreras, E., Nazari, M., Sutherland, R. J. and Mohajerani, M. H.

(2019). Low-cost solution for rodent home-cage behaviour monitoring. *PLoS One* **14**,

e0220751.

Tähti, H. and Antti, S. (1977). Respiratory and circulatory differences between induced and

spontaneous arousals in hibernating hedgehogs (Erinaceus europaeus L.). *Ann. Zool. Fennici* **14**,

198–203.

Takahashi, T. M., Sunagawa, G. A., Soya, S., Abe, M., Sakurai, K., Ishikawa, K., Yanagisawa,

M., Hama, H., Hasegawa, E., Miyawaki, A., et al. (2020). A discrete neuronal circuit induces

a hibernation-like state in rodents. *Nature* **583**, 109–114.

Upton, B. A., D'Souza, S. P. and Lang, R. (2021). QPLOT Neurons—Converging on a

Thermoregulatory Preoptic Neuronal Population. *Front. Neurosci.* **15**, null.

Utz, J. C. and van Breukelen, F. (2013). Prematurely induced arousal from hibernation alters key

aspects of warming in golden-mantled ground squirrels, Callospermophilus lateralis. *J. Therm.*

Biol. **38**, 570–575.

van Breukelen, F. and Martin, S. L. (2015). The Hibernation Continuum: Physiological and

Molecular Aspects of Metabolic Plasticity in Mammals. *Physiology* **30**, 273–281.

Williams, C. T., Barnes, B. M., Richter, M. and Buck, C. L. (2012). Hibernation and Circadian

Rhythms of Body Temperature in Free-Living Arctic Ground Squirrels. *Physiol. Biochem. Zool.*

85, 397–404.

Williams, C. T., Radonich, M., Barnes, B. M. and Buck, C. L. (2017). Seasonal loss and

resumption of circadian rhythms in hibernating arctic ground squirrels. *J. Comp. Physiol. B* **187**,

693–703.

Figure legends

Figure 1: Physiological monitoring of the golden hamster during hibernation

A. Representative core body temperature trace (T_b °C, black line) and ambient temperature (T_a

°C, blue line) over the whole experiment. Hamsters were kept on long photoperiod (LP

(16L:8D), T_a = 21°C) for 6 weeks then, moved to short photoperiod and warm conditions

(SP= 8L:16D, T_a = 21°C) for 4 weeks. Finally, the T_a was lowered to 6.5°C for the remainder

of the experiment. The vertical dotted lines indicate when the changes in light or temperature

conditions occurred. Note the drop in Tb in the first week of SP, this was due to the anipill

shifting into the testis sack, which resolved itself.

B. Body temperature (T_b°C) traces of 18 hamsters synchronized to first torpor bout entry,

illustrating test-drops prior to T-A cycling and first T-A cycle.

C. Body temperature (T_b °C) of entry to torpor from 18 individuals to explore variation in entry dynamics. Red line is a regression analysis for the steepest part of the slope, showing an average maximal cooling rate of -3.24°C per hour.

D. Raleigh plots showing the time of entry and arousal from 28 individuals. Photoperiod is represented by grey for dark and yellow for light. The black dots are an individual arousal or entry event. The black arrow indicates the mean time of entry or arousal, and the light grey arrows indicates the standard deviation. The length of the arrow represents mean resultant length as an indicator of concentration around the mean, therefore reflecting the statistical significance. The Raleigh statistical test was used to test for a time-of-day preference in arousal and entry. Only the entry was significant $p<0.01$.

E. A schematic of the remote monitoring system for arousal from torpor. IPTT-300 was used to monitor iBAT temperature dynamics. A live video feed using a raspberry pi and camera module measured ventilation frequency (VF). iBAT temperature is continuously read every 3-5 seconds and all data is parsed to a networked database immediately accessible to local networked devices. The computer screen in the schematic shows the video feeds for VF and the green line is the iBAT temperature from an animal that has completed an arousal. In addition, Anipills were used to monitor core body temperature.

F. The change in iBAT temperature (red line) and core body temperature (green line) from torpor baseline from 10 representative arousals (Left axis Δ°C). Right axis shows actual iBAT temperature. Mean values are shown with standard deviations in grey shading.

G. A closer analysis of iBAT temperature dynamics during arousal of four representative arousals. The colour of the dots reflects ventilation rate. The change in iBAT temperature from torpor baseline is plotted on the left axis (Δ°C) and the actual temperature of the iBAT

 is plotted on the right axis (°C). The dashed red line indicates 0.5°C and 3°C change from

torpor baseline. Inset graph top left: Zoom in on the first hour of arousal.

Figure 2: *c-fos* expression in the POA and DMH during T-A cycling

A. A schematic showing interscapular BAT temperature (T_{iBAT}, red) and core body temperature

(T_b, grey) and the sampling points taken during the T-A-cycle. Interbout euthermic animals

sampled either 1 or 12 hours after arousal (IBE-1h, IBE-12h). Entry to torpor animals

sampled when T_b and T_{iBAT} are less than 25°C (ENT<25°C). Torpid animals sampled 40

hours into the torpor bout (T-40h). Early arousal animals defined by deviation in T_{iBAT} from

the torpor baseline of either 0.5°C or 3°C (AΔ0.5°C and AΔ3°C).

B. Brain regions of interest defined by the golden hamster brain atlas. Left panel: Bregma +0.9

in the pre-optic hypothalamic area (POA): MnPO; median preoptic area, AVPE; anteroventral

periventricular area, MPO; medial preoptic area, LPO; lateral preoptic area, OVLT; organum

vasculosum lamina terminalis, OC; optic chiasm, AC; anterior commissure. Right: Bregma -

2.6 in the medio basal hypothalamus: DMH; dorsomedial hypothalamus, VMH; ventromedial

hypothalamus, Arc; Arcuate nucleus, 3V; third ventricle, ME; median eminence, PT; Pars

tuberalis.

C. Representative images of in-situ hybridization using *c-fos* (white) and sytox orange to show

nuclei (red) in the preoptic area (POA), indicated for white outline, for each group defined in

A. Scalebar 500 µm.

D. Quantification of the percentage of positive cells in the preoptic area (POA). Results of one-

way ANOVA and post hoc testing by Dunnett's multiple comparisons test are shown; * p-

value ≤0.05.

E. Representative images of in-situ hybridization using *c-fos* (white) and sytox orange to show

nuclei (red) in the dorsomedial hypothalamus (DMH), indicated by white outline, for each

group defined in A. Scalebar 500 µm.

F. Quantification of the percentage of positive cells in the dorsomedial hypothalamus (DMH).

Results of one-way ANOVA and post hoc testing by Dunnett's multiple comparisons test are

shown; ** p-value ≤0.005.

Figure 3*: c-fos* expression in the choroid plexus, pars tuberalis and tanycytes during early

arousal from torpor

A. A schematic indicating Bregma -2.6 and the location of α-tanycytes (αTC), Pars tuberalis

(PT) and choroid plexus (CP). ME; median eminence. Grey dotted lines indicate the regions

of the hypothalamic nuclei. Note the CP is in the lateral ventricles and is therefore displayed

in the inset relative to the hypothalamic region shown.

B. Representative images of in-situ hybridization using *c-fos* (white) and sytox orange to show

nuclei (red) in the choroid plexus (CP), for IBE-12h and AΔ3°C. Scalebar 500 µm.

C. Quantification of the percentage of positive cells in the choroid plexus (CP). Results of one-

way ANOVA and post hoc testing by Dunnett's multiple comparisons test are shown; * p-

value ≤0.05, ** p-value ≤0.005, *** p-value ≤0.0005

D. Representative images of in-situ hybridization using *c-fos* (white) and sytox orange to show

nuclei (red) in the pars tuberalis (PT), for each group defined in Figure 2A. The

representative image for AΔ0.5°C contains a white dotted line box indicating the zoomed in

image shown in I. Scalebar 500 µm.

E. Quantification of the percentage of positive cells in the pars tuberalis (PT). Results of one-

way ANOVA and post hoc testing by Dunnett's multiple comparisons test are shown; ****

p-value <0.0001

F. Representative images of in-situ hybridization using *c-fos* (white) and sytox orange to show

nuclei (red) in the α-tanycytes (αTC) region, for each group defined in Figure 2A. Scalebar

200 µm.

G. Quantification of the percentage of positive cells in the α-tanycytes (αTC) region. Results of

one-way ANOVA and post hoc testing by Dunnett's multiple comparisons test are shown; **

p-value ≤0.005.

H. Vimentin staining (green) for tanycyte process in the mediobasal hypothalamus. *c-fos* (white)

and sytox orange to show nuclei (red). The hypothalamic nuclei are indicated by white dotted

lines. DMH; dorsomedial hypothalamus, VMH; ventromedial hypothalamus, Arc; Arcuate

nucleus. Scalebar 500 µm.

I. A zoomed in image of the pars tuberalis (PT) shown in D during early arousal (AΔ0.5°C).

Vimentin staining (green) for tanycyte processes, *c-fos* shown in white, and sytox orange to

show nuclei is red. 3V; third ventricle. Scalebar 50 µm.

Supplementary Figure 1: Physiological monitoring of hibernation in the golden hamster

A. Percentage of animals hibernating in response to SP-cold conditions over time.

B. Core body temperature (T_b) in response to the transition from long photoperiod (LP) at

ambient temperature (T_a) 21°C, to short photoperiod (SP) at Ta 21°C (first dotted horizontal

line), to SP and T_a 6.5°C. Prior to T-A cycling diel patterning of mean core body temperature

was observed at all stages of the experiment, except the first 4 weeks of cold where the T_b is

reducing (see inset periodograms). Periodograms were generated using ActogramJ and the

chi-square method.

C. Raleigh plots showing the time of test-drop entry and arousal from 24 individuals.

Photoperiod is represented by grey for dark and yellow for light. The black dots are an

individual arousal or entry event. The black arrow indicates the mean time of entry or arousal,

and the light grey arrows indicates the standard deviation. The length of the arrow represents

mean resultant length as an indicator of concentration around the mean, therefore reflecting

the statistical significance. The Raleigh statistical was used to test for a time-of-day

preference in arousal and entry. Both entry and arousal were significant $p < 0.05$.

D. Top: Duration (hours) of inter bout euthermia after previous corresponding torpor bout for 22

individuals; listed on the side with individual colours shown in each box plot. Bottom: Torpor

bout duration (hours) for each torpor bout made.

Supplementary Figure 2: Validation of *c-fos* probe and numbers of *c-fos* positive cells in the

ARC and VMH

A. In-situ hybridization of the suprachiasmatic nucleus (SCN) in the golden hamster for *c-fos*

RNA (brown staining). Left: no light pulse during the dark phase. Right: 20-minute light

pulse during the dark phase. 3V; 3rd ventricle, indicated by black dotted line.

B. Quantification of the percentage of positive cells in the arcuate nucleus (ARC) region. Results

of one-way ANOVA and post hoc testing by Dunnett's multiple comparisons test are shown;

* p-value ≤ 0.05.

C. Quantification of the percentage of positive cells in the ventromedial hypothalamus (VMH)

region. Results of one-way ANOVA and post hoc testing by Dunnett's multiple comparisons

test are shown; ** p-value ≤ 0.005.

Supplementary Figure 3: *c-fos* expression in the OVLT, ME and beta tanycytes

A. Quantification of the percentage of positive cells in the Vascular organ of lamina terminalis

(OVLT) region. Results of one-way ANOVA and post hoc testing by Dunnett's multiple

comparisons test are shown; ** p-value ≤ 0.005.

B. Representative images of in-situ hybridization using *c-fos* (white) and sytox orange to show nuclei (red) in the OVLT region, for IBE-12h and AΔ3°C. Vimentin staining is shown in green. Scalebar 200 µm.

C. Quantification of the percentage of positive cells in the median eminence (ME). Results of one-way ANOVA and post hoc testing by Dunnett's multiple comparisons test are shown; ns = not significant.

D. Representative images of in-situ hybridization using *c-fos* (white) and sytox orange to show nuclei (red) in the ME region, for IBE-12h and AΔ3°C. Vimentin staining is shown in green. Scalebar 100 µm.

E. Quantification of the percentage of positive cells in the β-tanycytes (βTC) region. Results of one-way ANOVA and post hoc testing by Dunnett's multiple comparisons test are shown; * p-value ≤0.05.

F. Representative images of in-situ hybridization using *c-fos* (white) and sytox orange to show nuclei (red) in the β-tanycytes (βTC) region, for IBE-12h and AΔ3°C. Vimentin staining is shown in green. Scalebar 500 µm.

Author Contributions

Conceptualization: S.H.W., D.G.H., F.A.F.M., Methodology: F.A.F.M., F.C.M., V.J.M, Software: F.A.F.M., Validation: F.A.F.M., F.C.M., Formal analysis: F.A.F.M., S.H.W., D.G.H., Investigation: F.A.F.M, F.C.M., V.J.M., Resources: S.H.W., Data curation: F.A.F.M., Writing - original draft: F.A.F.M., S.H.W., D.G.H., Writing - Review & editing: All authors, Visualization: F.A.F.M., Supervision: S.H.W., D.G.H., Project administration: S.H.W., Funding acquisition: S.H.W.

Supplementary Figure 1

A

B

ARC ISH *c-fos* positive

C

VMH ISH *c-fos* positive

Supplementary Figure 3