SntB triggers the antioxidant pathways to regulate development and aflatoxin biosynthesis in *Aspergillus flavus*

Dandan Wu [1,#], Chi Yang [1,2,#], Yanfang Yao [1], Dongmei Ma [3], Hong Lin [1], Ling Hao [1], Wenwen Xin [4], Kangfu Ye [1], Minghui Sun [3], Yule Hu [1], Yanling Yang[1,*], Zhenhong Zhuang [1,*]

1. Key Laboratory of Pathogenic Fungi and Mycotoxins of Fujian Province, Key Laboratory of Biopesticide and Chemical Biology of Education Ministry, Proteomic Research Center, and School of Life Sciences, Fujian Agriculture and Forestry University, Fuzhou, 350002, China;

2. Institute of Edible Mushroom, Fujian Academy of Agricultural Sciences, Fuzhou, 350012, China;

3. College of Animal Sciences (College of Bee Science), Fujian Agriculture and Forestry University, Fuzhou 350002, China.

4. State Key Laboratory of Pathogen and Biosecurity, Institute of Microbiology and Epidemiology, Academy of Military Medical Sciences (AMMS), Beijing 100071, China;

* Corresponding author. E-mail addresses: ZH_Zhuang@fafu.edu.cn (Zhenhong Zhuang); 1034319020@qq.com (Yanling Yang).

[#] These authors contributed equally to this work

Abstract

The epigenetic reader SntB was identified as an important transcriptional regulator of growth, development, and secondary metabolite synthesis in *Aspergillus flavus*. However, the underlying molecular mechanism is still unclear. In this study, *sntB* gene deletion (Δ*sntB*), complementary (Com-*sntB*), and HA tag fused to *snt2* (*snt2*-HA) strains were constructed by using the homologous recombination method, respectively. Our results revealed that deletion of *sntB* inhibited the processes of mycelia growth, conidial production, sclerotia formation, aflatoxin synthesis, and ability to colonize host compared to wild type (WT), and the defective phenotype of knockout strain Δ*sntB* can be restored by its complementary strain Com-*sntB*. Chromatin immunoprecipitation sequencing (ChIP-seq) of *sntB*-HA and WT and RNA sequencing (RNA-seq) of Δ*sntB* and WT strains revealed that SntB played key roles in oxidative stress response of *A. flavus*. The function of *catC* (encode a catalase) gene was further analyzed based on the integration results of ChIP-seq and RNA-seq. In Δ*sntB* strain, the relative expression level of *catC* was significantly higher than in WT strain, while a secretory lipase encoding gene (G4B84_008359) was down-regulated. Under the stress of oxidant menadione sodium bisulfite (MSB), the deletion of *sntB* obvious down-regulated the expression level of *catC*. After deletion of *catC* gene, the mycelia growth, conidial production, and sclerotia formation were inhibited, while aflatoxin synthesis was increased compared to the WT strain. Results also showed that the inhibition rate of MSB to Δ*catC* strain was significantly lower than that of WT group and AFB1 yield of the Δ*catC* strain was significantly decreased

than that of WT strain under the stress of MSB. Our study revealed the potential

machinery that SntB regulated fungal morphogenesis, mycotoxin anabolism, and

fungal virulence through the axle of from SntB to fungal virulence and mycotoxin

bio-synthesis, i.e. H3K36me3 modification-SntB-Peroxisomes-Lipid

hydrolysis-fungal virulence and mycotoxin bio-synthesis. The results of the study

shad light into the SntB mediated epigenetic regulation pathway of fungal mycotoxin

anabolism and virulence, which provided potential strategy for control the

contamination of *A. flavus* and its aflatoxins.

Key words

Aspergillus flavus, SntB, ChIP-seq, RNA-seq, CatC, oxidative response

1. Introduction

Aspergillus flavus is one of the common asexual species, a saprophytic fungus and the second largest pathogenic fungus after *Aspergillus fumigatus*, widely distributed in soil, air, water, plants, and agricultural products in nature [1]. Aflatoxins produced by *A. flavus* has strong toxicity, and is extremely harmful to human society. Animal and human health can be negatively affected by aflatoxins, which are carcinogenic, teratogenic, and mutagenic [2]. Among aflatoxins, AFB1 is the most frequently occurring and the most toxic and carcinogenic, which is converted to AFB1-8 and 9-epoxide in the liver and formed adducts with the guanine base of DNA and thus results in acute and chronic diseases in both human and household animals [3]. According to the Food and Agriculture Organization of the United Nations, 25% of the food crops in the world are contaminated with aflatoxins [4], which are often detected in grains, nuts, and spices [5, 6]. It is urgent to control the contamination of *A. flavus* and its main mycotoxin, AFB1.

In recent decades, the biosynthetic pathway of aflatoxins was investigated in detail benefit from the sequence of *A. flavus* genome [7]. This pathway consists of a complex set of enzymatic reactions [8-11]. In general, these enzymes are encoded by clusters of genes, which are regulated by cluster-specific genes: *aflR* and *aflS* [12, 13]. The initial stage of aflatoxins biosynthesis is catalyzed by polyketide synthase (PKSA) to form the polyketone backbone [14]. Aflatoxins synthesis also regulated by environmental stimuli including pH, light, nutrient sources, and oxidative stress response, which may result in the modulation of the expression of genes involved in toxin production [15-17].

Besides the biosynthetic pathway and its internal gene regulation, protein

post-translational modifications (PTMs), an important mean of epigenetics, represent

an important role in the regulation of aflatoxins synthesis, including

2-hydroxyisobutyrylation, succinylation, acetylation, and methylation [18-24], in which

Snt2 (also called *sntB*, an epigenetic reader) is deeply involved. Despite

advancements in the field, our understanding of the molecular mechanisms of

aflatoxin production in *A. flavus* is still fragmentary.

The epigenetic reader encoded by *sntB* in *A. nidulans* was identified as a

transcriptional regulator of the sterigmatocystin biosynthetic gene cluster and

deletion of *sntB* gene in *A. flavus* results in loss of aflatoxin production [25], increasing

global levels of H3K9K14 acetylation and impairing several developmental processes

[20]. The homolog gene in yeast, *SNT2* coordinates the transcriptional response to

hydrogen peroxide stress [26, 27]. In *Penicillium expansum*, SntB regulated the

development, patulin and citrinin production, and virulence on apples [28]. In *A.*

nidulans, SntB combined with a H3K4 histone demethylase KdmB, a cohesin

acetyltransferase (EcoA), and a histone deacetylase (RpdA) to form a chromatin

binding complex and bound to regulatory genes and coordinated fungal development

with mycotoxin synthesis [29]. *sntB* also regulated the virulence in *Fusarium*

oxysporum, and respiration in *F. oxysporum* and *Neurospora crassa* [30, 31].

However, the specific regulatory mechanism of *sntB* in *A. flavus* remains unclear. In

this study, we identified the regulatory network of *sntB* by Chromatin

immunoprecipitation sequencing (ChIP-seq) and RNA sequencing (RNA-seq), which

shed light on its impact on fungal biology.

Experimental Procedures

A. flavus **strains, media, and culture conditions**

A. flavus Δ*ku70* Δ*pyrG* was used as a host strain for genetic manipulations. All strains

used in this study are listed in the **Table 1**. potato dextrose agar (PDA, 39 g/L, BD,

Difco, Franklin, NJ, USA), complete medium (CM, 6 g/L tryptone, 6 g/L yeast extract,

10 g/L glucose), and potato dextrose (PDB, 24 g/L, BD, Difco, Franklin, NJ, USA)

were used for mycelial growth and sporulation determination, sclerotia production,

and mycotoxin production analysis, respectively. All experiments were technically

repeated three times and biologically repeated three times.

Table 1. *A. flavus* strains used in this study

Strain name	Related genotype	Source
A. flavus CA14	Δ*ku70*, Δ*pyrG*	Kindly presented form Prof. Chang[1]
Wild-type (WT)	Δ*ku70*, Δ*pyrG*::*AfpyrG*	This study
Δ*sntB*	Δ*ku70*, Δ*sntB*::*AfpyrG*	This study
Com-*sntB*	Δ*ku70*, Δ*pyrG*; Δ*sntB*::*AfpyrG*:: *sntB*	This study
Δ*catC*	Δ*ku70*, Δ*catC*::*AfpyrG*	This study

The construction of mutant strains

All mutant strains, including *snt2* and *catC* gene knock-out strain (Δ*snt2* and Δ*catC*),

the complementation strain for the Δ*sntB* strain (Com-*sntB*), and HA tag fused to *snt2*

strain (*snt2*-HA), were constructed following the protocol of homologous

recombination [32] and the detail protocol was as described in our previous study [33].

The related primers were listed in Table 2. The constructed strains were confirmed by

t diagnostic PCRs and by southern bolt [34]. The construction of *snt2*-HA was further

determined by western blot with anti-HA antibody as described previously [33].

Table 2. Primers used for strain construction in this study

Primer name	Sequence (5' → 3')	Fragment amplified
sntB-p1	CTTCTCGAATTCCCCTTCATGACACTCTCC	
sntB-p2	GCTAAATCAGGATGGGTTGGAGGGTGAC	
sntB-p3	GCATCTCCTTTGTGTTGTTTGGACCGTGT	
sntB-p4	CAACCAACCACTGACGTCGACCAC	
sntB-p5	GTCACCCTCCAACCCATCCTGATTTAGCGCCTCAAACAATGCTCTTCACCC	For the construction of knockout strain
sntB-p6	ACACGGTCCAAACAACACAAAGGAGATGCGTCTGAGAGGAGGCACTGATGC	
sntB-p7	TAGATCACCCAGCGGGCCACAA	
sntB-p8	GACTCAAATGGAAATCCCGTCGTGCC	
sntB-F	GCAGCAACACCACGTGAGGCCCAATTC	a fragment from *sntB* ORF
sntB-R	CCAGGTCACAGGGCATAGAACACACTCGTA	
P1020-F	ATCGGCAATACCGTCCAGAAGC	Verify the fragment of *AfpyrG*
P801-R	CAGGAGTTCTCGGGTTGTCG	
sntB-*C*-p2	GGGTGAAGAGCATTGTTTGAGGCCCTCCAACCTTACTCCGTACACAATTCTAG	For the construction of complementary strain of *snt2*
s*ntB*-*C*-p3	GCATCAGTGCCTCCTCTCAGACGGCACGACGGGATTTCCATTTGAGTC	
sntB-*C*-p4	CCAATTTCCTGATGATTGTGATGTGTGTCC	
SC-P-F	GCCTCAAACAATGCTCTTCACCC	

SC-P-R	GTCTGAGAGGAGGCACTGATGC	
catC-P1	CTTGAGACGCAGGACGAA	For the
catC-P2	GGGTGAAGAGCATTGTTTGAGGCTGATGTG GGTTGTATGAATG	construction of knockout strain of
catC-P3	GCATCAGTGCCTCCTCTCAGACTGGATGCGG GTGAATACTG	*catC*
catC-P4	ACAAGCTGTCATGCGTGG	
catC-P5	TGGGAGTCTCGAACACAC	
catC-P6	GAAAACCCCGCAACAGAC	
catC-F	AGCCTATTTCGGACCCCT	
catC-R	CAGTCTCCTTTCGGCATC	

Phenotypic analysis and aflatoxin analysis

126 The spores (10^7 conidia/mL) of wild type (WT), $\Delta sntB$, and Com-*sntB* strains were

127 used. The details of experiment were according to our previous study [35]. Hyphal

128 septum was stained according to the described method [36]. Each fungal strain was

129 evaluated on four plates, and each experiment was repeated three times.

Fungal colonization on crop kernels

According to our previous experimental protocol [35], the colonizing ability of WT,

$\Delta sntB$, and Com-*sntB* fungal strains on peanut and corn kernels was analyzed. The

crop kernels were disinfected with 0.05% sodium hypochlorite and soaked for 30 min

in a solution containing 10^5 conidia/mL fungal spores. Afterwards, the seeds were

placed in a petri dish and cultured at 29°C for 6 d. Finally, the number of conidia was

calculated and AFB1 product was analyzed by TLC.

Animal invasion assay

The animal invasion assay using silkworms was according to previous study [32, 37].

Silkworms (*Bombyx mori*) were randomly separated into four groups (10 larvae/group)

when silkworm larva reaches about 1 g in weight. Each silkworm was injected with 5

μL saline, or 5 μL conidial suspension (10^6 spores/mL) from WT, Δ*sntB*, and

Com-*sntB* strains. The survival rate of silkworms was calculated. Dead silkworms

were transferred into fresh 9 cm Petri dishes and inoculated for 5 d in the dark. The

conidia number and AFB1 production from each group were measured.

RNA-seq analysis

To reveal the potential complex regulatory network of the *sntB*, RNA-seq analysis

was carried out on the WT and Δ*sntB* strains. RNA-seq analysis was carried out by

APPLIED PROTEIN TECHNOLOY, Shanghai (www. aptbiotech.com) [38]. Data

processing was according pervious study [39]. Differentially expressed gene (DEGs)

were assigned as genes with |log2FoldChange|>1 and adjusted *P*-adj< 0.05. Gene

Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways

were used to analyze the functions of DEGs.

ChIP-seq and data analysis

ChIP-seq analysis was carried out on the WT and *sntB*-HA strains. The conidia

(10^4/ml) for each strain were inoculated in 100 mL PDB shaking at 180 rpm under

29°C for 72h, and subjected to ChIP -seq analysis by Wuhan IGENEBOOK

Biotechnology Co.,Ltd (http://www.igenebook.com). ChIP experiment was according

previous study [32]. Raw sequencing with low-quality reads discarded and reads

contaminated with adaptor sequences trimmed were filtered by Trimmomatic (v0.36)

[40]. The clean reads were mapped to the reference genome of *A. flavus* by

Burrows-Wheeler Alignment tool (BWA, version 0.7.15) [41]. MACS2 (v2.1.1) and

Bedtools (v2.25.0) were used for peak calling and peak annotation, respectively.

Differential binding peaks were identified by   Fisher's test with q value <0.05. Genes

less than 2000 bp away were associated to the corresponding peak. GO and KEGG

enrichment analyses of annotated genes were implemented in EasyGO [42] and

KOBAS (v2.1.1) [43], with a corrected *P*-value cutoff of 0.05.

Quantitative RT-PCR (qRT-PCR) analysis

The fungal spores (10^6/mL) were cultured in PDB medium for 48 h, and then

mycelium was ground into powder with liquid nitrogen. Total RNA was prepared by

TRIpure total RNA Extraction Reagent (Bestek, China) according to the protocol used

by Zhang [44]. qRT-PCR was performed according to previously described [45], and the

primers were shown in **Table 3**.

Table 3. Primers used for RT-qPCR in this study

Primer name	Sequence (5' → 3')	Fragment amplified
sntB-qF	ACTCATCAGAGCCCCTATGGGCC AGTC	
sntB-qR	GCACTAAGAACGCGATCGACAG AATAGACAC	
catC-qF	GAAAGAGTTGTCCATGCCA	
catC-qR	CAGAAAACGGGTGTGTGAT	
brlA-F	GCCTCCAGCGTCAACCTTC	
brlA-R	TCTCTTCAAATGCTCTTGCCTC	
abaA-F	TCTTCGGTTGATGGATGATTTC	
abaA-R	CCGTTGGGAGGCTGGGT	
nsdC-F	GCCAGACTTGCCAATCAC	
nsdC-R	CATCCACCTTGCCCTTTA	Sclerotium related genes
nsdD-F	GGACTTGCGGGTCGTGCTA	
nsdD-R	AGAACGCTGGGTCTGGTGC	

sclR-F	CAATGAGCCTATGGGAGTGG	
sclR-R	ATCTTCGCCCGAGTGGTT	
aflC-F	GTGGTGGTTGCCAATGCG	
aflC-R	CTGAAACAGTAGGACGGGAGC	
aflP-F	CGATGTCTATCTTCTCCGATCTATTC	Toxin synthesis structure genes
aflP-R	TCTCAGTCTCCAGTCTATTATCTACC	
aflO-F	CTTTCGGCAGTGACCTAACC	
aflO-R	TCTTGAACTATAAGGCGACCA	
aflR-F	AAAGCACCCTGTCTTCCCTAAC	
aflR-R	GAAGAGGTGGGTCAGTGTTTGTAG	Toxin synthesis regulatory genes
aflS-F	GCTCAGACTGACCGCCGCTC	
aflS-R	GCTCAGACTGACCGCCGCTC	
18S rRNA-F	CTGAAGACTAACTACTGCGAAAGC	
18S rRNA-R	GAGCGGGTCATCATAGAAACAC	RNA extraction quality testing
β-tublin-F	TTGAGCCCTACAACGCCACT	
β-tublin-R	TGGTTCAGGTCACCGTAAGAGG	
actin-F	ACGGTGTCGTCACAAACTGG	Internal reference gene
actin-R	CGGTTGGACTTAGGGTTGATAG	

Oxidative stress assays

To evaluate the role of *sntB* in fungal resistance to oxidative stress, a series concentration (0, 0.12, 0.24, and 0.36 mM) of menadione sodium bisulfite (MSB) were added to the medium. 10^6 fungal spores for each strain were inoculated on the medium and cultured in dark at 37℃. The diameters of colonies were measured 3 d after inoculation and the inhibition rate was calculated as previously described [33].

The AFB1 product was analyzed by TLC after the strains were cultured in YES

medium in dark at 29°C for 7 d.

Statistical analysis

All data in this study were expressed as mean ± standard deviation. The statistical

analysis was performed using the software GraphPad Prism8 (GraphPad Software, La

Jolla, CA, USA). The difference was considered to be statistically significant when

$P<0.05$.

Results

The phenotype of SntB in *A. flavus*

The role of SntB in *A. flavus* has been previously characterized by analyzing both

Δ*sntB* and overexpression *sntB* genetic mutants [20]. To further investigate the intrinsic

mechanism of this regulator on the development and aflatoxin biosynthesis in *A.*

flavus, the *sntB* deletion strain (Δ*sntB*) and the complementary strain (Com-*sntB*)

were constructed by the method of homologous recombination and verified by

diagnostic PCR and southern blotting (Figure S1). The expression levels of *sntB* in

WT, Δ*sntB*, and Com-*sntB* strains was further detected by qRT-PCR and the result

showed that the expression of *sntB* was absent in the gene-deletion strains, and it fully

recovered in the Com-*sntB* (Figure S1), which reflected that the *sntB* and Com-*sntB*

had been constructed, and could be used in the subsequent experiments of this study.

The phenotype analysis of this study revealed that the deletion of *sntB* gene

significantly inhibited the growth of mycelium, hyphae morphology, the length of

fungal cell (between two adjacent septa), the number of conidiation, sclerotium

formation, and aflatoxin biosynthesis, while the above phenotypes of both

development and mycotoxin bio-synthesis were fully recovered in the Com-*sntB*

strain (Figure 1). To reveal the signaling pathways of *sntB* in conidiation, sclerotium

formation, and aflatoxin biosynthesis, qRT-PCR analysis was performed to assess the

expression levels of sporulation related transcriptional factor genes, *steA*, *WetA*, *fluG*,

and *veA,* sclerotia formation related transcriptional factor genes, *nsdC*, *nsdD*, and *sclR*

[46, 47], and the AFs synthesis gene cluster structural genes *aflC*, *aflR,* and *aflP*, and the

main regulatory gene *aflR* and *aflS*. As shown in Figure S2, the relative expression

levels of these genes were significantly lower in the Δ*sntB* strain compared to that of

the WT strain, and recovered in the Com-*sntB* strain. These results indicated that SntB

regulates the conidiation, sclerotium formation, and aflatoxin biosynthesis by the

canonical signaling pathways mediated by these regulators.

Figure 1. The functions of SntB in *A. flavus*. (A) The colonies of WT, Δ*sntB*, and

Com-*sntB* strains grown on PDA at 37°C in dark for 4 d. (B) The colony diameter

statistics of the above fungal strains. (C) Microscopic examination revealed the

difference in mycelia of each fungi strain at 37°C in dark, scale=200 μm. (D)

Microscopic examination of the hyphal septum of each strain at 37°C in dark,

scale=50 μm. (E) The spore production statistics. (F) All the above fungal strains were

point-inoculated on CM medium and grown for 7 d at 37°C. (G) The number of

sclerotia of the above fungal strains. ND=Not detectable. (H) AFB1 production of the

above fungal strains was detected by TLC after the strains incubating at 29°C in PDB

medium for 7 d.

SntB plays important role in virulence of *A. flavus* to both plant and animal hosts

In order to explore the effect of SntB on the fungal colonization ability, peanut and

maize kernels were infected with spore solution of each fungal strain. Compared with

WT, the conidiation yield of Δ*sntB* on the infected host was significantly reduced

($P<0.001$) and no AFB1 could be detected on the hosts infected by Δ*sntB*, while in the

Com-*sntB* strain, the capacity to produce conidia and AFB1 on both crop kernels

recovered (Figure 2A-2C and Figure S3A). The role of *sntB* in fungal virulence to

animals was also investigated. As Figure 2D-2E shown, the survival rate of silkworms

injected by spores of Δ*sntB* strain was significantly higher than that from WT infected

larvae. There were less fungal mycelium, conidia, and AFB1 production on the dead

silkworms injected by Δ*sntB* compared to the silkworms from the WT injection

groups, but when the gene was reintroduced (i.e. the Com-*sntB* group), similar to

what found in the WT group, the survival rate of silkworms obviously dropped and

more fungal mycelium, conidia, and AFB1 produced on the dead silkworms (Figure

2F-2H). All the above results revealed that SntB plays an essential role in virulence of

A. flavus.

The capacity of fungal infection is closely related to secreted hydrolases, such as

amylase, lipase, protease, etc. In order to explore the effect of SntB on the activity of

hydrolases, the activities of amylase in the above each fungal strain were further

determined. The results showed that the colonies of Δ*sntB* produced almost no

degradation transparent circle after adding iodine solution compared with that of WT

and Com-*sntB*, which indicated that the activity of α-amylase in Δ*sntB* strain was

significantly reduced ($P<0.001$) (Figure S3B and S3C). This suggests that SntB plays

an important role in the fungal pathogenicity by changing the hydrolases activity of *A.*

flavus.

Figure 2. The role of SntB on the ability of *A. flavus* to colonize host. (A) Phenotypic

of corn and peanut kernels colonized by Δ*sntB*, Com-*sntB*, and WT strains at 29°C in

dark for 7 d. (B) Statistical of the number of conidia on the surface of peanut and

maize kernels. (C) TLC analysis to detect the yield of AFB1 in kernels infected by the

above fungal strains after 7 d incubation. (D) Photographs of the silkworms infected

by the above fungal strains. (E) The survival rate of silkworms 1 week after injection

of the above strains. (F) Photographs of the dead silkworms infected by *A. flavus* after

6 d incubation. (G) The spore production statistics of the above fungal strains on the

dead silkworms shown in (F). (H) TLC analysis of AFB1 levels produced in infected

dead silkworms in (F).

SntB chords global gene expression

To explore the downstream signaling pathways regulated by SntB, samples with three

biological replicates of WT and Δ*sntB* strains were submitted for RNA-Seq. In the

assay, the bases score Q30 was more than 93.19% (Table S1) and mapping ratio was

from 95.17% to 95.80% (Table S2). To further confirm the quality of RNA-Seq, PCA

(Principal Components Analysis) and Pearson correlation analysis were performed.

Correlation analysis revealed that the samples were clustered by groups (Figure 3A).

A plot of PC1 (47.70%) and PC2 (23.20%) scores showed a clear separation between

the groups (Figure 3B). A total of 1,446 and 1,034 genes were significantly up- and

down-regulated, respectively, in the Δ*sntB* compared to the WT strain (Figure 3C and

Supplementary Table S3). GO enrichment analysis identified 93 enriched GO terms

(*P*<0.05) (Figure 3D and Supplementary Tables S4). In the biological process

category, the most enriched terms were "oxidation-reduction process (GO: 0055114)".

In the molecular function category, "catalytic activity (GO: 0003824),"

"oxidoreductase activity (GO: 0016491)," and "cofactor binding (GO: 0048037)"

were the most significantly enriched terms. Whereas terms associated with "Set3

complex (GO: 0034967)", "mitochondrial crista junction (GO: 0044284)" and

"extracellular region (GO: 0005576)" were significantly enriched in the cellular

component category. Additionally, all the DEGs were mapped according to the KEGG

database, and 42 significantly enriched pathways were identified ($P<0.05$)

(Supplementary Table S5). Among them, "metabolic pathways (ko01100)," "aflatoxin

biosynthesis (ko00254)," and "microbial metabolism in diverse environments

(ko01120)" were the most significantly enriched (Figure 3E and Supplementary Table

S5).

Figure 3. SntB chords global gene expression in *A. flavus*. (A) The Pearson

correlation results shown by heatmap. (B) Principal component analysis (PCA) on six

fungal samples, including three Δ*sntB* (KOsnt2) and three WT fungal strains. (C)

Volcano map reflecting the distribution of the differential expression genes. (D) Gene

Ontology (GO) analyses of the differentially expressed genes. (E) Kyoto encyclopedia

of genes and genomes (KEGG) analyses of the differentially expressed peaks related

genes.

Characterization the binding regions of SntB by chromatin immunoprecipitation

sequencing (ChIP-seq)

To characterize the chromatin regions targeted by SntB, ChIP-seq studies were carried

out with both HA tag fused *sntB* strain (*sntB*-HA) and WT strain. The *sntB*-HA strain

was constructed by homologous recombination through fused HA to the 3' end of

sntB (Figure 4A). In the ChIP-seq assay, more than 94.66% of bases score Q30 and

above in each sample (Table S6), and reaching 52.50% to 94.48% of mapping ratio

(Table S7). The principal component analysis (PCA) (Figure S4A) and heat map

(Figure S4B) reflected that the quality of samples was competent for subsequent

ChIP-seq analysis. There were 1,510 up-enriched differently accumulated peaks

(DAPs) in *sntB*-HA fungal strain compared to the WT strain, which were distributed

on the whole *A. flavus* genome (Figure 4B, and Table S8). Most of the up-enriched

peaks were located in the promoter (82.85%) region (Figure 4C). The genes of the

DAPs were further subjected to GO and KEGG analysis. The most strikingly enriched

GO terms in the biological process category were "cell communication

(GO:0007154)", "response to stimulus (GO:0050896)", and "response to external

stimulus (GO:0009605)". Whereas terms associated with "DNA-binding transcription

factor activity (GO:0003700)", "DNA-binding transcription factor activity, RNA

polymerase II-specific (GO:0000981)", and "sequence-specific DNA binding

(GO:0043565)" were the most significantly enriched molecular function category

(Figure 4D and Table S9). And these genes were mostly enriched in "Methane

metabolism" and "MAPK signaling pathway - yeast" pathways (P value <0.05)

(Figure 4E and Table S10).

Figure 4. Characterization the binding regions of SntB. (A) Verification of the

construction of sntB-HA strain using Western blot. (B) The distribution of differently

accumulated peaks on the genome. (C) Vennpie map of the differently accumulated

peaks distribution on gene functional elements. (D) GO analyses of the of the

differently accumulated peaks related genes. (E) KEGG analyses of the differently

accumulated peak related genes.

Integration of the results of ChIP-seq and RNA-seq assays

After the overlapping the results from both different sequence methods (ChIP-seq and

RNA-seq), 238 DEGs were found (Figure 5A). According to the GO annotation, these

DEGs were significantly enriched in 8 GO terms, including "cellular response to

reactive oxygen species (GO:0034614)", "reactive oxygen species metabolic process

(GO:0072593)", and "cellular response to oxygen-containing compound

(GO:1901701)" (Figure 5B). It was further noted that the DEGs were significantly

assigned to "Carbon metabolism (afv01200)", "Peroxisome (afv04146)", and

"Glyoxylate and dicarboxylate metabolism (afv00630)" KEGG pathways (Figure 5C

and Table S11-S12). These results revealed that SNT2 is essential for *A. flavus* to

maintain the homeostasis of intracellular reactive oxygen species. Studies had shown

that SNT2 could response to oxidative stress in yeast [27] and *Magnaporthe oryzae* [48].

As Figure 5D and Figure 5E showed, due to the deletion of the *sntB* gene, Δ*sntB*

exhibited a severe MSB sensitivity phenotype compared to that of the wild-type (WT)

strain, and the phenotype recovered in the complementary strain (Com-Δ*sntB*). The

results showed that the inhibition rate of oxidant MSB to Δ*sntB* would be significantly

enhanced with the increase of MSB concentration. These results showed that SntB

deeply participates in the regulation of oxidative stress pathway. As the most abundant

peroxisomal enzyme, catalases catalyze decomposition of hydrogen peroxide [49]. As

the most abundant peroxisomal enzyme, catalases catalyze decomposition of

hydrogen peroxide [49]. To further study the mechanism of SntB mediated oxidative

response of *A. flavus*, the *catC* (encode a catalase) gene was selected based on the

above integration results. The relative expression levels of *catC* in wild-type and

Δ*sntB* strains under MSB treatment were measured. As Figure 5F shown, the

expression level of *catC* was significantly higher in Δ*sntB* strain than in WT strain, it

suggested that to compensate the absence of *sntB*, *catC* is up-regulated to respond the

higher intracellular oxidative level. However, under the stress of oxidant MSB, the

deletion of *sntB* obvious suppressed the expression level of *catC* compared to that of

WT strain, which reflected that the absence of *sntB* significantly impaired the capacity

of *catC* to further respond to extra external oxidative stress. These results revealed

that SntB is deeply involved in CatC mediated oxidative response in *A. flavus*.

Figure 5. Integration of the results of ChIP-seq and RNA-seq assays. (A) Venn

diagrams of ChIP-seq and RNA-seq. (B) GO analyses of the common genes. (C)

KEGG analyses of the common genes. (D) The phenotype of WT, ΔsntB, and

Com-sntB strains cultured in PDA containing a series concentration of MSB for 3 d.

(E) Statistical analysis of the growth inhibition rate of MSB to all the above fungal

strains according to Panel D. (F) The relative expression level of sod1 in WT and

367   Δ*sntB* strains with or without MSB treatment.

369   CatC is important for *A. flavus* response to oxidative stress

The functions of *catC* gene in *A. flavus* were further explored by knockout of the *catC*

(Figure S5). As shown in Figure 6A-6C, the diameter of Δ*catC* strain was

significantly smaller than that of WT, and the conidia number in the Δ*catC* strain

decreased significantly compared to that of WT. The sclerotia production of Δ*catC*

strain was also significantly less than that of WT strain (Figure 6D and 6E). In view of

Catc is involved in the oxidative response pathways of *A. flavus* (Figure 4), both

Δ*catC* and WT strains were treated by a serial concentration of MSB, and the results

showed that the inhibition rates of MSB to Δ*catC* strain was significantly lower than

that of WT group (Figure 6F and 6G). Catalase is a major peroxisome protein and

plays a critical role in removing peroxisome-generated reactive oxygen species. This

result echoed that deletion of *sntB* increased intracellular oxidative level and the

inhibition rate of MSB, and up-regulated the expression of *catC* (Figure 5D-5F). The

role of CatC in the bio-synthesis of AFB1 was also assessed (Figure 6H). The results

showed that a relatively large amount of AFB1 was produced by the Δ*catC* strain

compared to the WT. But when under the stress of MSB, AFB1 yield of the WT strain

was significantly more than that of Δ*catC* strain. All the above results revealed that

the *catC* plays an important role in SntB mediated regulation pathway on fungal

morphogenesis, oxidative stress responding and AFB1 production.

Figure 6. The functions of *catC* in *A. flavus*. (A) The colonies of WT and Δ*catC*

strains grown on PDA at 37°C in dark for 4 d. (B) The colony diameter statistics of

the above fungal strains. (C) The spore production statistics of the above fungal

strains. (D) All above fungal strains were point-inoculated on CM medium and grown

for 7 d at 37°C. (E) The number of sclerotia of the above fungal strains. (F) The

phenotype of above strains cultured in PDA medium containing a series concentration

of MSB for 3 d. (G) Statistical analysis of the growth inhibition rate of MSB to all the

above fungal strains according to (F). (H) AFB1 production of the above fungal

strains was detected by TLC after the strains incubating at 29°C in PDB medium for 7

d.

SntB regulates fungal virulence through peroxidase mediated lipolysis

Biogenesis of peroxisomes was reported to promote lipid hydrolysis, produce the

production of glycerol, and further change fungal pathogenicity [50]. Due to deeply

involved in oxidative response of *A. flavus*, we wondered if SntB also takes part in the

regulation of the production of lipid and glycerol. As shown in Table S13, one gene

(G4B84_008359) in lipase activity GO term was significantly down-regulated in

Δ*sntB* strain, which encoded a secretory lipase belonged to the virulence factors

reported in *Pseudomonas aeruginosa* [51]. The lipase activity was assayed by

examining the ability to cleave glycerol tributyrate substrate [52]. The results showed

that the colony diameter of Δ*sntB* strain on PDA medium with tributyrin were

significantly smaller than that of the control, and the colony diameters of WT and

Com-*sntB* strains on PDA medium were obviously bigger than those on 0.3%

tributyrin PDA medium. The relative inhibition rate of tributyrin to the colony growth

of Δ*sntB* strains was significantly higher than that of WT and Com-*sntB* strain (Figure

S3D and S3E). Our previous study revealed that H3 lysine 36 trimethylation

(H3K36me3) modification on the chromatin region of the sntB is regulated by AshA

and SetB [32]. H3K36me3 usually promote gene transcription [53, 54]. Our study

revealed the potential machinery associated with SntB mediated regulation on fungal

morphogenesis, mycotoxin anabolism and fungal virulence, which lurks the axle of

from SntB to fungal virulence and mycotoxin bio-synthesis through lipid catabolism

(i.e. H3K36me3 modification-SntB-Peroxisomes-Lipid hydrolysis-fungal virulence

and mycotoxin bio-synthesis).

Figure 7. SntB regulate peroxisome biogenesis, fatty acid utilization, and fungal

pathogenicity *in A. flavus*. (A) The phenotype of each strain on PDA medium

containing 0.3% tributyrin, (B) Statistics of inhibition rates. The asterisk *** above

the bars represents significantly different (p<0.001). (C) Mechanistic diagram of the

bio-functions of SntB in *A. flavus*.

Discussion

SntB is a conserved regulator in many species, including *A. nidulans* [25],

Saccharomyces cerevisiae [27], *Schizzosaccharomyces pombe* [55], *Fusarium*

oxysporum [30], *Neurospora crassa* [31], and *A. flavus* [20, 25]. SntB can regulate the

production of uncharacterized secondary metabolites, including aspergillicin A1 and

aspergillicin F2 [56]. MoSnt2 protein was required for regulation of

infection-associated autophagy in *Magnaporthe oryzae* [48]. In *A. flavus*, the functions

of *sntB* gene were analyzed by both Δ*sntB* and overexpression *sntB* genetic mutants[20].

SntB deletion impaired several developmental processes, such as sclerotia formation

and heterokaryon compatibility, secondary metabolite synthesis, and ability to

colonize host seeds, which were consistent with our results (Figure 1 and 2). Unlike, a

complementation strain was constructed in this study which further clarified and

confirmed the function of *sntB* gene. In this study, the potential mechanism under

these effects were further analyzed by detection the related transcriptional factor

genes of sporulation (*steA*, *WetA*, *fluG*, and *veA*), sclerotia formation (*nsdC*, *nsdD*, and

sclR), and the AFs synthesis related genes *aflC*, *aflD*, *aflO*, *aflP*, and *aflR* (Figure S2).

In the RNA-seq data, we also found some DEGs related to AFs synthesis (*aflB*, *aflE*,

aflH, *aflK*, *aflN*, *aflO*, *aflP*, *aflQ*, *aflR*, *aflS*, *aflV*, and *aflW*) (Figure S6). And all these

genes were down-regulated, which was consistent with that the AFs production in

Δ*sntB* was significantly decreased compared to WT and Com-*sntB* (Figure 1F). These

results inferred that SntB regulated the morphogenesis and the production of *A. flavus*

through the above canonical signal pathways.

For the process of *A. flavus* invading hosts, in view of it is a notorious pathogen for

plant and animal, we established both crop and insect models, especially silkworm

represented animal mode profoundly revealed the critical role of SntB in fungal

virulence (Figure 2). The results of crop kernel models showed that the number of

spores of Δ*sntB* on kernels of both peanut and maize was dramatically lower than that

of strains WT and Com-*sntB* (Figure 2A and 2B) and almost no AFB1 was detected

on maize and peanut kernels inoculated with Δ*sntB*, while plenty of AFB1 were

detected from the kernels infected by WT fungal strain, and AFB1 biosynthesis

capacity of Com-*sntB* strains recovered compared to the Δ*sntB* and WT fungal strains

(Figure 2C). These results were corroborated by previous study [20]. It was also found

in this study that the survival rate of silkworms injected by spores of Δ*sntB* strain was

significantly higher than the silkworms injected with spores from WT and Com-*sntB*

fungal strains (Figure 2D and 2E). What's more, we also assayed the effect of SntB on

the activity amylase, which was closely related to the capacity of fungal infection [57].

As shown in Figure S3C, after adding iodine solution, the Δ*sntB* strain almost did not

produce a degradation transparent ring compared with wild-type and complementary

strains, indicating the amylase activity of the *sntB* gene knockout strain was

significantly decreased (*P*<0.001). Our results comprehensively reveal the important

function of SntB in the growth, development, secondary metabolite synthesis, and

virulence of *A. flavus*.

SntB was reported as an important epigenetic reader. In *A. flavus*, SntB was reported

to regulate global histone modifications (acetylation and methylation) and interact

with EcoA and RpdA to form a conserved chromatin regulatory complex [20, 29]. Loss

of *sntB* in *Magnaporthe orzyae* also led to an increase in H3 acetylation [48]. In our

RNA-seq data, we also found a set domain containing histone-lysine

N-methyltransferase (Ash1, G4B84_009862) was down-regulated in Δ*sntB* strain

compared to WT (Table S3), which was reported to regulate mycotoxin metabolism

and virulence via H3K36 methylation in *A. flavus* [32]. Besides, SntB is reported to be

a transcriptional regulator in *A. nidulans* [25] and *Fusarium oxysporum* [30]. So, we use

RNA-seq and ChIP-seq to study the transcriptional response of *sntB* in *A. flavus*. By

integration analysis the results of ChIP-seq and RNA-seq assays, we found that the

enriched GO terms and KEGG pathways of the DEGs were related to oxidative

response (Figure 5A-5C). These results reflected that Snt2 plays an important role in

fungal responding to oxidative stress, which is consistent with previous reports that

Snt2 could response to oxidative stress in yeast [27], *Fusarium oxysporum* [30], and

Magnaporthe oryzae [48].

As a harmful by-product of oxidative metabolism, reactive oxygen species (ROS) is

unavoidable and essential for fungus development [58, 59]. ROS has also been shown to

be required for aflatoxin production [60-62]. Several oxidative stress-responsive

transcription factors have been identified as regulating aflatoxin production, including

AtfB, AP-1, and VeA [61, 63-65]. Previous studies shown that Snt2 protein coordinates

the transcriptional response to hydrogen peroxide-mediated oxidative stress in the

yeast [26, 27] and is involved in fungal respiration and reactive oxidative stress in

Fusarium oxysporum and *Neurospora crassa* [30, 31]. Several GO terms ("cellular

response to reactive oxygen species", "reactive oxygen species metabolic process",

and "cellular response to oxygen-containing compound") and KEGG pathways

(Peroxisome) were enriched by the DEGs screened out by integration of ChIP-seq and

RNA-seq data in this study (Figure 5). This is the first time to show that the SntB in *A.*

flavus is important in oxidative stress response, through which SntB participates the

regulating of aflatoxin bio-synthesis and fungal development.

Fungal defense against ROS is mediated by superoxide dismutases (SOD), catalases

(CAT), and glutathione peroxidases (GPX). The effect of MSB on cellular growth and

antioxidant enzyme induction in *A. flavus* was previously explored [66-68]. Once in the

cell, menadione may release superoxide anion [69], which was scavenged by

superoxide dismutases (SOD) and transformed into hydrogen peroxide, or react with

nitric oxide to form peroxynitrite [70]. This study found that after knock out *sntB* gene,

the strain growth was significantly inhibited by MSB (Figure 5F and 5G). Some genes

encoded superoxide dismutases and catalases were reported to associated with AF/ST

synthesis [71], including *mnSOD*, *sod1*, *sod2*, *catA*, *catB*, and *hyr1*. In our RNA-seq

data, 7 related genes were screened out (Table S14), among which, bZIP transcription

factor *Atf21* (G4B84_008675), *catlase C* (G4B84_009242), *catlase A*

(G4B84_010740), superoxide dismutases *sod2* (G4B84_003204), peroxisomal

membrane protein *PmpP24* (G4B84_001452) were up-regulated, while *catlase B*

(G4B84_008381) and superoxide dismutase *sod1* (G4B84_009129) were

down-regulated in Δ*sntB* strain, respectively.

Some studies reported the correlation between ROS formation, aflatoxin production,

and antioxidant enzyme activation. Aflatoxin B1 biosynthesis and the activity of total

superoxide dismutase were effectively inhibited by cinnamaldehyde, whereas the

activities of catalase and glutathione peroxidase were opposite [72]. The expression of

catA, *cat2* and *sod1*, and CAT enzymatic activity were opposite correlated to AFB1

biosynthesis under AFs inhibitor piperine treatment [73]. Deletion the gene

of *sod* (GenBank accession no: CA747446) reduced AFs production [74], which was

most similar to *sod2* (G4B84_003204). The mitochondria-specific *sod* and the

genes *aflA*, *aflM*, and *aflP* belonging to the AFs gene cluster were reported to

co-regulated [75]. Ethanol can inhibit fungal growth and AFB_1 production in *A.*

flavus and enhanced levels of anti-oxidant enzymatic genes

including *Cat*, *Cat1*, *Cat2*, *CatA*, and Cu, Zn superoxide dismutase gene *sod1*. All

these reports indicated that the expression of anti-oxidant enzymatic genes was

opposite correlated to AFB1 biosynthesis.

In our study, 7 genes related to oxidative response were obviously differentially

expressed in transcriptome data (Table S14). Among these DEGs, 5 out of 7 genes

were up-regulated in Δ*sntB* strain. Based on the AFs production in Δ*sntB* was

significantly decreased compared to WT and Com-*sntB* (Figure 1F), the most

up-regulated gene in Δ*sntB* strain, *catC* (G4B84_000242), was selected for further

analysis. We found that the deletion of *sntB* significantly up-regulated the *catC* gene,

however, the expression of the *catC* gene was almost undetectable under MSB

treatment (Figure 5F). Results also showed that the inhibition rates of MSB to Δ*catC*

strain was significantly lower than that of WT group and AFB1 yield of the Δ*catC*

strain was significantly decreased than that of WT strain under the stress of MSB

(Figure 6F-6H). These results indicated that SntB is profoundly involved in the *catC*

mediated oxidative response.

Peroxisomes are intimately associated with the metabolism of lipid droplets [76] and

the histone lysine methyltransferase ASH1 promotes peroxisome biogenesis, inhibits

lipolysis, and further affects pathogenesis of *Metarhizium robertsii* [50]. Set2 histone

methyltransferase family in *A. flavus*, AshA and SetB, were found to regulate

mycotoxin metabolism and virulence via H3K36me3, including the chromatin region

of the *sntB* [32]. By ChIP-seq and RNA-seq, SntB was found to essential for *A. flavus*

to maintain the homeostasis of intracellular reactive oxygen species (Figure 5A) and

several anti-oxidant enzymes were up-regulated in Δ*sntB* strain (Table S14). In

addition, we also found only one down-regulated DEG (G4B84_008359) in lipase

activity GO term in our RNA-seq data (Table S13), which encodes a secretory lipase

and belongs to the virulence factors reported in *Pseudomonas aeruginosa* [51]. These

results suggested that SntB plays a pivotal role in regulating peroxisome biogenesis to

promote lipolysis involving in fungal pathogenesis.

Overall, we explored and clarified the bio-function of the SntB and found that SntB

responses to oxidative stress through related oxidoreductase represented by CatC in *A.*

flavus (Figure 7). Our study revealed the potential machinery associated with SntB

mediated regulation on fungal morphogenesis, mycotoxin anabolism and fungal

virulence, which lurks the axle of from SntB to fungal virulence and mycotoxin

bio-synthesis (i.e. SntB -Peroxisomes-Lipid hydrolysis-fungal virulence and

mycotoxin bio-synthesis). The work of this study provided a novel perspective for

developing new prevention and control strategies against pathogenic fungi.

Compliance with ethical requirements

This article does not contain any studies with human or animal subjects.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Data Availability

All data needed to evaluate the conclusions are present in the paper and/or the

Supporting Information. Raw data of the ChIP and RNA-seq were submitted to

GSE247683.

Funding

This work was funded by the grants of the National Natural Science Foundation of

China (No. 32070140), the Nature Science Foundation of Fujian Province (No.

2021J02026), and the State Key Laboratory of Pathogen and Biosecurity (Academy of

Military Medical Science) (SKLPBS2125).

Acknowledgement

We especially thank Professor Jun Yuan, Xiuna Wang, Yu Wang and Xinyi Nie for

their support in instrument maintenance and reagent ordering.

References:

[1] Hedayati M.T., Pasqualotto A.C., Warn P.A.*, et al.* Aspergillus flavus: human

pathogen, allergen and mycotoxin producer [J]. Microbiology (Reading), 2007,

153(Pt 6): 1677-1692.

[2] Guan Y., Chen J., Nepovimova E.*, et al.* Aflatoxin Detoxification Using

Microorganisms and Enzymes [J]. Toxins (Basel), 2021, 13(1).

[3] Mishra H.N., Das C. A review on biological control and metabolism of

aflatoxin [J]. Crit Rev Food Sci Nutr, 2003, 43(3): 245-264.

[4] Vardon P.J., Mclaughlin C., Nardinelli C. Potential economic costs of

mycotoxins in the United States [J]. 2003.

[5] Riba A., Bouras N., Mokrane S.*, et al.* Aspergillus section Flavi and aflatoxins

in Algerian wheat and derived products [J]. Food Chem Toxicol, 2010, 48(10):

2772-2777.

[6] Plaz Torres M.C., Bodini G., Furnari M.*, et al.* Nuts and Non-Alcoholic Fatty

Liver Disease: Are Nuts Safe for Patients with Fatty Liver Disease? [J].

Nutrients, 2020, 12(11).

[7] Cleveland T.E., Yu J., Fedorova N.*, et al.* Potential of Aspergillus flavus

genomics for applications in biotechnology [J]. Trends Biotechnol, 2009,

27(3): 151-157.

[8] Yabe K., Nakajima H. Enzyme reactions and genes in aflatoxin biosynthesis

[J]. Appl Microbiol Biotechnol, 2004, 64(6): 745-755.

[9]   Yu J. Current understanding on aflatoxin biosynthesis and future perspective in reducing aflatoxin contamination [J]. Toxins (Basel), 2012, 4(11): 1024-1057.

[10]   Roze L.V., Hong S.Y., Linz J.E. Aflatoxin biosynthesis: current frontiers [J]. Annu Rev Food Sci Technol, 2013, 4: 293-311.

[11]   Ehrlich K.C. Predicted roles of the uncharacterized clustered genes in aflatoxin biosynthesis [J]. Toxins (Basel), 2009, 1(1): 37-58.

[12]   Chang P.K. The Aspergillus parasiticus protein AFLJ interacts with the aflatoxin pathway-specific regulator AFLR [J]. Mol Genet Genomics, 2003, 268(6): 711-719.

[13]   Price M.S., Yu J., Nierman W.C.*, et al.* The aflatoxin pathway regulator AflR induces gene transcription inside and outside of the aflatoxin biosynthetic cluster [J]. FEMS Microbiol Lett, 2006, 255(2): 275-279.

[14]   Xu J., Luo X. Molecular biology of aflatoxin biosynthesis [J]. Wei Sheng Yan Jiu, 2003, 32(6): 628-631.

[15]   Yu J.H., Keller N. Regulation of secondary metabolism in filamentous fungi [J]. Annu Rev Phytopathol, 2005, 43: 437-458.

[16]   Georgianna D.R., Payne G.A. Genetic regulation of aflatoxin biosynthesis: from gene to genome [J]. Fungal Genet Biol, 2009, 46(2): 113-125.

[17]   Montibus M., Pinson-Gadais L., Richard-Forget F.*, et al.* Coupling of transcriptional response to oxidative stress and secondary metabolism regulation in filamentous fungi [J]. Crit Rev Microbiol, 2015, 41(3): 295-308.

[18]   Lv Y., Wang J., Yang H.*, et al.* Lysine 2-hydroxyisobutyrylation orchestrates

cell development and aflatoxin biosynthesis in Aspergillus flavus [J]. Environ

Microbiol, 2022, 24(9): 4356-4368.

[19]   Yang G., Yue Y., Ren S.*, et al.* Lysine acetylation contributes to development,

aflatoxin biosynthesis and pathogenicity in Aspergillus flavus [J]. Environ

Microbiol, 2019, 21(12): 4792-4807.

[20]   Pfannenstiel B.T., Greco C., Sukowaty A.T.*, et al.* The epigenetic reader SntB

regulates secondary metabolism, development and global histone

modifications in Aspergillus flavus [J]. Fungal Genet Biol, 2018, 120: 9-18.

[21]   Ren S., Yang M., Yue Y.*, et al.* Lysine Succinylation Contributes to Aflatoxin

Production and Pathogenicity in Aspergillus flavus [J]. Mol Cell Proteomics,

2018, 17(3): 457-471.

[22]   Wang Y., Yang M., Ge F.*, et al.* Lysine Succinylation of VBS Contributes to

Sclerotia Development and Aflatoxin Biosynthesis in Aspergillus flavus [J].

Mol Cell Proteomics, 2023, 22(2): 100490.

[23]   Lv Y. Proteome-wide profiling of protein lysine acetylation in Aspergillus

flavus [J]. PLoS One, 2017, 12(6): e0178603.

[24]   Wang J., Liang L., Wei S.*, et al.* Histone 2-Hydroxyisobutyryltransferase

Encoded by Afngg1 Is Involved in Pathogenicity and Aflatoxin Biosynthesis

in Aspergillus flavus [J]. Toxins (Basel), 2022, 15(1).

[25]   Pfannenstiel B.T., Zhao X., Wortman J.*, et al.* Revitalization of a Forward

Genetic Screen Identifies Three New Regulators of Fungal Secondary

Metabolism in the Genus Aspergillus [J]. mBio, 2017, 8(5).

[26] Singh R.K., Gonzalez M., Kabbaj M.H.*, et al.* Novel E3 ubiquitin ligases that regulate histone protein levels in the budding yeast Saccharomyces cerevisiae [J]. PLoS One, 2012, 7(5): e36295.

[27] Baker L.A., Ueberheide B.M., Dewell S.*, et al.* The yeast Snt2 protein coordinates the transcriptional response to hydrogen peroxide-mediated oxidative stress [J]. Mol Cell Biol, 2013, 33(19): 3735-3748.

[28] Tannous J., Barda O., Luciano-Rosario D.*, et al.* New Insight Into Pathogenicity and Secondary Metabolism of the Plant Pathogen Penicillium expansum Through Deletion of the Epigenetic Reader SntB [J]. Front Microbiol, 2020, 11: 610.

[29] Karahoda B., Pardeshi L., Ulas M.*, et al.* The KdmB-EcoA-RpdA-SntB chromatin complex binds regulatory genes and coordinates fungal development with mycotoxin synthesis [J]. Nucleic Acids Res, 2022, 50(17): 9797-9813.

[30] Denisov Y., Freeman S., Yarden O. Inactivation of Snt2, a BAH/PHD-containing transcription factor, impairs pathogenicity and increases autophagosome abundance in Fusarium oxysporum [J]. Mol Plant Pathol, 2011, 12(5): 449-461.

[31] Denisov Y., Yarden O., Freeman S. The transcription factor SNT2 is involved in fungal respiration and reactive oxidative stress in Fusarium oxysporum and Neurospora crassa [J]. Physiological and Molecular Plant Pathology, 2011,

666 76(2): 137-143.

667 [32] Zhuang Z., Pan X., Zhang M.*, et al.* Set2 family regulates mycotoxin
668 metabolism and virulence via H3K36 methylation in pathogenic fungus
669 *Aspergillus flavus* [J]. Virulence, 2022, 13(1): 1358-1378.

670 [33] Pan X., Hao L., Yang C.*, et al.* SWD1 epigenetically chords fungal
671 morphogenesis, aflatoxin biosynthesis, metabolism, and virulence of
672 *Aspergillus flavus* [J]. Journal of Hazardous Materials, 2023: 131542.

673 [34] Hu Y., Yang G., Zhang D.*, et al.* The PHD Transcription Factor Rum1
674 Regulates Morphogenesis and Aflatoxin Biosynthesis in [J]. Toxins, 2018,
675 10(7).

676 [35] Pan X., Hao L., Yang C.*, et al.* SWD1 epigenetically chords fungal
677 morphogenesis, aflatoxin biosynthesis, metabolism, and virulence of
678 *Aspergillus flavus* [J]. J Hazard Mater, 2023, 455: 131542.

679 [36] Kaminskyj S.G. Septum position is marked at the tip of Aspergillus nidulans
680 hyphae [J]. Fungal Genet Biol, 2000, 31(2): 105-113.

681 [37] Zhang Y., Fan J., Ye J.*, et al.* The fungal-specific histone acetyltransferase
682 Rtt109 regulates development, DNA damage response, and virulence in
683 Aspergillus fumigatus [J]. Mol Microbiol, 2021, 115(6): 1191-1206.

684 [38] Wen M., Lan H., Sun R.*, et al.* Histone deacetylase SirE regulates
685 development, DNA damage response and aflatoxin production in Aspergillus
686 flavus [J]. Environmental Microbiology, 2022.

687 [39] Hao L., Zhang M., Yang C.*, et al.* The epigenetic regulator Set9 harmonizes

fungal development, secondary metabolism, and colonization capacity of

Aspergillus flavus [J]. International Journal of Food Microbiology, 2023, 403:

110298.

[40] Liao Y., Smyth G.K., Shi W. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features [J]. Bioinformatics,

2013, 30(7): 923-930.

[41] Li H., Durbin R. Fast and accurate short read alignment with

Burrows-Wheeler transform [J]. Bioinformatics, 2009, 25(14): 1754-1760.

[42] Zhou X., Su Z. EasyGO: Gene Ontology-based annotation and functional

enrichment analysis tool for agronomical species [J]. BMC Genomics, 2007, 8:

246.

[43] Xie C., Mao X., Huang J.*, et al.* KOBAS 2.0: a web server for annotation and

identification of enriched pathways and diseases [J]. Nucleic Acids Res, 2011,

39(Web Server issue): W316-322.

[44] Mengjuan Z., Guanglan L., Xiaohua P.*, et al.* The PHD transcription factor

Cti6 is involved in the fungal colonization and aflatoxin B1 biological

synthesis of Aspergillus flavus [J]. IMA Fungus, 2021, 12(1): 12.

[45] Hu Y., Yang G., Zhang D.*, et al.* The PHD Transcription Factor Rum1

Regulates Morphogenesis and Aflatoxin Biosynthesis in Aspergillus flavus [J].

Toxins (Basel), 2018, 10(7): 301.

[46] Cary J.W., Harris-Coward P.Y., Ehrlich K.C.*, et al.* NsdC and NsdD affect

Aspergillus flavus morphogenesis and aflatoxin production [J]. Eukaryot Cell,

710   2012, 11(9): 1104-1111.

711   [47] Yang G., Hu Y., Fasoyin O.E.*, et al.* The Phosphatase CDC14 Regulates

712   Development, Aflatoxin Biosynthesis and Pathogenicity [J]. Frontiers In

713   Cellular and Infection Microbiology, 2018, 8: 141.

714   [48] He M., Xu Y., Chen J.*, et al.* MoSnt2-dependent deacetylation of histone H3

715   mediates MoTor-dependent autophagy and plant infection by the rice blast

716   fungus Magnaporthe oryzae [J]. Autophagy, 2018, 14(9): 1543-1561.

717   [49] Okumoto K., Tamura S., Honsho M.*, et al.* Peroxisome: Metabolic Functions

718   and Biogenesis [J]. Adv Exp Med Biol, 2020, 1299: 3-17.

719   [50] Wang L., Lai Y., Chen J.*, et al.* The ASH1-PEX16 regulatory pathway controls

720   peroxisome biogenesis for appressorium-mediated insect infection by a fungal

721   pathogen [J]. Proc Natl Acad Sci U S A, 2023, 120(4): e2217145120.

722   [51] Papadopoulos A., Busch M., Reiners J.*, et al.* The periplasmic chaperone Skp

723   prevents misfolding of the secretory lipase A from Pseudomonas aeruginosa

724   [J]. Front Mol Biosci, 2022, 9: 1026724.

725   [52] Sieber M.H., Thummel C.S. The DHR96 nuclear receptor controls

726   triacylglycerol homeostasis in Drosophila [J]. Cell Metab, 2009, 10(6):

727   481-490.

728   [53] Zhao W., Neyt P., Van Lijsebettens M.*, et al.* Interactive and noninteractive

729   roles of histone H2B monoubiquitination and H3K36 methylation in the

730   regulation of active gene transcription and control of plant growth and

731   development [J]. New Phytol, 2019, 221(2): 1101-1116.

732   [54]   Kooistra S.M., Helin K. Molecular mechanisms and potential functions of

733   histone demethylases [J]. Nat Rev Mol Cell Biol, 2012, 13(5): 297-311.

734   [55]   Roguev A., Shevchenko A., Schaft D.*, et al.* A comparative analysis of an

735   orthologous proteomic environment in the yeasts Saccharomyces cerevisiae

736   and Schizosaccharomyces pombe [J]. Mol Cell Proteomics, 2004, 3(2):

737   125-132.

738   [56]   Greco C., Pfannenstiel B.T., Liu J.C.*, et al.* Depsipeptide Aspergillicins

739   Revealed by Chromatin Reader Protein Deletion [J]. ACS Chem Biol, 2019,

740   14(6): 1121-1128.

741   [57]   Mellon J.E., Cotty P.J., Dowd M.K. Aspergillus flavus hydrolases: their roles

742   in pathogenesis and substrate utilization [J]. Appl Microbiol Biotechnol, 2007,

743   77(3): 497-504.

744   [58]   Aguirre J., Ríos-Momberg M., Hewitt D.*, et al.* Reactive oxygen species and

745   development in microbial eukaryotes [J]. Trends Microbiol, 2005, 13(3):

746   111-118.

747   [59]   Gessler N.N., Aver'yanov A.A., Belozerskaya T.A. Reactive oxygen species in

748   regulation of fungal development [J]. Biochemistry (Mosc), 2007, 72(10):

749   1091-1109.

750   [60]   Jayashree T., Subramanyam C. Oxidative stress as a prerequisite for aflatoxin

751   production by Aspergillus parasiticus [J]. Free Radic Biol Med, 2000, 29(10):

752   981-985.

753   [61]   Roze L.V., Chanda A., Wee J.*, et al.* Stress-related transcription factor AtfB

integrates secondary metabolism with oxidative stress response in aspergilli

[J]. J Biol Chem, 2011, 286(40): 35137-35148.

[62] Yang L., Fountain J.C., Wang H.*, et al.* Stress Sensitivity Is Associated with

Differential Accumulation of Reactive Oxygen and Nitrogen Species in Maize

Genotypes with Contrasting Levels of Drought Tolerance [J]. Int J Mol Sci,

2015, 16(10): 24791-24819.

[63] Reverberi M., Zjalic S., Ricelli A.*, et al.* Modulation of antioxidant defense in

Aspergillus parasiticus is involved in aflatoxin biosynthesis: a role for the

ApyapA gene [J]. Eukaryot Cell, 2008, 7(6): 988-1000.

[64] Sakamoto K., Arima T.H., Iwashita K.*, et al.* Aspergillus oryzae atfB encodes

a transcription factor required for stress tolerance in conidia [J]. Fungal Genet

Biol, 2008, 45(6): 922-932.

[65] Baidya S., Duran R.M., Lohmar J.M.*, et al.* VeA is associated with the

response to oxidative stress in the aflatoxin producer Aspergillus flavus [J].

Eukaryot Cell, 2014, 13(8): 1095-1103.

[66] Wang X., Zha W., Liang L.*, et al.* The bZIP Transcription Factor AflRsmA

Regulates Aflatoxin B(1) Biosynthesis, Oxidative Stress Response and

Sclerotium Formation in Aspergillus flavus [J]. Toxins (Basel), 2020, 12(4).

[67] Vig I., Benkő Z., Gila B.C.*, et al.* Functional characterization of genes

encoding cadmium pumping P(1B)-type ATPases in Aspergillus fumigatus and

Aspergillus nidulans [J]. Microbiol Spectr, 2023, 11(5): e0028323.

[68] Zaccaria M., Ludovici M., Sanzani S.M.*, et al.* Menadione-Induced Oxidative

Stress Re-Shapes the Oxylipin Profile of Aspergillus flavus and Its Lifestyle

[J]. Toxins (Basel), 2015, 7(10): 4315-4329.

[69]   Criddle D.N., Gillies S., Baumgartner-Wilson H.K.*, et al.* Menadione-induced

reactive oxygen species generation via redox cycling promotes apoptosis of

murine pancreatic acinar cells [J]. J Biol Chem, 2006, 281(52): 40485-40492.

[70]   Ferreira G.F., Baltazar Lde M., Santos J.R.*, et al.* The role of oxidative and

nitrosative bursts caused by azoles and amphotericin B against the fungal

pathogen Cryptococcus gattii [J]. J Antimicrob Chemother, 2013, 68(8):

1801-1811.

[71]   Caceres I., Khoury A.A., Khoury R.E.*, et al.* Aflatoxin Biosynthesis and

Genetic Regulation: A Review [J]. Toxins (Basel), 2020, 12(3).

[72]   Sun Q., Shang B., Wang L.*, et al.* Cinnamaldehyde inhibits fungal growth and

aflatoxin B1 biosynthesis by modulating the oxidative stress response of

Aspergillus flavus [J]. Appl Microbiol Biotechnol, 2016, 100(3): 1355-1364.

[73]   Caceres I., El Khoury R., Bailly S.*, et al.* Piperine inhibits aflatoxin B1

production in Aspergillus flavus by modulating fungal oxidative stress

response [J]. Fungal Genet Biol, 2017, 107: 77-85.

[74]   He Z.M., Price M.S., Obrian G.R.*, et al.* Improved protocols for functional

analysis in the pathogenic fungus Aspergillus flavus [J]. BMC Microbiol, 2007,

7: 104.

[75]   Hong S.Y., Roze L.V., Wee J.*, et al.* Evidence that a transcription factor

regulatory network coordinates oxidative stress response and secondary

metabolism in aspergilli [J]. Microbiologyopen, 2013, 2(1): 144-160.

[76]   Joshi A.S., Cohen S. Lipid Droplet and Peroxisome Biogenesis: Do They Go

800   Hand-in-Hand? [J]. Frontiers in Cell and Developmental Biology, 2019, 7.

Supplementary figure

Figure S1. The construction of mutant strains. (A) PCR verification of gDNA in WT,

ΔsntB and Com-sntB strains ("ORF" represents the sntB gene fragment, "AP"

represents the amplification of the fusion fragment upstream with primers sntB-p1

and P801, and "BP" represents the downstream of the fusion fragment from primers

P1020 and sntB-p4). (B) Southern blot analysis result of WT, ΔsntB candidate strains

(The genome DNA of each strain was digested with restrictive endonuclease EcoR I

and hybridized with the probe of 1533 bp). (C) qRT-PCR verification of the

expression level of the sntB gene in WT and sntB gene mutant strains.

Figure S2. The expression of genes related to sporulation, sclerotia production, and

aflatoxin synthesis**.** (A) The expression of sporulation-related genes *steA*, *WetA*,

fluG and *veA* in each strain at 48 h. (B) The Expression of sclerotia-associated genes

nsdC, *nsdD*, and *sclR* in each strain at 48 h. (C) The Expression of

aflatoxin-associated genes in each strain at 48 h. The asterisk *** above the bars

represents significantly different (p<0.001)

Figure S3. The changes of number of conidia, amylase, and lipase in WT, ΔsntB and

Com-sntB strains. (A) Statistics of the number of conidia on the corn seed. (B) The
phenotype of each strain on starch screening medium supplemented with 0.1% of

soluble starch in darkness at 29℃ for 3 d, followed by addition of iodine solution. (C)

HC value of the clear circle (Outer diameter / inner diameter) of the salient analysis of
the mapThe asterisk ** above the bars represents significantly different (p<0.01).

Figure S4. Sequence information of ChIP-seq. (A) Heatmap. (B) PCA. (C) Peak

distribution on the genome of SNTB-HA group. (D) Peak distribution on the genome

of WT group.

Figure S5. PCR verification of gDNA in WT and ΔcatC

Figure S6. Heatmap of the DEGs related to oxidative response in transcriptome data

draw by TBtools.

Supplementary table captions

Table S1. Sequencing data statistics in RNA-seq.

Table S2. Alignment results of each sample in RNA-seq.

Table S3. The information of DEGs in transcriptome data.

Table S4. GO enriched in the DEGs in transcriptome data.

Table S5. KEGG pathways enriched of the DEGs in transcriptome data.

Table S6. Sequencing data statistics in ChIP-seq.

Table S7. Alignment results of each sample in ChIP-seq.

Table S8. The information of up regulated peak in ChIP-seq data.

Table S9. GO enriched in the up regulated genes in ChIP-seq data.

Table S10. KEGG enriched in the up regulated genes in ChIP-seq data.

Table S11. GO enriched in the 238 common DEGs in RNA-seq and ChIP-seq data.

Table S12. KEGG enriched in the 238 common DEGs in RNA-seq and ChIP-seq data.

Table S13. The information of DEGs related to lipase activity in transcriptome data.

Table S14. The information of DEGs related to oxidative response in transcriptome

data.