**Distinct roles of H3K27me3 and H3K36me3 in vernalization response, maintenance and resetting in winter wheat**

Xuemei Liu[1,2,#], Xuelei Lin[1,#,*], Min Deng[1,2,#], Bingxin Shi[1], Jinchao Chen[1,2], Haoran Li[1], Shujuan Xu[3,4], Xiaomin Bie[5], Xiansheng Zhang[5], Kang Chong[2,3], Jun Xiao[1,2,6,*]

[1] Key Laboratory of Plant Cell and Chromosome Engineering, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing 100101, China

[2] University of Chinese Academy of Sciences, Beijing 100049, China

[3] Key Laboratory of Plant Molecular Physiology, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China

[4] John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK

[5] National Key Laboratory of Wheat Improvement, College of Life Sciences, Shandong Agricultural University, Tai'an 271018, China

[6] Centre of Excellence for Plant and Microbial Science (CEPAMS), JIC-CAS, Beijing, 100101, China

[#]Equal contributions: Xuemei Liu, Xuelei Lin, Min Deng

[*]Correspondence: linxuelei@genetics.ac.cn; jxiao@genetics.ac.cn

## Author Contributions

J.X. designed and supervised the research, J.X., X.M.L and X.L.L wrote the manuscript. X.L.L. and M.D. did the sample collection, RT-PCR, CUT&Tag, ATAC-seq and RNA-seq experiments; M.D. and J.C.C. performed plasmid construction; X.L.L did in situ hybridization; H.R.L did western blot; X.M.B and X.S.Z generated *Tasdg8* mutants; B.X.S and J.C.C performed the genotyping of crispr/cas9 mutants; X.M.L performed all the data analysis. X.M.L, X.L.L. and J.X. prepared all the figures. S.J.X. and K.C. suggested and polished the manuscript. All authors discussed the results and commented on the manuscript.

**Competing Interest Statement**:

The authors declare no competing interests.

**Classification**: Biological Sciences, Plant Biology

**Keywords**: vernalization, H3K27me3, H3K36me3, winter wheat, resetting, maintenance

## Abstract

Winter plants rely on vernalization, a vital process for adapting to cold and ensuring

successful reproduction. However, understanding the role of histone modifications in guiding the vernalization process in winter wheat is limited. In this study, we investigate the transcriptome and chromatin dynamics in the shoot apex throughout the life cycle of winter wheat in the field. Two core histone modifications, H3K27me3 and H3K36me3, exhibit opposite pattern on the key vernalization gene *VERNALIZATION1* (*VRN1*), correlated with its induction during cold exposure. Additionally, H3K36me3 remains high at *VRN1* after cold exposure, maintaining its active state. Mutations in FERTILIZATION-INDEPENDENT ENDOSPERM (TaFIE) and SET DOMAIN GROUP 8 (TaSDG8), writer complex components of H3K27me3 and H3K36me3, respectively, affect flowering time. Interestingly, *VRN1* loses its high expression after cold exposure memory in the absence of H3K36me3. During embryo development, *VRN1* is silenced with the removal of H3K36me3 in both winter and spring alleles. H3K27me3 is selectively added to the winter allele, influencing the cold exposure requirement for the next generation. Integrating gene expression with H3K27me3 and H3K36me3 patterns identified potential regulators of flowering. This study reveals distinct roles of H3K27me3 and H3K36me3 in controlling vernalization response, maintenance, and resetting in winter wheat.

## Significance Statement

Vernalization, initially observed in cereals, lacks a comprehensive understanding of its underlying mechanism, particularly regarding chromatin-mediated transcriptional regulation in winter wheat. By delving into the transcriptome and chromatin dynamics in the shoot apex throughout winter wheat's life cycle, we pinpointed two crucial histone modifications, H3K27me3 and H3K36me3, each playing distinct roles at different vernalization stages. H3K27me3 is implicated in establishing and resetting the extended cold exposure requirement for winter wheat, gradually diminishing during vernalization. On the other hand, H3K36me3 is crucial for maintaining *VRN1*'s active state post-cold exposure, contributing to the memory of the vernalization treatment. Additionally, the integration of transcriptome and histone modification profiles unveiled potential novel regulators of flowering in winter wheat.

## Introduction

Winter wheat undergoes vernalization, a process involving prolonged exposure to low temperatures, to shift from vegetative to reproductive growth and enhance grain production (1). This crucial process ensures proper flowering timing and maximizes grain yields (2–5). The regulation of vernalization in winter wheat is intricate, involving a network of regulatory genes, with *VRN1*, *VRN2*, and *VRN3* being extensively studied (6–8). The interactions among these genes play a central role in controlling the vernalization response and requirement (9, 10). The activation of *VRN1* is pivotal for initiating the floral transition,

and the polymorphism at the *VRN1* locus determines the need and duration of cold exposure (11–14). Research has revealed that *VRN1* expression is subject to regulation at both transcriptional (15, 16) and post-transcriptional levels (17), as well as through epigenetic modifications (17–20).

Chilling temperatures have been demonstrated to induce histone modifications that facilitate the activation of *VRN1*. Specifically, the removal of tri-methylation from lysine 27 on histone H3 (H3K27me3) at the *VRN1* locus enables its activation. Additionally, histone marks associated with active transcription, such as H3K4me3, increase at the promoter and first intron regions of *VRN1*, promoting gene expression (17, 18). Beyond H3K27me3 and H3K4me3, H3K36me3 plays a crucial role in reversing H3K27me3 at the FLOWERING LOCUS C (*FLC*) locus in *Arabidopsis* during the vernalization process (21, 22). After cold exposure, the transcriptional status of key genes is maintained mitotically until spring when the flowering transition occurs (2, 23). In *Arabidopsis*, the silenced status of *FLC* is preserved through the spreading of H3K27me3 across the entire gene body after returning to warm temperatures (24). In barley and wheat, winter-activated *VRN1* expression remains high post-cold exposure (4), but the connection to active histone modifications and the specific histone modification involved is yet to be studied.

Crucially, the epigenetic status of key regulators in flowering regulation in response to vernalization must be reset to maintain the vernalization requirement for the next generation (25–27). In *Arabidopsis*, the coupling of H3K27me3 to *FLC* silence persists in both male and female gametes after vernalization (28, 29). The repressed state of *FLC* is maternally transmitted to early embryos, where the reactivation of *FLC* occurs with the function of the pioneer transcription factor (TF) *LEAFY COTYLEDON 1* (*LEC1*) and other B3-domain TFs, including *LEC2*, *FUSCA3* (*FUS3*), and *ABSCISIC ACID INSENSITIVE* (*ABI3*) (30–32). Notably, active demethylation of H3K27me3 by histone demethylase EARLY FLOWERING 6 (ELF6) is also necessary during this process (25). For wheat, the timing of when active *VRN1* is reset and the factors involved remain largely unknown.

In addition to *VRN1*, other candidates are proposed to play a role in the regulation of the vernalization response in wheat, based on their expression patterns. These include potential *FLC* orthologs *ODDSOC2* (*TaOS2*) and *TaOS1*, although their impact on flowering time appears to be marginal, as indicated by genetic data from mutations (33, 34). Furthermore, a global analysis of the H3K27me3 and H3K4me3 patterns in *Brachypodium* at different vernalization stages (V0, V30, V30N) has identified additional loci involved in mediating the epigenetic memory of flowering competency acquired during the vernalization process (35). Thus, both transcriptional and histone modification patterns

may serve as indicators for the identification of potential regulators for vernalization-promoted flowering.

In this study, we delved into the role of the chromatin landscape in driving the vernalization response, memory, and resetting in winter wheat by profiling chromatin accessibility, core histone modifications, and the histone variant H2A.Z in the shoot apex throughout an entire growth season in the field. We observed antagonistic modifications of H3K27me3 and H3K36me3 at the *VRN1* locus during the entire life cycle of winter wheat. Through assessing flowering time, *VRN1* transcription levels, and relative histone modifications in genome-edited Polycomb Repressive Complexes 2 (PRC2) and SDG8 mutants, we unveiled distinct functions of H3K27me3 and H3K36me3 in mediating the vernalization response, memory, and resetting processes. Moreover, integrating transcriptome patterns with the profiles of H3K27me3 and H3K36me3 allowed the identification of additional *VRN1*- and *FLC*-pattern-like genes for potential regulation of flowering in wheat.

## Results

**Profiling the transcriptome and chromatin landscape in the shoot apex across the life-cycle of winter wheat in field**

Winter wheat is traditionally sown in early October and experiences a prolonged cold winter from November to early March, with temperatures ranging from -10°C to 5°C in Beijing, China (Fig. 1A). As temperatures rise above 10°C in mid-March, wheat jointing stage occurs. We conducted a comprehensive study on the transcriptome and chromatin dynamics during the vernalization response, maintenance, and resetting stages of winter wheat cultivar Kenong 9204 (KN9204), using samples of shoot apex and developing seed/embryo from field-grown wheat (Fig. 1A). Our efforts resulted in the generation of 21 RNA-seq libraries, 14 ATAC-seq libraries, and 94 CUT&Tag libraries, targeting various histone modifications and variant (H3K27ac, H3K27me3, H3K36me3, H3K4me3, and H2A.Z) (Fig. 1B). Quality assessments revealed excellent reproducibility for both transcriptome and epigenome data (Fig. S1A, D, E), and the distribution of histone modifications and chromatin accessibility profiles across the genome and genes aligned with their known roles in transcriptional regulation, as previously reported (36, 37) (Fig. S1B, C).

Differential expression analysis was conducted for S0 (pre-vernalization), S1 and S2 (during vernalization), and S3 and S4 (fully vernalized), revealing 19,370 differentially expressed genes (DEGs) (Table S1). Subsequent K-means clustering classified DEGs into five clusters with distinct expression patterns (Fig. 1C). Notably, genes in cluster C5 exhibited high expression before cold exposure, while genes in clusters C1 and C3 were

highly induced during cold treatment, and those in clusters C4 and C2 were predominantly expressed in the fully vernalized state. Gene Ontology (GO) enrichment analysis shed light on the functions of stage-specific highly expressed genes (Fig. 1D). Before vernalization, wheat developments fast, genes were associated with chromatin assembly, plant organ formation, hormone biosynthetic, and response to light intensity (C5). During vernalization, winter wheat initially acquired cold tolerance, with genes related to fluid transport, pentose metabolic, carbohydrate phosphorylation, oxidative photosynthetic carbon pathway are highly expressed in the S1 stage (C1), and cold acclimation genes upregulated in the S2 stage (C3). Notably, key cold-responsive modules, including the *INDUCER OF CBF EXPRESSION-C REPEAT BINDING FACTOR-COLD RESPONSIVE* (*ICE-CBF-COR*) cluster (38), were highly expressed at the S2 stage (Fig. 1E), indicating early cold tolerance acquisition. In the vernalization maturation stage (39), genes related to amino acid catabolic, protein maturation, protein complex oligomerization, and amino acid transport were highly expressed (C4, C2), including the crucial flowering time regulator, *VRN1* (6) (Fig. 1E). While both *VRN1* and *TaCOR14/15* responded to low temperatures, their expression patterns differed (Fig. 1E). *TaCOR14/15* exhibited high expression levels only during early cold exposure, followed by a decline, whereas *VRN1*'s expression increased gradually with prolonged cold exposure until the vernalization effect was fully established (Fig. 1E). These distinct expression patterns corresponded to their functions in cold tolerance and the measurement of cold duration for flowering promotion (3). In summary, the transcriptome effectively captures the various stages of vernalization in winter wheat, highlighting the differentiation between short-duration cold tolerance genes and long-duration vernalization genes.

**Dynamics of chromatin states during vernalization establishment and maintenance**
We utilized chromHMM (40) to explore the dynamic alterations in chromatin states throughout vernalization, integrating multiple histone modifications and variant data. Models with 5 to 25 states were trained, and 11 chromatin states were selected based on their similarities (Fig. S2A). Each chromatin state was assigned a descriptive label according to enriched histone modifications, distance to genes, and expression levels of nearby genes (Fig. 2A). These 11 states were grouped into four based on transcriptional activity (Fig. 2A). Group G1, marked by high gene expression, included states E1, E2, E3, and E4, characterized by the presence of H3K36me3, associated with active transcription (Fig. S2B). In contrast, G3, featuring low gene expression, comprised states E8, E9, and E10, enriched in H3K27me3, a repressive modification inversely correlated with transcription (Fig. S2B). E8 displayed a bivalent chromatin state (41), showing both H3K27me3 and active histone marks H3K4me3 and H3K27ac. E4, rich in H3K27ac, H3K4me3, and H2A.Z, exhibited the highest chromatin accessibility, while E1, E2, and E3,

with high transcription levels, displayed lower accessibility (Fig. 2A), aligning with the idea that transcriptional regulatory regions are generally more accessible than gene bodies (42).

Exploring the transition of chromatin states during vernalization establishment (S0-S4) revealed more significant changes in G2 and G3 compared to G1 and G4 (Fig. 2B). E3, E5, and E9 exhibited the most substantial changes (Fig. 2B), with E9 in the fully vernalized S4 stage displaying a distinct profile (Fig. 2C). Most regions in E9 state at S0, S1, S2, S3 remained in E9 state at S4 (Fig. S2C). However, genes with chromatin state changed at S4, especially transitioning into E1 or E2 state, were mainly associated with flowering development, auxin signaling, and flower organ specification (Fig. S2D), indicating a crucial role for H3K27me3 in vernalization establishment. After extended exposure to winter cold, wheat plants initiated bolting in late March, relying on the 'memory' of the vernalized state (3). Notably, E9 state remained more stable from S4 to S5 and S6 (Fig. 2B, C), suggesting that H3K27me3 may not be involved in maintaining vernalization memory.

Further examination of the core regulator, VRN1, revealed quantitative induction in response to cold exposure from S0 to S4, maintaining high expression in S6 after cold treatment in the shoot apex (Fig. 2D). This pattern aligns with its role in promoting flowering and regulating spikelet development (6, 43). Before vernalization, *VRN1* exhibited minimal expression, with its promoter and first intron (*TraesCS5A02G391700*) covered by H3K27me3 within chromatin state E9 (Fig. 2E). Continuous exposure to low temperatures led to a gradual decrease in H3K27me3 levels at *VRN1*, accompanied by a progressive increase of H3K36me3 levels, resulting in a shift from E9 to E2 and E4 chromatin states and continuous upregulation of transcription (Fig. 2E, Fig. S2E, F). A similar inverse pattern of H3K27me3 and H3K36me3 was observed in the core vernalization gene *FLC* in *Arabidopsis thaliana* (22). Moreover, upon return to normal temperatures after vernalization, H3K27me3 enveloped the entire *FLC* locus to inhibit transcription in *Arabidopsis* (21). Similarly, in wheat, upon transitioning to normal temperatures (S5 and S6 stages) after vernalization, the presence of H3K36me3 coincided with high *VRN1* expression (Fig. 2E, Fig. S2E, F). These findings highlight the changing chromatin states during vernalization in winter wheat, emphasizing the importance of H3K27me3 and H3K36me3 in regulating key flowering genes like *VRN1*, with the conserved roles across various plant species.

**Modulation of H3K27me3 or H3K36me3 affects vernalization-induced flowering in winter wheat**

Giving the potential pivotal role of H3K27me3 and H3K36me3 in orchestrating the
vernalization response, we conducted a genetic analysis to investigate their influence on
flowering by manipulating these histone modifications. Through CRISPR-Cas9 genome
editing, we generated mutations targeting the "writer" complex responsible for H3K27me3
and H3K36me3 in the winter wheat KN199 (Fig. S3A, B), a derived cultivar from KN9204
with moderate genetic transformation efficiency (44). Knocking out all three subgenomes
of *TaFIE*, a core component of the PRC2 (45), proved embryonic lethal (37). Therefore, we
opted for a mild mutation, the *Tafie-cr-87* line, which exhibited a frameshift mutation in the
A and D subgenomes and two amino acid deletions in the B subgenome of *TaFIE* (Fig.
S3A) (37). Notably, the global H3K27me3 level was reduced in *Tafie-cr-87* compared to
KN199 at shoot apex (Fig. S3C).
239
We assessed the flowering times of *Tafie-cr-87* and KN199 under various vernalization
treatments, including 0 day (V0), 20 days (V20), and 30 days (V30) of exposure to 4°C.
Under V0 and V20 conditions, *Tafie-cr-87* displayed earlier flowering compared to KN199,
while no differences were observed under V30 conditions with saturated cold exposure
(Fig. 3A, B). Subsequently, we examined the expression levels of *VRN1* in *Tafie-cr-87* and
KN199 under different vernalization treatments. *VRN1* expression was significantly
elevated in *Tafie-cr-87* compared to KN199 at the seedling stage under V0 and V20
conditions, while no differences were observed under V30 conditions (Fig. 3C, Fig. S3E).
Additionally, H3K27me3 levels were reduced at the *VRN1* promoter region in *Tafie-cr-87*
compared to KN199 under V0 and V20 conditions, but H3K27me3 diminished in both
plants under V30 conditions (Fig. 3D, Fig. S3F). These findings suggest that PRC2-
H3K27me3 primarily inhibits *VRN1* expression without or under insufficient vernalization
treatment, with the repressive effect diminishing with prolonged cold exposure.
253
In *Arabidopsis*, SDG8 functions as a methyltransferase for H3K36me3 in regulating
flowering (46). In wheat, three homologues of *SDG8* form a triads, with A-, B- and D-
subgenome *TaSDG8* exhibiting high expression in the shoot apex (Fig. S3D). To
investigate the role of H3K36me3 in vernalization-induced flowering, we generated two
*TaSDG8-a/b/d* mutant lines using CRISPR-Cas9 with reduced global H3K36me3 level (Fig.
S3B, C), and evaluated their flowering time with vernalization treatment. Both *Tasdg8-cr-3*
and *Tasdg8-cr-5* lines exhibited delayed flowering (Fig. 3E, F). Assessment of *VRN1*
expression levels (Fig. 3G, Fig. S3G) and H3K36me3 status (Fig. 3H, Fig. S3H) in KN199
and *Tasdg8-cr-3/5* lines under V30 and V30+10 (exposure to 4°C for 30 days and convert
to normal temperature for 10 days) conditions revealed that *VRN1* levels decreased along
with reduced H3K36me3 levels in *Tasdg8-cr-3/5* lines under V30+10 compared to V30
conditions. In contrast, increased expression and H3K36me3 level was observed in KN199

under these conditions (Fig. 3G, H). Thus, the accumulation of H3K36me3 in *VRN1*,
induced by prolonged cold exposure, contributes to the memory of vernalization treatment
post-cold exposure in winter wheat.
269
**Resetting H3K27me3 and H3K36me3 at *VRN1* during embryogenesis to restore**
**vernalization requirement in winter wheat**
Following flowering, the vernalization memory must be wiped clean in the offspring to
prevent the inheritance of the vernalized state across generations (25, 27). In *Arabidopsis*,
the resetting of the vernalization requirement takes place during embryogenesis and
involves the reactivation of *FLC* (25, 30). This process is associated with the reset of
epigenetic modifications, such as H3K27me3 at the *FLC* locus (27). Given that VRN1 is
the pivotal locus determining the vernalization requirement in wheat, we examined the
expression pattern of *VRN1* during embryogenesis using in situ hybridization in winter
wheat KN9204 (Fig. 4A). Prior to fertilization, we observed a robust signal of *VRN1*
transcript in the embryo sac. However, *VRN1* expression gradually decreased with
increasing days after pollination (DAPs), becoming barely detectable at DAP6 in the
developing embryo and entirely undetectable at DAP14 (Fig. 4A). This indicates that *VRN1*
expression is repressed during embryogenesis, mirroring the resetting process of *FLC* in
*Arabidopsis* but with an opposite trend.
285
To delve deeper into the resetting of the vernalization "memory" in wheat, we utilized
developing seeds at DAP0, DAP2, and DAP6 to conduct CUT&Tag analysis for four histone
modifications (H3K27ac, H3K27me3, H3K36me3, H3K4me3) (Fig. 1A, B). Consistent with
*in situ* hybridization findings, we validated a transcriptional decline of *VRN1* during
embryogenesis (Fig. 4B). Particularly noteworthy was the shift in the chromatin state at the
*VRN1-5A* locus from an active to a repressed state during embryogenesis (Fig. 4C).
Notably, H3K36me3 levels significantly decreased at proximal promoter and within the
*VRN1-5A* gene body, especially in the first intron region, from DAP0 to DAP6. Conversely,
the repressive histone mark H3K27me3 gradually increased, encompassing the entire
genic region, proximal promoter, and 3'UTR regions of *VRN1-5A* at DAP6 (Fig. 4C).
Though there are various among different subgenomes of *VRN1* triads, the general pattern
is similar (Fig. S4A). This underscores the significance of the reciprocal patterns of
H3K27me3 and H3K36me3 in the transcriptional resetting of *VRN1* during embryogenesis.
Additionally, we identified a similar pattern of reduced *VRN1* expression in spring wheat
Chinese Spring (CS) for both the recessive *vrn1-5A vrn1-5B* and dominant *Vrn-5D* alleles
from DAP0 to DAP8 (Fig. 4D) (37). Coinciding with reduced transcription, H3K36me3
levels declined in all three sub-genomes (Fig. 4E, Fig. S4B). However, we did not observe
the accumulation of H3K27me3 during embryogenesis at DAP8 (Fig. 4E, Fig. S4B). This

difference is likely linked to the variation that *VRN1* can be detected at the seedling stage, especially three-leaf stage, in CS but merely in KN9204 (Fig. S4C).

To ascertain whether the restoration of H3K27me3 at the *VRN1* locus during embryogenesis contributes to the repression of *VRN1* prior to vernalization, we compared mRNA levels and H3K27me3 statuses at the *VRN1* locus in KN199 and *Tafie-cr-87* during embryogenesis (DAP2, DAP8, DAP16) and the three-leaf seedling stages before cold exposure. Both KN199 and *Tafie-cr-87* exhibited a decline in *VRN1* mRNA levels during embryogenesis, with slightly or no significant difference between the two genotypes at DAP16 (Fig. 4F, Fig. S4D, F). However, H3K27me3 accumulated in KN199 but remained less enriched in *Tafie-cr-87* at 8 and DAP16 (Fig. 4G, Fig. S4E, G). At the three-leaf seedling stage, *Tafie-cr-87* displayed higher *VRN1* expression levels than KN199 (Fig. 4H, Fig. S4H, J), accompanied by lower H3K27me3 levels (Fig. 4I, Fig. S4I, K). In conclusion, the resetting of H3K27me3 and H3K36me3 at the *VRN1* locus during embryogenesis plays a critical role in maintaining the repression of *VRN1* prior to cold exposure, particularly the accumulation of H3K27me3 during embryogenesis.

**Discovery of novel regulators for vernalization response in winter wheat**
Drawing insights from the transcriptional profiles of *VRN1* and *FLC* throughout the life cycle of wheat and *Arabidopsis* (27) and considering the reciprocal regulation by H3K27me3 and H3K36me3 in vernalization response, maintenance, and resetting (21, 22), we hypothesized that genes exhibiting analogous patterns of transcriptional and chromatin state changes might also play a role in vernalization-regulated flowering.

The transcriptional level of *VRN1* steadily increases during vernalization (from S0 to S4) and remains elevated post-vernalization (from S5 to S6) (Fig. 2E). Consequently, we conducted an exploration for *VRN1*-pattern-like genes using transcriptome, H3K27me3, and H3K36me3 data spanning S0 to S6. This effort resulted in the identification of 212 *VRN1*-pattern-like genes, discerned from the DEGs of C2 and C4 via K-means clustering (Table S4). Similar to *VRN1*, these genes exhibit consistent upregulation during vernalization and maintain heightened transcriptional activity after vernalization (Fig. 5A), all while being subject to regulation by both H3K27me3 and H3K36me3 (Fig. 5B). GO enrichment analysis reveals their involvement in flower development, signal transduction, auxin signaling, and shoot system development (Fig. 5C). Notably, the transcription factors within the *VRN1*-pattern-like genes are enriched in the MIKC_MADS family (Fig. 5D), particularly in the same subclade as *VRN1* (Fig. S5, Table S5). Among these genes, 93 harbor SNPs in their promoters or gene bodies significantly associated with heading time (Table S6), and 4 genes with severe TILLING mutations (Wang et al., 2023) (Table S7)

exhibit aberrant flowering (Fig. 5E). These include the known flowering regulators TaFUL3
(43) and TaSOC1 (47), which display strikingly similar transcriptional and epigenetic
profiles to *VRN1* (Fig. 5F). Notably, seven SNPs (red line in Fig. 5F) associated with the
heading time are located in the promoter and gene body of *TaFUL3_2A* (Fig. 5G, Table
S8).

FLC, a flowering suppressor, plays a central role in mediating vernalization-regulated
flowering in *Arabidopsis* (48). Consequently, efforts in previous research aimed to pinpoint
its homologous gene in wheat (34, 49). We also found that these genes show high
sequence similarity to *FLC* (highlighted in green) (Fig. S6), and the transcriptional inhibition
of these genes by vernalization (Fig. S5, Table S5). However, the H3K27me3 and
H3K36me3 level are not aligned with the transcription during vernalization (Fig. 6A).

Giving the importance of H3K27me3 and H3K36me3 in shaping *FLC* transcription pattern
(21, 22), we embarked on a quest to identify *FLC*-pattern-like genes in other gene families
using transcriptome, H3K27me3, and H3K36me3 data. Through this exploration, we
discovered 585 *FLC*-pattern-like genes within DEGs of C5 through K-means clustering
(Table S4). These genes undergo transcriptional inhibition by vernalization (Fig. 6B) and
are associated with increased H3K27me3 and decreased H3K36me3 (Fig. 6C). They are
linked to various biological processes, including steroid metabolic processes, toxin
biosynthesis, seedling development, and hormone biosynthesis (Fig. 6D). Within this set,
249 genes (Table S6) harbor SNPs significantly associated with flowering time, while 13
genes (Table S7) with severe TILLING mutations can exert an impact on wheat flowering
time (Fig. 6E). Furthermore, *TaTOE1* has been previously reported to influence wheat
flowering (50) and we identified eight SNPs associated with heading time within its
promoter and genebody (Fig. 6F, G, Table S9).

In sum, through a thorough examination of transcriptional and epigenetic feature
similarities, combined with available genotype and phenotype at population level and
TILLING mutant information, we have identified potential novel regulators governing
vernalization-regulated flowering in winter wheat.

**Discussion**
Plants respond to seasonal temperature changes, notably extended winter cold,
orchestrating crucial developmental shifts such as flowering (1). Unlike in *Arabidopsis*,
where cold silences the flowering repressor *FLC*, cereals like wheat activate the flowering
promoter *VRN1* during winter (3, 51). Despite their opposite roles in flowering regulation,
our in-depth analysis of transcriptome and epigenetic profiles revealed a conserved

epigenetic regulatory model for mediating *VRN1* transcriptional dynamic in winter wheat, akin to *FLC* regulation in *Arabidopsis*.

*FLC*, active pre-cold, is gradually silenced, kept low post-warm return (52, 53). It remains low throughout gametophyte generation, reactivates during embryogenesis, and stays high during seedling development, resetting the vernalization requirement for the next generation (25–27). In our study, transcriptome analysis of the shoot apex during the life cycle of winter wheat KN9204 indicated that *VRN1* is initially inactive in seedlings before cold treatment. During the cold winter, *VRN1* gradually activates and remains high in the following spring in the inflorescence tissue (Fig. 2). Subsequently, *VRN1* is re-silenced during embryogenesis to reset the vernalization requirement (Fig. 4).

In *Arabidopsis*, *FLC*'s chromatin status correlates with its transcription. Pre-cold, H3K4me3 and H3K36me3 mark *FLC*, shifting to H3K27me3 at the nucleation site during cold, spreading across the entire gene body post-warm return, serving as a memory of the vernalization treatment. Similarly, we observed a conserved yet mirrored pattern of H3K27me3 and H3K36me3 dynamics at the *VRN1* locus in winter wheat (Fig. 2). H3K27me3 decreases, H3K36me3 but not H3K4me3 accumulates during cold, persisting post-warm return (Fig. 2). *Tasdg8-cr* lines fail to sustain active *VRN1* expression post-vernalization, resulting in delayed flowering (Fig. 3). During embryogenesis, H3K36me3 reduction correlates with *VRN1* transcription drop in both winter and spring wheat, while H3K27me3 accumulates selectively in winter wheat, crucial for inhibiting unlicensed *VRN1* transcription pre-cold in *Tafie-cr-87* (Fig. 4). It would be intriguing to unravel the mechanisms underlying the dynamic patterns of H3K27me3 and H3K36me3 at the *VRN1* locus throughout the life cycle of winter wheat. Furthermore, exploring the association between the dynamics of the epigenetic landscape and transcriptional patterns across different winter and spring *VRN1* alleles presents a compelling avenue for future analysis.

Giving *FLC*'s crucial role in vernalization responses across species, substantial efforts have been made to find *FLC* counterparts in monocots (34, 54). Based on sequence similarity, TaOS2/TaAGL33 has been identified as a potential floral repressor in wheat, exhibiting a response to cold exposure, albeit with a less pronounced early flowering phenotype (33, 55). Similarly, in barley and *Brachypodium*, BdOS2 and HvOS2 also function as floral repressors (33, 56). Our analysis, however, didn't reveal a correlated H3K27me3 and H3K36me3 pattern at the TaOS2 locus during development (Fig. 6), suggesting it might not be an epigenetic regulation target for vernalization. Recognizing the mitotic memory in vernalization treatment, we suggest that epigenetic modification patterns could be vital markers for identifying genes similar to *FLC* and *VRN1*. By

418    integrating expression patterns and H3K27me3/H3K36me3 profiles during vernalization
419    (Fig. 5,6), we identified potential factors like TaFUL3 and TaTOE1, associated with
420    flowering and showing natural variations linked to heading time diversity in extensive wheat
421    populations, meriting further analysis.
422
423    As global warming alters temperature conditions (57), impacting the growth areas for wheat
424    varieties, it becomes crucial to understand and adapt vernalization responses to ensure
425    optimal grain yield. Fine-tuning the major vernalization response locus *VRN1*, along with
426    the identification of additional genetic loci controlling the vernalization response, could
427    enhance wheat's resilience to environmental change.
428

## Materials and Methods

### Growth conditions, vernalization treatment and experimental materials

431    The winter wheat cultivar KN9204 used in this study were grown during the normal growing
432    seasons at the Experimental Station of Institute of Genetics and Developmental Biology,
433    Chinese Academy of Sciences, Beijing, China at the beginning of October each year. They
434    experienced natural vernalization during the winter and flowered the following spring.
435

The stage-specific apexes of wheat (S0-S6) were dissected under the stereomicroscope
based on the anatomic and morphological features at the right time (Fig. 1A), and
immediately frozen in liquid nitrogen and stored at -80 °C. About 10 to 50 spikes were
pooled. The stamens were removed before the pollen maturation. Then, we conducted
artificial pollination and recorded the number of days to ensure the accurate time of seed
development. The seeds at DAP0, DAP2, DAP6, DAP8, and DAP16 stage were sampled
for later use of total RNA extraction and nuclei isolation. The dissection method of embryo
and embryo sac follow the instructions in the previous article (37). Embryo sacs containing
egg/embryo and surrounding tissue in DAP0 and DAP2 were dissected in a 5% Sucrose
solution containing 0.1% RNase inhibitor with fine forceps using the dissecting microscope.
For DAP6 and later stages, the independent embryo was detached from the embryo sac
using the same method. Embryos and embryo sacs sampled from eight to ten spikes were
pooled for one biological replicate in early stages and three to five were pooled for one
biological replicate in late stages. The RNA-seq (three replicates), ATAC-seq and
CUT&Tag (two replicates) experiments at twelve development stages were carried out.
451

For artificial vernalization treatment, seeds of winter wheat and spring wheat (*Triticum*
*aestivum* cv. KN9204, KN199 and CS), seeds of *TaFIE* or *TaSDG8* transgenic lines were
surface sterilized in 2% NaClO for 20 minutes, then rinsed overnight with flowing water.
The seeds were germinated and then transferred to soil, and grown at 4°C in the dark for

0, 20, 30 days (V0, V20, V30) and then grown at room temperature for 10 days (V30+10).
Wheat leaves from these samples were used to extract RNA for RT–qPCR or isolate nuclei
for CUT&Tag–qPCR. Then were transferred in a greenhouse (20°C–22°C, 16-h light/8-h
dark). Finally, the heading time is counted, which is the time from sowing to heading minus
the corresponding 4°C vernalization treatment time.

**TILLING mutation library and identification of lines with heading time difference**
The previously published KN9204 TILLING mutation library (58) was used to screen
mutants with differences in heading time from wild-type KN9204. Each mutant line was
planted in a 1.5 m single-row plot with 10cm between plants and 25 cm between rows and
managed according to local practices. Heading time was calculated as the days from
sowing to half of the spikes in a row heading.

**RNA-seq, ATAC-seq and CUT&Tag experiment**
Total RNA was extracted using HiPure Plant RNA Mini Kit (Magen, R4111-02). RNA-seq
libraries' construction and sequencing platform were the same as previous description
(37), by Annoroad Gene Technology.

ATAC-seq and CUT&Tag experiments followed the previously described method (37). Tn5
transposase used and tagmentation assay is done following the manual (Vazyme, TD501-
01). Libraries were purified with AMPure beads (Beckman, A63881) and sequenced using
the Illumina Novaseq platform at Annoroad Gene Technology. Antibodies used for histone
modifications are listed in Table S10.

**RNA-seq data analysis**
Fastp (v0.23.1) (59) was used to process raw reads to remove adapters, low quality bases
and reads. Then the cleaned reads were aligned to IWGSC RefSeq v1.0 (55) using hisat2
(v2.0.5) (60). Samtools (v1.5) (61) was used to sort the aligned reads and converted it into
bam files. The gene count matrix was calculated by featureCounts (v2.0.1) (62) with
parameter "-p –P –B -C" The counts matrixs were used as inputs to identify differentially
expressed genes using R package DESeq2 (v1.28.1) (63), with a threshold of p.adj < 0.05
and abs(log2FoldChange) > 1. TPM (Transcripts Per Kilobase Million) values generated
from the counts matrix were used to characterize gene expression and used for
hierarchical clustering analysis. K-means cluster analysis of differentially expressed genes
were performed in R (v4.0.2) using Z-scaled TPM and the gene expression heatmap was
displayed using R package ComplexHeatmap (v2.4.3) (64). The gene ontology enrichment
analysis was done on using R package clusterProlifer (v3.16.1) (65) and the gene ontology
annotation was performed using PANTHER (66). The identification and family classification

of wheat TFs were predicted on the PlantTFdb website ([http://planttfdb.gao-lab.org/](http://planttfdb.gao-lab.org/)) (67). Enrichment analysis of TF families for VRN1-like genes was achieved using the enricher function in R package clusterProlifer (v3.16.1) (65).

**ATAC-seq and CUT&Tag data analysis**

Raw reads were processed by fastq (v0.23.1) with "--detect_adapter_for_pe" parameter to remove adapters, trim low quality bases and filter bad reads. Clean reads were aligned to IWGSC RefSeq v1.0 using bwa mem algorithm (v 0.7.17) (68) with "-M" parameter. The aligned reads were sorted and filtered by samtools (v1.5) with "-F 1804 -f 2 -q 30" parameter. Picard (v 2.23.3) was used to remove the duplicated reads. The de-duplicated bam files from two biological replicates were merged by samtools (v1.5) and converted into bigwig files using bamCoverage provided by deeptools (v3.5.0) (69) with parameters "-bs 10 --effectiveGenomeSize 14600000000 --normalizeUsing RPKM --smoothLength 50". The bigwig files were visualized using deeptools (v3.5.0) and IGV (v2.8.0.01) (70).

The peak calling was done using merged bam files by macs2 (v 2.2.7.1) (71). For ATAC-seq data, the parameter of peak calling using macs2 is "-q 0.05 -f BAMPE --nomodel --extsize 200 --shift -100 -g 14600000000". For narrow peak of H3K27ac, H3K4me3 and H2A.Z, the peak calling parameter is "-p 1e-3 -f BAMPE -g 14600000000 --keep-dup all". For broad peak of H3K27me3 and H3K36me3, the peak calling parameter is "-f BAMPE -g 14600000000 --keep-dup all --broad --broad-cutoff 0.05". Peaks from S0 to S6 of the same data type were merged using bedtools (v2.29.2) (72) to generate the reference peaks. Peak was annotated to the wheat genome using the R package ChIPseeker (v1.24.0) (73).

For quantification of ATAC-seq and CUT&Tag data, reads count under reference peak and the normalized CPM (DBA_SCORE_TMM_READS_EFFECTIVE_CPM) values were generated using R package DiffBind (v2.16.2). The CPM values of reference peak were used to do hierarchical clustering analysis.

**Chromatin state analysis**

ChromHMM (v1.22) (40) was used to train models containing 5 to 25 chromatin states using all CUT&Tag data. CompareModels was used to calculate correlation between different models. OverlapEnrichment was used to calculate the enrichment of each state at different locations in genome.

**Association analysis between genotype and heading time**

The genotype and heading time data from 788 wheat cultivars was obtained from the

532  published data (74). We extracted SNPs located in the promoter and genebody of VRN1-
533  pattern-like and FLC-pattern-like genes. The population was divided into two sub-
534  population by each single SNP, and t-test was performed to test the significance of the
535  difference in heading time between two sub-population. The significance threshold is set
536  to 1e-6.
537
**Phylogenetic tree of MADS family**
All the proteins in *Arabidopsis thaliana*, wheat, rice, barley, and *Brachypodium distachyon*
that were predicted by hmmsearch (v3.3.2) (75) to contain the MADS domain were
considered to be the proteins of the MADS family. The longest protein of the MADS family
gene was aligned by muscle (v3.8.31) (76), and then trimal (v1.4.rev15) (77) was used to
cut the aligned sequences. Only the sequence that existed in 80% of the genes was
retained. RAxML (v8.2.12) (78) was used to establish the phylogenetic tree of the MADS
family.
546
**Construction and genotyping of CRISPR/Cas9 lines**
The construction and identification of *Tafie-cr-87* mutants was reported in previous study
(37). For mutant *Tasdf8-cr-3/5*, plasmid constructs pJIT163-Ubi-Cas9 and Pu6-gRNA39
were used (79, 80). Gene-specific primers were designed around the gRNA target site to
identify the *Tasdg8-cr-3/5* mutants. PCR products were genotyped by Sanger sequencing.
Primers used for construction and identification are listed in Table S11.
553
**In situ hybrization**
RNA in situ hybridization was carried out as described previously (81). Fresh young spikes
and seeds were fixed in formalin-acetic acid-alcohol overnight at 4 °C, dehydrated through
a standard ethanol series, embedded in Paraplast Plus tissue-embedding medium (Sigma-
Aldrich, P3683), and sectioned at 8-10 μm width using a microtome (Leica Microsystems,
RM2235). Digoxigenin-labeled RNA probes based on the sequence of *VRN1* were
synthesized using a DIG northern Starter Kit (Roche, 11277073910), according to the
manufacturer's instructions. Primers used for probe synthesis are listed in Table S11.
562
**RT-qPCR**
First-strand cDNA was synthesized from 2 μg of DNase I-treated total RNA using the
TransScript First Strand cDNA Synthesis SuperMix Kit (TransGen, AT301-02). RT-qPCR
was performed using the ChamQ Universal SYBR qPCR Master Mix (Vazyme, Q711-02)
by QuantStudio5 (Applied biosystems). The expression of interested genes was
normalized to Tubulin for calibration, and the relative expression level is calculated via the
2-ΔΔCT analysis method (82). Primers used for RT-qPCR are listed in Table S11.

**CUT&Tag-qPCR**

CUT&Tag-qPCR was performed as previously reported with minor modifications (83). Briefly, the DNA products of CUT&Tag were divided, 6μL was used as "Input", and the other 18μL undergoing library PCR amplification and purification was used as "IP products." Finally, the "Input" and "IP products" were dilute 30 times for qPCR assay. The primers for specific regions are provided in Table S11.

**Data availability**

The omics data at S0, S2, S3, DAP0, DAP2, DAP6 stages are reported for the first time in this study and the raw sequence data have been deposited in the Genome Sequence Archive of the National Genomics Data Center, China National Center of Bioinformation/Beijing Institute of Genomics, Chinese Academy of Sciences (PRJCA022101). Other omics data were download from our previous report (84).

**Acknowledgments**

This research was supported by National Natural Science Foundation (31921005), Beijing Natural Science Foundation Outstanding Youth Project (JQ23026), National Key Research and Development Program of China (2021YFD1201500), the Major Basic Research Program of Shandong Natural Science Foundation (ZR2019ZD15).

**References**

1.   P. Chouard, Vernalization and its Relations to Dormancy. *Annual Review of Plant Physiology* **11**, 191–238 (1960).

2.   R. Amasino, Seasonal and developmental timing of flowering. *Plant J* **61**, 1001–1013 (2010).

3.   S. Xu, K. Chong, Remembering winter through vernalisation. *Nature Plants* **4**, 997–1009 (2018).

4.   A. Distelfeld, C. Li, J. Dubcovsky, Regulation of flowering in temperate cereals. *Current Opinion in Plant Biology* **12**, 178–184 (2009).

5.   I. R. Henderson, C. Shindo, C. Dean, The Need for Winter in the Switch to Flowering. *Annual Review of Genetics* **37**, 371–392 (2003).

6.   L. Yan, *et al.*, Positional cloning of the wheat vernalization gene VRN1. *Proceedings of the National Academy of Sciences* **100**, 6263–6268 (2003).

7.   L. Yan, *et al.*, The wheat VRN2 gene is a flowering repressor down-regulated by vernalization. *Science* **303**, 1640–1644 (2004).

606     8.    L. Yan, *et al.*, The wheat and barley vernalization gene VRN3 is an orthologue of FT. *Proc*
607           *Natl Acad Sci U S A* **103**, 19581–19586 (2006).

608     9.    A. Chen, J. Dubcovsky, Wheat TILLING Mutants Show That the Vernalization Gene VRN1
609           Down-Regulates the Flowering Repressor VRN2 in Leaves but Is Not Essential for
610           Flowering. *PLOS Genetics* **8**, e1003134 (2012).

611    10.    J. Xiao, *et al.*, Wheat genomic study for genetic improvement of traits in China. *Sci. China*
612           *Life Sci.* **65**, 1718–1775 (2022).

613    11.    D. Fu, *et al.*, Large deletions within the first intron in VRN-1 are associated with spring
614           growth habit in barley and wheat. *Mol Genet Genomics* **273**, 54–65 (2005).

615    12.    I. Diaz, *et al.*, The GAMYB protein from barley interacts with the DOF transcription factor
616           BPBF and activates endosperm-specific genes during seed development. *Plant J* **29**,
617           453–464 (2002).

618    13.    N. Kippes, *et al.*, Identification of the VERNALIZATION 4 gene reveals the origin of spring
619           growth habit in ancient wheats from South Asia. *Proc Natl Acad Sci U S A* **112**, E5401-
620           5410 (2015).

621    14.    A. B. Shcherban, E. A. Salina, Evolution of VRN-1 homoeologous loci in allopolyploids
622           of Triticum and their diploid precursors. *BMC Plant Biol* **17**, 188 (2017).

623    15.    Y. Chen, B. F. Carver, S. Wang, F. Zhang, L. Yan, Genetic loci associated with stem
624           elongation and winter dormancy release in wheat. *Theor Appl Genet* **118**, 881–889
625           (2009).

626    16.    G. Li, *et al.*, Vernalization requirement duration in winter wheat is controlled by TaVRN-
627           A1 at the protein level. *Plant J* **76**, 742–753 (2013).

628    17.    J. Xiao, *et al.*, O-GlcNAc-mediated interaction between VER2 and TaGRP2 elicits
629           TaVRN1 mRNA accumulation during vernalization in winter wheat. *Nat Commun* **5**, 4572
630           (2014).

631    18.    S. N. Oliver, E. J. Finnegan, E. S. Dennis, W. J. Peacock, B. Trevaskis, Vernalization-
632           induced flowering in cereals is associated with changes in histone methylation at the
633           VERNALIZATION1 gene. *Proc Natl Acad Sci U S A* **106**, 8386–8391 (2009).

634    19.    S. N. Oliver, W. Deng, M. C. Casao, B. Trevaskis, Low temperatures induce rapid changes
635           in chromatin state and transcript levels of the cereal VERNALIZATION1 gene. *J Exp Bot*
636           **64**, 2413–2422 (2013).

637    20.    S. Xu, *et al.*, The vernalization-induced long non-coding RNA VAS functions with the
638           transcription factor TaRF2b to promote TaVRN1 expression for flowering in hexaploid
639           wheat. *Mol Plant* **14**, 1525–1538 (2021).

640    21.    H. Yang, *et al.*, Distinct phases of Polycomb silencing to hold epigenetic memory of cold
641           in Arabidopsis. *Science* **357**, 1142–1145 (2017).

22.    H. Yang, M. Howard, C. Dean, Antagonistic roles for H3K36me3 and H3K27me3 in the

cold-induced epigenetic switch at Arabidopsis FLC. *Curr Biol* **24**, 1793–1797 (2014).

23.    S. Sung, R. M. Amasino, Vernalization and epigenetics: how plants remember winter.
*Curr Opin Plant Biol* **7**, 4–10 (2004).

24.    S. Berry, C. Dean, Environmental perception and epigenetic memory: mechanistic insight
through FLC. *Plant J* **83**, 133–148 (2015).

25.    P. Crevillén, *et al.*, Epigenetic reprogramming that prevents transgenerational
inheritance of the vernalized state. *Nature* **515**, 587–590 (2014).

26.    E. J. Finnegan, M. Robertson, C. A. Helliwell, Resetting FLOWERING LOCUS C Expression
After Vernalization Is Just Activation in the Early Embryo by a Different Name. *Frontiers*
*in Plant Science* **11** (2021).

27.    X. Luo, Y. He, Experiencing winter for spring flowering: A molecular epigenetic
perspective on vernalization. *J Integr Plant Biol* **62**, 104–117 (2020).

28.    C. C. Sheldon, *et al.*, Resetting of FLOWERING LOCUS C expression after epigenetic
repression by vernalization. *Proc Natl Acad Sci U S A* **105**, 2214–2219 (2008).

29.    J. Choi, *et al.*, Resetting and regulation of Flowering Locus C expression during
Arabidopsis reproductive development. *Plant J* **57**, 918–931 (2009).

30.    Z. Tao, *et al.*, Embryonic epigenetic reprogramming by a pioneer transcription factor in
plants. *Nature* **551**, 124–128 (2017).

31.    Z. Tao, *et al.*, Embryonic resetting of the parental vernalized state by two B3 domain
transcription factors in Arabidopsis. *Nat. Plants* **5**, 424–435 (2019).

32.    G. Xu, Z. Tao, Y. He, Embryonic reactivation of FLOWERING LOCUS C by ABSCISIC ACID-
INSENSITIVE 3 establishes the vernalization requirement in each Arabidopsis
generation. *Plant Cell* **34**, 2205–2221 (2022).

33.    N. Sharma, *et al.*, A Flowering Locus C Homolog Is a Vernalization-Regulated Repressor
in Brachypodium and Is Cold Regulated in Wheat. *Plant Physiol* **173**, 1301–1315 (2017).

34.    A. Kennedy, K. Geuten, The Role of FLOWERING LOCUS C Relatives in Cereals. *Front*
*Plant Sci* **11**, 617340 (2020).

35.    Q. Huan, Z. Mao, K. Chong, J. Zhang, Global analysis of H3K4me3/H3K27me3 in
Brachypodium distachyon reveals VRN3 as critical epigenetic regulation point in
vernalization and provides insights into epigenetic memory. *New Phytologist* **219**,
1373–1387 (2018).

36.    X. Liu, *et al.*, Uncovering the transcriptional regulatory network involved in boosting
wheat regeneration and transformation. *Nat. Plants*, 1–18 (2023).

37.    L. Zhao, *et al.*, Dynamic chromatin regulatory programs during embryogenesis of
hexaploid wheat. *Genome Biology* **24**, 7 (2023).

38.  J. Guo, Y. Ren, Z. Tang, W. Shi, M. Zhou, Characterization and expression profiling of the
ICE-CBF-COR genes in wheat. *PeerJ* **7**, e8190 (2019).

39.  W. Yong, *et al.*, Vernalization-induced flowering in wheat is mediated by a lectin-like
gene VER2. *Planta* **217**, 261–270 (2003).

40.  J. Ernst, M. Kellis, ChromHMM: automating chromatin-state discovery and
characterization. *Nat Methods* **9**, 215–216 (2012).

41.  B. E. Bernstein, *et al.*, A Bivalent Chromatin Structure Marks Key Developmental Genes
in Embryonic Stem Cells. *Cell* **125**, 315–326 (2006).

42.  S. L. Klemm, Z. Shipony, W. J. Greenleaf, Chromatin accessibility and the regulatory
epigenome. *Nat Rev Genet* **20**, 207–220 (2019).

43.  C. Li, *et al.*, Wheat VRN1, FUL2 and FUL3 play critical and redundant roles in spikelet
development and spike determinacy. *Development* **146**, dev175398 (2019).

44.  K. Wang, H. Liu, L. Du, X. Ye, Generation of marker-free transgenic hexaploid wheat via
an Agrobacterium-mediated co-transformation strategy in commercial Chinese wheat
varieties. *Plant Biotechnology Journal* **15**, 614–623 (2017).

45.  J. Xiao, D. Wagner, Polycomb repression in the regulation of growth and development
in Arabidopsis. *Current Opinion in Plant Biology* **23**, 15–24 (2015).

46.  A. Pajoro, E. Severing, G. C. Angenent, R. G. H. Immink, Histone H3 lysine 36 methylation
affects temperature-induced alternative splicing and flowering in plants. *Genome Biol*
**18**, 102 (2017).

47.  N. Shitsukawa, *et al.*, Wheat SOC1 functions independently of WAP1/VRN1, an
integrator of vernalization and photoperiod flowering promotion pathways. *Physiologia*
*Plantarum* **130**, 627–636 (2007).

48.  S. D. Michaels, R. M. Amasino, FLOWERING LOCUS C encodes a novel MADS domain
protein that acts as a repressor of flowering. *Plant Cell* **11**, 949–956 (1999).

49.  N. Sharma, K. Geuten, B. S. Giri, A. Varma, The molecular mechanism of vernalization in
Arabidopsis and cereals: role of Flowering Locus C and its homologs. *Physiol Plant* **170**,
373–383 (2020).

50.  M. Zikhali, L. U. Wingen, M. Leverington-Waite, S. Specel, S. Griffiths, The identification
of new candidate genes Triticum aestivum FLOWERING LOCUS T3-B1 (TaFT3-B1) and
TARGET OF EAT1 (TaTOE1-B1) controlling the short-day photoperiod response in
bread wheat. *Plant Cell Environ* **40**, 2678–2690 (2017).

51.  T. S. Ream, D. P. Woods, R. M. Amasino, The molecular basis of vernalization in different
plant groups. *Cold Spring Harb Symp Quant Biol* **77**, 105–115 (2012).

52.  S. Sung, R. M. Amasino, REMEMBERING WINTER: Toward a Molecular Understanding of
Vernalization. *Annual Review of Plant Biology* **56**, 491–508 (2005).

53.    J. Song, A. Angel, M. Howard, C. Dean, Vernalization – a cold-induced epigenetic switch.
715           *Journal of Cell Science* **125**, 3723–3731 (2012).

716    54.    P. Ruelens, *et al.*, FLOWERING LOCUS C in monocots and the tandem origin of
717           angiosperm-specific MADS-box genes. *Nat Commun* **4**, 2280 (2013).

718    55.    International Wheat Genome Sequencing Consortium (IWGSC), Shifting the limits in
719           wheat research and breeding using a fully annotated reference genome. *Science* **361**,
720           eaar7191 (2018).

721    56.    A. G. Greenup, *et al.*, ODDSOC2 Is a MADS Box Floral Repressor That Is Down-Regulated
722           by Vernalization in Temperate Cereals. *Plant Physiology* **153**, 1062–1073 (2010).

723    57.    C. B. Field, V. Barros, T. F. Stocker, Q. Dahe, Eds., *Managing the Risks of Extreme Events*
724           *and Disasters to Advance Climate Change Adaptation: Special Report of the*
725           *Intergovernmental Panel on Climate Change* (Cambridge University Press, 2012)
726           https:/doi.org/10.1017/CBO9781139177245 (December 19, 2023).

727    58.    D. Wang, *et al.*, Boosting wheat functional genomics via an indexed EMS mutant library
728           of KN9204. *Plant Comm* **4** (2023).

729    59.    S. Chen, Y. Zhou, Y. Chen, J. Gu, fastp: an ultra-fast all-in-one FASTQ preprocessor.
730           *Bioinformatics* **34**, i884–i890 (2018).

731    60.    D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment
732           and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915 (2019).

733    61.    P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008
734           (2021).

735    62.    Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for
736           assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

737    63.    M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion
738           for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

739    64.    Z. Gu, R. Eils, M. Schlesner, Complex heatmaps reveal patterns and correlations in
740           multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

741    65.    T. Wu, *et al.*, clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
742           *Innovation (Camb)* **2**, 100141 (2021).

743    66.    H. Mi, A. Muruganujan, D. Ebert, X. Huang, P. D. Thomas, PANTHER version 14: more
744           genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools.
745           *Nucleic Acids Res* **47**, D419–D426 (2019).

746    67.    F. Tian, D.-C. Yang, Y.-Q. Meng, J. Jin, G. Gao, PlantRegMap: charting functional
747           regulatory maps in plants. *Nucleic Acids Res* **48**, D1104–D1113 (2020).

748    68.    H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler
749           transform. *Bioinformatics* **25**, 1754–1760 (2009).

69.  F. Ramírez, *et al.*, deepTools2: a next generation web server for deep-sequencing data
analysis. *Nucleic Acids Res* **44**, W160-165 (2016).

70.  H. Thorvaldsdóttir, J. T. Robinson, J. P. Mesirov, Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. *Brief Bioinform* **14**, 178–192
(2013).

71.  Y. Zhang, *et al.*, Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

72.  A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).

73.  G. Yu, L.-G. Wang, Q.-Y. He, ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. *Bioinformatics* **31**, 2382–2383 (2015).

74.  S. Cheng, *et al.*, Harnessing Landrace Diversity Empowers Wheat Breeding for Climate
Resilience. 2023.10.04.560903 (2023).

75.  L. S. Johnson, S. R. Eddy, E. Portugaly, Hidden Markov model speed heuristic and
iterative HMM search procedure. *BMC Bioinformatics* **11**, 431 (2010).

76.  R. C. Edgar, Muscle5: High-accuracy alignment ensembles enable unbiased assessments
of sequence homology and phylogeny. *Nat Commun* **13**, 6968 (2022).

77.  S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973
(2009).

78.  A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

79.  Y. Wang, *et al.*, Simultaneous editing of three homoeoalleles in hexaploid bread wheat
confers heritable resistance to powdery mildew. *Nat Biotechnol* **32**, 947–951 (2014).

80.  Q. Shan, *et al.*, Targeted genome modification of crop plants using a CRISPR-Cas
system. *Nat Biotechnol* **31**, 686–688 (2013).

81.  R. Cui, *et al.*, Functional conservation and diversification of class E floral homeotic genes
in rice (Oryza sativa). *Plant J* **61**, 767–781 (2010).

82.  K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402–408 (2001).

83.  C. Tian, *et al.*, Impaired histone inheritance promotes tumor progression. *Nat Commun*
**14**, 3429 (2023).

84.  X. Lin, *et al.*, Systematic mining and functional analysis of factors regulating wheat spike
development for breeding selection. 2022.11.11.516122 (2023).

85.  S. Schilling, A. Kennedy, S. Pan, L. S. Jermiin, R. Melzer, Genome-wide analysis of MIKC-
type MADS-box genes in wheat: pervasive duplications, functional conservation and

putative neofunctionalization. *New Phytol* **225**, 511–529 (2020).

**Figures**

**Fig. 1. Profiling the transcriptome and epigenome in the shoot apex during**
**vernalization in winter wheat.**

**A**. Temperature at different sampling time points. DAP: Days After Pollination.

**B.** Data types and sampling time points in this study.

**C.** K-means clustering heatmap of DEGs. The color represents the level of transcription.

**D.** GO enrichment analysis of DEGs in all clusters.

**E.** Transcription patterns of *TaICE2*, *TaCOR14*, *TaCOR15* and *VRN1* during vernalization.

798
799 **Fig. S1 Quality control for transcriptome and epigenome data.**

800 **A.** Pearson correlation analysis of all transcriptome data. The color represents Pearson
801 correlation coefficient.

**Fig. 2 The chromatin state changes during vernalization establishment and**

**maintenance.**

**A.** 11 chromatin states informed from all CUT&Tag data. Each row represents a chromatin

state and each column respectively represents histone marker enrichment (Emission

Probability), genomic coverage, genomic location distribution (Enrichment), gene

expression level and chromatin accessibility level.

**B.** Comparison of variable bases fraction between different chromatin state. The letters

represent the results of multiple comparisons, with a significance level of 0.05.

**C. The fraction of bases that change chromatin state between different stages.**

**D.** In situ hybridization of *VRN1* at S0, S2, S4, and S6 stages.

**E.** The chromatin state, H3K27me3, H3K36me3 and transcriptional level changes of

*VRN1_5A* from S0 to S6.

**Fig. S2 Changes of individual chromatin state during vernalization.**

**A.** Correlation of different chromatin state models trained by chromHMM. The color represents the level of correlation.

**B.** The levels of H3K27me3 (top) and H3K36me3 (bottom) in genes with different transcription levels, take S0 as an example. Different colors represent genes at different levels of transcription. All genes in the S0 period were ranked from lowest to highest expression, with 0-20% representing the top 20% with the lowest expression and 80-100% representing the top 20% with the highest expression.

**C.** Changes in E9 state from S0-S3 to S4 stage.

**D.** The GO enrichment analysis of genes with altered E9 states from S0-S3 to S4 stage.

**E, F.** The chromatin state, H3K27me3, H3K36me3 and transcriptional level changes of *VRN1_5B* (E) and *VRN1_5D* (F) from S0 to S6.

**Fig. 3 Mutation of TaFIE and TaSDG8 affects vernalisation-induced flowering.**

**A, B.** Photos (A) and heading time (B) of wild type (KN199) and *Tafie* crispr/cas9 line (*Tafie-cr-87*) under different vernalization treatments. V0, V20, and V30 represent 0, 20, and 30 days of treatment at 4°C respectively.

**C**. The relative expression level of *VRN1_5A* in seedlings of wild type and *Tafie* crispr/cas9 line under different vernalization treatments.

**D**. The H3K27me3 level at different location of *VRN1_5A* in seedlings of wild type (KN199) and *Tafie* crispr/cas9 line (*Tafie-cr-87*) at heading stage under different vernalization treatments.

**E, F.** Photos (E) and heading time (F) of wild type (KN199) and *Tasdg8* crispr/cas9 lines (*Tasdg8-cr-3*, *Tasdg8-cr-5*) under 30 days of vernalization treatment at 4°C.

**G.** The relative expression level of *VRN1_5A* in seedlings of wild type (KN199) and *Tasdg8* crispr/cas9 (*Tasdg8-cr-3*, *Tasdg8-cr-5*) lines under different vernalization treatments. V30+10 represents 30 day days of treatment at 4°C and subsequent 10 days treatment at normal temperature.

**H**. The H3K36me3 level at different location of *VRN1_5A* in seedlings of wild type (KN199) and *Tasdg8* crispr/cas9 (*Tasdg8-cr-3*, *Tasdg8-cr-5*) lines under different vernalization

treatments.

**Fig. S3 Generation of *Tafie* and *Tasdg8* crispr/cas9 lines.**

**A.** Indication of genome editing sites and genotyping of *Tafie* crispr/cas9 line.

**B.** Indication of genome editing sites and genotyping of *Tasdg8* crispr/cas9 lines.

**C**. The western blot shows the H3K27me3, H3K27me2, and H3K36me3 level in wild type

(KN199) and *Tafie, Tasdg8* crispr/cas9 lines (*Tafie-cr-87 and Tasdg8-cr-3/5*). The number

show quantification of protein levels compared to the wild type (KN199) in each condition.

**D**. The phylogenetic tree and expression level of *TaSDG8*.

**E.** The relative expression level of *VRN1_5B* and *VRN1_5D* in seedlings of wild type

(KN199) and *Tafie* crispr/cas9 (*Tafie-cr-87*) line under different vernalization treatments.

**F.** The H3K27me3 level at different location of *VRN1_5B* and *VRN1_5D* in seedlings of

wild type (KN199) and *Tafie* crispr/cas9 line (*Tafie-cr-87*) at heading stage under different

vernalization treatments.

**G.** The relative expression level of *VRN1_5A* and *VRN1_5D* in seedlings of wild type

(KN199) and *Tasdg8* crispr/cas9 (*Tasdg8-cr-3* and *Tasdg8-cr-5*) lines under different

 vernalization treatments.

**H.** The H3K36me3 level at different location of *VRN1_5B* and *VRN1_5D* in seedlings of

wild type (KN199) and *Tasdg8* crispr/cas9 (*Tasdg8-cr-3*, *Tasdg8-cr-5*) lines under different

vernalization treatments.

**Fig. 4 H3K27me3 restoration associates with the reset of vernalization requirement**

**during embryogenesis.**

**A.** In situ hybridization of *VRN1* at DAP0, DAP2, DAP6, and DAP14 stages.

**B.** The expression level of *VRN1* during embryogenesis in winter wheat KN9204.

**C.** The H3K27me3 and H3K36me3 changes of *VRN1_5A* during embryogenesis in winter

wheat KN9204.

**D**. The expression level of *VRN1* during embryogenesis in spring wheat Chinese Spring.

**E.** The H3K27me3 and H3K36me3 changes of recessive allele *VRN1_5A* during

embryogenesis in spring wheat CS.

**F.** The relative expression level of *VRN1_5A* during embryogenesis of wild type (KN199)
and *Tafie* crispr/cas9 (*Tafie-cr-87*) line.
**G.** The H3K27me3 level at different location of *VRN1_5A* during embryogenesis of wild
type (KN199) and *Tafie* crispr/cas9 line (*Tafie-cr-87*).
**H.** The relative expression level of *VRN1_5A* in leaves of wild type (KN199) and *Tafie*
crispr/cas9 (*Tafie-cr-87*) line.
**I.** The H3K27me3 level at different location of *VRN1_5A* in leaves of wild type (KN199) and
*Tafie* crispr/cas9 line (*Tafie-cr-87*).
894

**Fig. S4 mRNA level and H3K27me3, H3K36me3 abundance of VRN1_5B and**
**VRN1_5D during embryogenesis A.** The H3K27me3 and H3K36me3 changes of

*VRN1_5B* and *VRN1_5D* during embryogenesis in winter wheat KN9204.

**B.** The H3K27me3 and H3K36me3 changes of recessive allele *VRN1_5B* and dominant

allele *VRN1_5D* during embryogenesis in spring wheat CS.

**C.** The relative expression level of *VRN1* in leaves of CS and KN9204.

**D, H**. The relative expression level of *VRN1_5B* in embryos and leaves of wild type

(KN199) and *Tafie* crispr/cas9 (*Tafie-cr-87*) line.

**E, I.** The H3K27me3 level at different location of *VRN1_5B* in embryos and leaves of wild

type (KN199) and *Tafie* crispr/cas9 line (*Tafie-cr-87*).

**F, J.** The relative expression level of *VRN1_5D* in embryos and leaves of wild type (KN199)

and *Tafie* crispr/cas9 (*Tafie-cr-87*) line.

**G, K.** The H3K27me3 level at different location of *VRN1_5D* in embryos and leaves of wild

type (KN199) and *Tafie* crispr/cas9 line (*Tafie-cr-87*).

**Fig. 5 Identification and characterization of *VRN1*-pattern-like gene.**

**A.** The transcription pattern of *VRN1*-pattern-like genes in C2 and C4. Other refers to other genes in C2 and C4.

**B.** The H3K27me3 and H3K36me3 dynamic of *VRN1*-pattern-like genes in C2 and C4. Other refers to other genes in C2 and C4.

**C.** The GO enrichment analysis of *VRN1*-pattern-like genes and other genes in C2 and C4.

**D.** The TF family enrichment analysis of *VRN1*-pattern-like genes and other genes in C2 and C4.

**E.** Different types of *VRN1*-pattern-like genes. T-test represents SNP sites on the promoter or genebody of the gene that are significantly associated with heading time. Mutant represents an abnormal flowering phenotype in plants with this gene mutation.

**F.** The H3K27me3, H3K36me3 and transcriptional level changes of *TaFUL3_2A* from S0 to S6.

**G**. The heading time of two haplotypes divided by seven SNPs (red line in F) within promoter and genebody of *TaFUL3_2A*.

930 **Fig. S5 Expression pattern of MADS genes from S0 to S6.**

931 VRN1 represents genes in *VRN1* subfamily, SEP represents genes in *SEP* subfamily, FLC

932 represents genes in *FLC* subfamily and other represents genes in other subfamily of

933 MADS.

935
**Fig. 6 Identification and characterization of *FLC*-pattern-like gene.**

**A.** The H3K27me3 and H3K36me3 dynamic of FLC orthologs reported in previous study
(85).

**B.** The transcription pattern of *FLC*-pattern-like genes in C5. Other refers to other genes in
C5.

**C.** The H3K27me3 and H3K36me3 dynamic of *FLC*-pattern-like genes in C5. Other refers

to other genes in C5.

**D.** The GO enrichment analysis of *FLC*-pattern-like genes.

**E.** Different types of *FLC*-pattern-like genes. T-test represents SNP sites on the promoter

or genebody of the gene that are significantly associated with heading time. Mutant

represents an abnormal flowering phenotype in plants with this gene mutation.

**F.** The heading time of two haplotypes divided by eight SNPs (red line in G) within promoter

and genebody of *TaTOE_1A*.

**G.** The H3K27me3, H3K36me3 and transcriptional level changes of *TaTOE_1A* from S0 to

S6.

**Fig. S6 Phylogenetic tree of the MADS family from *Arabidopsis thaliana*, wheat, rice,**

**barley and *Brachypodium distachyon*.**

Blue represents the subfamily of Arabidopsis *FLC*, red represents the subfamily of wheat

*VRN1*, yellow represents the subfamily of *SEP,* and green represents the subfamily of the

previously reported orthologs of *FLC* in wheat (85).

**Supplementary tables**

**Table S1** Differentially expressed genes from S0 to S4.

**Table S2** The expression level of *ICE-CBF-COR* cluster genes.

**Table S3** The changes of state E9 from S0-S3 to S4.

**Table S4** The VRN1-pattern-like and FLC-pattern-like genes.

**Table S5** The expression level of genes in MADS family.

**Table S6** SNPs located within the promoter of genebody of *VRN1*-pattern-like or *FLC*-pattern-like genes that were significantly associated with heading time in GWAS.

**Table S7** Mutant lines containing severe mutations in *VRN1*-pattern-like or *FLC*-pattern-like genes.

**Table S8** SNPs associated with heading time located on the promoter and genebody of *TaFUL3_*2A.

**Table S9** SNPs associated with heading time located on the promoter and genebody of *TaTOE_*1A.

**Table S10** Antibodies used in this study.

**Table S11** Primers used in this study.