Emergence of a senescent and inflammatory pulmonary CD4[+] T cell population prior to lung allograft failure

Sajad Moshkelgosha[1], Liran Levy[1,2], Shahideh Safavi[1,3], Sumiha Karunagaran[1], Gavin Wilson[1], Benjamin Renaud-Picard[1,4], Goodness Madu[1], Rashi Ramchandani[1], Jillian Oliver[1], Tatsuaki Watanabe[1,5], Ke Fan Bei[1], Betty Joe[1], Qixuan Li[6], Ella Huszti[6], May Cheung[1], David Hedley[7], Jonathan Yeung[1,8,9], Shaf Keshavjee[1,8,9], Tereza Martinu[1,8,10], Stephen Juvet[1,8,10]

1. Toronto General Hospital Research Institute, University Health Network, Toronto, ON
2. Sheba Medical Center, Sackler School of Medicine, Tel-Aviv University, Tel-Aviv, Israel
3. Barts Health NHS Trust, London, UK
4. Nouvel Hôpital Civil, Strasbourg, France
5. Department of Thoracic Surggery, Institute of Development, Aging and Cancer, Tohoku University, Sendai, Japan
6. Biostatistics Department, University Health Network, Toronto, ON
7. Division of Medical Oncology, Department of Medicine, University of Toronto
8. Toronto Lung Transplant Program, Ajmera Transplant Centre, University Health Network, Toronto, ON
9. Division of Thoracic Surgery, Department of Surgery, University of Toronto, Toronto, ON
10. Division of Respirology, Department of Medicine, University of Toronto, Toronto, ON

Address of Corresponding Author:
Stephen Juvet, MD, PhD
Toronto General Hospital: University Health Network (UHN)
200 Elizabeth St., Rm 11 PMB 126
Toronto, Ontario, M5G 2C4
EMAIL: stephen.juvet@uhn.ca
TEL: (416) 340-4800 - ext: 8178

Acknowledgments

The authors wish to thank Martin Oberle, Marcelo Cuesta, Iva Avramov and other members of the Toronto Lung Transplant Program Biobank for their assistance with sample procurement and storage. We also thank Dr. Cynthia Guidos and Dr. Dariush Davani for their critical input on the design of mass cytometry experiments. We also wish to thank Dr. Sarah Crome and Sarah Colpitts for their assistance and advice in planning CITE-Seq experiments.

This work was funded by Cystic Fibrosis Foundation grants JUVET18AB0 and JUVET23G0 (to S.J.), and MARTIN18I0 (to T.M. and S.J.) by an International Society for Heart and Lung Transplantation Norman Shumway Career Development Award (to S.J.), by a Canadian Society of Transplantation Research Training Award (to S.M.) and by the University Health Network Foundation.

Abbreviations:

ALAD: Acute lung allograft dysfunction

CLAD: Chronic lung allograft dysfunction

LT: Lung transplantation

BAL: Bronchoalveolar lavage

IMC: Imaging mass cytometry

tSNE: T-distributed stochastic neighbor embedding

CyTOF: Cytometry by Time of Flight

CITE-seq: Cellular indexing of transcriptomes and epitopes

Tph: peripheral helper T

FEV1: Forced expiratory volume in one second

TLTP: Toronto lung transplant program

GLMM: Generalized linear mixed model

ROC: Receiver-operating characteristic

DSA: Donor specific antibodies

HR: Hazard radio

CMV: Cytomegalovirus

ADT: Antibody derived tags

RICD: Restimulation induced cell death

Abstract

Lung transplantation is a life-saving therapy for end-stage pulmonary disease, but its long-term outlook is poor due to a high incidence of chronic lung allograft dysfunction (CLAD). CLAD results from alloimmune-mediated progressive fibrosis and culminates in death or the need for re-transplantation after a median of 6 years. Existing immunosuppression fails to prevent CLAD, suggesting the existence of alloimmune pathways resistant to these drugs. Here, we used mass cytometry to identify cell populations enriched in the bronchoalveolar lavage (BAL) of patients with subsequent allograft dysfunction. We show that $CD4^+CD57^+PD1^+$ T cells emerge in stable lung transplant recipients in the first year post-transplant, conferring heightened risks for CLAD and death or re-transplantation. $CD4^+CD57^+PD1^+$ T cells display features of senescence and secrete inflammatory cytokines. Cellular indexing by transcriptomes and epitopes (CITE-Seq) on BAL $CD4^+$ T cells revealed the existence of 2 oligoclonal $CD57^+$ subsets with putative cytotoxic and follicular helper functions. Finally, we observed that $CD4^+CD57^+PD1^+$ T cells are associated with lung allograft fibrosis in a mouse model and in human explanted CLAD lungs, where they localize near airway epithelium and B cells. Together, our findings reveal the existence of an inflammatory T cell population that predicts future lung allograft dysfunction and may represent a rational therapeutic target in lung transplant recipients.

The study of allograft rejection has led to important advances in our understanding of cellular immunity[1]. Unfortunately, solid organ transplantation continues to be hampered by high rates of chronic rejection and graft failure despite the availability of potent immunosuppressive drugs. In lung transplantation (LT), which remains the only life-prolonging therapy for many cases of end-stage lung disease, median graft survival is only 6 years[2] due to chronic lung allograft dysfunction (CLAD). CLAD is the result of a progressive scarring process due to accumulated injuries resulting from alloreactivity and innate immune activation caused by donor brain death, ischemia-reperfusion injury, post-transplant infections, gastroesophageal reflux, and air pollution[3]. Both acute T cell-mediated rejection and antibody-mediated rejection are well-recognized CLAD risk factors[4,5].

LT recipients often grapple with episodes of acute lung allograft dysfunction (ALAD)[6]. We defined ALAD as a marked decrement in lung function – quantified by a decline in the forced expiratory volume in one second (FEV_1) by 10% or more from its prior value. Despite exhaustive history, physical examination, and diagnostic testing including thoracic imaging and bronchoscopy, the cause of ALAD often is not definitively identified. Recurring ALAD events may be the result of rejection or uncontrolled alloimmunity that cannot be identified due to the poor diagnostic accuracy of transbronchial biopsies [7,8].

Bronchoalveolar lavage (BAL) is a technique in which saline instilled via a bronchoscope wedged in a pulmonary segmental airway is aspirated into a collection trap. BAL permits microbiologic and cytologic analysis of the distal lung compartments including the small airways and alveolar spaces. We have been studying cellular elements of the BAL using advanced multiparametric analytical approaches for the last several years[6,9-13]. Using single-cell RNA sequencing, we recently identified specific alveolar macrophage populations associated with ALAD episodes and CLAD[6], which may represent both a mechanistic link between inflammatory insults such as infection and rejection, as well as a potential target for therapeutic intervention.

Prior studies by several groups have examined specific BAL T cell populations in LT recipients. Early work examined the question of whether changes in the CD4:CD8 ratio

among BAL T cells might differentiate infection from rejection[14]. Later studies examined

whether T cell activation[15], differentiation[16], regulatory T cells[17] and apoptosis[18] could

distinguish CLAD, acute rejection and infection[19]. High variability between LT recipients

meant that such distinctions were difficult[19] and not useful in an individual patient.

Moreover, since acute rejection and infection and CLAD can all coexist, and there are

large overlaps in the distributions of pre-specified BAL immune cell populations between

these clinical entities, these hypothesis-directed analyses have not led to the

identification of useful biomarkers or mechanistic insights.

Here, we used mass cytometry[20] – a high-dimensional technique in which the

expression of up to 40 cell-associated proteins can be analyzed simultaneously,

coupled with unbiased differential discovery approaches – to identify $CD4^+CD57^+PD1^+$

T cells as a predictor of incident ALAD and CLAD. Apart from their usefulness as a

biomarker, our data implicate $CD4^+CD57^+PD1^+$ T cells in the mechanism of allograft

loss. We find the cells in chronically rejected human and mouse lung allografts, show

that they are clonally expanded, express transcripts associated with cytotoxic and

helper functions, and they secrete inflammatory cytokines. Further, they are senescent,

hypoproliferative, and resistant to cell death *in vitro.* Our observations therefore suggest

the existence of a specific alloimmune pathway that can escape conventional

immunosuppression that may represent a valuable therapeutic target.

Results

Identification of a T cell subset associated with lung allograft dysfunction

Early detection of alloimmune responses in the graft – before ALAD onset – is likely to

provide both important prognostic information as well as an opportunity for early

intervention. In pursuit of a greater understanding of ALAD mechanisms, we

prospectively collected and viably cryopreserved BAL cells from 50 Toronto Lung

Transplant Program (TLTP) LT recipients presenting for surveillance bronchoscopy at 3

months post-transplant, and obtained two more follow up samples from each recipient

to form a longitudinal cohort **(Figure 1A; Supp Fig 1)**. We performed mass cytometry[20]

using a panel of 36 heavy metal-conjugated antibodies **(Supp Table 1)**. Conventional

flow cytometric gating revealed considerable inter- and intra-patient heterogeneity in

BAL cell composition (data not shown). Unsupervised multidimensional clustering using

FlowSOM [21] demonstrated similar variation **(Figure 1B, Supp Fig 2)**. In order to

facilitate identification of cell populations associated with allograft dysfunction, we

randomly divided the cohort into discovery and validation subsets (n=25 each; **Supp**

Table 2). Longitudinal samples in the discovery group were then used to search for

differences in the BAL cellular composition between stable patients and those

experiencing ALAD within 30 days of each bronchoscopy. Initially we employed cluster

identification, characterization and regression (CITRUS[22]) on all BAL T cells from all the

time-points for all patients in the cohort. This analysis revealed that $CD4^+CD57^+PD1^+$ T

cells were the top differentially represented cell population between patients with and

without ALAD within 30 days of at least one bronchoscopy **(Supp Fig 3)**. CITRUS does

not adequately account for the paired design of our cohort (multiple samples per

patient), nor does it incorporate time as a factor in the analysis. We therefore also

applied a generalized linear mixed model (GLMM) pipeline described for CyTOF

analysis[23]. To facilitate this analysis, we used FlowSOM to identify 20 clusters of T cells

(Figure 1C). T cell subsets with greater representation in samples associated with

ALAD compared to samples associated with stability were identified using this approach

(Figure 1D). The expression profile of T cell markers revealed that the $CD4^+$

$CD57^+PD1^+$ T cell population was significantly differentially represented in ALAD

(Figure 1E, Supp Fig 4). Collectively, these data suggested that there is a strong

association between CD4$^+$CD57$^+$PD1$^+$ T cells and incident ALAD.

CD4$^+$CD57$^+$PD1$^+$ T cells appear in the BAL prior to the onset of allograft

dysfunction

Next, we assessed the relationship of BAL CD4$^+$CD57$^+$PD1$^+$ T cells to ALAD onset. We

observed that while patients with stable lung allograft function had low frequencies of

CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell compartment (**Figure 2A**), these cells typically

accumulated in higher frequencies prior to the onset of ALAD **(Figure 2B)**. Receiver-

operating characteristic (ROC) curve analysis showed that having $\geq$ 7.8% CD57$^+$PD1$^+$

cells in the BAL CD4$^+$ T cell compartment optimally discriminated concurrent ALAD from

stable patients, with an area under the ROC curve of 0.76 **(Figure 2C)**. Using this

criterion, 10 of 11 (91%) LT recipients with ALAD within 30 days of any of their

bronchoscopies demonstrated $\geq$ 7.8% CD57$^+$PD1$^+$ in the CD4$^+$ T cell compartment of at

least one of their BAL samples; in contrast, only 3 of 14 (21%) stable LT recipients had

CD57$^+$PD1$^+$ cells exceeding this threshold **(Figure 2D)**. Moreover, having this

biomarker conferred a reduced time-to-ALAD in univariable survival analysis **(Figure**

2E). We then applied the 7.8% threshold to the validation subset (n=25) and observed a

similar reduction in time-to-ALAD from first appearance of the cells in LT recipients with

at least one BAL sample in which it was exceeded, compared to LT recipients who did

not have any BAL samples in which the threshold was exceeded **(Figure 2F)**. Together,

these data illustrated that in LT recipients developing subsequent allograft dysfunction,

CD57$^+$PD1$^+$ cells frequently exceeded the 7.8% threshold in BAL samples obtained

prior to ALAD and CLAD **(Supp Fig 5).**

Anti-human leukocyte antigen donor-specific antibodies (DSA) are an adverse

prognostic factor in lung transplantation[24]. Patients with $\geq$7.8% CD57$^+$PD1$^+$ cells in the

BAL CD4$^+$ T cell compartment were more likely to develop de novo DSA than those

without **(Figure 2G)**. BAL CD4$^+$CD57$^+$PD1$^+$ T cells were not simply a surrogate for other

conventional clinical and bronchoscopic parameters, since clinical symptoms, biopsy-

proven vascular or airway rejection, BAL pathogens and antimicrobial treatments did not

differ significantly between patients with and without $\geq$ 7.8% CD57$^+$PD1$^+$ cells in the

BAL CD4$^+$ T cell compartment **(Supp Table 3)**. Similarly, although the number of events was small, having CD57$^+$PD1$^+$ cells above the 7.8% threshold was not clearly associated with an immediate post-bronchoscopy augmentation of immunosuppression or change in calcineurin inhibitor **(Supp Table 4)**. Taken together, these observations indicate that BAL CD4$^+$CD57$^+$PD1$^+$ T cells may be a useful biomarker of incident lung allograft dysfunction that adds significant diagnostic utility to conventional clinical monitoring.

CD4$^+$CD57$^+$PD1$^+$ T cells confer an increased risk for CLAD and death and are present in chronically rejected human lung allografts

To examine whether the presence of CD4$^+$CD57$^+$PD1$^+$ T cells in the BAL samples of LT recipients increase the risk for CLAD, we followed all 50 individuals in our cohort for up to 2000 days post-LT (median follow-up 1788 days). Using the appearance of ≥7.8% CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell compartment or the last analyzed BAL as the index event, we found that time to CLAD was significantly shorter in those patients whose BAL samples exceeded the 7.8% threshold (log rank p=0.0071, **Supp Figure 6**). We then performed Cox proportional hazards analysis, excluding two patients who developed CLAD prior to their final BAL samples (**Supp Fig 1**). Treating the appearance of ≥7.8% CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell compartment as a time-dependent covariate, Cox proportional hazards analysis revealed a hazard ratio (HR) of 3.61 (95% CI 1.55-8.39) for time-to-CLAD. In this and other survival analyses, patients were followed until 31 December 2022 (median follow up time from transplantation: 2245 days). The increased risk for CLAD associated with CD4$^+$CD57$^+$PD1$^+$ cells persisted in bivariable models adjusted for age at transplant (HR 3.67, 95% CI 1.57-8.55), sex (HR 3.45, 95% CI 1.47-8.07), and primary cytomegalovirus (CMV) mismatch (donor seropositive, recipient seronegative; HR 4.02, 95% CI 1.70-9.52).

Similarly, a Cox proportional hazards model revealed that ≥7.8% CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell compartment was associated with a HR of 4.82 (95% CI 1.8-13.1) for time-to-death or re-transplantation. The effect persisted in bivariable models adjusting for age at transplant (HR 4.82, 85% CI 1.8-13.1), sex (HR 4.84, 1.8-13.2) and primary CMV mismatch (HR 4.92, 95% CI 1.8-13.4). These time-to-CLAD and time-to-

death or re-transplantation analyses are presented in **Supp Table 5**. We then examined

time-to-death or re-transplantation in a survival analysis landmarked at 12 months post-

transplant. LT recipients with $\geq$ 7.8% CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell

compartment had an increased risk of death or re-transplantation (p=0.00079, **Figure**

3A).

These observations suggested that CD4$^+$CD57$^+$PD1$^+$ T cells may lie on the mechanistic

pathway leading to lung allograft fibrosis. Therefore, we sought to identify them in CLAD

lung tissue obtained at the time of re-transplantation[25] using imaging mass cytometry

(IMC[26]). CLAD and normal lung tissue sections were stained with a panel of 35 heavy

metal conjugated antibodies **(Supp Table 6)** and subjected to IMC. Unsupervised

clustering identified CD4$^+$CD57$^+$PD1$^+$ T cells as one of the cell populations present

(cluster 50 in **Figure 3B**), with more cells in CLAD compared to normal tissue (**Figure**

3C-D). Image reconstruction and proximity analysis revealed that CD4$^+$CD57$^+$PD1$^+$ T

cells were frequently located adjacent to distinct B cell populations, including trafficking

B cells (CD20$^+$CCR6$^+$, Cluster 8) and plasma cells (IgM$^+$CD138$^+$, cluster 30) and

airways **(Figure 3E-F)**. The enrichment of CD4$^+$CD57$^+$PD-1$^+$ T cells in CLAD tissue

compared to normal lung tissue may indicate their involvement in disease development.

CD4$^+$CD57$^+$PD1$^+$ T cells infiltrate mouse lung allografts and associate with the

severity of chronic rejection

We have developed a minor alloantigen-mismatched mouse orthotopic lung transplant

model in which a minimal graft storage period prior to transplantation results in mild or

absent allograft fibrosis at 28 days post-transplant, while prolonged warm and cold

ischemic storage prior to transplantation leads to B cell- and alloimmune-dependent

fibrosis with features of human chronic lung allograft rejection[27] (**Figure 4A-B**). We

asked whether CD4$^+$CD57$^+$PD-1$^+$ T cells might be found within lung allografts in this

model, and whether their abundance correlated with the degree of allograft fibrosis.

Allografts were scored in blinded fashion using our published scoring system[28], which

includes assessment of the proportion of the allograft affected by interstitial

inflammation and fibrosis. As we have observed previously[27,28], pathology was variable,

with some animals exhibiting higher degrees of fibrosis and inflammation (**Fig 4C**). Flow

cytometry performed on single-cell suspensions of digested lung allograft tissue
revealed that CD57[+]PD1[+] T cells were common among lung allograft CD4[+] T cells, but
rare in normal lung tissue (**Figure 4D**). It also showed that allografts with higher
degrees of pathology – as determined by the presence of both interstitial inflammation
and parenchymal fibrosis – had higher frequencies of CD57[+]PD1[+] T cells compared to
those with a low degree of pathology (**Figure 4E-F**). Further, CD4[+]CD57[+]PD1[+] T cells
were readily detectable in allograft sections by immunofluorescence (**Figure 4G**).
Together, these data support the concept that CD4[+]CD57[+]PD1[+] T cells accumulate in
rejecting lung allografts and correlate with the severity of inflammation and fibrosis in
mice.

Identification of two transcriptionally distinct CD4[+]CD57[+]PD1[+] cell subsets using
CITE-Seq

To characterize gene expression in CD4[+]CD57[+]PD1[+] T cells, we enriched CD3[+] T cells
from four ALAD BAL samples and performed cellular indexing by transcriptomes and
epitopes (CITE-Seq[29]) with 5' end sequencing. We applied a panel of oligonucleotide-
conjugated antibodies, including CD4, CD57 and PD1, to ALAD BAL samples
containing at least 7.8% CD57[+]PD1[+] cells in the CD4[+] T cell compartment as
determined by flow cytometry. The conjugated oligonucleotides or antibody-derived tags
(ADTs) identified in CITE-Seq allowed confident linkage of transcriptomes with protein
expression.

BAL CD4[+] T cells from patients with ALAD fell into 9 subclusters in multidimensional
space (**Figure 5A**). Based on CD57 and PD1 protein expression, CD4[+]CD57[+]PD1[+] T
cells assembled into two main areas, cluster 4 and the closely adjacent clusters 1 and 2
(Figure 5A, Supp Fig 7A-B). We noted that CD27 gene and protein expression was
enriched in cluster 4 compared to clusters 1 and 2 **(Figure 5B).** The CD27[-] population
had a gene expression profile similar to peripheral helper T (Tph) cells described in
rheumatoid arthritis[30]; differentially expressed genes in this population included CXCL13,
TOX, IL-21, CD40L, TIGIT, ICOS and TNFSRF18 **(Figure 5C, Supp Fig 7C)**. These
cells are characterized by the absence of the conventional follicular helper T cell
markers CXCR5 and Bcl6, and by expression of the chemokine CXCL13 [30,31].

Accumulating studies have shown that Tph cells are involved in various acute and

chronic inflammatory diseases [30,32], and can promote B cell differentiation into

plasmablasts through interactions involving IL-21 and SLAMF5 [30]. Therefore, the Tph-

like features of CD27$^-$ CD4$^+$CD57$^+$PD1$^+$ T cells, along with their proximity to B cells

(Figure 3E-F), their association with de novo DSA **(Figure 2G)** suggest that they may

provide B cell help.

The CD27$^+$ population expresses several cytotoxicity-related transcripts including

granzymes and NKG7[33] **(Figure 5C)**. They also express CXCR6, which has been

associated with tissue resident memory CD8$^+$ T cell localization to airways[34] and is in

keeping with our observation of CD4$^+$CD57$^+$PD1$^+$ T cells in proximity to airways **(Figure**

3F).

We next examined TCR CDR3 sequences in the CITE-Seq data, which revealed that

CD57$^+$PD1$^+$CD27$^-$ cells are clonally expanded to a greater extent than

CD57$^+$PD1$^+$CD27$^+$ and other BAL CD4$^+$ T cells **(Figure 5D)**. This suggested that they

had recently undergone antigen-driven expansion in the lung. Analysis of the top 20

expanded TCR clonotypes did not infer any known antigen. However, the high similarity

among the top 20 expanded TCRs may suggest that they may be alloreactive **(Supp**

Figure 8). Pseudotime trajectory analysis revealed that CD57$^+$PD1$^+$CD27$^-$ cells and

their CD27$^+$ counterparts may share a common origin **(Figure 5E)**. Together, the TCR

sequence data and pseudotime analysis suggested that BAL CD4$^+$CD57$^+$PD1$^+$ T cells

either differentiate into Tph or cytotoxic effector cells. The former have undergone

significant clonal expansion, whereas the latter have a distinct TCR repertoire and less

clonal expansion, suggesting that they may be responding to different antigens in the

lung microenvironment.

Finally, both CD27$^-$ and CD27$^+$ subsets demonstrated expression of previously defined

senescence and exhaustion transcripts including TIGIT, KLRG1, LAG3, HAVCR2/TIM-3

and EOMES which have been described in different acute and chronic inflammatory

conditions[35-37]. This observation is in keeping with reports that have described different

senescent CD57$^+$ and/or PD1$^+$ T cell populations in other disease states [38,39].

Functional analysis of CD4⁺CD57⁺PD1⁺ T cells reveals features of inflammation and senescence

Considering the potential distinct functional properties – Tph, cytotoxic, and senescent – of CD4⁺CD57⁺PD1⁺ T cells, we set out to characterize their functions *in vitro*. For this purpose, CD4⁺ T cells from BAL samples of 15 lung transplant recipients experiencing ALAD were sorted to obtain CD4⁺CD57⁺PD1⁺ T cells and their CD4⁺CD57⁻PD1⁻ counterparts **(Figure 6A)**. Fluorescence-activated cell sorting data demonstrated that in all cases, CD57⁺PD1⁺ cell frequency was ≥7.8% of BAL CD4⁺ T cells **(Figure 6B)**. Based on the hypothesis that CD4⁺CD57⁺PD-1⁺ T cells may be senescent, we predicted that sorted cells would exhibit decreased proliferation, enhanced cytokine secretion, and resistance to TCR restimulation-induced cell death (RICD) when compared to CD57⁻PD1⁻ cells.

Supernatants from cells stimulated overnight were collected and subjected to a multiplexed cytokine assay. This revealed that CD4⁺CD57⁺PD1⁺ cells – compared to CD4⁺CD57⁻PD1⁻ cells sorted from the same BAL samples – expressed more IL-21, IL-4, CXCL8 and granzyme A **(Figure 6C)**. In keeping with our CITE-Seq data, these findings suggested that CD4⁺CD57⁺PD1⁺ T cells may have Tph or cytotoxic properties – or both. Stimulated CD4⁺CD57⁺PD1⁺ T cells and their CD57⁻PD1⁻ counterparts were then tested for proliferation, based on dilution of CFSE **(Figure 6D).** Proliferation data revealed that CD4⁺CD57⁺PD1⁺ T cells exhibited significantly less proliferation compared to CD4⁺CD57⁻PD1⁻ cells **(Figure 6E)**, in keeping with T cell senescence [38,39]. Finally, CD4⁺CD57⁺PD1⁺ T cells were less susceptible to RICD compared to CD4⁺CD57⁻PD1⁻ cells **(Figure 6F, G)**. Together these data imply that CD4⁺CD57⁺PD1⁺ T cells are senescent but pro-inflammatory T cells that are resistant to antigen-induced death. These observations may help to explain the emergence of ALAD and CLAD in lung transplant recipients despite potent triple-drug immunosuppression, which would be predicted to be less effective against these cells.

Discussion

In this study, we used a high dimensional mass cytometric discovery approach to

identify a T cell population in lung transplant recipient BAL that is associated with a

heightened risk for future allograft dysfunction, failure, and death. The risk conferred by

the appearance of $CD4^+CD57^+PD1^+$ T cells in the BAL remained elevated after bivariate

adjustments for age, sex, and donor and recipient CMV serostatus mismatch.

In keeping with previous work examining lung transplant recipient BAL, our samples

contained a diversity of leukocyte populations. The prior studies demonstrated

statistically significant relationships between BAL $CD8^+$ T cells, NK cells, eosinophils,

Tregs and outcomes in lung transplantation, but this has not led to meaningful

improvements in clinical care because wide overlaps in the frequency distributions of

these populations have meant that they lack discriminatory power in individual

patients[14,15,19,40-43]. In contrast, while $CD4^+CD57^+PD1^+$ T cells will require further

validation in larger studies, their prevalence among $CD4^+$ T cells discriminated patients

who remained stable from those about to undergo ALAD with high precision, in a

discovery cohort and confirmed in an independent validation cohort. Our observation

therefore represents a promising biomarker that may, in the future, permit the selection

of lung transplant recipients for targeted preventative interventions.

A review of clinical and laboratory variables present at the time of each bronchoscopy

did not reveal any clear association between the appearance of $\geq 7.8\%$ $CD57^+PD1^+$ T

cells in the BAL $CD4^+$ T cell compartment and other indicators of lung allograft

inflammation or dysfunction. This observation likely reflects the relatively poor

performance of histopathologic and cytologic findings in the diagnosis of lung allograft

rejection. Indeed, interpretation of transbronchial biopsy histology is fraught, with high

rates of ungradable biopsies and substantial inter-observer disagreement among

pathologists[7,8]. It is therefore perhaps not surprising that bronchoscopies with high

frequencies of $CD4^+CD57^+PD1^+$ T cells showed no particular association with histologic

rejection. Likewise, we found that this T cell population is not a marker of lung allograft

infection, since there were no differences in pathogen isolation from the same BAL

samples. Our interpretation of these findings is that analysis of BAL $CD4^+CD57^+PD1^+$ T

cells may dramatically improve the utility of surveillance bronchoscopy after LT, which

will require further evaluation in larger prospective studies.

Beyond their value as a biomarker of future allograft dysfunction, several lines of

evidence support the notion that CD4$^+$CD57$^+$PD1$^+$ T cells are mechanistically relevant

to the pathogenesis of chronic lung allograft rejection and fibrosis. First, our CITE-Seq

data indicate that the majority of clonally expanded CD4$^+$ T cell populations in the BAL

of patients with ALAD were CD57$^+$PD1$^+$, which indicates that these cells have

undergone antigen-driven expansion in the lung. Further, in addition to their appearance

in BAL, we detected CD4$^+$CD57$^+$PD1$^+$ T cells in human explanted CLAD lung tissue

obtained at the time of re-transplantation. The cells were observed in proximity to B

cells and small airways; coupled with the finding that the transcriptome and secretome

of BAL CD4$^+$CD57$^+$PD1$^+$ T cells includes IL-21 and granzymes, this observation

suggests that they are likely participating in allograft injury and may subserve Tph

functions. In support of this assertion, in our ischemia-reperfusion injury-augmented

mouse model of human CLAD, we observed CD4$^+$CD57$^+$PD1$^+$ T cells in the lung

allografts of mice with marked fibrotic pathology. In contrast, allografts with milder

fibrosis did not contain many CD4$^+$CD57$^+$PD1$^+$ T cells. Interestingly, we have previously

shown that fibrosis in this model is B cell dependent[27], and so in future work it will be

important to consider the possible existence of a CD4$^+$CD57$^+$PD1$^+$ T cell-B cell axis that

drives chronic lung allograft rejection in mice and humans.

Accumulating evidence has revealed an association of CD4$^+$CD57$^+$ T cells – with or

without associated PD-1 expression – with autoimmune diseases, infectious diseases,

and cancers [44]. CD4$^+$CD57$^+$ T cells have been detected in the peripheral blood of kidney

transplant recipients undergoing rejection [45], particularly in costimulation blockade

resistant rejection [46,47]. Of note, the CD4$^+$CD57$^+$ T cells associated with rejection in

kidney transplantation were measured in the PBMC and exhibit low expression of PD1.

In another study, Brugière and colleagues observed that the risk of CLAD was

increased in patients with a high proportion of circulating CD4$^+$CD57$^+$ILT2$^+$ T cells,

which could be inhibited by HLA-G[48]. However, in that study, PD-1 expression was not

assessed, and the association with CLAD was not as strong as we report here.

Therefore, it is unclear whether any of these peripheral CD4$^+$CD57$^+$ populations bear a

functional resemblance or developmental relationship to those that we have described
within lung allografts. Moreover, although peripheral blood CD57[+] T cells are also
reported to reflect CMV infection and/or age[49], our patients' characteristics did not
suggest any association between age and/or CMV seropositivity with CD4[+]CD57[+]PD1[+]
T cells.

PD1 expression by CD8[+] T cells in tumor microenvironments is associated with cancer
progression, and generally reflects a state of exhaustion in which effector functions are
deactivated. BAL CD4[+]CD57[+]PD1[+] T cells also expressed several additional molecules
that have been associated with exhaustion, including TIM-3, LAG3 and CTLA-4. While
analysis of the cells *ex vivo* revealed that they were both hypoproliferative and resistant
to T cell RICD, their production of inflammatory cytokines – which is not a feature of
exhaustion – suggests that they are senescent rather than exhausted. Of note,
senescent CD4[+] and CD8[+] T cells in humans were described as having the phenotype
CD45RA[+]KLRG1[+]CD57[+]CD28[-]CD27[-] [50,51]. Animal and human data indicate that
senescent T cells participate in other chronic inflammatory, degenerative and metabolic
disorders[39].

Lung transplant recipients typically receive immunosuppressive drugs – calcineurin
inhibitors, corticosteroids, and antimetabolites – in much higher doses than other solid
organ transplant recipients, and yet also have the highest rates of acute and chronic
allograft rejection. With respect to T cells, these drugs act by inhibiting cell proliferation.
In this context, we speculate that CD4[+]CD57[+]PD1[+] T cells, which exhibit reduced
proliferation, escape this inhibition and can continue to produce cytokines. Further work
will be required to understand how these cells arise within the lung allograft, and
whether specific targeting of these cells may have a therapeutic benefit.

Methods

Study design and BAL collection

The samples analyzed in this study were collected with informed consent for research

use and were approved by University Health Network Research Ethics Board (15-9531).

BAL cells from 50 TLTP lung transplant recipients presenting for surveillance

bronchoscopy at 3 months post-transplant were obtained according to a standard

protocol [52]. In addition, for each LT recipient, two further samples were obtained from

each recipient to form a longitudinal cohort **(Supp Fig 1)**. Cells were centrifuged at 400g

for 5 min and resuspended in 1 ml of 10% dimethyl sulfoxide/90% FBS for viably

cryopreservation. Cryovials were placed in Mr. Frosty containers at -80^oC for 24 h,

then stored in liquid nitrogen until thawing for processing and mass cytometry. BAL cells

from a further 14 LT recipients with acute lung allograft dysfunction (ALAD, defined as a

decrease in the forced expiratory volume in one second [FEV1] by 10% or more from

the baseline FEV1 measurements) were used for sorting fresh $CD4^+CD57^+PD1^+$ T cells

and their $CD57^-PD1^-$ counterparts for functional analyses.

Mass cytometry

Metal-conjugated antibodies were purchased either preconjugated from Fluidigm or

purified from BioLegend, Thermo Fisher Scientific or BD Biosciences and subsequently

conjugated to metals using Maxpar Antibody Labeling Kits (Fluidigm) according to the

manufacturer's instructions **(Supp Table 1)**. Cryopreserved BAL cells were recovered

and divided into stimulated (with PMA 50ng/mL and ionomycin 500ng/mL) and

unstimulated conditions to allow analysis of cytokine secretion expression and surface

markers. Cells were incubated with a cocktail of antibodies against cell surface

molecules for 30 min at room temperature. After treatment with intracellular fixation

and permeabilization buffers (BD Biosciences), cells were incubated with metal-

conjugated antibodies against intracellular proteins. Cells were then washed and

stained with Intercalator-Ir (Fluidigm) diluted in PBS containing 1.6% paraformaldehyde

and stored at 4 °C until next-day acquisition. To assess variation between mass

cytometry runs, a control healthy volunteer PBMC sample was aliquoted, recovered and

stained and analyzed alongside the BAL samples.

Mass cytometry data analysis

Mass cytometry data were initially analyzed with manual gating to exclude debris, dead cells and doublets and to select CD45[+] cells using Cytobank software (Beckman Coulter). We then used the FlowSOM and viSNE algorithms (5,000 iterations, perplexity$\square$=$\square$100) for visualization of high-dimensional data [21,53]. Subsequently, we applied cluster identification, characterization, and regression (CITRUS) [22] on the CD3+ population from the discovery cohort (n=25, **Supp Table 2**) to identify cell subsets associated with ALAD within 30 days of the sample. CITRUS was run at least 5 times with "equal number" of cells per sample, looking for "abundance" as cluster characterization, with a 1% false discovery rate. To incorporate the time-course design of our study, we implemented a generalized linear mixed model analysis-adapted R pipeline as described by Nowicka and colleagues [23]. In this approach, different timepoints for each LT recipient were defined as a variable to allow intra-subject pairing. Finally, we used manual gating to confirm the results of CITRUS and generalized linear mixed model analysis.

Imaging Mass Cytometry

Lung tissue samples were obtained during re-transplantation procedures from 8 patients who were undergoing re-transplantation due to end-stage CLAD. A single biopsy was obtained from the upper lobe of each explanted sample. The harvested lung tissue was inflated and immersed in a 10% formalin solution for 24 hours prior to immersion in a 70% ethanol solution for an additional 24-hour period. Samples were then embedded in paraffin. Four-micron thick sections of paraffin-embedded lung tissue were prepared and affixed to slides for further analysis. For every specimen, we identified three distinct 1x1mm Regions of Interest (ROIs) situated around individual airways. In addition to the CLAD patient samples, we included control samples in our study. Controls included lung tissue samples from two lobectomy procedures for suspected cancer and excess tissue from two donor lungs collected at the end of the cold ischemic phase of preservation. Sections of CLAD and control lung tissue were stained with a panel of 35 heavy-metal-tagged antibodies **(Supp Table 6)**. Subsequently, the three ROIs chosen for each specimen were subjected to ablation using the Hyperion Imaging System (Fluidigm)

400 using a laser frequency of 200Hz, with laser power ranging from 3.5 to 4.5. The ablation
401 procedure generated files in both .txt and .mcd formats, which were subsequently
402 subjected to thorough analysis using the HistoCAT software platform[54].
403
Animal model
Orthotopic lung transplantation was performed using C57BL/10 donors and C57BL/6
recipients under different graft preservation conditions as described previously [27]. Briefly,
harvested lungs were stored either under minimal (60 minutes at 4ºC followed by 15 min
of warm anastomotic time) or prolonged (6h at 4ºC, 45 min at 37ºC, followed by 15 min
of warm anastomotic time) storage prior to reperfusion. Orthotopic lung transplant
recipients were sacrificed 28 days after LT for histological analysis and cellular
phenotyping. Grafts were flushed with saline and digested with collagenase A and
DNase I in a gentleMACS tissue dissociator (Miltenyi Biotec). Cells were stained with
cocktails of fluorochrome-conjugated monoclonal antibodies including AF647-
conjugated anti-mouse CD57 (polyclonal; Bioss) and AF488-conjugated anti-mouse
PD1 (polyclonal; Bioss). Flow cytometry data were analyzed using FlowJo (v10, FlowJo
LLC) and correlated with histological grading. We graded peribronchial fibrosis and
parenchymal fibrosis in a blinded fashion according to a published scoring system[28].
Immunofluorescent staining of sections of formalin-fixed and paraffin-embedded lung
tissue was conducted by antigen retrieval using 1x pH 9 Tris/EDTA buffer before
staining with unconjugated anti-mouse CD4. This was followed by a secondary antibody
cocktail including Alexa Fluor 555 anti-rabbit IgG (A-31572, Invitrogen) as well as
AF647-conjugated anti-mouse CD57 (polyclonal; Bioss) and AF488-conjugated anti-
mouse PD1 (polyclonal; Bioss). Nuclei were stained with DAPI and
immunofluorescence images were acquired using a Zeiss AxioObserver.
425
CITE-seq sample processing and acquisition
We identified four LT recipients scheduled for bronchoscopy with ALAD (FEV_1 decrease
by ≥10% from the prior value). Fresh BAL cells were collected by centrifugation at 400g
for 5 minutes. T cells were obtained using a $CD3^+$ T cell enrichment kit (EasySep™
Human T Cell enrichment, Stemcell Technologies) according to the manufacturer's

instructions. Enriched T cells were resuspended at 1×10^6 cells/mL in PBS + 0.05% BSA.

50 µL of the cell suspension were stained with BV711 conjugated anti-human CD4,

APC conjugated anti-human PD1 and PE-Cy7 anti-human CD57 and analyzed by flow

cytometry to confirm ≥ 7.8% $CD57^+PD1^+$ cells in the BAL $CD4^+$ T cell compartment. If

confirmed, the remaining cells were used for CITE-seq. Briefly, 1×10^6 cells were

resuspended in 45 µL PBS + 0.05% BSA before adding 5 µL of Fc blocking reagent and

incubated for 10 minutes at 4°C. TotalSeq™-C antibody cocktails were prepared during

the incubation by adding pre-determined concentrations of each antibody **(Supp Table**

6). The resulting cocktail was then added to cells and incubated for 30 minutes at 4°C.

After incubation, the cell pellets were resuspended in 1mL PBS + 0.05% BSA and

centrifuged for 5 minutes at 400 x g. After a total of 3 washes, the final cell density was

adjusted to 1000-1200 cells/µL and subjected to the 10X Genomics Chromium 5'

Immune Profiling v2.0 protocol for library preparation. A total of 2×10^4 cells from each

sample were loaded onto a 10X single cell G chip. After droplet generation, samples

were transferred to a pre-chilled 96 well plate (Eppendorf), heat sealed and incubated

overnight in a Veriti 96-well thermocycler (Thermo Fisher). The next day, sample cDNA

was recovered using Recovery Agent (10X Genomics) and subsequently cleaned up

using Silane DynaBeads (Thermo Fisher) mix as outlined by the user guide 5' Immune

Profiling v2.0. Purified cDNA was amplified for 11 cycles before being cleaned up using

SPRIselect beads (Beckman Coulter). Samples were diluted 9:1 (elution buffer

(Qiagen):cDNA) and run on a Bioanalyzer (Agilent Technologies) to determine cDNA

concentration. The library was then sequenced using an Illumina Novaseq 6000

sequencer using the following parameters Read 1: 28b, index 1: 10bp, Index 2: 10bp,

Read 2: 90bp.

CITE-seq analysis

The sequencing output was first demultiplexed and converted into FASTQ files using

Cell Ranger [55]. Reads were aligned to the human reference genome (grch38), and

count UMIs in the mRNA libraries, and CITE-seq-Count to count UMIs in the antibody-

derived tag (ADT) libraries. The matrix, barcodes, and gene information from Cell

Ranger were then used to create gene-cell matrices and a Seurat object [56]. For each

Seurat object, cells with poor quality were excluded based on the following quality

control metrics: unique gene counts over 5000 or below 200, UMI counts over 30,000,

and mitochondrial gene expression higher than 10%. Each sample was scaled and

normalized using Seurat's "SCTransform" function followed by adding the protein

expression levels to the Seurat object and normalization and scaling for ADT assay.

Next, CD4$^+$ T cells were subsetted based on high CD4 and low CD8 protein expression

(ADT-CD8 < 12 & ADT-CD4 > 12). Seurat objects of CD4+ cells from all four BAL

samples were merged into a single Seurat object. The SCTransform method was used

for normalization, scaling, and integration which corrects for unwanted sources of

variation and enhances the signal of genuine biological differences between cells[57].

Then, dimensionality reduction was applied to the integrated Seurat object, allowing

visualization and interpretation of the high-dimensional scRNAseq data. This was

followed by unsupervised clustering (resolution = 0.5) which led to the identification of

10 clusters. Then cells with higher expression of protein level of CD57 and PD1 were

highlighted on tSNE plot.

RNA velocity estimates the future state of individual cells by calculating the ratio of

unspliced to spliced mRNA for each gene. The spliced and unspliced counts from the

scRNAseq data were utilized to calculate RNA velocity using the velocyto package in

Python[58]. Splicing information extracted from velocyto was then added to all the

selected cells in the CD4$^+$ Seurat object. Subsequently, CD4$^+$ T cells underwent

pseudotime trajectory analysis using the scVelo pipeline[59]. This analysis inferred the

trajectory and the chronological sequence of cell states, highlighting the differentiation

paths taken by cells. The pseudotime values were visualized on the tSNE plots,

providing insights into the dynamic processes and cellular transitions at play.

Fluorescence-activated cell sorting

BAL cells collected from 15 ALAD samples and stained with fluorophore-tagged surface

antibodies for 30□min at room temperature. The following antibodies were used to stain

the cells: BV711-conjugated anti-human CD4 (BioLegend); APC-conjugated anti-human

PD1 (BioLegend); PE/Cy7-conjugated anti-human CD57, in addition to biotinylated

CD163, CD14, CD8, and CD56 (all from BioLegend). Cells washed with 500 µL of 1%

FBS in PBS and incubated with streptavidin-AF488 (1:200, BioLegend) for 20 minutes,

before washing and staining with propidium iodide to identify dead cells. Stained cells

were subjected to FACS sorting using a Sony SH800 instrument. After exclusion of PI-

positive cells and putative doublets based on forward and side scatter analysis, cells in

the dump channel (AF488) were eliminated. The remaining cells were gated to select

CD4$^+$ T cells and sorted for CD57$^+$PD1$^+$ and CD57$^-$PD1$^-$ cells.

Cell culture, multiplex protein analysis, and *in vitro* assays

Sorted cells were labeled with 1µM CFSE in serum free PBS prior to wash and culture

with plate bound anti-CD3 (5µM) and soluble anti-CD28 (1µM) antibodies in DMEM

medium containing 5.55 mM D-glucose, 4 mM L-glutamine, and 1 mM sodium pyruvate,

supplemented with 10% FBS, 50 U/ml penicillin, and 50 mg/ml streptomycin and 500

U/ml human recombinant IL-2 (Proleukin, Chiron). After 18 hours, 100µL supernatant

was replaced with fresh media. The collected supernatant was stored at -80^{o}C for later

multiplex analysis. Supernatants were thawed and subjected to a custom multiplex bead

kit (R&D Systems) to assess the levels of released analytes including IL-21, IL-4,

CXCL8, and granzyme A. Samples were prepared in a 1:1 dilution with assay buffer, as

suggested by the manufacturer. Diluted samples and standards were run in duplicate.

Biomarker concentrations were obtained using a Bio-Plex MAGPIX Multiplex Reader

(Bio-Rad). All samples and standards were run in duplicate according to manufacturers'

protocols.

After 5 days, CFSE dilution in the sorted cells – a measure of T cell proliferation – was

determined by flow cytometry. Finally, a restimulation-induced cell death assay[60] was

used to assess whether CD57$^+$PD1$^+$ cells are less susceptible to this mode of cell death

than their CD57$^-$PD1$^-$ counterparts. CD57$^+$PD1$^+$ and CD57$^-$PD1$^-$ cells were stimulated in

culture for 7 days before washing, and were then restimulated with anti-CD3 mAb

(OKT3, 100 ng/ml final) for 24 hours. Cells were then harvested and stained with

Annexin V and propidium iodide for evaluation of apoptotic cell death.

Statistics

All statistical tests were two-sided unless stated otherwise. The Wilcoxon rank-sum test was used to assess statistical differences between two groups. When assessing >2 groups simultaneously, the nonparametric Kruskal–Wallis test was used. A Fisher's exact test was applied to assess statistical differences between two categorical variables. Receiver Operating Characteristic (ROC) curves were used to assess classification accuracy, which was quantified by Area Under the Curve (AUC). Kaplan–Meier plots and two-sided log-rank tests were used to compare survival probabilities between groups. Cox Proportional Hazards (PH) regression models were used to adjust for potential confounders for time-to-CLAD or time-to-death/re-transplantation analysis. and Cox regression analyses were used to assess covariates with respect to time-to-allograft dysfunction. All statistical analyses were performed using R v.3.5.1+ or Prism 8+ (GraphPad Software).

Figure 1. Identification of a BAL CD3$^+$ T cell population associated with lung

allograft dysfunction. A. Study schematic. Fifty consecutive patients presenting for

bronchoscopy at 3 months post-transplant were enrolled. For each enrolled patient, BAL

cells at three time points (3mo, 6mo, 9mo or next available) were collected. BAL cells

were cryopreserved and subjected to mass cytometry (see supplementary table 1); all

samples from each patient were analyzed in the same mass cytometry experiment. **B.** t-

distributed stochastic neighbour embedding (tSNE) plots of BAL CD45$^+$ cells falling into

20 clusters (highlighted in different colours) identified by FlowSOM. Inset tSNE plot

highlighting expression of CD3$^+$ T cells is shown at lower right. **C.** t-SNE plot based on

arcsinh-transformed marker expression in 500 randomly selected cells per sample from

the discovery set (n=25, all 3 time points shown). Cells are coloured according to the 20

cell populations obtained with FlowSOM after the metaclustering step with

ConsensusClusterPlus. **D.** Differential analysis heatmap of arcsine-square-root

transformed cell frequencies that were subsequently normalized per cluster (rows, 20

clusters identified in the panel C). Statistically significant clusters are highlighted by

green boxes at right along with p-values. **E.** Analysis of molecules expressed by all cells

(average) compared with those expressed by the most differentially represented cell

population in ALAD patients (cluster 11) reveals that the latter is

CD45$^+$CD3$^+$CD4$^+$CD57$^+$PD1$^+$.

Figure 2. Relationship of BAL CD4$^+$CD57$^+$PD1$^+$ T cells with incident allograft

dysfunction. Representative examples of forced expiratory volume in one second

(FEV1) over time in weeks post-transplant for stable (**A**) and ALAD (**B**) lung transplant

recipients in the discovery cohort. Weeks during which bronchoscopy was performed

are highlighted in yellow; scatterplots showing expression of CD57 and PD1 in the CD4$^+$

T cell compartment of the BAL are shown for each time point. Green line indicates 80%

of best post-transplant baseline, below which CLAD may be diagnosed. **C.** Receiver-

operator characteristic (ROC) curve analysis of the performance characteristics of

CD57$^+$PD1$^+$ cells as a proportion of the BAL CD4$^+$ compartment for diagnoses of

subsequent ALAD. The area under the curve (AUC) = 0.76. The percentage of

CD57$^+$PD1$^+$ T cells providing optimal discrimination of ALAD and stable patients (7.8%)

is circled in red. **D.** The highest proportion of CD57$^+$PD1$^+$ T cells observed in all BAL

samples from each patient (discovery cohort, n=25) is shown, divided by the presence

or absence of ALAD within 30 days. Mann-Whitney test p=0.0018. Dotted line shows

the 7.8% threshold identified in ROC curve in C. **E.** Time to lung allograft dysfunction

from the emergence of CD57$^+$PD1$^+$ cells $\geq$ 7.8% of CD4$^+$ T cells in the BAL (red line), or

from the final BAL analyzed (blue line) in the discovery cohort (n=25). Log rank test p <

0.0001. **F.** Time to lung allograft dysfunction from the emergence of CD57$^+$PD1$^+$ cells $\geq$

7.8% of CD4$^+$ T cells in the BAL (red line), or from the final BAL analyzed (blue line) in

the validation cohort (n=25). Log rank test p < 0.02. G. Of 28 patients who developed de

novo DSA, 19 (67.9%) had CD57$^+$PD1$^+$ cells $\geq$ 7.8% of CD4$^+$ T cells in at least one BAL

sample; only 9 of 23 (39.1%) patients who did not develop de novo DSA had

CD57$^+$PD1$^+$ cells $\geq$ 7.8% of CD4$^+$ T cells in at least on BAL sample. Chi square test p =

0.0402.

Figure 3. CD4$^+$CD57$^+$PD1$^+$ T cells are associated with an increased risk of death or

re-transplantation and are found in CLAD lung tissue. A. Kaplan-Meier curves

showing time to death or re-transplantation from 12 months post-transplant. Log-rank

test p = 0.00079. **B.** Selected heatmap of protein expression for selected clusters and

markers to highlight protein profile of cluster 50. **C.** Imaging mass cytometry of normal

donor lung tissue (left) and CLAD lung tissue obtained at the time of re-transplantation

(right). E-cadherin$^+$ epithelial cells (green pseudocolour), red CD3$^+$ T cells (red

pseudocolour), CD4$^+$ cells (blue pseudocolour), and CD4$^+$CD57$^+$PD1$^+$ T cells (cluster 50,

pale blue pseudocolour) are shown. **D.** CD4$^+$CD57$^+$PD1$^+$ T cells are more numerous in

CLAD lung tissue. Data from 3 regions of interest (ROIs) from each of n=8 CLAD, and

n=4 normal lungs are shown. **E.** In CLAD lung tissue, CD4$^+$CD57$^+$PD1$^+$ T cells (cluster

50, white) are found in association with airways (green pseudocolour E-cadherin$^+$ cells)

and with two distinct B cell populations (cluster 8, pale blue pseudocolour and cluster 30,

purple pseudocolour). **F.** Cell-cell interaction heatmap shows association of

CD4$^+$CD57$^+$PD1$^+$ T cells (cluster 50) with epithelial cells (cluster 27) and B cells

(clusters 8 and 30). G. Liquid phase mass cytometry performed on single cell

suspensions obtained from CLAD lung tissue show variable presence of

CD4$^+$CD57$^+$PD1$^+$ T cells.

Figure 4. CD4$^+$CD57$^+$PD1$^+$ T cells are associated with fibrosis in a mouse model of chronic lung allograft rejection. A. C57BL/10 (B10) donor lungs were subjected to either prolonged (6h cold ischemic time (CIT) at 4°C followed by 1h warm ischemic time (WIT) at 37°C) or minimal (60min CIT followed by 15min WIT) storage prior to orthotopic left lung transplantation into minor alloantigen-mismatched C57BL/6 (B6) recipients. **B.** After 28 days, paraffin-embedded sections of lung tissue were subjected to Masson trichrome staining (200x magnification). **C.** Grafts were scored using a published scoring system[28] that quantifies the percentage of the graft affected by interstitial inflammation and parenchymal fibrosis. Grafts with high-grade pathology (High Path) are represented as triangles, and those with minimal pathology (Low Path) are represented as circles. **D.** Single-cell suspensions of naïve B6 lungs (No OLT) and lung allografts obtained at 28 days post-transplantation were subjected to flow cytometry. Analyses are gated on CD4$^+$ T cells. Percentages of cells in the CD57$^+$PD-1$^+$ gate are shown in the representative plots. **E.** Percentage of CD4$^+$ T cells expressing CD57 and PD-1 is shown for naïve B6 lungs (No OLT, squares), Low Path grafts (circles) and High Path grafts (triangles). Student's t test, p=0.0005. **F.** Percentage of CD4$^+$ T cells expressing CD57 and PD-1 is shown for Low Path grafts (circles) and High Path grafts (triangles). Student's t test, p=0.0205. **G.** Immunofluorescence of prolonged storage, High Path lung allograft showing PD1 (green), CD57 (red), CD4 (yellow) and DAPI (blue) and merged image.

Figure 5. Identification of distinct CD27$^+$ and CD27$^-$ subsets of CD4$^+$CD57$^+$PD1$^+$ T cells. CITE-Seq was performed on 4 BAL samples from patients with ALAD. **A.** After downsampling and integration, BAL CD4$^+$ T cells (CD3$^+$CD4$^+$CD8$^-$ identified by antibody-derived oligonucleotide tag, ADT) were displayed on a tSNE plot revealing 10 distinct cell clusters (left tSNE plot). Right panel displays CD57 and PD-1 expression (blue dots) on a feature plot showing all BAL CD4$^+$ cells. **B.** tSNE plot demonstrating CD27 RNA expression (left) and at the protein level (ADT-CD27, right). **C.** Dot plot of the selected top differentially expressed genes in BAL CD27$^+$ and CD27$^-$ CD4$^+$CD57$^+$PD-1$^+$ T cells. Purple colour indicates higher expression; dot size reflects the percentage of cells in each population expressing the gene. **D.** tSNE plot overlaid with colours reflecting TCR clonal expansion. Dotted red circle indicates highly

expanded clones within the CD27⁻ population.. **E.** tSNE plot overlaid with RNA velocity

trajectories (arrows). Dotted red circle shown in D is overlaid here indicating the likely

origin of CD4$^+$CD57$^+$PD-1$^+$ T cells.

Figure 6. Functional properties of CD4$^+$CD57$^+$PD1$^+$ T cells. A. CD4$^+$CD57$^+$PD1$^+$ T

cells can be identified by conventional flow cytometry. BAL cells were stained with CD4,

CD57 and PD1 antibodies and CD57$^+$PD1$^+$ cells were identified within the CD4$^+$

population, allowing quantification and cell sorting. **B**. Percentage of CD57$^+$PD1$^+$ cells

within the CD4$^+$ gate is shown for n=14 patients with ALAD. **C.** Sorted CD57$^+$PD1$^+$ and

CD57$^-$PD1$^-$ CD4$^+$ T cells from ALAD BAL samples were stimulated *in vitro* with anti-

CD3/anti-CD28 antibodies overnight. Culture supernatants were subjected to

multiplexed cytokine quantification. IL-21 (p=0.031), IL-4 (p=0.062), CXCL8 (IL-8,

p=0.031) and GZMA (p=0.062) were generally secreted in greater concentrations by

CD57$^+$PD1$^+$ T cells than their CD57$^-$PD1$^-$ counterparts, Wilcoxon matched-pairs signed

rank tests. **D.** Sorted CD57$^+$PD1$^+$ (blue) and CD57$^-$PD1$^-$ CD4$^+$ (pink) T cells from ALAD

BAL were examined for proliferation in response to CD3/CD28 stimulation by dilution of

CFSE. **E.** The percentage of CD57$^+$PD1$^+$ T cells undergoing proliferation was lower

compared to that of their CD57$^-$PD1$^-$ counterparts. Linked CD57$^+$PD1$^+$ and CD57$^-$PD1$^-$

cells are sorted from same LT recipient. Wilcoxon matched pairs signed rank test,

p=0.0041. **F.** Sorted CD57$^+$PD1$^+$ T cells (right) and their CD57$^-$PD1$^-$ counterparts (left)

were subjected to RICD. Annexin V and propidium iodide (PI) staining is shown. **G.**

Annexin V staining was lower in CD57$^+$PD1$^+$ T cells compared to their CD57$^-$PD1$^-$

counterparts. Linked CD57$^+$PD1$^+$ and CD57$^-$PD1$^-$ T cells are sorted from same LT

recipient. Wilcoxon matched pairs signed rank test, p=0.0159.

References

650   1   Ali, J. M., Bolton, E. M., Bradley, J. A. & Pettigrew, G. J. Allorecognition pathways in
651   transplant rejection and tolerance. *Transplantation* **96**, 681-688,
652   doi:10.1097/TP.0b013e31829853ce (2013).

653   2   Chambers, D. C. *et al.* The International Thoracic Organ Transplant Registry of the
654   International Society for Heart and Lung Transplantation: Thirty-eighth adult lung
655   transplantation report - 2021; Focus on recipient characteristics. *J Heart Lung Transplant*
656   **40**, 1060-1072, doi:10.1016/j.healun.2021.07.021 (2021).

657   3   Kawashima, M. & Juvet, S. C. The role of innate immunity in the long-term outcome of
658   lung transplantation. *Ann Transl Med* **8**, 412, doi:10.21037/atm.2020.03.20 (2020).

659   4   Bando, K. *et al.* Obliterative bronchiolitis after lung and heart-lung transplantation. An
660   analysis of risk factors and management. *The Journal of thoracic and cardiovascular*
661   *surgery* **110**, 4-13; discussion 13-14 (1995).

662   5   Heng, D. *et al.* Bronchiolitis obliterans syndrome: incidence, natural history, prognosis,
663   and risk factors. *J Heart Lung Transplant* **17**, 1255-1263 (1998).

664   6   Moshkelgosha, S. *et al.* Interferon-stimulated and metallothionein-expressing
665   macrophages are associated with acute and chronic allograft dysfunction after lung
666   transplantation. *J Heart Lung Transplant* **41**, 1556-1569,
667   doi:10.1016/j.healun.2022.05.005 (2022).

668   7   Bhorade, S. M. *et al.* Interobserver variability in grading transbronchial lung biopsy
669   specimens after lung transplantation. *Chest* **143**, 1717-1724, doi:10.1378/chest.12-2107
670   (2013).

671   8   Stephenson, A. *et al.* Interpretation of transbronchial lung biopsies from lung transplant
672   recipients: inter- and intraobserver agreement. *Can Respir J* **12**, 75-77 (2005).

673   9   Levy, L. *et al.* in *International Society for Heart and Lung Transplantation 39th Annual*
674   *Meeting* Vol. 38(S4) S163-164 (J Heart Lung Transplant, Orlando FL USA, 2019).

675   10   Levy, L. *et al.* Epithelial cell death markers in bronchoalveolar lavage correlate with
676   chronic lung allograft dysfunction subtypes and survival in lung transplant recipients-a
677   single-center retrospective cohort study. *Transpl Int* **32**, 965-973, doi:10.1111/tri.13444
678   (2019).

679   11   Moshkelgosha, S. *et al.* in *J Heart Lung Transpl* Vol. 37 S216 (2018).

680   12   Moshkelgosha, S. *et al.* in *International Society for Heart and Lung Transplantation* Vol.
681   38 S17 (Orlando, FL USA, 2019).

682   13   Moshkelgosha, S. *et al.* in *J Heart Lung Transpl* Vol. 39 S199-S200 (2020).

683   14   Whitehead, B. F. *et al.* Analysis of bronchoalveolar lavage from human lung transplant
684   recipients by flow cytometry. *Respiratory medicine* **89**, 27-34 (1995).

685   15   Reynaud-Gaubert, M. *et al.* Clinical utility of bronchoalveolar lavage cell phenotype
686   analyses in the postoperative monitoring of lung transplant recipients. *European journal*
687   *of cardio-thoracic surgery : official journal of the European Association for Cardio-*
688   *thoracic Surgery* **21**, 60-66 (2002).

689   16   Hodge, S. *et al.* Increased levels of T cell granzyme b in bronchiolitis obliterans syndrome
690   are not suppressed adequately by current immunosuppressive regimens. *Clinical and*
691   *experimental immunology* **158**, 230-236, doi:10.1111/j.1365-2249.2009.04008.x (2009).

17   Neujahr, D. C. *et al.* Dynamics of human regulatory T cells in lung lavages of lung
transplant recipients. *Transplantation* **88**, 521-527, doi:10.1097/TP.0b013e3181b0e719
(2009).
18   Hodge, S. J., Hodge, G. L., Reynolds, P. N. & Holmes, M. D. Differential rates of apoptosis
in bronchoalveolar lavage and blood of lung transplant patients. *J Heart Lung Transplant*
24, 1305-1314, doi:10.1016/j.healun.2004.10.009 (2005).
19   Greenland, J. R. *et al.* Bronchoalveolar lavage cell immunophenotyping facilitates
diagnosis of lung allograft rejection. *Am J Transplant* **14**, 831-840, doi:10.1111/ajt.12630
(2014).
20   Bandura, D. R. *et al.* Mass cytometry: technique for real time single cell multitarget
immunoassay based on inductively coupled plasma time-of-flight mass spectrometry.
Analytical chemistry **81**, 6813-6822, doi:10.1021/ac901049w (2009).
21   Van Gassen, S. *et al.* FlowSOM: Using self-organizing maps for visualization and
interpretation of cytometry data. *Cytometry A* **87**, 636-645, doi:10.1002/cyto.a.22625
(2015).
22   Bruggner, R. V., Bodenmiller, B., Dill, D. L., Tibshirani, R. J. & Nolan, G. P. Automated
identification of stratifying signatures in cellular subpopulations. *Proc Natl Acad Sci U S A*
111, E2770-2777, doi:10.1073/pnas.1408792111 (2014).
23   Nowicka, M. *et al.* CyTOF workflow: differential discovery in high-throughput high-
dimensional cytometry datasets. *F1000Res* **6**, 748, doi:10.12688/f1000research.11622.2
(2017).
24   Tikkanen, J. M. *et al.* De Novo DQ Donor-Specific Antibodies Are Associated with Chronic
Lung Allograft Dysfunction after Lung Transplantation. *American journal of respiratory*
and critical care medicine **194**, 596-606, doi:10.1164/rccm.201509-1857OC (2016).
25   Renaud-Picard, B. *et al.* in *J Heart Lung Transpl* Vol. 41   S106 (2022).
26   Giesen, C. *et al.* Highly multiplexed imaging of tumor tissues with subcellular resolution
by mass cytometry. *Nature methods* **11**, 417-422, doi:10.1038/nmeth.2869 (2014).
27   Watanabe, T. *et al.* A B cell-dependent pathway drives chronic lung allograft rejection
after ischemia-reperfusion injury in mice. *American journal of transplantation : official*
journal of the American Society of Transplantation and the American Society of
Transplant Surgeons **19**, 3377-3389, doi:10.1111/ajt.15550 (2019).
28   Martinu, T. *et al.* Spectrum of chronic lung allograft pathology in a mouse minor-
mismatched orthotopic lung transplant model. *Am J Transplant* **19**, 247-258,
doi:10.1111/ajt.15167 (2019).
29   Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single
cells. *Nature methods* **14**, 865-868, doi:10.1038/nmeth.4380 (2017).
30   Rao, D. A. *et al.* Pathologically expanded peripheral T helper cell subset drives B cells in
rheumatoid arthritis. *Nature* **542**, 110-114, doi:10.1038/nature20810 (2017).
31   Manzo, A. *et al.* Mature antigen-experienced T helper cells synthesize and secrete the B
cell chemoattractant CXCL13 in the inflammatory environment of the rheumatoid joint.
Arthritis Rheum **58**, 3377-3387, doi:10.1002/art.23966 (2008).
32   Huang, Y. *et al.* T peripheral helper cells in autoimmune diseases: What do we know?
Front Immunol **14**, 1145573, doi:10.3389/fimmu.2023.1145573 (2023).

33 Ng, S. S. *et al.* The NK cell granule protein NKG7 regulates cytotoxic granule exocytosis
and inflammation. *Nat Immunol* **21**, 1205-1218, doi:10.1038/s41590-020-0758-6 (2020).
34 Wein, A. N. *et al.* CXCR6 regulates localization of tissue-resident memory CD8 T cells to
the airways. *J Exp Med* **216**, 2748-2762, doi:10.1084/jem.20181308 (2019).
35 Milardi, G. *et al.* Follicular helper T cell signature of replicative exhaustion, apoptosis,
and senescence in common variable immunodeficiency. *Eur J Immunol* **52**, 1171-1189,
doi:10.1002/eji.202149480 (2022).
36 Llao-Cid, L. *et al.* EOMES is essential for antitumor activity of CD8(+) T cells in chronic
lymphocytic leukemia. *Leukemia : official journal of the Leukemia Society of America,*
Leukemia Research Fund, U.K **35**, 3152-3162, doi:10.1038/s41375-021-01198-1 (2021).
37 Wolf, Y., Anderson, A. C. & Kuchroo, V. K. TIM3 comes of age as an inhibitory receptor.
Nature reviews. Immunology **20**, 173-185, doi:10.1038/s41577-019-0224-6 (2020).
38 Liu, X., Hoft, D. F. & Peng, G. Senescent T cells within suppressive tumor
microenvironments: emerging target for tumor immunotherapy. *J Clin Invest* **130**, 1073-
1083, doi:10.1172/JCI133679 (2020).
39 Mittelbrunn, M. & Kroemer, G. Hallmarks of T cell aging. *Nat Immunol* **22**, 687-698,
doi:10.1038/s41590-021-00927-z (2021).
40 Hodge, G., Hodge, S., Reynolds, P. N. & Holmes, M. Airway infection in stable lung
transplant patients is associated with decreased intracellular T-helper type 1 pro-
inflammatory cytokines in bronchoalveolar lavage T-cell subsets. *Transplant infectious*
disease : an official journal of the Transplantation Society **10**, 99-105,
doi:10.1111/j.1399-3062.2007.00236.x (2008).
41 Hodge, G., Hodge, S., Chambers, D., Reynolds, P. N. & Holmes, M. Acute lung transplant
rejection is associated with localized increase in T-cell IFNgamma and TNFalpha
proinflammatory cytokines in the airways. *Transplantation* **84**, 1452-1458,
doi:10.1097/01.tp.0000290679.94163.e1 (2007).
42 Hodge, G., Hodge, S., Reynolds, P. N. & Holmes, M. Compartmentalization of
intracellular proinflammatory cytokines in bronchial intraepithelial T cells of stable lung
transplant patients. *Clinical and experimental immunology* **145**, 413-419,
doi:10.1111/j.1365-2249.2006.03143.x (2006).
43 Hodge, G., Hodge, S., Reynolds, P. N. & Holmes, M. Increased intracellular pro- and anti-
inflammatory cytokines in bronchoalveolar lavage T cells of stable lung transplant
patients. *Transplantation* **80**, 1040-1045 (2005).
44 Focosi, D., Bestagno, M., Burrone, O. & Petrini, M. CD57+ T lymphocytes and functional
immune deficiency. *J Leukoc Biol* **87**, 107-116, doi:10.1189/jlb.0809566 (2010).
45 Shaw, B. I. *et al.* Functional Characteristics and Phenotypic Plasticity of CD57(+)PD1(-)
CD4 T Cells and Their Relationship with Transplant Immunosuppression. *J Immunol* **206**,
1668-1676, doi:10.4049/jimmunol.2000736 (2021).
46 Sun, H. *et al.* TIGIT regulates apoptosis of risky memory T cell subsets implicated in
belatacept-resistant rejection. *Am J Transplant* **21**, 3256-3267, doi:10.1111/ajt.16571
(2021).
47 Espinosa, J. *et al.* CD57(+) CD4 T Cells Underlie Belatacept-Resistant Allograft Rejection.
Am J Transplant **16**, 1102-1112, doi:10.1111/ajt.13613 (2016).

48   Brugiere, O. *et al.* Chronic lung allograft dysfunction is associated with an early increase
of circulating cytotoxic CD4+CD57+ILT2+ T cells, selectively inhibited by the immune
check-point HLA-G. *J Heart Lung Transplant* **41**, 626-640,
doi:10.1016/j.healun.2022.01.013 (2022).
49   Hassouneh, F. *et al.* Functional Changes of T-Cell Subsets with Age and CMV Infection.
Int J Mol Sci **22**, doi:10.3390/ijms22189973 (2021).
50   Libri, V. *et al.* Cytomegalovirus infection induces the accumulation of short-lived,
multifunctional CD4+CD45RA+CD27+ T cells: the potential involvement of interleukin-7
in this process. *Immunology* **132**, 326-339, doi:10.1111/j.1365-2567.2010.03386.x
(2011).
51   Di Mitri, D. *et al.* Reversible senescence in human CD4+CD45RA+CD27- memory T cells. *J*
Immunol **187**, 2093-2100, doi:10.4049/jimmunol.1100978 (2011).
52   Martinu, T. *et al.* International Society for Heart and Lung Transplantation consensus
statement for the standardization of bronchoalveolar lavage in lung transplantation. *J*
Heart Lung Transplant, doi:10.1016/j.healun.2020.07.006 (2020).
53   Amir el, A. D. *et al.* viSNE enables visualization of high dimensional single-cell data and
reveals phenotypic heterogeneity of leukemia. *Nat Biotechnol* **31**, 545-552,
doi:10.1038/nbt.2594 (2013).
54   Schapiro, D. *et al.* histoCAT: analysis of cell phenotypes and interactions in multiplex
image cytometry data. *Nat Methods* **14**, 873-876, doi:10.1038/nmeth.4391 (2017).
55   Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells.
Nature communications **8**, 14049, doi:10.1038/ncomms14049 (2017).
56   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nature*
biotechnology **36**, 411-420, doi:10.1038/nbt.4096 (2018).
57   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
seq data using regularized negative binomial regression. *Genome Biology* **20**, 296,
doi:10.1186/s13059-019-1874-1 (2019).
58   La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498,
doi:10.1038/s41586-018-0414-6 (2018).
59   Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to
transient cell states through dynamical modeling. *Nature biotechnology* **38**, 1408-1414,
doi:10.1038/s41587-020-0591-3 (2020).
60   Katz, G. & Snow, A. L. Fluorescence-activated cell sorting-based quantitation of T cell
receptor restimulation-induced cell death in activated, primary human T cells. *Methods*
Mol Biol **979**, 15-23, doi:10.1007/978-1-62703-290-2_2 (2013).
814