Emergence of a senescent and inflammatory pulmonary CD4+ T cell population prior to lung allograft failure

Sajad Moshkelgosha[1], Liran Levy[1,2], Shahideh Safavi[1,3], Sumiha Karunagaran[1], Gavin Wilson[1], Benjamin Renaud-Picard[1,4], Goodness Madu[1], Rashi Ramchandani[1], Jillian Oliver[1], Tatsuaki Watanabe[1,5], Ke Fan Bei[1], Betty Joe[1], Qixuan Li[6], Ella Huszti[6], May Cheung[1], David Hedley[7], Jonathan Yeung[1,8,9], Shaf Keshavjee[1,8,9], Tereza Martinu[1,8,10], Stephen Juvet[1,8,10]

1. Toronto General Hospital Research Institute, University Health Network, Toronto, ON
2. Sheba Medical Center, Sackler School of Medicine, Tel-Aviv University, Tel-Aviv, Israel
3. Barts Health NHS Trust, London, UK
4. Nouvel Hôpital Civil, Strasbourg, France
5. Department of Thoracic Surggery, Institute of Development, Aging and Cancer, Tohoku University, Sendai, Japan
6. Biostatistics Department, University Health Network, Toronto, ON
7. Division of Medical Oncology, Department of Medicine, University of Toronto
8. Toronto Lung Transplant Program, Ajmera Transplant Centre, University Health Network, Toronto, ON
9. Division of Thoracic Surgery, Department of Surgery, University of Toronto, Toronto, ON
10. Division of Respirology, Department of Medicine, University of Toronto, Toronto, ON

Address of Corresponding Author:
Stephen Juvet, MD, PhD
Toronto General Hospital: University Health Network (UHN)
200 Elizabeth St., Rm 11 PMB 126
Toronto, Ontario, M5G 2C4
EMAIL: stephen.juvet@uhn.ca
TEL: (416) 340-4800 - ext: 8178

Acknowledgments

The authors wish to thank Martin Oberle, Marcelo Cuesta, Iva Avramov and other members of the Toronto Lung Transplant Program Biobank for their assistance with sample procurement and storage. We also thank Dr. Cynthia Guidos and Dr. Dariush Davani for their critical input on the design of mass cytometry experiments. We also wish to thank Dr. Sarah Crome and Sarah Colpitts for their assistance and advice in planning CITE-Seq experiments.

This work was funded by Cystic Fibrosis Foundation grants JUVET18AB0 and JUVET23G0 (to S.J.), and MARTIN18I0 (to T.M. and S.J.) by an International Society for Heart and Lung Transplantation Norman Shumway Career Development Award (to S.J.), by a Canadian Society of Transplantation Research Training Award (to S.M.) and by the University Health Network Foundation.

Abbreviations:

ALAD: Acute lung allograft dysfunction

CLAD: Chronic lung allograft dysfunction

LT: Lung transplantation

BAL: Bronchoalveolar lavage

IMC: Imaging mass cytometry

tSNE: T-distributed stochastic neighbor embedding

CyTOF: Cytometry by Time of Flight

CITE-seq: Cellular indexing of transcriptomes and epitopes

Tph: peripheral helper T

FEV1: Forced expiratory volume in one second

TLTP: Toronto lung transplant program

GLMM: Generalized linear mixed model

ROC: Receiver-operating characteristic

DSA: Donor specific antibodies

HR: Hazard radio

CMV: Cytomegalovirus

ADT: Antibody derived tags

RICD: Restimulation induced cell death

Abstract

Lung transplantation is a life-saving therapy for end-stage pulmonary disease, but its long-term outlook is poor due to a high incidence of chronic lung allograft dysfunction (CLAD). CLAD results from alloimmune-mediated progressive fibrosis and culminates in death or the need for re-transplantation after a median of 6 years. Existing immunosuppression fails to prevent CLAD, suggesting the existence of alloimmune pathways resistant to these drugs. Here, we used mass cytometry to identify cell populations enriched in the bronchoalveolar lavage (BAL) of patients with subsequent allograft dysfunction. We show that $CD4^+CD57^+PD1^+$ T cells emerge in stable lung transplant recipients in the first year post-transplant, conferring heightened risks for CLAD and death or re-transplantation. $CD4^+CD57^+PD1^+$ T cells display features of senescence and secrete inflammatory cytokines. Cellular indexing by transcriptomes and epitopes (CITE-Seq) on BAL $CD4^+$ T cells revealed the existence of 2 oligoclonal $CD57^+$ subsets with putative cytotoxic and follicular helper functions. Finally, we observed that $CD4^+CD57^+PD1^+$ T cells are associated with lung allograft fibrosis in a mouse model and in human explanted CLAD lungs, where they localize near airway epithelium and B cells. Together, our findings reveal the existence of an inflammatory T cell population that predicts future lung allograft dysfunction and may represent a rational therapeutic target in lung transplant recipients.

Introduction

The study of allograft rejection has led to important advances in our understanding of
cellular immunity[1]. Unfortunately, solid organ transplantation continues to be hampered
by high rates of chronic rejection and graft failure despite the availability of potent
immunosuppressive drugs. In lung transplantation (LT), which remains the only life-
prolonging therapy for many cases of end-stage lung disease, median graft survival is
only 6 years[2] due to chronic lung allograft dysfunction (CLAD). CLAD is the result of a
progressive scarring process due to accumulated injuries resulting from alloreactivity
and innate immune activation caused by donor brain death, ischemia-reperfusion injury,
post-transplant infections, gastroesophageal reflux, and air pollution[3]. Both acute T cell-
mediated rejection and antibody-mediated rejection are well-recognized CLAD risk
factors[4,5].

LT recipients often grapple with episodes of acute lung allograft dysfunction (ALAD)[6]. We
characterized ALAD as a marked decrement in lung function – quantified by a decline in
the forced expiratory volume in one second (FEV_1) by 10% or more from its prior value.
Despite exhaustive history, physical examination, and diagnostic testing including
thoracic imaging and bronchoscopy, the cause of ALAD often is not definitively identified.
Recurring ALAD events may be the result of rejection or uncontrolled alloimmunity that
cannot be identified due to the poor diagnostic accuracy of transbronchial biopsies [7,8].

Bronchoalveolar lavage (BAL) is a technique in which saline instilled via a bronchoscope
wedged in a pulmonary segmental airway is aspirated into a collection trap. BAL permits
microbiologic and cytologic analysis of the distal lung compartments including the small
airways and alveolar spaces. We have been studying cellular elements of the BAL using
advanced multiparametric analytical approaches for the last several years[6,9-13]. Using
single-cell RNA sequencing, we recently identified specific alveolar macrophage
populations associated with ALAD episodes and CLAD[6], which may represent both a
mechanistic link between inflammatory insults such as infection and rejection, as well as
a potential target for therapeutic intervention.

Prior studies by several groups have examined specific BAL T cell populations in LT
recipients. Early work examined the question of whether changes in the CD4:CD8 ratio

among BAL T cells might differentiate infection from rejection[14]. Later studies examined

whether T cell activation[15], differentiation[16], regulatory T cells[17] and apoptosis[18] could

distinguish CLAD, acute rejection and infection[19]. High variability between LT recipients

meant that such distinctions were difficult[19] and not useful in an individual patient.

Moreover, since acute rejection and infection and CLAD can all coexist, and there are

large overlaps in the distributions of pre-specified BAL immune cell populations between

these clinical entities, these hypothesis-directed analyses have not led to the identification

of useful biomarkers or mechanistic insights.

Here, we used mass cytometry[20] – a high-dimensional technique in which the expression

of up to 40 cell-associated proteins can be analyzed simultaneously, coupled with

unbiased differential discovery approaches – to identify $CD4^+CD57^+PD1^+$ T cells as a

predictor of incident ALAD and CLAD. Apart from their usefulness as a biomarker, our

data implicate $CD4^+CD57^+PD1^+$ T cells in the mechanism of allograft loss. We find the

cells in chronically rejected human and mouse lung allografts, show that they are clonally

expanded, express transcripts associated with cytotoxic and helper functions, and they

secrete inflammatory cytokines. Further, they are senescent, hypoproliferative, and

resistant to cell death *in vitro.* Our observations therefore suggest the existence of a

specific alloimmune pathway that can escape conventional immunosuppression that may

represent a valuable therapeutic target.

Results

49 **Identification of a T cell subset associated with lung allograft dysfunction**

50 Early detection of alloimmune responses in the graft – *before* ALAD occurs – is likely to

51 provide both important prognostic information as well as an opportunity for early

52 intervention. In pursuit of a greater understanding of ALAD mechanisms, we

53 prospectively collected and viably cryopreserved BAL cells from 50 Toronto Lung

54 Transplant Program (TLTP) LT recipients presenting for surveillance bronchoscopy at 3

55 months post-transplant, and obtained two more follow up samples from each recipient

56 to form a longitudinal cohort **(Figure 1A; Supp Fig 1)**. We performed mass cytometry[20]

57 using a panel of 36 heavy metal-conjugated antibodies **(Supp Table 1)**. Conventional

58 flow cytometric gating revealed considerable inter- and intra-patient heterogeneity in

59 BAL cell composition (data not shown). Unsupervised multidimensional clustering using

60 FlowSOM [21] demonstrated similar variation **(Figure 1B, C)**. In order to facilitate

61 identification of cell populations associated with allograft dysfunction, we randomly

62 divided the cohort into discovery and validation subsets (n=25 each; **Supp Table 2**).

63 Longitudinal samples in the discovery group were then used to search for differences in

64 the BAL cellular composition between stable patients and those experiencing ALAD

65 within 30 days of each bronchoscopy. Initially we employed cluster identification,

66 characterization and regression (CITRUS[22]) on all BAL T cells from all the time-points

67 for all patients in the cohort. This analysis revealed that $CD4^+CD57^+PD1^+$ T cells were

68 the top differentially represented cell population between patients with and without

69 ALAD within 30 days of at least one bronchoscopy **(Figure 1D-F)**. CITRUS does not

70 adequately account for the paired design of our cohort (multiple samples per patient),

71 nor does it incorporate time as a factor in the analysis. We therefore also applied a

72 generalized linear mixed model (GLMM) pipeline described for CyTOF analysis[23]. To

73 facilitate this analysis, we used FlowSOM to identify 20 clusters of T cells **(Figure 1G)**.

74 T cell subsets with greater representation in samples associated with ALAD compared

75 to samples associated with stability were identified using this approach **(Figure 1H)**,

76 revealing that the $CD4^+ CD57^+PD1^+$ T cell population – among several others – was

77 significantly differentially represented in ALAD. Collectively, these data suggested that

78 there is an association between $CD4^+CD57^+PD1^+$ T cells and incident ALAD.

CD4$^+$CD57$^+$PD1$^+$ T cells appear in the BAL prior to the onset of allograft dysfunction

Next, we assessed the relationship of BAL CD4$^+$CD57$^+$PD1$^+$ T cells to ALAD onset. We observed that while patients with stable lung allograft function had low frequencies of CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell compartment (**Figure 2A**), these cells typically accumulated in higher frequencies prior to the onset of ALAD **(Figure 2B)**. Receiver-operating characteristic (ROC) curve analysis showed that having $\geq$ 7.8% CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell compartment optimally discriminated concurrent ALAD from stable patients, with an area under the ROC curve of 0.76 **(Figure 2C)**. Using this criterion, 10 of 11 (91%) LT recipients with ALAD within 30 days of any of their bronchoscopies demonstrated $\geq$ 7.8% CD57$^+$PD1$^+$ in the CD4$^+$ T cell compartment of at least one of their BAL samples; in contrast, only 3 of 14 (21%) stable LT recipients had CD57$^+$PD1$^+$ cells exceeding this threshold **(Figure 2D)**. Moreover, having this biomarker conferred a reduced time-to-ALAD in univariable survival analysis **(Figure 2E)**. We then applied the 7.8% threshold to the validation subset (n=25) and observed a similar reduction in time-to-ALAD from first appearance of the cells in LT recipients with at least one BAL sample in which it was exceeded, compared to LT recipients who did not have any BAL samples in which the threshold was exceeded **(Figure 2F)**. Together, these data illustrated that in LT recipients developing subsequent allograft dysfunction, CD57$^+$PD1$^+$ cells frequently exceeded the 7.8% threshold in BAL samples obtained prior to ALAD and CLAD **(Supp Fig 2).**

Anti-human leukocyte antigen donor-specific antibodies (DSA) are an adverse prognostic factor in lung transplantation[24]. Patients with ≥7.8% CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell compartment were more likely to develop de novo DSA than those without **(Figure 2G)**. BAL CD4$^+$CD57$^+$PD1$^+$ T cells were not simply a surrogate for other conventional clinical and bronchoscopic parameters, since clinical symptoms, biopsy-proven vascular or airway rejection, BAL pathogens and antimicrobial treatments did not differ significantly between patients with and without $\geq$ 7.8% CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell compartment **(Supp Table 3)**. Similarly, although the number of events was small, having CD57$^+$PD1$^+$ cells above the 7.8% threshold was not clearly

109 associated with an immediate post-bronchoscopy augmentation of immunosuppression
110 or change in calcineurin inhibitor **(Supp Table 4)**. Taken together, these observations
111 indicate that BAL $CD4^+CD57^+PD1^+$ T cells may be a useful biomarker of incident lung
112 allograft dysfunction that adds significant diagnostic utility to conventional clinical
113 monitoring.

$CD4^+CD57^+PD1^+$ T cells confer an increased risk for CLAD and death and are
present in chronically rejected human lung allografts

To examine whether the presence of $CD4^+CD57^+PD1^+$ T cells in the BAL samples of LT
recipients increase the risk for CLAD, we followed all 50 individuals in our cohort for up
to 2000 days post-LT (median follow-up 1788 days). Using the appearance of ≥7.8%
$CD57^+PD1^+$ cells in the BAL $CD4^+$ T cell compartment or the last analyzed BAL as the
index event, we found that time to CLAD was significantly shorter in those patients whose
BAL samples exceeded the 7.8% threshold (log rank p=0.0071, **Supp Figure 3A**). We
then performed Cox proportional hazards analysis, excluding two patients who developed
CLAD prior to their final BAL samples (**Supp Fig 1**). Treating the appearance of ≥7.8%
$CD57^+PD1^+$ cells in the BAL $CD4^+$ T cell compartment as a time-dependent covariate,
Cox proportional hazards analysis revealed a hazard ratio (HR) of 3.61 (95% CI 1.55-
8.39) for time-to-CLAD. In this and other survival analyses, patients were followed until
31 December 2022 (median follow up time from transplantation: 2245 days). The
increased risk for CLAD associated with $CD4^+CD57^+PD1^+$ cells persisted in bivariable
models adjusted for age at transplant (HR 3.67, 95% CI 1.57-8.55), sex (HR 3.45, 95%
CI 1.47-8.07), and primary cytomegalovirus (CMV) mismatch (donor seropositive,
recipient seronegative; HR 4.02, 95% CI 1.70-9.52).

Similarly, a Cox proportional hazards model revealed that ≥7.8% $CD57^+PD1^+$ cells in the
BAL $CD4^+$ T cell compartment was associated with a HR of 4.82 (95% CI 1.8-13.1) for
time-to-death or retransplantation. The effect persisted in bivariable models adjusting for
age at transplant (HR 4.82, 85% CI 1.8-13.1), sex (HR 4.84, 1.8-13.2) and primary CMV
mismatch (HR 4.92, 95% CI 1.8-13.4). These time-to-CLAD and time-to-death or
retransplantation analyses are presented in **Supp Table 5**. We then examined time-to-
death or retransplantation in a survival analysis landmarked at 12 months post-transplant.

LT recipients with $\geq 7.8\%$ CD57$^+$PD1$^+$ cells in the BAL CD4$^+$ T cell compartment had an
increased risk of death or retransplantation (p=0.00079, **Figure 3A**).

These observations suggested that CD4$^+$CD57$^+$PD1$^+$ T cells may lie on the mechanistic
pathway leading to lung allograft fibrosis. Therefore, we sought to identify them in CLAD
lung tissue obtained at the time of retransplantation[25] using imaging mass cytometry
(IMC[26]). CLAD and normal lung tissue sections were stained with a panel of 35 heavy
metal conjugated antibodies **(Supp Table 6)** and subjected to IMC. Unsupervised
clustering identified CD4$^+$CD57$^+$PD1$^+$ T cells as one of the cell populations present
(cluster 50 in **Figure 3B, Supp Fig 3B**), with more cells in CLAD compared to normal
tissue (**Figure 3C-D**). Image reconstruction and proximity analysis revealed that
CD4$^+$CD57$^+$PD1$^+$ T cells were frequently located adjacent to distinct B cell populations,
including trafficking B cells (CD20+CCR6+, Cluster 8) and plasma cells (IgM+CD138+,
cluster 30) and airways **(Figure 3E-F)**.

CD4$^+$CD57$^+$PD1$^+$ T cells infiltrate mouse lung allografts and associate with the
severity of chronic rejection

We have developed a minor alloantigen-mismatched mouse orthotopic lung transplant
model in which a minimal graft storage period prior to transplantation results in mild or
absent allograft fibrosis at 28 days post-transplant, while prolonged warm and cold
ischemic storage prior to transplantation leads to B cell- and alloimmune-dependent
fibrosis with features of human chronic lung allograft rejection[27] (**Figure 4A-B**). We asked
whether CD4$^+$CD57$^+$PD-1$^+$ T cells might be found within lung allografts in this model, and
whether their abundance correlated with the degree of allograft fibrosis. Allografts were
scored in blinded fashion using our published scoring system[28], which includes
assessment of the proportion of the allograft affected by interstitial inflammation and
fibrosis. As we have observed previously[27,28], pathology was variable, with some animals
exhibiting higher degrees of fibrosis and inflammation (**Fig 4C**). Flow cytometry performed
on single-cell suspensions of digested lung allograft tissue revealed that CD57$^+$PD1$^+$ T
cells were common among lung allograft CD4$^+$ T cells, but rare in normal lung tissue
(**Figure 4D**). It also showed that allografts with higher degrees of pathology – as
determined by the presence of both interstitial inflammation and parenchymal fibrosis –

had higher frequencies of CD57⁺PD1⁺ T cells compared to those with a low degree of
pathology (**Figure 4E-F**). Further, CD4⁺CD57⁺PD1⁺ T cells were readily detectable in
allograft sections by immunofluorescence (**Figure 4G**). Together, these data support the
concept that CD4⁺CD57⁺PD1⁺ T cells accumulate in rejecting lung allografts and correlate
with the severity of inflammation and fibrosis in mice.

Identification of two transcriptionally distinct CD4⁺CD57⁺PD1⁺ cell subsets using
CITE-Seq

To characterize gene expression in CD4⁺CD57⁺PD1⁺ T cells, we enriched CD3⁺ T cells
from four ALAD BAL samples and performed cellular indexing by transcriptomes and
epitopes (CITE-Seq[29]) with 5' end sequencing. We applied a panel of oligonucleotide-
conjugated antibodies, including CD4, CD57 and PD1, to ALAD BAL samples containing
at least 7.8% CD57⁺PD1⁺ cells in the CD4⁺ T cell compartment as determined by flow
cytometry. The conjugated oligonucleotides or antibody-derived tags (ADTs) identified in
CITE-Seq allowed confident linkage of transcriptomes with protein expression.

BAL CD4⁺ T cells from patients with ALAD fell into 9 subclusters in multidimensional
space (**Figure 5A**). Based on CD57 and PD1 protein expression, CD4⁺CD57⁺PD1⁺ T
cells assembled into two main areas, cluster 4 and the closely adjacent clusters 1 and 2
(Figure 5A). We noted that CD27 gene and protein expression was enriched in cluster
4 compared to clusters 1 and 2 **(Figure 5B).** The CD27⁻ population had a gene
expression profile similar to peripheral helper T (Tph) cells described in rheumatoid
arthritis[30]; differentially expressed genes in this population included CXCL13, TOX, IL-
21, CD40L, TIGIT, ICOS and TNFSRF18 **(Figure 5C)**. These cells are characterized by
the absence of the conventional follicular helper T cell markers CXCR5 and Bcl6, and
by expression of the chemokine CXCL13 [30,31]. Accumulating studies have shown that
Tph cells are involved in various acute and chronic inflammatory diseases [30,32]. Tph
cells can promote B cell differentiation into plasmablasts through interactions involving
IL-21 and SLAMF5 [30]. Therefore, the Tph-like features of CD27⁻ CD4⁺CD57⁺PD1⁺ T
cells, along with their proximity to B cells **(Figure 3E-F)**, their association with de novo
DSA **(Figure 2G)** suggest that they may provide B cell help.

The CD27[+] population expresses several cytotoxicity-related transcripts including
granzymes and NKG7[33] **(Figure 5C)**. They also express CXCR6, which has been
associated with tissue resident memory CD8[+] T cell localization to airways[34] and is in
keeping with our observation of CD4[+]CD57[+]PD1[+] T cells in proximity to airways **(Figure
3F)**.

We next examined TCR CDR3 sequences in the CITE-Seq data, which revealed that
CD57[+]PD1[+]CD27[-] cells are clonally expanded to a greater extent than CD57[+]PD1[+]CD27[+]
and other BAL CD4[+] T cells **(Figure 5D)**. This suggested that they had recently
undergone antigen-driven expansion in the lung. Analysis of the top 20 expanded TCR
clonotypes did not infer any known antigen. However, the high similarity among the top
20 expanded TCRs may suggest that they are alloreactive **(Supp Figure 5).** Pseudotime
trajectory analysis revealed that CD57[+]PD1[+]CD27[-] cells and their CD27[+] counterparts
may share a common origin **(Figure 5E)**. Together, the TCR sequence data and
pseudotime analysis suggested that BAL CD4[+]CD57[+]PD1[+] T cells either differentiate into
Tph or cytotoxic effector cells. The former have undergone significant clonal expansion,
whereas the latter have a distinct TCR repertoire and less clonal expansion, suggesting
that they may be responding to different antigens in the lung microenvironment.

Finally, both CD27[-] and CD27[+] subsets demonstrated expression of previously defined
senescence and exhaustion transcripts including TIGIT, KLRG1, LAG3, HAVCR2/TIM-3
and EOMES which have been described in different acute and chronic inflammatory
conditions[35-37] . This observation is in keeping with reports that have described different
senescent CD57[+] and/or PD1[+] T cell populations in other disease states [38,39].

Functional analysis of CD4[+]CD57[+]PD1[+] T cells reveals features of inflammation
and senescence

Considering the potential distinct functional properties – Tph, cytotoxic, and senescent –
of CD4[+]CD57[+]PD1[+] T cells, we set out to characterize their functions *in vitro*. For this
purpose, CD4[+] T cells from BAL samples of 15 lung transplant recipients experiencing
ALAD were sorted to obtain CD4[+]CD57[+]PD1[+] T cells and their CD4[+]CD57[-]PD1[-]
counterparts **(Figure 6A)**. Fluorescence-activated cell sorting data demonstrated that in
all cases, CD57[+]PD1[+] cell frequency was ≥7.8% of BAL CD4[+] T cells **(Figure 6B)**. Based

on the hypothesis that CD4$^+$CD57$^+$PD-1$^+$ T cells may be senescent, we predicted that

sorted cells would exhibit decreased proliferation, enhanced cytokine secretion, and

resistance to TCR restimulation-induced cell death (RICD) when compared to CD57$^-$PD1$^-$

cells.

Supernatants from cells stimulated overnight were collected and subjected to a

multiplexed cytokine assay. This revealed that CD4$^+$CD57$^+$PD1$^+$ cells – compared to

CD4$^+$CD57$^-$PD1$^-$ cells sorted from the same BAL samples – expressed more IL-21, IL-4,

CXCL8 and granzyme A **(Figure 6C)**. In keeping with our CITE-Seq data, these findings

suggested that CD4$^+$CD57$^+$PD1$^+$ T cells may have Tph or cytotoxic properties – or both.

Stimulated CD4$^+$CD57$^+$PD1$^+$ T cells and their CD57$^-$PD1$^-$ counterparts were then tested

for proliferation, based on dilution of CFSE **(Figure 6D).** Proliferation data revealed that

CD4$^+$CD57$^+$PD1$^+$ T cells exhibited significantly less proliferation compared to CD4$^+$CD57$^-$

PD1$^-$ cells **(Figure 6E)**, in keeping with T cell senescence [38,39]. Finally, CD4$^+$CD57$^+$PD1$^+$

T cells were less susceptible to RICD compared to CD4$^+$CD57$^-$PD1$^-$ cells **(Figure 6F, G)**.

Together these data imply that CD4$^+$CD57$^+$PD1$^+$ T cells are senescent but pro-

inflammatory T cells that are resistant to antigen-induced death. These observations may

help to explain the emergence of ALAD and CLAD in lung transplant recipients despite

potent triple-drug immunosuppression, which would be predicted to be less effective

against these cells.

Discussion

In this study, we used a high dimensional mass cytometric discovery approach to
identify a T cell population in lung transplant recipient BAL that is associated with a
heightened risk for future allograft dysfunction, failure, and death. The risk conferred by
the appearance of CD4$^+$CD57$^+$PD1$^+$ T cells in the BAL remained elevated after bivariate
adjustments for age, sex, and donor and recipient CMV serostatus mismatch.

In keeping with previous work examining lung transplant recipient BAL, our samples
contained a diversity of leukocyte populations. The prior studies demonstrated
statistically significant relationships between BAL CD8$^+$ T cells, NK cells, eosinophils,
Tregs and outcomes in lung transplantation, but this has not led to meaningful
improvements in clinical care because wide overlaps in the frequency distributions of
these populations have meant that they lack discriminatory power in individual
patients[14,15,19,40-43]. In contrast, while CD4$^+$CD57$^+$PD1$^+$ T cells will require further
validation in larger studies, their prevalence among CD4$^+$ T cells discriminated patients
who remained stable from those about to undergo ALAD with high precision, in a
discovery cohort and confirmed in an independent validation cohort. Our observation
therefore represents a promising biomarker that may, in the future, permit the selection
of lung transplant recipients for targeted preventative interventions.

A review of clinical and laboratory variables present at the time of each bronchoscopy
did not reveal any clear association between the appearance of ≥ 7.8% CD57$^+$PD1$^+$ T
cells in the BAL CD4$^+$ T cell compartment and other indicators of lung allograft
inflammation or dysfunction. This observation likely reflects the relatively poor
performance of histopathologic and cytologic findings in the diagnosis of lung allograft
rejection. Indeed, interpretation of transbronchial biopsy histology is fraught, with high
rates of ungradable biopsies and substantial inter-observer disagreement among
pathologists[7,8]. It is therefore perhaps not surprising that bronchoscopies with high
frequencies of CD4$^+$CD57$^+$PD1$^+$ T cells showed no particular association with histologic
rejection. Likewise, we found that this T cell population is not a marker of lung allograft
infection, since there were no differences in pathogen isolation from the same BAL
samples. Our interpretation of these findings is that analysis of BAL CD4$^+$CD57$^+$PD1$^+$ T

cells may dramatically improve the utility of surveillance bronchoscopy after LT, which

will require further evaluation in larger prospective studies.

Beyond their value as a biomarker of future allograft dysfunction, several lines of

evidence support the notion that CD4$^+$CD57$^+$PD1$^+$ T cells are mechanistically relevant

to the pathogenesis of chronic lung allograft rejection and fibrosis. First, our CITE-Seq

data indicate that the majority of clonally expanded CD4$^+$ T cell populations in the BAL

of patients with ALAD were CD57$^+$PD1$^+$, which indicates that these cells have

undergone antigen-driven expansion in the lung. Further, in addition to their appearance

in BAL, we detected CD4$^+$CD57$^+$PD1$^+$ T cells in human explanted CLAD lung tissue

obtained at the time of re-transplantation. The cells were observed in proximity to B

cells and small airways; coupled with the finding that the transcriptome and secretome

of BAL CD4$^+$CD57$^+$PD1$^+$ T cells includes IL-21 and granzymes, this observation

suggests that they are likely participating in allograft injury and may subserve Tph

functions. In support of this assertion, in our ischemia-reperfusion injury-augmented

mouse model of human CLAD, we observed CD4$^+$CD57$^+$PD1$^+$ T cells in the lung

allografts of mice with marked fibrotic pathology. In contrast, allografts with milder

fibrosis did not contain many CD4$^+$CD57$^+$PD1$^+$ T cells. Interestingly, we have previously

shown that fibrosis in this model is B cell dependent[27], and so in future work it will be

important to consider the possible existence of a CD4$^+$CD57$^+$PD1$^+$ T cell-B cell axis that

drives chronic lung allograft rejection in mice and humans.

PD-1 expression by CD8$^+$ T cells in tumor microenvironments is associated with cancer

progression, and generally reflects a state of exhaustion in which effector functions are

deactivated. BAL CD4$^+$CD57$^+$PD1$^+$ T cells also expressed several additional molecules

that have been associated with exhaustion, including TIM-3, LAG3 and CTLA-4. While

analysis of the cells *ex vivo* revealed that they were both hypoproliferative and resistant

to T cell RICD, their production of inflammatory cytokines – which is not a feature of

exhaustion – suggests that they are senescent rather than exhausted. Of note,

senescent CD4$^+$ and CD8$^+$ T cells in humans were described as having the phenotype

CD45RA$^+$KLRG1$^+$CD57$^+$CD28$^-$CD27$^-$ [44,45] Animal and human data indicate that

senescent T cells participate in other chronic inflammatory, degenerative and metabolic

disorders[39].

Lung transplant recipients typically receive immunosuppressive drugs – calcineurin
inhibitors, corticosteroids, and antimetabolites – in much higher doses than other solid
organ transplant recipients, and yet also have the highest rates of acute and chronic
allograft rejection. With respect to T cells, these drugs act by inhibiting cell proliferation.
In this context, we speculate that $CD4^+CD57^+PD1^+$ T cells, which exhibit reduced
proliferation, escape this inhibition and can continue to produce cytokines. Further work
will be required to understand how these cells arise within the lung allograft, and
whether specific targeting of these cells may have a therapeutic benefit.

Methods

Study design and BAL collection

The samples analyzed in this study were collected with informed consent for research

use and were approved by University Health Network Research Ethics Board (15-9531).

BAL cells from 50 TLTP lung transplant recipients presenting for surveillance

bronchoscopy at 3 months post-transplant were obtained according to a standard

protocol [46]. In addition, for each LT recipient, two further samples were obtained from

each recipient to form a longitudinal cohort **(Supp Fig 1)**. Cells were centrifuged at 400g

for 5 min and resuspended in 1 ml of 10% dimethyl sulfoxide/90% FBS for viably

cryopreservation. Cryovials were placed in Mr. Frosty containers at -80°C for 24 h, then

stored in liquid nitrogen until thawing for processing and mass cytometry. BAL cells from

a further 14 LT recipients with acute lung allograft dysfunction (ALAD, defined as a

decrease in the forced expiratory volume in one second [FEV1] by 10% or more from

the baseline FEV1 measurements) were used for sorting fresh $CD4^+CD57^+PD1^+$ T cells

and their $CD57^-PD1^-$ counterparts for functional analyses.

Mass cytometry

Metal-conjugated antibodies were either purchased preconjugated from Fluidigm or

purchased purified from BioLegend, Thermo Fisher Scientific or BD Biosciences and

subsequently conjugated to metals using Maxpar Antibody Labeling Kits (Fluidigm)

according to the manufacturer's instructions **(Supp Table 1)**. Cryopreserved samples

were recovered and divided into stimulated (with PMA 50ng/mL and ionomycin

500ng/mL) and unstimulated conditions to allow analysis of cytokine secretion

expression and surface markers. Cells were incubated with a cocktail of antibodies

against cell surface molecules for 30 min at room temperature. After treatment with

intracellular fixation and permeabilization buffers (BD Biosciences), cells were incubated

with metal-conjugated antibodies against intracellular proteins. Cells were then washed

and stained with Intercalator-Ir (Fluidigm) diluted in PBS containing 1.6%

paraformaldehyde and stored at 4 °C until next-day acquisition. To address technical

variations between mass cytometry runs, a control healthy volunteer PBMC sample was

aliquoted, recovered and stained and analyzed alongside the BAL samples.

Mass cytometry data analysis

Mass cytometry data were initially analyzed with manual gating to exclude debris, dead cells and doublets and to select CD45+ cells using Cytobank software (Beckman Coulter). We then used the FlowSOM and viSNE algorithms (5,000 iterations, perplexity = 100) for visualization of high-dimensional data [21,47]. Subsequently, we applied cluster identification, characterization, and regression (CITRUS) [22] on the CD3+ population from the discovery cohort (n=25, **Supp Table 2**) to identify cell subsets associated with ALAD within 30 days of the sample. CITRUS was run at least 5 times with "equal number" of cells per sample, looking for "abundance" as cluster characterization, with a 1% false discovery rate. To incorporate the time-course design of our study, we implemented a generalized linear mixed model analysis-adapted R pipeline as described by Nowicka and colleagues [23]. In this approach, different timepoints for each LT recipient were defined as a variable to allow intra-subject pairing. Finally, we used manual gating to confirm the results of CITRUS and generalized linear mixed model analysis.

Imaging Mass Cytometry

Lung tissue samples were obtained during retransplantation procedures from 8 patients who were undergoing a second Lung Transplantation due to end-stage CLAD. A single biopsy was obtained from the upper lobe of each patient presenting with CLAD. The harvested lung tissue was initially inflated and immersed in a 10% formalin solution for 24 hours. Subsequently, it underwent immersion in a 70% ethanol solution for an additional 24-hour period before being embedded in paraffin. Sections of the paraffin-embedded lung tissue, with a thickness of 4µm, were prepared and affixed to slides for further analysis. For every specimen, we meticulously identified three distinct 1x1mm Regions of Interest (ROIs) situated around individual airways. In addition to the CLAD patient samples, we included control samples in our study. Four lung biopsies were procured: two from patients undergoing lobectomy procedures for suspected cancer and two from surplus donor lung tissue collected at the conclusion of the cold ischemic phase during lung transplantation. To facilitate our analysis, sections of the lung tissue were subjected to staining using a panel of 35 heavy-metal-tagged antibodies

(Supplementary Table 6). Subsequently, the three ROIs chosen for each specimen

were subjected to ablation using the Hyperion Imaging System developed by Fluidigm.

The ablation process employed a laser frequency of 200Hz, with laser power settings

ranging from 3.5 to 4.5. The ablation procedure generated files in both .txt and .mcd

formats, which were subsequently subjected to thorough analysis using the HistoCAT

software platform[48].

Animal model

Orthotopic lung transplantation was performed using C57BL/10 donors and C57BL/6

recipients under different graft preservation conditions as described previously [27].

Briefly, harvested lungs were stored either under minimal (60 minutes at 4ºC followed

by 15 min of warm anastomotic time) or prolonged (6h at 4ºC, 45 min at 37ºC, followed

by 15 min of warm anastomotic time) storage prior to reperfusion. Orthotopic lung

transplant recipients were sacrificed 28 days after LT for histological analysis and

cellular phenotyping. Grafts were flushed with saline and digested with collagenase A

and DNase I in a gentleMACS tissue dissociator (Miltenyi Biotec). Cells were stained

with cocktails of fluorochrome-conjugated monoclonal antibodies including AF647-

conjugated anti-mouse CD57 (polyclonal; Bioss) and AF488-conjugated anti-mouse

PD1 (polyclonal; Bioss). Flow cytometry data were analyzed using FlowJo (v10, FlowJo

LLC) and correlated with histological grading. We graded peribronchial fibrosis and

parenchymal fibrosis in a blinded fashion according to a published scoring system[28].

Immunofluorescent staining of sections of formalin-fixed and paraffin-embedded lung

tissue was conducted by antigen retrieval using 1x pH 9 Tris/EDTA buffer before

staining with unconjugated anti-mouse CD4. This was followed by a secondary antibody

cocktail including Alexa Fluor 555 anti-rabbit IgG (A-31572, Invitrogen) as well as

AF647-conjugated anti-mouse CD57 (polyclonal; Bioss) and AF488-conjugated anti-

mouse PD1 (polyclonal; Bioss). Nuclei were stained with DAPI and

immunofluorescence images were acquired using a Zeiss AxioObserver.

CITE-seq sample processing and acquisition

We identified four LT recipients scheduled for bronchoscopy with ALAD (FEV_1 decrease
by ≥10% from the prior value). Fresh BAL cells were collected by centrifugation at 400g
for 5 minutes. T cells were obtained using a $CD3^+$ T cell enrichment kit (EasySep™
Human T Cell enrichment, Stemcell Technologies) according to the manufacturer's
instructions. Enriched T cells were resuspended at $1x10^6$ cells/mL in PBS + 0.05% BSA.
50 μL of the cell suspension were stained with BV711 conjugated anti-human CD4,
APC conjugated anti-human PD1 and PE-Cy7 anti-human CD57 and analyzed by flow
cytometry to confirm ≥ 7.8% $CD57^+PD1^+$ cells in the BAL $CD4^+$ T cell compartment. If
confirmed, the remaining cells were used for CITE-seq. Briefly, $1x10^6$ cells were
resuspended in 45 μL PBS + 0.05% BSA before adding 5 μL of Fc blocking reagent and
incubated for 10 minutes at 4°C. TotalSeq™-C antibody cocktails were prepared during
the incubation by adding pre-determined concentrations of each antibody **(Supp Table
6)**. The resulting cocktail was then added to cells and incubated for 30 minutes at 4°C.
After incubation, the cell pellets were resuspended in 1mL PBS + 0.05% BSA and
centrifuged for 5 minutes at 400 x g. After a total of 3 washes, the final cell density was
adjusted to 1000-1200 cells/μL and subjected to the 10X Genomics Chromium 5'
Immune Profiling v2.0 protocol for library preparation. A total of 2×10^4 cells from each
sample were loaded onto a 10X single cell G chip. After droplet generation, samples
were transferred to a pre-chilled 96 well plate (Eppendorf), heat sealed and incubated
overnight in a Veriti 96-well thermocycler (Thermo Fisher). The next day, sample cDNA
was recovered using Recovery Agent (10X Genomics) and subsequently cleaned up
using Silane DynaBeads (Thermo Fisher) mix as outlined by the user guide 5' Immune
Profiling v2.0. Purified cDNA was amplified for 11 cycles before being cleaned up using
SPRIselect beads (Beckman Coulter). Samples were diluted 9:1 (elution buffer
(Qiagen):cDNA) and run on a Bioanalyzer (Agilent Technologies) to determine cDNA
concentration. The library was then sequenced using an Illumina Novaseq 6000
sequencer using the following parameters Read 1: 28b, index 1: 10bp, Index 2: 10bp,
Read 2: 90bp.
434
435
CITE-seq analysis

The sequencing output was first demultiplexed and converted into FASTQ files using

Cell Ranger [49]. Reads were aligned to the human reference genome (grch38), and

count UMIs in the mRNA libraries, and CITE-seq-Count to count UMIs in the antibody-

derived tag (ADT) libraries. The matrix, barcodes, and gene information from Cell

Ranger were then used to create gene-cell matrices and a Seurat object [50]. For each

Seurat object, cells with poor quality were excluded based on the following quality

control metrics: unique gene counts over 5000 or below 200, UMI counts over 30,000,

and mitochondrial gene expression higher than 10%. Each sample was scaled and

normalized using Seurat's "SCTransform" function followed by adding the protein

expression levels to the Seurat object and normalization and scaling for ADT assay.

Next, CD4$^+$ T cells were subsetted based on high CD4 and low CD8 protein expression

(ADT-CD8 < 12 & ADT-CD4 > 12). Seurat objects of CD4+ cells from all four BAL

samples were merged into a single Seurat object. The SCTransform method was used

for normalization, scaling, and integration which corrects for unwanted sources of

variation and enhances the signal of genuine biological differences between cells[51].

Then, dimensionality reduction was applied to the integrated Seurat object, allowing

visualization and interpretation of the high-dimensional scRNAseq data. This was

followed by unsupervised clustering (resolution = 0.5) which led to the identification of

10 clusters. Then cells with higher expression of protein level of CD57 and PD1 were

highlighted on tSNE plot.

RNA velocity estimates the future state of individual cells by calculating the ratio of

unspliced to spliced mRNA for each gene. The spliced and unspliced counts from the

scRNAseq data were utilized to calculate RNA velocity using the velocyto package in

Python[52]. Splicing information extracted from velocyto was then added to all the

selected cells in the CD4$^+$ Seurat object. Subsequently, CD4$^+$ T cells underwent

pseudotime trajectory analysis using the scVelo pipeline[53]. This analysis inferred the

trajectory and the chronological sequence of cell states, highlighting the differentiation

paths taken by cells. The pseudotime values were visualized on the tSNE plots,

providing insights into the dynamic processes and cellular transitions at play.

Fluorescence-activated cell sorting

BAL cells collected from 15 ALAD samples and stained with fluorophore-tagged surface

antibodies for 30 min at room temperature. The following antibodies were used to stain

the cells: BV711-conjugated anti-human CD4 (BioLegend); APC-conjugated anti-human

PD1 (BioLegend); PE/Cy7-conjugated anti-human CD57, in addition to biotinylated

CD163, CD14, CD8, and CD56 (all from BioLegend). Cells washed with 500 µL of 1%

FBS in PBS and incubated with streptavidin-AF488 (1:200, BioLegend) for 20 minutes,

before washing and staining with propidium iodide to identify dead cells. Stained cells

were subjected to FACS sorting using a Sony SH800 instrument. After exclusion of PI-

positive cells and putative doublets based on forward and side scatter analysis, cells in

the dump channel (AF488) were eliminated. The remaining cells were gated to select

CD4$^+$ T cells and sorted for CD57$^+$PD1$^+$ and CD57$^-$PD1$^-$ cells.

Cell culture, multiplex protein analysis, and *in vitro* assays

Sorted cells were labeled with 1µM CFSE in serum free PBS prior to wash and culture

with plate bound anti-CD3 (5µM) and soluble anti-CD28 (1µM) antibodies in DMEM

medium containing 5.55 mM D-glucose, 4 mM L-glutamine, and 1 mM sodium pyruvate,

supplemented with 10% FBS, 50 U/ml penicillin, and 50 mg/ml streptomycin and 500

U/ml human recombinant IL-2 (Proleukin, Chiron). After 18 hours, 100µL supernatant

was replaced with fresh media. The collected supernatant was stored at -80°C for later

multiplex analysis. Supernatants were thawed and subjected to a custom multiplex bead

kit (R&D Systems) to assess the levels of released analytes including IL-21, IL-4,

CXCL8, and granzyme A. Samples were prepared in a 1:1 dilution with assay buffer, as

suggested by the manufacturer. Diluted samples and standards were run in duplicate.

Biomarker concentrations were obtained using a Bio-Plex MAGPIX Multiplex Reader

(Bio-Rad). All samples and standards were run in duplicate according to manufacturers'

protocols.

After 5 days, CFSE dilution in the sorted cells – a measure of T cell proliferation – was

determined by flow cytometry. Finally, a restimulation-induced cell death assay[54] was

used to assess whether CD57$^+$PD1$^+$ cells are less susceptible to this mode of cell death

than their CD57$^-$PD1$^-$ counterparts. CD57$^+$PD1$^+$ and CD57$^-$PD1$^-$ cells were stimulated in

culture for 7 days before washing, and were then restimulated with anti-CD3 mAb

(OKT3, 100 ng/ml final) for 24 hours. Cells were then harvested and stained with

Annexin V and propidium iodide for evaluation of apoptotic cell death.

Statistics

All statistical tests were two-sided unless stated otherwise. The Wilcoxon rank-sum test was used to assess statistical differences between two groups. When assessing >2 groups simultaneously, the nonparametric Kruskal–Wallis test was used. A Fisher's exact test was applied to assess statistical differences between two categorical variables. Receiver Operating Characteristic (ROC) curves were used to assess classification accuracy, which was quantified by Area Under the Curve (AUC). Kaplan–Meier plots and two-sided log-rank tests were used to compare survival probabilities between groups. Cox Proportional Hazards (PH) regression models were used to adjust for potential confounders for time-to-CLAD or time-to-death/retransplantation analysis. and Cox regression analyses were used to assess covariates with respect to time-to-allograft dysfunction. All statistical analyses were performed using R v.3.5.1+ or Prism 8+ (GraphPad Software).

Figure legends

Figure 1. Identification of a bronchoalveolar lavage (BAL) CD4$^+$ T cell population

associated with lung allograft dysfunction. A. Study schematic. Fifty consecutive

patients presenting for bronchoscopy at 3 months post-transplant were enrolled. For

each enrolled patient, BAL cells at three time points (3mo, 6mo, 9mo or next available)

were collected. BAL cells were cryopreserved and subjected to mass cytometry (see

supplementary table 1); all samples from each patient were analyzed in the same mass

cytometry experiment. **B.** t-distributed stochastic neighbour embedding (tSNE) plots of

BAL CD45$^+$ cells falling into 20 clusters (highlighted in different colours) identified by

FlowSOM. **C.** tSNE plots highlighted by expression of 4 different lineage markers; CD3

(top left), CD11c (top right), CD56 (bottom left), CD19 (bottom right). **D.** CITRUS

(cluster identification, characterization, and regression) analysis of all BALs from n=25

discovery group patients showing cell populations differentiating patients developing

allograft dysfunction within 30 days from those whose allograft function remained stable.

Red circle indicates the statistically significantly differentially represented population. **E.**

Mass cytometry histograms of the top differentially represented population, which

expresses CD4, CD57, CD3 and PD1. **F.** Box-and-whisker plot showing the abundance

of CITRUS cluster 1076 – representing CD4$^+$CD57$^+$PD-1$^+$ T cells – which exhibited a

significant difference between ALAD and stable samples. **G.** t-SNE plot based on

arcsinh-transformed marker expression in 500 randomly selected cells per sample from

the discovery set (n=25, all 3 time points shown). Cells are coloured according to the 20

cell populations obtained with FlowSOM after the metaclustering step with

ConsensusClusterPlus. **H.** Differential analysis heatmap of arcsine-square-root

transformed cell frequencies that were subsequently normalized per cluster (rows, 20

clusters identified in the panel E). Statistically significant clusters are highlighted with

green and showing their p-values. **I.** Marker expression of clusters revealed that cluster

14 is significantly differentially represented between ALAD and stable patients

(highlighted with red rectangle), with a similar protein expression signature

(CD4$^+$CD57$^+$PD1$^+$) as identified by CITRUS.

Figure 2. Relationship of BAL CD4$^+$CD57$^+$PD1$^+$ T cells with incident allograft

dysfunction. Representative examples of forced expiratory volume in one second

(FEV1) over time in weeks post-transplant for stable (**A**) and ALAD (**B**) lung transplant recipients in the discovery cohort. Weeks during which bronchoscopy was performed are highlighted in yellow; scatterplots showing expression of CD57 and PD1 in the CD4$^+$ T cell compartment of the BAL are shown for each time point. Green line indicates 80% of best post-transplant baseline, below which CLAD may be diagnosed. **C.** Receiver-operator characteristic (ROC) curve analysis of the performance characteristics of CD57$^+$PD1$^+$ cells as a proportion of the BAL CD4$^+$ compartment for diagnoses of subsequent ALAD. The area under the curve (AUC) = 0.76. The percentage of CD57$^+$PD1$^+$ T cells providing optimal discrimination of ALAD and stable patients (7.8%) is circled in red. **D.** The highest proportion of CD57$^+$PD1$^+$ T cells observed in all BAL samples from each patient (discovery cohort, n=25) is shown, divided by the presence or absence of ALAD within 30 days. Mann-Whitney test p=0.0018. Dotted line shows the 7.8% threshold identified in ROC curve in C. **E.** Time to lung allograft dysfunction from the emergence of CD57$^+$PD1$^+$ cells $\geq$ 7.8% of CD4$^+$ T cells in the BAL (red line), or from the final BAL analyzed (blue line) in the discovery cohort (n=25). Log rank test p < 0.0001. **F.** Time to lung allograft dysfunction from the emergence of CD57$^+$PD1$^+$ cells $\geq$ 7.8% of CD4$^+$ T cells in the BAL (red line), or from the final BAL analyzed (blue line) in the validation cohort (n=25). Log rank test p < 0.02. G. Of 28 patients who developed de novo DSA, 19 (67.9%) had CD57$^+$PD1$^+$ cells $\geq$ 7.8% of CD4$^+$ T cells in at least one BAL sample; only 9 of 23 (39.1%) patients who did not develop de novo DSA had CD57$^+$PD1$^+$ cells $\geq$ 7.8% of CD4$^+$ T cells in at least on BAL sample. Chi square test p = 0.0402.

Figure 3. CD4$^+$CD57$^+$PD1$^+$ T cells are associated with an increased risk of death or re-transplantation and are found in CLAD lung tissue. A. Kaplan-Meier curves showing time to death or re-transplantation from 12 months post-transplant. Log-rank test p = 0.00079. **B.** Selected heatmap of protein expression for selected clusters and markers to highlight protein profile of cluster 50. **C.** Imaging mass cytometry of normal donor lung tissue (left) and CLAD lung tissue obtained at the time of retransplantation (right). E-cadherin$^+$ epithelial cells (green pseudocolour), red CD3$^+$ T cells (red pseudocolour), CD4$^+$ cells (blue pseudocolour), and CD4$^+$CD57$^+$PD1$^+$ T cells (cluster 50, pale blue pseudocolour) are shown. **D.** CD4$^+$CD57$^+$PD1$^+$ T cells are more numerous

in CLAD lung tissue. Data from 3 regions of interest (ROIs) from each of n=8 CLAD, and

n=4 normal lungs are shown. **E.** In CLAD lung tissue, CD4$^+$CD57$^+$PD1$^+$ T cells (cluster

50, white) are found in association with airways (green pseudocolour E-cadherin$^+$ cells)

and with two distinct B cell populations (cluster 8, pale blue pseudocolour and cluster

30, purple pseudocolour). **F.** Cell-cell interaction heatmap shows association of

CD4$^+$CD57$^+$PD1$^+$ T cells (cluster 50) with epithelial cells (cluster 27) and B cells

(clusters 8 and 30). G. Liquid phase mass cytometry performed on single cell

suspensions obtained from CLAD lung tissue show variable presence of

CD4$^+$CD57$^+$PD1$^+$ T cells.

Figure 4. CD4$^+$CD57$^+$PD1$^+$ T cells are associated with fibrosis in a mouse model of

chronic lung allograft rejection. A. C57BL/10 (B10) donor lungs were subjected to

either prolonged (6h cold ischemic time (CIT) at 4°C followed by 1h warm ischemic time

(WIT) at 37°C) or minimal (60min CIT followed by 15min WIT) storage prior to orthotopic

left lung transplantation into minor alloantigen-mismatched C57BL/6 (B6) recipients. **B.**

After 28 days, paraffin-embedded sections of lung tissue were subjected to Masson

trichrome staining (200x magnification). **C.** Grafts were scored using a published

scoring system[28] that quantifies the percentage of the graft affected by interstitial

inflammation and parenchymal fibrosis. Grafts with high-grade pathology (High Path)

are represented as triangles, and those with minimal pathology (Low Path) are

represented as circles. **D.** Single-cell suspensions of naïve B6 lungs (No OLT) and lung

allografts obtained at 28 days post-transplantation were subjected to flow cytometry.

Analyses are gated on CD4$^+$ T cells. Percentages of cells in the CD57$^+$PD-1$^+$ gate are

shown in the representative plots. **E.** Percentage of CD4$^+$ T cells expressing CD57 and

PD-1 is shown for naïve B6 lungs (No OLT, squares), Low Path grafts (circles) and High

Path grafts (triangles). Student's t test, p=0.0005. **F.** Percentage of CD4$^+$ T cells

expressing CD57 and PD-1 is shown for Low Path grafts (circles) and High Path grafts

(triangles). Student's t test, p=0.0205. **G.** Immunofluorescence of prolonged storage,

High Path lung allograft showing PD1 (green), CD57 (red), CD4 (yellow) and DAPI

(blue) and merged image.

Figure 5. Identification of distinct CD27⁺ and CD27⁻ subsets of CD4⁺CD57⁺PD1⁺ T cells. Sequencing by CITE-Seq was performed on 4 BAL samples from patients with ALAD. **A.** After downsampling and integration, BAL CD4⁺ T cells (CD3⁺CD4⁺CD8⁻ identified by antibody-derived oligonucleotide tag, adt) were displayed on a tSNE plot revealing 10 distinct cell clusters (left tSNE plot). Feature plot showing CD57 adt and PD1 adt labeling in CD4⁺ cells as indicated by blue dots inside. Three distinct clusters with detectable CD57 and PD1 protein expression are identified (1, 2, 4); however, clusters 1 and 2 have combined as they have similar gene profile. **B.** tSNE plot demonstrating CD27 expression in in RNA level (left) and protein level (adt, right). **C.** Dot plot of the selected top differentially expressed genes in BAL CD4⁺ T cells showing the differences between CD27+ and CD27- associated gene profile. CD27- cells contains LAG3⁺ CXCL13⁺ TNFRSF18⁺ population suggesting putative peripheral helper function while CD27⁺ cells expressing GZMK⁺ NKG7⁺ EOMES⁺ cells implying putative cytotoxic CD4⁺ T cells. **D.** tSNE plot highlighted by size of TCR expansion where yellow is large TCR expansion and purple is single TCR. Selected tSNE with red dotted circle shows highest frequency of expanded TCR in CD4+ cells. **E.** tSNE plot highlighted by velocity trajectory shown by arrows. Same high frequency of TCR mentioned in D, was identified red dotted circle implying cells from cluster 1, 2, and 4 originated from the same cells.

Figure 6. Functional properties of CD4⁺CD57⁺PD1⁺ T cells. A. CD4⁺CD57⁺PD1⁺ T cells can be identified by conventional flow cytometry. BAL cells were stained with CD4, CD57 and PD1 antibodies and CD57⁺PD1⁺ cells were identified within the CD4⁺ population, allowing quantification and cell sorting. **B**. Percentage of CD57⁺PD1⁺ cells within the CD4⁺ gate is shown for n=14 patients with ALAD. **C.** Sorted CD57⁺PD1⁺ and CD57⁻PD1⁻ CD4⁺ T cells from ALAD BAL samples were stimulated *in vitro* with anti-CD3/anti-CD28 antibodies overnight. Culture supernatants were subjected to multiplexed cytokine quantification. IL-21 (p=0.031), IL-4 (p=0.062), CXCL8 (IL-8, p=0.031) and GZMA (p=0.062) were generally secreted in greater concentrations by CD57⁺PD1⁺ T cells than their CD57⁻PD1⁻ counterparts, Wilcoxon matched-pairs signed rank test. **D.** Example of sorted CD57⁺PD1⁺ (blue) and CD57⁻PD1⁻ CD4⁺ (pink) T cells from ALAD BAL were examined for proliferation by dilution of CFSE. CD57⁻PD1⁻ CD4⁺

(pink) demonstrated potential to respond to CD3/CD28 stimulation by proliferation as

shown by several CFSE dilution picks. However, CD57$^+$PD1$^+$ cells did not proliferate

upon CD3/CD28 stimulation. **E.** Percentage of proliferation of CD57$^+$PD1$^+$ are

significantly lower compared to their CD57$^-$PD1$^-$ counterparts. Linked CD57$^+$PD1$^+$ and

CD57$^-$PD1$^-$ cells are sorted from same LT recipient. **F.** Example of sorted CD57$^+$PD1$^+$

and CD57$^-$PD1$^-$ counterpart T cells subjected to RICD assay demonstrated apoptosis in

response to CD3 restimulation in CD57$^-$PD1$^-$ while CD57$^+$PD1$^+$ are irresponsive. **G.**

Percentage of Annexin V as a marker apoptosis are significantly lower in CD57$^+$PD1$^+$

compared to their CD57$^-$PD1$^-$ counterparts. Linked CD57$^+$PD1$^+$ and CD57$^-$PD1$^-$ cells

are sorted from same LT recipient.

Supplementary figure legends

Supp Fig 1. Study flow diagram for the BAL CyTOF cohort.

Supp Fig 2. Animated association between lung allograft dysfunction and

appearance of CD57+PD1+ T cells in individuals. 50 LT recipient in CyTOF cohort

divided into two groups, discovery (1-25) and validation (26-50). For each individual LT

recipient, different events have been demonstrated. Events are included of appearance

CD57$^+$PD1$^+$ T cells (as blue cross), ALAD as blue triangle and CLAD as red triangle.

Orange line indicates time post-transplant with no exact timing relevance.

Supp Fig 3. CD4$^+$CD57$^+$PD1$^+$ T cells are associated with an increased risk of CLAD

and are found in CLAD lung tissue. A. Kaplan-Meier curves showing time to CLAD**.**

B. Imaging mass cytometry heatmap of 50 clusters (y axis) and their marker expression

in x axis.

Supp Fig 4. Identification of distinct CD27$^+$ and CD27$^-$ subsets of CD4$^+$CD57$^+$PD1$^+$

T cells. Sequencing by cellular indexing of transcriptomes and epitopes (CITE-Seq)

was performed on 4 BAL samples from patients with ALAD. **A.** After downsampling and

integration, BAL CD4$^+$ T cells (CD3$^+$CD4$^+$CD8$^-$ identified by antibody-derived

oligonucleotide tag, adt) were displayed on a tSNE plot revealing 10 distinct cell

clusters. Feature plot showing CD57 adt labeling in CD4+ cells as indicated by lavender

dots inside red circles. Three distinct clusters with detectable CD57 protein expression

are identified (1, 2, 4). **B.** Heat map of the top differentially expressed genes in BAL

CD4$^+$ T cells, with clusters identified in B highlighted in red boxes. Cluster 1 contains a

CD27$^-$ LAG3$^+$ CXCL13$^+$ TNFRSF18$^+$ population (putative follicular helper function).

Cluster 2 contains CD27-GZMA+CXCR6+CCL5+ cells (putative mucosa-associated

invariant T cells). Cluster 4 contains CD27$^+$GZMK$^+$NKG7$^+$EOMES$^+$ cells (putative

cytotoxic CD4+ T cells). **C.** Feature plot of selected genes related to high association

with Cluster 1, 2, and 4.

Supp Fig 5. TCR analysis indicates high similarities between Clusters 1 and 2. A.

Quantitative clonal expansion of TCRseq data suggested highest expansion from

cluster 2 (129 large clonal expansion). **B.** Top 20 expanded clones for each clusters

suggested high similarities between expanded clones of Cluster 1 and 2 while there is

limited similarities among those clones and top 20 clones of cluster 4.

<u>**Supplementary table legends**</u>

Supp Table 1. Antibody panel for CyTOF. Highlighted channels are intracellular

markers.

Supp Table 2. Clinical characteristics of participants. Patients were randomized into

a discovery group (n=25) and a validation group (n=25).

Supp Table 3. Clinical and bronchoscopic parameters associated with ≥ 7.8%

CD57$^+$PD1$^+$ T cells in the BAL CD4$^+$ T cell compartment. Fisher's exact tests were

performed. 1Missing samples excluded. 2Comparison of normal to abnormal cytology.

3Comparison of A0/AX with A1 or greater. 4Testing not done (n.d.) due to rarity of small

airway inflammation in the cohort. 5Comparison of any antimicrobial therapy pre- or

post-bronchoscopy vs. none. 6Testing not done (n.d.) due to small number of events.

*Eosinophilia noted in one of the samples with lymphocytosis. **Eosinophilia noted in

isolation in one case and in another sample with neutrophilia. †One patient received

both pre-bronchoscopy antibiotics and post-bronchoscopy antivirals.

Supp Table 4. Change in immunosuppression in patients with and without ≥ 7.8%

CD57$^+$PD1$^+$ T cells in the CD4$^+$ T cell compartment of at least one BAL

sample. Fisher's exact tests were performed. ATG, anti-thymocyte globulin.

Supp Table 5. Statistical analysis in patients with and without ≥ 7.8% CD57$^+$PD1$^+$

T cells in the CD4$^+$ T cell.

Supp Table 6. Antibody panel for IMC.

Supp Table 7. Antibody panel for CITE-seq.

References

Ali, J. M., Bolton, E. M., Bradley, J. A. & Pettigrew, G. J. Allorecognition pathways in transplant rejection and tolerance. *Transplantation* **96**, 681-688, doi:10.1097/TP.0b013e31829853ce (2013).

Chambers, D. C. *et al.* The International Thoracic Organ Transplant Registry of the International Society for Heart and Lung Transplantation: Thirty-eighth adult lung transplantation report - 2021; Focus on recipient characteristics. *J Heart Lung Transplant* **40**, 1060-1072, doi:10.1016/j.healun.2021.07.021 (2021).

Kawashima, M. & Juvet, S. C. The role of innate immunity in the long-term outcome of lung transplantation. *Ann Transl Med* **8**, 412, doi:10.21037/atm.2020.03.20 (2020).

Bando, K. *et al.* Obliterative bronchiolitis after lung and heart-lung transplantation. An analysis of risk factors and management. *The Journal of thoracic and cardiovascular surgery* **110**, 4-13; discussion 13-14 (1995).

Heng, D. *et al.* Bronchiolitis obliterans syndrome: incidence, natural history, prognosis, and risk factors. *J Heart Lung Transplant* **17**, 1255-1263 (1998).

Moshkelgosha, S. *et al.* Interferon-stimulated and metallothionein-expressing macrophages are associated with acute and chronic allograft dysfunction after lung transplantation. *J Heart Lung Transplant* **41**, 1556-1569, doi:10.1016/j.healun.2022.05.005 (2022).

Bhorade, S. M. *et al.* Interobserver variability in grading transbronchial lung biopsy specimens after lung transplantation. *Chest* **143**, 1717-1724, doi:10.1378/chest.12-2107 (2013).

Stephenson, A. *et al.* Interpretation of transbronchial lung biopsies from lung transplant recipients: inter- and intraobserver agreement. *Can Respir J* **12**, 75-77 (2005).

Levy, L. *et al.* in *International Society for Heart and Lung Transplantation 39th Annual Meeting* Vol. 38(S4)   S163-164 (J Heart Lung Transplant, Orlando FL USA, 2019).

Levy, L. *et al.* Epithelial cell death markers in bronchoalveolar lavage correlate with chronic lung allograft dysfunction subtypes and survival in lung transplant recipients-a single-center retrospective cohort study. *Transpl Int* **32**, 965-973, doi:10.1111/tri.13444 (2019).

Moshkelgosha, S. *et al.* in *J Heart Lung Transpl* Vol. 37   S216 (2018).

Moshkelgosha, S. *et al.* in *International Society for Heart and Lung Transplantation* Vol. 38   S17 (Orlando, FL USA, 2019).

Moshkelgosha, S. *et al.* in *J Heart Lung Transpl* Vol. 39   S199-S200 (2020).

Whitehead, B. F. *et al.* Analysis of bronchoalveolar lavage from human lung transplant recipients by flow cytometry. *Respiratory medicine* **89**, 27-34 (1995).

Reynaud-Gaubert, M. *et al.* Clinical utility of bronchoalveolar lavage cell phenotype analyses in the postoperative monitoring of lung transplant recipients. *European journal of cardio-thoracic surgery : official journal of the European Association for Cardio-thoracic Surgery* **21**, 60-66 (2002).

Hodge, S. *et al.* Increased levels of T cell granzyme b in bronchiolitis obliterans syndrome are not suppressed adequately by current immunosuppressive regimens. *Clinical and experimental immunology* **158**, 230-236, doi:10.1111/j.1365-2249.2009.04008.x (2009).

Neujahr, D. C. *et al.* Dynamics of human regulatory T cells in lung lavages of lung transplant recipients. *Transplantation* **88**, 521-527, doi:10.1097/TP.0b013e3181b0e719 (2009).

Hodge, S. J., Hodge, G. L., Reynolds, P. N. & Holmes, M. D. Differential rates of apoptosis in bronchoalveolar lavage and blood of lung transplant patients. *J Heart Lung Transplant* **24**, 1305-1314, doi:10.1016/j.healun.2004.10.009 (2005).

Greenland, J. R. *et al.* Bronchoalveolar lavage cell immunophenotyping facilitates diagnosis of lung allograft rejection. *Am J Transplant* **14**, 831-840, doi:10.1111/ajt.12630 (2014).

Bandura, D. R. *et al.* Mass cytometry: technique for real time single cell multitarget immunoassay based on inductively coupled plasma time-of-flight mass spectrometry. *Analytical chemistry* **81**, 6813-6822, doi:10.1021/ac901049w (2009).

Van Gassen, S. *et al.* FlowSOM: Using self-organizing maps for visualization and interpretation of cytometry data. *Cytometry A* **87**, 636-645, doi:10.1002/cyto.a.22625 (2015).

Bruggner, R. V., Bodenmiller, B., Dill, D. L., Tibshirani, R. J. & Nolan, G. P. Automated identification of stratifying signatures in cellular subpopulations. *Proc Natl Acad Sci U S A* **111**, E2770-2777, doi:10.1073/pnas.1408792111 (2014).

Nowicka, M. *et al.* CyTOF workflow: differential discovery in high-throughput high-dimensional cytometry datasets. *F1000Res* **6**, 748, doi:10.12688/f1000research.11622.2 (2017).

Tikkanen, J. M. *et al.* De Novo DQ Donor-Specific Antibodies Are Associated with Chronic Lung Allograft Dysfunction after Lung Transplantation. *American journal of respiratory and critical care medicine* **194**, 596-606, doi:10.1164/rccm.201509-1857OC (2016).

Renaud-Picard, B. *et al.* in *J Heart Lung Transpl* Vol. 41   S106 (2022).

Giesen, C. *et al.* Highly multiplexed imaging of tumor tissues with subcellular resolution by mass cytometry. *Nature methods* **11**, 417-422, doi:10.1038/nmeth.2869 (2014).

Watanabe, T. *et al.* A B cell-dependent pathway drives chronic lung allograft rejection after ischemia-reperfusion injury in mice. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* **19**, 3377-3389, doi:10.1111/ajt.15550 (2019).

Martinu, T. *et al.* Spectrum of chronic lung allograft pathology in a mouse minor-mismatched orthotopic lung transplant model. *Am J Transplant* **19**, 247-258, doi:10.1111/ajt.15167 (2019).

Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single cells. *Nature methods* **14**, 865-868, doi:10.1038/nmeth.4380 (2017).

Rao, D. A. *et al.* Pathologically expanded peripheral T helper cell subset drives B cells in rheumatoid arthritis. *Nature* **542**, 110-114, doi:10.1038/nature20810 (2017).

Manzo, A. *et al.* Mature antigen-experienced T helper cells synthesize and secrete the B cell chemoattractant CXCL13 in the inflammatory environment of the rheumatoid joint. *Arthritis Rheum* **58**, 3377-3387, doi:10.1002/art.23966 (2008).

Huang, Y. *et al.* T peripheral helper cells in autoimmune diseases: What do we know? *Front Immunol* **14**, 1145573, doi:10.3389/fimmu.2023.1145573 (2023).

Ng, S. S. *et al.* The NK cell granule protein NKG7 regulates cytotoxic granule exocytosis and inflammation. *Nat Immunol* **21**, 1205-1218, doi:10.1038/s41590-020-0758-6 (2020).

Wein, A. N. *et al.* CXCR6 regulates localization of tissue-resident memory CD8 T cells to the airways. *J Exp Med* **216**, 2748-2762, doi:10.1084/jem.20181308 (2019).

Milardi, G. *et al.* Follicular helper T cell signature of replicative exhaustion, apoptosis, and senescence in common variable immunodeficiency. *Eur J Immunol* **52**, 1171-1189, doi:10.1002/eji.202149480 (2022).

Llao-Cid, L. *et al.* EOMES is essential for antitumor activity of CD8(+) T cells in chronic lymphocytic leukemia. *Leukemia : official journal of the Leukemia Society of America, Leukemia Research Fund, U.K* **35**, 3152-3162, doi:10.1038/s41375-021-01198-1 (2021).

Wolf, Y., Anderson, A. C. & Kuchroo, V. K. TIM3 comes of age as an inhibitory receptor. *Nature reviews. Immunology* **20**, 173-185, doi:10.1038/s41577-019-0224-6 (2020).

Liu, X., Hoft, D. F. & Peng, G. Senescent T cells within suppressive tumor microenvironments: emerging target for tumor immunotherapy. *J Clin Invest* **130**, 1073-1083, doi:10.1172/JCI133679 (2020).

Mittelbrunn, M. & Kroemer, G. Hallmarks of T cell aging. *Nat Immunol* **22**, 687-698, doi:10.1038/s41590-021-00927-z (2021).

Hodge, G., Hodge, S., Reynolds, P. N. & Holmes, M. Airway infection in stable lung transplant patients is associated with decreased intracellular T-helper type 1 pro-inflammatory cytokines in bronchoalveolar lavage T-cell subsets. *Transplant infectious disease : an official journal of the Transplantation Society* **10**, 99-105, doi:10.1111/j.1399-3062.2007.00236.x (2008).

Hodge, G., Hodge, S., Chambers, D., Reynolds, P. N. & Holmes, M. Acute lung transplant rejection is associated with localized increase in T-cell IFNgamma and TNFalpha proinflammatory cytokines in the airways. *Transplantation* **84**, 1452-1458, doi:10.1097/01.tp.0000290679.94163.e1 (2007).

Hodge, G., Hodge, S., Reynolds, P. N. & Holmes, M. Compartmentalization of intracellular proinflammatory cytokines in bronchial intraepithelial T cells of stable lung transplant patients. *Clinical and experimental immunology* **145**, 413-419, doi:10.1111/j.1365-2249.2006.03143.x (2006).

Hodge, G., Hodge, S., Reynolds, P. N. & Holmes, M. Increased intracellular pro- and anti-inflammatory cytokines in bronchoalveolar lavage T cells of stable lung transplant patients. *Transplantation* **80**, 1040-1045 (2005).

Libri, V. *et al.* Cytomegalovirus infection induces the accumulation of short-lived, multifunctional CD4+CD45RA+CD27+ T cells: the potential involvement of interleukin-7 in this process. *Immunology* **132**, 326-339, doi:10.1111/j.1365-2567.2010.03386.x (2011).

Di Mitri, D. *et al.* Reversible senescence in human CD4+CD45RA+CD27- memory T cells. *J Immunol* **187**, 2093-2100, doi:10.4049/jimmunol.1100978 (2011).

Martinu, T. *et al.* International Society for Heart and Lung Transplantation consensus statement for the standardization of bronchoalveolar lavage in lung transplantation. *J Heart Lung Transplant*, doi:10.1016/j.healun.2020.07.006 (2020).

Amir el, A. D. *et al.* viSNE enables visualization of high dimensional single-cell data and reveals phenotypic heterogeneity of leukemia. *Nat Biotechnol* **31**, 545-552, doi:10.1038/nbt.2594 (2013).

Schapiro, D. *et al.* histoCAT: analysis of cell phenotypes and interactions in multiplex image cytometry data. *Nat Methods* **14**, 873-876, doi:10.1038/nmeth.4391 (2017).

Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nature communications* **8**, 14049, doi:10.1038/ncomms14049 (2017).

Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nature biotechnology* **36**, 411-420, doi:10.1038/nbt.4096 (2018).

Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data using regularized negative binomial regression. *Genome Biology* **20**, 296, doi:10.1186/s13059-019-1874-1 (2019).

La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498, doi:10.1038/s41586-018-0414-6 (2018).

Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to transient cell states through dynamical modeling. *Nature biotechnology* **38**, 1408-1414, doi:10.1038/s41587-020-0591-3 (2020).

Katz, G. & Snow, A. L. Fluorescence-activated cell sorting-based quantitation of T cell receptor restimulation-induced cell death in activated, primary human T cells. *Methods Mol Biol* **979**, 15-23, doi:10.1007/978-1-62703-290-2_2 (2013).

A

Donor C57BL/10   *Prolonged storage*   Recipient C57BL/6

CIT 4°C 6h   WIT 37°C 1h

or 60min   *or* 15min

Minimal storage

B Prolonged Storage   Minimal Storage

C

% Parenchymal Fibrosis vs % Interstitial Inflammation

High Path

Low Path

D

No OLT   Low Path   High Path

PD1-BV605 / CD57-APC

0.084%   6.81%   13.4%

E

% CD57+ PD1+ of total CD4+ T Cells

0.0005

No OLT   OLT

F

% CD57+ PD1+ of total CD4+ T Cells

0.0205

Low Path   High Path

G

DAPI   PD1   CD57   CD4   merge

50 µm

A

ADT-CD4⁺CD8⁻

ADT-PD1&CD57

CD4⁺
CD4⁺CD57⁺PD1⁺

B

CD27

ADT-CD27

C

CD27⁺

CD27⁻

CXCL13 TOX CD40LG IL21 LAG3 TNFRSF18 CCR6 KLRG1 TIGIT CXCR6 GZMK GZMA CST7 NKG7 EOMES CXCR3 CCL5 HAVCR2

Average Expression

0.4
0.0
-0.4

Percent Expressed

20
40
60
80

D

Large (20 < X <= 100)
Medium (5 < X <= 20)
Single (0 < X <= 1)
Small (1 < X <= 5)
NA

E