Role of Ethanolamine Utilization and Bacterial Microcompartment Formation in *Listeria monocytogenes* Intracellular Infection

Ayan Chatterjee[a], Karan Gautam Kaval[a]*, and Danielle A. Garsin[a]#

[a]Department of Microbiology and Molecular Genetics, The University of Texas Health Science Center at Houston, Houston, Texas, USA

Running Title: ethanolamine utilization in *L. monocytogenes*

#Address correspondence to Danielle A. Garsin, Danielle.A.Garsin@uth.tmc.edu

*Present address: Department of Molecular Biology, University of Texas Southwestern Medical Center, Dallas, Texas, USA

ABSTRACT

Ethanolamine (EA) affects the colonization and pathogenicity of certain human bacterial pathogens in the gastrointestinal tract. However, EA can also affect the intracellular survival and replication of host-cell invasive bacteria such as *Listeria monocytogenes* (LMO) and *Salmonella enterica* serovar Typhimurium (*S.* Typhimurium). The EA utilization (*eut)* genes can be categorized as regulatory, enzymatic, or structural, and previous work in LMO showed that loss of genes encoding functions for the enzymatic breakdown of EA inhibited LMO intracellular replication. In this work, we sought to further characterize the role of EA utilization during LMO infection of host cells. Unlike what was previously observed for *S.* Typhimurium, in LMO, an EA regulator mutant (*ΔeutV)* was equally deficient in intracellular replication compared to an EA metabolism mutant (*ΔeutB*)*,* and this was consistent across Caco-2, RAW 264.7 and THP-1 cell lines. The structural genes encode proteins that self-assemble into bacterial microcompartments (BMCs) that encase the enzymes necessary for EA metabolism. For the first time, native EUT BMCs were fluorescently tagged, and EUT BMC formation was observed in vitro, and in vivo. Interestingly, BMC formation was observed in bacteria infecting Caco-2 cells, but not the macrophage cell lines. Finally, the cellular immune response of Caco-2 cells to infection with *eut* mutants was examined, and it was discovered that *ΔeutB* and *ΔeutV* mutants similarly elevated the expression of inflammatory cytokines. In conclusion, EA sensing and utilization during LMO intracellular infection are important for optimal LMO replication and immune evasion but are not always concomitant with BMC formation.

INTRODUCTION

Ethanolamine (EA) is a metabolically useful compound that arises from the membrane lipid phosphatidylethanolamine contained in all bacterial and eukaryotic cells (1, 2). Because of the natural turnover of the epithelium and the cells of the microbiota, the gastrointestinal environment is particularly rich in EA with concentrations as high as 2mM (3, 4). As this compound can serve as a useful source of carbon and/or nitrogen, there is a correlation between bacteria that contain the genes to catabolize this EA and those found in the gut including species of *Enterococcus, Escherichia, Clostridium, Salmonella* and *Listeria* (reviewed by (5, 6)). However, expression of the *eut* genes has been documented in other host environments, including the intracellular one, and may be of relevance for *Listeria monocytogenes* (LMO) (7, 8).

LMO is a facultative, intracellular pathogen, capable of invading, surviving, and multiplying inside various host cell types including cells of the gastrointestinal epithelium and the innate immune cells. Upon attachment, invasion is mediated by engulfment and internalization into a vacuole. Depending on the host cell environment, LMO can disrupt the phagosome, escape into the host cytoplasm, replicate, and spread using actin-based motility (reviewed by (9)). In the host environment of human colon epithelial (Caco-2) cells, a microarray analysis found that the *eut* operon of LMO was upregulated (8). Furthermore, a study of LMO gene expression under several different host conditions, discovered strong upregulation of the *eut* genes in the intestines of infected mice and weaker, but still significant levels of expression in human blood (7).

The EA utilization genes are usually found together in an operon or locus and encode the

necessary regulators, enzymes, and structural proteins (5, 6)). Specifically, *Listeria*

monocytogenes possesses 17 genes involved in EA metabolism (Fig. 1A) (10). The regulation

of the *eut* genes for this bacterium and other Gram-positives such as *Enterococcus* involves an

EA-responsive sensor kinase, encoded by *eutW,* a response regulator of the ANTAR family,

encoded by *eutV,* and an AdoCbl-dependent riboswitch (10–13). To metabolize EA, an

ammonia lyase, composed of two sub-units encoded by *eutBC*, is required for the initial

catabolic step that generates acetaldehyde and ammonia. To concentrate these products for the

downstream reactions and to prevent interaction of toxic acetaldehyde with the cytoplasm,

some, including LMO, also encode for structural proteins that self-assemble into enclosed,

proteinaceous structures called bacterial microcompartments (BMCs) (reviewed by (5, 6)).

In addition to upregulation of the *eut* genes, a contribution of ethanolamine metabolism to LMO

intracellular replication in Caco-2 cells was noted; a *eutB* mutant was deficient compared to the

parental strain (8). Furthermore, in an intravenous (i.v.) mouse infection model, a *eutB* mutant

was deficient in dissemination to the spleen and liver (10). Since EutB is one of the subunits that

comprise the ammonia lyase, these data suggest EA catabolism promotes survival in both

gastrointestinal epithelial cells and during bloodstream infection (8, 10). Interestingly and in

contrast, replication of a *eutB* mutant strain of *Salmonella enterica* serovar Typhimurium was

not deficient in host cells. However, replication was diminished for a strain deleted for *eutR*,

which encodes the EA-sensing transcriptional regulator (14). This was attributed to the fact that

EutR regulates genes associated with pathogenicity in *S.* Typhymurium – the SPI-2

pathogenicity island. Further work by the Kendall group established the importance of the EutH

permease for *S.* Typhimurium (15). EutH is required for EA entry under acidic conditions such

as those encountered in the phagosome (16). Interestingly, a slight phenotype for a *eutH* mutant

in intracellular replication was also shown for LMO, but regulatory or metabolism mutants were

not directly compared (15). Whether a LMO *eutV* mutant strain would display a greater

deficiency than a *eutB* mutant strain in intracellular replication like *S.* Typhimurium is unknown.

As mentioned above, EA metabolism is frequently, but not always encased in an organelle-like

structure called a bacterial microcompartment (BMC) (reviewed by (6, 17, 18)). Rather than

having a selectively permeable membrane, these "organelles" have a selectively permeable

protein shell made up of structural components. A feature of the assembly is that some of the

enzymatic proteins have "encapsulation peptides" (EPs) that interact with the structural proteins

to enable BMC formation. Native EUT BMCs have been visualized in several species of bacteria

including LMO by transmission electron microscopy (TEM) (19). However, fluorescent-tagging

of EUT BMCs to our knowledge was only done in a heterologous system in which *eut* genes

encoding the structural proteins of *S.* Typhimurium were over-expressed in *E. coli* using a

constitutively active promoter, and BMCs were visualized by tagging an encapsulation peptide

with GFP (20–22). Here we report the fluorescent tagging of native EUT proteins under the

control of *eut* promoters to permit the visualization of endogenous BMC formation in vitro and in

host cells.

In this work, we sought to further characterize the role of EA utilization during LMO infection of

host cells. Unlike what was previously observed with *S.* Typhimurium (14), an EA metabolism

mutant (*ΔeutB*) was equally deficient in intracellular replication compared to an EA regulator

mutant (*ΔeutV),* and this was consistent across a number of cell types. To visualize EUT BMC

formation, native structural proteins were fluorescently tagged and expressed under *eut*

promoters. EUT BMC formation was observed in vitro, and for the first time, in vivo, in some,

but not all host cells tested. Interestingly, formation of EUT BMCs was associated with the

phagosomal phase of infection. Finally, it was found that LMO *ΔeutB* and *ΔeutV* mutants

similarly affected the cellular immune response; comparable increases in several inflammatory

cytokines were observed.

RESULTS

Establishment of conditions in which LMO can utilize ethanolamine (EA) as a nitrogen source dependent on the *eut* genes:

Previous work demonstrated that LMO can utilize EA as a source of carbon and/or nitrogen depending on the media conditions (19, 23). We optimized a similar chemically defined medium that we named MMWB (Modified MWB) (Table S1) (24–27). MMWB contained 4.1mM glutamine (Gln) as the nitrogen source, allowing for robust growth of LMO (Fig. 1B). EA could be substituted for Gln, which requires the bacteria to utilize EA as a nitrogen source. As expected, no growth was observed in the absence of Gln, underscoring the necessity of a nitrogen source (Fig. 1B). To test if EA could substitute as a nitrogen source in the medium, we replaced Gln with 16mM EA. However, no growth was observed until we additionally supplemented the medium with 200µM of AdoCbl (Fig. 1B). The data aligns with previous research demonstrating a requirement for AdoCbl as a co-factor in the metabolism of EA and in the induction of the *eut* genes (reviewed by (6)), and all subsequent experiments with media conditions labeled "EA" included AdoCbl. In conclusion, we show that LMO can grow in MMWB utilizing EA as a nitrogen source.

As described in the introduction and illustrated in Fig. 1A, the *eut* locus of LMO contains 17 genes that encode the regulators, enzymes, and structural shell proteins necessary for EA metabolism. To verify induction of these genes in MMWB + EA, expression of a subset was measured by qRT-PCR. As expected, all genes measured, *eutV, A, B, K,* and *M,* displayed increased expression following 4 hours of growth in MMWB + EA relative to MMWB + Gln. Of note, the expression was specific for *eut* genes associated with EA metabolism; genes associated with propanediol (PDU) metabolism, *pduA* and *pduQ,* were not induced (Fig 1C).

To further verify the role of the *eut* genes in LMO's use of EA as a nitrogen source in MMWB,

we generated strains containing *ΔeutV* (LMAC030) and *ΔeutB* (LMAC031) non-polar, in-frame,

deletion mutations in the parental background (Fig 1D and 1E). When grown in the presence of

Gln as the nitrogen source, these mutant strains exhibited growth comparable to the parental

strain. However, they did not grow when EA was provided as a nitrogen source, indicating that

the response regulator, EutV, and the ammonia lyase subunit, EutB, were indispensable for

growth in the presence of EA in vitro, as expected. To ensure that the defects in growth were

due to the introduced mutations, we complemented these strains. In order to retain EA-

inducibility, the complementation constructs included the regulatory sequences upstream of

eutV or *eutG.* Both these regions contain predicted promoters followed by ANTAR binding sites

that are the hallmarks of antitermination regulation by EutV (Fig 1A) (10). Growth in EA was

partially restored for the *ΔeutV* strain when *eutV,* under the control of the *eutV* promoter, was

introduced as a transgene into a chromosomal site outside of the EA locus (see Materials and

Methods for details), (Fig 1D and S1A). Note that successful complementation of *ΔeutV*

required using the combined *eutVW* sequence; attempts to complement with the *eutV* sequence

alone were unsuccessful. As the *eutV* and *eutW* genes overlap and appear to be co-transcribed,

it is possible that a *eutV* transcript alone is unstable. Growth in EA was also restored in the

ΔeutV background, but to a lesser degree, when the *eutG* promoter was utilized (Fig S1A). In

contrast, successful complementation was only achieved in the *ΔeutB* strain with the *eutG*

promoter driving the expression of *eutB*; use of the *eutV* promoter did not enable growth in EA

(Fig 1E and S1B). We speculate that these differences in the promoters' abilities to restore

growth in EA is due to differences in the levels of EutV and EutB that are required for EA

metabolism and achieved by the promoters. Overall, these results established conditions in

which LMO can utilize EA as a source of nitrogen dependent on the positive regulation of the

eut genes (EutV) and the ability to metabolize EA (EutB).

LMO EUT BMC formation is dependent on EA, EutV, and EutB:

EA utilization has been associated with BMC formation for many EA-utilizing bacteria including

L. monocytogenes (6, 19). Specifically, LMO EUT BMCs were previously visualized under

anaerobic conditions (19). To test for BMC formation in aerobic MMWB + EA conditions,

transmission electron microscopy (TEM) was utilized. Structures resembling BMCs were readily

visible in the parental LMO background following 6 hours growth in medium containing EA but

not Gln (Fig. 2A) with 35.3% of the cells examined (n=51) containing at least one visible BMC.

(Note that this count is likely an underestimate due to the fact that a given plane of BMC-

containing cell may not contain a BMC.) As dependence of the formation of these structures on

EutV and EutB had not been previously reported in LMO, we also examined *ΔeutV* and *ΔeutB*

deletion mutant strains grown in EA; both strains failed to form any visible BMCs (Fig. 2B and

2C). The complemented strains both restored BMC formation, as predicted, indicating that the

deletions did not have polar effects on the expression of the downstream structural genes (Fig.

2B, 2C and 1A).   Specifically, 37.5% and 38.4% of the *eutV* (n=24) and *eutB* (n=26)

complemented cells presented with at least one visible BMC.

In order to visualize EUT BMC formation in live cells, we introduced a *eutK::mCherry*

translational fusion under the control of the EA inducible *eutG* promoter into an ectopic,

chromosomal site in wild-type, *ΔeutV* and *ΔeutB* backgrounds (Materials and Methods). Tags

were not added to the endogenous structural genes, as it was previously reported that this can

disrupt BMC formation (28, 29). To ensure that introduction of the transgene did not disrupt the

ability of the strain to utilize EA, growth curves were performed, and the strain was found to

grow in EA with kinetics similar to the parental strain (Fig. S1C). Following six hours of

incubation in medium containing EA, distinct red punctate structures were observed

corresponding to BMC-like formations in about 85% of the cells (Fig. 3A and 3E). Notably, when

bacterial cells were cultivated in a medium containing Gln, these BMC-like structures were

rarely evident under confocal microscopy. (Fig. 3B and 3E). By performing live cell imaging of

this strain, the dynamics of these structures forming and dissipating was revealed (Mov. S1).

Consistent with the TEM observations, BMCs were not visible in the *ΔeutV* and *ΔeutB* strain

backgrounds (Fig. 3C, 3D and 3E). In addition to the *eutK::mCherry* translational fusion,

eutM::mCherry and a *eutK::mNeonGreen* fusions were additionally generated. They presented

with a similar punctate fluorescent pattern when grown in EA, indicating that the observed

localization was unlikely to be artifactual due to a specific fusion or tag (S2A and S2B). In

conclusion, these experiments demonstrate that EUT BMC formation is dependent on EA, EutV,

and EutB and can be visualized with fluorescent tags in *L. monocytogenes*.

Loss of EA utilization impairs intracellular replication of LMO:

As LMO invades the cells of the gastrointestinal epithelium in humans, the Caco-2 human

colorectal adenocarcinoma cell line is frequently used to model LMO intracellular infections (30,

31). Previous work on strain LMO EGDe documented that a strain with an insertion mutant in

eutB was deficient in Caco-2 intracellular replication (8). To verify the phenotype with LMO

10403S, we infected Caco-2 cells with the parental strain and the *ΔeutV* and *ΔeutB* strains.

Following a one-hour infection at an MOI of 5, the cells were washed, and medium containing

gentamycin was added for one hour to kill extracellular bacteria. At the zero-time point, defined

as when the extracellular bacteria were removed and the cells washed, the number of

intracellular bacteria between the strains was not significantly different, indicating no defect in

bacterial invasion (Fig. 4A). However, by three hours, there was approximately half-a-log less

CFUs of the *ΔeutV* and *ΔeutB* strains compared to the parent, but no significant difference in

the level of attenuation between the two mutant strains. Note, that at this time point there is a

drop in the number of bacteria compared to the zero-time point. It is thought that CFUs at the

zero-time point can be inflated due to bacterial adherence to the surface of Caco-2 cells despite

the wash step, and this phenomenon has been observed previously during LMO infection (30).

By six hours, the CFUs were increased over the zero-time point, indicative of intracellular

replication, but the attenuation of the *eut* mutants persisted (Fig. 4A).

Following enterocyte invasion, *L. monocytogenes* can infect recruited immune cells such as

macrophages, which can lead to systemic infections (9). A LMO mutant in *eutH,* which encodes

for an EA transporter thought to be necessary for EA utilization in acidic environments, was

previously shown to have a slight defect in intracellular replication in bone marrow-derived

macrophages (BMDMs), suggesting that EA utilization might play a role in this process (15). To

further test the efficiency of intracellular replication of the *eut* mutant strains in immune cells, we

infected RAW 264.7 and differentiated THP-1 cells, which are mouse and human derived

macrophages, respectively (Fig. 4B and 4C). Replication of the *ΔeutV* and *ΔeutB* strains was

attenuated by about 0.5 logs following three-four hours of infection, and the attenuation

persisted at the six-hour time point. Again, there was no significant difference in the degree of

deficiency between the two strains. In conclusion, the *ΔeutV* and *ΔeutB* strains are deficient in

intracellular replication to a similar degree in a range of cell types suggesting that EA utilization

positively contributes to LMO infection efficiency.

LMO forms EUT BMCs during intracellular infection of Caco-2 cells:

To test for BMC formation in the intracellular environment, Caco-2 cells were infected with the

strains expressing the *eutK::mCherry* fusion. Strains containing the *eutK::mCherry* translational

fusion were utilized for the in vivo experiments because they were brighter compared to the

eutM::mCherry strains (Fig. S2A) and were less prone than mNeonGreen to interference from

background autofluorescence (Fig. S2B). The samples were fixed at 2, 4, 6 and 8 hours post-

infection, and bacterial cells were visualized using an antibody specific to *L. monocytogenes.*

Host cell actin was visualized by staining with phalloidin. At 4, 6, and 8 hours, punctate, red

fluorescence inside some of the bacterial cells from the parental strain was visible, indicative of

BMC formation (Fig. 5A and Fig. S3A). As expected, little to no signal was observed in samples

from the *ΔeutV* and *ΔeutB* strains containing the *eutK::mCherry* fusion (Fig. 5B-C and Fig. S3A).

Time points later than 8 hours were not captured due to host cell lysis. However, as shown by

imaging the larger fields (Fig. S3A), not all bacteria within the host cells presented with visible

BMCs. By counting the number of bacteria presenting with and without BMCs in a given image,

it was determined that approximately 25% of the *eutK::mCherry* parental strain of LMO had

BMCs. However, the percentage of bacteria containing BMCs in a given image was variable

and became more so over time (Fig. S3B). As host cell actin will associate with cytoplasmic

LMO, escape from the phagosome can be visualized by the formation of actin clouds and tails

(Fig. 5D). Interestingly, we observed that bacteria associated with actin clouds or tails never

contained visible BMCs. Rather BMC formation was observed in some, but not all, bacteria

unassociated with actin (Figure 5D and S3A). Note that the signal from the antibody against

LMO (green) was weaker in the actin associated bacteria, perhaps due to actin interfering with

antibody binding. Therefore, we have outlined some of the bacteria with actin tails in Fig. 5D to

aid in visualization. Actin tails and clouds also formed during infection with both the *eut* mutant

strains, demonstrating that phagosomal escape is not blocked (Fig S3A). In conclusion, these

data show that in Caco-2 cells EUT BMCs form in LMO*.* Moreover, because BMCs were only

observed in some bacteria not associated with actin, the results suggest that BMC formation

may occur preferentially in the phagosome.

Confocal imaging was also performed on RAW 264.7 cells infected with the *eutK::mCherry*

strain at 2, 4, and 6 hours post-infection (Fig. S4A). Following staining and imaging as

described for the Caco-2 cells, actin tails and clouds were observed at later time points, typical

of intracellular LMO infection. However, despite the *eut* genes contributing to intracellular

replication in these cells (Fig. 3B), we were unable to detect any mCherry signal at any time

point (Fig. S4A). The same experiment was performed using M1 differentiated THP-1 cells,

reported to delay phagosomal escape (32). No mCherry signal was evident during the infection

of these cells at either early or late time points (Fig. S4B). However, normal infection kinetics

were observed with actin clouds and tails visible by five hours. These results suggest that BMCs

do not form in RAW 264.7 or THP-1 cells.

Loss of EA utilization increases host cell inflammatory responses:

To further investigate the role of EA utilization in LMO's interaction with host cells, we examined

the inflammatory response of Caco-2 cells following infection. Specifically, the expression of

genes encoding for inflammatory cytokines, vital signaling mediators that orchestrate and

amplify the immune response by recruiting immune cells and enhancing antimicrobial

capabilities, were measured (33, 34). Caco-2 cell cultures were subjected to infection with the

LMO parental or the *ΔeutV* and *ΔeutB* mutants as before, and host cell RNA was extracted at

four hours post-infection. The expression levels of specific inflammatory cytokines, interleukin-1

beta (IL-1β), tumor necrosis factor-alpha (TNF-α), interleukin-6 (IL-6), interleukin-8 (IL-8), and

chemokines CXCL2 and MCP-1 were examined by qRT-PCR as described previously for Caco-

2 cells (35). A 2-4-fold increase in the expression of these factors by Caco-2 cells infected with

the *ΔeutV* and *ΔeutB* mutant strains was observed in comparison to those infected with the

parental strain (Figure 6). There were no significant differences in the expression levels of these

cytokines between the *ΔeutV* and *ΔeutB* mutant strains, suggesting that the observed

differences are solely due to the loss of ethanolamine utilization. The stronger inflammatory

response coupled with the reduced intracellular replication observed during infection with the

mutant strains (Figure 3A) suggests that loss of ethanolamine utilization reduces LMO's ability

to evade the host immune system.

DISCUSSION

In this study, we demonstrated that the catabolism of EA contributes to the efficacy of

intracellular infection of LMO in a variety of host cells. In *S.* Typhimurium, by contrast, the ability

to sense, but not catabolize EA, was shown to enhance intracellular survival in macrophages; a

eutR regulatory mutant was deficient in intracellular survival, but a *eutB* metabolic mutant was

not. (14). Here, a direct comparison between LMO *eutV* regulatory and *eutB* metabolic mutants

revealed indistinguishable phenotypes. Both exhibited the same level of depressed intracellular

replication and caused similar changes to the host cell immune response (Fig. 3 and 6). The

difference could be related to the scope of the genes regulated by EutR and EutV. EutR is a

DNA binding transcriptional regulator. In addition to binding its promoter in the *eut* locus, EutR

directly binds the promoter of the *ssrB* gene in *S.* Typhimurium encoding a transcriptional

regulator. SsrB activates the expression of the Salmonella pathogenicity island 2 (SPI-2) genes

that are required for *S.* Typhimurium intracellular replication and survival (14). EutR in EHEC

also regulates virulence related genes outside of the *eut* locus (36, 37). In contrast, EutV is an

ANTAR (AmiR and NasR transcriptional antiterminator regulators) response regulator that binds

specific sequences in mRNA leaders that overlap terminators, preventing their formation (38,

39) In LMO, there is no evidence that EutV directly regulates genes other than those related to

EA utilization (10, 38). The observation that strains containing *eutV* and *eutB* deletions caused

identical effects on LMO intracellular replication as well as on host cell immune responses is

consistent with the phenotypes being caused by the loss of the ability to catabolize EA in this

bacterium, unlike in *S.* Typhimurium. How exactly the loss of EA metabolism causes these

changes in LMO's interactions with host cells is of future interest.

By fluorescently tagging BMC structural proteins, we were able to visualize BMC formation in

vitro and in vivo. We did observe BMCs forming in a subset of the bacteria inside Caco-2 cells.

Interestingly, these visible BMCs were only in seen in bacteria not associated with actin; there

were no examples of BMCs in bacteria associated with tails or clouds in our imaging

experiments. These data suggest that BMCs might only form during the phagosomal phase of

infection. However, attempts to confirm this observation with different approaches such as

antibodies against phagosomal markers and/or developing antibodies to visualize the BMCs by

immunolabeling, were unsuccessful.

Unlike the Caco-2 cells, no BMCs were observed during LMO infection of RAW 264.7 and

THP-1 cells. We postulate two possible explanations for the lack of visible BMC formation in

these monocyte/macrophage cell lines. One, BMCs were generated, but were unable to be

visualized due to low expression of the *eutK::mCherry* transgene. Two, the metabolism of

ethanolamine is occurring at a lower level in these cells that does not require BMC formation.

Recall that some bacterial species, like *Pseudomonas aeruginosa* and *Acinetobacter baumannii*

encode for the core EA metabolic enzymes, but lack the structural proteins (39, 40). Even in

bacteria such as *S. typhimurium* that contain the full complement of structural proteins, it has

been shown that they are still able to metabolize EA, albeit with less efficiency, when they are

deleted for the microcompartment-encoding genes (41, 42).   Perhaps there are differences in

the amount of free EA available in these different host cells that determine whether BMC

formation is necessary for EA utilization or not. Future work using transcriptomics and

metabolomics to further understand the host cell environments and the bacterial response may

provide more insight into these observations.

Another surprising finding in this study was the lack of BMC formation in the *ΔeutB* mutant. We

expected that no structures would form in the *ΔeutV* mutant that lacks the regulator necessary

for expression of all the *eut* genes but were surprised that a strain missing a single catalytic

enzyme failed to form BMCs. However, it has been reported that the shell proteins of the

carbon-fixing β-carboxysomes condense around key catalytic enzymes necessary for nucleation

of the BMC structure (18, 29). In contrast, the α-carboxysomes can assemble either shell-first

or undergo concomitant shell–core assembly, and the propanediol (PDU) metabolizing BMCs

form independent cargo and shell aggregates (43–45). While there have been studies in

heterologous systems to examine the requirements for EUT BMC assembly (20–22), the

dynamics have not been determined in a natural system. Therefore, it is possible that core

assembly requiring EutB is necessary for shell protein condensation. However, our previous

work in *E. faecalis* documented normal BMC formation by TEM in strains with *eutB* mutations

(46), raising the possibility of species-specific differences. Undoubtedly, it will be of interest to

further study EUT BMC formation and elucidate the assembly pathway(s). More broadly, the

field should investigate if the assembly of BMCs varies in a manner dependent on species and

environmental context, including host environments. In conclusion, EA utilization contributes

importantly to LMO intracellular replication, but may not always require EUT BMC assembly.

MATERIALS AND METHODS

Bacterial strains and media

Listeria monocytogenes 10403s (LMO) strain (source: Bei resources) was used in this study

and all mutations and recombinants were constructed in this background. All LMO strains were

grown in brain heart infusion (BHI) broth or on BHI agar (1.5% agar) plates supplemented with

streptomycin (50 µg /ml) in 37 °C. The chemically defined medium MMWB (Table S1) was

prepared by modifying the original recipe reported by Welshimer (24). Medium was always

prepared fresh, and filter sterilized with a 0.22micron filter. MMWB medium was supplemented

with either glutamine (Gln: 4.1mM) or ethanolamine (EA: 16mM) and VitB12 (200µM) as

needed. Cloning was done in either *E. coli* Top10 or DH5α. All *E. coli* strains were grown in

Luria-Bertani (LB) broth and were supplemented with antibiotics [Ampicillin (100 µg /ml) or

Kanamycin (50 µg/ml)] wherever required. For recombinant LMO strains, appropriate antibiotics

[Kanamycin (50 µg/ml) and streptomycin (50 µg/ml)] were used.

Generation of recombinant, *ΔeutV*, *ΔeutB* and the various complemented strains in LMO

All primers used in this study were purchased from Sigma-Aldrich and are listed in Table S2.

PCR reactions were performed using Platinum™ SuperFi II DNA Polymerase (ThermoFisher

Scientific: 12361010) according to the manufacturer's protocol. Restriction free cloning was

conducted using NEBuilder® HiFi DNA Assembly (NEB: E2621L). Isolation of plasmids and

purification of PCR products were conducted by using commercially available kits from either

Qiagen or Macherey-Nagel. Sequencing was conducted at Genewiz, and the results visualized

and verified using SnapGene (version 5.2.5.1). For creating the *ΔeutV* (LMAC030) and *ΔeutB*

(LMAC031)*,* the respective upstream and downstream regions were cloned into pMAD and

electroporated into LMO as reporter earlier (47). Electrocompetent cells of LMO were prepared

as described (48). The complemented strains *ΔeutV::eutV* (LMAC037) *and ΔeutB::eutB*

(LMAC034) were created by cloning the respective gene along with promoter region of either

eutV or *eutG* in pIMK plasmid followed by electroporation. For creating the recombinant strain

expressing mCherry tagged to EutK (LMAC038), the *eutK* coding sequence was driven by the

eutG promoter and fused to the mCherry coding sequence. It was then cloned into the pIMK

plasmid to create the construct AC038 followed by electroporation as described by Monk *et al*,

2008. Similarly, AC038 was electroporated into LMAC030 (*ΔeutV*) and LMAC031 (*ΔeutB*) to

generate strains LMAC040 (*eutK::mCherry*; *ΔeutV*) and LMAC043 (*eutK::mCherry*; *ΔeutB*),

respectively.

Growth curves

LMO and the mutants were grown overnight in 10 ml of BHI broth at 37°C with aeration.

Bacterial cells were harvested and washed thrice with PBS followed by a wash with MMWB

medium devoid of glutamine or EA. Cultures were then seeded on a 48 well plate from Falcon®

(Corning) containing the desired medium with a starting OD_{600} of 0.05. Plates were sealed with

Breathe-Easy® sealing membranes (Diversified Biotech) and the growth curve was monitored

on a Cytation™ 5 plate reader system (BioTek Instruments Inc) for 72 hours, with OD readings

taken every hour.

Transmission Electron Microscopy

Bacterial strains were grown in BHI broth to midlog phase at 37°C with aeration. Bacterial cells

were then harvested, washed, and processed as mentioned above. The strains were

resuspended in MMWB medium containing either Gln (4.1mM) or EA (16mM) and Vit B12

(200µM) as needed. Cells were then harvested, fixed, processed and sliced (12). Imaging was

conducted in JEOL 1400 TEM system, equipped with the Gatan 2k x 2k CCD camera, and

accompanying software.

Cell culture and infection

All the cell lines were purchased from Sigma. The murine macrophage RAW264.7 cell line and

Human colon adenocarcinoma Caco-2 cell line was cultured in DMEM (Gibco: 11995065)

supplemented with 10% fetal bovine serum (FBS) (Gibco: 10082147) at 37°C in a humidified

atmosphere at 5% CO_2. For culturing the Caco-2 cell line, DMEM medium was additionally

supplemented with non-essential amino acids (NEAA) (Gibco-11140050). The human

monocytic leukemia THP-1 cell line was thawed and cultured in RPMI media supplemented with

20% FBS and 2-mercaptoethanol (Gibco: 21985023) to a final concentration of 0.05 mM at

37°C in a humidified atmosphere at 5% CO_2. Once the culture was established, the FBS

concentration was reduced to 10% for further subculturing. PMA was used at a final

concentration of 1000ng/ml for 48 hours to differentiate the THP-1 cells. This was followed by

incubating the differentiated THP-1 cell in PMA free media for 24hr. Polarization of the

differentiated THP-1 macrophages to M1 phenotype was done by adding 10ng/ml LPS and

50ng/ml IFN-γ (32). For RAW264.7 and THP-1 macrophage intracellular survival assay, 5×10^5

cells were seeded into 12 well plates the day before the experiment. For Caco-2 cells, 5×10^5

cells were seeded into 12 well plates and allowed to grow for one week until tight junctions were

formed (49). LMO strains were grown overnight in BHI broth at 37°C. The next day the overnight

cultures were inoculated into fresh BHI to a starting OD_{600} of 0.05 and grown for 2 h at 37°C till

an OD_{600} of ~0.2. Cell lines were infected with a MOI of 5. The infection was conducted for 1

hour. The wells were then washed with Dulbecco's phosphate-buffered saline (PBS)

(prewarmed at 37°C), and extracellular bacteria were killed by incubating with DMEM containing

40 µg/ml gentamicin for 1 hour. Medium was changed after 1 hour and intracellular bacteria was

harvested by lysing the host cells with 1 ml of ice-cold PBS containing 0.1% Triton X-100.

Several serial dilutions were plated on BHI agar plates and kept at 37°C for the bacterial

colonies to appear. Colonies were counted manually as well as using the software openCFU

(version 3.9.0) (50).

cDNA preparation and qRT PCR

All RNA extraction was carried out using NucleoSpin RNA Plus kit (Macherey Nagel) according

to manufacturer's protocol. cDNA was synthesized from RNA using the RevertAid reverse

transcriptase for first strand cDNA synthesis (ThermoFisher Scientific: K1622). qRT-PCR was

performed using SYBR green based iTaq Universal SYBR Green Supermix (Biorad: 1725121),

using appropriate primers (Table S2) on a Biorad C1000 thermocycle equipped with CFX96

Optical Reaction Module. The relative expression of various genes was normalized to that of the

endogenous reference 16S rRNA gene and the fold change in expression was calculated using

the comparative Ct method. For host qRT-PCR, the normalization was done using ACTH as

endogenous reference. p-values were calculated using the Two-way ANOVA with Bonferroni's

multiple comparison test.

Confocal microscopy

To visualize the expression of EutK-mCherry, the recombinant LMO strains LMAC038

(*eutK::mCherry)*, LMAC040 ((*eutK::mCherry*; *ΔeutV*) and LMAC043 (*eutK::mCherry; ΔeutB*),

were initially grown overnight in BHI broth at 37°C. Bacterial cells were then washed with sterile

PBS and reconstituted in MMWB medium containing either Gln (4.1mM) or EA (16mM) and

VitB12 (200μM) as needed and incubated at 37°C on a shaker (180 rpm). Cells were harvested

by centrifugation (5000x g) at desired timepoints and were fixed with 4% paraformaldehyde for

10 minutes at 25°C. Cells were then washed thrice with PBS and smear dried on a Poly-L-lysine

coated slide (Electron Microscopy Science: 63410), mounted with Prolong Diamond Antifade

Mountant (Invitrogen: P36965), covered with #1.5H glass cover slips (Electron Microscopy

Sciences), and imaged using the Olympus FLUOVIEW FV3000 confocal microscope equipped

with the Fluoview FV315-SW software. Images were acquired in Z-stacks using a step size of

0.20 µm, which were then processed using the Olympus cellSens Dimension software.

Caco-2, RAW264.7 and THP-1 cells were thawed and seeded as mentioned earlier on 6 well

plates containing #1.5H glass cover slips (Electron Microscopy Sciences). Cells were grown and

maintained as described above. The infection was conducted for 1 hour at an MOI of 5, followed

by gentamycin treatment as described above. After infection, the cells were fixed with 4%

paraformaldehyde for 10 minutes at 25°C followed by permeabilization with 0.01% Triton X-100

in PBS for 10 minutes at 25°C. The cells were then treated with 5% BSA in PBS for 90 minutes

at 25°C followed by treatment with anti-Listeria antibody (Abcam: ab35132) at 25°C for 2 hours.

The next day, cells were then washed with PBS and treated with Alexa488 conjugated

secondary antibody (Invitrogen: A27034) for 90 minutes at 25°C. Cells were then washed and

stained for actin using Alexa Fluor plus 405 Phalloidin (Invitrogen: A30104) for 20 minutes at

25°C. Finally, cells were again washed with PBS, mounted, and imaged as described above.

Statistical Analysis

Statistical Analyses were done using GraphPad Prism, v9.5.1. P values were calculated using

two-way ANOVA with Bonferroni's multiple comparison test.

Data Availability

No large datasets were generated that required deposition into public repositories. All data is in

the manuscript with the source data underlying the Figures is in Table S3.

ACKNOWLEDGEMENTS

We thank S. Kolodziej and P. Navarro for sectioning samples for TEM and C. Wu and M.B. Willis for help with strain design and troubleshooting. A special thanks to A. Nagar for advice on culturing and imaging cell lines. Finally, we thank the following members of the LMO community for strains, protocols and advice: D. Portnoy, J.D. Sauer, S. Halbedel and M. O'Riordan. This work was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award number R21AI167124 to DAG. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Author contributions: A.C., K.K., and D.A.G. designed research; A.C and K.K performed experiments; A.C and D.A.G analyzed data and wrote the paper.

BIBLIOGRAPHY

1.   Randle CL, Albro PW, Dittmer JC. 1969. The phosphoglyceride composition of Gram-

negative bacteria and the changes in composition during growth. Biochim Biophys Acta

187:214–220.

2.   White DA. 1973. Phospholipid composition of mammalian tissues, p. 441–482. *In* Ansell,

GB, Hawthorne, JN, Dawson, RMC (eds.), Form and function of phospholipids. Elsevier

Publishing Company, New York.

3.   Bertin Y, Girardeau JP, Chaucheyras-Durand F, Lyan B, Pujos-Guillot E, Harel J, Martin

C. 2011. Enterohaemorrhagic Escherichia coli gains a competitive advantage by using

ethanolamine   as   a   nitrogen   source   in   the   bovine   intestinal   content.   Environ

Microbiol2010/09/21. 13:365–377.

4.   Thiennimitr P, Winter SE, Winter MG, Xavier MN, Tolstikov V, Huseby DL, Sterzenbach

T, Tsolis RM, Roth JR, Baumler AJ. 2011. Intestinal inflammation allows Salmonella to

use ethanolamine to compete with the microbiota. Proc Natl Acad Sci U S A2011/10/05.

108:17480–17485.

5.   Garsin DA. 2010. Ethanolamine utilization in bacterial pathogens: roles and regulation.

Nat Rev Microbiol 8:290–295.

6.   Kaval KG, Garsin DA. 2018. Ethanolamine Utilization in Bacteria. mBio 9.

7.   Toledo-Arana A, Dussurget O, Nikitas G, Sesto N, Guet-Revillet H, Balestrino D, Loh E,

Gripenland J, Tiensuu T, Vaitkevicius K, Barthelemy M, Vergassola M, Nahori MA,

Soubigou G, Regnault B, Coppee JY, Lecuit M, Johansson J, Cossart P. 2009. The

Listeria transcriptional landscape from saprophytism to virulence. Nature.

8.   Joseph B, Przybilla K, Stuhler C, Schauer K, Slaghuis J, Fuchs TM, Goebel W. 2006.

Identification of Listeria monocytogenes genes contributing to intracellular replication by

expression profiling and mutant screening. J Bacteriol 188:556–568.

9.   Pizarro-Cerda J, Cossart P. 2018. Listeria monocytogenes: cell biology of invasion and

intracellular growth. Microbiol Spectr2018/12/14. 6.

10.   Mellin JR, Koutero M, Dar D, Nahori MA, Sorek R, Cossart P. 2014. Riboswitches.

Sequestration of a two-component response regulator by a riboswitch-regulated

noncoding RNA. Science (1979) 345:940–943.

11.   DebRoy S, Gao P, Garsin DA, Harvey BR, Kos V, Nes IF, Solheim M. 2014.

Transcriptional and Post Transcriptional Control of Enterococcal Gene Regulation, p. . *In*

Gilmore, MS, Clewell, DB, Ike, Y, Shankar, N (eds.), Enterococci: From Commensals to

Leading Causes of Drug Resistant Infection. Boston.

12.   DebRoy S, Gebbie M, Ramesh A, Goodson JR, Cruz MR, van Hoof A, Winkler WC,

Garsin DA. 2014. Riboswitches. A riboswitch-containing sRNA controls gene expression

by sequestration of a response regulator. Science (1979) 345:937–940.

13.   Chen J, Gottesman S. 2014. RNA. Riboswitch regulates RNA. Science (1979) 345:876–

877.

14.   Anderson CJ, Clark DE, Adli M, Kendall MM. 2015. Ethanolamine Signaling Promotes

Salmonella Niche Recognition and Adaptation during Infection. PLoS Pathog

11:e1005278.

15.   Anderson CJ, Satkovich J, Koseoglu VK, Agaisse H, Kendall MM. 2018. The

Ethanolamine Permease EutH Promotes Vacuole Adaptation of Salmonella enterica and

Listeria monocytogenes during Macrophage Infection. Infect Immun2018/03/14. 86.

16.   Penrod JT, Mace CC, Roth JR. 2004. A pH-sensitive function and phenotype: evidence

539   that EutH facilitates diffusion of uncharged ethanolamine in Salmonella enterica. J

540   Bacteriol 186:6885–6890.

541   17.   Bobik TA, Lehman BP, Yeates TO. 2015. Bacterial microcompartments: widespread

542   prokaryotic organelles for isolation and optimization of metabolic pathways. Mol Microbiol

543   98:193–207.

544   18.   Kerfeld CA, Aussignargues C, Zarzycki J, Cai F, Sutter M. 2018. Bacterial

545   microcompartments. Nat Rev Microbiol2018/03/06. 16:277–290.

546   19.   Zeng Z, Boeren S, Bhandula V, Light SH, Smid EJ, Notebaart RA, Abee T. 2021.

547   Bacterial Microcompartments Coupled with Extracellular Electron Transfer Drive the

548   Anaerobic Utilization of Ethanolamine in Listeria monocytogenes. mSystems 6.

549   20.   Quin MB, Perdue SA, Hsu SY, Schmidt-Dannert C. 2016. Encapsulation of multiple cargo

550   proteins   within   recombinant   Eut   nanocompartments.   Appl   Microbiol

551   Biotechnol2016/07/28. 100:9187–9200.

552   21.   Choudhary S, Quin MB, Sanders MA, Johnson ET, Schmidt-Dannert C. 2012.

553   Engineered protein nano-compartments for targeted enzyme localization. PLoS One

554   7:e33342.

555   22.   Held M, Kolb A, Perdue S, Hsu SY, Bloch SE, Quin MB, Schmidt-Dannert C. 2016.

556   Engineering formation of multiple recombinant Eut protein nanocompartments in E. coli.

557   Sci Rep2016/04/12. 6:24359.

558   23.   Kutzner E, Kern T, Felsl A, Eisenreich W, Fuchs TM. 2016. Isotopologue profiling of the

559   listerial N-metabolism. Mol Microbiol 100.

560   24.   WELSHIMER HJ. 1963. VITAMIN REQUIREMENTS OF LISTERIA MONOCYTOGENES.

561   J Bacteriol 85.

562   25.   Premaratne RJ, Lin WJ, Johnson EA. 1991. Development of an improved chemically

563   defined minimal medium for Listeria monocytogenes. Appl Environ Microbiol 57.

564   26.   Tsai HN, Hodgson DA. 2003. Development of a Synthetic Minimal Medium for Listeria

565   monocytogenes. Appl Environ Microbiol 69.

566   27.   FRIEDMAN ME, KAUTTER DA. 1962. Effect of nutrition on the respiratory virulence of

567   Listeria monocytogenes. J Bacteriol 83.

568   28.   Savage DF, Afonso B, Chen AH, Silver PA. 2010. Spatially ordered dynamics of the

569   bacterial carbon fixation machinery. Science (1979) 327:1258–1261.

570   29.   Cameron JC, Wilson SC, Bernstein SL, Kerfeld CA. 2013. Biogenesis of a bacterial

571   organelle: the carboxysome assembly pathway. Cell 155:1131–1140.

572   30.   Gaillard JL, Berche P, Mounier J, Richard S, Sansonetti P. 1987. In vitro model of

573   penetration and intracellular growth of Listeria monocytogenes in the human enterocyte-

574   like cell line Caco-2. Infect Immun1987/11/01. 55:2822–2829.

575   31.   Mounier J, Ryter A, Coquis-Rondon M, Sansonetti PJ. 1990. Intracellular and cell-to-cell

576   spread of Listeria monocytogenes involves interaction with F-actin in the enterocytelike

577   cell line Caco-2. Infect Immun1990/04/01. 58:1048–1058.

578   32.   Ouadrhiri Y, Scorneaux B, Sibille Y, Tulkens PM. 1999. Mechanism of the intracellular

579   killing and modulation of antibiotic susceptibility of Listeria monocytogenes in THP-1

580   macrophages activated by gamma interferon. Antimicrob Agents Chemother 43.

581   33.   Borish LC, Steinke JW. 2003. Borish, L. C., & Steinke, J. W. (2003). 2. Cytokines and

582   chemokines. The Journal of Allergy and Clinical Immunology, 111(2 Suppl), S460–S475.

583   http://doi.org/10.1067/mai.2003.1082. Cytokines and chemokines. J Allergy Clin Immunol

584   111.

585   34.   Kany S, Vollrath JT, Relja B. 2019. Cytokines in inflammatory disease. Int J Mol Sci

586   https://doi.org/10.3390/ijms20236008.

35.   Xie M, Ding C, Guo L, Chen G, Zeng H, Liu Q. 2018. Evaluation of Caco-2 cells response
to Listeria monocytogenes virulence factors by RT-PCR. Microb Pathog 120.

36.   Kendall MM, Gruber CC, Parker CT, Sperandio V. 2012. Ethanolamine controls
expression of genes encoding components involved in interkingdom signaling and
virulence in enterohemorrhagic Escherichia coli O157:H7. mBio 3.

37.   Garsin DA. 2012. Ethanolamine: a signal to commence a host-associated lifestyle?
mBio2012/07/05. 3:e00172–12.

38.   Ramesh A, DebRoy S, Goodson JR, Fox KA, Faz H, Garsin DA, Winkler WC. 2012. The
mechanism for RNA recognition by ANTAR regulators of gene expression. PLoS
Genet2012/06/12. 8:e1002666.

39.   Tsoy O, Ravcheev D, Mushegian A. 2009. Comparative genomics of ethanolamine
utilization. J Bacteriol.

40.   Villa EA, Escalante-Semerena JC. 2022. Acinetobacter baumannii Catabolizes
Ethanolamine in the Absence of a Metabolosome and Converts Cobinamide into
Adenosylated Cobamides. mBio 13.

41.   Penrod JT, Roth JR. 2006. Conserving a volatile metabolite: a role for carboxysome-like
organelles in Salmonella enterica. J Bacteriol 188:2865–2874.

42.   Brinsmade SR, Paldon T, Escalante-Semerena JC. 2005. Minimal functions and
physiological conditions required for growth of salmonella enterica on ethanolamine in the
absence of the metabolosome. J Bacteriol 187:8039–8046.

43.   Oltrogge LM, Chaijarasphong T, Chen AW, Bolin ER, Marqusee S, Savage DF. 2020.
Multivalent interactions between CsoS2 and Rubisco mediate α-carboxysome formation.
Nat Struct Mol Biol https://doi.org/10.1038/s41594-020-0387-7.

44.   Yang M, Wenner N, Dykes GF, Li Y, Zhu X, Sun Y, Huang F, Hinton JCD, Liu LN. 2022.
Biogenesis of a bacterial metabolosome for propanediol utilization. Nat Commun 13.

45.   Kennedy NW, Mills CE, Abrahamson CH, Archer AG, Shirman S, Jewett MC, Mangan

NM, Tullman-Ercek D. 2022. Linking the Salmonella enterica 1,2-Propanediol Utilization

Bacterial Microcompartment Shell to the Enzymatic Core via the Shell Protein PduB. J

Bacteriol 204.

46.   Kaval KG, Singh K V, Cruz MR, DebRoy S, Winkler WC, Murray BE, Garsin DA. 2018.

Loss of Ethanolamine Utilization in Enterococcus faecalis Increases Gastrointestinal

Tract Colonization. mBio2018/05/10. 9.

47.   Arnaud M, Chastanet A, Debarbouille M. 2004. New vector for efficient allelic

replacement in naturally nontransformable, low-GC-content, gram-positive bacteria. Appl

Environ Microbiol2004/11/06. 70:6887–6891.

48.   Monk IR, Gahan CG, Hill C. 2008. Tools for functional postgenomic analysis of listeria

monocytogenes. Appl Environ Microbiol2008/04/29. 74:3921–3934.

49.   Natoli M, Leoni BD, D'Agnano I, Zucco F, Felsani A. 2012. Good Caco-2 Cell Culture

Practices. Toxicology in Vitro 26.

50.   Geissmann Q. 2013. OpenCFU, a New Free and Open-Source Software to Count Cell

Colonies and Other Circular Objects. PLoS One 8.

 FIGURE LEGENDS

Fig. 1: MMWB plus EA supports LMO10403s growth dependent on *eutV* and *eutB*. (A)

Graphical representation of the organization and regulation of the *eut* operon in LMO. See text

for details. (B) Growth curve of LMO when grown in MMWB containing either glutamine (Gln) or

ethanolamine (EA) as a source of nitrogen. (C) Relative expression of the *eut* genes when LMO

bacteria were grown in the presence of EA with respect to Gln. (D) Growth curve comparison of

ΔeutV and ΔeutV complemented *(ΔeutV::eutVW)* when grown in the presence of Gln or EA. (E)

Growth curve comparison of *ΔeutB and ΔeutB* complemented *(ΔeutB::eutB)* when grown in the

presence of Gln or EA. Shown is the average of three replicates and error bars represent the

standard deviation (SD).

Fig 2: EUT BMCs form in medium containing EA. TEM images showing the formation of EUT

BMCs when LMO was grown in MMWB with (A) Gln or EA. No such structures were visible in

the *ΔeutV* (B) and *ΔeutB* (C) strains grown under the same conditions. Complementation of

ΔeutV and *ΔeutB* restored the formation of BMC structures (B and C right panel respectively).

Scale bar represents 500nm and yellow asterisks indicate the BMCs.

Fig 3: EUT BMCs can be fluorescently tagged. Confocal images of the LMO *eutK::mCherry*

strain when grown in in MMWB with either EA (A) or Gln (B) as a nitrogen source. (B) Confocal

images of LMO strains containing the *eutK::mCherry* transgene (red) in *ΔeutV* (C) and *ΔeutB*

(D) backgrounds following growth in MMWB with EA. Scale bar represents 10μm. (E)

Percentage of cells with visible BMCs. Shown is the average from replicates containing an n of

20-60 cells and the error bars represent the SD.

Fig. 4: *eut* mutants are deficient in intracellular replication: Bacterial CFUs were quantified

in Caco-2 cells (A), RAW264.7 macrophages (B) and M1 polarized THP-1 macrophages (C). At

the indicated time points post-infection, the cells were washed, lysed, and serial dilutions of the

lysates were plated as described in methods. The mean of three replicates were calculated with

the error bars indicating the standard deviation. 2-way ANOVA with Bonferroni multiple testing

correction was used to calculate the significance of the indicated comparisons. *$P < 0.05$; **$P <$

0.01; ***$P < 0.001$; ****$P < 0.0001$.

Fig. 5: LMO forms visible BMCs in Caco-2 cells dependent on *eutV* and *eutB*. Caco-2 cells

were infected with LMO strains containing the *eutK::mCherry* transgene (red) in the wild type

(A), *ΔeutV* (B) and *ΔeutB* (C) backgrounds. The cells were fixed at the indicated time points and

stained with phalloidin to visualize actin (blue) and α-LMO antibody to visualize the bacteria

(green). Representative confocal images are shown. Scale bars represents 1μm. (D) Close-up

of boxed view of LMO containing *eutK::mCherry* at the 6 hour time point from Fig. S3. Red

arrows indicate BMCs. Yellow arrows indicate examples of bacteria surrounded by an actin

cloud. White arrows point to examples of actin tails. Five bacteria associated with an actin tail

have been outlined in white for easier visualization.

Fig 6: *eut* mutants increase pro-inflammatory signatures during Caco-2 infection.

Following infection of Caco-2 cells with the indicated strains, host cell RNA was isolated and

qRT-PCR analysis revealed a significant elevation in the expression of genes encoding

inflammatory cytokines interleukin-1 beta (IL-1β), tumor necrosis factor-alpha (TNF-α),

interleukin-6 (IL-6), interleukin-8 (IL-8), and chemokines CXCL-2 and MCP-1. The mean of three

replicates were calculated with the error bars indicating the standard deviation. 2-way ANOVA

with Bonferroni multiple testing correction was used to calculate the significance of the indicated

comparisons. $*P < 0.05$; $**P < 0.01$; $***P < 0.001$; $****P < 0.0001$.

(A)

WT + Gln   WT + EA

(B)

ΔeutV + EA   *ΔeutV::eutV* + EA

(C)

ΔeutB + EA   *ΔeutB::eutB* + EA

(A) *eutK::mCherry*

	BMC	Bacteria	BMC Bacteria Actin
2h			
4h			
6h			
8h			

(B) *eutK::mCherry; ΔeutV*

	BMC	Bacteria	BMC Bacteria Actin
2h			
4h			
6h			
8h			

(C) *eutK::mCherry; ΔeutB*

	BMC	Bacteria	BMC Bacteria Actin
2h			
4h			
6h			
8h			

(D)