Unraveling the Role of P450 Reductase in Herbicide Metabolic Resistance Mechanism

Running title: **Role of P450 Reductase in Herbicide Metabolic Resistance Mechanism**

Carlos Alberto Gonsiorkiewicz Rigon[1], Satoshi Iwakami[2], Todd A. Gaines[1], Franck E. Dayan[1]

[1] Colorado State University, Department of Agricultural Biology, Fort Collins, CO 80523.

Carlos.Rigon@colostate.edu; Todd.Gaines@colostate.edu; Franck.Dayan@colostate.edu

[2] Kyoto University, Graduate School of Agriculture, Kyoto 606-8502, Japan

iwakami.satoshi.2v@kyoto-u.ac.jp

Corresponding Author: Franck Dayan, franck.dayan@colostate.edu

HIGHLIGHTS

Knocking out cytochrome P450 reductase 1 in herbicide tolerant Arabidopsis reduces herbicide

resistance, highlighting CPRs as targets for managing herbicide resistance evolution in weeds.

ABSTRACT

Plants require cytochrome P450 reductase (CPR) to supply two electrons for cytochrome P450

monooxygenase enzymes (P450) to react with an organic substrate. The transfer of electrons to

the P450 active site in the P450 catalytic site relies on a robust and intricate CPR:P450

complex in the endoplasmic reticulum membrane. Transgenic Arabidopsis plants carrying

CYP81A12 from *Echinochloa phyllopogon*, which metabolizes a broad spectrum of herbicides,

were crossed with CPR knockout *atr1* or *atr2* mutant lines. Homozygous gene knockout was

confirmed using PCR, and gene copy number of *CYP81A12* was determined using ddPCR.

Arabidopsis lines expressing *CYP81A12* in combination with *atr1* or *atr2* knockout were used

for herbicide dose-response and metabolism studies. Knocking out *ATR1* in transgenic

Arabidopsis *CYP81A12* significantly reduced herbicide resistance. Transgenic mutant plants

(*CYP81A12 atr1-b*) had a 3.6-, 5.6-, 6.8- and at least 26- fold reduction in resistance to

mesotrione, 2,4-D, penoxsulam and chlorsulfuron, respectively, in the dose-response assay.

Knockouts of the *ATR2* also decreased herbicide resistance, but to a lower magnitude than

ATR1. These results corroborate ½ MS medium assay, and herbicide resistance reduction was

observed for additional tested herbicides, bensulfuron, propoxycarbazone and bentazon. Our

findings highlight the importance of CPRs in metabolic herbicide resistance in plants, by

identifying that a single CPR knockout can reverse herbicide sensitivity . The different CPRs

found in weeds have potential as target genes to manage metabolic herbicide resistance

evolution. We further provide an in-depth exploration of the evolutionary implications in weed

management arising from the results.

Key Words: CPR, CPR:P450 complex, electron, metabolism, monooxygenase, evolution

INTRODUCTION

Plants are sessile organisms, which remain permanently anchored to their site of germination. To overcome their lack of mobility, plants have evolved complex detoxification mechanisms to deal with environmental and biotic stresses. These natural detoxification processes by chance enable enhanced herbicide metabolic pathways in certain crops, conferring natural tolerance to herbicides (Jeschke *et al.*, 2019). This strategic approach has yielded substantial advantages in the commercialization of herbicides intended to eliminate weeds while safeguarding crops; however, the prolonged application of herbicides in agroecosystems has selected weeds that exhibit the same mechanisms of resistance (Powles & Yu, 2010) . Among these, the enhanced capacity of herbicide metabolism is one of the most important (Rigon *et al.*, 2020). These selected weeds have an enhanced activity of enzyme families that metabolize the herbicides before reaching the target site (Gaines *et al.*, 2020). This mechanism of resistance threatens weed management due to its broad substrate specificity to many herbicides from different mechanisms of action (Rigon *et al.*, 2020). Metabolic herbicide resistance mechanism is divided in three phases, known as Phase I – oxidation, Phase II – conjugation, and Phase III – sequestration and degradation. Phase I is mediated primarily by cytochrome P450 monooxygenase (P450) enzymes. These enzymes are membrane-bound, localized in the endoplasmic reticulum, and are involved in the biosynthesis of many molecules including antioxidants, secondary metabolites, and in detoxification of many xenobiotic compounds (Pandian *et al.*, 2020). P450s are the most important enzymes in phase I of herbicide metabolism and plants possess a high number of P450 genes that might play a crucial role in herbicide resistance in metabolic resistant weed species (Dimaano & Iwakami, 2021).

Cytochrome P450 reductase (CPR) is a membrane-bound enzyme localized in the endoplasmic reticulum that plays a critical role in the biosynthesis of a wide range of compounds in living organisms, including plants, animals, and bacteria. The structure of P450 reductase is divided in four domains, including *N*-terminal flavin mononucleotide (FMN)-binding domain, a connecting domain, an flavin adenine dinucleotide (FAD)-binding domain, and a *C*-terminal NADPH-binding domain (Wang *et al.*, 1997, Jensen & Møller, 2010) (Fig. **1**). CPR enzymes arose from of a fusion of different genes encoding FAD-binding, ferredoxin-NADP$^+$, and an FMN-flavodoxin (Porter & Kasper, 1986). Plants require CPR to supply two electrons for P450 enzymes to conduct each monooxygenase reaction with an organic substrate (Lau & O'Keefe,

1996, Paine *et al.*, 2005). The CPR reduction includes multiple steps for electron transfer. The

first step is reduction from NADPH to FAD, then FMN, and finally to the P450 acceptor (Munro

et al., 2001) (Fig. **1**). The transfer of electrons to the P450 active site in the P450 catalytic cycle

relies on a robust and intricate CPR:P450 complex in the membrane (Mukherjee *et al.*, 2021),

making CPR function an essential component of specialized metabolism.

In contrast to the huge number of P450 genes within plant genomes, for example, 246 in

Arabidopsis thaliana (Arabidopsis) (Paquette *et al.*, 2009), 328 in *Oryza* spp. (rice) (Nelson *et*

al., 2004), 867 in *Echinochloa crus-galli* (barnyardgrass) (Wu *et al.*, 2022), 694 in *E. colona*

(jungle rice) (Wu *et al.*, 2022), and 323 in *Conyza canadensis* (horseweed) (Peng *et al.*, 2014),

CPRs generally have only two or three copies in plant genomes. There are two CPR genes in

Arabidopsis*,* referred to as *Arabidopsis thaliana* cytochrome P450 reductase 1 (*ATR1*), 2 (*ATR2*)

(Urban *et al.*, 1997) and one hypothetical reductase 3 (*ATR3)* that encodes a diflavin reductase

essential for embryogenesis, but that is unable to reduce P450 *in vitro* (Mizutani & Ohta, 1998,

Varadarajan *et al.*, 2010). The isoforms are grouped in two distinct groups with different roles.

Class I is *ATR1*, which is involved in primary and basal metabolism. Class II is *ATR2*, which is

inducible by environmental stimuli such as wounding, pathogen attack, or high light intensity

exposure (Rana *et al.*, 2013, Parage *et al.*, 2016).

In current research efforts, there has been a growing focus on the identification of P450

enzymes responsible for herbicide metabolism in weeds (Rigon *et al.*, 2020). An important

example is the CYP81A12 enzyme derived from a multiple-resistant population of *Echinochloa*

phyllopogon. This enzyme is capable of metabolizing herbicides from diverse modes of action

when expressed heterologously in *E. coli* or Arabidopsis (Iwakami *et al.*, 2014, Dimaano *et al.*,

2020). Considering the dependence of P450 enzymes on P450 reductase for their functionality,

the validation of this interdependence and the identification of pivotal cytochrome P450

reductases could potentially offer a valuable mechanism for weed management and the

mitigation of metabolic herbicide resistance evolution in weeds.

The study hypothesized that *ATR1* and *ATR2* are the primary electron donors for P450

enzymes and that silencing these genes would reduce the activity of P450s involved in herbicide

metabolism. The main objective was to assess the potential of targeting cytochrome P450

reductase to diminish herbicide metabolism mediated by *CYP81A12*.

MATERIALS AND METHODS

Plant material

Mutant SALK lines of Arabidopsis were ordered from the Arabidopsis Biological Resource Center (ABRC - https://abrc.osu.edu/). The lines SALK_111558, SALK_208483C have a T-DNA insertion in Arabidopsis cytochrome P450 reductase 1 (*ATR1* - at4g24520), hereafter named as *atr1-a* and *atr1-b*, respectively. The lines SALK_026053 and SALK_152766 have T-DNA insertion in Arabidopsis cytochrome P450 reductase 2 (*ATR2* - at4g30210), hereafter named as *atr2-a* and *atr2-b*.

Seeds from mutants and wild-type Columbia_0 (Col_0) were sown in soil and kept in a growth chamber at 22/18°C with a light intensity of 120 µE m^{-2} s^{-1} for 16 h. When seedlings had four true leaves, they were transplanted into individual pots and kept in the growth chamber in the same conditions. All the following experiments used the same methods to grow the Arabidopsis plants.

Characterization and confirmation of T-DNA insertion

PCR assays were performed to identify and confirm the presence of T-DNA insertion in the *ATR1* and *ATR2* genes in the mutant lines and to verify homozygosity. The methods are described in Methods **S1**.

Generating transgenic herbicide-resistant Arabidopsis with mutant *atr1* or *atr2* genes and double knockout lines

Aiming to generate transgenic Arabidopsis lines with a loss of function in *ATR1* or *ATR2*, a transgenic Arabidopsis line carrying the P450 *CYP81A12* from *Echinochloa phyllopogon* (synonym *E. oryzicola*, late watergrass) (Iwakami *et al.*, 2014) was used. The transgenic Arabidopsis metabolizes five different herbicide modes of action including acetolactate synthase (ALS) inhibitors (bensulfuron-methyl, pyrazosulfuron-ethyl, chlorsulfuron, azimsulfuron, imazamox, propoxycarbazone-sodium and penoxsulam), 4-hydroxyphenylpyruvate dioxygenase (HPPD) inhibitor (mesotrione), protoporphyrinogen oxidase (PPO) inhibitor (pyraclonil), deoxyxylulose phosphate synthase (DXS) inhibitor (clomazone), and photosystem II (PSII) inhibitor (bentazon) (Dimaano *et al.*, 2020). This population hereafter is called Arabidopsis

CYP81A12. This line was crossed with mutant lines *atr1-a*, *atr1-b*, *atr2-a*, and *atr2-b* to generate

transgenic CYP81A12 carrying mutant *atr1* or *atr2*, hereafter are named as *CYP81A12 atr1-a*,

CYP81A12 atr1-b, *CYP81A12 atr2-a*, and *CYP81A12 atr2-b*. The methods of crossing and

genotyping populations are described in Methods **S2**. Aiming to generate double knockout lines,

i.e. same line with loss of function in *atr1* and *atr2*, mutants plants were crossed and the methods

are described in Methods **S3**.

Gene expression analysis in *atr1* and *atr2* mutant lines

Mutant and transgenic mutant lines were grown in a growth chamber with the same

conditions as specified before. Approximately 100 mg of leaf tissue were collected from plants

in the rosette stage. Sampling consisted of cutting the youngest fully expanded leaf for each

biological replication, placing in 2 mL tubes, and incubating immediately in liquid nitrogen and

afterwards kept at -80°C. Plants of lines Col_0, transgenic *CYP81A12*, and transgenic mutants

CYP81A12 atr1-a, *CYP81A12 atr1-b*, *CYP81A12 atr2-a*, and *CYP81A12 atr2-b* were kept for

fresh mass measurement at 28 d after transplant to analyze any fitness cost in the mutant lines.

RNA extraction was conducted using Direct-zol™ RNA miniprep from Zymo Research

following manufacturer instructions. DNase I treatment was conducted during the RNA

extraction protocol. Quantification and quality control were performed using Nanodrop 2000.

cDNA synthesis was conducted using iScript™ from Zymo Research using 1 µg of RNA. Gene

expression was analyzed on CFX96 Real Time System using SsoAdvanced™ Universal SYBR®

Green Supermix. The reactions consisted of 5 µL of cDNA from samples diluted in Nuclease-

Free Water in 1:20, 2.5 µL of each primer at 5 µM, and 5 µL of SsoAdvanced universal SYBR®

Green supermix (2×). Thermal cycler conditions consisted of an initial cycle of 30 s at 95°C,

followed by a sequence of 35 cycles starting at 95°C for 10 s and 72°C for 30 s. Melt-curve

analysis was performed by adding steps incrementing temperature 65–95°C by 0.5°C.

Primer sets were specifically designed to target various regions of the genes, both upstream

and downstream of the T-DNA insertion site. This systematic design allowed for accurate

analysis of how the gene was affected by the T-DNA insertion. For *ATR1* gene, four sets of

primers were designed, called ATR1_A with forward primers in the 5'UTR and reverse primers

in the 1st exon (spanning the T-DNA insertion position in *atr1-a* line), ATR1_B with forward

and reverse primers in the 1st exon, ATR1_C with the forward primer in exon 4th and reverse

primer in exon 5th (spanning the T-DNA insertion position in *atr1-b* line), and ATR1_D with

forward in the 8[th] exon and reverse in the 9[th] exon (Fig. **3**). For *ATR2* gene, five sets of primers

were designed, ATR2_A with forward and reverse primers localized in the 1[st] exon, ATR2_B

with forward primer in the 3[th] exon and reverse primer in the 4[th] exon (spanning the T-DNA

insertion position in *atr2-a* line), ATR2_C with forward primer localized in the 9[th] and reverse in

the 10[th] exon, ATR2_D with forward primer in the 12[th] exon and reverse in the 13[th] (spanning the

T-DNA insertion position in *atr2-b* line), and ATR2_E with forward and reverse primers

designed in the 14[th] exon. The primer locations in *ATR1* and *ATR2* are represented in Fig. **3a** and

b, respectively. Primer sequences for gene expression analysis and their characteristics are listed

in Tab. **S2**.

Herbicide sensitivity of Arabidopsis lines in ½ MS medium

Plant growth of different lines of Arabidopsis were evaluated on ½ Murashige and Skoog

(MS) solid medium containing different doses of bensulfuron, chlorsulfuron, penoxsulam,

propoxycarbazone, mesotrione and bentazon. Seeds of Arabidopsis wild type (Col_0), transgenic

Arabidopsis *CYP81A12*, mutants and transgenic mutant *CYP81A12 atr1-a*, *CYP81A12 atr1-b*,

CYP81A12 atr2-a, and *CYP81A12 atr2-b* were sterilized by chloride gas for 7 h by mixing 100

mL of bleach with 3% hydrochloric acid. Technical grade herbicides were diluted in ethanol

(bensulfuron, penoxsulam, propoxycarbazone, bentazon) or methanol (chlorsulfuron,

mesotrione) to mix with ½ MS medium before it solidified. Petri dishes containing ½ MS with

different herbicides doses were divided into 10 sections, where six seeds of each Arabidopsis

line were placed. The plates were incubated in a growth chamber at 25°C, 14-h photoperiod, and

110 μE m^{-2} s^{-1} light intensity until the day of evaluation. The number of green leaves were

counted 14 d after plating, and the data were normalized to percentage of untreated control.

Whole-plant dose response

Arabidopsis plants of Col_0, transgenic *CYP81A12*, and transgenic mutant *CYP81A12*

atr1-a, *CYP81A12 atr1-b*, *CYP81A12 atr2-a*, and *CYP81A12 atr2-b* were grown with the same

conditions as specified before. Plants in rosette stage (14 d after transplanting) were sprayed with

increasing doses of chlorsulfuron (Telar ® XP, 750 g Kg^{-1}), penoxsulam (Ricer, 217 g a.i. kg^{-1}),

mesotrione (Callisto 480g a.i. L^{-1}), and 2,4-D (Shredder 2,4-D LV4, 455g a.e. L^{-1}). The

herbicides were applied in an automated spray chamber (Greenhouse Spray Chamber, model

Generation IV) using a TJ8002E nozzle, calibrated to deliver 200 L ha^{-1} at a pressure of 280 kPa

and speed of 1.2 m s^{-1}. The experimental design was completely randomized in 6 $\times$ 11 with four

replications. Factor A was Arabidopsis lines. Factor B was increasing herbicides dose for each

herbicide. Chlorsulfuron and penoxsulam doses were in range from 0 to 51.2 g a.i. ha^{-1};

mesotrione doses were from 0 to 400 g a.i. ha^{-1}; and 2,4-D doses were from 0 to 128 g a.e. ha^{-1}.

The shoot fresh mass was measured at 28 d after herbicide treatment.

Herbicide metabolism analysis

Seedlings of Arabidopsis lines Col_0, transgenic *CYP81A12*, and transgenic mutant

CYP81A12 atr-a, *CYP81A12 atr-b*, *CYP81A12 atr2-a*, and *CYP81A12 atr2-b* were transplanted

into pots filled with potting soil and grown under the same conditions as previously described. At

rosette stage, mesotrione or chlorsulfuron were applied were applied at 100 and 50 g ha^{-1},

respectively. The doses were chosen based on the whole-plant dose-response curve experiment

described above. After herbicide application the plants were kept in the growth chamber at same

conditions and whole plants were collected at 6, 12, 24, 48, and 96 h for mesotrione and adding

144 h and 196 h for plants treated with chlorsulfuron. At each time of tissue collection, plant

samples were weighed, washed with 80% acetone three times, dried with towel paper, placed in

15 mL Falcon tubes, and kept at -80°C ultra-freezer until the extraction. The extraction was

conducted by grinding the tissue with a pestle, then added 10X of plant mass with 80% methanol

and 0.1% formic acid. The solution was homogenized for 30 s using PowerGen 125 (Fisher

Scientific) and centrifuged for 10 min at 4,700$\times$g (Sorvall Legend X1R, Thermo Scientific,

Waltham, MA). The supernatant was filtered with Econofltr Nylon 13 mm 0.2 µm (Agilent

Technologies, Santa Clara, CA) and injected in the LC-MS/MS.

LC-MS/MS system consisted of a Nexera X2 UPLC with 2 LC-30AD pumps, a SIL-30AC

MP autosampler, a DGU-20A5 Prominence degasser, a CTO-30A column oven, and SPD-M30A

diode array detector coupled to an 8040 quadrupole mass-spectrometer (Shimadzu Scientific

Instruments, Columbia, MD). For mesotrione, the MS was in positive mode [M+H]$^{+}$ with MRM

optimized for mesotrione 340>227.95 and set for 100 ms dwell time with a Q1 pre-bias of -

16.0V, a collision energy of -18.0V and a Q3 pre-bias of -16.0V and for hydroxy-mesotrione

356>55>228 and set for 100 ms dwell time with a Q1 pre-bias of -16V, a collision energy of -

18.0V and a Q3 pre-bias of -16.0V. For chlorsulfuron, the MS was in positive mode with MRM

optimized for chlorsulfuron 357.9>141.1>167.05 and set for 100 ms dwell time with a Q1 pre-

bias of -25.0V, a collision energy of -20.0V and a Q3 pre-bias of -15.0V and for hydroxy-

chlorsulfuron 374.0>127.0>182.0 and set for a collision energy of -35.0. The samples were

chromatographed on a 100×4.6 mm Kinetex 2.6 µm biphenyl column (Phenomenex, Torrance,

CA) maintained at 40°C. For both herbicides, solvent A consisted of water with 0.1% formic

acid and solvent B was methanol with 0.1% formic acid. The solvent program started at 50% B

and increased to 95% B in 5 min and maintained at 95% for 4 min. The solvent was returned to

50% B and maintained there for 3 min before the next injection. The flow rate was set at 0.4 mL

min^{-1} and each sample were analyzed with 5 µL injection volumes.

Identification and phylogenetic analysis of CPRs in different weeds

The presence of CPR genes and copy number was searched in several important weed

species. The genomes of *Alopecurus myosuroides* (Cai *et al*., 2023), *Chenopodium formosanum*

(Jarvis *et al., 2022)*, *Bassia scoparia* (Hall *et al.*, 2023), *Eleusine indica* (Zhang *et al*., 2023),

Bromus tectorum (Revolinski *et al.,* 2023), *Echinochloa crus-galli* (Wu *et al*., 2022), *E.*

oryzicola (Wu *et al.,* 2022), *E. colona* (Wu *et al.*, 2022), *Ipomoea purpurea* (Gupta *et al.*, 2023) ,

Poa annua (Robbins *et al.*, 2023), and *Conyza canadensis* (Laforest *et al.*, 2020) were accessible

through WeedPedia (https://weedpedia.weedgenomics.org) (Montgomery *et al.*, 2023) . CPRs

were identified with a blast search using *ATR1* (AT4G24520) or *ATR2* (AT4G30210) against the

genomes or utilized the InterPro ID for NADPH-cytochrome P450 reductase (IPR023208).

Additionally, the genomes of *Amaranthus palmeri* and *A. tuberculatus* were available

(Montgomery *et al.*, 2020), and the COGE web browser (https://genomevolution.org/coge) was

used for blasting purposes. The *CPR* genes in *Amaranthus* species were identified using a

comparative analysis approach. Firstly, *ATR1* and *ATR2* sequences were subjected to a sequence

similarity search against the *Amaranthus hypochondriacus* (grain amaranth) genome (Lightfoot

et al., 2017), which is accessible through the Phytozome database. Subsequently, the identified

sequences were further compared against the Palmer amaranth and common waterhemp

genomes. Other already known plant *CPR* genes were extracted from GeneBank NCBI (Benson

et al., 2013), such as corn (*Zea mays*, CPR2a: NP_001159331.1, CPR2b: NP_001146345.2), rice

(*Oryza sativa*, CPR2a: XP_015651232.1, CPR2b: XP_015650780.1), soybean (*Glycine max*,

CPR1: NP_001236742.2, CPR2: XP_003549436.1), cotton (*Gossypium hirsutum*, CPR1:

ACN54323.1, CPR2: NP_001314398.2), and Barrel medic (*Medicago truncatula*, CPR1:

XP_003602898.1, CPR2: XP_003610109.1). Multiple protein alignment was performed using

ClustalO. Tree construction with Neighbor-joining method was performed using Geneious

Prime® 2023.0.1. The *N*-terminal, transmembrane, FMN binding, FAD binding, and NADP$^+$

binding domains were identified. The transmembrane was identified using DeepTMHMM tool

(Hallgren *et al.*, 2022) (https://dtu.biolib.com/DeepTMHMM).

Statistical analysis

Gene expression analysis of *ATR1* and *ATR2* genes in mutant plants was performed using

ALS as a normalization gene (Chen *et al.*, 2017). Expression analyses were performed utilizing

four biological and two technical replicates for each line. The mean Ct values and the standard

deviation were calculated by treatment. A melt curve analysis confirmed the presence of a single

amplified product for each reaction. Relative transcript abundance was calculated using $2^{(-\Delta\Delta Ct)}$

method for each gene and biotype (Livak & Schmittgen, 2001). The reference population used

for the calculation was Col_0. Dunnett's test ($p < 0.05$) was used to compare the relative

expression between Col_0 and mutant lines.

Number of green leaves at 14 d after seeds were planted and shoot fresh weight (% of

untreated control) at 28 d after herbicide application (DAA) were used as response variable for

the ½ MS medium and whole-plant dose-response experiment, respectively. Data were converted

into percentages relative to untreated treatment. The drc package (Ritz *et al.*, 2015) from

statistical software R v.3.5.3 (R Core Team, 2023) was used for data analysis. The data were

tested for the best nonlinear model by comparing different models by *ANOVA()* and using

function *modelFit()* testing for lack-of-fit. Based on the analysis, the data were adjusted using the

three-parameter log-logistic model with the function *modelFit()*. For the whole-plant dose

response experiment, the herbicide doses causing 50% reduction in shoot fresh weight (GR$_{50}$)

were determined using the function *summary()* and the resistance index (RI) was calculated by

the function *EDcomp()* by dividing the GR$_{50}$ from transgenic *CYP81A12* or transgenic

CYP81A12 mutants (*atr1* or *atr2*) by the GR$_{50}$ of Col_0. For the dose-response in ½ MS medium

the calculation was the same, using the GR$_{50}$ obtained by the variable green leaves. The *x* and *y*

data from the dose-response curves of both experiments were extracted from R using the

function *write.csv()* and plotted with the treatment means by GraphPad Prism version 9 for

Windows (San Diego, California).

The metabolism data obtained by LC-MS/MS were submitted to analysis of variance

(ANOVA) and the means were compared using Dunnett's test ($p < 0.05$). The peak area of the

herbicides and metabolites of all the different Arabidopsis lines were compared to the Col_0 line.

RESULTS

Arabidopsis mutants

310   SALK lines *atr1-a*, *atr1-b*, and *atr2-a* were segregating for T-DNA insertion and
heterozygous plants were selected to self and seed kept for seed production. *atr2-b* was
homozygous for T-DNA insertion and four plants were self-pollinated to produce seeds. Progeny
plants from *atr1-a*, *atr1-b*, and *atr2-a* were genotyped again and individuals homozygous for the
T-DNA insertion were selected to self and seeds were produced. Homozygous T-DNA insertion
in *ATR1* in *atr1-a* and *atr1-b* lines and homozygous T-DNA insertion in *ATR2* in *atr2-a* and
atr2-b lines were confirmed by PCR. Primers LP and RP for both genes amplified only when
Col_0 DNA was in the reaction. No SALK line amplified the WT version of the respective gene
and only amplified the T-DNA reaction, confirming the homozygosity for T-DNA insertion. The
amplifications were confirmed by electrophoresis in 1% agarose gel (Fig. **S3**), followed by
Sanger sequencing. Based on Sanger sequencing results, the insertion of T-DNA was in the 5'
UTR and in the 4$^{\text{th}}$ intron in *atr1-a and atr1-b* mutant lines, respectively, for *ATR1,* and in the 4$^{\text{th}}$
intron and the 12$^{\text{th}}$ exon in *atr2-a* and *atr2-b* mutant lines, respectively, for *ATR2* (Fig. **S3**).

Transgenic Arabidopsis *CYP81A12* with *atr1* and *atr2* knockdown

324   F1 plants from crosses were genotyped for T-DNA insertion. Around 20 plants were
genotyped from each cross and around 90% of the plants were heterozygous for *ATR1* T-DNA
insertion for cross A and B and heterozygous for *ATR2* T-DNA insertion in cross C and D,
indicating success in the crosses. These plants were genotyped for *CYP81A12* presence by PCR
and plants that had the gene were selected for self-pollination to generate F2. F2 plants
segregated 1:2:1 for *ATR1* T-DNA insertion for cross A and B and for *ATR2* T-DNA insertion
for cross C and D (Tab. **S3**). Homozygous mutants were submitted for PCR amplification of
CYP81A12 and when confirmed, samples were submitted for ddPCR. Four, one, three, and five
plants out of 15, 12, 9 and 15 homozygous had *CYP81A12*. These plants were analyzed by
ddPCR. When only one gene copy was identified in comparison to *ALS* gene, the plants were
self-pollinated, and their progenies were analyzed again using ddPCR.

The ddPCR results indicated the same copy number of *CYP81A12* and *ALS* for the parental

transgenic line *CYP81A12* (Tab. **S4**). Plants selected based on previous genotyping (as indicated

above from crosses A, B, C, and D) had a ratio of *ALS* and *CYP81A12* from 0.9 to 1.1,

suggesting two copies of each gene, one copy per haploid genome. These plants were selected

and used for further experiments.

atr1 and _atr2_ double knockouts

Genotyping analysis of F2 plants obtained from various crosses between *atr1* and *atr2*

mutants demonstrated an absence of double knockout genotypes (Tab. **S5**). A total of 416 F2

plants were genotyped, revealing seven out of the nine possible genotypes. Among the

genotyped plants, 46.2% exhibited heterozygosity for both genes, 5% displayed heterozygosity

for one gene and wildtype for the other, 6.2% exhibited homozygous mutation in one gene and

heterozygosity in the other, while 42.6% exhibited homozygous mutation in one gene and

wildtype in the other. Notably, double wildtype and double mutant genotypes were not observed.

The segregation pattern observed in the F2 generation deviated from the expected Mendelian

segregation for two independent genes, indicating the presence of linked genes. Further

investigation revealed that the *atr1* and *atr2* genes are localized on chromosome four of

Arabidopsis and are approximately two megabases (MBp) apart, providing supporting evidence

that the two genes may be linked.

ATR1 and _ATR2_ gene expression in mutant and transgenic mutant lines

Mutant *atr1-a* showed a reduced transcription of *ATR1* when the primer set used was

spanning the T-DNA insertion (Fig. **4a**); however, normal expression of *ATR1* gene was

observed when primer set used was downstream of the T-DNA insertion (Fig. **4b-d**), indicating

that the T-DNA insertion in 5' UTR did not decrease *ATR1* expression in *atr1-a* line. Mutant line

atr1-b, which had T-DNA insertion in the 4th intron, had significant reduction in *ATR1*

transcripts. This transcript disruption was confirmed in the *ATR1* transcript when the primer set

used spanned the T-DNA insertion and a section of downstream sequence (Fig. **4b-d**).

Interestingly, this line had increased transcript abundance when the primer set was upstream of

the T-DNA (Fig. **4a**). These results indicate a possible compensation in the gene expression by

the plant for the disrupted gene.

Mutants for *ATR2* showed similar results. Mutant *atr2-a* had an upregulation of transcript

upstream of the T-DNA insertion in this line (Fig. **4e**); however, this line had significant

reduction of *ATR2* expression downstream of the T-DNA insertion, especially when the set of

primers used was spanning the T-DNA insertion (Fig. **4f-i**). Mutant *atr2-b* had T-DNA insertion

in the 12th exon and had a reduction in the transcription upstream and downstream of the T-DNA

insertion (Fig. **4e-i**). T-DNA insertion in the *ATR1* mutant *atr1-b* and in the *ATR2* mutants *atr2-a*

and *atr2-b* had knockdown of expression of their respective mRNA. *CYP81A12* expression was

analyzed, and all the transgenic mutant lines had the same transcription production as the

transgenic *CYP81A12* (Fig. **4j**).

Knocking down *ATR1* or *ATR2* did not cause obvious effects on plant growth. Transgenic

CYP81A12 had a lower fresh mass with a smaller height of the plants in comparison to Col_0

(**Fig. S2**).

***atr1* mutant increases herbicide sensitivity in *CYP81A12* line**

Based on whole-plant dose response, transgenic Arabidopsis *CYP81A12* carrying *atr1* and

atr2 were more sensitive to all herbicides tested, including penoxsulam, chlorsulfuron,

mesotrione, and 2,4-D (Fig. **5**, Fig. **S4-7**). The transgenic Arabidopsis *CYP81A12* had a

resistance index of 6.8, 23.6, and 20.9 for penoxsulam, mesotrione, and 2,4-D, respectively (Tab.

1). When *ATR1* was knocked out (*CYP81A12 atr1-b*), the resistance index (RI) decreased to 1.0,

5.7, and 4.2, respectively to penoxsulam, mesotrione, and 2,4-D. The chlorsulfuron doses used in

this study were not able to control the transgenic Arabidopsis *CYP81A12* and *CYP81A12 atr1-a*

(Fig. **5** and Fig. **S4-7**). The resistance to chlorsulfuron of these lines was very high. The lines

CYP81A12 atr2-a and *CYP81A12 atr2-b* showed that the highest doses imparted a slight plant

control, and the model was fitted to the three-parameter model. These lines had a RI of 97.7 and

86.7 for chlorsulfuron (Tab. **1**). The line *CYP81A12 atr1-b* reduced the RI to 3.7 (Tab. **1**).

CYP81A12 atr1-b had the highest decrease in the RI to all herbicides. *ATR2* knockdown lines

(*CYP81A12 atr2-a* and *CYP81A12 atr2-b*) also reduced the RI, but not with the same magnitude

as *atr1-b* (Tab. **1**).

The control of different herbicide doses on Arabidopsis lines grown in ½ MS medium

yielded similar results (Fig. **6** and Tab. **2**). Among the mutants, the *CYP81A12 atr1-b* line

exhibited the most significant reversal of resistance to bensulfuron, penoxsulam, chlorsulfuron,

propoxycarbazone, mesotrione, and bentazon, as evidenced by the highest magnitude of response

(Fig. **6** and Fig. **S7**). Notably, *CYP81A12 atr1-b* displayed symptoms similar to the Col_0 line at lower herbicide doses (Fig. **6**). In contrast, *CYP81A12 atr1-a*, *CYP81A12 atr2-a*, and *CYP81A12 atr2-b* did not exhibit reduced herbicide resistance compared to *CYP81A12*. Specifically, *CYP81A12* displayed resistance indices (RI) of 111.0, 35.7, 2.1, 4.2, 4.6, and 107.3 for bensulfuron, chlorsulfuron, penoxsulam, propoxycarbazone, mesotrione, and bentazon, respectively, while *CYP81A12 atr1-a* demonstrated reduced RIs of 22.3, 10, 0.9, 1.2, 0.9, and 0.9 for the same herbicides, reflecting similar GR_{50} values as Col_0 for penoxsulam, propoxycarbazone, mesotrione, and bentazon (Tab. **2**).

Reduction of herbicide metabolism

Transgenic Arabidopsis line *CYP81A12* had a smaller peak area for the parental mesotrione than Col_0 in all time points after herbicide application, indicating faster herbicide metabolism (Fig. **7a-b**). The transgenic lines *CYP81A12 atr1-a*, *CYP81A12 atr2-a* and *CYP81A12 atr2-b* had similar profiles, with smaller peak area of parental mesotrione than Col_0. The only Arabidopsis line that showed the same peak area as Col_0 was *CYP81A12 atr1-b*. Hydroxylation of mesotrione was higher than Col_0 for all lines in all times points, except for *CYP81A12 atr1-b* (Fig. **7a-b**), which had a very small amount of hydroxy-mesotrione, highlighting that knockdown of *ATR1* prevents mesotrione metabolism in *CYP81A12 atr1-b*. Similar results were observed for chlorsulfuron metabolism (Fig. **S8**). Transgenic *CYP81A12* had a significantly smaller peak area of parental chlorsulfuron than Col_0, inferring faster herbicide metabolism. The transgenic mutant line *CYP81A12 atr1-a* was not different in area peak of parental chlorsulfuron and hydroxy-chlorsulfuron when compared to Col_0, indicating a loss of the metabolism capacity.

NADPH-cytochrome P450 reductase in weeds

Thirteen different weed species were analyzed, and their corresponding CPRs (cytochrome P450 reductase) were identified (Fig. **8**). The number of CPR copies varied within each weed species, ranging from two to three copies (Tab. **S6**). Through the construction of a phylogenetic tree, CPRs can be categorized into two distinct classes, namely CPR I and CPR II (Fig. **8**). These findings align with previous research, which suggests that CPR Class I is exclusive to dicot species, while Class II is present in both monocots and dicots (Jensen & Møller, 2010, Istiandari *et al.*, 2021).

To differentiate between the CPRs within weed species, they were denoted as "a", "b", or
"c" and individual copies were numbered as ".1", ".2", or ".3" to represent one, two, or three
copies, respectively, in the present study. Kochia (*Bassia scoparia*), common waterhemp,
common morningglory (*Ipomea purpurea*), and Djulis (*Chenopodium formosanum*) have both
CPR 1 and CPR 2 (Fig. **8**). Common morning glory, a tetraploid, possesses two copies of CPR 1
and one copy of CPR 2, whereas the hexaploid Djulis has three copies of CPR 1 and three copies
of CPR 2. The diploid kochia hasone copy of each CPR class. On the other hand, the diploids
common waterhemp and Palmer amaranth possess one copy of CPR1 and two copies of CPR2
and horseweed possess three copies of only CPR class II. Class II Palmer amaranth CPRs have
93% identity, while the CPRs of horseweed (Class II) had a 74-76% identity among their three
copies. Djulis, a hexaploid species with 27 chromosomes, exhibits three copies of each CPR
class. Notably, each copy is localized within a different subgenome (B, C, and D).
Monocot plants exhibit diverse CPR compositions within CPR Class II, forming a distinct
branch. Although their classification into specific classes is not documented in the literature,
various monocot species possess CPR variants within Class II and we propose a division of CPR
variants in grasses into CPRa, CPRb and CPRc class II (Fig. **8**). For instance, blackgrass and
goosegrass each possess three different CPR II variants: Am_CPRa, Am_CPRb, and Am_CPRc
for blackgrass, and Ei_CPRa, Ei_CPRb, and Ei_CPRc for goosegrass. These CPR variants are
categorized into separate branches based on phylogenetic analysis. Similarly, annual bluegrass, a
tetraploid species, possesses three CPR variants, with two copies of each CPR variant present in
subgenomes A or B. Similarly, late watergrass, an allotetraploid species, exhibits two copies of
each of the three CPR variants, distributed across its subgenomes. Jungle rice, an allohexaploid
species, possesses nine CPR variants, with copies of each CPR variant located in subgenomes D,
E, or F. On the other hand, barnyardgrass, also an allohexaploid species, diverges from jungle
rice as it possesses eight CPR variants. Among these variants, two CPRs (CPRa and CPRc) have
three copies each, while one CPR (CRPb) has two copies. It is noteworthy that the diploid downy
brome is the sole monocot weed species analyzed in this study with only two CPR variants,
whereas the remaining monocot weed species all possess three distinct CPR variants with
different copy numbers.
All CPR genes had a well conserved domain (Fig. **S9**). Among the identified CPRs in
weed genomes, three CPRs, namely Eco_CPRc.1, Ec_CPRc.3, and Eo_CPRc.1, stand out with

significantly larger protein sizes compared to the others. These three CPRs have 1520, 1905, and

1948 amino acids, respectively, whereas the remaining CPRs range from 689 to 823 amino acids.

Eo_CPRc.1, Ec_CPRc.3 had a bigger *N*-terminal and additionally Eco_CPRc.1 had a long *C*-

terminal extensions matching phosphatidylinositol 3-kinase, which could be an assembly error.

Additionally, these three CPRs exhibit a similarity ranging from 87% to 94% among themselves.

Additionally, there is close genetic relationship between subgenome D of jungle rice and

subgenome C of barnyardgrass (Aoki & Yamaguchi, 2008, Wu *et al.*, 2022). This suggests a

common diploid ancestor, explaining the high similarity observed in CPRs among these three

species.

DISCUSSION

Cytochrome P450 reductases (CPR) are a class of enzymes that play a crucial role in the

electron transfer system from NADPH to P450s (Mizutani & Ohta, 1998, Jensen & Møller,

2010). While there are numerous P450 genes, the number of CPR genes is much smaller (Backes

& Kelley, 2003). In Arabidopsis, two CPR genes, class I *ATR1* and class II *ATR2*, have been

identified. *ATR1* is consistently expressed, while *ATR2* is induced in response to stress (Mizutani

& Ohta, 1998, Jensen & Møller, 2010). Since P450s are implicated in metabolic herbicide

resistance in weeds, it is important to understand the specific role of P450 reductases in the

herbicide-metabolizing capabilities of these P450s.

Through the generation of transgenic mutant Arabidopsis lines, i.e., transgenic Arabidopsis

carrying *CYP81A12* from late watergrass able to metabolize a wide range of herbicides and with

knockdown of either the *ATR1* or *ATR2* gene, our observations indicate that *ATR1* plays a

predominant role in facilitating electron transfer from NADPH to *CYP81A12* in this particular

case. Upon knocking down the *ATR1* gene in the transgenic *CYP81A12* plants (*CYP81A12 atr1-

b*), reductions in herbicide resistance for 2,4-D and mesotrione were observed. Conversely, the

CYP81A12 atr1-a line, with a T-DNA insertion in the 5' UTR of *ATR1*, did not exhibit a

reduction in herbicide resistance due to insufficient gene expression reduction (Fig. **4**).

Knockouts of the *ATR2* gene (*CYP81A12 atr2a* and *atr2-b*) also led to a decrease in herbicide

resistance, although not to the same extent as *ATR1*.

In Arabidopsis, *ATR1* is commonly referred to as *CPR1*, while *ATR2* is known as *CPR2* in

other species. Among the identified CPRs in weeds, it was observed that only CPR class II is

present in grasses. In our current study, the *ATR1* of Arabidopsis was the most crucial in

providing electrons to *CYP81A12*, probably due to its best interaction efficiency of CPR:P450

complex or due its higher basal activity (TAIR, 2024). We hypothesize that different CPRs may

exhibit varying interactions with various P450s, including those involved in herbicide

metabolism. A recent investigation utilizing human CPRs indicated that mutations in the FMN

binding domain significantly enhanced CYP activity. This suggests that amino acid variations

within this domain play a significant role in the diverse interactions between CPRs and P450s

(Esteves *et al.*, 2020). Another study indicates that distinct CPR classes sourced from various

leguminous plants (such as *Medicago truncatula*, *Lotus japonicus*, and *Glycyrrhiza uralensis*)

exhibit diverse interactions with distinct P450 enzyme families within transgenic yeast.

Consequently, this variation leads to differing levels of hydroxylation activity on the substrate.

(Cheng *et al.*, 2023). Based on our results, it is evident that the presence of CPR in weed species

exhibits variation. Notably, all examined grasses possess at least three different homologs of

CPRs class II, CPRa, CPRb and CPRc, except for downy brome, which has two. One of the three

different CPR class II homologs might have a higher activity and contribute the most towards

reducing certain P450 genes in these species. This discrepancy in the number of CPRs among

grasses suggests duplication events that occurred prior to species differentiation within the grass

family (Jensen & Møller, 2010). A detailed analysis of protein sequences revealed substantial

differences primarily in the *N-* and *C*-terminal regions. However, the FMN-binding and FAD-

binding domains in all CPRs displayed a high degree of conservation due to their specialized

functions (Fig. **S9**) and because of that, it is hypothesized that electrostatic interactions are

considered to play a crucial role in guiding the formation of P450:CPR complexes.

The presence of varying numbers of CPRs in different weed species suggests distinct

genetic pressures these weeds have encountered over time, leading to the acquisition of diverse

functions. If herbicide application has selected resistant plants with enhanced herbicide-

metabolizing activity, we hypothesize that herbicide use may have also selected for higher CPR

activity to support the increased activity of P450s by facilitating electron transfer; however, this

aspect needs further investigation. Additionally, in our study, we observed that in Arabidopsis,

ATR2 is unable to complement the electron transfer to *CYP81A12* when *atr1* is knocked out.

However, this may not be the case for other weed species that possess different CPR isoforms.

For instance, jungle rice has three CPR isoforms, each with three copies. These CPRs could

potentially be paralogs with distinct functions, interacting differentially with various P450

enzyme families, or they may have complementary functions.

Cytochrome P450 monooxygenases play pivotal roles in both primary and secondary

metabolism in plants (Werck-Reichhart & Feyereisen, 2000). This significance is exemplified by

studies such as the one involving homozygous mutants for CYP51A2, which exhibit defects in

early embryogenesis, underscoring the enzyme's critical role in plant growth and development

(Kim *et al.*, 2005). In our study, we were not able to generate *ATR1* and *ATR2* double knockouts

through crosses, likely due to their inherent linkage. The hypothetical scenario of *ATR1* and

ATR2 double knockout raises intriguing possibilities, suggesting the hypothesis that a double

mutation might render plants inviable due to reduced P450 activity. For most weed species

examined in this study, the CPRs are dispersed across different chromosomes and do not exhibit

linkage. Nevertheless, our laboratory is actively pursuing the generation of a double knockout

Arabidopsis line using CRISPR/Cas 9 to gain further insights into the role of *ATR1* and *ATR2* in

metabolic resistance mechanisms in weeds.

Research on the role of CPR in the evolution of pesticide resistance has primarily focused

on insects, with limited investigations into the functions of CPRs in plants. In insect studies,

inhibiting CPR in deltamethrin-resistant *Cimex lectularius* increased susceptibility to

deltamethrin, indicating that CPR could be targeted to reduce P450 activity (Zhu *et al.*, 2012).

Similarly, downregulating CPR in resistant *Tetranychus cinnabarinus* (carmine spider mite) and

T. urticae (spider mite) through RNA interference reduced P450 activity and increased

susceptibility to fenpropathrin and various acaricides, respectively (Adesanya *et al.*, 2020). In

plants, which possess one, two, or three CPR isoforms, no research exists regarding the diverse

functions of CPRs for herbicide evolution. For instance, Arabidopsis *ATR2*, which is co-

expressed with lignin biosynthetic genes, showed significant alterations in lignin composition

when mutated, leading to reduced activity of P450 enzymes such as cinnamate 4-hydroxylase, p-

coumarate 3-hydroxylase, and ferulate 5-hydroxylase (Ehlting *et al.*, 2005, Sundin *et al.*, 2014).

The crystal structure of Arabidopsis *ATR2* revealed that mutations in the interflavin electron

transfer region can impact ATR2 activity (Niu *et al.*, 2017), offering opportunities to exploit

these reliable enzymes for reducing P450 activity against herbicide metabolism.

Protein alignments revealed the presence of conserved domains shared across various

CPRs, along with significant variation in the FMN binding domain (Fig. **6**). There were

conserved regions among CPRs in both dicot and monocot weeds, presenting a valuable

opportunity to target and reduce P450 activity. Inhibiting CPR activity, either through chemical

inhibitors or RNA-targeting approaches, offers a potential means to restore herbicide

effectiveness against weeds exhibiting metabolic herbicide resistance. Combining CPR inhibition

with herbicide treatments can serve as a strategy to control metabolic herbicide-resistant weed

populations and, additionally, delay the evolution of metabolic herbicide resistance in sensitive

weed populations.

The study of cytochrome P450 reductases and their interactions with cytochrome P450

enzymes is crucial for understanding metabolic herbicide resistance in weeds. Our findings

highlight the importance of CPRs, particularly *ATR1* in this case of transgenic Arabidopsis, in

transferring electrons to P450s involved in herbicide metabolism. The presence of different

numbers of CPRs in weed species indicates the diverse genetic pressures these weeds have

undergone. Our results suggest that CPR is a potential target to reduce herbicide metabolism.

Identifying the most important CPRs in weeds that are closely involved in herbicide metabolism

is very promising as a new tool for weed management using gene silencing approaches (Zabala-

Pardo *et al.*, 2022). Further research on CPRs and their functions in weeds will contribute to our

understanding of herbicide resistance evolution and the development of effective weed

management strategies.

ACKNOWLEDGEMENTS

This work was supported by funding from Cotton Inc.

AUTHOR CONTRIBUTIONS

CAGR, FED, and TAG designed the research. CAGR performed experiments. FED, TAG and

SAI contributed experimental materials. CAGR, FED analyzed the data. CAGR, SAI, FED, and

TAG wrote the paper. All authors contributed to editing and approving the paper.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare.

576
577

REFERENCES

Adesanya, AW, Cardenas, A, Lavine, MD, Walsh, DB, Lavine, LC & Zhu, F. 2020. RNA interference of NADPH-cytochrome P450 reductase increases susceptibilities to multiple acaricides in *Tetranychus urticae. Pesticide Biochemistry and Physiology,* **165**: 104550.

Aoki, D & Yamaguchi, H. 2008. Genetic relationship between *Echinochloa crus-galli* and *Echinochloa oryzicola* accessions inferred from internal transcribed spacer and chloroplast DNA sequences. *Weed Biology and Management,* **8**: 233-242.

Backes, WL & Kelley, RW. 2003. Organization of multiple cytochrome P450s with NADPH-cytochrome P450 reductase in membranes. *Pharmacology & Therapeutics,* **98**: 221-233.

Benson, DA, Cavanaugh, M, Clark, K, Karsch-Mizrachi, I, Lipman, DJ, Ostell, J & Sayers, EW. 2013. GenBank. *Nucleic Acids Research,* **41**: D36-42.

Cai, L, Comont, D, MacGregor, D, Lowe, C, Beffa, R, Neve, P & Saski, C. 2023. The blackgrass genome reveals patterns of non-parallel evolution of polygenic herbicide resistance. *New Phytologist,* **237**: 1891-1907.

Chen, J, Huang, Z, Huang, H, Wei, S, Liu, Y, Jiang, C, Zhang, J & Zhang, C. 2017. Selection of relatively exact reference genes for gene expression studies in goosegrass (Eleusine indica) under herbicide stress. *Scientific Reports,* **7**: 46494.

Cheng, J, Li, G, Wang, X, Yang, C, Xu, F, Qian, Z & Ma, X. 2023. Cloning and functional characterization of NADPH-cytochrome P450 reductases in *Aconitum vilmorinianum. Molecules,* **28**: 7409.

Dimaano, NG & Iwakami, S. 2021. Cytochrome P450-mediated herbicide metabolism in plants: current understanding and prospects. *Pest Management Science,* **77**: 22-32.

Dimaano, NG, Yamaguchi, T, Fukunishi, K, Tominaga, T & Iwakami, S. 2020. Functional characterization of cytochrome P450 *CYP81A* subfamily to disclose the pattern of cross-resistance in *Echinochloa phyllopogon. Plant Molecular Biology,* **102**: 403-416.

Ehlting, J, Mattheus, N, Aeschliman, DS, Li, E, Hamberger, B, Cullis, IF, Zhuang, J, Kaneda, M, Mansfield, SD, Samuels, L, et al. 2005. Global transcript profiling of primary stems from *Arabidopsis thaliana* identifies candidate genes for missing links in lignin biosynthesis and transcriptional regulators of fiber differentiation. *The Plant Journal,* **42**: 618-40.

Esteves, F, Campelo, D, Gomes, BC, Urban, P, Bozonnet, S, Lautier, T, Rueff, J, Truan, G & Kranendonk, M. 2020. The role of the FMN-domain of human cytochrome P450 oxidoreductase in its promiscuous interactions with structurally diverse redox partners. *Frontiers in Pharmacology,* **11**: 299.

Gaines, TA, Duke, SO, Morran, S, Rigon, CAG, Tranel, PJ, Küpper, A & Dayan, FE. 2020. Mechanisms of evolved herbicide resistance. *Journal of Biological Chemistry,* **295**: 10307–10330.

Gupta, S, Harkess, A, Soble, A, Van Etten, M, Leebens-Mack, J & Baucom, RS. 2023. Interchromosomal linkage disequilibrium and linked fitness cost loci associated with selection for herbicide resistance. *New Phytologist,* **238**: 1263-1277.

Hall, N, Chen, J, Saski, C, Westra, P, Gaines, T & Patterson, E. 2023. FHY3/FAR1 transposable elements generate adaptive genetic variation in the *Bassia scoparia* genome. *bioRxiv*: 2023.05.26.542497.

Hallgren, J, Tsirigos, K, Pedersen, MD, Almagro Armenteros, JJ, Marcatili, P, Nielsen, H, Krogh, A & Winther, O. 2022. DeepTMHMM predicts alpha and beta transmembrane proteins using deep neural networks. bioRxiv.

Istiandari, P, Yasumoto, S, Srisawat, P, Tamura, K, Chikugo, A, Suzuki, H, Seki, H, Fukushima, EO & Muranaka, T. 2021. Comparative analysis of NADPH-cytochrome P450 reductases from legumes for heterologous production of triterpenoids in transgenic *Saccharomyces cerevisiae*. *Frontiers in Plant Science,* **12**: 762546.

Iwakami, S, Endo, M, Saika, H, Okuno, J, Nakamura, N, Yokoyama, M, Watanabe, H, Toki, S, Uchino, A & Inamura, T. 2014. Cytochrome P450 *CYP81A12* and *CYP81A21* are associated with resistance to two acetolactate synthase inhibitors in *Echinochloa phyllopogon*. *Plant Physiology,* **165**: 618-629.

Jarvis, DE, Sproul, JS, Navarro-Domínguez, B, Krak, K, Jaggi, K, Huang, YF, Huang, TY, Lin, TC, Jellen, EN & Maughan, PJ. 2022. Chromosome-scale genome assembly of the hexaploid Taiwanese goosefoot "Djulis" (*Chenopodium formosanum*). *Genome Biology and Evolution,* **14**.

Jensen, K & Møller, BL. 2010. Plant NADPH-cytochrome P450 oxidoreductases. *Phytochemistry,* **71**: 132-41.

Jeschke, P, Witschel, M, Krämer, W & Schirmer, U. 2019. *Modern Crop Protection Compounds,* Wiley-VCH Verlag GmbH & Co. KGaA, Wiley.

Kim, HB, Schaller, H, Goh, CH, Kwon, M, Choe, S, An, CS, Durst, F, Feldmann, KA & Feyereisen, R. 2005. Arabidopsis cyp51 mutant shows postembryonic seedling lethality associated with lack of membrane integrity. *Plant Physiology,* **138**: 2033-47.

Laforest, M, Martin, SL, Bisaillon, K, Soufiane, B, Meloche, S & Page, E. 2020. A chromosome-scale draft sequence of the Canada fleabane genome. *Pest Management Science,* **76**: 2158-2169.

Lau, SM & O'Keefe, DP. 1996. Analysis of herbicide metabolism by monocot microsomal cytochrome P450. *Methods Enzymology,* **272**: 235-42.

Lightfoot, DJ, Jarvis, DE, Ramaraj, T, Lee, R, Jellen, EN & Maughan, PJ. 2017. Single-molecule sequencing and Hi-C-based proximity-guided assembly of amaranth (*Amaranthus hypochondriacus*) chromosomes provide insights into genome evolution. *BMC Biology,* **15**: 74.

Livak, KJ & Schmittgen, TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(T)(-Delta Delta C) method. *Methods,* **25**: 402-408.

Mizutani, M & Ohta, D. 1998. Two isoforms of NADPH: Cytochrome P450 reductase in *Arabidopsis thaliana*: gene structure, heterologous expression in insect cells, and differential regulation. *Plant Physiology,* **116**: 357-367.

Montgomery, JS, Giacomini, D, Waithaka, B, Lanz, C, Murphy, BP, Campe, R, Lerchl, J, Landes, A, Gatzmann, F, Janssen, A, et al. 2020. Draft genomes of *Amaranthus tuberculatus*, *Amaranthus hybridus*, and *Amaranthus palmeri*. *Genome Biology and Evolution,* **12**: 1988-1993.

Montgomery, JS, Morran, S, MacGregor, DR, McElroy, JS, Neve, P, Neto, C, Vila-Aiub, MM, Sandoval, MV, Menéndez, AI, Kreiner, JM, et al. 2023. The international weed genomics consortium: community resources for weed genomics research. *Genome Biology,* **in press**: 2023.07.19.549613.

Mukherjee, G, Nandekar, PP & Wade, RC. 2021. An electron transfer competent structural ensemble of membrane-bound cytochrome P450 1A1 and cytochrome P450 oxidoreductase. *Communications Biology,* **4**: 55.

Munro, AW, Noble, MA, Robledo, L, Daff, SN & Chapman, SK. 2001. Determination of the redox properties of human NADPH-cytochrome P450 reductase. *Biochemistry,* **40**: 1956-1963.

Nelson, DR, Schuler, MA, Paquette, SM, Werck-Reichhart, D & Bak, S. 2004. Comparative genomics of rice and Arabidopsis. Analysis of 727 cytochrome P450 genes and pseudogenes from a monocot and a dicot. *Plant Physiology,* **135**: 756-772.

Niu, G, Zhao, S, Wang, L, Dong, W, Liu, L & He, Y. 2017. Structure of the *Arabidopsis thaliana* NADPH-cytochrome P450 reductase 2 (*ATR2*) provides insight into its function. *The FEBS Journal,* **284**: 754-765.

Paine, MJI, Scrutton, NS, Munro, AW, Gutierrez, A, Roberts, GCK & Wolf, CR. 2005. Electron transfer partners of cytochrome. *In:* NEW YORK, N. S. (ed.) *Cytochrome P450, structure, mechanism, and biochemistry.* New York.

Pandian, BA, Sathishraj, R, Djanaguiraman, M, Prasad, PVV & Jugulam, M. 2020. Role of cytochrome P450 enzymes in plant stress response. *Antioxidants,* **9**: 454.

Paquette, SM, Jensen, K & Bak, S. 2009. A web-based resource for the Arabidopsis P450, cytochromes b5, NADPH-cytochrome P450 reductases, and family 1 glycosyltransferases (http://www.P450.kvl.dk). *Phytochemistry,* **70**: 1940-7.

Parage, C, Foureau, E, Kellner, F, Burlat, V, Mahroug, S, Lanoue, A, Dugé de Bernonville, T, Londono, MA, Carqueijeiro, I, Oudin, A, et al. 2016. Class II cytochrome P450 reductase governs the biosynthesis of alkaloids. *Plant Physiology,* **172**: 1563-1577.

Peng, YH, Lai, Z, Lane, T, Nageswara-Rao, M, Okada, M, Jasieniuk, M, O'Geen, H, Kim, RW, Sammons, RD, Rieseberg, LH, et al. 2014. De novo genome assembly of the economically important weed horseweed using integrated data from multiple sequencing platforms. *Plant Physiology,* **166**: 1241-54.

Porter, TD & Kasper, CB. 1986. NADPH-cytochrome P-450 oxidoreductase: flavin mononucleotide and flavin adenine dinucleotide domains evolved from different flavoproteins. *Biochemistry,* **25**: 1682-1687.

Powles, SB & Yu, Q. 2010. Evolution in Action: Plants Resistant to Herbicides. *Annual Review of Plant Biology,* **61**: 317-347.

R Core Team. 2023. R: A language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria. https://www.R-project.org/. Vienna, Austria.

Rana, S, Lattoo, SK, Dhar, N, Razdan, S, Bhat, WW, Dhar, RS & Vishwakarma, R. 2013. NADPH-cytochrome P450 reductase: molecular cloning and functional characterization of two paralogs from *Withania somnifera* (L.) Dunal. *PLos One,* **8**: 16.

Revolinski, SR, Maughan, PJ, Coleman, CE & Burke, IC. 2023. Preadapted to adapt: underpinnings of adaptive plasticity revealed by the downy brome genome. *Communications biology,* **6**: 326.

Rigon, CAG, Gaines, TA, Küpper, A & Dayan, FE. 2020. Metabolism-based herbicide resistance, the major threat among the non-target site resistance mechanisms. *Outlooks on Pest Management,* **31**: 162-168.

Ritz, C, Baty, F, Streibig, JC & Gerhard, D. 2015. Dose-response analysis using R. *PLos One,* **10**: 13.

Robbins, MD, Bushman, BS, Huff, DR, Benson, CW, Warnke, SE, Maughan, CA, Jellen, EN, Johnson, PG & Maughan, PJ. 2023. Chromosome-scale genome assembly and annotation of allotetraploid annual bluegrass (*Poa annua* L.). *Genome Biology and Evolution,* **15**.

Sundin, L, Vanholme, R, Geerinck, J, Goeminne, G, Höfer, R, Kim, H, Ralph, J & Boerjan, W. 2014. Mutation of the unducible *Arabidopsis thaliana* cytochrome P450 alters lignin composition and improves saccharification. *Plant Physiology,* **166**: 1956-1971.

TAIR. 2024. *The Arabidopsis Information Resource (TAIR)* [Online]. Available: www.arabidopsis.org [Accessed Mar 23, 2024 2024].

Urban, P, Mignotte, C, Kazmaier, M, Delorme, F & Pompon, D. 1997. Cloning, yeast expression, and characterization of the coupling of two distantly related *Arabidopsis thaliana* NADPH-cytochrome P450 reductases with P450 *CYP73A5*. *Journal of Biological Chemistry,* **272**: 19176-19186.

Varadarajan, J, Guilleminot, J, Saint-Jore-Dupas, C, Piégu, B, Chabouté, ME, Gomord, V, Coolbaugh, RC, Devic, M & Delorme, V. 2010. ATR3 encodes a diflavin reductase essential for Arabidopsis embryo development. *New Phytologist,* **187**: 67-82.

Wang, M, Roberts, DL, Paschke, R, Shea, TM, Masters, BSS & Kim, J-JP. 1997. Three-dimensional structure of NADPH–cytochrome P450 reductase: Prototype for FMN- and FAD-containing enzymes. *Proceedings of the National Academy of Sciences,* **94**: 8411-8416.

Werck-Reichhart, D & Feyereisen, R. 2000. Cytochromes P450: a success story. *Genome Biology,* **1**: reviews3003.1.

Wu, D, Shen, E, Jiang, B, Feng, Y, Tang, W, Lao, S, Jia, L, Lin, H-Y, Xie, L, Weng, X, et al. 2022. Genomic insights into the evolution of *Echinochloa* species as weed and orphan crop. *Nature Communications,* **13**: 689.

Zabala-Pardo, D, Gaines, T, Lamego, FP & Avila, LA. 2022. RNAi as a tool for weed management: challenges and opportunities. *Advances in Weed Science,* **40**: e020220096.

Zhang, C, Johnson, NA, Hall, N, Tian, X, Yu, Q & Patterson, EL. 2023. Subtelomeric 5-enolpyruvylshikimate-3-phosphate synthase copy number variation confers glyphosate resistance in *Eleusine indica*. *Nature Communications,* **14**: 4865.

Zhu, F, Sams, S, Moural, T, Haynes, KF, Potter, MF & Palli, SR. 2012. RNA interference of NADPH-cytochrome P450 reductase results in reduced insecticide resistance in the bed bug, *Cimex lectularius*. *Plos One,* **7**: e31037.

Table 1. Parameters of the logistic equation and resistance index (RI) for shoot fresh weight (% of untreated control) of wild-type (Col_0), transgenic Arabidopsis *CYP81A12*, transgenic Arabidopsis mutants *CYP81A12 atr1-a*, *CYP81A12 atr1-b*, *CYP81A12 atr2-a* and *CYP81A12 atr2-b* treated with different doses of penoxsulam, chlorsulfuron, mesotrione and 2,4-D.

Line	Herbicide	b	GR50 Dose (g ha^{-1})	Lower CI (95%)	Upper CI (95%)	SE	RI	*p*-value
Col_0		0.5	2.3	0.9	3.7	0.7	-	
CYP81A12		0.7	15.7	6.7	24.6	4.5	6.8	<0.001
CYP81A12 atr1-a		1.8	17.1	11.2	23.1	3.0	7.4	0.01
CYP81A12 atr1-b	PENO	0.7	2.3	1.2	3.5	0.6	1.0	0.9
CYP81A12 atr2-a		2.9	6.7	5.0	8.4	0.8	2.9	0.04
CYP81A12 atr2-b		0.7	3.4	1.5	5.2	0.9	1.5	0.4
Col_0		1.3	0.9	0.7	1.1	0.1	-	-
CYP81A12		-	-	-	-	-	-	-
CYP81A12 atr1-a		-	-	-	-	-	-	-
CYP81A12 atr1-b	CHLO	2.4	3.4	2.5	4.4	0.4	3.7	<0.01
CYP81A12 atr2-a		2.6	87.9	1.8	174.1	42.5	97.7	0.03
CYP81A12 atr2-b		0.6	77.5	1.1	153.9	37.7	86.7	0.04
Col_0		3.7	4.4	3.85	5.1	0.3	-	-
CYP81A12		15	93.6	58.6	128.6	17.7	20.9	<0.001
CYP81A12 atr1-a		5.5	89.3	78.9	99.7	5.2	19.9	<0.001
CYP81A12 atr1-b	MESO	4.3	26.8	23	30.5	1.9	5.7	<0.001
CYP81A12 atr2-a		2.8	41.8	35.5	48.1	3.2	9.3	<0.001
CYP81A12 atr2-b		2.9	59.1	50.3	68.1	4.5	13.2	<0.001
Col_0		0.5	5.9	2.7	9.3	1.6	-	-
CYP81A12		1.2	141.7	96.5	186.5	22.7	23.6	<0.001
CYP81A12 atr1-a		1.6	103.3	73.7	132.8	14.9	17.2	<0.001
CYP81A12 atr1-b	2,4-D	2.9	25.3	20.15	30.4	2.6	4.2	<0.001
CYP81A12 atr2-a		0.8	50.1	29.6	70.5	10.3	8.3	<0.001
CYP81A12 atr2-b		3.8	130.1	108.1	151.9	11.1	21.6	<0.001

b: slope; d: upper limit was fixed to 100; GR50: herbicide dose that causes a 50% reduction in shoot fresh weight variable; CI: confidence interval of parameter GR50 ($\alpha = 0.05$); RI: resistance index = GR50 ratio between lines with Col_0 (control). PENO – penoxsulam, CHLO – chlorsulfuron, MESO – mesotrione.

Table 2. Parameters of the logistic equation and resistance index (RI) for leaf number of Arabidopsis seedlings (% of untreated control) of wild-type, *atr1-a*, *atr1-b*, *atr2-a*, *atr2-b,* and Arabidopsis transgenic *CYP81A12*, Arabidopsis transgenic mutants *CYP81A12 atr1-a*, *CYP81A12 atr1-b*, *CYP81A12 atr2-a*, *CYP81A12 atr2-b* growing in ½ MS medium supplemented with different doses of herbicides.

Lines		b	ED50 dose (nM)	Lower CI (95%)	Upper CI (95%)	SE	RI	*p*-value
Col_0		35.7	9.5	9.1	9.8	0.17	-	-
atr1-a		32.9	3.4	-16.3	23.0	10.0	0.3	0.5
atr1-b		31.5	9.0	8.2	9.9	0.4	0.9	0.4
atr2-a		27.7	9.1	8.3	9.8	0.4	0.9	0.4
atr2-b	BSM	26.6	6.7	-12.9	26.4	9.9	0.8	0.8
CYP81A12		18.3	1055.5	1034.3	1071.8	9.5	111.0	<0.001
CYP81A12 atr1-a		3.6	1368.4	1321.6	1360.9	9.9	141.4	<0.001
CYP81A12 atr1-b		24.9	185.2	192.4	231.7	10.0	22.3	<0.001
CYP81A12 atr2-a		2.0	1547.8	1528.2	1567.5	9.9	163.2	<0.001
CYP81A12 atr2-b		5.6	942.6	923.1	962.2	9.9	99.4	<0.001
Col_0		3.6	3.7	0.73	6.73	1.5	-	-
atr1-a		1.3	1.2	0.8	1.5	0.2	0.3	<0.01
atr1-b		1.7	2.4	1.4	3.3	0.5	0.6	0.03
atr2-a		7.3	3.6	-1.1	8.4	2.4	0.9	0.19
atr2-b	CHL	2.1	1.9	1.4	2.3	0.2	0.5	<0.01
CYP81A12		1.2	132.2	81.9	182.6	25.08	35.7	0.02
CYP81A12 atr1-a		2.7	121.6	93.9	149.2	13.8	32.8	<0.01
CYP81A12 atr1-b		2.3	37.2	27.74	46.66	4.7	10.0	<0.01
CYP81A12 atr2-a		5.4	129.9	8.5	251.4	60.5	35.1	0.1
CYP81A12 atr2-b		0.7	76.1	17.5	134.6	29.1	20.5	0.07
Col_0		3.6	13.8	11.8	15.7	0.95	-	-
atr1-a		1.4	3.8	2.6	4.9	0.5	0.3	<0.001
atr1-b		0.9	3.3	0.5	6.1	1.4	0.2	0.02
atr2-a		1.0	2.7	1.1	4.37	0.8	0.2	<0.001
atr2-b	PEN	3.1	7.0	5.9	8.1	0.5	0.5	<0.001
CYP81A12		8.8	29.6	13.4	45.8	8.0	2.1	<0.001
CYP81A12 atr1-a		10.3	26.9	10.3	43.5	8.21	1.9	<0.01
CYP81A12 atr1-b		11.6	11.9	-8.8	32.8	10.4	0.9	0.25
CYP81A12 atr2-a		10.1	27.8	9.8	45.8	8.9	2.0	<0.01
CYP81A12 atr2-b		8.5	23.7	13.2	34.3	5.2	1.7	<0.001
Col_0		2.4	3.5	2.8	4.0	0.3	0.9	-
atr1-a	PRO	2.1	3.4	1.8	5.1	0.8	0.9	0.9
atr1-b		3.1	3.2	2.5	3.9	0.3	0.9	0.7

Line	Treatment	b	GR50	CI lower	CI upper	SE	RI	P
atr2-a		8.1	3.8	0.2	7.4	1.8	1.1	0.9
atr2-b		2.4	2.4	1.6	3.2	0.4	0.8	0.2
CYP81A12		2.4	15.5	12.1	19.0	1.7	4.2	<0.001
CYP81A12 atr1-a		1.9	13.0	10.9	15.1	1.0	3.7	<0.001
CYP81A12 atr1-b		2.1	4.3	3.7	4.9	0.3	1.2	0.2
CYP81A12 atr2-a		2.4	15.7	10.3	21.1	2.6	4.3	<0.001
CYP81A12 atr2-b		1.8	11.1	7.6	14.5	1.7	3.2	<0.001
Col_0		4.5	159.2	131.8	186.7	13.9	-	-
atr1-a		5.4	129.3	106.8	151.8	11.9	0.8	0.3
atr1-b		6.5	160.4	114.8	206.1	23.2	1	0.9
atr2-a		6.3	124.3	84.97	163.6	20.1	0.8	0.1
atr2-b	MES	3.8	117.7	100.4	135.1	8.3	0.7	0.03
CYP81A12		7.9	740.7	604.7	876.78	69.19	4.6	<0.0001
CYP81A12 atr1-a		2.9	480.2	415.1	545.2	33.1	3	<0.0001
CYP81A12 atr1-b		5.1	142.8	112.9	172.72	15.2	0.9	0.4
CYP81A12 atr2-a		6.1	564	492.2	634	36.1	3.5	<0.001
CYP81A12 atr2-b		2.9	585.2	495.7	675.1	45.6	3.6	<0.0001
Col_0		5.4	0.1	0.09	0.11	0.004	-	-
atr1-a		3.9	0.01	0.01	0.02	0.001	0.1	<0.001
atr1-b		3.1	0.03	0.02	0.05	0.007	0.3	<0.001
atr2-a		2.3	0.03	0.02	0.04	0.005	0.3	<0.001
atr2-b	BEN	2.2	0.02	0.01	0.03	0.002	0.2	<0.001
CYP81A12		5	11.24	9.57	12.9	0.82	107.3	<0.001
CYP81A12 atr1-a		3.7	12.84	10.7	14.9	1	132.3	<0.001
CYP81A12 atr1-b		5.22	0.08	-0.07	0.2	0.07	0.9	0.8
CYP81A12 atr2-a		8.43	11.7	3	20.5	4.35	111.6	<0.001
CYP81A12 atr2-b		3.4	12.2	10.2	14.3	1.01	114.6	<0.001

b: slope; d: upper limit was fixed to 100; GR50: herbicide dose that causes a 50% reduction in shoot fresh weight variable; CI: confidence interval of parameter GR50 ($\alpha = 0.05$); RI: resistance index = ED50 ratio between lines with Col_0 (control).

Fig. 1. Process of electron transfer from NADPH-cytochrome P450 reductase to the catalytic site of cytochrome P450 monooxygenase. Two electrons are sequentially transferred from NADPH to the heme group of P450 through the cofactors FAD and FMN, each transferred in single-electron steps. Cytochrome P450, once activated by the electron transfer, catalyzes the oxidative breakdown of a substrate by inserting one oxygen atom into the chemical compound. Created with Biorender.

Fig. 2. Schemes representing the Arabidopsis crosses to generate transgenic Arabidopsis *CYP81A12* with loss of function in *atr1* or *atr2*. F1 plants were genotyped for T-DNA insertion and *CYP81A12* presence. F2 plants were genotyped for homozygous T-DNA insertion and two copies of *CYP81A12* using ddPCR. M - male, F - female. Created with BioRender.com.

Fig. 3. T-DNA location in *ATR1* (A) and *ATR2* (B) genes in SALK lines and primers localization for real time qPCR gene expression analysis. Yellow triangle locates the T-DNA insertion in the genes, *atr1-a* and *atr2-b* have T-DNA insertion in the 5'UTR and 4[th] intron, respectively, and *atr2-a* and *atr2-b* have T-DNA insertion in the 3[rd] and 12[th] exon, respectively. Boxes indicate exons or untranslated regions (UTR), black line indicates introns. Created with BioRender.com.

Fig. 4. Relative gene expression of *ATR1* and *ATR2* gene in different mutant and transgenic mutant Arabidopsis lines. a-d) *ATR1* gene expression was analyzed in four different locations using different primer sets ATR1_A, ATR1_B, ATR1_C and ATR1_D. e-i) *ATR2* gene expression was analyzed in five different locations using primer set ATR2_A, ATR2_B, ATR2_C, ATR2_D and ATR2_E. j) Expression of *CYP81A12* with only one primer set. The primers set localization is illustrated in Figure 3. ATR1 and ATR2 gene expression was calculated using Col_0 as control. *CYP81A12* gene expression was calculated using transgenic Arabidopsis CYP81A12 as control. *Significant by Dunnett's Test.

Fig. 5. Arabidopsis lines submitted to increasing doses of penoxsulam, chlorsulfuron (ALS-inhibitors), mesotrione (HPPD-inhibitor) and 2,4-D (auxinic). Each treatment was applied on four plants, represented in the figure. Arabidopsis lines - Wild-type (Col_0), transgenic (*CYP81A12*) and transgenic mutant line *CYP81A12 atr1-b*.

Fig. 6. Herbicide sensitivity of wild type (Col_0), mutant *atr1-a*, *atr1-b*, *atr2-a*, *atr2-b*, transgenic Arabidopsis *CYP81A12* and transgenic mutant Arabidopsis *CYP81A12 atr1-a, CYP81A12 atr1-b*, *CYP81A12 atr2-a, CYP81A12 atr2-b*. Pictures at 14 d-old seedlings growing in ½ MS media containing different herbicide concentrations. a) Plate map, ½ MS control untreated and basta (25mg L^{-1}); b) ½ MS plates with different doses of bensulfuron-methyl, chlorsulfuron, penoxsulam, propoxycarbazone, mesotrione, and bentazon.

Fig. 7. LC-MS/MS analysis of mesotrione metabolites found in different Arabidopsis lines. Arabidopsis lines analyzed were wild-type (Col_0), transgenic *CYP81A12* and transgenic mutants *CYP81A12 atr1-a*, *CYP81A12 atr1-b, CYP81A12 atr2-a, CYP81A12 atr2-b.* A) MS chromatogram 24 h after herbicide application. The retention time for parental mesotrione and hydroxy-mesotrione is 10.3- and 5.9-min, respectively. Peak at retention time 7.7 min is present in all samples, even in untreated plants. B) Peak area of mesotrione and hydroxy-mesotrione at 6, 24, 48 and 96 h after mesotrione application on different Arabidopsis lines. * *p-value* <0.05 by Dunnett's Test.

Fig. 8. Phylogenetic tree of 69 CPR protein sequences from 13 different weed species and others 6 plant species with known CPRs sequences and classification. The phylogenetic tree shows branching of class I (red) and II (different blue shades) CPRs. Different CPR were named as "a", "b" or "c" and the copies were numbered as ".1", ".2" or ".3" for one, two or three copies, respectively. * Different types of CPR have different copy number in the same species. #CPR, number of cytochrome P450 reductases; #cp, number of copies. Am - *Alopecurus myosuroides* (blackgrass), At - *Arabidopsis thaliana* (Arabidopsis), Ap – *Amaranthus palmeri* (Palmer amaranth), Atu – *Amaranthus tuberculatus* (waterhemp), Bs - *Bassia scoparia* (kochia), Bt - *Bromus tectorum* (downy brome), Cc - *Conyza canadensis* (horseweed), Cf - *Chenopodium formosanum* (Djulis), Ec - *Echinochloa crus-galli* (barnyardgrass), Eo - *E. oryzicola* (late watergrass), Eco - *E. colona* (jungle rice), Ei - *Eleusine indica* (goosegrass), Gh - *Gossypium hirsutum* (cotton), Gm - *Glycine max* (soybean), Ip - *Ipomoea purpurea* (common morning-

glory), Mt - *Medicago truncatula* (Barrel medic), Os - *Oryza sativa* (rice), Pa - *Poa annua* (annual bluegrass), Zm – Zea mays (corn).

SUPPORTING INFORMATION

Fig. S1. Two-step PCR for genotyping mutant lines.

Fig. S2. Phenotype analysis of transgenic mutant lines.

Fig. S3. T-DNA insertion genotyping for *ATR1* (a and b) and *ATR2* gene (c and d) in different Arabidopsis mutant lines.

Fig. S4. Pictures of different lines of Arabidopsis submitted to different doses of mesotrione and 2,4-D.

Fig. S5. Pictures of different lines of Arabidopsis submitted to different doses of penoxsulam and chlorsulfuron.

Fig. S6. Different lines of Arabidopsis in response to increasing doses of different herbicides.

Fig. S7. Dose-response of different lines of Arabidopsis growing in different herbicides concentration in ½ MS plates.

Fig. S8. LC-MS/MS analysis of chlorsulfuron metabolites found in different Arabidopsis lines.

Fig. S9. Multiple protein alignment of 67 CPR protein sequences from 13 different weed species and others six plant species with colors indicating conserved residue between sequences.

Table S1. Primers and their characteristics used to genotype by PCR *ATR1* and *ATR2* genes with the T-DNA insertion and *CYP81A12* in F1 or F2 plants.

Table S2. Primers and their characteristics used for gene expression analysis of *ATR1*, *ATR2*, and CYP81A12.

Table S3. PCR genotyping for T-DNA insertion in F2 segregating plants derived from crosses between *atr1* or *atr2* mutants with transgenic Arabidopsis *CYP81A12*.

Table S4. ddPCR results of parental transgenic Arabidopsis *CYP81A12* and four selected F3 plants from crosses A (*CYP81A12* × *atr1-a*), B (*CYP81A12* × *atr1-b*), C (*CYP81A12* × *atr2-a*), and D (*CYP81A12* × *atr2-b*).

Table S5. Genotype analysis of F2 segregating populations derived from crosses between *atr1* and *atr2* Arabidopsis mutants.

Table S6. Weed species and their information about NADPH-cytochrome P450 reductase composition.