Unraveling the Role of P450 Reductase in Herbicide Metabolic Resistance Mechanism

Carlos Alberto Gonsiorkiewicz Rigon[1], Satoshi Iwakami[2], Todd A. Gaines[1], Franck E. Dayan[1]

[1] Colorado State University, Department of Agricultural Biology, Fort Collins, CO 80523

[2] Kyoto University, Graduate School of Agriculture, Kyoto 606-8502, Japan

Corresponding Author: Franck Dayan, franck.dayan@colostate.edu

ORCID:

Rigon: 0000-0001-8565-7383

Iwakami: 0000-0003-4012-9899

Gaines: 0000-0003-1485-7665

Dayan: 0000-0001-6964-2499

SUMMARY

- Plants require cytochrome P450 reductase (CPR) to supply two electrons for cytochrome P450 monooxygenase enzymes (P450) to react with an organic substrate. The transfer of electrons to the P450 active site in the P450 catalytic site relies on a robust and intricate CPR:P450 complex in the endoplasmic reticulum membrane.

- Transgenic Arabidopsis plants carrying *CYP81A12* from *Echinochloa phyllopogon*, which metabolize a broad spectrum of herbicides, were crossed with CPR knockout *atr1* or *atr2* mutant lines. Homozygous gene knockout was confirmed using PCR, and gene copy number of *CYP81A12* was determined using ddPCR. Arabidopsis lines expressing *CYP81A12* in combination with *atr1* or *atr2* knockout were used for herbicide dose-response and metabolism studies.

- Knocking out *ATR1* in transgenic Arabidopsis *CYP81A12* significantly reduced herbicide resistance. Transgenic mutant plants (*CYP81A12 atr1-b*) had a 3.6-, 5.6-, 6.8- and at least 26-fold reduction in resistance to mesotrione, 2,4-D, penoxsulam and chlorsulfuron, respectively, in the dose-response assay. Knockouts of the *ATR2* also decreased herbicide resistance, but to a lower magnitude than *ATR1*. These results were confirmed through the ½ MS medium assay, and herbicide resistance reduction was observed for additional tested herbicides, bensulfuron, propoxycarbazone and bentazon.

- Our findings highlight the importance of CPRs in metabolic herbicide resistance in plants, specifically identifying that *ATR1* is the most important in the CPR:P450 complex in Arabidopsis for herbicide metabolism. The different CPRs found in weeds have potential as target genes to manage herbicide resistance evolution. We further provide an in-depth exploration of the evolutionary implications in weed management arising from the results.

Key Words: CPR, CPR:P450 complex, electron, metabolism, monooxygenase.

INTRODUCTION

Plants are sessile organisms, which remain permanently anchored to their site of

germination. To overcome their lack of mobility, plants have evolved complex detoxification

mechanisms to deal with environmental and biotic stresses. These natural detoxification

processes by chance enable enhanced herbicide metabolic pathways in certain crops, conferring

natural tolerance to herbicides (Jeschke *et al.*, 2019). This strategic approach has yielded

substantial advantages in the commercialization of herbicides intended to eliminate weeds while

safeguarding crops; however, the prolonged application of herbicides in agroecosystems has

selected weeds that exhibit the same mechanisms of resistance (Powles & Yu, 2010). Among

these, the enhanced capacity of herbicide metabolism is one of the most important (Rigon *et al.*,

2020). These selected weeds have an enhanced activity of enzyme families that metabolize the

herbicides before reaching the target site (Gaines *et al.*, 2020). This mechanism of resistance

threatens weed management due to its broad substrate specificity to many herbicides from

different mechanisms of action (Rigon *et al.*, 2020). Metabolic herbicide resistance mechanism is

divided in three phases, known as Phase I – oxidation, Phase II – conjugation, and Phase III –

sequestration and degradation. Phase I is mediated primarily by cytochrome P450

monooxygenase (P450) enzymes. These enzymes are membrane-bound, localized in the

endoplasmic reticulum, and are involved in the biosynthesis of many molecules including

antioxidants, secondary metabolites, and in detoxification of many xenobiotic compounds

(Pandian *et al.*, 2020). P450s are the most important enzymes in phase I of herbicide metabolism

and plants possess a high number of P450 genes that might play a crucial role in herbicide

resistance in metabolic resistant weed species (Dimaano & Iwakami, 2021).

Cytochrome P450 reductase (CPR) is a membrane-bound enzyme localized in the

endoplasmic reticulum that plays a critical role in the biosynthesis of a wide range of compounds

in living organisms, including plants, animals, and bacteria. The structure of P450 reductase is

divided in four domains, including *N*-terminal flavin mononucleotide (FMN)-binding domain, a

connecting domain, an flavin adenine dinucleotide (FAD)-binding domain, and a *C*-terminal

NADPH-binding domain (Wang *et al.*, 1997, Jensen & Møller, 2010) (Fig. **1**). CPR enzymes

arose from of a fusion of different genes encoding FAD-binding, ferredoxin-NADP$^+$, and an

FMN-flavodoxin (Porter & Kasper, 1986). Plants require CPR to supply two electrons for P450

enzymes to conduct each monooxygenase reaction with an organic substrate (Lau & O'Keefe,

1996, Paine *et al.*, 2005). The CPR reduction includes multiple steps for electron transfer. The

first step is reduction from NADPH to FAD, then FMN, and finally to the P450 acceptor (Munro

et al., 2001) (Fig. **1**). The transfer of electrons to the P450 active site in the P450 catalytic cycle

relies on a robust and intricate CPR:P450 complex in the membrane (Mukherjee *et al.*, 2021),

making CPR function an essential component of specialized metabolism.

In contrast to the huge number of P450 genes within plant genomes, for example, 246 in

Arabidopsis thaliana (Arabidopsis) (Paquette *et al.*, 2009), 328 in *Oryza* spp. (rice) (Nelson *et*

al., 2004), 867 in *Echinochloa crus-galli* (barnyardgrass) (Wu *et al.*, 2022), 694 in *E. colona*

(jungle rice) (Wu *et al.*, 2022), and 323 in *Conyza canadensis* (horseweed) (Peng *et al.*, 2014),

CPRs generally have only two or three copies in plant genomes. There are two CPR genes in

Arabidopsis*,* referred to as *Arabidopsis thaliana* cytochrome P450 reductase 1 (*ATR1*), 2 (*ATR2*)

(Urban *et al.*, 1997) and one hypothetical reductase 3 (*ATR3)* that encodes a diflavin reductase

essential for embryogenesis, but that is unable to reduce P450 *in vitro* (Mizutani & Ohta, 1998,

Varadarajan *et al.*, 2010). The isoforms are grouped in two distinct groups with different roles.

Class I is *ATR1*, which is involved in primary and basal metabolism. Class II is *ATR2*, which is

inducible by environmental stimuli such as wounding, pathogen attack, or high light intensity

exposure (Rana *et al.*, 2013, Parage *et al.*, 2016).

In current research efforts, there has been a growing focus on the identification of P450

enzymes responsible for herbicide metabolism in weeds (Rigon *et al.*, 2020). An important

example is the CYP81A12 enzyme derived from a multiple-resistant population of *Echinochloa*

phyllopogon. This enzyme is capable of metabolizing herbicides from diverse modes of action

when expressed heterologously in *E. coli* or Arabidopsis (Iwakami *et al.*, 2014, Dimaano *et al.*,

2020). Considering the dependence of P450 enzymes on P450 reductase for their functionality,

the validation of this interdependence and the identification of pivotal cytochrome P450

reductases could potentially offer a valuable mechanism for weed management and the

mitigation of metabolic herbicide resistance evolution in weeds.

The hypothesis of this study was that *ATR1* and *ATR2* are the main source of electrons for

P450 enzymes and that silencing these genes would decrease the activity of P450s that are

involved in herbicide metabolism. The main objective was to evaluate the function of *ATR1* and

ATR2 in transgenic Arabidopsis over-expressing the P450 *CYP81A12*.

MATERIALS AND METHODS

Plant material

Mutant SALK lines of Arabidopsis were ordered from the Arabidopsis Biological Resource Center (ABRC - https://abrc.osu.edu/). The lines SALK_111558, SALK_208483C have a T-DNA insertion in Arabidopsis cytochrome P450 reductase 1 (*ATR1* - at4g24520), hereafter named as *atr1-a* and *atr1-b*, respectively. The lines SALK_026053 and SALK_152766 have T-DNA insertion in Arabidopsis cytochrome P450 reductase 2 (*ATR2* - at4g30210), hereafter named as *atr2-a* and *atr2-b*.

Seeds from mutants and wild-type Columbia_0 (Col_0) were sown in soil and kept in a growth chamber at 22°C/18°C with a light intensity of 120 µE m^{-2} s^{-1} for 16 h. When seedlings had four true leaves, they were transplanted into individual pots and kept in the growth chamber in the same conditions. All the following experiments used the same methods to grow the Arabidopsis plants.

Characterization and confirmation of T-DNA insertion

PCR assays were performed to identify and confirm the presence of T-DNA insertion in the *ATR1* and *ATR2* genes in the mutant lines and to verify homozygosity. To amplify the wild type (WT) version of *ATR1* or *ATR2* genes, different primer sets named LP (forward) and RP (reverse) spanning the T-DNA insertion specific for each gene and line were used. The primers to identify the T-DNA insertion for all the mutant lines were RP primers, specific for each gene and line, with LBb1.3 (Fig. **S1**). The primer LBb1.3 is a primer designed in the right border of the T-DNA sequence (Fig. **S1**). Primer sequences of LBb1.3, LP and RP for *atr1-a*, *atr1-b* and *atr2-a* were provided by SIGnAL (http://signal.salk.edu/). Primer sets for *atr2-b* were designed using Primer3plus. The primer sequences and their characteristics are described in Tab. **S1**.

Leaf tissue was collected from individual plants of mutants and Col_0 and incubated in 1.2 mL tubes when the plants were at rosette stage. Around 20 plants for each mutant line were genotyped. The tissues were frozen at -20°C until further analyses. Frozen materials were ground using a Qiagen Tissuelyzer and DNA was extracted using a modified hexadecyltrimethylammonium bromide (CTAB) method. DNA quality and concentration were measured using Nanodrop2000.

PCR reactions were performed with Econotaq Plus 2× Master Mix. PCR mixtures

consisted of 12.5 µL of Econotaq Master Mix, 6.5 µL nuclease-free water, and 2.5 µL of forward

and reverse primers (10 µM). Each sample was submitted to two different reactions, called "WT

PCR" and "T-DNA PCR". The former consisted of LP and RP primers to amplify the wild-type

form or heterozygous and the latter was composed of RP and LBb1.3 primers to amplify the T-

DNA insertion if present in the sample. PCR cycling conditions were composed by initial

denaturation of 94°C for 2 min, followed by 35 cycles consisting of denaturation at 94°C for 30

s, annealing at 55°C for "WT PCR" and 48°C for "T-DNA PCR" for 30 s, followed by extension

at 72°C for 70 s, with a final extension time at 72°C for 5 min. Schemes of the primers and PCRs

are represented in Fig. **S1**. The DNA amplification was confirmed by running in 1% agarose gel

electrophoresis and PCR products were sent for Sanger sequencing to confirm the T-DNA

insertion.

Generating transgenic herbicide-resistant Arabidopsis with mutant *atr1* or *atr2* genes

Aiming to generate a transgenic Arabidopsis line with a loss of function in *ATR1* or *ATR2*,

a transgenic Arabidopsis line carrying the P450 *CYP81A12* from *Echinochloa phyllopogon*

(synonym *E. oryzicola*, late watergrass) (Iwakami *et al.*, 2014) was used. The transgenic

Arabidopsis metabolizes five different herbicide modes of action including acetolactate synthase

(ALS) inhibitors (bensulfuron-methyl, pyrazosulfuron-ethyl, chlorsulfuron, azimsulfuron,

imazamox, propoxycarbazone-sodium and penoxsulam), 4-hydroxyphenylpyruvate dioxygenase

(HPPD) inhibitor (mesotrione), protoporphyrinogen oxidase (PPO) inhibitor (pyraclonil),

deoxyxylulose phosphate synthase (DXS) inhibitor (clomazone), and photosystem II (PSII)

inhibitor (bentazon) (Dimaano *et al.*, 2020). This population hereafter is called Arabidopsis

CYP81A12.

Transgenic Arabidopsis *CYP81A12* and mutant lines *atr1-a*, *atr1-b*, *atr2-a*, and *atr2-b*

were grown with the same conditions as indicated before. When plants were in the flowering

stage, around four weeks after transplanting, the mutant lines were crossed with the transgenic

Arabidopsis *CYP81A12*. The crossing consisted of cutting out the stamens from the immature

flowers of Arabidopsis *CYP81A12* and the next day pollinating the carpel with pollen from

different mutant line flowers. The cross was repeated three additional times with new flowers

within an interval of a week. The crossed flowers were labeled, and the seeds were collected

when siliques turned brown. In total, four crosses were performed, cross A – transgenic

CYP81A12 × *atr1-a*, cross B – transgenic *CYP81A12* ×*atr1-b*, cross C – transgenic *CYP81A12*

×*atr2-a*, and cross D – transgenic *CYP81A12* × *atr2-b* (Fig. **2**).

F1 plants were grown, and tissue collected at 2 weeks after transplanting for DNA

extraction. Extracted DNA was used to genotype for T-DNA insertion and presence of

CYP81A12 gene. T-DNA insertion was genotyped as indicated before, using two PCR reactions,

WT PCR and T-DNA PCR. PCR amplification for the whole *CYP81A12* was performed to

genotype for the presence of the gene using genomic DNA. Primer sequences for *CYP81A12* are

listed in Tab. **1**. Plants that were heterozygous for T-DNA insertion and had the presence of

CYP81A12 were selected and self-pollinated. Seeds were collected when mature. F2 plants were

grown with the same conditions as described before, and tissue collected for genotyping.

Genomic DNA was extracted, and T-DNA insertion was genotyped using the same two PCR

reactions as indicated before. It was expected that 25% of the F2 population were homozygous

for T-DNA insertion. These plants were selected to genotype for the presence of *CYP81A12* and

those plants that had the gene were selected. These plants were genotyped for copy number

analysis of *CYP81A12* using QX200 Droplet Digital PCR system (ddPCR) due to its precision

and sensitivity for detection of low copy numbers. ddPCR consisted of a master mix composed

of 10 µL of 2×QX200 EvaGreen® Supermix, 2.5 µL of each primer (5 µM), and 5 µL of

genomic DNA (5 ng µL^{-1}).

Selected samples were processed for droplet generation using QX200™ droplet generator,

followed by PCR reaction with initial cycle of 5 min at 95°C followed by a sequence of 40

cycles starting at 95°C for 30 s and 60°C for 1 min. Droplet reading of PCR products was

conducted in the QX200 Droplet Reader. The *ALS* gene was used as a reference gene containing

only two copies (one copy per haplotype). Primer sequences for *ALS* and *CYP81A12* used for

ddPCR are listed in Tab. **S2**. The samples were analyzed by QuantaSoft AP Software, which

measures the number of positive and negative droplets for each gene in each well. Droplets are

assigned as positive or negative by thresholding based on their fluorescence amplitude. The

software calculates the starting concentration of each target DNA molecule by modeling a

Poisson distribution; the formula used for Poisson modeling is: Copies per droplet = -ln(1-p)

where p= fraction of positive droplets. Concentrations are provided in units of copies per

microliter of input sample with 95% confidence intervals. The ratio of the number of copies per

droplet of *ALS* and *CYP81A12* was calculated. After genotyping, the Arabidopsis transgenic

CYP81A12 carrying mutant *atr1* or *atr2* hereafter are named as *CYP81A12 atr1-a*, *CYP81A12*

atr1-b, *CYP81A12 atr2-a*, and *CYP81A12 atr2-b*.

Generating double knockout *atr1 atr2* Arabidopsis lines

To obtain an Arabidopsis line with a double knockout of *ATR1* and *ATR2* within the same

plant, mutant lines *atr1-a*, *atr1-b*, *atr2-a*, and *atr2-b* were cultivated under the same specified

conditions as previously indicated. Upon reaching the flowering stage, crosses were performed

as described earlier. F1 plants were subjected to genotyping to determine homozygosity for

ATR1 and *ATR2* using the same PCR reactions as previously mentioned. Heterozygous plants

were selected for self-pollination. F2 seeds were then sown in soil, and subsequent plants were

genotyped for *atr1* and *atr2* mutations.

Gene expression analysis in *atr1* and *atr2* mutant lines

Mutant and transgenic mutant lines were grown in a growth chamber with the same

conditions as specified before. Approximately 100 mg of leaf tissue were collected from plants

in the rosette stage, around two weeks after transplanting. Sampling consisted of cutting the

youngest fully expanded leaf for each biological replication, placing in 2 mL tubes, and

incubating immediately in liquid nitrogen and afterwards kept at -80°C. Plants of lines Col_0,

transgenic *CYP81A12*, and transgenic mutants *CYP81A12 atr1-a*, *CYP81A12 atr1-b*, *CYP81A12*

atr2-a, and *CYP81A12 atr2-b* were kept for fresh mass measurement at 28 d after transplant to

analyze any fitness cost in the mutant lines.

RNA extraction was conducted using Direct-zol™ RNA miniprep from Zymo Research

following manufacturer instructions. DNase I treatment was conducted during the RNA

extraction protocol. Quantification and quality control were performed using Nanodrop 2000.

cDNA synthesis was conducted using iScript™ from Zymo Research using 1 μg of RNA. Gene

expression was analyzed on CFX96 Real Time System using SsoAdvanced™ Universal SYBR®

Green Supermix. The reactions consisted of 5 μL of cDNA from samples diluted in Nuclease-

Free Water in 1:20, 2.5 μL of each primer at 5 μM, and 5 μL of SsoAdvanced universal SYBR®

Green supermix ($2\times$). Thermal cycler conditions consisted of an initial cycle of 30 s at 95°C,

followed by a sequence of 35 cycles starting at 95°C for 10 s and 72°C for 30 s. Melt-curve

analysis was performed by adding steps incrementing temperature 65–95°C by 0.5°C.

Primer sets were specifically designed to target various regions of the genes, both upstream

and downstream of the T-DNA insertion site. This systematic design allowed for accurate

analysis of how the gene was affected by the T-DNA insertion, facilitating the identification of

gene knockout. For *ATR1* gene, four sets of primers were designed, called ATR1_A with

forward primers in the 5'UTR and reverse primers in the 1st exon (spanning the T-DNA insertion

position in *atr1-a* line), ATR1_B with forward and reverse primers in the 1st exon, ATR1_C with

the forward primer in exon 4th and reverse primer in exon 5th (spanning the T-DNA insertion

position in *atr1-b* line), and ATR1_D with forward in the 8th exon and reverse in the 9th exon

(Fig. **3**). For *ATR2* gene, five sets of primers were designed, ATR2_A with forward and reverse

primers localized in the 1st exon, ATR2_B with forward primer in the 3th exon and reverse primer

in the 4th exon (spanning the T-DNA insertion position in *atr2-a* line), ATR2_C with forward

primer localized in the 9th and reverse in the 10th exon, ATR2_D with forward primer in the 12th

exon and reverse in the 13th (spanning the T-DNA insertion position in *atr2-b* line), and ATR2_E

with forward and reverse primers designed in the 14th exon. The primer locations in *ATR1* and

ATR2 are represented in Fig. **3a** and **b**, respectively. Primer sequences for gene expression

analysis and their characteristics are listed in Tab. **S2**.

Herbicide sensitivity of Arabidopsis lines in ½ MS medium

Plant growth of different lines of Arabidopsis were evaluated on ½ Murashige and Skoog

(MS) solid medium containing different doses of bensulfuron, chlorsulfuron, penoxsulam,

propoxycarbazone, mesotrione and bentazon. Seeds of Arabidopsis wild type (Col_0), transgenic

Arabidopsis *CYP81A12*, mutant *atr1-a*, *atr1-b*, *atr2-a*, *atr2-b* and transgenic mutant *CYP81A12*

atr1-a, *CYP81A12 atr1-b*, *CYP81A12 atr2-a*, and *CYP81A12 atr2-b* were sterilized by chloride

gas for 7 h by mixing 100 mL of bleach with 3% hydrochloric acid. Technical grade herbicides

were diluted in ethanol (bensulfuron, penoxsulam, propoxycarbazone, bentazon) or methanol

(chlorsulfuron, mesotrione) to mix with ½ MS medium before it solidified. Petri dishes

containing ½ MS with different herbicides doses were divided into 10 sections, where six seeds

of each Arabidopsis line were placed. The plates were incubated in a growth chamber at 25°C,

14-h photoperiod, and 110 µE m^{-2} s^{-1} light intensity until the day of evaluation. The number of

green leaves were counted 14 d after plating, and the data were normalized to percentage of

untreated control.

Whole-plant dose response

Arabidopsis plants of Col_0, transgenic *CYP81A12*, and transgenic mutant *CYP81A12 atr1-a*, *CYP81A12 atr1-b*, *CYP81A12 atr2-a*, and *CYP81A12 atr2-b* were grown with the same conditions as specified before. Plants in rosette stage (14 d after transplanting) were sprayed with increasing doses of chlorsulfuron (Telar ® XP, 750 g Kg^{-1}), penoxsulam (Ricer, 217 g a.i. kg^{-1}), mesotrione (Callisto 480g a.i. L^{-1}), and 2,4-D (Shredder 2,4-D LV4, 455g a.e. L^{-1}). The experimental design was completely randomized in 6×11 with four replications. Factor A was Arabidopsis lines as described above. Factor B was increasing herbicides dose for each herbicide. Chlorsulfuron and penoxsulam doses were 0, 0.1, 0.2, 0.4, 0.8, 1.6, 3.2, 6.4, 12.8, 25.6, and 51.2 g a.i. ha^{-1}; mesotrione doses were 0, 3, 6, 12, 25, 50, 100, 200, and 400 g a.i. ha^{-1}; and 2,4-D doses were 0, 1, 2, 4, 8, 16, 32, 64, and 128 g a.e. ha^{-1}. The shoot fresh mass was measured at 28 d after herbicide treatment by cutting the plant close to the soil surface.

Herbicide metabolism analysis

Seedlings of Arabidopsis lines Col_0, transgenic *CYP81A12*, and transgenic mutant *CYP81A12 atr-a*, *CYP81A12 atr-b*, *CYP81A12 atr2-a*, and *CYP81A12 atr2-b* were transplanted into pots filled with potting soil and grown under the same conditions as previously described. At rosette stage, mesotrione (Calisto ®, 480 g L^{-1} mesotrione, Syngenta, Switzerland) or chlorsulfuron (Telar ® XP, 750 g Kg^{-1} chlorsulfuron, Bayer CropScience, Germany) were applied in an automated spray chamber (Greenhouse Spray Chamber, model Generation IV) using a TJ8002E nozzle, calibrated to deliver 200 L ha^{-1} at a pressure of 280 kPa and speed of 1.2 m s^{-1}. Herbicides were applied at 100 g ha^{-1} for mesotrione and 50 g ha^{-1} for chlorsulfuron. The doses were chosen based on the whole-plant dose-response curve experiment described above as a good dose to discriminate resistant and sensitive plants. After herbicide application the plants were kept in the growth chamber at same conditions and whole plants were collected at 6, 12, 24, 48, and 96 h for mesotrione and adding 144 h and 196 h for plants treated with chlorsulfuron. At each time of tissue collection, plant samples were weighed, washed with 80% acetone for three times, dried with towel paper, placed in 15 mL Falcon tubes, and kept at –70°C ultra-freezer until the extraction. The extraction was conducted by grinding the tissue with a pestle, then 10X of plant mass with 80% methanol and 0.1% formic acid. The solution was homogenized for 30 s using PowerGen 125 (Fisher Scientific) and centrifuged for 10 min at

4,700×g (Sorvall Legend X1R, Thermo Scientific, Waltham, MA). The supernatant was filtered

with Econofltr Nylon 13 mm 0.2 µm (Agilent Technologies, Santa Clara, CA) and injected in the

LC-MS/MS.

LC-MS/MS system consisted of a Nexera X2 UPLC with 2 LC-30AD pumps, a SIL-30AC

MP autosampler, a DGU-20A5 Prominence degasser, a CTO-30A column oven, and SPD-M30A

diode array detector coupled to an 8040 quadrupole mass-spectrometer (Shimadzu Scientific

Instruments, Columbia, MD). For mesotrione, the MS was in positive mode $[M+H]^+$ with MRM

optimized for mesotrione 340>227.95 and set for 100 ms dwell time with a Q1 pre-bias of -

16.0V, a collision energy of -18.0V and a Q3 pre-bias of -16.0V and for hydroxy-mesotrione

356>55>228 and set for 100 ms dwell time with a Q1 pre-bias of -16V, a collision energy of -

18.0V and a Q3 pre-bias of -16.0V. For chlorsulfuron, the MS was in positive mode with MRM

optimized for chlorsulfuron 357.9>141.1>167.05 and set for 100 ms dwell time with a Q1 pre-

bias of -25.0V, a collision energy of -20.0V and a Q3 pre-bias of -15.0V and for hydroxy-

chlorsulfuron 374.0>127.0>182.0 and set for a collision energy of -35.0. The samples were

chromatographed on a 100×4.6 mm Kinetex 2.6 µm biphenyl column (Phenomenex, Torrance,

CA) maintained at 40°C. For both herbicides, solvent A consisted of water with 0.1% formic

acid and solvent B was methanol with 0.1% formic acid. The solvent program started at 50% B

and increased to 95% B in 5 min and maintained at 95% for 4 min. The solvent was returned to

50% B and maintained there for 3 min before the next injection. The flow rate was set at 0.4 mL

min^{-1} and each sample were analyzed with 5 µL injection volumes.

Identification and phylogenetic analysis of CPRs in different weeds

The presence of CPR genes and copy number was searched in several important weed

species. The genomes of *Alopecurus myosuroides* (Cai *et al*., 2023), *Chenopodium formosanum*

(Jarvis *et al.*, 2022), *Bassia scoparia* (Hall *et al.*, 2023), *Eleusine indica* (Zhang *et al.*, 2023),

Bromus tectorum (Revolinski *et al.*, 2023), *Echinochloa crus-galli* (Wu *et al.*, 2022), *E.*

oryzicola (Wu *et al.*, 2022), *E. colona* (Wu *et al.*, 2022), *Ipomoea purpurea* (Gupta *et al.*, 2023)*,*

Poa annua (Robbins *et al*., 2023), and *Conyza canadensis* (Laforest *et al*., 2020) were accessible

through WeedPedia (https://weedpedia.weedgenomics.org) (Montgomery *et al.*, 2023). CPRs

were identified with a blast search using *ATR1* (AT4G24520) or *ATR2* (AT4G30210) against the

genomes or utilized the InterPro ID for NADPH-cytochrome P450 reductase (IPR023208).

Additionally, the genomes of *Amaranthus palmeri* and *A. tuberculatus* were available

(Montgomery *et al.*, 2020), and the COGE web browser (https://genomevolution.org/coge) was

used for blasting purposes. The *CPR* genes in *Amaranthus* species were identified using a

comparative analysis approach. Firstly, *ATR1* and *ATR2* sequences were subjected to a sequence

similarity search against the *Amaranthus hypochondriacus* (grain amaranth) genome (Lightfoot

et al., 2017), which is accessible through the Phytozome database. Subsequently, the identified

sequences were further compared against the Palmer amaranth and common waterhemp

genomes. Other already known plant *CPR* genes were extracted from GeneBank NCBI (Benson

et al., 2013), such as corn (*Zea mays*, CPR2a: NP_001159331.1, CPR2b: NP_001146345.2), rice

(*Oryza sativa*, CPR2a: XP_015651232.1, CPR2b: XP_015650780.1), soybean (*Glycine max*,

CPR1: NP_001236742.2, CPR2: XP_003549436.1), cotton (*Gossypium hirsutum*, CPR1:

ACN54323.1, CPR2: NP_001314398.2), and Barrel medic (*Medicago truncatula*, CPR1:

XP_003602898.1, CPR2: XP_003610109.1). Multiple protein alignment was performed using

ClustalO. Tree construction with Neighbor-joining method was performed using Geneious

Prime® 2023.0.1. The *N*-terminal, transmembrane, FMN binding, FAD binding, and NADP$^+$

binding domains were identified. The transmembrane was identified using DeepTMHMM tool

(Hallgren *et al.*, 2022) (https://dtu.biolib.com/DeepTMHMM).

Statistical analysis

Gene expression analysis of *ATR1* and *ATR2* genes in mutant plants was performed using

ALS as a normalization gene due to its stability under different conditions and herbicide

applications (Chen *et al.*, 2017). Expression analyses were performed utilizing four biological

and two technical replicates for each line. The mean Ct values and the standard deviation were

calculated by treatment. A melt curve analysis confirmed the presence of a single amplified

product for each reaction, based on presence of a single melting temperature consistent across

samples. Relative transcript abundance was calculated using $2^{(-\Delta\Delta Ct)}$ method for each gene and

biotype (Livak & Schmittgen, 2001). The reference population used for the calculation was

Col_0. Dunnett's test ($p < 0.05$) was used to compare the relative expression between Col_0 and

mutant lines.

For the dose-response experiments *in vitro* and *in planta*, the response variables evaluated

were number of green leaves at 14 d after seeds were planted and shoot fresh weight (% of

untreated control) at 28 d after herbicide application (DAA) for the whole-plant dose-response

experiment. Data were converted into percentages relative to untreated treatment for both

experiments. The drc package (Ritz *et al.*, 2015) from statistical software R v.3.5.3 (R Core

Team, 2023) was used for data analysis. The data were tested for the best nonlinear model by

comparing different models by *ANOVA()* and using function *modelFit()* testing for lack-of-fit.

Based on the analysis, the data were adjusted using the three-parameter log-logistic model with

the function *modelFit()* from the drc package (Ritz *et al.*, 2015). For the whole-plant dose

response experiment, the herbicide doses causing 50% reduction in shoot fresh weight (GR_{50})

were determined using the function *summary()* and the resistance index (RI) was calculated by

the function *EDcomp()* by dividing the GR_{50} from transgenic *CYP81A12* or transgenic

CYP81A12 mutants (*atr1* or *atr2*) by the GR_{50} of Col_0. For the dose-response in plates the

calculation was the same, using the GR_{50} obtained by the variable green leaves analyzed. The *x*

and *y* data from the dose-response curves of both experiments were extracted from R using the

function *write.csv()* and plotted with the treatment means by GraphPad Prism version 9 for

Windows (San Diego, California).

The metabolism data obtained by LC-MS/MS were submitted to analysis of variance

(ANOVA) and the means were compared using Dunnett's test ($p < 0.05$). The peak area of the

herbicides and metabolites of all the different Arabidopsis lines were compared to the Col_0 line.

RESULTS

Arabidopsis mutants

SALK lines *atr1-a*, *atr1-b*, and *atr2-a* were segregating for T-DNA insertion and

heterozygous plants were selected to self and seed kept for seed production. *atr2-b* was

homozygous for T-DNA insertion and four plants were self-pollinated to produce seeds. Progeny

plants from *atr1-a*, *atr1-b*, and *atr2-a* were genotyped again and individuals homozygous for the

T-DNA insertion were selected to self and seeds were produced. Homozygous T-DNA insertion

in *ATR1* in *atr1-a* and *atr1-b* lines and homozygous T-DNA insertion in *ATR2* in *atr2-a* and

atr2-b lines were confirmed by PCR. Primers LP and RP for both genes amplified only when

Col_0 DNA was in the reaction. No SALK line amplified the WT version of the respective gene

and only amplified the T-DNA reaction, confirming the homozygosity for T-DNA insertion. The

amplifications were confirmed by electrophoresis in 1% agarose gel (Fig. **S3**), followed by

Sanger sequencing. Based on Sanger sequencing results, the insertion of T-DNA was in the 5'

UTR and in the 4th intron in *atr1-a and atr1-b* mutant lines, respectively, for *ATR1,* and in the 4th

intron and the 12th exon in *atr2-a* and *atr2-b* mutant lines, respectively, for *ATR2* (Fig. **S3**).

Transgenic Arabidopsis *CYP81A12* with *atr1* and *atr2* knockdown

F1 plants from crosses were genotyped for T-DNA insertion. Around 20 plants were

genotyped from each cross and around 90% of the plants were heterozygous for *ATR1* T-DNA

insertion for cross A and B and heterozygous for *ATR2* T-DNA insertion in cross C and D,

indicating success in the crosses. These plants were genotyped for *CYP81A12* presence by PCR

and plants that had the gene were selected for self-pollination to generate F2. F2 plants

segregated 1:2:1 for *ATR1* T-DNA insertion for cross A and B and for *ATR2* T-DNA insertion

for cross C and D (Tab. **S3**). Homozygous mutants were submitted for PCR amplification of

CYP81A12 and when confirmed, samples were submitted for ddPCR. Four, one, three, and five

plants out of 15, 12, 9 and 15 homozygous had *CYP81A12*. These plants were analyzed by

ddPCR. When only one gene copy was identified in comparison to *ALS* gene, the plants were

self-pollinated, and their progenies were analyzed again using ddPCR.

The ddPCR results indicated the same copy number of *CYP81A12* and *ALS* for the parental

transgenic line *CYP81A12* (Tab. **S4**). Plants selected based on previous genotyping (as indicated

above from crosses A, B, C, and D) had a ratio of *ALS* and *CYP81A12* from 0.9 to 1.1,

suggesting two copies of each gene, one copy per haploid genome. These plants were selected

and used for further experiments.

***atr1* and *atr2* double knockouts**

Genotyping analysis of F2 plants obtained from various crosses between *atr1* and *atr2*

mutants demonstrated an absence of double knockout genotypes (Tab. **S5**). A total of 416 F2

plants were genotyped, revealing seven out of the nine possible genotypes. Among the

genotyped plants, 46.2% exhibited heterozygosity for both genes, 5% displayed heterozygosity

for one gene and wildtype for the other, 6.2% exhibited homozygous mutation in one gene and

heterozygosity in the other, while 42.6% exhibited homozygous mutation in one gene and

wildtype in the other. Notably, double wildtype and double mutant genotypes were not observed.

The segregation pattern observed in the F2 generation deviated from the expected Mendelian

segregation for two independent genes, indicating the presence of linked genes. Further

investigation revealed that the *atr1* and *atr2* genes are localized on chromosome four of

Arabidopsis and are approximately two megabases (MBp) apart, providing supporting evidence
that the two genes may be linked.

ATR1 and *ATR2* gene expression in mutant and transgenic mutant lines

Different primer sets were able to detect the loss of function of *ATR1* and *ATR2* genes in
the mutant lines. Mutant *atr1-a* showed a reduced transcription of *ATR1* when the primer set
used was spanning the T-DNA insertion (Fig. **4a**); however, normal expression of *ATR1* gene
was observed when primer set used was downstream of the T-DNA insertion (Fig. **4b-d**),
indicating that the T-DNA insertion in 5' UTR did not decrease *ATR1* expression in *atr1-a* line.
Mutant line *atr1-b*, which had T-DNA insertion in the 4th intron, had significant reduction in
ATR1 transcripts. This transcript disruption was confirmed in the *ATR1* transcript when the
primer set used spanned the T-DNA insertion and a section of downstream sequence (Fig. **4b-d**).
Interestingly, this line had increased transcript abundance when the primer set was upstream of
the T-DNA (Fig. **4a**). These results indicate a possible compensation in the gene expression by
the plant for the disrupted gene.

Mutants for *ATR2* showed similar results. Mutant *atr2-a* had an upregulation of transcript
upstream of the T-DNA insertion in this line (Fig. **4e**); however, this line had significant
reduction of *ATR2* expression downstream of the T-DNA insertion, especially when the set of
primers used was spanning the T-DNA insertion (Fig. **4f-i**). Mutant *atr2-b* had T-DNA insertion
in the 12th exon and had a reduction in the transcription upstream and downstream of the T-DNA
insertion (Fig. **4e-i**). T-DNA insertion in the *ATR1* mutant *atr1-b* and in the *ATR2* mutants *atr2-a*
and *atr2-b* had knockdown of expression of their respective mRNA. *CYP81A12* expression was
analyzed, and all the transgenic mutant lines had the same transcription production as the
transgenic *CYP81A12* (Fig. **4j**).

Knocking down *ATR1* or *ATR2* did not cause obvious effects on plant growth. Transgenic
CYP81A12 had a lower fresh mass with a smaller height of the plants in comparison to Col_0
(**Fig. S2**).

atr1 mutant increases herbicide sensitivity in *CYP81A12* line

Based on whole-plant dose response, transgenic Arabidopsis *CYP81A12* carrying *atr1* and
atr2 were more sensitive to all herbicides tested, including penoxsulam, chlorsulfuron,
mesotrione, and 2,4-D (Fig. **5**, Fig. **S4-7**). The transgenic Arabidopsis *CYP81A12* had a

resistance index of 6.8, 23.6, and 20.9 for penoxsulam, mesotrione, and 2,4-D, respectively (Tab.

1). When *ATR1* was knocked out (*CYP81A12 atr1-b*), the resistance index (RI) decreased to 1.0,

5.7, and 4.2, respectively to penoxsulam, mesotrione, and 2,4-D. The chlorsulfuron doses used in

this study were not able to control the transgenic Arabidopsis *CYP81A12* and *CYP81A12 atr1-a*

(Fig. **5** and Fig. **S4-7**). The resistance to chlorsulfuron of these lines was very high. The lines

CYP81A12 atr2-a and *CYP81A12 atr2-b* showed that the highest doses imparted a slight plant

control, and the model was fitted to the three-parameter model. These lines had a RI of 97.7 and

86.7 for chlorsulfuron (Tab. **1**). The line *CYP81A12 atr1-b* reduced the RI to 3.7 (Tab. **1**).

CYP81A12 atr1-b had the highest decrease in the RI to all herbicides. *ATR2* knockdown lines

(*CYP81A12 atr2-a* and *CYP81A12 atr2-b*) also reduced the RI, but not with the same magnitude

as *atr1-b* (Tab. **1**).

The control of different herbicide doses on Arabidopsis lines grown in ½ MS medium

yielded similar results (Fig. **6** and Tab. **2**). Among the mutants, the *CYP81A12 atr1-b* line

exhibited the most significant reversal of resistance to bensulfuron, penoxsulam, chlorsulfuron,

propoxycarbazone, mesotrione, and bentazon, as evidenced by the highest magnitude of response

(Fig. **6** and Fig. **S7**). Notably, *CYP81A12 atr1-b* displayed symptoms similar to the Col_0 line at

lower herbicide doses (Fig. **6**). In contrast, *CYP81A12 atr1-a*, *CYP81A12 atr2-a*, and *CYP81A12*

atr2-b did not exhibit reduced herbicide resistance compared to *CYP81A12*. Specifically,

CYP81A12 displayed resistance indices (RI) of 111.0, 35.7, 2.1, 4.2, 4.6, and 107.3 for

bensulfuron, chlorsulfuron, penoxsulam, propoxycarbazone, mesotrione, and bentazon,

respectively, while *CYP81A12 atr1-a* demonstrated reduced RIs of 22.3, 10, 0.9, 1.2, 0.9, and 0.9

for the same herbicides, reflecting similar GR_{50} values as Col_0 for penoxsulam,

propoxycarbazone, mesotrione, and bentazon (Tab. **2**).

Reduction of herbicide metabolism

Metabolism of mesotrione and chlorsulfuron was analyzed in the different Arabidopsis

lines after herbicide application. The transgenic Arabidopsis line *CYP81A12* had a smaller peak

area for the parental mesotrione than Col_0 in all time points after herbicide application,

indicating faster herbicide metabolism (Fig. **7a-b**). The transgenic lines *CYP81A12 atr1-a*,

CYP81A12 atr2-a and *CYP81A12 atr2-b* had similar profiles, with smaller peak area of parental

mesotrione than Col_0. The only Arabidopsis line that showed the same peak area as Col_0 was

CYP81A12 atr1-b. Hydroxylation of mesotrione was higher than Col_0 for all lines in all times

points, except for *CYP81A12 atr1-b* (Fig. **7a-b**), which had a very small amount of hydroxy-

mesotrione, highlighting that knockdown of *ATR1* prevents mesotrione metabolism in

CYP81A12 atr1-b. Similar results were observed for chlorsulfuron metabolism (Fig. **S8**).

Transgenic *CYP81A12* had a significantly smaller peak area of parental chlorsulfuron than

Col_0, inferring faster herbicide metabolism. The transgenic mutant line *CYP81A12 atr1-a* was

not different in area peak of parental chlorsulfuron and hydroxy-chlorsulfuron when compared to

Col_0, indicating a loss of the metabolism capacity.

NADPH-cytochrome P450 reductase in weeds

Thirteen different weed species were analyzed, and their corresponding CPRs (cytochrome

P450 reductase) were identified (Fig. **8**). The number of CPR copies varied within each weed

species, ranging from two to three copies (Tab. **S6**). Through the construction of a phylogenetic

tree, CPRs can be categorized into two distinct classes, namely CPR I and CPR II (Fig. **8**). These

findings align with previous research, which suggests that CPR Class I is exclusive to dicot

species, while Class II is present in both monocots and dicots (Jensen & Møller, 2010, Istiandari

et al., 2021).

To differentiate between the CPRs within weed species, they were denoted as "a", "b", or

"c" and individual copies were numbered as ".1", ".2", or ".3" to represent one, two, or three

copies, respectively, in the present study. Kochia (*Bassia scoparia*), common waterhemp,

common morningglory (*Ipomea purpurea*), and Djulis (*Chenopodium formosanum*) have both

CPR 1 and CPR 2 (Fig. **8**). Common morning glory, a tetraploid, possesses two copies of CPR 1

and one copy of CPR 2, whereas the hexaploid Djulis has three copies of CPR 1 and three copies

of CPR 2. The diploids kochia and common waterhemp possess one copy of each CPR class. On

the other hand, the diploids Palmer amaranth and horseweed possess two copies of CPR 2, with

horseweed having an additional third copy, while neither species has any copies of CPR 1. Class

II Palmer amaranth CPRs have 93% identity, while the CPRs of horseweed (Class II) had a 74-

76% identity among their three copies. Djulis, a hexaploid species with 27 chromosomes,

exhibits three copies of each CPR class. Notably, each copy is localized within a different

subgenome (B, C, and D).

Monocot plants exhibit diverse CPR compositions within CPR Class II, forming a distinct

branch. Although their classification into specific classes is not documented in the literature,

various monocot species possess CPR variants within Class II. For instance, blackgrass and

goosegrass each possess three different CPR II variants: Am_CPRa, Am_CPRb, and Am_CPRc

for blackgrass, and Ei_CPRa, Ei_CPRb, and Ei_CPRc for goosegrass. These CPR variants are

categorized into separate branches based on phylogenetic analysis. Similarly, annual bluegrass, a

tetraploid species, possesses three CPR variants, with two copies of each CPR variant present in

subgenomes A or B. Similarly, late watergrass, an allotetraploid species, exhibits two copies of

each of the three CPR variants, distributed across its subgenomes. Jungle rice, an allohexaploid

species, possesses nine CPR variants, with copies of each CPR variant located in subgenomes D,

E, or F. On the other hand, barnyardgrass, also an allohexaploid species, diverges from jungle

rice as it possesses eight CPR variants. Among these variants, two CPRs have three copies each,

while one CPR has two copies. It is noteworthy that the diploid downy brome is the sole

monocot weed species analyzed in this study with only two CPR variants, whereas the remaining

monocot weed species all possess three distinct CPR variants with different copies numbers.

All CPR genes had a well conserved domain (Fig. **S9**). Among the identified CPRs in

weed genomes, three CPRs, namely Eco_CPRc.1, Ec_CPRc.3, and Eo_CPRc.1, stand out with

significantly larger protein sizes compared to the others. These three CPRs have 1520, 1905, and

1948 amino acids, respectively, whereas the remaining CPRs range from 689 to 823 amino acids.

Eo_CPRc.1, Ec_CPRc.3 had a bigger *N*-terminal and additionally Eco_CPRc.1 had a long *C*-

terminal when comparing to the other CPRs. Additionally, these three CPRs exhibit a similarity

ranging from 87% to 94% among themselves. Additionally, there is close genetic relationship

between subgenome D of jungle rice and subgenome C of barnyardgrass (Aoki & Yamaguchi,

2008, Wu *et al.*, 2022). This suggests a common diploid ancestor, explaining the high similarity

observed in CPRs among these three species.

DISCUSSION

Cytochrome P450 reductases (CPR) are a class of enzymes that play a crucial role in the

electron transfer system from NADPH to P450s (Mizutani & Ohta, 1998, Jensen & Møller,

2010). While there are numerous P450 genes, the number of CPR genes is much smaller (Backes

& Kelley, 2003). In Arabidopsis, two CPR genes, class I *ATR1* and class II *ATR2*, have been

identified. *ATR1* is consistently expressed, while *ATR2* is induced in response to stress (Mizutani

& Ohta, 1998, Jensen & Møller, 2010). Since P450s are implicated in metabolic herbicide

resistance in weeds, it is important to understand the specific role of P450 reductases in the

herbicide-metabolizing capabilities of these P450s.

Through the generation of transgenic mutant Arabidopsis lines, i.e., transgenic Arabidopsis

carrying *CYP81A12* from late watergrass able to metabolize a wide range of herbicides and with

knockdown of either the *ATR1* or *ATR2* gene, our observations indicate that *ATR1* plays a

predominant role in facilitating electron transfer from NADPH to *CYP81A12*. Upon knocking

down the *ATR1* gene in the transgenic *CYP81A12* plants (*CYP81A12 atr1-b*), a 3.6-fold and 5.6-

fold reduction in herbicide resistance for 2,4-D and mesotrione were observed, respectively.

Conversely, the *CYP81A12 atr1-a* line, with a T-DNA insertion in the 5' UTR, did not exhibit a

reduction in herbicide resistance due to insufficient gene expression reduction (Fig. **4**).

Knockouts of the *ATR2* gene (*CYP81A12 atr2a* and *atr2-b*) also led to a decrease in herbicide

resistance, although not to the same extent as *ATR1*.

In Arabidopsis, *ATR1* is commonly referred to as *CPR1*, while *ATR2* is known as *CPR2* in

other species. Among the identified CPRs in weeds, it was observed that only CPR class II is

present in grasses. In our current study, the *ATR1* of Arabidopsis was the most crucial in

providing electrons to *CYP81A12*. We hypothesize that different CPRs may exhibit varying

interactions with various P450s, including those involved in herbicide metabolism. A recent

investigation utilizing human CPRs indicated that mutations in the FMN binding domain

significantly enhanced CYP activity. This suggests that amino acid variations within this domain

play a significant role in the diverse interactions between CPRs and P450s (Esteves *et al.*, 2020).

Based on our results, it is evident that the presence of CPR in weed species exhibits variation.

Notably, all examined grasses possess at least three different homologs of CPRs class II, except

for downy brome, which has two. This discrepancy in the number of CPRs among grasses

suggests duplication events that occurred prior to species differentiation within the grass family

(Jensen & Møller, 2010). A detailed analysis of protein sequences revealed substantial

differences primarily in the *N*- and *C*-terminal regions. However, the FMN-binding and FAD-

binding domains in all CPRs displayed a high degree of conservation due to their specialized

functions (Fig. **S9**) and because of that, it is hypothesized that electrostatic interactions are

considered to play a crucial role in guiding the formation of P450:CPR complexes.

The presence of varying numbers of CPRs in different weed species suggests distinct

genetic pressures these weeds have encountered over time, leading to the acquisition of diverse

functions. If herbicide application has selected resistant plants with enhanced herbicide-

metabolizing activity, we hypothesize that herbicide use may have also selected for higher CPR

activity to support the increased activity of P450s by facilitating electron transfer; however, this

aspect needs further investigation. Additionally, in our study, we observed that in Arabidopsis,

ATR2 is unable to complement the electron transfer to *CYP81A12* when *atr1* is knocked out.

However, this may not be the case for other weed species that possess different CPR isoforms.

For instance, jungle rice has three CPR isoforms, each with three copies. These CPRs could

potentially be paralogs with distinct functions, interacting differently with various P450

enzyme families, or they may have complementary functions. Another relevant factor is that

Arabidopsis *ATR1* and *ATR2* are linked, leading to the hypothesis that double knockout plants

would be inviable. However, the absence of double wildtype plants in the F2 generation

dismissed this hypothesis. For most weed species examined in this study, the CPRs are dispersed

across different chromosomes and not exhibit linkage. Nevertheless, our laboratory is actively

pursuing the generation of a double knockout Arabidopsis line using CRISPR/Cas 9 to gain

further insights into the role of *ATR1* and *ATR2* in metabolic resistance mechanisms in weeds.

Research on the role of CPR in the evolution of pesticide resistance has primarily focused

on insects, with limited investigations into the functions of CPRs in plants. In insect studies,

inhibiting CPR in deltamethrin-resistant *Cimex lectularius* increased susceptibility to

deltamethrin, indicating that CPR could be targeted to reduce P450 activity (Zhu *et al.*, 2012).

Similarly, downregulating CPR in resistant *Tetranychus cinnabarinus* (carmine spider mite) and

T. urticae (spider mite) through RNA interference reduced P450 activity and increased

susceptibility to fenpropathrin and various acaricides, respectively (Adesanya *et al.*, 2020). In

plants, which possess one, two, or three CPR isoforms, no research exists regarding the diverse

functions of CPRs for herbicide evolution. For instance, Arabidopsis *ATR2*, which is co-

expressed with lignin biosynthetic genes, showed significant alterations in lignin composition

when mutated, leading to reduced activity of P450 enzymes such as cinnamate 4-hydroxylase, p-

coumarate 3-hydroxylase, and ferulate 5-hydroxylase (Ehlting *et al.*, 2005, Sundin *et al.*, 2014).

The crystal structure of Arabidopsis *ATR2* revealed that mutations in the interflavin electron

transfer region can impact ATR2 activity (Niu *et al.*, 2017), offering opportunities to exploit

these reliable enzymes for reducing P450 activity against herbicide metabolism.

Protein alignments revealed the presence of conserved domains shared across various

CPRs, along with significant variation in the FMN binding domain (Fig. **6**). There were

conserved regions among CPRs in both dicot and monocot weeds, presenting a valuable

opportunity to target and reduce P450 activity. Inhibiting CPR activity, either through chemical

inhibitors or RNA-targeting approaches, offers a potential means to restore herbicide

effectiveness against weeds exhibiting metabolic herbicide resistance. Combining CPR inhibition

with herbicide treatments can serve as a strategy to control metabolic herbicide-resistant weed

populations and, additionally, delay the evolution of metabolic herbicide resistance in sensitive

weed populations.

The study of cytochrome P450 reductases and their interactions with cytochrome P450

enzymes is crucial for understanding metabolic herbicide resistance in weeds. Our findings

highlight the importance of CPRs, particularly *ATR1* in Arabidopsis, in transferring electrons to

P450 involved in herbicide metabolism. The presence of varying numbers of CPRs in different

weed species indicates the diverse genetic pressures these weeds have undergone. Inhibiting

CPR activity could serve as a promising approach to mitigate herbicide resistance, control weed

populations, and prolong the activity of herbicides. Further research on CPRs and their functions

in plants will contribute to our understanding of herbicide resistance evolution and the

development of effective weed management strategies.

ACKNOWLEDGEMENTS

This work was supported by funding from Cotton Inc.

AUTHOR CONTRIBUTIONS

CAGR, FED, and TAG designed the research. CAGR performed experiments. FED,

TAG and SAI contributed experimental materials. CAGR, FED analyzed the data. CAGR, SAI,

FED, and TAG wrote the paper. All authors contributed to editing and approving the paper.

Table 1. Parameters of the logistic equation and resistance index (RI) for shoot fresh weight (%
of untreated control) of wild-type (Col_0), transgenic Arabidopsis *CYP81A12*, transgenic
Arabidopsis mutants *CYP81A12 atr1-a*, *CYP81A12 atr1-b*, *CYP81A12 atr2-a* and *CYP81A12*
atr2-b treated with different doses of penoxsulam, chlorsulfuron, mesotrione and 2,4-D.

| Line | Herbicide | b | GR50 | | | | RI | *p*-value |
			Dose (g ha^{-1})	Lower CI (95%)	Upper CI (95%)	SE		
Col_0		0.5	2.3	0.9	3.7	0.7	-	
CYP81A12		0.7	15.7	6.7	24.6	4.5	6.8	<0.001
CYP81A12 atr1-a	PENO	1.8	17.1	11.2	23.1	3.0	7.4	0.01
CYP81A12 atr1-b		0.7	2.3	1.2	3.5	0.6	1.0	0.9
CYP81A12 atr2-a		2.9	6.7	5.0	8.4	0.8	2.9	0.04
CYP81A12 atr2-b		0.7	3.4	1.5	5.2	0.9	1.5	0.4
Col_0		1.3	0.9	0.7	1.1	0.1	-	-
CYP81A12		-	-	-	-	-	-	-
CYP81A12 atr1-a	CHLO	-	-	-	-	-	-	-
CYP81A12 atr1-b		2.4	3.4	2.5	4.4	0.4	3.7	<0.01
CYP81A12 atr2-a		2.6	87.9	1.8	174.1	42.5	97.7	0.03
CYP81A12 atr2-b		0.6	77.5	1.1	153.9	37.7	86.7	0.04
Col_0		3.7	4.4	3.85	5.1	0.3	-	-
CYP81A12		15	93.6	58.6	128.6	17.7	20.9	<0.001
CYP81A12 atr1-a	MESO	5.5	89.3	78.9	99.7	5.2	19.9	<0.001
CYP81A12 atr1-b		4.3	26.8	23	30.5	1.9	5.7	<0.001
CYP81A12 atr2-a		2.8	41.8	35.5	48.1	3.2	9.3	<0.001
CYP81A12 atr2-b		2.9	59.1	50.3	68.1	4.5	13.2	<0.001
Col_0		0.5	5.9	2.7	9.3	1.6	-	-
CYP81A12		1.2	141.7	96.5	186.5	22.7	23.6	<0.001
CYP81A12 atr1-a	2,4-D	1.6	103.3	73.7	132.8	14.9	17.2	<0.001
CYP81A12 atr1-b		2.9	25.3	20.15	30.4	2.6	4.2	<0.001
CYP81A12 atr2-a		0.8	50.1	29.6	70.5	10.3	8.3	<0.001
CYP81A12 atr2-b		3.8	130.1	108.1	151.9	11.1	21.6	<0.001

638 b: slope; d: upper limit was fixed to 100; GR50: herbicide dose that causes a 50% reduction in
639 shoot fresh weight variable; CI: confidence interval of parameter GR50 ($\alpha = 0.05$); RI: resistance
640 index = GR50 ratio between lines with Col_0 (control). PENO – penoxsulam, CHLO –
641 chlorsulfuron, MESO – mesotrione.

647 **Table 2**. Parameters of the logistic equation and resistance index (RI) for leaf number of

648 Arabidopsis seedlings (% of untreated control) of wild-type, *atr1-a*, *atr1-b*, *atr2-a*, *atr2-b,* and

649 Arabidopsis transgenic *CYP81A12*, Arabidopsis transgenic mutants *CYP81A12 atr1-a*,

650 *CYP81A12 atr1-b*, *CYP81A12 atr2-a*, *CYP81A12 atr2-b* growing in ½ MS medium

651 supplemented with different doses of herbicides.

Lines		b	ED50 dose (nM)	Lower CI (95%)	Upper CI (95%)	SE	RI	*p*-value
Col_0		35.7	9.5	9.1	9.8	0.17	-	-
atr1-a		32.9	3.4	-16.3	23.0	10.0	0.3	0.5
atr1-b		31.5	9.0	8.2	9.9	0.4	0.9	0.4
atr2-a		27.7	9.1	8.3	9.8	0.4	0.9	0.4
atr2-b	BSM	26.6	6.7	-12.9	26.4	9.9	0.8	0.8
CYP81A12		18.3	1055.5	1034.3	1071.8	9.5	111.0	<0.001
CYP81A12 atr1-a		3.6	1368.4	1321.6	1360.9	9.9	141.4	<0.001
CYP81A12 atr1-b		24.9	185.2	192.4	231.7	10.0	22.3	<0.001
CYP81A12 atr2-a		2.0	1547.8	1528.2	1567.5	9.9	163.2	<0.001
CYP81A12 atr2-b		5.6	942.6	923.1	962.2	9.9	99.4	<0.001
Col_0		3.6	3.7	0.73	6.73	1.5	-	-
atr1-a		1.3	1.2	0.8	1.5	0.2	0.3	<0.01
atr1-b		1.7	2.4	1.4	3.3	0.5	0.6	0.03
atr2-a		7.3	3.6	-1.1	8.4	2.4	0.9	0.19
atr2-b	CHL	2.1	1.9	1.4	2.3	0.2	0.5	<0.01
CYP81A12		1.2	132.2	81.9	182.6	25.08	35.7	0.02
CYP81A12 atr1-a		2.7	121.6	93.9	149.2	13.8	32.8	<0.01
CYP81A12 atr1-b		2.3	37.2	27.74	46.66	4.7	10.0	<0.01
CYP81A12 atr2-a		5.4	129.9	8.5	251.4	60.5	35.1	0.1
CYP81A12 atr2-b		0.7	76.1	17.5	134.6	29.1	20.5	0.07
Col_0		3.6	13.8	11.8	15.7	0.95	-	-
atr1-a		1.4	3.8	2.6	4.9	0.5	0.3	<0.001
atr1-b		0.9	3.3	0.5	6.1	1.4	0.2	0.02
atr2-a		1.0	2.7	1.1	4.37	0.8	0.2	<0.001
atr2-b	PEN	3.1	7.0	5.9	8.1	0.5	0.5	<0.001
CYP81A12		8.8	29.6	13.4	45.8	8.0	2.1	<0.001
CYP81A12 atr1-a		10.3	26.9	10.3	43.5	8.21	1.9	<0.01
CYP81A12 atr1-b		11.6	11.9	-8.8	32.8	10.4	0.9	0.25
CYP81A12 atr2-a		10.1	27.8	9.8	45.8	8.9	2.0	<0.01
CYP81A12 atr2-b		8.5	23.7	13.2	34.3	5.2	1.7	<0.001
Col_0		2.4	3.5	2.8	4.0	0.3	0.9	-
atr1-a	PRO	2.1	3.4	1.8	5.1	0.8	0.9	0.9
atr1-b		3.1	3.2	2.5	3.9	0.3	0.9	0.7

Line		b	GR50	CI low	CI high	SE	RI	p
atr2-a		8.1	3.8	0.2	7.4	1.8	1.1	0.9
atr2-b		2.4	2.4	1.6	3.2	0.4	0.8	0.2
CYP81A12		2.4	15.5	12.1	19.0	1.7	4.2	<0.001
CYP81A12 atr1-a		1.9	13.0	10.9	15.1	1.0	3.7	<0.001
CYP81A12 atr1-b		2.1	4.3	3.7	4.9	0.3	1.2	0.2
CYP81A12 atr2-a		2.4	15.7	10.3	21.1	2.6	4.3	<0.001
CYP81A12 atr2-b		1.8	11.1	7.6	14.5	1.7	3.2	<0.001
Col_0		4.5	159.2	131.8	186.7	13.9	-	-
atr1-a		5.4	129.3	106.8	151.8	11.9	0.8	0.3
atr1-b		6.5	160.4	114.8	206.1	23.2	1	0.9
atr2-a		6.3	124.3	84.97	163.6	20.1	0.8	0.1
atr2-b	MES	3.8	117.7	100.4	135.1	8.3	0.7	0.03
CYP81A12		7.9	740.7	604.7	876.78	69.19	4.6	<0.0001
CYP81A12 atr1-a		2.9	480.2	415.1	545.2	33.1	3	<0.0001
CYP81A12 atr1-b		5.1	142.8	112.9	172.72	15.2	0.9	0.4
CYP81A12 atr2-a		6.1	564	492.2	634	36.1	3.5	<0.001
CYP81A12 atr2-b		2.9	585.2	495.7	675.1	45.6	3.6	<0.0001
Col_0		5.4	0.1	0.09	0.11	0.004	-	-
atr1-a		3.9	0.01	0.01	0.02	0.001	0.1	<0.001
atr1-b		3.1	0.03	0.02	0.05	0.007	0.3	<0.001
atr2-a		2.3	0.03	0.02	0.04	0.005	0.3	<0.001
atr2-b	BEN	2.2	0.02	0.01	0.03	0.002	0.2	<0.001
CYP81A12		5	11.24	9.57	12.9	0.82	107.3	<0.001
CYP81A12 atr1-a		3.7	12.84	10.7	14.9	1	132.3	<0.001
CYP81A12 atr1-b		5.22	0.08	-0.07	0.2	0.07	0.9	0.8
CYP81A12 atr2-a		8.43	11.7	3	20.5	4.35	111.6	<0.001
CYP81A12 atr2-b		3.4	12.2	10.2	14.3	1.01	114.6	<0.001

b: slope; d: upper limit was fixed to 100; GR50: herbicide dose that causes a 50% reduction in shoot fresh weight variable; CI: confidence interval of parameter GR50 ($\alpha = 0.05$); RI: resistance index = ED50 ratio between lines with Col_0 (control).

Fig. 1. Process of electron transfer from NADPH-cytochrome P450 reductase to the catalytic site

of cytochrome P450 monooxygenase. Two electrons are sequentially transferred from NADPH

to the heme group of P450 through the cofactors FAD and FMN, each transferred in single-

electron steps. Cytochrome P450, once activated by the electron transfer, catalyzes the oxidative

breakdown of a substrate by inserting one oxygen atom into the chemical compound. Created

with Biorender.

Fig. 2. Schemes representing the Arabidopsis crosses to generate transgenic Arabidopsis
CYP81A12 with loss of function in *atr1* or *atr2*. F1 plants were genotyped for T-DNA insertion
and *CYP81A12* presence. F2 plants were genotyped for homozygous T-DNA insertion and two
copies of *CYP81A12* using ddPCR. M - male, F - female. Created with BioRender.com.

Fig. 3. T-DNA location in *ATR1* (A) and *ATR2* (B) genes in SALK lines and primers localization for real time qPCR gene expression analysis. Yellow triangle locates the T-DNA insertion in the genes, *atr1-a* and *atr2-b* have T-DNA insertion in the 5'UTR and 4th intron, respectively, and *atr2-a* and *atr2-b* have T-DNA insertion in the 3rd and 12th exon, respectively. Boxes indicate exons or untranslated regions (UTR), black line indicates introns. Created with BioRender.com.

Fig. 4. Relative gene expression of *ATR1* and *ATR2* gene in different mutant and transgenic

mutant Arabidopsis lines. a-d) *ATR1* gene expression was analyzed in four different locations

using different primer set ATR1_A, ATR1_B, ATR1_C and ATR1_D. e-i) *ATR2* gene

expression was analyzed in five different locations using primer set ATR2_A, ATR2_B,

ATR2_C, ATR2_D and ATR2_E. j) Expression of *CYP81A12* with only one primer set. The

primers set localization is illustrated in Figure 3. ATR1 and ATR2 gene expression was

calculated using Col_0 as control. *CYP81A12* gene expression was calculated using transgenic

Arabidopsis CYP81A12 as control. *Significant by Dunnett's Test.

Fig. 5. Arabidopsis lines submitted to increasing doses of penoxsulam, chlorsulfuron (ALS-
inhibitors), mesotrione (HPPD-inhibitor) and 2,4-D (auxinic). Each treatment was applied on
four plants, represented in the figure. Arabidopsis lines - Wild-type (Col_0), transgenic
(*CYP81A12*) and transgenic mutant line *CYP81A12 atr1-b.*

Fig. 6. Sensitivity of wild type (Col_0), mutant *atr1-a*, *atr1-b*, *atr2-a*, *atr2-b*, transgenic

Arabidopsis *CYP81A12* and transgenic mutant Arabidopsis *CYP81A12 atr1-a, CYP81A12 atr1-*

b, *CYP81A12 atr2-a, CYP81A12 atr2-b*. Pictures at 14 d-old seedlings growing in ½ MS media

containing different herbicide concentrations. a) Plate map, ½ MS control untreated and basta

(25mg L^{-1}); b) ½ MS plates with different doses of bensulfuron-methyl, chlorsulfuron,

penoxsulam, propoxycarbazone, mesotrione, and bentazon.

Fig. 7. LC-MS/MS analysis of mesotrione metabolites found in different Arabidopsis lines.

Arabidopsis lines analyzed were wild-type (Col_0), transgenic *CYP81A12* and transgenic

mutants *CYP81A12 atr1-a*, *CYP81A12 atr1-b, CYP81A12 atr2-a, CYP81A12 atr2-b.* A) MS

chromatogram 24 h after herbicide application. The retention time for parental mesotrione and

hydroxy-mesotrione is 10.3- and 5.9-min, respectively. Peak at retention time 7.7 min is present

in all samples, even in untreated plants. B) Peak area of mesotrione and hydroxy-mesotrione at 6,

24, 48 and 96 h after mesotrione application on different Arabidopsis lines. * *p-value* <0.05 by

Dunnett's Test.

Fig. 8. Phylogenetic tree of 67 CPR protein sequences from 13 different weed species and others

6 plant species with know CPRs sequences and classification. The phylogenetic tree shows

branching of class I (red) and II (blue) CPRs. Different CPR were named as "a", "b" or "c" and

the copies were numbered as ".1", ".2" or ".3" for one, two or three copies, respectively. *

Different type of CPR has different copy number in the same species. #CPR, number of

cytochrome P450 reductases; #cp, number of copies. Am - *Alopecurus myosuroides* (blackgrass),

At - *Arabidopsis thaliana* (Arabidopsis), Ap – *Amaranthus palmeri* (Palmer amaranth), Atu –

Amaranthus tuberculatus (waterhemp), Bs - *Bassia scoparia* (kochia), Bt - *Bromus tectorum*

(downy brome), Cc - *Conyza canadensis* (horseweed), Cf - *Chenopodium formosanum* (Djulis),

Ec - *Echinochloa crus-galli* (barnyardgrass)*, Eo - E. oryzicola* (late watergrass), Eco - *E. colona*

(jungle rice), Ei - *Eleusine indica* (goosegrass), Gh - *Gossypium hirsutum* (cotton), Gm - *Glycine*

max (soybean), Ip - *Ipomoea purpurea* (common morning-glory), Mt - *Medicago truncatula*

746   (Barrel medic), Os - *Oryza sativa* (rice), Pa - *Poa annua* (annual bluegrass), Zm – Zea mays

747   (corn).

REFERENCES

Adesanya, AW, Cardenas, A, Lavine, MD, Walsh, DB, Lavine, LC & Zhu, F. 2020. RNA interference of NADPH-cytochrome P450 reductase increases susceptibilies to multiple acaricides in *Tetranychus urticae. Pesticide Biochemistry and Physiology,* **165**: 104550.

Aoki, D & Yamaguchi, H. 2008. Genetic relationship between *Echinochloa crus-galli* and *Echinochloa oryzicola* accessions inferred from internal transcribed spacer and chloroplast DNA sequences. *Weed Biology and Management,* **8**: 233-242.

Backes, WL & Kelley, RW. 2003. Organization of multiple cytochrome P450s with NADPH-cytochrome P450 reductase in membranes. *Pharmacology & Therapeutics,* **98**: 221-233.

Benson, DA, Cavanaugh, M, Clark, K, Karsch-Mizrachi, I, Lipman, DJ, Ostell, J & Sayers, EW. 2013. GenBank. *Nucleic Acids Research,* **41**: D36-42.

Cai, L, Comont, D, MacGregor, D, Lowe, C, Beffa, R, Neve, P & Saski, C. 2023. The blackgrass genome reveals patterns of non-parallel evolution of polygenic herbicide resistance. *New Phytologist,* **237**: 1891-1907.

Chen, J, Huang, Z, Huang, H, Wei, S, Liu, Y, Jiang, C, Zhang, J & Zhang, C. 2017. Selection of relatively exact reference genes for gene expression studies in goosegrass (Eleusine indica) under herbicide stress. *Scientific Reports,* **7**: 46494.

Dimaano, NG & Iwakami, S. 2021. Cytochrome P450-mediated herbicide metabolism in plants: current understanding and prospects. *Pest Management Science,* **77**: 22-32.

Dimaano, NG, Yamaguchi, T, Fukunishi, K, Tominaga, T & Iwakami, S. 2020. Functional characterization of cytochrome P450 *CYP81A* subfamily to disclose the pattern of cross-resistance in *Echinochloa phyllopogon. Plant Molecular Biology,* **102**: 403-416.

Ehlting, J, Mattheus, N, Aeschliman, DS, Li, E, Hamberger, B, Cullis, IF, Zhuang, J, Kaneda, M, Mansfield, SD, Samuels, L, et al. 2005. Global transcript profiling of primary stems from *Arabidopsis thaliana* identifies candidate genes for missing links in lignin biosynthesis and transcriptional regulators of fiber differentiation. *The Plant Journal,* **42**: 618-40.

Esteves, F, Campelo, D, Gomes, BC, Urban, P, Bozonnet, S, Lautier, T, Rueff, J, Truan, G & Kranendonk, M. 2020. The role of the FMN-domain of human cytochrome P450 oxidoreductase in its promiscuous interactions with structurally diverse redox partners. *Frontiers in Pharmacology,* **11**.

Gaines, TA, Duke, SO, Morran, S, Rigon, CAG, Tranel, PJ, Küpper, A & Dayan, FE. 2020. Mechanisms of evolved herbicide resistance. *Journal of Biological Chemistry,* **295**: 10307–10330.

Gupta, S, Harkess, A, Soble, A, Van Etten, M, Leebens-Mack, J & Baucom, RS. 2023. Interchromosomal linkage disequilibrium and linked fitness cost loci associated with selection for herbicide resistance. *New Phytol,* **238**: 1263-1277.

Hall, N, Chen, J, Saski, C, Westra, P, Gaines, T & Patterson, E. 2023. FHY3/FAR1 transposable elements generate adaptive genetic variation in the *Bassia scoparia* genome. *bioRxiv*: 2023.05.26.542497.

Hallgren, J, Tsirigos, K, Pedersen, MD, Almagro Armenteros, JJ, Marcatili, P, Nielsen, H, Krogh, A & Winther, O. 2022. DeepTMHMM predicts alpha and beta transmembrane proteins using deep neural networks. bioRxiv.

Istiandari, P, Yasumoto, S, Srisawat, P, Tamura, K, Chikugo, A, Suzuki, H, Seki, H, Fukushima, EO & Muranaka, T. 2021. Comparative analysis of NADPH-cytochrome

819   P450 reductases from legumes for heterologous production of triterpenoids in transgenic
820    *Saccharomyces cerevisiae*. *Frontiers in Plant Science,* **12**: 762546.

821 **Iwakami, S, Endo, M, Saika, H, Okuno, J, Nakamura, N, Yokoyama, M, Watanabe, H,**
822    **Toki, S, Uchino, A & Inamura, T. 2014**. Cytochrome P450 *CYP81A12* and *CYP81A21*
823    are associated with resistance to two acetolactate synthase inhibitors in *Echinochloa*
824    *phyllopogon*. *Plant Physiology,* **165**: 618-629.

825 **Jarvis, DE, Sproul, JS, Navarro-Domínguez, B, Krak, K, Jaggi, K, Huang, YF, Huang, TY,**
826    **Lin, TC, Jellen, EN & Maughan, PJ. 2022**. Chromosome-scale genome assembly of the
827    hexaploid Taiwanese goosefoot "Djulis" (*Chenopodium formosanum*). *Genome Biol*
828    *Evol,* **14**.

829 **Jensen, K & Møller, BL. 2010**. Plant NADPH-cytochrome P450 oxidoreductases.
830    *Phytochemistry,* **71**: 132-41.

831 **Jeschke, P, Witschel, M, Krämer, W & Schirmer, U. 2019**. *Modern Crop Protection*
832    *Compounds,* Wiley-VCH Verlag GmbH & Co. KGaA, Wiley.

833 **Laforest, M, Martin, SL, Bisaillon, K, Soufiane, B, Meloche, S & Page, E. 2020**. A
834    chromosome-scale draft sequence of the Canada fleabane genome. *Pest Manag Sci,* **76**:
835    2158-2169.

836 **Lau, SM & O'Keefe, DP. 1996**. Analysis of herbicide metabolism by monocot microsomal
837    cytochrome P450. *Methods Enzymol,* **272**: 235-42.

838 **Lightfoot, DJ, Jarvis, DE, Ramaraj, T, Lee, R, Jellen, EN & Maughan, PJ. 2017**. Single-
839    molecule sequencing and Hi-C-based proximity-guided assembly of amaranth
840    (*Amaranthus hypochondriacus*) chromosomes provide insights into genome evolution.
841    *BMC Biology,* **15**: 74.

842 **Livak, KJ & Schmittgen, TD. 2001**. Analysis of relative gene expression data using real-time
843    quantitative PCR and the 2(T)(-Delta Delta C) method. *Methods,* **25**: 402-408.

844 **Mizutani, M & Ohta, D. 1998**. Two isoforms of NADPH: Cytochrome P450 reductase in
845    *Arabidopsis thaliana*: gene structure, heterologous expression in insect cells, and
846    differential regulation. *Plant Physiology,* **116**: 357-367.

847 **Montgomery, JS, Giacomini, D, Waithaka, B, Lanz, C, Murphy, BP, Campe, R, Lerchl, J,**
848    **Landes, A, Gatzmann, F, Janssen, A, et al. 2020**. Draft genomes of *Amaranthus*
849    *tuberculatus*, *Amaranthus hybridus*, and *Amaranthus palmeri*. *Genome Biology and*
850    *Evolution,* **12**: 1988-1993.

851 **Montgomery, JS, Morran, S, MacGregor, DR, McElroy, JS, Neve, P, Neto, C, Vila-Aiub,**
852    **MM, Sandoval, MV, Menéndez, AI, Kreiner, JM, et al. 2023**. The International Weed
853    Genomics Consortium: Community Resources for Weed Genomics Research. *bioRxiv*:
854    2023.07.19.549613.

855 **Mukherjee, G, Nandekar, PP & Wade, RC. 2021**. An electron transfer competent structural
856    ensemble of membrane-bound cytochrome P450 1A1 and cytochrome P450
857    oxidoreductase. *Communications Biology,* **4**: 55.

858 **Munro, AW, Noble, MA, Robledo, L, Daff, SN & Chapman, SK. 2001**. Determination of the
859    redox properties of human NADPH-cytochrome P450 reductase. *Biochemistry,* **40**: 1956-
860    1963.

861 **Nelson, DR, Schuler, MA, Paquette, SM, Werck-Reichhart, D & Bak, S. 2004**. Comparative
862    genomics of rice and Arabidopsis. Analysis of 727 cytochrome P450 genes and
863    pseudogenes from a monocot and a dicot. *Plant Physiology,* **135**: 756-772.

Niu, G, Zhao, S, Wang, L, Dong, W, Liu, L & He, Y. 2017. Structure of the *Arabidopsis*
thaliana NADPH-cytochrome P450 reductase 2 (*ATR2*) provides insight into its function.
The FEBS Journal, **284**: 754-765.
Paine, MJI, Scrutton, NS, Munro, AW, Gutierrez, A, Roberts, GCK & Wolf, CR. 2005.
Electron transfer partners of cytochrome. *In:* NEW YORK, N. S. (ed.) *Cytochrome P450,*
structure, mechanism, and biochemistry. New York.
Pandian, BA, Sathishraj, R, Djanaguiraman, M, Prasad, PVV & Jugulam, M. 2020. Role of
cytochrome P450 enzymes in plant stress response. *Antioxidants,* **9**: 454.
Paquette, SM, Jensen, K & Bak, S. 2009. A web-based resource for the Arabidopsis P450,
cytochromes b5, NADPH-cytochrome P450 reductases, and family 1 glycosyltransferases
(http://www.P450.kvl.dk). *Phytochemistry,* **70**: 1940-7.
Parage, C, Foureau, E, Kellner, F, Burlat, V, Mahroug, S, Lanoue, A, Dugé de Bernonville,
T, Londono, MA, Carqueijeiro, I, Oudin, A, et al. 2016. Class II cytochrome P450
reductase governs the biosynthesis of alkaloids. *Plant Physiology,* **172**: 1563-1577.
Peng, YH, Lai, Z, Lane, T, Nageswara-Rao, M, Okada, M, Jasieniuk, M, O'Geen, H, Kim,
RW, Sammons, RD, Rieseberg, LH & Stewart, CN. 2014. De novo genome assembly
of the economically important weed horseweed using integrated data from multiple
sequencing platforms. *Plant Physiology,* **166**: 1241-54.
Porter, TD & Kasper, CB. 1986. NADPH-cytochrome P-450 oxidoreductase: flavin
mononucleotide and flavin adenine dinucleotide domains evolved from different
flavoproteins. *Biochemistry,* **25**: 1682-1687.
Powles, SB & Yu, Q. 2010. Evolution in Action: Plants Resistant to Herbicides. *Annual Review*
of Plant Biology, **61**: 317-347.
R Core Team. 2023. R: A language and environment for statistical computing. R foundation for
statistical computing, Vienna, Austria. https://www.R-project.org/. Vienna, Austria.
Rana, S, Lattoo, SK, Dhar, N, Razdan, S, Bhat, WW, Dhar, RS & Vishwakarma, R. 2013.
NADPH-cytochrome P450 reductase: molecular cloning and functional characterization
of two paralogs from *Withania somnifera* (L.) Dunal. *PLos One,* **8**: 16.
Revolinski, SR, Maughan, PJ, Coleman, CE & Burke, IC. 2023. Preadapted to adapt:
underpinnings of adaptive plasticity revealed by the downy brome genome.
Communications biology, **6**: 326.
Rigon, CAG, Gaines, TA, Küpper, A & Dayan, FE. 2020. Metabolism-based herbicide
resistance, the major threat among the non-target site resistance mechanisms. *Outlooks on*
Pest Management, **31**: 162-168.
Ritz, C, Baty, F, Streibig, JC & Gerhard, D. 2015. Dose-response analysis using R. *PLos One,*
10: 13.
Robbins, MD, Bushman, BS, Huff, DR, Benson, CW, Warnke, SE, Maughan, CA, Jellen,
EN, Johnson, PG & Maughan, PJ. 2023. Chromosome-scale genome assembly and
annotation of allotetraploid annual bluegrass (*Poa annua* L.). *Genome Biol Evol,* **15**.
Sundin, L, Vanholme, R, Geerinck, J, Goeminne, G, Höfer, R, Kim, H, Ralph, J & Boerjan,
W. 2014. Mutation of the unducible *Arabidopsis thaliana* cytochrome P450 alters lignin
composition and improves saccharification. *Plant Physiology,* **166**: 1956-1971.
Urban, P, Mignotte, C, Kazmaier, M, Delorme, F & Pompon, D. 1997. Cloning, yeast
expression, and characterization of the coupling of two distantly related *Arabidopsis*
thaliana NADPH-cytochrome P450 reductases with P450 *CYP73A5*. *Journal of*
Biological Chemistry, **272**: 19176-19186.

Varadarajan, J, Guilleminot, J, Saint-Jore-Dupas, C, Piégu, B, Chabouté, ME, Gomord, V,
Coolbaugh, RC, Devic, M & Delorme, V. 2010. ATR3 encodes a diflavin reductase
essential for Arabidopsis embryo development. *New Phytol,* **187**: 67-82.
Wang, M, Roberts, DL, Paschke, R, Shea, TM, Masters, BSS & Kim, J-JP. 1997. Three-
dimensional structure of NADPH–cytochrome P450 reductase: Prototype for FMN- and
FAD-containing enzymes. *Proceedings of the National Academy of Sciences,* **94**: 8411-
8416.
Wu, D, Shen, E, Jiang, B, Feng, Y, Tang, W, Lao, S, Jia, L, Lin, H-Y, Xie, L, Weng, X, et
al. 2022. Genomic insights into the evolution of *Echinochloa* species as weed and orphan
crop. *Nature Communications,* **13**: 689.
Zhang, C, Johnson, NA, Hall, N, Tian, X, Yu, Q & Patterson, EL. 2023. Subtelomeric 5-
enolpyruvylshikimate-3-phosphate synthase copy number variation confers glyphosate
resistance in *Eleusine indica*. *Nature Communications,* **14**: 4865.
Zhu, F, Sams, S, Moural, T, Haynes, KF, Potter, MF & Palli, SR. 2012. RNA interference of
NADPH-cytochrome P450 reductase results in reduced insecticide resistance in the Bed
bug, *Cimex lectularius*. *Plos One,* **7**: e31037.
926
927

928 **SUPPORTING INFORMATION**
929

930 **Fig. S1.** Two-step PCR for genotyping mutant lines.
931
932 **Fig. S2.** Phenotype analysis of transgenic mutant lines.
933
934 **Fig. S3.** T-DNA insertion genotyping for *ATR1* (a and b) and *ATR2* gene (c and d) in different
935 Arabidopsis mutant lines.
936
937 **Fig. S4.** Pictures of different lines of Arabidopsis submitted to different doses of mesotrione and
938 2,4-D.
939
940 **Fig. S5.** Pictures of different lines of Arabidopsis submitted to different doses of penoxsulam and
941 chlorsulfuron.
942
943 **Fig. S6.** Different lines of Arabidopsis in response to increasing doses of different herbicides.
944
945 **Fig. S7.** Dose-response of different lines of Arabidopsis growing in different herbicides
946 concentration in ½ MS plates.
947
948 **Fig. S8.** LC-MS/MS analysis of chlorsulfuron metabolites found in different Arabidopsis lines.
949

Fig. S9. Multiple protein alignment of 67 CPR protein sequences from 13 different weed species and others six plant species with colors indicating conserved residue between sequences.

Table S1. Primers and their characteristics used to genotype by PCR *ATR1* and *ATR2* genes with the T-DNA insertion and *CYP81A12* in F1 or F2 plants.

Table S2. Primers and their characteristics used for gene expression analysis of *ATR1*, *ATR2*, and CYP81A12.

Table S3. PCR genotyping for T-DNA insertion in F2 segregating plants derived from crosses between *atr1* or *atr2* mutants with transgenic Arabidopsis *CYP81A12*.

Table S4. ddPCR results of parental transgenic Arabidopsis *CYP81A12* and four selected F3 plants from crosses A (*CYP81A12* × *atr1-a*), B (*CYP81A12* × *atr1-b*), C (*CYP81A12* × *atr2-a*), and D (*CYP81A12* × *atr2-b*).

Table S5. Genotype analysis of F2 segregating populations derived from crosses between *atr1* and *atr2* Arabidopsis mutants.

Table S6. Weed species and their information about NADPH-cytochrome P450 reductase composition.