Live-imaging reveals Coordinated Cell Migration and Cardiac Fate Determination

during Mammalian Gastrulation.

Shayma Abukar[1,2], Peter A. Embacher[3], Alessandro Ciccarelli[4], Sunita Varsani-Brown[4], Jamie

A. Dean[3], James Briscoe[4], Kenzo Ivanovitch[1,*]

[1]Developmental Biology of Birth Defects, Institute of Child Health, University College

London, 30 Guilford Street, London WC1N 1EH, UK

[2]Institute of Cardiovascular Science, University College London, Gower Street, London

WC1E 6BT, UK

[3]Department of Medical Physics and Biomedical Engineering, University College London,

Gower St, London WC1E 6BT, UK

[4]The Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK

[*] Corresponding author

E-mail: k.ivanovitch@ucl.ac.uk (K.I.)

Abstract

Heart development involves the specification of distinct sets of cardiac progenitors at various

times and locations during ontogeny. Here, we used live imaging in mice from the initiation

of gastrulation to heart tube formation stages to investigate the origin and migratory paths of

cardiac progenitors. We tracked individual mesodermal cells, reconstructing the lineage tree

of the cells and fates for up to four generations. Our findings revealed independent unipotent

progenitors originating at specific times that exclusively contribute to the left

ventricle/atrioventricular canal (LV/AVC) or atrial myocytes. LV/AVC progeny

differentiated into myocytes early, forming the cardiac crescent, while atrial progenitors

differentiated later and contributed to the venous poles of the heart tube during

morphogenesis. We also identified short-lived bipotent and tripotent mesodermal progenitors

that contribute to a diverse array of mesodermal cell types, illustrating early plasticity during

gastrulation. Sister cells generated by multipotent progenitors dispersed more and adopted

more diverse migratory trajectories within the anterior mesoderm space than those from

unipotent progenitors. Together the data reveal the regulation of individual cell directionality

and cardiac fate allocation within the seemingly unorganised migratory pattern of mesoderm

cells.

Introduction

How cell fate specification and morphogenesis are coordinated in time and space to generate tissues and organs of unique forms and functions is central to developmental biology. This is evident during gastrulation when mesodermal cells acquire diverse cardiac fates and engage in complex cell movements to generate spatial patterns, such as a cohesive cardiac crescent, which transforms into a primitive heart tube.

Clonal analysis establishes lineage relationships between progenitors and their derivatives. In the gastrulating mouse embryo, tracking the derivatives of single progenitors led to the finding that progenitors are assigned to specific anatomical locations in the heart prior to the formation of the heart fields [1-4]. Unipotent *Mesp1+* progenitors are solely destined for the left ventricle (LV) and atria myocardium can be distinguished from unipotent endocardium progenitors [2]. Additional clonal analysis of *Hand1+* progenitors located at the embryonic/extraembryonic boundary in the early gastrulating embryo identified bipotent and tripotent progenitors. These progenitors generated LV/AVC myocytes in addition to pericardium, epicardium, and extraembryonic tissues [1].

One limitation of clonal analysis, however, is that the history of the cells is *deduced* by analysing descendants at the endpoint. It does not allow the identification of the progenitors' initial locations or subsequent migratory paths in the embryo. Single-cell tracking in live-imaging is needed for this and is the most rigorous approach to reconstituting cell lineages, identifying when cardiac progenitors become lineage-restricted during gastrulation and enabling migration analysis [5].

A recent live-imaging analysis uncovered the dynamics of mesodermal cell migration during
mouse gastrulation [6]. This analysis revealed that cells dispersed extensively in the embryo,
with clearly separate movements of daughter cells, suggesting cell identity may not be fixed
but instead influenced by the position of the cells at the end of the migration period.
However, as Dominguez et al. discussed, the motility of mesodermal cells is unlikely to be
completely random. There may be some regulation of directionality of individual cell
migration to ensure progeny migrate to their correct locations and establish spatial patterns,
including the cardiac crescent and distinct LV/AVC and atrial progenitor domains [7-9].
Indeed, a previous migration analysis showed mesodermal cell migrate with directionality
during mouse gastrulation [10]. Thus, early mammalian mesoderm migration may exhibit
some degree of determinism. This is reminiscent of an evolutionarily distinct species, the
ascidian, in which a small number of genealogically related and determined heart progenitors
migrate with predetermined directionally [11]. However, it is not known whether progenitors
adopt more stereotypical migratory trajectories, once committed to specific cardiac fates in
the context of mammalian gastrulation.

Here, through long-term live imaging and single-cell tracking in mice, spanning from the
initiation of gastrulation to the stages of heart tube formation, our goal was to reconstitute the
lineage tree of cells and assess how the migratory paths of cells relate to their eventual
cardiac fate within the seemingly unorganised migration pattern.

Results

Development and characterisation of *cTnnT-2a-eGFP* mice

To track cardiomyocytes in vivo, we developed a knock-in mouse reporter line *cTnnT-2a-eGFP* where the *eGFP* sequence is inserted downstream of the endogenous *cardiac troponin*

T (*cTnnT*) loci. A virus-derived 2a self-cleaving peptide inserted between the *cTnnT* and

eGFP coding sequence ensures co-expression of both cTnnT and eGFP proteins (Fig.1 A)

[12]. The *cTnnT-2a-eGFP* line was maintained as homozygotes. Animals are viable and

indistinguishable from heterozygotes. Whole-mount immunostaining for cTnnT confirmed

specific eGFP expression in cTnnT+ cardiomyocytes at E8 -heart tube stage- and E12.5

(Fig.1 B-C).

We first analysed cardiac differentiation dynamics in real time using multiphoton live-

imaging and the cardiomyocyte *cTnnT-2a-eGPF* reporter line (Fig.1 D). We combined the

cTnnT-2a-eGPF reporter with the *Bre:H2BCerulean* BMP reporter [7], which expressed

cerulean in the lateral plate mesoderm. We found initial sparse GFP positive cells appearing

within the Bre-Cerulean positive lateral plate mesoderm at E7.5 - consistent with initial

sparse cTnnT protein distribution found in the lateral plate mesoderm [13]. This was followed

by the establishment of the cardiac crescent epithelium-like structure and primitive heart tube

(Fig.1 D). We conclude that the *cTnnT-2a-eGFP* reporter faithfully identifies cardiomyocytes

among a population of lateral plate mesodermal cell derivatives.

Establishing long term light-sheet live microscopy for cardiac lineages analysis

Using live-imaging and single-cell tracking in conjunction with the cardiomyocyte *cTnnT-2a-*

eGPF live-reporter, we set out to reconstitute the lineage trees of mesodermal cells and

identify the initial mesodermal progenitors that contribute solely to the heart tube in the

gastrulating mouse embryo (Fig.2 A). To achieve this required culturing and imaging early

mouse embryos from the onset of gastrulation to the heart tube stage (i.e. ~25-35 hours).

We found the Viventis LS1 open-top light-sheet microscope allowed the culture of early

mouse embryos over long periods of embryonic development (>24 hours). Incubation media

was stable and could be exchanged during acquisition. A large media volume (~1ml)

improved embryonic viability for long-term imaging. Embryos cultured from E6.5 and for 35

hours developed normally; a cardiac crescent formed and generated a heart tube

corresponding to E8 embryos.

To permanently label mesodermal cells and their progeny at a density suitable for live cell

tracking, we used an inducible $T^{2a\text{-}cre/ERT2}$ mouse combined with $R26^{tdomato}$ reporter and

administered intermediate doses of tamoxifen (0.002mg/bw) at stages encompassing E5 to

E6.5 [7]. We first administered tamoxifen earlier – (at E5) - and cultured embryos in

tamoxifen-free culture media, from before the start of the gastrulation period and onset of

T/Bra expression in the primitive streak and mesoderm – (at E6) -. In these conditions, no

tdTomato expressing cells could be identified in the intra-embryonic mesoderm over ~11

hours of live-imaging acquisition (Figure 2-supplementary figure 1 A). This confirms that

creERT2 activity in $T^{2a\text{-}cre/ERT2}$ embryos requires T/Bra expression [7]. In the absence of

tamoxifen, rare tdTomato-positive cells were identified in only one embryo (not shown),

confirming that tdTomato widespread mesodermal expression in $T^{2a\text{-}cre/ERT2}$ $R26^{tdTomato}$ embryos

requires tamoxifen.

We generated three light-sheet live-imaging datasets spanning 23 to 35 hours of mouse

embryonic development from gastrulation – (E6.5-E7) - to heart tube stage (Fig.2 B and

Videos 1-3). Embryos were imaged for 1 minute at 2 minutes intervals. Raw data amounts to

5-7 terabytes per experiment, representing up to half a million images. To correct for drift

during acquisition, BigSticher [14] was used to register the datasets in 4D as previously

described [6]. Movies were synchronised according to the timing of appearance of the cardiac

crescent and heart tube inflows (Figure 2-supplementary figure 2 A-K).

Each movie contains up to ~1000 time points, and a small percentage (<1%) of linkage

inaccuracy between cells could lead to lineage misinterpretation that propagates over the

course of the movie. Automated cell tracking methods have seen advancements [15, 16].

However, achieving the level of precision necessary to reconstitute cell lineages remains

challenging. To obtain accurate cell lineages, we manually tracked single cells by visualising

them at successive time points from the beginning to the end of the movie using Massive

Muti-view Tracker (MaMut) [5] (Videos 1-3). We interrupted a track when it was impossible

to identify the same cell across two successive time points unequivocally. A total of 61

mother cells were tracked for up to 4 generations.

We determined the identity of the final daughters based on their location in the heart tube.

The heart tube is formed by an inner endocardial layer ensuring the presence of a circulatory

system, a myocardial layer formed by cTnnT positive cardiomyocytes and derived from the

splanchnic mesoderm, and an outer layer derived from the somatic mesoderm called the

pericardium. We could discriminate these cell types in our live-imaging datasets (Videos 4-

6). Moreover, endocardial cells had distinct spindle-like shapes, displaying protrusions and

transmigrating across the myocardium (Fig. 4D and Video 6, n=4). Cardiomyocytes were

further distinguished by their higher levels of *cTnnT-2a-eGFP* reporter expression (Material

and Methods and Figure 2-supplementary figure 3 A-D). Finally, the locations of the

myocyte *cTnnT-2a-eGFP*+ cells within the heart tube indicated their fates. The heart tube has

an inverted Y shape, and the two arms of the Y -or inflows-, positioned inferiorly, are fated to

become the atria, with the stem of the Y becoming the left ventricle (LV) and atrioventricular

canal (AVC) [17]. In what follows, we describe the lineage trees and timing for mesodermal

progenitors' specification into distinct cardiac lineages.

Distinct mesoderm contributes to the heart tube and inflows myocardium.

We first addressed the timing of LV/AVC and atria myocyte lineage segregation. From our 3

datasets, we identified 29 progenitors contributing to at least one *cTnnT-2a-eGFP*+ myocyte

(Fig. 3A). We analysed their progenies' locations in the heart tube and found they established

a clonal boundary located at the junction between the LV/AVC and atria myocyte

compartments suggesting that atrial and LV/AVC progenitors have distinct mesodermal

origins (Fig.3 Bi-iii and Figure 3-supplementary Figure 1 Ai-vi). Among these 29

progenitors, 14 contributed to the LV/AVC and 13 to the atria. None contributed to clones

spanning the LV/AVC and atria compartments. Two additional progenitors generated *cTnnT-*

2a-eGFP+ cells in deeper z-locations within the heart tube (~ -250 μm).

The LV/AVC and atrial myocyte lineages have distinct temporal origins in the mesoderm.

Early mesoderm contributed to the LV/AVC, while late mesodermal cells generated atrial

myocytes. (Fig. 3Bi). This observation is in line with the previous hypothesis that atrial and

LV/AVC compartments have distinct spatial and temporal origins during gastrulation in the

mouse [7-9].

The LV/AVC progenitors are born first and differentiate into *cTnnT-2a-eGFP*+ myocytes

before other cardiomyocytes. This establishes the initial cardiac crescent within a ~10-hour

period with the first LV/AVC progeny differentiating at ~E6.5 + 15 hours and the last at

~E6.5 + 25-27 hours. Atrial progenitors are born the latest, differentiate the latest - from

~E6.5 + 25-27 hours -, and are recruited to posterior regions during the folding of the cardiac

crescent into the heart tube. This establishes the inflows (Fig.3 D-E and H-I). A subset of

mesoderm progenitors located in the inflow regions did not become *cTnnT-2a-eGFP*+ (n=6);

however, these cells may become positive at later stages.

Myocytes developed concurrently within each lineage (Fig.3 F). The time intervals between

the first and the last daughter to transition into cTnnT-2a-eGFP+ myocytes were 2 hours and

30 minutes on average for LV/AVC lineages, and 2 hours and 35 minutes on average, for

atrial lineages. Notably, the differentiation timing varied among lineages, with some

displaying greater synchrony than others. For instance, in 3 out of 29 lineages, the mother cell

generated LV/AVC cTnnT-2a-eGFP+ myocyte daughters in more than 5 hours. In contrast,

in 8 out of 26 lineages, all daughters transitioned into cTnnT-2a-eGFP+ myocytes in less than

one hour.

Together, the live-imaging analysis shows that the heart tube is established by at least two

sets of independent LV/AVC and atrial myocyte progenitors generated from early and late

mesoderm and differentiating into myocytes at different embryonic stages.

Identification of unipotent, bipotent and tripotent cardiac progenitors.

We next sought to address if mesodermal progenitors contributing to the LV/AVC myocytes

had the potential to generate additional mesodermal lineages. We identified unipotent

LV/AVC progenitors contributing only to *cTnnT-2a-eGFP*+ myocytes (n=9/14) (Fig.3 A, C

and H-H'). However, challenging the notion that mesodermal lineages are locked into

specific fates early [18], we found bipotent progenitors contributing to LV/AVC myocytes

and the endocardium (n=2/14) (Fig.3 A and Fig.4 C, C') as well as the LV/AVC myocytes

and extraembryonic mesoderm (n=1/14) [1]. We also found tripotent progenitors contributing

to LV/AVC myocytes, extra-embryonic mesoderm, and pericardium (n=1/14) and LV/AVC

myocytes, endocardium and extra-embryonic mesoderm (n=1/14) (Fig.3 A and Fig. 4 -

supplementary Figure 1 A, A').

LV/AVC myocyte progenitors were located in the proximal mesoderm, intermingled with

additional progenitors contributing to exclusively non-myocyte *cTnnT-2a-eGFP-* cardiac

lineages (n=8). We identified unipotent pericardial and endocardial progenitors (n=5/8)

(Fig.3 A and Fig.4 A, A') and bipotent progenitors generating endocardial and extra-

embryonic mesodermal cells (n=2/8) (Fig.3 A and Fig.4 B, B') and endocardial and

embryonic mesodermal cells (n=1/8) (Fig.3 A). An additional bipotent progenitor located in

proximity to an LV/AVC myocyte progenitor (cell-cell distance: 29 μm), generated highly

migratory endothelial-like daughter cells located in the embryonic mesoderm and extra-

embryonic mesoderm (Fig.3 A and Fig.4 E, E').

Bipotent progenitors contributing to cardiac and extra-embryonic mesodermal cells were

preferentially located at the extra embryonic/embryonic border (Fig.3 C). However, other

bipotent/tripotent progenitors were intermingled with unipotent LV/AVC progenitors within

the proximal mesoderm, with no clear spatial pattern identifiable. The initial cells' locations

in the early proximal mesoderm seems, therefore, not to correlate with specific mesodermal

fates. However, tracking a greater number of cells would be required to fully address this

question. In line with previous live-imaging analyses [6, 7], the distal mesoderm migrated to

more medial locations (Fig.3 C -dark red cells), known to provide progeny for the right

ventricle, outflow tract and branchiomeric muscles [19-22].

In the late mesoderm, we identified 6 unipotent atrial myocyte progenitors out of the 13

progenitors contributing to the *cTnnT-2a-eGFP+* atria myocytes. Longer tracks

encompassing later stages are required to determine if the remaining progenitors contribute to

atria *cTnnT-2a-eGFP+* myocytes entirely or also to additional lineages. Moreover, we found

additional progenitors contributing exclusively to *cTnnT-2a-eGFP-* daughters located in the

inflow's regions of the heart tube and posterior lateral plate mesoderm (n=16 lineages,

identified as meso GFP- in Fig.3 A). Additional markers or longer movies will be required to

determine the identity of these cells. A subset of these cells had spindle-like shapes and were

identified as endothelial-like cells (n=6, Fig.3 A).

Together, the live imaging analysis of lineages shows that the early mesodermal cells harbour

a previously underappreciated plasticity and diversity of fates during gastrulation [1]. Yet,

their ability to alternate fates seems to be rapidly reduced. 20 out of 35 initial mother cells

contributing to cardiac fates were unipotent, and in all cardiac lineage trees with two or three

fates (n=8), progeny become lineage-restricted early, during migration, before the onset of

cTnnT-2a-eGFP+ expression in the embryo (Fig.3 G). All 6 bipotent progenitors analysed

generated unipotent progenitors after the first cell division; and the two tripotent progenitors

generated unipotent daughters after the first and second cell division (Fig.3 A). These results

are consistent with previous clonal analyses suggesting that early mesodermal progenitors are

rapidly specified into discrete fates after the initiation of gastrulation [2-4].

Migration analysis in lineages reveal hidden patterns of cell migration.

Previous live analysis of cell trajectories during gastrulation revealed apparently chaotic

individual cell movements during migration [6]. Consistent with this analysis, we found that

mesodermal cells dispersed extensively during migration with strong separating movements

between daughter cells (Fig.5 A). We analysed distances between the first two daughters

(coordinates were taken 10 minutes after the first cell division - Timepoint 0 -, and last time

point before the daughters' subsequent cell division - Timepoint 1 -) and granddaughters

(final time point at which all granddaughter cells exist, we only considered branches lasting

at least 4 hours into the cell cycle to allow sufficient cell migration - Timepoint 2 -) in each

lineage. Distances between daughters and granddaughters gradually increased, reaching

considerable distances within a single lineage (up to 365 μm) (Fig.5 A). We noted, however,

that distances were highly heterogenous; a proportion of the progeny generated less

dispersive daughters with separating distances of less than 50 μm between them (11 out of 27

cardiac lineages).

One possibility for the observed heterogeneity in these distances is that the daughters

generated by unipotent progenitors exhibit less dispersive migratory paths than those

generated by bipotent progenitors (Fig.5 B). To test if this correlation was true, we analysed

cell movements in lineages, taking advantage of our lineage analysis from the live-imaging

data.

We first analysed if distances between daughters and granddaughters generated by bipotent

cardiac progenitors were greater than those generated by unipotent cardiac progenitors.

Immediately after the first cell division, we found no distance differences between sisters

generated from unipotent or bipotent progenitors. Distances between sisters with shared

cardiac fates became smaller on average compared to distances between sisters of distinct

cardiac fates (Fig.5 C). A similar difference was found when cell coordinates were sampled

before the onset of *cTnnT-2a-eGFP* expression (Fig.5 D-H). Thus, unipotent cardiac

progenitors generated less dispersed daughter and granddaughter cells than bipotent

progenitors. Distances between sisters could be high in non-cardiac lineages generating only

extra embryonic mesoderm, occasionally reaching distances of over 300 μm (Fig.5 G).

The observation that sister cells sharing the same fate end up in the same position in the

embryo may be due to these cells following the same migratory paths. Alternatively, pairs of

sister cells could independently follow distinct migratory trajectories and, by chance,

converge into similar embryonic territories. To address this question, we analysed if sisters

harbouring a shared cardiac fate migrated in closer proximity than sisters with divergent

fates. We used a dynamic time warping (DTW) algorithm (Fig.5 I and Material and Methods)

that accounts for instances where cells may exhibit similar behaviours but with temporal

shifts [23, 24]. We selected time periods such that sisters' trajectories started and ended at the

same time and calculated the cumulative distances between the two trajectories that yielded

the optimal alignment in each pair of sister cells (i.e. a DTW distance).

On average, unipotent progenitors generated sisters with lower DTW distances than bipotent

progenitors (Fig.5 J-L). We observed four case (out of 20) where a unipotent progeny

generated sisters with a distinct migratory trajectory (log DTW value > 8.5), and two cases

(out of 10) where bipotent progenitors produced sisters with similar migratory paths but

distinct fates (log DTW value < 8.5) (Fig.5 J). This suggests that sisters can occasionally

diverge in their migratory paths yet adopt a similar cardiac fate; or adopt similar paths and

contribute to different fates. (Fig.5 J-L and Figure 5 -supplementary 1 A-B). However, the

majority of sister pairs with the same cardiac fate exhibited notably similar migratory

trajectories (Fig .5 N-S). To rigorously assess this observation, we employed a permutation

test. We created 100.000 permutations by pooling all log DTW distances and randomly

assigning them to either unipotent or bipotent conditions. In each permutation, we computed

the difference in average log DTW values between sister cells with shared fates and those

with distinct fates. This iterative process generated a null distribution, allowing us to test

whether unipotent progenitors are producing sisters with more similar trajectories compared

to bipotent progeny (Fig.5 Supplementary Figure 1C). Our analysis revealed that this was the

case (p-value= 0.00027). Plotting the DTW values over time shows that the migratory

trajectories of sister cells sharing the same fates are similar throughout their entire migratory

periods and any observed similarity is not attributed to systematic smaller DTW-

contributions towards the end of the trajectory (Fig.5 M). Moreover, the DTW values

diverged early in sister cells with distinct fates, indicating that their migratory trajectories are

rapidly distinct at the beginning of their migration when the different fates head off towards

different destinations (Fig.5 M). The results suggest that pairs of sisters with shared fate

maintain closer proximity throughout their entire migration periods, exhibiting more

analogous migratory paths compared to sister cells that adopt distinct fates (Fig.5 M and N-

S).

Discussion

Our findings illustrate the progression of cardiac mesodermal lineages during gastrulation

(Fig.6). Using live-imaging and single cell tracking, we reconstructed cardiac mesodermal

lineages and migratory paths of the cells over extended periods encompassing gastrulation

and heart tube morphogenesis (~35 hours). Culturing embryos in large volumes of media

culture (~1ml) using an open top light sheet microscope was critical to achieve these

experiments.

Our live-imaging suggests that the cardiac crescent (or first heart field -FHF- and juxta-

cardiac field partially overlapping the FHF) [25-28], are predominantly destined to contribute

to the LV/AVC. In contrast, the atria arise at different times during gastrulation [1, 2, 7-9].

These findings suggest that an early segregation of the ventricular and atrial cells has been

conserved during evolution; an early segregation of these progenitor populations was

previously shown at single cell resolution in the zebrafish [29-32]. They also align with *in*

vitro differentiation experiments demonstrating that modulating pathways known to induce

mesoderm can generate molecularly distinct mesoderm favouring the generation of

ventricular or atrial-like cardiomyocytes respectively [33-37].

We found that the early proximal mesoderm harbours multipotent progenitors generating

extraembryonic mesoderm, pericardial, endocardial and endothelial-like cells in addition to

LV/AVC myocytes. These progenitors became rapidly restricted into unique cardiac fates

during migration - prior to the establishment of the cardiac crescent and onset of myocyte

differentiation. The observation of short-lived multipotent cardiac progenitors is consistent

with clonal analysis results of *Hand1+* and *Mesp1+* mesodermal progenitors [1-3].

Previous migration analysis noted opposing cell density and motility gradients in the

mesoderm [6]. According to this model, cells continually exchange neighbours and disperse

widely until their movements gradually diminish, eventually settling in positions and fates as

gastrulation concludes. Our live-imaging analysis builds upon these findings, offering a more

detailed and prolonged evaluation of the migratory paths of cells in relation to their future

fates. The analysis revealed that progenitors contributing to LV/AVC and atrial myocytes

remain as separated cell populations throughout migration, establishing two distinct

progenitor domains in the heart tube without mixing. During ontogeny, the pericardial and

myocardial layers, along with the subjacent plexus of elongated endocardial cells, emerge in

close proximity within the cardiac crescent [38]. The migratory trajectories of progeny

contributing to these three distinct cardiac fates were more deterministic than previously

recognised. Sister cells contributing to the same fate tended to exhibit similar migratory

paths. In the future, it will be crucial to discern whether early mesodermal progenitors exhibit

similar migratory paths because of similarities in their initial internal state and their ability to

interpret environmental cues in a cell-specific manner.

Previous studies in zebrafish indicated that G-protein-coupled receptor signalling, a hallmark

of chemokine signalling, regulates heart progenitor movements during gastrulation [39-41].

Moreover, BMP, Nodal, and FGF morphogen gradients regulate cell migration independently

of cell fate [42-46]. Thus, it seems that mesodermal cells respond to morphogen cues with

precision, providing determinism to the morphogenetic cell behaviours. Simultaneously, they

demonstrate plasticity regarding their final fate [47]. While progenitors are seen giving rise to

only one cardiac cell type, they could potentially generate additional cardiac fates when no

longer constrain by positional cues. We propose that achieving a delicate balance between

determinism and plasticity is essential to ensure robust morphogenesis. This balance enables

cells to follow specific developmental pathways while also maintaining the flexibility needed

to adapt to changing external cues.

Together, our live-imaging analysis of migration and cardiac lineages provide evidence that

some regulation of directionality of cell movements and fate allocation may exist early within

the mesoderm. The findings have broader implications for our understanding of

organogenesis since they address how initial differences between progenitors and signalling

cues may ultimately affect the fate and movements of cells.

Material and Methods

Experimental model and subject details

All animal procedures were performed in accordance with the Animal (Scientific Procedures)

Act 1986 under the UK Home Office project licenses PP8527846 (Crick) and

PP3483414 (UCL) and PIL IA66C8062.

Mouse strains

The $T^{nEGFP-CreERT2/+}$(MGI:5490031) and $Foxa2^{nEGP-CreERT2/+}$ (MGI:5490029) lines were obtained from

Hiroshi Sasaki. The $R26^{\;Tomato\;Ai14/\;Tomato\;Ai14}$ ($Gt(ROSA)26Sor^{tm14(CAGtdTomato)Hze}$ (MGI:3809524), were

obtained from the Jackson Laboratory. The $(no\;gene)^{Tg(BRE:H2B;Turquoise)Jbri}$ BMP reporter line was

generated by the Briscoe laboratory previously [7].

Generation of the cTnnT-2a-eGFP line

The C -terminal tagging of cardiac troponin cTnnt2 with eGFP was generated in the Genetic

Modification Service using CRISPR-Cas9 strategy. This editing was performed by co-

transfection of a Cas9-gRNA vector and a donor vector comprising the T2A self-cleaving

peptide and eGFP into B6N 6.0 embryonic stem cells using Lipofectamine 2000. The donor

vector contained a 786bp insert of T2A-eGFP with 1kb homology arms either side. The guide

sequence used was 5'-TTTCATCTATTTCCAACGCC -3'. Two correctly targeted ESC

clones were microinjected into blastocysts which were then transfered into the uterus of

pseudo pregnant BRAL (C57BL/6 albino) females. Generated chimeras were crossed to

BRAL and F0 chimera offspring were initially screened for the proper integration of T2A -

eGFP by Sanger sequencing. The F0 were again crossed to BRAL to confirm germline

transmission of the mutation and F1 mice were validated by Sanger sequencing.

Immunostaining

Embryos were dissected in 1x Phosphate buffered saline (PBS, Invitrogen), fixed for 4 hours

in 4% PFA at room temperature, washed with 1x PBS-0.1% Triton (0.1% PBS-T) and

permeabilised in 0.5% PBS-T. Embryos were blocked with 1% donkey serum for 1hr,

incubated overnight at 4deg with antibodies diluted in 0.1% PBS-T: mouse anti-cTnnT

(1:250, Thermo Fischer Scientific Systems, MS295P0). Embryos were washed in 0.1% PBS-

T at room-temperature and incubated overnight at 4deg with secondary antibodies coupled to

555 fluorophores (1:200, Molecular Probes). After washing with 0.1% PBS-T, embryos were

then incubated overnight at 4deg with DAPI (1:1000). Embryos were washed in 0.1% PBS-T

and mounted in vectashield. Confocal images were obtained on an inverted Sp5 confocal

microscope with a 20X oil objective (for early E6.5-E7.5 embryos) or a 10X air objective

(0.4 NA) (for E12.5 hearts) at a 2048 × 2048 pixels dimension with a z-step of 1 to 5 μm.

Embryos were systematically imaged throughout from top to bottom. Images were processed

using Fiji software [48].

Embryo culture

Embryos were dissected at E6.5 in a solution of DMEM (Dulbecco's Modified Eagle

Medium- D5921 Sigma-Aldrich) with 10% FBS (Fetal Bovine Serum- A5256701 Thermo

Fisher), 25 mM HEPES-NaOH (pH 7.2), penicillin, and streptomycin. The dissection was

done on a stereoscope microscope with a Tokai Hit thermoplate at 37 °C. Dissection was

completing within 5 minutes and immediately transfer to media culture to preserve the

embryo's developmental potential. Media culture was a mixture of 75% freshly prepared rat

serum (filtered through a 0.2-mm filter) and 25% DMEM (containing 1 mg/ml D-glucose and

pyruvate, without phenol red and L-glutamine- D5921 Sigma-Aldrich), supplemented with

1× glutamax, 100 units/ml penicillin, 100 μg/ml streptomycin, and 11 mM HEPES. The rat

serum was prepared according to established protocols [49, 50], stored at −80 °C, heat-

inactivated at 56 °C for 30 minutes, and filtered through a 0.22-μm filter before use. All

media were equilibrated with a mixture of 5% O_2, 5% CO_2, and 90% N_2, and warmed to

37 °C before adding embryos.

Multiphoton microscopy

To hold embryos in position during time-lapse acquisition, we made bespoke plastic holders

with holes of different diameters (0.3 to 05 mm) to ensure a good fit for the ectoplancental

cone similarly to the traps developed by Nonaka and colleagues [51]. Embryos were mounted

with their anterior side facing up. To avoid evaporation, the medium was covered with

mineral oil (Sigma-Aldrich; M8410). Before starting the time-lapse acquisition, embryos

were precultured for at least 2 hours in the microscopy culture set up. For the acquisition, we

used the multiphoton Olympus FVMPE-RS equipped with a 5% CO_2 incubator and a heating

chamber maintaining 37°C. The objective lens used was a HCX APO L 20x/1.00 W dipping

objective, which allowed a 2-mm working distance for imaging mouse embryos. A

SpectraPhysics MaiTai DeepSee pulsed laser was set at 880 nm and used for one-channel

two-photon imaging. Image settings was: output power: 250 mW, pixel dwell time: 7 μs, line

averaging: two and image dimension: 610 × 610 μm (1,024 × 1,024 pixels). The z step was 6

μm.

Light sheet microscopy

For all light sheet acquisitions, we imaged *TnEGFP-CreERT2/+*; *R26 Tomato Ai14/ Tomato*

Ai14, *cTnnT-2a-GFP+/-* embryos. Tamoxifen (T5648 SIGMA) was dissolved in corn oil.

Oral gavage was performed (0.02 mg/body weight) at indicated embryonic stages. Embryos

were dissected at least 12 hours after. Before the time-lapse acquisition, embryos were

precultured for at least 2 hours in the microscopy culture set-up. To hold embryos in position

during the acquisition, we embedded part of the ectoplacental cone in Matrigel growth factor

reduced phenol red-free (Corning Cat. No 356231) diluted two times with culture medium in

a dedicated open-top FEP sample chamber containing an array of four chambers. Typically, 1

embryo was mounted per well, totalling 4 per experiment. Pixel size was 2 μm x 0.347 x

0.347 (z, x, y). The Viventis LS1 used a single view and dual illumination light sheet.

Detection was done using a Nikon 25X NA 1.1 water immersion objective with final 18.7X

magnification generating a field of view of 800 x 800 μm. Ilumination with the 488 and 561

lasers was sequential. Image acquisition was performed every 2 mins and in stacks totalling

500 μm. Exposure times for GFP and tdTomato detection was 50ms and 100ms. Light sheet

thickness was 3.3 μm. Embryos were incubated at 37°C in 8% CO_2 with humidification

throughout.

Cell tracking

The original images underwent conversion into HDF5/XML file formats and registration

using BigSticher [14], following the methodology outlined in prior work [6]. Subsequently,

these processed images were imported into the MaMuT Fiji plugins for manual cell tracking

[5]. The MaMut viewer windows allowed visualization of each tracked cell at any brightness,

scale, and rotation parameters. Cell identification is initiated at the initial time point, with

manual tracking occurring every five-time points, except during mitosis. During mitosis,

tracking was performed at every time-point to capture daughter cell separation accurately.

The TrackScheme lineage browser was employed to visualize the reconstructed cell lineage

tree. Tracked cells were depicted as nodes interconnected by edges, and cell divisions were

illustrated as split branches. For subsequent analysis, the tracked cells' spots, tracks, cell

division time and GFP intensities were exported in csv format.

Threshold analysis

GFP values were normalized based on the highest GFP value within each movie's tracking

data. To discern GFP-positive cells from the background, we implemented the following

approach: GFP background intensities were measured at regular intervals throughout each

movie and linear interpolation was applied between time points to establish a background

value for every time-point. The threshold for distinguishing GFP-positive and GFP-negative

cells was determined by using a multiple of the mean background. The threshold was set to

the lowest multiplier, ensuring that every endocardial cell in each movie were classified as

GFP-negative.

Lineage analysis

Each reconstructed lineage tree starts with a mother cell. The mother divides giving rise to

the two daughter cells D1 and D2. Then D1 divides giving rise to daughters D11 an D12.

D11 divides giving rise to daughters D111 and D112. Lineage trees were generated in R

version 4.3.2 using the 'ggtree' version 3.10.0 package [52] with Newick format and imported

into R through the 'read.tree' function from the 'ape' package version 5.7.1. The branch

lengths of the trees are proportional to the cell cycle length. The trees were coloured using the

'scale_colour_gradient' function in the 'ggtree' package, which imparts a colour gradient

based on the normalized GFP intensities of the cells or paintsubtree function in phytools

version 2.03 package [53]in R version 4.3.2 (https://www.R-project.org).

Homotypic and Heterotypic Distances Calculation

Trajectories were reconstructed in 3D using matplotlib version 3.7.2 package in Python

3.11.53. We computed Euclidean distances to quantify the spatial relationships among cells,

categorizing them into 'homotypic distances' when comparing cells of the same fate and

'heterotypic distances' when comparing cells of different fates. Progenitors' coordinates were

sampled before cTnnT-2a-eGFP signal was observed. We only considered branches lasting at

least 4 hours into the cell cycle to allow sufficient cell migration. To analyse the evolution of

homotypic and heterotypic distances over time, we focused on daughters' coordinates at three

distinct time points: T0, T1, and T2. T0: 20 minutes after the mother's initial cell division.

T1: last time point before the daughters' subsequent cell division. T2: final time point at

which all granddaughter cells exist. If no third generation is present, we only considered

branches lasting at least 4 hours into the cell cycle to allow sufficient cell migration.

Midpoint Calculations

The midpoints between D1's and D2's progeny were first determined by averaging their

coordinates. Progenitors' coordinates were sampled before cTnnT-2a-eGFP signal was

observed. Subsequently, a midpoint distance between D1 and D2's progeny's midpoints was

calculated.

Dynamic time warping (DTW)

We analyzed migratory trajectories within cardiac lineages by comparing the paths of sister

cells from the time immediately following the division of their mother cell to the final time

point before the first sister cell division. Trajectory similarity was assessed using the 'dtw-

python' (version 1.3.0) Python package, measuring the dynamic time warping distance

(DTW) between the cells. We opted for DTW distance over Euclidean distance, as DTW

accommodates cells with spatially similar yet temporally nonsynchronous trajectories [52,

53]. We applied the 'symmetricP1' step pattern, a slope-constrained step pattern [22, 23]. To

determine whether DTW distances between sister cells, whether sharing the same fate or

having distinct fates, exhibited significant differences, we performed a permutation test using

the 'scipy' Python package (Version 1.11.1) [54]. This involved 100,000 resamplings and the

calculation of differences between log mean DTW distances for unipotent and bipotent cells

in each permutation. The resulting p-value was computed as the proportion of permuted

values greater than or equal to the observed values + 1, divided by the total number of

permutations + 1. Our null hypothesis assumed no difference in log mean DTW distances

between sister cells generated by unipotent and bipotent progenitors, while the alternative

hypothesis suggested a significant difference.

Figure Legends

Figure 1. Development and Characterisation of the cTnnT-2a-GFP line. (A) Schematics illustration of a CRISPR-cas9-mediated strategy to insert a 2a-GFP cassette in front of the stop codon into the cTnnT2 gene. **(B-C)** Representative immunofluorescence staining for cTnnT2 in hearts at different stages in cTnnT-2a-GFP mice. Scale bar at E8: 100 μm. Scale bar at E12.5: 200 μm **(D)** Image sequence from time-lapse video of an *TnGPF-CreERT2/+; Bre:H2B-Turquoise; cTnnT-2a-GFP* embryo. Scale bar: 100 μm. IF: Inflows, HT: heart tube, OFT: outflow tract, LV: left ventricle, RV: right ventricle, RA: right atria, LA: left atria, YS: Yolk Sac, Ant: Anterior, Post: posterior,

Figure 2. **Long-term live-imaging from gastrulation to heart tube formation**. **(A)** Schematic of the cell tracking procedure. Tracks carry information on (i) reporter expression, (ii) lineage relationships and (iii) trajectories. **(B)** Image sequences from three time-lapse videos of *TnGPF-CreERT2/+; R26tdtomato+/-; cTnnT-2a-GFP* embryos resulting from the administration of tamoxifen (0.02mg/body weight) at indicated times. LV/AVC: left ventricle and atrio ventricular canal. Scale bar: 100 μm

Figure 3. Independent LV/AVC and Atria progenitors contribute to distinct regions of the heart tube. (A) Reconstruction of mesodermal cell lineages. Cell types are indicated at the endpoints. Lineages are coloured according to their normalised GFP intensities. **(B)** Only the progenitors contributing to at least one cTnnT-2a-GFP+ progenitors are represented. (i) Fate map depicting early (green) and late (magenta) mesoderm contributing to distinct regions of the heart tube (ii) Each colour represents a distinct clone. (iii) cTnnT-2a-GFP+ cells are shown in red. **(C)** Fate map showing all the early mesodermal progenitors coloured coded by fate. **(D)** Birth date of progenitors contributing to cTnnT-2a-GFP+ LV/AVC and

cTnnT-2a-GFP+ Atria myocytes. The cells already present at the beginning of the movie are

highlighted in a grey-dotted box. **(E)** Timepoints at which LV/AVC and atrial progenitors'

GFP intensities are above a certain threshold value and defined as GFP-positive cells

(Material and Methods) For LV/AVC progenitors, the mean differentiation time is 19.0 hours

+/-2.2 and for atrial progenitors it is 30.2 hours +/-2.4; mean +/- SD. **(F)** Time between the

first and last progenitor to become cTnnT-2a-GFP+ in every lineage tree contributing to

myocytes. Mean time for LV/AVC is 2.5 hours +/-1.7 (SD) and 2.3 hours +/-2.6 -SD), p=

0.44. The statistical analysis was done using a Mann-Whitney test. **(G)** Birth date of

unipotent progenitors for LV/AVC and atrial myocytes, endocardium and pericardium. The

cells already present at the beginning of the movie are highlighted in a grey-dotted box. **(H-**

I) Image sequences from time-lapse video of *TnGPF-CreERT2/+; R26tdtomato+/-; cTnnT-*

2a-GFP embryos depicting an LV/AVC (H) and Atrial (I) progenitors. The corresponding

lineage tree coloured by the normalised GFP intensity of the track cells are shown in H' and

I'. Arrows point to the cells in the lineage tree in H' and I'. Scale bar: 100 μm

Figure 4. Lineage analysis in time-lapse movies demonstrates unipotent and bipotent

mesodermal progenitors exist. (A-C and E) Image sequences from time-lapse videos

of *TnGPF-CreERT2/+; R26tdtomato+/-; cTnnT-2a-GFP* embryos depicting an unipotent

pericardium (A) and an Endocardial and Extra-embryonic mesoderm (B), an endocardial and

myocyte (C) and endothelial-like and Extra-embryonic bipotent progenitors (E). The

corresponding lineage tree coloured by the normalised GFP intensity of the track cell are

shown in A'-C' and E'. Arrows point to the cells represented in the lineage tree in A'-C' and

E'. **(D)** Image sequences from time-lapse videos of *TnGPF-CreERT2/+; R26tdtomato+/-;*

cTnnT-2a-GFP embryos depicting an endocardial progenitor migrating inside the forming

heart tube. Scale bar: 100 μm

Figure 5. Cell migration analysis in lineages reveals hidden patterns. (A) Dispersion

analysis over time shows the distances between all daughters and granddaughters in all the

cardiac lineages at time points T0, T1, and T2, as explained in the results and material

method sections. The average distances at T0 (n=28), T1 (n=28) and T2 (n=136) are 15.2 μm

+/- 3.1; 39.08 μm +/- 46.26 and 60.25 μm +/- 67.75, mean +/- SD; T0 vs T1, p=0.027; T1 vs

T2, p=0.040. Results were statistically analysed using an uncorrected Dunn's test following a

Kruskal-Wallis test (P<0.001). **(B)** Schematic diagram of the hypothesis. If cells migrate

randomly, then the unipotent and bipotent progeny will generate equally dispersed daughters

by the end of the migration. Alternatively, unipotent progeny will generate less dispersed

daughter cells if heterogeneity in migratory trajectories exist. Midpoint distance depicts the

Euclidian distance between the D1's daughters' midpoint and the D2's daughters' midpoint.

Homotypic and heterotypic distances are between cells of the same or different fate. **(C)**

Dispersion analysis comparing homotypic and heterotypic distances between all daughters

and granddaughters in all the cardiac lineages at time points T0, T1, and T2. The average

homotypic distances at T0 (n=20), T1 (n=20) and T2 (n=105) are 14.45 μm +/- 3.1, 24.67 μm

+/-11.69 and 32.90 μm +/- 19.75, mean +/- SD. The average heterotypic distances at T0

(n=11), T1 (n=11) and T2 (n=37) are 16.11 μm +/-2.86, 63.40 μm +/- 67.13 and 104.4 μm +/-

89.49, mean +/- SD. Results were statistically analysed using an uncorrected Dunn's test

following a Kruskal-Wallis test (P<0.001). Timepoint 0: 9 hours and 40 minutes on average

+/- 7.4; Timepoint 1: 17 hours and 30 minutes on average +/- 8.4; Timepoint 2: 22 hours and

43 minutes on average +/- 6; +/- SD). **(D)** Dispersion analysis comparing homotypic and

heterotypic distances between all daughters and granddaughters for all the cardiac lineages.

Coordinates were sampled before the onset of cTnnT-2a-GFP expression in the embryos. The

average homotypic and heterotypic distances are 34.82 μm +/-20.38 (n=110) and 137.2 μm

+/-118.4 (n=70); mean +/- SD. Results were statistically analysed using a Mann-Whitney

test. Timepoints on average were 22 hours +/- 5, +/-SD (**E-F**) Homotypic distances in all

lineages generating at least one cTnnT-2a-GFP+ LV/AVC (E) or cTnnT-2a-GFP+ atria (F)

myocytes. (**G**) Homotypic distances per fate and in lineages developing in only one fate.

Coordinates were sampled before the onset of cTnnT-2a-GFP expression in the embryos. The

average homotypic distances for LV/AVC, Atria, Pericardium, Endocardium and ExMeso

fates were 33.37 μm +/-20.89 (n=90), 27.30 μm +/- 29.40 (n=34), 36.67 μm +/- 15.14 (n=24),

41.66 μm +/- 21.31 (n=18) and 79.09 μm +/- 94.06 (n=38), mean +/- SD. Results were

statistically analysed using a Kruskal-Wallis and Dunn's multiple comparisons tests. (**H**)

Midpoint analysis comparing distances between D1 and D2's daughters' midpoints in cases

where D1 and D2 generate the same fate (unipotent) or different fates (bipotent). Analysis

was done for all the cardiac lineages. Coordinates were sampled before the onset of cTnnT-

2a-GFP expression in the embryos. The average midpoints distances for unipotent and

bipotent progenitors were 32.89 μm +/-20.03 (n=20) and 97.35 μm +/-88.49 (n=16), mean +/-

SD. (**I**) Dynamic time warping allows quantifying similarities between two migration

paths. (**J**) Dynamic time warping score comparing D1 and D2 cell tracks up to their next cell

division in cases where D1 and D2 have the same cardiac fate (unipotent) or distinct fates

(bipotent). Results were statistically analysed using a Mann- Whitney test. (**K-L**) log DTW

score for all lineages generating only cTnnT-2a-GFP+ LV/AVC (K) or cTnnT-2a-GFP+ atria

(L) myocytes. (**M**) Average DTW values over time for D1 and D2 daughters generated by

unipotent (red) and bipotent (bleu) progenitors. Shaded blue and red represent the standard

error. At step 7 (indicated by an orange arrow), the DTW values for daughters of shared and

distinct fates are statistically distinct. Results were statistically analysed using a Mann-

Whitney test. (**N-S**) Examples of trajectories and corresponding midpoint distance calculation

(O'-T') and lineage trees (O''-T''). The red lines in lineage trees indicate the time points at

which the cells' coordinates were sampled for the midpoint analysis. The red lines in the

trajectory diagram represents the distances between D1 and D2's daughters' midpoints.

Figure 6. Working Model of early cardiac development. (A) Early proximal mesodermal

cells are initially highly plastic (or multi-potent) but rapidly become committed to specific

cardiac fates as they initiate migration towards specific embryonic regions in response to

environmental cues. LV/AVC and Atrial progenitors are generated from early and late

mesoderm, respectively. LV/AVC progenitors differentiate first into myocytes and establish

the cardiac crescent. Atria progenitors differentiate later and generate the heart tube's

inflows. (**B**) Pair of sisters sharing the same fate have more identical migration paths than

sisters with distinct fates. PS: primitive streak. LV/AVC: left ventricle/atrioventricular canal.

ExM: Extra-embryonic mesoderm.

Figure 2-supplementary figure 1. Tamoxifen induces tdTomato expression in *TnGPF-*

***CreERT2/+; R26tdtomato+/-; cTnnT-2a-GFP* embryos at the correct embryonic stages.**

(**A**) Image sequence *TnGPF-CreERT2/+; R26tdtomato+/-; cTnnT-2a-GFP* embryos resulting

from the administration of tamoxifen (0.02mg/bw) at E5.5, and culture from E6 to E7 in the

absence of tamoxifen. Arrows point to tdTomato-positive cells present in the Extra-

embryonic mesoderm. PS: primitive streak. Scale bar: 100 μm

Figure 2-supplementary figure 2. GFP signal Quantification. (A, D and G)

Quantification of the cTnnT-2a-GFP signal in the cardiac crescent and inflows over time. (**B-**

C, E-J, H-F) We quantified cTnnT-2a-GFP expression by drawing "circles" in the cardiac

crescent (B, E and H) and inflows (C, J and F) at different time points. We synchronised the

three movies according to their onset of cTnnT-2a-GFP expression in the initial cardiac

crescent and subsequent heart tube inflows generated during morphogenesis.

Figure 2-supplementary figure 3. Threshold analysis (A-C) Green lines represent the

threshold set to discern GFP-positive cells from the background**. (D)** Red round circles

represent the areas in which the background GFP intensity was measured.

Figure 3-supplementary figure 1. Independent progenitors contribute to the LV/AVC

and Atria. (A) Only the progenitors contributing to at least one cTnnT-2a-GFP+ progenitors

are represented. In i-iii is movie 2 and in iv-xi is movie 3. (i and iv) Fate map depicting early

(green) and late (magenta) mesoderm contributing to distinct regions of the heart tube. (ii and

v) Each colour represents a distinct clone. (iii and vi) cTnnT-2a-GFP+ cells are shown in red.

Scale bar: 100 μm

Figure 4-supplementary figure 1. Tripotent mesodermal progenitors exist. (A) Image

sequences from time-lapse video of *TnGPF-CreERT2/+; R26tdtomato+/-; cTnnT-2a-*

GFP embryos depicting a progenitor contributing to three cell types. The corresponding

lineage tree coloured by the normalised GFP intensity of the track cells are shown in A'.

Arrows point to the cells represented in the lineage tree in A'. Scale bar: 100 μm

Figure 5-supplementary figure 1. Cardiac progenitor trajectories analysis. (A-B)

Examples of trajectories and corresponding midpoint distance calculation (A'-A') and lineage

trees (A''-B'''). The red lines in lineage trees indicate the time points at which the cells'

coordinates were sampled for the midpoint analysis. The red lines in the trajectory diagram

represents the distances between D1 and D2 daughters' midpoints. (**C**) Histogram showing

the distribution of the permutated tests obtained by calculating the log mean difference

between randomly shuffled unipotent and bipotent dynamic time warping (DTW) distances.

The data were binned at 30μm intervals. A red dashed line marks the observed log mean

difference.

Video 1-3: Live-imaging from gastrulation to heart tube stage. Embryos are $T^{nGPF\text{-}CreERT2/+}$;

$R26^{tdTomatofl/+}$; cTnnT-2a-GFP. Pregnant mice received tamoxifen at around E6 (Videos 1-2)

and E6.5 (Video 3). Images are Maximum projection of 250 sections acquired every 2 μm.

Interval between frames: 2 minutes. Related to Fig.2 B.

Video 4: Example of a myocyte progenitor visualised along the xy and yz axis. From video 2.

Scale bar: 50 μm.

Video 5: Example of an endocardial progenitor visualised along the xy and yz axis. From

video 1. Scale bar: 50 μm.

Video 6: Example of a pericardial progenitor visualised along the xy and yz axis. From video

3. Scale bar: 50 μm.

Acknowledgments

We thank Dr Martin Dominguez for their initial help with BigSticher and MaMut. We thank

Dr Rosie Marshall and Prof. Andrew Copp for their help with rat serum preparation. We also

thank the Light Microscopy facilities and Biological Research Facilities at the Francis Crick

Institute and UCL for their help with imaging, image analysis and transgenic colonies. We

thank Drs Nancy Stathopoulou and Florencia Cavodeassi for comments on the manuscript.

We thank all members of the Ivanovitch and Alexandre labs for help, advice, reagents, and

feedback.

Author Contribution

S.A. Conceptualization, Formal analysis, Investigation, Visualization, Writing – review &

editing, P.A.E. Methodology, Writing – review & editing, A.C. Methodology, Writing –

review & editing, S.V.B. Methodology, J.A.D. Methodology, Supervision, Writing – review

662 & editing, J.B. Funding acquisition, Supervision, Writing – review & editing, K.I. Funding

663 acquisition, Project Administration, Conceptualization, Formal analysis, Investigation,

664 Visualization, Supervision, Writing – original draft, Writing – review & editing.

Funding

This work was funded by BHF (FS/IBSRF/21/25085) to K.I. S.A. received a 4-year BHF

PhD Studentships (FS/4yPhD/F/22/34181). P.A.E. and J.A.D. were supported by the

Radiation Research Unit at the Cancer Research UK City of London Centre Award

[C7893/A28990]. This work was also supported by the Francis Crick Institute, which

receives its core funding from Cancer Research UK, the UK Medical Research Council and

Wellcome Trust (all under FC001051) to J.B. Research infrastructure within the Institute of

Child Health, UCL is supported by the NIHR Great Ormond Street Hospital Biomedical

Research Centre. The views expressed are those of the authors and not necessarily those of

the NHS, the NIHR or the Department of Health.

Data Availability

676 All relevant data are within the paper and its Supporting Information files.

Competing interests

678 The authors have declared that no competing interests exist.

References

679   1.   Zhang Q, Carlin D, Zhu F, Cattaneo P, Ideker T, Evans SM, et al. Unveiling
680   Complexity and Multipotentiality of Early Heart Fields. Circ Res. 2021;129(4):474-87. Epub
681   20210624. doi: 10.1161/CIRCRESAHA.121.318943. PubMed PMID: 34162224; PubMed
682   Central PMCID: PMCPMC9308985.
683

684   2.   Lescroart F, Chabab S, Lin X, Rulands S, Paulissen C, Rodolosse A, et al. Early
685   lineage restriction in temporally distinct populations of Mesp1 progenitors during mammalian
686   heart development. Nat Cell Biol. 2014;16(9):829-40. Epub 20140824. doi:
687   10.1038/ncb3024. PubMed PMID: 25150979; PubMed Central PMCID: PMCPMC6984965.
688

688   3.   Devine WP, Wythe JD, George M, Koshiba-Takeuchi K, Bruneau BG. Early
689   patterning and specification of cardiac progenitors in gastrulating mesoderm. Elife. 2014;3.
690   Epub 20141008. doi: 10.7554/eLife.03848. PubMed PMID: 25296024; PubMed Central
691   PMCID: PMCPMC4356145.
692

692   4.   Cohen-Gould L, Mikawa T. The fate diversity of mesodermal cells within the heart
693   field during chicken early embryogenesis. Dev Biol. 1996;177(1):265-73. doi:
694   10.1006/dbio.1996.0161. PubMed PMID: 8660893.
695

695   5.   Wolff C, Tinevez JY, Pietzsch T, Stamataki E, Harich B, Guignard L, et al. Multi-
696   view light-sheet imaging and tracking with the MaMuT software reveals the cell lineage of a
697   direct developing arthropod limb. Elife. 2018;7. Epub 20180329. doi: 10.7554/eLife.34410.
698   PubMed PMID: 29595475; PubMed Central PMCID: PMCPMC5929908.
699

699   6.   Dominguez MH, Krup AL, Muncie JM, Bruneau BG. Graded mesoderm assembly
700   governs cell fate and morphogenesis of the early mammalian heart. Cell. 2023;186(3):479-96
701   e23. doi: 10.1016/j.cell.2023.01.001. PubMed PMID: 36736300; PubMed Central PMCID:
702   PMCPMC10091855.
703

703   7.   Ivanovitch K, Soro-Barrio P, Chakravarty P, Jones RA, Bell DM, Mousavy Gharavy
704   SN, et al. Ventricular, atrial, and outflow tract heart progenitors arise from spatially and
705   molecularly distinct regions of the primitive streak. PLoS Biol. 2021;19(5):e3001200. Epub
706   20210517. doi: 10.1371/journal.pbio.3001200. PubMed PMID: 33999917; PubMed Central
707   PMCID: PMCPMC8158918.
708

708   8.   Bardot E, Calderon D, Santoriello F, Han S, Cheung K, Jadhav B, et al. Foxa2
709   identifies a cardiac progenitor population with ventricular differentiation potential. Nat
710   Commun. 2017;8:14428. Epub 20170214. doi: 10.1038/ncomms14428. PubMed PMID:
711   28195173; PubMed Central PMCID: PMCPMC5316866.
712

9.   Gonzalez DM, Schrode N, Ebrahim TAM, Broguiere N, Rossi G, Drakhlis L, et al.
Dissecting mechanisms of chamber-specific cardiac differentiation and its perturbation
following retinoic acid exposure. Development. 2022;149(13). Epub 20220708. doi:
10.1242/dev.200557. PubMed PMID: 35686629; PubMed Central PMCID:
PMCPMC9340554.

10. Saykali B, Mathiah N, Nahaboo W, Racu ML, Hammou L, Defrance M, et al. Distinct
mesoderm migration phenotypes in extra-embryonic and embryonic regions of the early
mouse embryo. Elife. 2019;8. Epub 20190405. doi: 10.7554/eLife.42434. PubMed PMID:
30950395; PubMed Central PMCID: PMCPMC6450669.

11. Christiaen L, Davidson B, Kawashima T, Powell W, Nolla H, Vranizan K, et al. The
transcription/migration interface in heart precursors of Ciona intestinalis. Science.
2008;320(5881):1349-52. doi: 10.1126/science.1158170. PubMed PMID: 18535245.

12. Szymczak AL, Workman CJ, Wang Y, Vignali KM, Dilioglou S, Vanin EF, et al.
Correction of multi-gene deficiency in vivo using a single 'self-cleaving' 2A peptide-based
retroviral vector. Nat Biotechnol. 2004;22(5):589-94. Epub 20040404. doi: 10.1038/nbt957.
PubMed PMID: 15064769.

13. Ivanovitch K, Temino S, Torres M. Live imaging of heart tube development in mouse
reveals alternating phases of cardiac differentiation and morphogenesis. Elife. 2017;6. Epub
20171205. doi: 10.7554/eLife.30668. PubMed PMID: 29202929; PubMed Central PMCID:
PMCPMC5731822.

14. Horl D, Rojas Rusak F, Preusser F, Tillberg P, Randel N, Chhetri RK, et al.
BigStitcher: reconstructing high-resolution image datasets of cleared and expanded samples.
Nat Methods. 2019;16(9):870-4. Epub 20190805. doi: 10.1038/s41592-019-0501-0. PubMed
PMID: 31384047.

15. Malin-Mayor C, Hirsch P, Guignard L, McDole K, Wan Y, Lemon WC, et al.
Automated reconstruction of whole-embryo cell lineages by learning from sparse
annotations. Nat Biotechnol. 2023;41(1):44-9. Epub 20220905. doi: 10.1038/s41587-022-
01427-7. PubMed PMID: 36065022; PubMed Central PMCID: PMCPMC7614077.

16. Sugawara K, Cevrim C, Averof M. Tracking cell lineages in 3D by incremental deep
learning. Elife. 2022;11. Epub 20220106. doi: 10.7554/eLife.69380. PubMed PMID:
34989675; PubMed Central PMCID: PMCPMC8741210.

17. Moorman A, Webb S, Brown NA, Lamers W, Anderson RH. Development of the
heart: (1) formation of the cardiac chambers and arterial trunks. Heart. 2003;89(7):806-14.
doi: 10.1136/heart.89.7.806. PubMed PMID: 12807866; PubMed Central PMCID:
PMCPMC1767747.

18. Wei Y, Mikawa T. Fate diversity of primitive streak cells during heart field formation
in ovo. Dev Dyn. 2000;219(4):505-13. doi: 10.1002/1097-0177(2000)9999:9999<::AID-
DVDY1076>3.0.CO;2-6. PubMed PMID: 11084650.

19. Kelly RG, Brown NA, Buckingham ME. The arterial pole of the mouse heart forms
from Fgf10-expressing cells in pharyngeal mesoderm. Dev Cell. 2001;1(3):435-40. doi:
10.1016/s1534-5807(01)00040-5. PubMed PMID: 11702954.

20. Mjaatvedt CH, Nakaoka T, Moreno-Rodriguez R, Norris RA, Kern MJ, Eisenberg
CA, et al. The outflow tract of the heart is recruited from a novel heart-forming field. Dev
Biol. 2001;238(1):97-109. doi: 10.1006/dbio.2001.0409. PubMed PMID: 11783996.

21. Waldo KL, Kumiski DH, Wallis KT, Stadt HA, Hutson MR, Platt DH, et al.
Conotruncal myocardium arises from a secondary heart field. Development.
2001;128(16):3179-88. doi: 10.1242/dev.128.16.3179. PubMed PMID: 11688566.

22. Zaffran S, Kelly RG, Meilhac SM, Buckingham ME, Brown NA. Right ventricular
myocardium derives from the anterior heart field. Circ Res. 2004;95(3):261-8. Epub
20040624. doi: 10.1161/01.RES.0000136815.73623.BE. PubMed PMID: 15217909.

23. Sakoe H, Chiba S. Dynamic-Programming Algorithm Optimization for Spoken Word
Recognition. Ieee T Acoust Speech. 1978;26(1):43-9. doi: Doi 10.1109/Tassp.1978.1163055.
PubMed PMID: WOS:A1978EN09300006.

24. Shokoohi-Yekta M, Hu B, Jin H, Wang J, Keogh E. Generalizing DTW to the multi-
dimensional case requires an adaptive approach. Data Min Knowl Discov. 2017;31(1):1-31.
Epub 20160215. doi: 10.1007/s10618-016-0455-0. PubMed PMID: 29104448; PubMed
Central PMCID: PMCPMC5668684.

25. Liang X, Wang G, Lin L, Lowe J, Zhang Q, Bu L, et al. HCN4 dynamically marks the
first heart field and conduction system precursors. Circ Res. 2013;113(4):399-407. Epub
20130606. doi: 10.1161/CIRCRESAHA.113.301588. PubMed PMID: 23743334; PubMed
Central PMCID: PMCPMC4017870.

26. Spater D, Abramczuk MK, Buac K, Zangi L, Stachel MW, Clarke J, et al. A HCN4+
cardiomyogenic progenitor derived from the first heart field and human pluripotent stem
cells. Nat Cell Biol. 2013;15(9):1098-106. Epub 20130825. doi: 10.1038/ncb2824. PubMed
PMID: 23974038.

27. Tyser RCV, Ibarra-Soria X, McDole K, Arcot Jayaram S, Godwin J, van den Brand
TAH, et al. Characterization of a common progenitor pool of the epicardium and
myocardium. Science. 2021;371(6533). Epub 20210107. doi: 10.1126/science.abb2986.
PubMed PMID: 33414188.

28. Watanabe Y, Wang Y, Tanaka Y, Iwase A, Kawamura T, Saga Y, et al. Hey2
enhancer activity defines unipotent progenitors for left ventricular cardiomyocytes in juxta-
cardiac field of early mouse embryo. Proc Natl Acad Sci U S A. 2023;120(37):e2307658120.
Epub 20230905. doi: 10.1073/pnas.2307658120. PubMed PMID: 37669370; PubMed Central
PMCID: PMCPMC10500178.

29. Stainier DY, Lee RK, Fishman MC. Cardiovascular development in the zebrafish. I.
Myocardial fate map and heart tube formation. Development. 1993;119(1):31-40. doi:
10.1242/dev.119.1.31. PubMed PMID: 8275863.

30. Yutzey KE, Bader D. Diversification of cardiomyogenic cell lineages during early
heart development. Circ Res. 1995;77(2):216-9. doi: 10.1161/01.res.77.2.216. PubMed
PMID: 7614708.

31. Keegan BR, Meyer D, Yelon D. Organization of cardiac chamber progenitors in the
zebrafish blastula. Development. 2004;131(13):3081-91. Epub 20040602. doi:
10.1242/dev.01185. PubMed PMID: 15175246.

32.   Garcia-Martinez V, Schoenwolf GC. Primitive-streak origin of the cardiovascular
system in avian embryos. Dev Biol. 1993;159(2):706-19. doi: 10.1006/dbio.1993.1276.
PubMed PMID: 8405690.

33.   Dark N, Cosson MV, Tsansizi LI, Owen TJ, Ferraro E, Francis AJ, et al. Generation
of left ventricle-like cardiomyocytes with improved structural, functional, and metabolic
maturity from human pluripotent stem cells. Cell Rep Methods. 2023;3(4):100456. Epub
20230424. doi: 10.1016/j.crmeth.2023.100456. PubMed PMID: 37159667; PubMed Central
PMCID: PMCPMC10163040.

34.   Mendjan S, Mascetti VL, Ortmann D, Ortiz M, Karjosukarso DW, Ng Y, et al.
NANOG and CDX2 pattern distinct subtypes of human mesoderm during exit from
pluripotency. Cell Stem Cell. 2014;15(3):310-25. Epub 20140718. doi:
10.1016/j.stem.2014.06.006. PubMed PMID: 25042702.

35.   Lee JH, Protze SI, Laksman Z, Backx PH, Keller GM. Human Pluripotent Stem Cell-
Derived Atrial and Ventricular Cardiomyocytes Develop from Distinct Mesoderm
Populations. Cell Stem Cell. 2017;21(2):179-94 e4. doi: 10.1016/j.stem.2017.07.003.
PubMed PMID: 28777944.

36.   Schmidt C, Deyett A, Ilmer T, Haendeler S, Torres Caballero A, Novatchkova M, et
al. Multi-chamber cardioids unravel human heart development and cardiac defects. Cell.
2023. Epub 20231116. doi: 10.1016/j.cell.2023.10.030. PubMed PMID: 38029745.

37.   Yang D, Gomez-Garcia J, Funakoshi S, Tran T, Fernandes I, Bader GD, et al.
Modeling human multi-lineage heart field development with pluripotent stem cells. Cell Stem
Cell. 2022;29(9):1382-401 e8. doi: 10.1016/j.stem.2022.08.007. PubMed PMID: 36055193.

38.   DeRuiter MC, Poelmann RE, VanderPlas-de Vries I, Mentink MM, Gittenberger-de
Groot AC. The development of the myocardium and endocardium in mouse embryos. Fusion
of two heart tubes? Anat Embryol (Berl). 1992;185(5):461-73. doi: 10.1007/BF00174084.
PubMed PMID: 1567022.

39.   Zeng XX, Wilm TP, Sepich DS, Solnica-Krezel L. Apelin and its receptor control
heart field formation during zebrafish gastrulation. Dev Cell. 2007;12(3):391-402. doi:
10.1016/j.devcel.2007.01.011. PubMed PMID: 17336905.

40.   Scott IC, Masri B, D'Amico LA, Jin SW, Jungblut B, Wehman AM, et al. The g
protein-coupled receptor agtrl1b regulates early development of myocardial progenitors. Dev
Cell. 2007;12(3):403-13. doi: 10.1016/j.devcel.2007.01.012. PubMed PMID: 17336906.

41.   Kupperman E, An S, Osborne N, Waldron S, Stainier DY. A sphingosine-1-phosphate
receptor regulates cell migration during vertebrate heart development. Nature.
2000;406(6792):192-5. doi: 10.1038/35018092. PubMed PMID: 10910360.

42.   Ciruna B, Rossant J. FGF signaling regulates mesoderm cell fate specification and
morphogenetic movement at the primitive streak. Dev Cell. 2001;1(1):37-49. doi:
10.1016/s1534-5807(01)00017-x. PubMed PMID: 11703922.

43.   Nutt SL, Dingwell KS, Holt CE, Amaya E. Xenopus Sprouty2 inhibits FGF-mediated
gastrulation movements but does not affect mesoderm induction and patterning. Genes Dev.
2001;15(9):1152-66. doi: 10.1101/gad.191301. PubMed PMID: 11331610; PubMed Central
PMCID: PMCPMC312687.

44.   Yang X, Dormann D, Munsterberg AE, Weijer CJ. Cell movement patterns during
gastrulation in the chick are controlled by positive and negative chemotaxis mediated by
FGF4 and FGF8. Dev Cell. 2002;3(3):425-37. doi: 10.1016/s1534-5807(02)00256-3.
PubMed PMID: 12361604.

45.   von der Hardt S, Bakkers J, Inbal A, Carvalho L, Solnica-Krezel L, Heisenberg CP, et
al. The Bmp gradient of the zebrafish gastrula guides migrating lateral cells by regulating
cell-cell adhesion. Curr Biol. 2007;17(6):475-87. Epub 20070301. doi:
10.1016/j.cub.2007.02.013. PubMed PMID: 17331724.

46.   Guo Q, Li JY. Distinct functions of the major Fgf8 spliceform, Fgf8b, before and
during mouse gastrulation. Development. 2007;134(12):2251-60. Epub 20070516. doi:
10.1242/dev.004929. PubMed PMID: 17507393; PubMed Central PMCID:
PMCPMC2518685.

47.   Tam PP, Parameswaran M, Kinder SJ, Weinberger RP. The allocation of epiblast cells
to the embryonic heart and other mesodermal lineages: the role of ingression and tissue
movement during gastrulation. Development. 1997;124(9):1631-42. doi:
10.1242/dev.124.9.1631. PubMed PMID: 9165112.

48.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al.
Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.
Epub 20120628. doi: 10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central
PMCID: PMCPMC3855844.

49.   Cockroft DL, Brook FA, Copp AJ. Inositol deficiency increases the susceptibility to
neural tube defects of genetically predisposed (curly tail) mouse embryos in vitro.
Teratology. 1992;45(2):223-32. doi: 10.1002/tera.1420450216. PubMed PMID: 1615432.

50.   Takahashi M, Makino S, Kikkawa T, Osumi N. Preparation of rat serum suitable for
mammalian whole embryo culture. J Vis Exp. 2014;(90):e51969. Epub 20140803. doi:
10.3791/51969. PubMed PMID: 25145996; PubMed Central PMCID: PMCPMC4672955.

51.   Ichikawa T, Nakazato K, Keller PJ, Kajiura-Kobayashi H, Stelzer EH, Mochizuki A,
et al. Live imaging and quantitative analysis of gastrulation in mouse embryos using light-
sheet microscopy and 3D tracking tools. Nat Protoc. 2014;9(3):575-85. Epub 20140213. doi:
10.1038/nprot.2014.035. PubMed PMID: 24525751.

52.   Yu G. Using ggtree to Visualize Data on Tree-Like Structures. Curr Protoc
Bioinformatics. 2020;69(1):e96. doi: 10.1002/cpbi.96. PubMed PMID: 32162851.

53.   Revell LJ. phytools: an R package for phylogenetic comparative biology (and other
things). Methods Ecol Evol. 2012;3(2):217-23. doi: 10.1111/j.2041-210X.2011.00169.x.
PubMed PMID: WOS:000302538500001.

54.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al.
SciPy 1.0: fundamental algorithms for scientific computing in Python. Nature Methods.
2020;17(3):261-72. doi: 10.1038/s41592-019-0686-2. PubMed PMID:
WOS:000510821500001.

Figure1

Figure2

Figure3

Figure 4

A Movie 1_unipotent Pericardial progenitor
T/Bra expression reporter T/Bra lineage cTnnT2 reporter

E6.5 | 0 | 13h50 | 21h20 | 30h | time

ii — Mother — 0
iii — Grand daughters
iv XY v XZ vi XZ — D111 D112 D121 D122 D21 D22
vii XY viii XY — 4th generation Pericardium

A'
Mother — GFP
0, 8, 16, 24, 32 hrs
D2, D1, D22, D21, D12, D11, D121, D112, D222, D221, D212, D211, D122, D1212, D1211, D111
CC inflows
Pericardium

B Movie 3_ Bipotent Endocardial and ExE meso progenitor
T/Bra expression reporter T/Bra lineage cTnnT2 reporter

E7 | 0 | 3h30 | 15h37 | 21h10

i XY — Mother
ii — Daughters
iii iv — D21 and D22 Ex-meso D12 and D11 Endocardium
v XY vi YZ vii YZ — D111 and D112 Endocardium

B'
Mother — GFP
0, 8, 16, 24 hrs
D2, D1, D22, D21, D12, D11, D212, D211, D222, D221, D122, D121, D111, D112
CC inflows
ExE meso Endocardium

C Movie 2_ bipotent endocardium/myocyte progenitor
T/Bra expression reporter T/Bra lineage cTnnT2 reporter

1h40 | 1h50 | 10h | 15h48

Mother — Daughters — GrandDaughters
XY / YZ — D21 myocytes D12 endocardium

C'
Mother — GFP
8, 16, 24, 32 hrs
D2, D1
CC inflows
Endocardium
LV/AVC

D Movie 2_ endocardium progenitor
T/Bra lineage cTnnT2 reporter

0 | 2h30 | 3h08 | 3h30
3h34 | 3h50 | 3h58 | 7h22

E Movie1_ Bipotent endothelial-like / ExE mesoderm progenitor
T/Bra expression reporter T/Bra lineage cTnnT2 reporter

E6.5 | 0 | 4h40 | 10h30 | 15h | 18h20

Mother — Daughters — D2 D1 — D21 D22 D11 D12

E'
Mother — GFP
8, 16, 24, 32 hrs
D2, D1, D22, D21, D12, D11, D122, D121, D112, D111
CC inflows
ExE meso Endothelial-like

Figure5

Figure6

Figure2_supplementary figure 1

A

$T^{nGFP-CreERT2/+}; R26^{tdtomato/+}; cTnnT-2a-eGFP$

T/Bra expression reporter   T/Bra lineage

Figure2_supplementary figure 2

Figure2_supplementary figure 3

Figure3_supplementary figure 1

A

Figure4_supplementary figure 1

A

Movie 3_ tripotent exMeso/endocardium/myocyte progenitor

T/Bra expression reporter   T/Bra lineage   cTnnT2 reporter

0   1h42   9h   19h20   25h20

Mother   1st generation   2nd generation

D111 D112 D121 D122 (myocytes)

D111 D121 D122 D12 (myocytes)

D222 D221 (endocardium)

D222 D221 (endocardium)

D211 D212 ExMeso

D2111 D2112 D2121 D2122 (ExMeso)

3rd generation

B

Mother

CC

inflows

D2   D1

Endocardium   ExE m   myocytes

GFP

Figure5_supplementary figure 1