Title: <u>Inverse folding of protein complexes with a structure-informed language model enables</u>

<u>unsupervised antibody evolution</u>

Authors: Varun R. Shanker[1,2,3], Theodora U.J. Bruun[2,3,4], Brian L. Hie[3,4]*, Peter S. Kim[3,4,5]*

Affiliations:

[1]Stanford Biophysics Program, Stanford University School of Medicine, Stanford, CA 94305,

USA

[2]Stanford Medical Scientist Training Program, Stanford University School of Medicine, Stanford

CA 94305, USA

[3]Sarafan ChEM-H, Stanford University, Stanford, CA 94305, USA

[4]Department of Biochemistry, Stanford University School of Medicine, Stanford, CA 94305,

USA

[5]Chan Zuckerberg Biohub, San Francisco, CA 94158, USA.

*Correspondence: B.L.H. (brianhie@stanford.edu), P.S.K. (kimpeter@stanford.edu)

Abstract

Large language models trained on sequence information alone are capable of learning high level

principles of protein design. However, beyond sequence, the three-dimensional structures of

proteins determine their specific function, activity, and evolvability. Here we show that a general

protein language model augmented with protein structure backbone coordinates and trained on

the inverse folding problem can guide evolution for diverse proteins without needing to

explicitly model individual functional tasks. We demonstrate inverse folding to be an effective

unsupervised, structure-based sequence optimization strategy that also generalizes to multimeric

complexes by implicitly learning features of binding and amino acid epistasis. Using this

approach, we screened ~30 variants of two therapeutic clinical antibodies used to treat SARS-

CoV-2 infection and achieved up to 26-fold improvement in neutralization and 37-fold

improvement in affinity against antibody-escaped viral variants-of-concern BQ.1.1 and XBB.1.5,

respectively. In addition to substantial overall improvements in protein function, we find inverse

folding performs with leading experimental success rates among other reported machine

learning-guided directed evolution methods, without requiring any task-specific training data.

Introduction

Evolution generates diverse proteins at the level of biological sequences by exploring a vast search space of potential mutations and acquiring those that improve fitness. However, it is the three-dimensional structure encoded by these sequences that ultimately determines the function and activity of a protein. Consequently, as proteins accumulate mutations, they undergo corresponding structural changes, which in turn facilitate functional adaptations[1].

In the laboratory, this tendency for greater sequence change to cause structural divergence poses a major challenge to engineering better proteins via a stepwise evolutionary process. Mutations added in sequential rounds of artificial evolution are increasingly likely to destabilize the structure and therefore diminish the protein's evolvability[2]. Identifying beneficial mutations is further challenged by the fact that almost all mutations to a prototypical protein are deleterious, or at best neutral, and only a rare subset are beneficial on its fitness landscape[3–8]. In total, these phenomena can often reduce the evolutionarily accessible paths and make evolution more susceptible to local fitness optima[9,10], further complicating attempts to increase fitness.

To address both the structural constraints of protein design and the high dimensionality of the mutational search space, we utilized a general protein language model augmented with structural information and trained across millions of non-redundant single sequence-structure pairs on the inverse folding objective[11]. Most simply, the inverse folding problem considers the task opposite of that performed by many of the recent powerful structure-prediction tools, including AlphaFold and ESMFold[12,13]: recovery of a protein's native sequence, given its three-dimensional backbone coordinates (**Figure 1a**). This is accomplished by predicting the identity of an amino acid given both the preceding amino acid sequence (referred to as autoregressive modeling) and the entire structure's backbone coordinates (**Methods**). Thus, sequences assigned

high likelihood scores by the inverse folding language model are expected to fold into the

backbone of the input structure with high confidence (**Figure 1b**).

Our inverse folding framework for protein design does not model an explicit protein

function or definition of protein fitness. Rather, using a structure-guided paradigm, we indirectly

explore the underlying fitness landscape by focusing exploration to regions where the backbone

fold of the protein is preserved. We hypothesize constraining evolution to regimes of high

inverse folding likelihood can serve as an effective prior for high-fitness variants, and thereby

improve the efficiency of evolution (**Figure 1c**).

We reasoned that this approach may be particularly valuable for the evolution of human

antibodies, which are used clinically to treat a broad range of diseases[14]. Antibodies are used

therapeutically to bind to a target antigen mediating pathogenesis, and modify or disrupt its

function[15] . A central concept of this study is to use the complete structure of the antibody-

antigen complex to guide evolution. By conditioning the inverse folding model on the entire

antibody-antigen complex, we sought to enable the discovery of mutations that preserve or

enhance the stability of the entire complex, and thus that improve antibody function.

Indeed, we show that as an unsupervised machine learning-guided evolution strategy,

inverse folding is capable of identifying high fitness mutations across several protein families

and tasks, performing better than sequence-only methods. We found that inverse folding

generalizes to protein complexes with improved antibody variant prediction when antigen

structural information is also included as input. To demonstrate the practical utility of this

method, we improved the potency of mature, clinical SARS-CoV-2 monoclonal antibody

therapies, in a low-throughput setting, against both their original viral target as well as viral

escape variants that reduced their efficacy, namely variants-of-concern (VOC) BQ.1.1 and

83 XBB.1.5. We achieved up to 26-fold improvement in the neutralization potency of Ly-1404

84 (Bebtelovimab) against BQ.1.1, and 11-fold for SA58, testing only a total of 31 and 25 antibody

85 variants, respectively. We also achieved 27-fold improvement in affinity against BQ.1.1 and 37-

86 fold improvement in affinity against XBB.1.5. Notably, all experimentally tested combinations

87 of inverse folding-recommended mutations showed improved activity, with many designs

88 comprising multiple synergistic mutations. With our approach, we report experimental success

89 rates that surpass all previous machine learning-guided protein evolution methods[8,16–28], including

90 those based on supervision with task-specific training data. These findings highlight the

91 advantage of an unsupervised, structure-based paradigm to identify efficient evolutionary

92 trajectories.

94 **Results**

95 *Inverse folding enriches sequence exploration for high function protein variants across diverse*

96 *tasks*

97   We evaluated whether inverse folding can be used to guide protein evolution, without

98 needing to explicitly model specific functional tasks, by assessing its ability to identify mutations

99 resulting in high levels of protein activity for a desired functional property, or fitness measure.

100 Accordingly, for 10 proteins from diverse families among four organisms, and with functions

101 ranging from enzyme catalysis (TPMT) to oncogenesis (HRAS) to transcriptional regulation

102 (GAL4), we used inverse folding likelihoods to score variants profiled in large datasets from

103 deep mutational scanning experiments[29–38]against a target backbone of the wild-type protein[39–48]

104 (**Methods**, **Supplementary Table 1**).

From the thousands of tested variants for each of the 10 proteins, we identified numerous

with experimentally determined protein activities ranking in the top percentiles of the entire

screen within just the set of top ten inverse-folding predictions (**Figure 1d**). Our analysis also

demonstrates that conditioning on structural information serves to improve predictive capabilities

of protein language models as we successfully identified mutations in the top fifth percentile for

9 out of the 10 proteins using inverse folding compared to just 2 proteins using a state-of-the-art

general protein language model trained only on sequence information and specifically for variant

prediction (ESM-1v)[49] (**Figure 1d**). This improvement in prediction also holds with increasingly

relaxed thresholds for classification as high-fitness variants.

These results suggest that inverse folding offers a promising alternative to brute force

experimental searches for beneficial mutations. Notably, some of the top mutations predicted by

inverse folding are also the same ones recovered from exhaustive experimental exploration. For

example, for restriction enzyme haeIIIM, variant Q18E is recommended within the top five

inverse folding predictions and experimentally ranks as the second-best substitution (and > 5

standard deviations above the mean) out of the nearly 2000 substitutions screened to the

endonuclease[38]. Another key advantage of our task-independent framework, in addition to being

broadly applicable across diverse proteins, is the ability to improve a single protein for multiple

desired properties without needing to develop specialized high-throughput assays to screen each

independently. From just the top 10 inverse folding predictions for MAPK1, we identify

substitutions Q105M and Y64D, which are experimentally shown to confer resistance to two

different oncogenic-targeting MAPK1 kinase inhibitors[32].

Inverse folding is a state-of-the-art zero-shot mutational effect predictor for antibodies

To analyze the effectiveness of augmenting a general protein language model with structural information, specifically for antibody variant prediction, we compared the inverse folding likelihoods of sequences across entire mutational landscapes against the corresponding experimental fitness values from three existing mutagenesis datasets. The first two of the datasets profile the scFv equilibrium dissociation constants (K_D) of all possible evolutionary intermediates between the inferred germline and somatic sequence of naturally affinity-matured influenza broadly neutralizing antibodies (bnAbs) CR9114 and CR6261, which bind the conserved stem epitope of influenza surface protein hemagglutinin (HA)[50]. For both bnAbs, only mutations in the heavy chain, which is responsible for antigen binding, were characterized. The profiled mutational landscape of CR9114 includes all possible combinations of 16 substitutions while that of CR6261 includes all possible combinations of 11 substitutions, totaling $2^{16} = 65,536$ and $2^{11} = 2,048$ variant antibody sequences respectively. Each of these libraries were screened for binding against two distinct influenza HA subtypes (H1 and H3 for CR9114 and H1 and H9 for CR6261). The third dataset assesses the effects of all possible single amino acid substitutions with a deep mutational scan profiling 4,275 mutations in the variable regions for both heavy chain (VH) and light chain (VL) of antibody G6.31 to binding with its ligand, vascular endothelial growth factor A (VEGF-A)[51].

For each dataset, we computed the Spearman correlation between the log likelihood estimated by the inverse folding model and the experimentally determined binding measure for a given antigen, across all sequences in the mutational library. We scored the inverse folding likelihood of each candidate sequence in the library using the backbone coordinates of a structure with the mature antibody bound to its target antigen[52–54].

Across all five experimental binding datasets, we found that inverse folding performs

better than both a sequence-only language model, ESM-1v[49], and a site-independent model of

mutational frequency curated with extensive antibody sequence alignments, abYsis[55]. In nearly

all experimental scenarios, supplementing sequence information with the backbone coordinates

of the antibody alone, without providing antigen information, as input to inverse folding is

sufficient to outperform other sequence-only methods. A notable feature of the autoregressive

architecture is that it computes the joint likelihood over all positions in a sequence, making it

well-suited to score combinatorial sequence changes. We find that inverse folding can capture

complex epistatic interactions, or potential interdependence among individual amino acids, as it

performs well on the CR9114 and CR6261 libraries composed of sequences with multiple

mutations (**Figure 2a,b**).

We achieved the greatest improvement in performance on all five experimental screens

by incorporating the structure of both the antibody and antigen (**Figure 2a**), indicating that the

inverse folding model can implicitly learn features of binding (**Figure 2c**). This result is

particularly significant, given that the inverse folding model is only trained on single-chain

protein structures, while the antibody-antigen complexes we use as inputs are composed of either

three (G6.31) or four (CR9114, CR6261) protein chains. The most substantial contribution of

antigen information is observed in the case of CR9114-H1, for which the correlation increases

from 0.17 with only antibody information to 0.65 with sequence and backbone coordinates of the

entire complex.

Remarkably, we could still predict effects of mutations on binding for a cross-reactive

antibody while using a different antigen as input to the model. (**Figure 2a,b**). Despite using a

complex with HA from H5N1 influenza as input to score CR9114 variants, we obtain

correlations of 0.65 and 0.50 with experimental binding data for H1 and H3, respectively. This is

particularly striking since, for example, H5 and H1 only share 63% sequence identify across both

HA subunits (**Supplementary Figure 3**). This same cross-reactive predictive capability is

observed for CR6261, which is tested experimentally against H1 and H9 while we use an input

structure with HA from 1918 H1N1 influenza (**Figure 2a**). Although inverse folding cannot learn

explicit chemical rules of binding (e.g., hydrogen bonding or disulfide bridge formation) since it

does not have access to amino acid side chain atomic coordinates, these results suggest that

structural principles like interface packing or potential steric interference are not only implicitly

accessible from residue identities, but are also informative for binding prediction.

Our model's top recommended mutations are made independently of a specific definition

of fitness; they simply represent a set of variants with a high likelihood of folding into the input

backbone structure. Therefore, our model's recommendations may also help identify mutations

that improve other useful biochemical properties beyond affinity. Impressively, for example, the

top inverse folding-recommended mutation to the VL of G6.31 is F83A, which was identified in

the original screening study[51] to be particularly interesting as it confers a three-fold increase in

VEGF-A binding affinity and a 5°C improvement in melting temperature, despite being 25Å

from the antigen and in the antibody framework region. It was determined that the VL F83A

substitution induces more compact packing and the site serves as a conformational switch that

affects biological activity at the antibody-antigen interface by modulating both interdomain and

elbow angle dynamics[51].

Engineering therapeutic antibodies for increased potency and resilience

Finally, we aimed to assess if the structure-augmented language model's predictive capabilities could not only resolve trends on large sets of experimental data, but also enable efficient and successful directed evolution campaigns while testing only a small number (on the order of tens) of variants. To do so, we considered the task of improving the potency and resilience (effectiveness against a virus as it mutates over time) of two mature, clinical monoclonal antibody therapies.

- Ly-1404 (Bebtelovimab) was isolated from a COVID-19 convalescent donor and binds to the receptor binding domain (RBD) of the SARS-CoV-2 Spike protein[56]. It was approved by the U.S. F.D.A. on February 11, 2022 given its activity against both the original Wuhan and Omicron SARS-CoV-2 variants and was the last remaining approved monoclonal antibody therapy withstanding against viral evolution[57] until its discontinuation on November 30, 2023 due to antibody evasion by VOC BQ.1.1.[58]

- SA58 (BD55-5840) was isolated from a vaccinated individual and is one of two RBD-targeting neutralizing antibodies (NAb) in a rationally developed antibody cocktail. SA58 alone retained efficacy against all Omicron subvariants, including *in vivo* protection against BA.5[59,60] and was shown to be effective as a post-exposure prophylaxis in a clinical study[61].

For both antibody engineering campaigns, we used the inverse folding language model to compute likelihoods of all ~4,300 possible single-residue substitutions in the VH or VL regions of the antibody. In the first round of evolution, we selected only the top ten predictions at unique residues in each chain for experimental validation. An important practical benefit of our method

is the ability to optimize against measures of fitness most relevant to the protein's downstream

function, rather than being limited to indirect and less accurate surrogate measures that are more

amenable to high-throughput screening[4,16]. We leverage this advantage to directly evolve these

antibodies for their ability to more potently neutralize SARS-CoV-2 pseudotyped lentivirus.

Variants recommended by the inverse folding language model were assessed by

comparing the half-maximal inhibitory concentration (IC_{50}) relative to the wild-type antibody.

Remarkably, although we chose to only test 20 single-site substitutions for each of the two

clinical monoclonal antibody therapies, approximately one-third of them improved neutralizing

potency. Notably, several of these variants improve neutralization IC_{50} by approximately 2-fold

with just a single amino acid change (**Figure 3a**, **Supplementary Data 1**).

Prompted by recent evidence showing that conservation of the overall RBD structure is

robust to SARS-CoV-2 evolution[62], we next sought to determine whether we could also evolve

the previously mature antibodies against SARS-CoV-2 BQ.1.1, the variant responsible for

diminished therapeutic efficacy. Although the antibodies were previously effective, a change in

antigen conceptually represents a fundamental shift in the underlying fitness landscape (**Figure

3b**). From the same set of 20 single amino acid substitutions to Ly-1404, we found that nearly

half improve neutralization of variant BQ.1.1. In addition to a high success rate, we also found

multiple of these mutations provided a large magnitude of improvement. Several single amino

acid substitutions to Ly-1404 individually result in over a 3-fold improvement while the most

beneficial mutation to SA58 results in a nearly 7-fold improvement (**Figure 3c**).

Taken together, approximately two-third and one-third of tested single amino acid

substitutions to Ly-1404 and SA58, respectively, were beneficial for neutralization of either the

original strain or BQ.1.1. These results reinforce that, despite all being predicted to have the

same backbone fold, inverse folding variants feature functional diversity and can be used for

distinct notions of protein fitness. Interestingly, for both antibodies, the most beneficial mutation,

is not shared by the each of the strains tested (**Supplementary Figure 4**).

A common challenge in directed evolution is contending with the combinatorial

explosion of possible sequences which emerges from trying to combine a set of individually

beneficial mutations. In the second round of evolution, we simply use the inverse folding model

again to acquire up to five top-scoring unique combinations of mutations to each antibody chain

(**Methods**). Notably, across both evolutionary trajectories, all 15 antibody designs with multiple

mutations have IC_{50} values better than wild-type, with many designs showing synergistic effects

upon combination. For example, just a single amino acid mutation in each of the two chains of

SA58 leads to over an 11-fold improvement (**Figure 3c,d**). Similarly, the most potent evolved

design of Ly-1404 is a combination of seven of the eight beneficial single amino acid

substitution to the VH and improves neutralization 26-fold (**Figure 3d**). Critically, these

improvements to neutralizing potency against BQ.1.1 do not sacrifice potency against the

original strains. We found that the top SA58 design against BQ.1.1 after the second round of

evolution also improves BA.1 neutralization nearly 3-fold (**Supplementary Data 1**).

Additional characterization of evolved antibodies

To further characterize the basis for enhanced neutralization of SARS-CoV-2 VOC

BQ.1.1, we tested the binding affinity of all variant antibodies to RBD as bivalent IgG using

biolayer interferometry (BLI) to obtain the apparent dissociation constant ($K_{D,app}$). For Ly-1404,

all 23 variants with improved neutralization also have improved binding affinity up to ~27-fold.

Interestingly, we found four additional inverse folding-recommended mutations, which were

neutral or deleterious to neutralization, also improved binding affinity. Across all variants there

is a Spearman correlation of 0.47 between fold-change in IC_{50} and fold-change in $K_{D,app}$ (**Figure**

4a).

We similarly screened the SA58 variants for binding to the RBD of BQ.1.1. However,

since the K_D of the wildtype antibody as IgG was already sub-picomolar, further improvements to

binding were below the limit of quantitation and indistinguishable using this measure. Given this

strong binding affinity of wildtype SA58 to BQ.1.1 RBD, we also screened this same set of

variants against emerging VOC XBB.1.5 and observe improvements in $K_{D,app}$ up to 37-fold

(**Figure 4c**).

By testing several top affinity-matured designs in a polyspecificity assay, we also

confirmed that improvements in binding are not mediated by generalized enhancements of non-

specific interactions (**Supplementary Figure 5a**). In this assay, we observed no substantial

changes in off-target binding of the evolved antibodies to membrane soluble proteins,

particularly within a therapeutically viable range (as defined by controls of clinically approved

antibodies with recorded high and low polyspecificity). Furthermore, we found no correlation

between fold-change in polyspecificity and affinity fold-change (**Supplementary Figure 5b**).

Analysis of evolutionary exploration

Confronted by the large number of possible mutations, traditional experimental-based

methods for antibody affinity maturation often restrict the mutational search space to only a few

regions of the antibody. Specifically, binding optimization efforts are typically focused within

the complementarity determining regions (CDR), which are hotspots for natural somatic

hypermutation. However, using our unbiased approach to consider all regions of the variable

domain allows for many discoveries that may be less intuitive to a rational designer. For

example, the most beneficial substitutions to Ly-1404, VH F24Y and VH V90S, are located

within framework regions and positioned distally from the binding interface (**Supplementary**

Figure 6, Supplementary Table 2). Interestingly, they both improve neutralization of BQ.1.1

by over 3-fold and are not deleterious to Wuhan neutralization. In other cases, inverse folding

also successfully predicts beneficial substitutions using residues rarely observed among human

antibody sequences. Substitution VL N95V in SA58, which improves neutralization

approximately 7-fold against BQ.1.1, is mediated by the incorporation of a valine observed in

only 0.7% of human antibody sequences at that position and enhances antibody-antigen contact.

While inverse folding is capable of successfully making novel predictions, in some instances it

also does suggest reverting residues to ones frequently selected for in natural somatic

hypermutation. Mutation VL F51Y in Ly-1404 changes a phenylalanine observed in just 5% of

sequences to a tyrosine observed in 86% of sequences. However, this variant results in no change

to Wuhan neutralization. Overall, these results highlight the novelty and value in augmenting a

language model with structural information to evolve antibodies and proteins complexes.

Discussion

The discovery of mutations that improve protein function is inherently challenging due to

the large sequence search space and complex rules that govern the relationship between sequence

and function, such as stability or environmental selection pressures. We show that a general

inverse folding protein language model informed with the sequence and backbone structural

coordinates of a protein can considerably improve directed evolution efforts by serving as an

improved prior compared to sequence-only deep learning methods. Importantly, we highlight

that inverse folding can interrogate protein fitness landscapes indirectly, without needing to

explicitly model individual functional tasks or properties, making it broadly applicable to

proteins across diverse settings ranging from enzyme catalysis to antibiotic and chemotherapy

resistance (**Figure 1d**). We also demonstrate inverse folding generalizes to multimeric proteins,

despite being trained only on single-chain proteins, through its ability to implicitly learn features

of binding. This result is particularly remarkable considering inverse folding has no access to

amino acid side chain atoms, coordinates, or bond information.

Equipped with these capabilities, we use inverse folding to evolve clinical therapeutic

antibodies and identify several mutations which act synergistically to improve antibody potency

and resilience against emerging variants of concern. In the context of pandemics and emergency-

use situations, where monoclonal antibody therapies are limited in supply and vulnerable to

resistance from viral evolution, the ability to rapidly make improvements in potency with a

general method could have major clinical and economic implications.

In comparison to fourteen other promising machine learning-guided protein design

methods[8,16–28], we find that inverse folding has the strongest performance to date, even without

requiring any assay-labeled fitness data to use as training data for task-specific model supervision

(**Figure 5**, **Supplementary Data 5**). By eliminating the reliance on any initial data collection,

inverse folding has the potential to accelerate entire evolutionary campaigns.

Computational methods like the one we propose have the opportunity to democratize

protein engineering efforts. Not only is our approach more efficient than conventional resource-

intensive techniques that experimentally test the effects of all single-residue changes on

biochemical functions like binding affinity, but consequently it enables directed evolution based

on properties that are not easily measured at scale or that are incompatible with high-throughput

screening. Overcoming these limitations, we anticipate our structure-based paradigm will be

useful for evolving proteins across many domains.

Figure 1: Guiding evolution of diverse proteins via inverse folding

(A) The inverse folding problem refers to the prediction of a protein's native amino acid

sequence, given its three-dimensional backbone structure, which is conceptually analogous to the

opposite problem solved by structure prediction tools like AlphaFold[12]. **(B)** A hybrid

autoregressive model[11] integrates amino acid values and backbone structural information to

evaluate the joint likelihood over all positions in a sequence. Amino acids from the protein

sequence are tokenized (red), combined with geometric features extracted from a structural

encoder (green), and modeled with an encoder-decoder transformer (purple). Sequences assigned

high likelihoods by the model represent high confidence in folding into the input backbone

structure. **(C)** Our structure-guided framework for protein design indirectly explores the

underlying fitness landscape, without modeling a specific definition of fitness or requiring any

task-specific training data, by constraining the search space to regions where the backbone fold

preserved. **(D)** High fitness sensitivity analysis reveals that multimodal input improves language

model performance compared to sequence-only input across 10 proteins from diverse protein

families (left). 'Fraction High fitness' is the fraction of the top ten single amino acid substitutions

recommended by each model that are ranked in the top indicated percentile of all experimentally

screened variants. A representative plot (right) demonstrates this metric for assessing enrichment

of high-fitness MAPK1 mutations, with successfully predicted mutations highlighted (blue) on

the empirical cumulative density function (ECDF) of the experimental data (black). The three

different thresholds, as defined by percentiles, are also shown as dashed lines. Inverse folding

predictions are more enriched, on average, for high fitness variants across various tested

thresholds for high fitness classification. bla, Beta-lactamase TEM; CALM1, Calmodulin-1;

haeIIIM, Type II methyltransferase M.HaeIII; HRAS, GTPase HRas; MAPK1, Mitogen-

activated protein kinase; TMPT, Thiopurine S-methyltransferase; TPK1, Thiamin

pyrophosphokinase 1; UBI4, Polyubiquitin; UBE2I, SUMO-conjugating enzyme UBC9

Figure 2: Inverse folding of antibody-antigen complexes resolves mutational landscapes by implicitly learning features of binding and protein epistasis

(A) Spearman correlation using inverse folding as well as sequence-based modeling approaches ESM-v[49] and abYsis[55] reported for three antibodies screened with corresponding influenza A HA subtypes H1, H3, and H9. Bars are colored by the type of model used: IF, Inverse Folding (green); LM, Language Model (orange); and MSA, Multiple Sequence Alignment (purple). Inverse folding was evaluated in three different settings: i) providing the entire antibody variable region and antigen complex (Ab-Ag) ii) providing only the antibody variable region (Ab only), and iii) providing only the single antibody variable region of the chain responsible for binding or being mutated (Ab VH only or Ab VH/VL only). Inverse folding implicitly learns features of binding and protein epistasis. For example, when scoring combinatorial mutations to CR9114

against H1, we find that the model has much higher performance (Spearman $\rho = 0.65$ for H1, 0.5

for H3) than a masked language model ESM-1v (Spearman $\rho = 0.08$ for H1, 0.09 for H3) and a

site-independent, alignment-based model abYsis (Spearman $\rho = 0.08$ for H1, 0.1 for H3). This

performance improvement is also consistent across the other combinatorial landscapes tested. **(B)**

Scatter plots showing inverse folding predictions against experimentally determined dissociation

constants of CR6261 against HA-H1(left) and HA-H9 (right). The germline and mature

sequences are highlighted on all plots as indicated in the legend. For visualization, all scatter

plots omit points on the lower limit of quantitation. Further analysis of assay limit on predictive

performance is shown in **Supplementary Figure 2**. **(C)** Conceptual schematic representation of

protein language performance improvements with improved priors. Providing sequence and

structural information of both the antibody and antigen enables inverse folding to most

efficiently identify complex destabilizing mutations and enrich for high fitness antibody variants.

Figure 3. Inverse folding-guided evolution of antibodies improves neutralization potency

and resilience

(A) Each point represents the fold-change in IC50 of pseudovirus neutralization for antibody

variants with single amino acid mutations. Antibodies are tested against the viral strain

represented in the input structure (Ly1404- Wuhan, SA58-BA.1 Omicron). A dashed line is

shown at fold-change of 1 corresponding to no change. 35% of Ly-1404 variants and 30% of

SA58 variants improved antibody potency (defined as 1.1-fold or higher improvement in IC50

compared to wild-type). Among this subset of beneficial mutations, we identify single amino

acid mutations that provide a 1.6-fold improvement in Ly-1404 IC50 and a 2.6-fold

improvement in SA58 IC50. **(B)** Conceptual representation of viral evolution. Selection for

immune evasion drives antibody escape, which fundamentally represents a dynamic change in

the underlying fitness landscape for the antibody. This antigenic drift displaces a potent antibody

from a peak on the previous fitness landscape (left) to a new starting point at lower activity

(right). **(C)** Strip plots visualizing antibody evolution across two rounds. Each point shows the

corresponding fold-change in IC_{50} of pseudovirus neutralization for a designed variant and is

colored according to the number of mutations it has (1-8). Consistent with preserving backbone

fold, all 55 designed variants across both antibody evolutionary campaigns could be expressed.

All round 1 variants are only composed of only single amino acid changes while beneficial

mutations are combined in round 2. All round 2 variants have improved neutralization activity

compared to their respective wild-type antibody (dotted line). **(D)** Pseudovirus neutralization

curves are shown for the most potent evolved antibody variant, consisting of mutations annotated

to the left. The top Ly-1404 variant, bearing seven amino acid substitutions in VH, achieves a

26-fold improvement in neutralization against BQ.1.1 (top). The top SA58 variant, bearing single

amino acid mutations in both VH and VL, achieves an 11-fold improvement in neutralization

against BQ.1.1 (bottom). **(E)** Residues at which mutations improve neutralization against either

the structure-encoded strain, BQ.1.1, or both viral strains are highlighted with spheres for

antibodies Ly-1404 (PDB 7MMO) and SA58 (PDB 7Y0W). Notably, beneficial mutations are

identified both within the binding interface as well distal to the antigen. Neutralization enhancing

mutations are labeled in **Supplementary Figure 6**.

Figure 4: Antibodies evolved for high potency also exhibit improved affinity

(A) Ly-1404 antibody variants show a Spearman correlation of 0.47 between apparent affinity

fold-change and potency fold-change. Improved affinity is observed to be necessary but not

sufficient for improved neutralization activity. Four variants exhibit improved affinity but do not

enhance neutralization. All variants with improved neutralization also display improved affinity.

The top inverse folding Ly-1404 design with a 27-fold improvement in neutralization has a 9.5-

fold improvement in affinity to BQ.1.1 RBD, as measured using BLI. **(C)** SA58 antibodies

evolved for improved potency against BQ.1.1 also exhibit improved affinity against VOC

XBB.1.5, up to 37-fold. **(B, D)** Representative traces of BLI binding assays for Ly-1404 and

SA58 variants with improved affinity.

Figure 5: Comparison to other machine learning-guided directed evolution methods

'Fraction improved' refers to the hit rate of variants tested which are improved relative to a

wildtype protein used as a starting point for directed evolution or a reference protein used as a

design template. Higher hit rates indicate more efficient experimental exploration. Inverse

folding achieves the highest hit rate with the lowest number of assay-labeled training data points

to-date[8,16–28].

Methods

Inverse folding model description and scoring of sequences

As input to the inverse folding model, we provide a protein structure $\mathbf{Y} \in \mathbb{R}^{N \times 3 \times 3}$, where N is the number of amino acids, and each amino acid is featurized by the three-dimensional physical coordinates of all three atoms in the protein backbone: the α-carbon, β-carbon, and nitrogen atoms in the protein backbone (hence the dimensionality $N \times 3 \times 3$). The inverse folding model learns the probability distribution p of a protein sequence $\mathbf{x} = (x_1, \dots, x_N) \in \mathcal{X}^N$ (where $\mathcal{X}$ is the alphabet of amino acids) given a structure $\mathbf{Y}$ via the chain rule of probability

$$p(\mathbf{x}|\mathbf{Y}) = p(x_1|\mathbf{Y})p(x_2|x_1, \mathbf{Y}) \dots p(x_N|x_1, \dots, x_{N-1}, \mathbf{Y}).$$

The probability distribution at each position is defined over $\mathcal{X}$, such that it is a 20-dimensional vector with all constituent entries summing to 1.

Thus, for a given sequence $\hat{\mathbf{x}} = (\hat{x}_1, \dots, \hat{x}_N)$ and its corresponding given structure $\hat{\mathbf{Y}}$, we can score the probability of $\hat{\mathbf{x}}$ folding into $\mathbf{Y}$ under the inverse folding model by computing the value of $p(\mathbf{x} = \hat{\mathbf{x}}|\mathbf{Y})$, which we can do autoregressively as

$$p(\mathbf{x} = \hat{\mathbf{x}}|\hat{\mathbf{Y}}) = p(x_1 = \hat{x}_1|\hat{\mathbf{Y}}) \dots p(x_N = \hat{x}_N|\hat{x}_1, \dots, \hat{x}_{N-1}, \hat{\mathbf{Y}}).$$

This is evaluated output is a likelihood between 0 and 1, inclusive. The computed score $p(\mathbf{x} = \hat{\mathbf{x}}|\hat{\mathbf{Y}})$ is used as prediction for "fitness" (e.g., binding affinity or enzymatic activity). Importantly, the inverse folding model does not have any explicit access to "fitness" during either training or evaluation, which we refer to as "zero shot" fitness prediction.

We use the inverse folding model checkpoint of ESM-IF1 GVP-Transformer as of April 10, 2022[11].

Diverse proteins benchmarking experiment with scanning mutagenesis data

We analyzed the effectiveness of using the inverse folding language model, ESM-IF1

model to identify high fitness variants from protein mutational scans as a proxy for the ability to

guide evolution without explicitly modeling a protein's function. We also compared its

performance to ESM-1v, a sequence-only general protein language model. To do so, we used all

deep mutational scanning (DMS) datasets from the benchmarking study by Livesey and Marsh[29]

profiling over 100 variants and reported to have 90% or higher coverage of DMS results across

the corresponding curated PDB structure (**Supplementary Table 1**). From this set of 12

proteins, Cas9 was excluded because its sequence length was larger than the maximum allowable

length of 1024 amino acids by ESM-1v and ccdB was excluded because the experimental values

were discretized within a small range. For each of the 10 mutagenesis datasets, all the sequence

likelihood of all variants with coverage in the structure were determined using inverse folding.

For ESM-1v, the average masked marginals likelihood score across all five models in the ESM-

1v group was used. The experimental data distribution was binarized for high-fitness

classification using a percentile-based threshold. The enrichment of high fitness variants was

then determined by using the metric of fraction high fitness as defined by the fraction of the top

10 model-predicted variants with experimental values above the high fitness threshold. The

analysis was performed at three different percentile thresholds, top 5^{th} percentile (95^{th} percentile),

top 10^{th} percentile (90^{th} percentile), and top 20^{th} percentile (80^{th} percentile), to determine

sensitivity of the result based on the stringency of the selected cutoff parameter.

Benchmarking of antibody mutagenesis

We use three antibody mutagenesis datasets[50,51] to benchmark the performance of

modeling variant effects on antibody binding using inverse folding against two sequence-only

methods, ESM-1v[49] and abYsis[55]. Variant sequences were scored using the inverse folding model

with three different forms of structure input: i) variable region of mutated antibody chain only ii)

variable regions of both antibody chains iii) variable regions of both antibody chains in complex

with antigen. The autoregressive scoring of sequences with inverse folding enables evaluation of

sequences with multiple mutations. The Spearman correlation was determined between the log

likelihood scores across all sequences and corresponding reported experimental binding

measurements: $-\log(K_D)$ for CR9114 and CR6261; log(binding enrichment) g6.31. The following

structures were used for input backbone coordinates of the VH, VL, and antigen: PDB 4FQI[52],

CR9114-H5; PDB 3GBN[53], CR6261-H1; PDB 2FJG, g6.31-VEGF.

ESM-1v and abYsis were scored using the variant sequence of the antibody variable

region. For variants with multiple mutations, the average effect of all mutant amino acids in the

sequence was considered, namely

497
$$p(\mathbf{x}) = \frac{1}{|\mathcal{M}|} \sum_{i \in M} [\log p(\mathbf{x}_i = \mathbf{x}_i^{\mathrm{mt}}) - \log p(\mathbf{x}_i = \mathbf{x}_i^{\mathrm{wt}})]$$

where $\mathcal{M}$ is defined as the set of all mutations in the input sequence **x**. For abYsis, individual

mutation likelihoods were determined using the frequency of amino acids at each position based

on multiple sequence alignment provided by the webtool version 3.4.1

(http://www.abysis.org/abysis/index.html). We aligned VH and VL protein sequences using the

default settings provided in the 'Annotate' tool, with the database of 'Homo sapiens' sequences

as of April 1, 2023.

Acquisition of antibody amino acid substitutions using inverse folding

We select amino acid substitutions recommended by the inverse folding model to test in

our directed evolution campaigns for Ly-1404 and SA58. For a given wild-type antibody

variable region sequence, $\mathbf{x} = (x_1, \dots, x_N) \in \mathcal{X}^N$, where $\mathcal{X}$ is the set of amino acids and N is the

sequence length, we score all possible single amino acid substitutions against a corresponding

structure of the variable regions of both antibody chains in complex with the RBD of SARS-

CoV-2 Spike protein, $\hat{\mathbf{Y}}$ by computing $p(\mathbf{x} = \hat{\mathbf{x}} | \hat{\mathbf{Y}})$. Protein structures used are reported in

Supplementary Table 1. We then select the set of top ten predicted single amino acid

substitutions at unique residues in each antibody variable region to test in the first round of

evolution.

515   After testing individual amino acid mutations in a pseudovirus neutralization screen, in

Round 2, beneficial mutations (defined as IC_{50} fold-change > 1.1) were combined to assess the

combinatorial effects and potential for further neutralization improvement. We tested up to four

combinations of single amino acid mutations on each chain (two total mutations to the antibody).

We also used the inverse folding model to score a library of all possible combinations of the

beneficial mutations to an antibody chain (For example, VH Ly-1404 has 8 beneficial mutations

resulting in 255 total candidate sequences), and selected the top five scoring designs (or less if

there were a fewer number of total possible combinations). Lastly, we tested a maximum of two

variants consisting of the best single-chain designs together. In total, 31 variants were tested for

Ly-1404 and 25 variants were tested for SA58.

Antibody cloning

527   We cloned the antibody sequences into the CMV/R plasmid backbone for expression

under a CMV promoter. The heavy chain or light chain sequence was cloned between the CMV

promoter and the bGH poly(A) signal sequence of the CMV/R plasmid to facilitate improved

protein expression. Variable regions were cloned into the human IgG1 backbone; Ly-1404

variants were cloned with a lambda light chain, whereas variants of SA58 were cloned with a

kappa light chain. The vector for both heavy and light chain sequences also contained the

HVM06_Mouse (UniProt: P01750) Ig heavy chain V region 102 signal peptide

(MGWSCIILFLVATATGVHS) to allow for protein secretion and purification from the

supernatant. VH and VL segments were ordered as gene blocks from Integrated DNA

Technologies and were cloned into linearized CMV/R backbones with 5× In-Fusion HD Enzyme

Premix (Takara Bio).

Antigen cloning

RBD sequences were cloned into a pADD2 vector between the rBeta-globin intron and β-

globin poly(A). All RBD constructs contain an AviTag and 6×His tag. RBD sequences were

based off wild-type Wuhan-Hu-1 (GenBank: BCN86353.1), Omicron BA.1

(GenBank: UFO69279.1), BQ.1.1 (GenBank: OP412163.1), XBB.1.5 (GenBank: OP790748.1).

DNA preparation

Plasmids were transformed into Stellar competent cells (Takara Bio), and transformed

cells were plated and grown at 37 °C overnight. Colonies were mini-prepped per the

manufacturer's recommendations (GeneJET, K0502, Thermo Fisher Scientific) and sequence

confirmed (Sequetech) and then maxi-prepped per the manufacturer's protocols (ZymoPure II

Plasmid Maxiprep Kit, Zymo Research). Plasmids were sterile filtered using a 0.22-μm syringe

filter and stored at 4 °C.

Protein expression

All proteins were expressed in Expi293F cells (Thermo Fisher Scientific, A14527).

Proteins containing a biotinylation tag (AviTag) were also expressed in the presence of a BirA

enzyme, resulting in spontaneous biotinylation during protein expression. Expi293F cells were

cultured in media containing 66% FreeStyle/33% Expi media (Thermo Fisher Scientific) and

grown in TriForest polycarbonate shaking flasks at 37 °C in 8% carbon dioxide. The day before

transfection, cells were pelleted by centrifugation and resuspended to a density of 3×10^6 cells

per milliliter in fresh media. The next day, cells were diluted and transfected at a density of

approximately $3–4 \times 10^6$ cells per milliliter. Transfection mixtures were made by adding the

following components: maxi-prepped DNA, culture media and FectoPRO (Polyplus) would be

added to cells to a ratio of 0.5 µg: 100 µl: 1.3 µl: 900 µl. For example, for a 100-ml transfection,

50 µg of DNA would be added to 10 ml of culture media, followed by the addition of 130 µl of

FectoPRO. For antibodies, we divided the transfection DNA equally among heavy and light

chains; in the previous example, 25 µg of heavy chain DNA and 25 µg of light chain DNA would

be added to 10 ml of culture media. After mixing and a 10-min incubation, the example

transfection cocktail would be added to 90 ml of cells. The cells were harvested 3–5 days after

transfection by spinning the cultures at 10,000g for 10 min. Supernatants were filtered using a

0.45-µm filter.

Antibody purification

We purified antibodies using a 5-ml MabSelect Sure PRISM column on the ÄKTA pure

fast protein liquid chromatography (FPLC) instrument (Cytiva). The ÄKTA system was

equilibrated with line A1 in 20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid

(HEPES) pH 7.4, 150 mM sodium chloride (NaCl), line A2 in 100 mM glycine pH 2.8, line B1

in 0.5 M sodium hydroxide, Buffer line in 20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic

acid (HEPES) pH 7.4, 150 mM sodium chloride (NaCl) and Sample lines in water. The protocol

washes the column with A1, followed by loading of the sample in the Sample line until air is

detected in the air sensor of the sample pumps, followed by five column volume washes with A1,

elution of the sample by flowing of 20 ml of A2 directly into a 50-ml conical containing 2 ml of

1 M tris(hydroxymethyl)aminomethane (Tris) pH 8.0, followed by five column volumes of A1,

B1 and A1 and then a wash step of the fraction collector with A1. We concentrated the eluted

samples using 50-kDa cutoff centrifugal concentrators, followed by buffer exchange using a PD-

10 column (Sephadex) that had been pre-equilibrated into 20 mM 4-(2-hydroxyethyl)-1-

piperazineethanesulfonic acid (HEPES) pH 7.4, 150 mM sodium chloride (NaCl). Purified

antibodies were used directly in experiments or flash-frozen and stored at −20 °C.

Antigen purification

All RBD antigens were His-tagged and purified using HisPur Ni-NTA resin (Thermo

Fisher Scientific, 88222). Cell supernatants were diluted with 1/3 volume of wash buffer (20 mM

imidazole, 20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) pH 7.4,

150 mM sodium chloride (NaCl), and the Ni-NTA resin was added to diluted cell supernatants.

For all antigens, the samples were then incubated at 4 °C while stirring overnight.

Resin/supernatant mixtures were added to chromatography columns for gravity flow purification.

The resin in the column was washed with wash buffer (20 mM imidazole, 20 mM HEPES pH

7.4, 150 mM NaCl), and the proteins were eluted with 250 mM imidazole, 20 mM HEPES pH

7.4, 150 mM NaCl. Column elutions were concentrated using centrifugal concentrators at 10-

kDa cutoff, followed by size-exclusion chromatography on an ÄKTA pure system (Cytiva).

ÄKTA pure FPLC with a Superdex 200 Increase (S200) gel filtration column was used for

purification. Then, 1 ml of sample was injected using a 2-ml loop and run over the S200, which

had been pre-equilibrated in degassed 20 mM HEPES, 150 mM NaCl before use and flash-frozen

before storage at −20 °C.

BLI binding experiments

All reactions were run on an Octet RED96 at 30 °C, and samples were run in 1× PBS

with 0.1% BSA and 0.05% Tween 20 (Octet buffer). IgGs were assessed for binding to

biotinylated antigens using streptavidin biosensors (Sartorius/ForteBio). Antigen was loaded at a

concentration of 200nM. Tips were then washed and baselined in wells containing only Octet

buffer. Samples were then associated in wells containing IgG at 100 nM concentration. A control

well with loaded antigen but that was associated in a well containing only 200 μl of Octet buffer

was used as a baseline subtraction for data analysis. Association and dissociation binding curves

were fit in Octet System Data Analysis Software version 9.0.0.15 using a 1:2 bivalent model for

IgGs to determine apparent K_d. Fold-change in apparent K_d were determined by computing the

ratio of wildtype K_d to variant K_d. Averages of K_d fold-change values from at least two

independent experiments are reported to two significant figures in **Supplementary Data 2**. To

estimate measurement error, we computed the standard deviation for each

antibody−antigen K_d pair.

Polyspecificity Particle assay

Polyspecificity reagent (PSR) was obtained as described by Xu et al[63]. Soluble membrane

proteins were isolated from homogenized and sonicated Expi 293F cells followed by

biotinylation with Sulfo-NHC-SS-Biotin (Thermo Fisher Scientific, 21331) and stored in PBS at

−80 °C. The PolySpecificity Particle (PSP) assay was performed as described in Makowski et

al.[64]. Protein A magnetic beads (Invitrogen, 10001D) were washed three times in PBSB (PBS

with 1 mg ml^{-1} BSA) and diluted to 54 µg ml^{-1} in PBSB. Then, 30 µl of the solution containing

the beads was incubated with 85 µl of antibodies at 15 μg ml^{-1} overnight at 4 °C with rocking.

The coated beads were then washed twice with PBSB using a magnetic plate stand (Invitrogen,

12027) and resuspended in PBSB. We then incubated 50 µl of 0.1 mg ml^{-1} PSR with the washed

beads at 4 °C with rocking for 20 min. Beads were then washed with PBSB and incubated with

0.001× streptavidin-APC (BioLegend, 405207) and 0.001× goat anti-human Fab fragment FITC

(Jackson ImmunoResearch, 109-097-003) at 4 °C with rocking for 15 min. Beads were then

washed and resuspended with PBSB. Beads were profiled via flow cytometry using a Sony

SH800 cell sorter. Data analysis was performed with FlowJo software version 10.9.0 to obtain

median fluorescence intensity (MFI) values, which are reported for each antibody across three or

more replicate wells. Elotuzumab (Fisher Scientific) and ixekizumab (Fisher Scientific) are also

included in each assay as controls.

Lentivirus production

We produced SARS-CoV-2 Spike (Wuhan-Hu-1, BA.1, and BQ.1.1 variants)

pseudotyped lentiviral particles. Viral transfections were done in HEK293T cells (American

Type Culture Collection, CRL-3216) using BioT (BioLand) transfection reagent. Six million

cells were seeded in D10 media (DMEM + additives: 10% FBS, L-glutamate, penicillin,

streptomycin and 10 mM HEPES) in 10-cm plates one day before transfection. A five-plasmid

system was used for viral production, as described in Crawford et al[65]. The Spike vector

contained the 21-amino-acid truncated form of the SARS-CoV-2 Spike sequence from the

Wuhan-Hu-1 strain of SARS-CoV-2 (GenBank: BCN86353.1), BA.1 variant of concern

(GenBank: OL672836.1), or BQ.1.1 variant of concern (GenBank: OP412163.1. The other viral

plasmids, used as previously described[65], are pHAGE-Luc2-IRS-ZsGreen (NR-52516), HDM-

Hgpm2 (NR-52517), pRC-CMV-Rev1b (NR-52519) and HDM-tat1b (NR-52518). These

plasmids were added to D10 medium in the following ratios: 10 µg pHAGE-Luc2-IRS-ZsGreen,

3.4 µg FL Spike, 2.2 µg HDM-Hgpm2, 2.2 µg HDM-Tat1b and 2.2 µg pRC-CMV-Rev1b in a

final volume of 1,000 µl.

After adding plasmids to medium, we added 30 µl of BioT to form transfection

complexes. Transfection reactions were incubated for 10 min at room temperature, and then 9 ml

of medium was added slowly. The resultant 10 ml was added to plated HEK cells from which the

medium had been removed. Culture medium was removed 24 h after transfection and replaced

with fresh D10 medium. Viral supernatants were harvested 72 h after transfection by spinning at

300g for 5 min, followed by filtering through a 0.45-µm filter. Viral stocks were aliquoted and

stored at −80 °C.

Pseudovirus neutralization

The target cells used for infection in SARS-CoV-2 pseudovirus neutralization assays are

from a HeLa cell line stably overexpressing human angiotensin-converting enzyme 2 (ACE2) as

well as the protease known to process SARS-CoV-2: transmembrane serine protease 2

(TMPRSS2). Production of this cell line is described in detail by Rogers et al[66]. with the addition

of stable TMPRSS2 incorporation. ACE2/TMPRSS2/HeLa cells were plated 1 day before

infection at 8,000 cells per well. Ninety-six-well, white-walled, white-bottom plates were used

for neutralization assays (Thermo Fisher Scientific).

On the day of the assay, purified IgGs in 1× PBS were made into D10 medium (DMEM +

additives: 10% FBS, L-glutamate, penicillin, streptomycin and 10 mM HEPES). A virus mixture

was made containing the virus of interest (for example, SARS-CoV-2) and D10 media. Virus

dilutions into media were selected such that a suitable signal would be obtained in the virus-only

wells. A suitable signal was selected such that the virus-only wells would achieve a

luminescence of at least >1,000,000 relative light units (RLU). Then, 60 μl of this virus mixture

was added to each of the antibody dilutions to make a final volume of 120 μl in each well. Virus-

only wells were made, which contained 60 μl of D10 and 60 μl of virus mixture. Cells-only wells

were made, which contained 120 μl of D10 media.

The antibody/virus mixture was left to incubate for 1 h at 37 °C. After incubation, the

medium was removed from the cells on the plates made one day prior. This was replaced with

100 μl of antibody/virus dilutions and incubated at 37 °C for approximately 48 h. Infectivity

readout was performed by measuring luciferase levels. Medium was removed from all wells, and

cells were lysed by the addition of 100 μl of BriteLite assay readout solution (PerkinElmer) into

each well. Luminescence values were measured using an Infinite 200 PRO Microplate Reader

(Tecan) using i-control version 2.0 software (Tecan) after shaking for 30 sec. Each plate was

normalized by averaging the cells-only (0% infection) and virus-only (100% infection)

wells. Neutralization titer was defined as the sample dilution at which the RLU was decreased by

50% as compared with the RLU of virus-only control wells after subtraction of background

RLUs in wells containing cells only. Normalized values were fitted with a three-parameter

nonlinear regression inhibitor curve in GraphPad Prism 9.1.0 to determine the half-maximal

inhibitory concentration (IC_{50}) and are reported in **Supplementary Data 1**. Neutralization assays

were performed in biological duplicates with technical duplicates.

Computing frequency of changes to antibody protein sequences

We computed the frequency of residues involved in affinity-enhancing substitutions using

the abYsis webtool, which also computes the frequency of amino acids at each position based on

a multiple sequence alignment. We aligned VH and VL protein sequences using the default

settings provided in the 'Annotate' tool, using the database of 'All' sequences as of April 1,

2023. We also used the Kabat region definition provided by abYsis webtool version 3.4.1 to

annotate the framework regions and CDRs within the VH and VL sequences which are reported

in **Supplementary Table 2**.

Comparing efficiency of machine learning-guided directed evolution methods

To compare inverse folding against other machine learning methods for protein

evolution, we compared the fraction of variants tested in the protein engineering campaign to the

number of assay-labeled training data points used to inform the predictions. Data was sourced

from Biswas et al.[17] and made contemporaneous by the addition of recently published studies as

indicated in **Supplementary Data 5**. The fraction improved, or hit rate, refers to experimentally

tested predictions which have improved functional activity relative to either a wildtype protein

that is used as a starting point for directed evolution or the protein used as a reference template

for design.

Acknowledgments

We would like to thank D. Xu, S. Kim, and the members of the Kim lab for helpful discussions

on this project. We are also grateful for assistance from D. Xu with protein graphics. V.R.S

acknowledges the support of the Stanford University Medical Scientist Training Program grants

(T32-GM007365 and T32-GM145402). V.R.S. and T.U.J.B. are both supported by the Sarafan

ChEM-H Chemistry/Biology Interface Training Program. T.U.J.B. is also supported by the

Knight-Hennessy Graduate Scholarship Fund and a CIHR Doctoral Foreign Study Award

(FRN:170770). B.L.H is supported by the Stanford Science Fellows Program. This work was

supported by the Virginia & D.K. Ludwig Fund for Cancer Research and the Chan Zuckerberg

Biohub.

Author contributions

Conceptualization, methodology, interpretation: V.R.S., B.L.H., P.S.K.; Computational

experiments and software development: V.R.S.; Antibody and antigen cloning, expression, and

purification: V.R.S., T.U.J.B.; Lentivirus production and pseudovirus neutralization: T.U.J.B;

Binding assays: V.R.S.; Writing (original draft): V.R.S with assistance from B.L.H and P.S.K.;

Writing (final draft): all authors

Competing interests

V.R.S., B.L.H., and P.S.K. are named as inventors on a patent application applied for by

Stanford University and the Chan Zuckerberg Biohub entitled "Antibody Compositions and

Optimization Methods".

References

1. Chothia, C. & Lesk, A. M. The relation between the divergence of sequence and structure in proteins. *EMBO J.* **5**, 823–826 (1986).

2. Bloom, J. D., Labthavikul, S. T., Otey, C. R. & Arnold, F. H. Protein stability promotes evolvability. *Proc. Natl. Acad. Sci.* **103**, 5869–5874 (2006).

3. Axe, D. D., Foster, N. W. & Fersht, A. R. A Search for Single Substitutions That Eliminate Enzymatic Function in a Bacterial Ribonuclease. *Biochemistry* **37**, 7157–7166 (1998).

4. Romero, P. A. & Arnold, F. H. Exploring protein fitness landscapes by directed evolution. *Nat. Rev. Mol. Cell Biol.* **10**, 866–876 (2009).

5. Shafikhani, S., Siegel, R. A., Ferrari, E. & Schellenberger, V. Generation of large libraries of random mutants in Bacillus subtilis by PCR-based plasmid multimerization. *BioTechniques* **23**, 304–310 (1997).

6. Guo, H. H., Choe, J. & Loeb, L. A. Protein tolerance to random amino acid change. *Proc. Natl. Acad. Sci.* **101**, 9205–9210 (2004).

7. Rennell, D., Bouvier, S. E., Hardy, L. W. & Poteete, A. R. Systematic mutation of bacteriophage T4 lysozyme. *J. Mol. Biol.* **222**, 67–88 (1991).

8. Ogden, P. J., Kelsic, E. D., Sinai, S. & Church, G. M. Comprehensive AAV capsid fitness landscape reveals a viral gene and enables machine-guided design. *Science* **366**, 1139–1143 (2019).

9. Poelwijk, F. J., Kiviet, D. J., Weinreich, D. M. & Tans, S. J. Empirical fitness landscapes reveal accessible evolutionary paths. *Nature* **445**, 383–386 (2007).

10. Wittmann, B. J., Yue, Y. & Arnold, F. H. Informed training set design enables efficient machine learning-assisted directed protein evolution. *Cell Syst.* **12**, 1026-1045.e7 (2021).

11. Hsu, C. *et al*. Learning inverse folding from millions of predicted structures. in *Proceedings*

of the 39th International Conference on Machine Learning 8946–8970 (PMLR, 2022).

12. Jumper, J. *et al*. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583–589 (2021).

13. Lin, Z. *et al*. Evolutionary-scale prediction of atomic-level protein structure with a language

model. *Science* **379**, 1123–1130 (2023).

14. Carter, P. J. & Lazar, G. A. Next generation antibody drugs: pursuit of the 'high-hanging

fruit'. *Nat. Rev. Drug Discov.* **17**, 197–223 (2018).

15. Schroeder, H. W. & Cavacini, L. Structure and function of immunoglobulins. *J. Allergy Clin.*

Immunol. **125**, S41–S52 (2010).

16. Makowski, E. K. *et al*. Co-optimization of therapeutic antibody affinity and specificity using

machine learning models that generalize to novel mutational space. *Nat. Commun.* **13**, 3788

(2022).

17. Biswas, S., Khimulya, G., Alley, E. C., Esvelt, K. M. & Church, G. M. Low-N protein

engineering with data-efficient deep learning. *Nat. Methods* **18**, 389–396 (2021).

18. Bedbrook, C. N. *et al*. Machine learning-guided channelrhodopsin engineering enables

minimally invasive optogenetics. *Nat. Methods* **16**, 1176–1184 (2019).

19. Wu, Z., Kan, S. B. J., Lewis, R. D., Wittmann, B. J. & Arnold, F. H. Machine learning-

assisted directed protein evolution with combinatorial libraries. *Proc. Natl. Acad. Sci.* **116**,

8852–8858 (2019).

20. Cadet, F. *et al*. A machine learning approach for reliable prediction of amino acid

interactions and its application in the directed evolution of enantioselective enzymes. *Sci.*

Rep. **8**, 16757 (2018).

21. Saito, Y. *et al*. Machine-Learning-Guided Mutagenesis for Directed Evolution of Fluorescent

Proteins. *ACS Synth. Biol.* **7**, 2014–2022 (2018).

22. Biswas, S. *et al*. Toward machine-guided design of proteins. 337154 Preprint at

https://doi.org/10.1101/337154 (2018).

23. Romero, P. A., Krause, A. & Arnold, F. H. Navigating the protein fitness landscape with

Gaussian processes. *Proc. Natl. Acad. Sci.* **110**, E193–E201 (2013).

24. Bedbrook, C. N., Yang, K. K., Rice, A. J., Gradinaru, V. & Arnold, F. H. Machine learning

to design integral membrane channelrhodopsins for efficient eukaryotic expression and

plasma membrane localization. *PLOS Comput. Biol.* **13**, e1005786 (2017).

25. Liao, J. *et al*. Engineering proteinase K using machine learning and synthetic genes. *BMC*

Biotechnol. **7**, 16 (2007).

26. Repecka, D. *et al*. Expanding functional protein sequence spaces using generative adversarial

networks. *Nat. Mach. Intell.* **3**, 324–333 (2021).

27. Shin, J.-E. *et al*. Protein design and variant prediction using autoregressive generative

models. *Nat. Commun.* **12**, 2403 (2021).

28. Hie, B. L. *et al*. Efficient evolution of human antibodies from general protein language

models. *Nat. Biotechnol.* 1–9 (2023) doi:10.1038/s41587-023-01763-2.

29. Livesey, B. J. & Marsh, J. A. Using deep mutational scanning to benchmark variant effect

predictors and identify disease mutations. *Mol. Syst. Biol.* **16**, e9380 (2020).

30. Weile, J. *et al*. A framework for exhaustively mapping functional missense variants. *Mol.*

Syst. Biol. **13**, 957 (2017).

31. Bandaru, P. *et al*. Deconstruction of the Ras switching cycle through saturation mutagenesis.

eLife **6**, e27810 (2017).

32. Brenan, L. *et al*. Phenotypic Characterization of a Comprehensive Set of MAPK1/ERK2

Missense Mutants. *Cell Rep*. **17**, 1171–1183 (2016).

33. Matreyek, K. A. *et al*. Multiplex assessment of protein variant abundance by massively

parallel sequencing. *Nat. Genet*. **50**, 874–882 (2018).

34. Mishra, P., Flynn, J. M., Starr, T. N. & Bolon, D. N. A. Systematic Mutant Analyses

Elucidate General and Client-Specific Aspects of Hsp90 Function. *Cell Rep*. **15**, 588–598

(2016).

35. Roscoe, B. P. & Bolon, D. N. A. Systematic exploration of ubiquitin sequence, E1 activation

efficiency, and experimental fitness in yeast. *J. Mol. Biol*. **426**, 2854–2870 (2014).

36. Kitzman, J. O., Starita, L. M., Lo, R. S., Fields, S. & Shendure, J. Massively parallel single-

amino-acid mutagenesis. *Nat. Methods* **12**, 203–206, 4 p following 206 (2015).

37. Stiffler, M. A., Hekstra, D. R. & Ranganathan, R. Evolvability as a Function of Purifying

Selection in TEM-1 β-Lactamase. *Cell* **160**, 882–892 (2015).

38. Rockah-Shmuel, L., Tóth-Petróczy, Á. & Tawfik, D. S. Systematic Mapping of Protein

Mutational Space by Prolonged Drift Reveals the Deleterious Effects of Seemingly Neutral

Mutations. *PLOS Comput. Biol*. **11**, e1004421 (2015).

39. Hewitt, W. M. *et al*. Insights Into the Allosteric Inhibition of the SUMO E2 Enzyme Ubc9.

Angew. Chem. Int. Ed. **55**, 5703–5707 (2016).

40. Cho, L. T.-Y. *et al*. An Intracellular Allosteric Modulator Binding Pocket in SK2 Ion

Channels Is Shared by Multiple Chemotypes. *Structure* **26**, 533-544.e3 (2018).

41. Klink, B. U., Goody, R. S. & Scheidig, A. J. A newly designed microspectrofluorometer for

kinetic studies on protein crystals in combination with x-ray diffraction. *Biophys. J*. **91**, 981–

992 (2006).

42. Ward, R. A. *et al*. Structure-Guided Design of Highly Selective and Potent Covalent

Inhibitors of ERK1/2. *J. Med. Chem.* **58**, 4790–4801 (2015).

43. Wu, H. *et al*. Structural basis of allele variation of human thiopurine-S-methyltransferase.

Proteins Struct. Funct. Bioinforma. **67**, 198–208 (2007).

44. Meyer, P. *et al*. Structural basis for recruitment of the ATPase activator Aha1 to the Hsp90

chaperone machinery. *EMBO J.* **23**, 511–519 (2004).

45. Grishin, A. M. *et al*. Structural Basis for the Inhibition of Host Protein Ubiquitination by

Shigella Effector Kinase OspG. *Structure* **22**, 878–888 (2014).

46. Hong, M. *et al*. Structural Basis for Dimerization in DNA Recognition by Gal4. *Structure* **16**,

1019–1026 (2008).

47. Minasov, G., Wang, X. & Shoichet, B. K. An Ultrahigh Resolution Structure of TEM-1 β-

Lactamase Suggests a Role for Glu166 as the General Base in Acylation. *J. Am. Chem. Soc.*

124, 5333–5340 (2002).

48. Didovyk, A. & Verdine, G. L. Structural Origins of DNA Target Selection and Nucleobase

Extrusion by a DNA Cytosine Methyltransferase *. *J. Biol. Chem.* **287**, 40099–40105 (2012).

49. Meier, J. *et al*. Language models enable zero-shot prediction of the effects of mutations on

protein function. 2021.07.09.450648 Preprint at https://doi.org/10.1101/2021.07.09.450648

(2021).

50. Phillips, A. M. *et al*. Binding affinity landscapes constrain the evolution of broadly

neutralizing anti-influenza antibodies. *eLife* **10**, e71393 (2021).

51. Koenig, P. *et al*. Mutational landscape of antibody variable domains reveals a switch

modulating the interdomain conformational dynamics and antigen binding. *Proc. Natl. Acad.*

Sci. **114**, E486–E495 (2017).

52. Dreyfus, C. *et al*. Highly Conserved Protective Epitopes on Influenza B Viruses. *Science*

337, 1343–1348 (2012).

53. Ekiert, D. C. *et al*. Antibody Recognition of a Highly Conserved Influenza Virus Epitope.

Science **324**, 246–251 (2009).

54. Fuh, G. *et al*. Structure-Function Studies of Two Synthetic Anti-vascular Endothelial Growth

Factor Fabs and Comparison with the Avastin™ Fab *. *J. Biol. Chem.* **281**, 6625–6631

(2006).

55. Swindells, M. B. *et al*. abYsis: Integrated Antibody Sequence and Structure—Management,

Analysis, and Prediction. *J. Mol. Biol.* **429**, 356–364 (2017).

56. Westendorf, K. *et al*. LY-CoV1404 (bebtelovimab) potently neutralizes SARS-CoV-2

variants. *Cell Rep.* **39**, 110812 (2022).

57. Takashita, E. *et al*. Efficacy of Antibodies and Antiviral Drugs against Omicron BA.2.12.1,

BA.4, and BA.5 Subvariants. *N. Engl. J. Med.* **387**, 468–470 (2022).

58. Research, C. for D. E. and. FDA Announces Bebtelovimab is Not Currently Authorized in

Any US Region. *FDA* (2022).

59. Cao, Y. *et al*. BA.2.12.1, BA.4 and BA.5 escape antibodies elicited by Omicron infection.

Nature **608**, 593–602 (2022).

60. Cao, Y. *et al*. Rational identification of potent and broad sarbecovirus-neutralizing antibody

cocktails from SARS convalescents. *Cell Rep.* **41**, (2022).

61. Song, R. *et al*. Post-exposure prophylaxis with SA58 (anti-SARS-COV-2 monoclonal

antibody) nasal spray for the prevention of symptomatic COVID-19 in healthy adult

workers: a randomized, single-blind, placebo-controlled clinical study*. *Emerg. Microbes*

Infect. **12**, 2212806 (2023).

62. Starr, T. N. *et al*. Shifting mutational constraints in the SARS-CoV-2 receptor-binding

domain during viral evolution. *Science* **377**, 420–424 (2022).

63. Xu, Y. *et al*. Addressing polyspecificity of antibodies selected from an in vitro yeast

presentation system: a FACS-based, high-throughput selection and analytical tool. *Protein*

Eng. Des. Sel. **26**, 663–670 (2013).

64. Makowski, E. K., Wu, L., Desai, A. A. & Tessier, P. M. Highly sensitive detection of

antibody nonspecific interactions using flow cytometry. *mAbs* **13**, 1951426 (2021).

65. Crawford, K. H. D. *et al*. Protocol and Reagents for Pseudotyping Lentiviral Particles with

SARS-CoV-2 Spike Protein for Neutralization Assays. *Viruses* **12**, 513 (2020).

66. Rogers, T. F. *et al*. Isolation of potent SARS-CoV-2 neutralizing antibodies and protection

from disease in a small animal model. *Science* **369**, 956–963 (2020).

Supplementary Figures, Tables, Information, & Data

Supplementary Table 1: List of proteins, protein structures, and assay information for deep

mutational scanning experiments

Supplementary Table 2: Analysis of neutralization-enhancing mutations

Supplementary Information: Antibody sequences

Supplementary Data 1: Neutralization data with IC_{50} values of evolved antibodies across both

evolutionary campaigns

Supplementary Data 2: Binding data with IgG K_D values of evolved antibodies

Supplementary Data 3: Antibody variant prediction benchmarking results

Supplementary Data 4: MFI values for polyspecificity experiments

Supplementary Data 5: Efficiency comparison of machine learning-guided directed evolution

methods

899
900

Supplementary Figure 1: Inverse folding identifies high fitness variants across proteins

with diverse functions

In addition to higher hit rates of high fitness variants, inverse folding generally identifies variants

with greater magnitude of improvements in fitness. The top ten predicted variants with

experimental fitness values ranking in the 20th percentile of all variants profiled in the deep

mutational screen are shown. The grey curve shows the empirical cumulative distribution

function (ECDF) of all experimental fitness values determined in the screen. The dotted lines

correspond to the three percentile-based thresholds used in the sensitivity analysis (**Figure 1d**) to

classify high fitness variants. bla, Beta-lactamase TEM; CALM1, Calmodulin-1; haeIIIM, Type

II methyltransferase M.HaeIII; HRAS, GTPase HRas; MAPK1, Mitogen-activated protein

912   kinase; TMPT, Thiopurine S-methyltransferase; TPK1, Thiamin pyrophosphokinase 1; UBI4,

913   Polyubiquitin; UBE2I, SUMO-conjugating enzyme UBC9

Supplementary Figure 2: Impact of lower limit of quantitation of binding assay on

predictive performance

(**A**) Scatter plots showing CR6261 variant sequences scored with inverse folding compared to

experimental binding data and inclusive of the assay's lower limit of quantitation, which is

918   omitted for visualization in **Figure 3b**. **(B)** Comparative bar plots showing the impact of

919   removing sequences with experimental measurements bounded artificially by the assay to

920   dataset-wide correlation. While Spearman correlations shown in Figure 3a are computed without

921   any modification to the data, trends in prediction and comparison among modeling methods are

922   robust to filtering sequences affected by this assay artifact.

HA of H5N1 influenza, PDB 4FQI
HA of H1N1 infleunza, PDB 7SCO

Backbone RMSD =2.1Å (1416 atoms)
Hamming Distance = 183/499 Amino Acids

Supplementary Figure 3: Structural and sequence similarity of H5 and H1

For cross-reactive antibodies, inclusion of the antigen structure is informative even for predicting

binding to a different antigen. In Figure 3a, we report a correlation of 0.65 between inverse

folding log likelihoods of CR9114 variants and experimental affinity measurements to H1

despite using a structure solved with CR9114 in complex with H5. Inverse folding uses both the

protein sequence and backbone structure coordinates as input. Across both HA subunits, H5 and

H1 have considerable sequence differences and a 2.1 Å root mean square deviation (RMSD)

across the entire protein backbone.

Supplementary Figure 4: Functional diversity of inverse folding-recommended mutations

Among the 20 single amino acid substitutions tested for Ly-1404, 14 of 20 = 70% improve

neutralization against at least one of the two strains tested. Similarly, 7 of 20 = 35% of the single

amino acid substitutions tested for SA58 improve neutralization. While some variants improve

function against both pseudovirus strains, others overwhelmingly only improve against one. This

suggests that focusing sequence exploration to structurally compatible mutations does not

compromise functional diversity.

Supplementary Figure 5: Polyspecificity of evolved antibodies

(A) The median fluorescence intensity (MFI) signal obtained from flow cytometry is shown for

several evolved antibodies with improved affinity and compared to two clinical monoclonal

945 antibodies with high and low polyspecificity used to define a clinically viable range. **(B)** Fold-

946 change in polyspecificity signal is plotted against fold-change in affinity as IgG against BQ.1.1

947 for Ly-1404 and XBB.1.5 for SA58. There is no correlation between the improvements in on-

948 target improvements in affinity and off-target nonspecific changes in polyspecificity (Spearman ρ

949 $= 0.007$).

Supplementary Figure 6: Mapping neutralization-enhancing substitutions

Neutralization-enhancing mutations are labeled on the structure of the wild-type antibody in

complex with the RBD of SARS-CoV-2 spike protein (Ly-1404: PDB 7mmo; SA58: PDB

7y0w). Notably, several mutations are identified to have significant beneficial impacts on binding

neutralization and affinity (**Supplementary Data 1 & 2**) despite located away from the binding

interface.

Supplementary Table 1. Summary of the DMS datasets used in this analysis, including functional assay, method of mutagenesis, and structure used for inverse folding scoring. We also note the specific DMS assay from each study we use for calculating correlation with inverse folding log likelihoods.

Protein(s) (Uniprot ID)	Organism	Functional Assay	Mutagenesis Method	Utilized assay	PDB Structure	Total coverage of DMS (%)	Access date*	Reference
UBE2I (P63279)	Human	POPCode, a variant of multiple-site directed mutagenesis.	Competitive growth assay in yeast.	score	5F6E chain A	100	12/10/2018	(Weile *et al*, 2017)
TPK1 (Q9H3S3)				score	3S4Y chain A	92.46		
CALM1 (P0DP23)				score	5V03 chain R	100		
HRas (P01112)	Human	Systematic site-directed mutagenesis.	Two-hybrid assay.	unregulated	2CE2 chain X	100	12/10/2018	(Bandaru *et al*, 2017)
MAPK1 (P28482)	Human	Systematic site-directed mutagenesis.	Competitive growth assay.	VRT	4ZZN chain A	99.44	12/10/2018	(Brenan *et al*, 2016)
TPMT (P51580)	Human	Systematic site-directed mutagenesis.	Fluorescence of a GFP fusion protein.	score	2BZG chain A	92.9	12/10/2018	(Matreyek *et al*, 2018)
UBI4(b) (P0CG63)	Yeast	Site directed mutagenesis by cassette ligation.	Fluorescence activated cell sorting (FACS).	Relative_E1-activity_limiting	4Q5E chain B	100	12/10/2018	(Roscoe & Bolon, 2014)
GAL4 (P04386)	Yeast	Systematic site-directed mutagenesis.	Two-hybrid assay.	Nonselection_24	3COQ chain B	90.64	12/10/2018	(Kitzman *et al*, 2015)
bla(b) (P62593)	E. coli	Systematic site-directed mutagenesis.	Antibiotic resistance.	Ampicillin_2500	1M40 chain A	100	12/10/2018	(Stiffler *et al*, 2015)
haeIIIM (P20589)	H. aegyptius	Random mutagenesis.	Competitive growth assay.	DMS_G3	3UBT chain B	99.37	12/10/2018	(Rockah-Shmuel *et al*, 2015)

*Access date is as reported in *Livesey & Marsh*, 2020 study from which these data were sourced and this table was adapted

Supplementary Table 2. Single amino acid substitutions with beneficial effects on neutralization are reported alongside the region of the variable domain they are located within, as well as the wild-type and mutant amino acid frequencies in observed human antibody sequences.

Ly-1404

Chain Mutated	Design	Region	WT Amino Acid Frequency	Mutant Amino Acid Frequency
HC	D88Q	HFR3	0.03333	0.00382
HC	V90S	HFR3	0.03316	0.05155
HC	S62N	CDR-H2	0.13159	0.16299
HC	V81T	HFR3	0.03432	0.00205
HC	F24Y	HFR1	0.01738	0.00002
HC	I31T	CDR-H1	0.00933	0.09048
HC	H99Y	HFR3	0.01593	0.00138
HC	T70S	HFR3	0.88405	0.06153
HC	I105L	CDR-H3	0.02764	0.05760
LC	A98I	CDR-L3	0.02297	0.03198
LC	Q39K	LFR2	0.92316	0.00238
LC	T5Q	LFR1	0.89340	0.00933
LC	K47E	LFR2	0.52285	0.01490
LC	M49L	LFR2	0.05585	0.77076

SA58

Chain Mutated	Design	Region	WT Amino Acid Frequency	Mutant Amino Acid Frequency
HC	T53L	CDR-H2	0.03814	0.00963
HC	A61S	CDR-H2	0.59797	0.13159
HC	E10Q	HFR1	0.24182	0.01366
LC	N95V	CDR-L3	0.13399	0.00685
LC	S85A	LFR3	0.01109	0.00698
LC	S54T	CDR-L2	0.65138	0.05372
LC	M4V	LFR1	0.29424	0.03348