Olfactory ensheathing cells from adult female rats are hybrid glia that promote neural repair

Patricia E. Phelps*, Sung Min Ha, Rana R. Khankan, Mahlet A. Mekonnen, Giovanni Juarez, Kaitlin L. Ingraham Dixie, Yen-Wei Chen and Xia Yang*

Department of Integrative Biology and Physiology
UCLA, Los Angeles, CA 90095

*Corresponding author:
Patricia E. Phelps, Ph.D.
UCLA, Dept. of Integrative Biology and Physiology
Terasaki Life Sciences Building
610 Charles Young Dr. East
Los Angeles, CA 90095-7239
Telephone: (310) 825-7264
Email: pphelps@physci.ucla.edu

*Corresponding author:
Xia Yang, PhD.
UCLA, Dept. of Integrative Biology and Physiology
Terasaki Life Sciences Building
610 Charles Young Dr. East
Los Angeles, CA 90095-7239
Email: xyang123@ucla.edu

Abbreviated title: OECs are hybrid glia
Number of pages: 52
Number of figures: 9
Number of Supplemental figures: 3
Number of Supplemental tables: 4
Key words: scRNA-seq, OECs, Reelin (*Reln*), Connective tissue growth factor (*Ctgf/Ccn2*), neural repair

Acknowledgments: This work was supported by the National Institute of Neurological Disorders and Stroke: RO1 NS076976 (PEP), R01 NS111378 (XY), R01 NS117148 (XY), and by NICHD of the National Institutes of Health under award number P50HD103557. Additional support came from the UCLA QCBio Collaboratory Postdoc Fellowship (SMH), Eureka fellowship (RRK), MARC U*STAR fellowship T34 GM008563 (GJ), and UCLA Graduate Year Fellowship (KLID). We thank Drs. Julie Miller and Stephanie White for their generosity and considerable assistance with western blots, Drs. Hui Zhong and VR Edgerton for their long-term collaboration on spinal cord injury studies, and Sun Young Lee for excellent technical assistance.

Data availability: Raw scRNA-seq data and expression matrix is available at GEO (accession number GSE215247). The scRNA-seq results also can be viewed interactively at the Single Cell Portal. OEC vs Leftovers:
https://singlecell.broadinstitute.org/single_cell/study/SCP1237/leftovers-and-oecs.
OEC subclusters:https://singlecell.broadinstitute.org/single_cell/study/SCP1489/oec-subcluster-batch-corrected

Conflict of Interest: The authors declare no competing interests.

50
Abstract

Olfactory ensheathing cells (OECs) are unique glial cells found in both central and peripheral

nervous systems where they support continuous axonal outgrowth of olfactory sensory neurons

to their targets. Previously we reported that following severe spinal cord injury, OECs

transplanted near the injury site modify the inhibitory glial scar and facilitate axon regeneration

past the scar border and into the lesion. To better understand the mechanisms underlying the

reparative properties of OECs, we used single-cell RNA-sequencing of OECs from adult rats to

study their gene expression programs. Our analyses revealed five diverse OEC subtypes, each

expressing novel marker genes and pathways indicative of progenitor, axonal regeneration,

secreted molecules, or microglia-like functions. We found substantial overlap of OEC genes with

those of Schwann cells, but also with microglia, astrocytes, and oligodendrocytes. We confirmed

established markers on cultured OECs, and localized select top genes of OEC subtypes in

olfactory bulb tissue. We also show that OECs secrete Reelin and Connective tissue growth

factor, extracellular matrix molecules which are important for neural repair and axonal

outgrowth. Our results support that OECs are a unique hybrid glia, some with progenitor

characteristics, and that their gene expression patterns indicate functions related to wound

healing, injury repair and axonal regeneration.

Introduction

Olfactory sensory neurons (OSNs) are remarkable as they undergo life-long neurogenesis within the adult olfactory epithelium (Graziadei & Monti Graziadei, 1985), and their axons are exclusively associated with a specialized type of glia, olfactory ensheathing cells (OECs; Doucette, 1990). OECs provide the outgrowth and guidance factors needed for OSN axons to reach their glomerular targets and exhibit regenerative abilities in response to injury (Doucette, 1991; Li et al., 2005; Williams et al., 2004). OECs have received considerable interest due to their ability to support axonal outgrowth even after severe or complete spinal cord transection in rodents, dogs and humans (Granger et al., 2012; Khankan et al., 2016; Ramon-Cueto et al., 2000; Tabakow et al., 2014; Takeoka et al., 2011; Thornton et al., 2018).

The regenerative capacity of adult spinal cord neurons is inhibited by the lesion site environment that includes a reactive glial scar, together with invading meningeal fibroblasts and multiple immune cells (Burda & Sofroniew, 2014; Wanner et al., 2013). The lesion core is composed of non-neural tissue that inhibits axonal outgrowth due to the upregulation and secretion of chondroitin sulfate proteoglycans that impede axon regeneration (Cregg et al., 2014). Some studies of severe or complete spinal cord transection in rats followed by the transplantation of OECs provided evidence that OECs surround axon bundles which regenerate into and occasionally across the injury site (Khankan et al., 2016; Ramon-Cueto et al., 2000; Takeoka et al., 2011; Thornton et al., 2018). In combination with *in vitro* studies, detailed analyses of injury sites suggest that OECs: 1) function as phagocytes to clear degenerating axonal debris and necrotic cells (Khankan et al., 2016; Nazareth et al., 2020; Su et al., 2013), 2) modulate the immune response to injury (Khankan et al., 2016; Vincent et al., 2007), and 3) interact favorably with the glial scar and lesion core that form after injury *in vitro*, (Lakatos et

al., 2003; Lakatos et al., 2000) and *in vivo*, (Khankan et al., 2016; Thornton et al., 2018). OEC

transplantation following complete spinal cord transection creates an environment that is

conducive to neural repair (Khankan et al., 2016; Takeoka et al., 2011; Thornton et al., 2018).

OECs secrete molecules that stimulate axon outgrowth such as Brain-derived neurotrophic factor

(Bdnf), Nerve growth factor (Ngf), and Laminin (Ruitenberg et al., 2003; Runyan & Phelps,

2009; Woodhall et al., 2001) and enhance cell-to-cell mediated interactions between neurites and

OECs in an inhibitory environment (Chung et al., 2004; Khankan et al., 2015; Windus et al.,

2007). Khankan et al. (2015) showed that neurites grew 2-3 times longer in growth-inhibitory

areas of multicellular 'scar-like' cultures when they were directly associated with olfactory bulb-

derived OECs (OB-OECs), but the molecular mechanisms are unknown.

The outcomes of OEC transplantation studies after spinal cord injury vary substantially in

the literature due to many technical differences in their experimental designs. The source of

OECs has a great impact on the outcome, with OB-OECs showing more promise than lamina

propria-derived OECs, and purified, freshly-prepared OECs being required for optimal OEC

survival. Other important variables include the severity of the injury (hemisection to complete

spinal cord transection), the age of the spinal cord injured host (early postnatal versus adult), and

OEC transplant strategies (delayed or acute transplantation, cell transplants with or without a

matrix; Franssen et al., 2007). Franssen et al. (2007) evaluated OEC transplantation studies, and

reported that 41 out of 56 studies showed positive effects, such as stimulation of regeneration,

positive interactions with the glial scar and remyelination of axons. More recent reviews and

meta-analyses of the effects of OEC transplantation following different spinal cord injury models

reported that OECs significantly improved locomotor function (Watzlawick et al., 2016;

Nakjavan-Shahraki et al., 2018).

Together with collaborators, we conducted six spinal cord injury studies in adult rats with

a completely transected, thoracic spinal cord model followed by OB-OEC transplantation

(Kubasak et al., 2008; Takeoka et al., 2011; Ziegler et al., 2011; Khankan et al., 2016; Thornton

et al., 2018; Dixie, 2019). Results from five of the six studies showed physiological and/or

anatomical evidence of axonal regeneration into and occasionally across the injury site. In 6-8-

month-long studies, Takeoka et al. (2011) and Ziegler et al. (2011) reported physiological

evidence of motor connectivity across the transection in OEC- but not media-transplanted rats.

These experiments used transcranial electric stimulation of the motor cortex or brainstem to

detect motor-evoked potentials (MEPs) with EMG electrodes in hindlimb muscles at 4- and 7-

months post-transection. After 7 months, 70% of OEC-treated rats responded to stimulation with

hindlimb MEPs (motor cortex, 5/20; brainstem 12/20; Takeoka et al, 2011). A complete re-

transection above the original transection was carried out one month later and all MEPs in OEC-

injected rats were eliminated. These results provide physiological evidence of axon conductivity

across the injury site in OEC-treated rats. Additionally, three of our long-term studies evaluated

anatomical axonal outgrowth of the descending serotonergic Raphespinal pathway into and

through the injury site. Significantly more serotonergic-labeled axons crossed the rostral

inhibitory scar border (Takeoka et al., 2011) or occupied a larger area within the injury site core

(Thornton et al., 2018, Dixie, 2019) in OEC-transplanted rats than in fibroblast or media

controls. In addition, significantly more neurofilament-labeled axons were found within the

lesion core of OEC-transplanted versus control rats (Thornton et al., 2018, Dixie, 2019).

The question addressed in the present study is how OB-OECs perform the diverse

functions associated with neural repair. We hypothesize that there exist OB-OEC subtypes, and

that their respective gene programs and secreted molecules underlie their multifaceted reparative

activities. Here we identify the OB-OEC subtypes and their molecular programs that contribute

to injury repair, such as the growth-stimulating secreted and cell adhesion molecules involved in

OB-OEC interactions that promote neurite and axonal outgrowth. We used single-cell RNA-

sequencing (scRNA-seq) to characterize the immunopurified female rat OB-OECs (hereafter

called OECs unless stated otherwise) that are similar to those transplanted into our previous

female rat spinal cord injury studies. We first compared the genes expressed by purified OECs

with those expressed by the 'leftover cells', i.e., the cells which are not selected by the panning

procedure, to determine if the purification process selected for specific OEC subtypes. After

confirming the characteristic OEC markers, our scRNA-seq data revealed five OEC subtypes

which expressed unique marker genes and were further characterized and confirmed

experimentally. Finally, we examined interesting extracellular matrix (ECM) molecules secreted

by OECs, Reelin (*Reln*) and Connective tissue growth factor (*Ccn2*/*Ctgf*), both of which may

facilitate axonal outgrowth into the inhibitory injury core environment.

Results

One particularly challenging aspect facing neural repair of spinal cord injury (SCI) is to facilitate

axonal outgrowth and eventually regeneration across the inhibitory injury site. A common

finding in our studies testing the effects of OEC transplantation following SCI is that OECs

modify the injury site so that some axons project past the glial scar and into the lesion core

surrounded by OECs (Dixie, 2019; Khankan et al., 2016; Takeoka et al., 2011; Thornton et al.,

2018). Figure 1 shows an enlargement of an injury site which contains green fluorescent protein

(GFP)-labeled OECs in the glial fibrillary acidic protein (GFAP)-negative injury core between

the two GFAP-positive spinal cord stumps. The OECs in the injury core are associated with a

few serotonergic- (Figure 1a-f) and many neurofilament-labeled axons (Figure 1g-l). We note,

however, that such bridge formation is rare following severe spinal cord injury in adult mammals

and was detected in 2 out of 8 OEC-transplanted rats versus 0/11 media or fibroblast-

transplanted controls in this study (Dixie, 2019). Combined with the 1/5 OEC-transplanted rats

with axons crossing the injury and 0/5 fibroblast controls in our previous study (Thornton, 2018),

we observed bridges in 3/13 OEC-transplanted rats vs 0/16 controls (p=0.042, two-sample

proportion test). Bridge formation, in conjunction with the additional physiological and

anatomical evidence of axonal connections across the injury site presented in our previous

studies, strongly supports the capacity of OECs in neural repair. To learn more about possible

mechanisms by which OECs support axon regeneration, we prepared immunopurified OECs and

examined their gene expression and diversity with scRNA-seq.

Unbiased scRNA-seq of immunopurified OECs and 'leftover' controls distinguishes OECs,

microglia, and fibroblasts.

Using scRNA-seq, we sequenced a total of 65,481 cells across 7 samples (n=3 OEC preparations

and 4 'leftover' controls) that passed quality control, with an average of 9,354 cells per sample.

Cell clusters were visualized using t-distributed stochastic neighbor embedding (tSNE). The

tSNE plot showed that cells from immunopanned OEC samples were separated from the cells

from the leftover samples (Figure 2a). Cell clustering analysis defined a total of seven clusters

(Figure 2b), each of which showed expression patterns of marker genes that distinguish one

cluster from the others (Figure 2c). Based on previously reported cell type marker genes for

fibroblasts and major glial cell types including OECs, astrocytes, oligodendrocytes, and

microglia, we found elevated expression of OEC marker genes in clusters 2, 3 and 7, microglia

marker genes in clusters 4, 6, and 7, and fibroblast marker genes in clusters 0, 1, and 5 (Figure

2d, Suppl. Figure 1). In contrast, except a few select genes, the majority of markers for astrocytes

and oligodendrocytes showed low expression (Figure 2d). The cell cluster tSNE plot with cell

type labels in Figure 2e, and the distinct expression patterns of cell type markers are confirmed

in a dot plot in Figure 2f. As expected, many cells enriched in the leftover controls were defined

as fibroblasts and microglia based on the corresponding known marker genes, whereas cells from

the immunopanned OEC samples showed high expression of OEC marker genes (Figure 2a vs.

2e). After cell type assignment, a dot plot clearly depicts distinct expression of cell type-specific

marker genes (Figure 2f). Our scRNA-seq results support the expected cell type composition

following the immunopanning procedure. All marker genes for each major cell type, including

statistics, are in Suppl. Table 1.

We next asked if there were differences between cultures of immunopurified OECs and

OECs grown in 'leftover' cultures. While our previous spinal cord injury studies transplanted 90-

95% purified OECs, other labs culture and implant less purified OECs and thus many of the

neighboring cells, i.e., primarily fibroblasts and microglia, would be included. Our comparison

showed that purified OECs express higher levels of *Cryab* (stress protection, inflammation

inhibition), *Nqo1* (antioxidant protection, stress adaptation), and *Postn* (cell proliferation,

survival, migration) genes (Figure 2g), whereas OECs in leftover cultures with fibroblasts and

microglia express higher levels of *Apoe* (lipid transport and glial function), *Calcb* (inflammation

and pain modulation), and *Mgp* (ECM calcium inhibitor; Figure 2g) than purified OECs. These

results indicate immunopurified OECs may have better stress response and proliferative and

survival capacity than the controls.

Immunofluorescence verification of typical OEC marker genes revealed by scRNA-seq

To validate the OEC markers revealed by scRNA-seq, we re-cultured extra cells not used for

sequencing for immunocytochemical confirmation. Because our adult OECs are immunopurified

with anti-nerve growth factor receptor p75 (Ngfr[p75]), we expected and found OEC samples

heavily enriched in Ngfr[p75] in both immunofluorescence (Figure 3a) and scRNA-seq (Figure 3b).

Next, we detected the high expression of two common glial markers, Blbp (*Fabp7;* Figure 3c,

3d) and S100β (Figure 3e, 3f), in OECs. Sox10 is expressed in the nuclei of neural crest-derived

cells such as OECs and Schwann cells, and our cultures showed high expression levels in

scRNA-seq and immunocytochemistry (Figure 3c, 3g, 3k, 3m). Three cell adhesion molecules,

L1-Cam (Figure 3g, 3h; Runyan & Phelps, 2009; Witheford et al., 2013), N-Cam (Figure 3i, 3j),

and N-Cadherins (Figure 3k, 3l), are also expressed in purified OEC cultures. In addition, other

previously reported OEC genes are now verified by scRNA-seq, including the pro-myelinating

Ciliary neurotrophic factor (*Cntf*, Figure 3n; Asan et al., 2003; Roet et al., 2011), the laminin

receptor α7 integrin (*Itga* 7, Figure 3o; Ingram et al., 2016), and the myelin related gene P0

(*Mpz,* Figure 3p; Sasaki et al., 2006). Thus, well-characterized OEC markers were detected by

scRNA-seq and most are widely distributed among OEC clusters.

      In addition to confirming select genes, we also compared the 209 significant OEC

markers identified from our scRNA-seq data with the 309 genes from the meta-analysis of five

OEC microarray studies on cultured early-passage OB-OECs versus other tissues (Roet et al.,

2011). We found 63 genes overlapping between our OEC markers and the published markers,

representing a 15.3-fold overlap enrichment (p-value: 4.3 e-58), further supporting the reliability

of our scRNA-seq findings.

235 **Subclustering analysis of OECs revealed refined OEC subtypes and their top marker genes.**

236 Next, we extracted OECs and performed a subclustering analysis with OECs alone, and found

237 five distinct subclusters (Figure 4a). All subclusters except for cluster 3 expressed previously

238 identified markers of OECs as well as subsets of marker genes for astrocytes, oligodendrocytes,

239 and Schwann cells (Figures 4b, c). The percentage of OECs within each cluster that express a

240 gene, the percentage of OECs in all other subtypes that express a gene and their p values are

241 found in Suppl. Table 2. Interestingly, cluster 3 showed expression of both microglia and OEC

242 markers. The top 20 genes expressed by each OEC subtype were investigated to evaluate

243 potential functional differences between the subtypes (Figure 4c; Suppl. Table 3), and their

244 enriched pathways (Figure 4d). Most references for the specific genes within the clusters are

245 found in Suppl. Table 3.

246 *Cluster 0 is rich in matricellular proteins.* Pathway enrichment analysis showed that

247 ECM organization and collagen biosynthesis were significantly enriched in the marker genes for

248 the largest OEC subcluster 0 (Figure 4d). Two of the top 10 genes are members of the Ccn

249 family of matricellular proteins, *Ccn2*/*Ctgf* (Connective tissue growth factor; Figure 5a) and

250 *Ccn3/NOV* (nephroblastoma overexpressed gene; Figure 5b, Suppl. Table 3). Both are secreted

251 ECM proteins which regulate the activity of growth factors and cytokines, function in wound

252 healing, cell adhesion, and injury repair. The presence of *Ccn3* in OECs is novel, while

253 *Ccn2/Ctgf* is well established (Lamond & Barnett, 2013; Roet et al., 2011). In figures 5c, c1-2,

254 we show that *Ccn2* and *Sox10* are highly expressed in our cultured OECs. Because Mokalled et

255 al. (2016) reported that *Ctgfα* is a critical factor required for spontaneous axon regeneration

256 following spinal cord injury site in zebrafish, we asked if GFP-labeled OECs transplanted 2

257 weeks following a complete spinal cord transection in adult rats, also expressed Ccn2/Ctgf. We

found high levels of Ctgf expression in GFP-OECs (n=4 rats) that bridged much of the injury site

and also detected Ctgf on near-by cells (Figure 5d, d1-2). GFP-labeled fibroblast transplantations

(n=3 rats) served as controls and also expressed Ctgf. Other top ECM genes are *Serpinf1* (Serine

protease inhibitor)*, Fbln2* (Fibulin2)*, Fn1* (Fibronectin-1), and *Col11a1* (Collagen, type XI,

alpha 1; Suppl. Table 3).

The top-ranked marker gene for cluster 0 is *Pmp22* (Peripheral myelin protein 22), a

tetraspan membrane glycoprotein that is highly expressed in myelinating Schwann cells, and

contributes to the membrane organization of compact peripheral myelin. The third-ranked gene,

Gldn (Gliomedin) is secreted by Schwann cells and contributes to the formation of the nodes of

Ranvier. *Fst* (Follistatin) also plays a role in myelination and is expressed in areas of adult

neurogenesis.

Many of the top cluster 0 genes are found in other glial cells: *Fbln2* is secreted by

astrocytes, *Ccn2/Ctgf* is expressed by astrocytes and Schwann cells, *Nqo1* (NAD(P)H

dehydrogenase, quinone 1 enzyme) is found in Bergmann glia, astrocytes, and oligodendrocytes,

and *Marcks* (Myristoylated alanine-rich C-kinase substrate protein) regulates radial glial function

and is found in astrocytes and oligodendrocytes. In addition, *Cd200* is expressed by astrocytes

and oligodendrocytes, and binds to its receptor, *Cd200R*, on microglial cells. Thus, the OECs in

cluster 0 express high levels of genes found in other glia, many of which are related to ECM and

myelination.

Cluster 1 contains classic OEC markers. Three of the top five genes in cluster 1, *Pcsk1*

(proprotein convertase PC1), *Gal* (neuropeptide and member of the corticotropin-releasing factor

family) and *Ucn2* (Urocortin-2) were reported in the meta-analysis by Roet et al. (2011). Galanin

is expressed by neural progenitor cells, promotes neuronal differentiation in the subventricular

zone and is involved in oligodendrocyte survival.

A well-known oligodendrocyte and Schwann gene associated with myelin, *Cnp* (2-3-

cyclic nucleotide 3-phosphodiesterase), was high in OEC cluster 1 in addition to the cytokine

Il11 (Interleukin 11) that is secreted by astrocytes and enhances oligodendrocyte survival,

maturation and myelin formation. The chondroitin sulfate proteoglycan Versican (*Vcan*) is

another top gene expressed by astrocytes and oligodendrocyte progenitor cells.

Many of the genes and pathways that characterize cluster 1 are involved in nervous

system development and axon regeneration (*Atf3, Btc, Gap43, Ngfr^{p75}, Bdnf, Pcsk1*), and are

found in other glia (Suppl. Table 3). Interestingly, after nerve injury, *Atf3* (cyclic AMP-

dependent transcription factor 3) is upregulated by Schwann cells in the degenerating distal

nerve stump and downregulated after axon regeneration is complete (Hunt et al., 2004). *Btc*

(Betacellulin), part of the Epidermal growth factor (Egf) family and a ligand for Egfr, also is

expressed by Schwann cells after nerve injury in a pattern similar to that of *Atf3*, as is Brain-

derived neurotrophic factor (*Bdnf*). In addition, the proprotein convertase PC1 (*Pcsk1*) cleaves

pro-Bdnf into its active form in Schwann cells and therefore also contributes to axon outgrowth.

Together, the OECs in cluster 1 secrete a number of important growth factors associated with

regeneration, including a novel one, *Atf3*, reported here (Suppl. Table 3).

Cluster 2 represents a distinctive proliferative OEC subtype. Most top genes in cluster 2

(Figure 6a-f) regulate the cell cycle (*Mki67, Stmn1, Cdk1, Cks2, Cdkn3, Cdca8, Cdca3*), are

associated with mitosis (*Cenpf, Spc24, Tpx, Ckap2, Prc1, Ube2c, Top2a*), or contribute to DNA

replication and repair (*Dut, Fam111a*, Suppl. Table 3). The top ranked gene, *Stmn1* (Stathmin 1;

Figure 6e), is a microtubule destabilizing protein that is widely expressed in other OEC clusters

and in areas of adult neurogenesis. Pathway enrichment confirms the proliferative property of

cluster 2 with pathways such as G1/S transition of mitotic cell cycle and DNA replication

(Figure 4d). These results show that there are distinct OEC progenitors in our cultures within the

olfactory nerve layer of the adult olfactory bulb.

MKi67 is a well-known proliferation gene concentrated in cluster 2 (Figure 6f). Previous

studies suggested the presence of two morphologically distinct OECs – the Schwann-cell-like,

spindle-shaped OECs with high Ngfr^{p75} expression and the astrocyte-like, large flat OECs with

low Ngfr^{p75} (Franceschini & Barnett, 1996). To determine if the proliferative OECs differ in

appearance from adult OECs, and whether there is concordance between our OEC subtypes

based on gene expression markers and previously described morphology-based OEC subtyping

(Franceschini & Barnett, 1996), we analyzed OECs identified with the anti-Ki67 nuclear marker

and anti-Ngfr^{p75}(Figure 6g-h). Of the Ki67-positive OECs in our cultures, 24% ± 8% were

strongly Ngfr^{p75}-positive and spindle-shaped, whereas 76% ± 8% were flat and weakly Ngfr^{p75}-

labeled (n=4 cultures, *p*= 0.023). Here we show that a large percentage (~3/4^{ths}) of proliferative

OECs are characterized by large, flat morphology and weak Ngfr^{p75} expression resembling the

previously described morphology-based astrocyte-like subtype. Our results indicate the two

types of OEC classifications share certain degrees of overlap, indicating similarities but also

differences between the two classification methods.

Cluster 3 resembles both microglia and OECs. In addition to the typical OEC markers

(*Ngfr^{p75}*, *S100b*, and *Sox10*), cluster 3 also expressed numerous microglia markers (Figure 4b). In

fact, all top 20 genes in cluster 3 are expressed in microglia, macrophages, and/or monocytes

(Suppl. Table 3). Microglia markers in the top 20 genes include *Tyrobp,* which functions as an

adaptor for a variety of immune receptors, *Cst3* (Cystatin C)*, Anxa3* (Annexin A3)*, Aif1 (*Iba-1),

and *Cd68*. OECs are known to endocytose bacteria and degrade axons, and therefore participate

in the innate immune function (Khankan et al., 2016; Leung et al., 2008; Nazareth et al., 2015;

Vincent et al., 2007). We also observed relatively high expression of genes involved in lysosome

functions, such as *Cst3, Ctsz, Laptm5, Ctsb*, and *Lyz2*, compared to those in other OEC clusters

(Figure 4e). Three additional top genes are involved in the complement system (*C1qA, C1qb,*

and *C1qc)*, and cluster 3 pathways are highly enriched for inflammatory response and neutrophil

immunity (Figure 4d).

Smithson and Kawaja (2010) identified unique microglial/macrophages that

immunolabeled with Iba-1 (*Aif1*) and Annexin A3 (*Anxa3*) in the olfactory nerve and outer nerve

layer of the olfactory bulb. These authors proposed that Iba1-Anxa3 double-labeled cells were a

distinct population of microglia/macrophages that protected the olfactory system against viral

invasion into the cranial cavity. Based on our scRNA-seq data we offer an alternative

interpretation that at least some of these Iba-1-Anxa3 cells may be a hybrid OEC-microglial cell

type. Supporting this interpretation, there are a number of reports that suggest OECs frequently

function as phagocytes (e.g., Khankan et al., 2016; Nazareth et al., 2020; Su et al. 2013).

Cluster 4 has characteristics of astrocytes and oligodendrocytes. Cluster 4 is quite small.

Its top two marker genes, *Sidt2*, a lysosomal membrane protein that digests macromolecules for

reutilization and *Ckb*, a brain-type creatine kinase which fuels ATP-dependent cytoskeletal

processes in CNS glia, are found in other OEC clusters. Four top genes, however, appear almost

exclusively in cluster 4: 1) *Npvf* (Neuropeptide VF precursor/Gonadotropin-inhibitory hormone)

inhibits gonadotropin secretion in several hypothalamic nuclei, and is expressed in the retina, 2)

Stmn2 (Stathmin2) is a tubulin-binding protein that regulates microtubule dynamics, 3) *Vgf* (non-

acronymic) is synthesized by neurons and neuroendocrine cells and promotes

oligodendrogenesis, and 4) *Mt3* (Metallothionein-3) is a small zinc binding protein associated

with growth inhibition and copper and zinc homeostasis (Suppl. Table 3).

Other genes expressed by cluster 4 are either expressed by other glia and/or are ECM

molecules. Five genes are associated with both oligodendrocytes and astrocytes *(Ckb, C1ql1,*

Gria2, *Stmn2, Ntrk2*, Suppl. Table 3). Among these, the astrocytic gene *C1ql1* is a member of

the Complement component 1q family and regulates synaptic connectivity by strengthening

existing synapses and tagging inactive synapses for elimination. Astrocytic Thrombospondin 2

(*Thbs2*) also controls synaptogenesis and growth. Other genes reported in astrocytes include *Ptn*

(Pleiotrophin), *Igfbp3* (Insulin-like growth factor binding protein 3), and *Mt3,* whereas *Gpm6b*

(Glycoprotein M6b) is involved in the formation of nodes of Ranvier in oligodendrocyte and

Schwann cell myelin. Cluster 4 also has a significant enrichment for genes involved in ECM

organization (Figure 4d; *Actn1*, *Col81a*, *Bgn*, *Fn1* and *Timp2*), together with clusters 0 and 1, as

well as genes involved in positive regulation of cell migration together with cluster 3 (Figure 4f).

Trajectory and potential interactions among OEC subclusters

We performed pseudotime trajectory analysis using the Slingshot algorithm to infer lineage

trajectories, cell plasticity and lineages by ordering cells in pseudotime based on their

transcriptional progression reflected in our scRNA-seq data. Transcriptional progression refers to

the changes in gene expression profiles of cells as they undergo differentiation or transition

through different states. The trajectory analysis results suggest that there are potential transitions

between specific OEC subclusters. Our results show that there are two distinct trajectories

(Figure 4g). The first trajectory involves clusters 2, 0 and 4, whereas the second involves clusters

2, 0, 1, and 3. Although the directionality of these trajectories is indistinguishable, the fact that

cluster 2 is enriched for cell cycle genes and is the converging point of both trajectories, suggests

that cluster 2 proliferates into the other OEC clusters. The predicted trajectories based on our

scRNAseq data suggest plasticity in the cell clusters.

We also modeled potential autocrine/paracrine ligand-receptor interactions between OEC

subclusters using NicheNet (Figure 4h). This analysis revealed that clusters 0, 1, and 2 have

more ligands going to other OEC clusters, whereas clusters 3 and 4 generally receive ligands

from other clusters. The network also showed that clusters 0, 1 and 4 express genes encoding

neurotropic factors such as Bdnf, Cntf and Neurturin (*Ntrn*), that are involved in neuronal

survival and neurite outgrowth. *Gal* from cluster 1 is a mediator for glia-glia, and glia-neuron

communication (Gresle et al., 2015; Ubink et al., 2003). We found that cluster 4 secretes colony

stimulating factor 1 (*Csf1*) which also shows potential regulation of cluster 3 through *Csf1r*.

Overall, the trajectory and network analyses support potential cell plasticity and lineage

relationships and cell-cell communications across OEC subclusters.

Spatial confirmation of defined OEC subclusters within the olfactory nerve layer *in situ*

Here we confirm, at the protein level, that some of the top genes derived from cultured OECs

clusters are expressed in sections of the olfactory system from 8-10-week-old female Sprague

Dawley rats. Our results illustrate the spatial distribution of 2-3 top genes from clusters 0 to 3 in

the olfactory nerve layer (Figure 7; corresponding tSNE plots are in Suppl. Figure 2 for

comparison). In cluster 0 we show high levels of Peripheral myelin protein 22 (Pmp22) in OECs

(Figure 7a) with the remainder of the olfactory bulb unlabeled. The small protein Gliomedin

(Gldn) is a component of Schwann cell microvilli and facilitates the formation of peripheral

nodes of Ranvier (Eshed et al., 2005). Here we detect numerous small dot-like structures that

overlay the olfactory nerve layer (Figures 7b, c). The detection of high levels of Gldn in

nonmyelinating OECs without nodes of Ranvier is surprising and suggests that it may also

function as a glial ligand associated with olfactory sensory neuron axons. For cluster 1, we

examined the expression of the important axonal growth factor Activating transcription factor 3

(Atf3) and confirmed that OECs express it widely (Figure 7d). Growth associated protein 43

(Gap43), a well-established marker of OECs and immature olfactory sensory neurons, is highly

expressed in the olfactory nerve layer and at a lower level in the glomeruli of the olfactory bulb

(Figure 7e). Nestin (Nes), an intermediate filament associated with neural stem cells, is widely

expressed in the olfactory nerve layer (Figure 7f).

Compared to the two large clusters, the remaining clusters contain more specialized

OECs. The top marker in the proliferative cluster 2 is Stathmin1 (Stmn1), encoding a

microtubule destabilizing protein found in areas of adult neurogenesis (Boekhoorn et al., 2014).

Stathmin 1 immunoreactivity (Figure 7g, Suppl. Figure 2) is detected in all OEC clusters and

immature olfactory sensory neurons, and therefore fills the olfactory nerve layer in a pattern

similar to that seen in Gap43 (Figure 7e). Two markers associated with the cell cycle, Ube2c

(Figure 7h) and Cdk1 (not shown), identify cluster 2 cells in the olfactory nerve layer, as well as

the early progenitor marker Mki67 (Figure 7i). Cluster 3 is strongly associated with OEC

immune function and the expression of Apolipoprotein E (Apoe), Annexin A3 (Anxa3), and

Allograft inflammatory factor 1 (Aif1/Iba1) were confirmed. Apoe (Figure 7j) is detected

throughout the olfactory nerve layer and includes some cellular structures. Expression of Anxa3

(Figure 7k) and Aif1/Iba1(Figure 7l) is similar – distinct small cells labeled in the olfactory

nerve layer. Our smallest group, cluster 4, has been difficult to confirm with selected antibodies

targeting the top marker genes, likely due to the limited number of cells in this cluster. Overall,

our *in-situ* experiments confirmed the presence and distribution of 4 out of the 5 OEC subtypes

detected by scRNA-seq.

OECs synthesize and secrete Reelin

Reelin (*Reln*) is detected in OECs in this study (Figures 2f, 8a, g) and reported in previous

microarray studies (Guerout et al., 2010; Roet et al., 2011). However, the presence of the *Reln*

gene within OECs is disputed in the literature (Dairaghi et al., 2018; Schnaufer et al., 2009). *Reln*

codes for a large ECM glycoprotein that is secreted by neurons (D'Arcangelo et al., 1995;

Kubasak et al., 2004), but also by Schwann cells (Panteri et al., 2006). In the canonical Reelin-

signaling pathway, Reelin binds to the very-low-density lipoprotein receptor (Vldlr) and

apolipoprotein E receptor 2 (ApoER2) and induces Src-mediated tyrosine phosphorylation of the

intracellular adaptor protein Disabled-1 (Dab1). Both Reelin and Dab1 are highly expressed in

embryos and contribute to correct neuronal positioning. A recent study claimed that peripheral

OECs express high levels of Dab1 and Vldlr (Dairaghi et al., 2018), and that Reelin was

expressed only by mesenchymal cells located near the cribriform plate, not in OECs (Dairaghi et

al., 2018; Schnaufer et al., 2009). Our scRNA-seq findings of OECs demonstrated low

expression levels of *Dab1* and *Apoer2* (Figure 8b-c), moderate expression of *Vldlr* (Figure 8d)

and substantial levels of *Reln* in OEC subclusters 0-3 (Figure 8a). Interestingly, OECs also

contain proteolytic enzymes that cleave the 400Kd secreted Reelin. Tissue plasminogen activator

(tPA; Figure 8e) cleaves Reln at its C-terminus, and the matrix metalloproteinase ADAMTS-4

cuts at both the N- and C-cleavage sites of Reln (Figure 8f; Krstic et al., 2012).

The confusion about Reelin expression in OECs likely stemmed from the use of standard

immunochemical methods to confirm its presence. Our attempts to detect Reelin expression in

the adult olfactory nerve layer (ONL) or in cultured OECs were unsuccessful using several

antibodies and multiple protocols, despite consistent localization of Reelin within mitral cells of

the olfactory bulb. To understand this discrepancy, we examined Reelin expression in embryonic

sections which included both the peripheral and OB-OECs. Images of E16.5 $Reln^{+/+}$ olfactory

system show Reelin expression in peripheral OECs and in neurons of the olfactory bulb (Figure

8g, green), but not on OB-OECs in the ONL or in sections of $Reln^{-/-}$ mice (Figure 8g, g2, h). In

comparison, Blbp expression is uniformly expressed in the ONL of both peripheral and OB-

OECs (Figure 8g1, h1, red). Enlargements of the region where axon bundles from olfactory

sensory neurons join the ONL show that Blbp is present in OB-OECs throughout the ONL, but

Reelin is detected only on peripheral OECs (Figure 8g, g1-2).

To determine if adult rat OB-OECs synthesize and secrete Reelin *in vitro*, we treated

primary Ngfr^{p75}-purified OECs with or without Brefeldin-A, a fungal metabolite that specifically

inhibits protein transport from the endoplasmic reticulum to the Golgi apparatus (Misumi et al.,

1986). In Brefeldin-treated primary OEC cultures, spindle-shaped GFP-labeled OECs contain

Reelin in their cytoplasm (Figure 9a, arrows). Large contaminating Reln-immunonegative cells

present in primary cultures serve as an intrinsic control (Figure 9a, arrowhead). Reelin

expression was also confirmed in Brefeldin-treated Ngfr^{p75}-purified OEC cultures (Figure 9b-c),

a finding which supports our scRNA-seq results.

To further confirm that cultured OECs express and secrete Reelin, we performed western

blotting. The G10 antibody recognizes full-length Reelin protein, which is approximately 400

kDa, and its two cleaved fragments at ~300 and ~180 kDa (Lambert de Rouvroit et al., 1999).

Both the rat ONL and $Reln^{+/+}$ mouse olfactory bulbs showed the characteristic large molecular

weight band at 400 kDa and another at 150 kDa (Figure 9d, lanes 1-2, Reln). As expected, these

bands were absent in mutant $Reln^{-/-}$ olfactory bulbs (Figure 9d, lane 3). Additionally, Reelin-

positive bands were present in blots of OEC whole cell lysates (WCL), representing Reelin

synthesized in OECs, and OEC conditioned media (CM; Figure 9d, lanes 4-8 Reln), representing

Reelin secreted by OECs. A 4-15% gradient gel allowed for the visualization of all three Reelin

isoforms (Figure 9e). Brain extracts of $Reln^{+/+}$ (positive control) and $Reln^{-/-}$ (negative control)

mice confirm the specificity of Reelin detection (Figure 9e, lanes 1-2, 9f). All three Reelin

isoforms at their corresponding molecular weights are present in rat ONL and OEC CM samples

(Figure 9e, lanes 3-6, 9f). Quantification of the band densities from 4-15% gels showed that

cultured OECs derived from the rat ONL and CM produce significantly higher levels of 400 and

300 kDa Reelin compared to $Reln^{-/-}$ mice (Figure 9f, $p<0.05$).

Discussion.

Our scRNA-seq study of immunopurified OECs identified five OEC subtypes with distinct gene

expression patterns, pathways, and properties of each subtype. The well-established OEC

markers are widely distributed throughout most clusters derived from cultured OECs. We also

experimentally confirmed a number of the top markers at the protein level in the OEC subtypes

on sections of the rat olfactory system and established that OECs secrete both Reelin and Ctgf,

ECM molecules reported to be important for neural repair and axon outgrowth. Additional

proregenerative OEC marker genes such as *Atf3, Btc,* and *Pcsk1* also were identified. Our

findings provide an unbiased and in-depth view of the heterogeneity of OEC populations with

demonstrated injury repair capacity, and the potential mechanisms that underlie their

regenerative and protective properties.

This scRNA-seq analysis confirmed the immunopurification of the OECs compared with

the mixture of OECs grown with numerous fibroblasts and microglia. Although some may

debate if purified OECs are needed to maximize neural repair, this study suggests that

immunopurified OECs have higher expression of anti-stress genes than unpurified controls.

Several of our earlier studies, which transplanted similarly purified OECs, also showed evidence

of neural regeneration following severe spinal cord injury (Khankan et al., 2016; Takeoka et al.,

2011; Thornton et al., 2018). In addition to the purity of OECs, we found significant overlap of

the marker genes between our study and those from the meta-analysis of five OEC microarray

studies on cultured OB-OECs (Roet et al., 2011), supporting the reproducibility of our scRNA-

seq findings. When we compared our data from cultured OB-OECs to previous scRNA-seq

studies of the olfactory system tissue which included peripheral OECs (Durante et al, 2000), or a

mixture of peripheral and OB-OECs (Tepe et al., 2018), there was little overlap. Most likely, the

substantial differences were due to the dissection areas, techniques used for OEC preparation, the

species studied, as well as sex and age of the subjects. It is important to note that our OB-OEC

populations were repeatedly shown to induce injury repair, and their subtypes and markers

support their diverse reparative functions, whereas the neural repair function of the cell

populations in the other scRNA-seq studies using different tissue preparations is unclear.

504   OECs are hybrid glia that promote neural repair.

Based on intrinsic gene expression patterns revealed by scRNA-seq data, there is expression of

select markers for Schwann cells, astrocytes, oligodendrocytes and microglia within our five

clusters of OECs. We observed transcriptionally distinct clusters that highly expressed select

markers for proliferation (cluster 2), microglia (cluster 3), and astrocytes and oligodendrocytes

(cluster 4). The microglia/macrophage-like cells reported by Smithson and Kawaja (2010) do

resemble our transcriptome-based subtyping of OEC cluster 3. Our results further support the

activities of OECs in modulating immune responses, based on the expression of genes involved

in the endocytosis of bacteria, the regulation of microglia activation, and the expression of

numerous lysosomal pathways (Leung et al., 2008). OECs in cluster 4 are distinctive by their

expression of several astrocytic markers that regulate synaptogenesis (Thrombospondin 2) and

synaptic connectivity, and remove inactive synaptic connections (Complement component 1q

family; Christopherson et al., 2005; Eroglu & Barres, 2010). Additionally, our trajectory and

network analyses indicate potential relationships between the OEC subclusters, which warrant

further experimental testing.

Both OECs and Schwann cells originate from the neural crest, express similar

immunological markers, and share many transcriptional similarities (Barraud, et al., 2010; Forni

et al., 2011; Vincent et al., 2005). OECs share phenotypic characteristics with myelin-producing

Schwann cells and oligodendrocytes, but more closely resemble Schwann cells (Doucette, 1991;

Radtke et al., 2011; Ramon-Cueto & Valverde, 1995). Gliomedin is a required ECM protein that

initiates the assembly of nodes of Ranvier along Schwann cells (Eshed et al., 2005), yet despite

the fact that OECs do not have nodes of Ranvier, they express high levels of Gliomedin. This

novel Gliomedin expression in OECs suggests a possible function as a glial ligand for the

unmyelinated axon bundles that OECs ensheath. Following peripheral nerve injury, the loss of

contact between axons and myelinating Schwann cells induces a nonmyelinating phenotypic

change in Schwann cells (Fu & Gordon, 1997). These dedifferentiated nonmyelinating Schwann

cells are proregenerative and upregulate the expression of Glial fibrillary acidic protein (Gfap;

(Jessen et al., 1990), Ngfr[p75] receptor (You et al., 1997), cell adhesion molecules L1 and N-Cam

(Martini & Schachner, 1988), and neurotrophic factors, including Bdnf (Meyer et al., 1992) and

Glial cell-line derived neurotrophic factor (Trupp et al., 1995). Genes for these proregenerative

factors are detected in OECs in the present study, as well as the transcription factor *Mitf*, that

controls a network of genes related to plasticity and repair in Schwann cells (Daboussi et al.,

2023).

Our *in vitro* and *in vivo* experimental validation of numerous marker genes at the protein

level strongly support their validity. Based on our tissue localization, the two large clusters (0

and 1) were found throughout the olfactory nerve layer, whereas cluster 2 (progenitors) and

cluster 3 (microglial) are scattered within the olfactory nerve layer. Future functional studies of

these subclusters will benefit from both the marker genes and their localizations uncovered in

this study.

Reelin and CTGF are distinctive markers for OECs.

Reelin is a developmentally expressed protein detected in specific neurons, in addition to OECs

and Schwann cells. The canonical Reelin-signaling pathway involves neuronal-secreted Reelin

binding to Vldlr and ApoER2 expressed on Dab1-labeled neurons. Following Reelin binding,

Dab1 is phosphorylated by Src family kinases which initiates multiple downstream pathways.

Very little is known, however, about Reelin secreted by glia. Panteri et al. (2006) reported that

Schwann cells express low levels of Reelin in adults, and that it is upregulated following a

peripheral nerve crush, as is reported above for many neurotrophic factors. Reelin loss in

Schwann cells reduced the diameter of small myelinated axons but did not affect unmyelinated

axons (Panteri et al., 2005). In the olfactory system, OECs ensheath the Dab1-labeled,

unmyelinated axons of olfactory sensory neurons which are continuously generated and die

throughout life. OEC transplantation following spinal cord injury would provide an exogenous

source of Reelin that could phosphorylate Dab1-containing neurons or their axons. Dab1 is

expressed at high levels in the axons of some projection neurons, such as the corticospinal

pathway (Abadesco et al., 2014). Future experiments are needed to explore the function that

glial-secreted Reelin may have on axonal regeneration.

Adult mammals show little evidence of spontaneous axonal regeneration after a severe

spinal cord injury in contrast to transected neonatal rats (Bregman, 1987; Bregman et al., 1993)

and young postnatal opossums (Lane et al., 2007). In immature mammals, axons continue to

project across or bridge the spinal cord transection site during development. Lower organisms

such as fish, show even more evidence of regeneration following severe SCI. Mokalled et al.

(2016) reported that glial secretion of Ctgfα/Ccn2 was both necessary and sufficient to stimulate

a glial bridge for axon regeneration across the zebrafish transection site. Cells in the injury site

that express Ctgf include ependymal cells, endothelial cells, and reactive astrocytes (Conrad et

al., 2005; Mokalled et al., 2016; Schwab et al., 2001). Here we show that, although rare, Ctgf-

positive OECs can contribute to glial bridge formation in adult rats. The most consistent finding

among our severe SCI studies combined with OEC transplantation is the extent of remodeling of

the injury site and axons growing into the inhibitory lesion site, together with OECs and

astrocytes. The formation of a glial bridge across the injury was critical to the spontaneous axon

generation seen in zebrafish (Mokalled et al., 2016) and likely contributed to the axon

regeneration detected in our OEC transplanted, transected rats (Dixie, 2019; Khankan et al.,

2016; Takeoka et al., 2011; Thornton et al., 2018).

Limitations in these OEC scRNA-seq studies

We recognize that this study is a single snapshot of OEC gene expression derived from adult

female rats before they are transplanted above and below the spinal cord transection site. We

would expect the gene expression of transplanted OECs to change in each new environment, i.e.

as they migrate into the injury site, integrate into the glial scar, and wrap around axons. Based on

our past studies, OECs survived in an outbred Sprague-Dawley female rat model for ~ 4 weeks

(Khankan et al., 2016) and in an inbred Fischer 344 female model for 5-6 months (Dixie, 2019).

As spinal cord injury transplant procedures are further enhanced and OEC survival improves,

these hybrid glial cells should be examined at multiple time points after transplantation to better

evaluate their proregenerative characteristics.

We acknowledge that in humans, males account for ~80% of spinal cord injuries

(National Spinal Cord Injury Statistical Center, 2024) and sustain more serious urinary tract

issues than females. We examined females in the current study due to practical experimental

considerations, but it is necessary to examine males in future studies. We also did not modulate

any of the genes or proteins in the identified OEC subtypes to test their causal and functional

roles, thus our findings remain correlative. Future gene/protein modulation studies are necessary

to understand the functional roles of the individual OEC subtypes in the context of their

reparative functions to determine which pathways and subtypes are more critical and can be

enhanced for neural repair. Our current findings build the foundation for these future studies to

help resolve the role of OECs in spinal cord injury repair.

Extensive differences between OEC preparations contribute to the large variation in

results from OEC treatments following spinal cord injury. This scRNA-seq study focused

entirely on OB-OECs, and the next step would be to carry out similar studies on the peripheral,

lamina-propria-derived OECs to discern the differences between the two OEC populations. Such

comparative studies using scRNA-seq will help define the underlying mechanisms and resolve

the variability in results from OEC-based therapy. Detailed studies of the composition of

different OEC transplant types will contribute to identifying the most reparative cell

transplantation treatments.

Conclusion
606

To understand how immunopurified OECs repair spinal cord injuries, we carried out an unbiased

scRNA-seq analysis and found that OECs have a wide range of subtypes including a proliferative

state, microglia-like cells and neural regenerative clusters. We demonstrated that these multi-

functional OECs are clearly distinguishable from each other by expression of unique genes and

pathways within subtypes based on the scRNAseq data and our *in vivo* tissue studies. These

subtypes of OECs produce important ECM genes, including *Reln* and *Ctgf* and many other

signaling molecules which are involved in a wide range of intercellular communication as well

as axon regeneration. Our single-cell resolution investigation offers a comprehensive molecular

landscape of OECs and numerous novel targets for future injury repair and

functional/mechanistic studies.

Material and Methods

Animals.

All of the experimental procedures used on animals were approved by the Chancellor's Animal

Research Committee at UCLA and carried out in accordance with the National Institutes of

Health guidelines. Animals were housed in the Terasaki Life Science Building vivarium under

standard conditions with free access to food and water.

Sprague-Dawley rats: We bred our male GFP-expressing Sprague-Dawley rats (Perry et al,

1999) with unlabeled female Sprague-Dawley rats (Charles Rivers Laboratory), which were used

previously to obtain OECs from female rats for two of our spinal cord injury studies (Khankan et

al., 2016; Thornton et al., 2018) in order to unambiguously identify the transplanted GFP-

positive OECs near the injury sites. A total of 8, 8-10-week-old GFP-labeled female rats were

used to obtain GFP-OECs for scRNA-seq. Only females were used in order to match the sex of

previous SCI studies conducted exclusively on female rats (Dixie, 2019; Khankan et al., 2016;

Takeoka et al., 2011; Thornton et al., 2018). Following complete thoracic spinal cord transection,

an adult rat is unable to urinate voluntarily and therefore urine must be manually "expressed"

twice a day throughout the experiment. Females have a shorter urethra than males, and thus their

bladders are easier to empty completely. An additional 4, 8-10-week GFP-labeled rats were used

to isolate Reelin protein for western blotting and 6, 8-10-week unlabeled female Sprague-

Dawley rats were used to confirm gene expression in the five OEC subtypes. An overdose of

ketamine-xylazine or pentobarbital was used for euthanasia before either the rat olfactory bulbs

were removed or the rats were perfused with 4% paraformaldehyde or 2% PLP (2%

Paraformaldehyde; 0.075M Lysine-HCL-monohydride; 0.010M Sodium periodate; 0.1M sodium

phosphate) followed by a 1-2 hr postfix.

Reeler mice: To study Reelin expression in OECs, heterozygous ($Reln^{+/-}$) mice, originally

purchased from the Jackson Laboratory (B6C3Fe-*ala-Reln^{rl}*, Bar Harbor, ME), were used for

breeding. Genotypes of these mice were identified by PCR (D'Arcangelo et al., 1996). To obtain

timed pregnancies, $Reln^{+/-}$ mice were bred and checked every morning for plugs. Pregnant mice

were heavily anesthetized with pentobarbital (100 mg/kg) and embryos were extracted at

embryonic day 16.5. Heads were immersed in 2% PLP, washed in PB, cryoprotected, and

embedded in Shandon M1 embedding matrix (Thermo Scientific, 1310).

Preparation of olfactory bulb-derived OEC cultures from rats and mice.

Olfactory bulbs were collected from 8-10-week-old GFP-labeled Sprague-Dawley rats or $Reln^{+/+}$

and $Reln^{-/-}$ mice. The leptomeninges were removed to reduce fibroblast contamination. Methods

to prepare OEC primary cultures were adapted from Ramon-Cueto et al. (2000) and follow those

reported by Runyan and Phelps (2009) for mouse OECs and Khankan et al. (2015) for rat OECs.

For scRNA-seq, this protocol was replicated on two separate dates and used a total of 8 adult

female Sprague-Dawley rats. Primary cultures were generated from both olfactory bulbs from

each rodent. The first two layers of the olfactory bulb were dissected, isolated, and then washed

in Hank's balanced salt solution (HBSS, Gibco, Rockville, MD) prior to tissue centrifugation at

365 g for 5 mins. The tissue pellet was resuspended in 0.1% trypsin and HBSS without

Ca^{2+}/Mg^{2+} (Gibco, Rockville, MD), then placed in a 37°C water bath, and mixed intermittently

for 10 mins. A mixture of 1:1 Dulbecco's modified eagle medium (DMEM) and Ham's F12 (D/F

medium, Gibco) supplemented with 10-15% Fetal Bovine Serum (FBS, Hyclone, Logan, UT)

and 1% Penicillin/Streptomycin (P/S, Gibco; D/F-FBS-P/S medium) was used to inactivate

trypsin prior to centrifugation. Dissociated cells were rinsed, centrifuged 3 times, and plated into

12.5 cm^2 culture flasks pre-coated with 0.05mg/ml poly-L-lysine (PLL, Sigma, St. Louis, MO).

Cells were maintained at 37°C for 5-8 days and D/F-FBS-P/S medium was changed every 2

days.

Immunopurification was carried out using hydrophobic petri dishes coated overnight with

Biotin-SP-conjugated AffiniPure goat anti-rabbit IgG (1:1000; Jackson ImmunoResearch, West

Grove, PA) in 50 mM Tris buffer at 4°C followed by another overnight incubation at 4°C with

either mouse anti-Ngfr^{p75} for rat OECs (Suppl. Table 4, Chandler et al., 1984) or rabbit anti-

Ngfr^{p75} for mouse OECs (Suppl. Table 4) in 25mM Phosphate-buffered saline (PBS). Dishes

were rinsed 3 times with 25 mM PBS and treated with a mixture of PBS and 0.5% Bovine serum

albumin (BSA) for 1 hr at room temperature. Prior to the addition of cells, antibody-treated

dishes were washed with PBS and DMEM.

In preparation for immunopanning, two pairs of adult rat olfactory bulbs (for scRNA-seq)

or four pairs of mouse bulbs (for western blots) were dissociated with 0.25% trypsin-EDTA at

37°C for 3 mins and D/F-FBS-P/S medium was added to inactivate trypsin. Following a medium

rinse and centrifugation, resuspended cells were seeded onto pre-treated mouse or rabbit anti-

Ngfr^{p75} petri dishes and incubated at 37°C for 10 mins. Unbound rat cells were removed with

medium and saved as 'leftover' control cells for scRNA-seq. A cell scraper was used to recover

bound cells that were then subjected to a second round of immunopanning in which purified cells

from 2 rats on each experimental day were combined yielding 2 samples of purified Ngfr^{p75}-

positive rat OECs and 2 samples of 'leftover' control cells. All cells for sequencing were replated

into PLL-coated flasks. Immunopurified Ngfr^{p75}-positive mouse OECs were plated on PLL-

coated 4-chamber polystrene-vessel culture slides (BD Falcon, San Jose, CA) or used for western

blots. Purified OECs and 'leftover' controls were incubated at 37°C with 5% CO_2 for 7-8 days in

D/F-FBS-P/S medium supplemented with a mitogen mixture of pituitary extract (20 μg/ml,

Gibco) and forskolin (2 μM, Sigma). Media was changed every 2 days and mitogens were

withdrawn 2 days prior to harvesting these cells for scRNA-seq.

Brefeldin-A treatment. Rat OECs that contained GFP could be visualized directly. Primary and

purified GFP-expressing OEC cultures were rinsed 3 times with D/F medium prior to treatment

with Brefeldin-A to block protein transport from the endoplasmic reticulum to the Golgi

apparatus and thus block secretion. Brefeldin-A (5 μg/ml; Epicenter, Madison, WI) was mixed

with D/F medium and added to OEC cultures for 2 hrs at 37ºC while the control cultures

received D/F medium only. Cultures were fixed with cold 4% paraformaldehyde in 0.1 M PB for

15 mins at RT, rinsed 3 times with PBS, and stored in PBS with sodium azide at 4ºC until

immunostaining.

Detection of Reelin expression in rat and mouse OEC cultures, tissue sections and western

blots.

Cultured rat OECs treated with or without Brefeldin-A were labeled with mouse anti-Reelin G10

and rabbit anti-Ngfr[p75] (Suppl. Table 4). Culture slides were rinsed with phosphate-buffered

saline (PBS; 0.1M phosphate buffer; 0.9% NaCl), blocked with 5% normal goat serum for 1 h,

and incubated with the appropriate primary antibodies overnight. The following day, slides were

rinsed 3 times with PBS, incubated with species-appropriate Alexa Fluor 594 and/or 647 (1:500,

1:100; Jackson ImmunoResearch) for 1 hr at RT, and then cover slipped with Fluorogel

(Electron Microscopy Sciences, Hatfield, PA).

Sagittal sections of the mouse E16.5 olfactory system were cut 40 μm thick, slide

mounted, and stored in PB with 0.06% sodium azide. For Reelin and Blbp detection, sections

were incubated overnight at RT in goat-anti-Reelin (Suppl. Table 4), rinsed thoroughly, and

incubated with donkey anti-goat 488 (1:100; Jackson Immunoresearch, #705-545-0030).

Sections were then incubated in rabbit polyclonal anti-Blbp (Suppl. Table 4) overnight followed

by 1 hr in donkey anti-rabbit 555 (1:800; Life Technologies, #A31572) and coverslipped with

Fluorogel. Confocal images were obtained from a Zeiss Laser Scanning Microscope (LSM800)

with solid-state lasers 488 and 561 nm for double-labeled images.

Tissue harvesting, protein isolation, and western blotting. Olfactory bulbs from adult $Reln^{+/+}$ and

$Reln^{-/-}$ mice and olfactory nerve layers from adult Sprague Dawley rats were freshly dissected as

described for primary OEC cultures. Olfactory tissue was collected, placed on ice, and

homogenized in Ripa lysis buffer plus protease inhibitor cocktail (Sigma). Protein concentrations

of brain and olfactory extracts were determined using the RC DC Protein Assay kit (Bio-Rad,

Hercules, CA) as described (Miller et al., 2008). Whole-cell lysates (WCL) and conditioned

medium (CM) were obtained from purified OEC cultures maintained in D/F medium. CM was

collected from OEC cultures before cells were incubated with Ripa lysis buffer and collected

with a cell scraper. Protein samples were stored at -80ºC.

Protein samples (50 μg) were heated to 90–100°C for 3 mins, resolved on 10% SDS-

PAGE or 4–15% mini-protean TGX gels (Bio-Rad) run in Tris-Glycine-SDS (TGS) buffer (Bio-

Rad) at 60V, and then transferred onto 0.22 μm PVDF/nitrocellulose membranes (Bio-Rad) in

TGS with 20% methanol at 25V overnight at 4°C. Precision plus protein standards in dual color

(Bio-Rad) were used to determine molecular weight. Membranes were cut, blocked with 5%

non-fat dry skimmed milk (Bio-Rad) in Tris-buffered saline containing 0.1% Tween-20 (TBST)

for 1 hr at room temperature on a shaker, and then incubated overnight at 4°C TBST plus 2.5%

milk containing anti-Reelin G10 or loading control mouse anti-GAPDH antibodies (Suppl. Table

4). Following primary antibody incubation, blots were washed 3 times in TBST for 10 mins

while shaking and then probed with horseradish peroxidase-conjugated secondary anti-mouse

IgG (1:2,000 for Reelin, 1:5,000 for GAPDH) in TBST. Immunoblots were developed using a

chemiluminescence HRP detection kit (GE Healthcare, Piscataway, NJ,) and imaged with a

Typhoon scanner (GE Healthcare).

For quantification, ImageJ software (NIH) was used to analyze the densitometric data.

Western blot images at 400, 300, and 150 kDa resolution were converted to grayscale followed

by manually defining a Region of Interest (ROI) frame that captured the entire band in each lane

using the "Rectangular" tool. The area of each selected band was measured by employing the

same ROI frame around the band to record the integrated density, "Grey Mean

Value". Background measurements were similarly quantified, and background subtraction was

performed by deducting the inverted background from the inverted band value. For relative

quantification, target protein bands were normalized to the corresponding loading control

(GAPDH) to derive normalized protein expression (fold change). Band intensities were

quantified in triplicate for each sample. Data were analyzed with the Mann-Whitney U test to

compare normalized protein expression between the $Reln^{-/-}$ group and the other groups. A one-

sided p-value was calculated to test the hypothesis that protein expression levels in the other

groups are greater than those in the $Reln^{-/-}$ group. Statistical significance was determined at p <

0.05. Analysis was performed using GraphPad Prism (version 9).

Single-cell RNA-sequencing

Cell collection and culture. Flasks with purified OECs and leftover cells were rinsed with HBSS

and dissociated with trypsin-EDTA as above. Cells were washed in D/F-FBS-P/S medium and

further resuspended with siliconized pipettes. Media and cells were washed and centrifuged at

365 *g* for 10 min, and resuspended. After the final wash, cells for sequencing were resuspended

in PBS+BSA in volumes of 1200 cells/µL. On each of 2 different culture days, 4 flasks of cells

(2 purified OECs and 2 'leftovers') were generated for scRNA-seq, yielding 8 samples in total (4

purified OECs and 4 'leftovers'). To confirm our scRNA-seq results directly with

immunocytochemistry, we plated extra cells from the second preparation on PLL-coated

coverslips in 24-well plates with 500µL D/F-FBS-P/S medium. Medium was changed at 2 days,

cultures were fixed at 3 days in cold 4% paraformaldehyde for 20 mins, washed 3X with PB and

stored in PB with 0.06% sodium azide.

769   scRNA-seq was performed using the 10X Genomics Chromium scRNA-seq system

which incorporates microfluidic capture-based encapsulation, barcoding, and library preparation

(10X Genomics, Pleasanton, CA). About 10,000 cells per OEC or leftover preparation were

loaded into a Chromium Chip B along with partitioning oil, reverse transcription reagents, and a

mix of hydrogel beads containing 3,500,000 unique 10X barcodes. Paired-end sequencing was

performed on a Novaseq S4 system, using the v3 Illumina platform, at 20-50K reads/cell. The

paired-end reads were processed using the Cell Ranger pipeline and *Rattus*

norvegicus (Rnor_6.0) from Ensembl as the reference genome (Yates et al., 2020) to produce a

gene expression matrix. Among the 8 samples, one OEC sample showed a low fraction of reads

and RNA content in cells, which may indicate high level of ambient RNA (Suppl. Figure 3a).

This sample was removed from further analysis, leaving 3 OEC samples and 4 leftover samples

for downstream analysis.

scRNA-seq data quality control, normalization, and integration. Gene expression matrix from

each of the 7 remaining samples that passed quality control was loaded into the Seurat R package

ver 3.0 (Stuart et al., 2019). Cells were called based on the number of genes (200-5000), unique

molecular identifiers (UMIs; 5000-20000), and percent of mitochondrial genes (<20%; Suppl.

Figure 3b). After quality control, log normalization was performed within each sample using

NormalizeData function with default parameters. Top 2,000 variable genes were selected using

FindVariableFeatures. The seven samples were integrated together with *FindIntegrationAnchors*

and *IntegrateData* functions which incorporate canonical correlation analysis to align cells with

similar transcriptomic patterns across samples.

Cell clustering and cell type identification. The integrated Seurat object was used for the

principal component analysis (PCA). The top 15 PCs were used to construct the k-nearest

neighbor graph, followed by Louvain algorithm to cluster cells based on similar gene expression

patterns. Cell clusters were visualized using tSNE plots.

To assign cell type identities to each cluster, marker genes of each cluster were found

using *FindAllMarkers* with average log fold change > 0.5 and minimum percent difference >

0.25. Cell cluster marker genes were subsequently compared to known OEC cell type markers

curated from previous studies that used different species, sex, age, dissection areas, and cell

preparation methods (Durante et al., 2020; Tepe et al., 2018). Additional marker genes for

fibroblasts and multiple glial cell types including astrocytes, oligodendrocytes, and microglia

were also used to compare with those of the cell clusters. Cell clusters whose marker genes

match known cell type markers were labeled with the corresponding cell type identity. Gene

expression heatmaps of top markers were generated to visualize the distinct expression patterns

of the marker genes for different cell clusters.

OEC subtype identification. A subset of cells expressing well-known OEC markers (e.g., $Ngfr^{p75}$,

S100β, and *Sox10*) was labeled as OECs and then selected for further subclustering analysis to

identify potential OEC subtypes. To do this, OECs from the original Seurat object were extracted

using the *subset* function and the top 10 PCs of each OEC subset were used for clustering

analysis and tSNE visualization. Marker genes for each OEC subcluster were identified using the

FindAllMarkerGene function. Pseudotime trajectory analysis across the OEC subtypes was

performed using Slingshot version 1.8.0 (Street et al., 2018).

Pathway enrichment analysis. To annotate the biological functions of the marker genes of

individual cell clusters or OEC subclusters, we carried out pathway enrichment analysis using

EnrichR ver 3.0, where KEGG, Reactome, and GO Biological Process were used as reference

pathway databases (Chen et al., 2013). Top pathways were ranked by multiple-testing adjusted

P-values.

Potential cell-cell ligand-receptor interactions across OEC subtypes. In order to characterize

potential interactions mediated by secreted ligand-receptor pairs between subtypes of OECs, we

implemented NicheNet (Browaeys et al., 2020), which curates ligand-receptor interactions from

various publicly available resources. The NicheNet ligand-receptor model was integrated with

markers from each OEC subtype to find potentially activated ligand-receptor pairs. Differentially

expressed ligands identified by markers from source OEC subtypes and activated receptors

identified by NicheNet in target OEC subtypes were used to assess ligand-receptor interaction.

Ligand-receptor interaction networks between cell types were visualized using Cytoscape

(Shannon et al., 2003).

Confirmation of OEC gene expression *in vitro* and *in vivo*.

 Immunocytochemistry experiments to confirm gene expression revealed by scRNA-seq were

conducted on extra cultured OECs and 'leftover' controls which were replated and cultured for 3

additional days. OECs samples were treated with the following primary antibodies against OEC

marker genes identified from scRNA-seq with experimental parameters recorded in Suppl. Table

4: 1) polyclonal and monoclonal anti-Blbp, 2) anti-N-Cadherins, 3) anti-green fluorescent

protein, 4) anti-Ki67, 5) anti-L1, 6) anti-NCAM, 7) monoclonal anti-Ngfr^{p75}, 8) anti-S100β, and

9) anti-SOX10. Prior to adding the primary antibodies, the cells were incubated in 0.3% Triton-X

100 detergent in PBS. Cells were blocked in 5% normal donkey serum (Jackson

ImmunoResearch Laboratories, # 017-000-121) with 0.3% Triton-X before adding the primary

antibodies and left to incubate at RT overnight on the rotator. Following multiple washes,

species-appropriate secondary antibodies with fluorescence for 488, 555, and 647 were used to

visualize the immunoreaction (1:200 – 1:500, Jackson ImmunoResearch Labs donkey-anti-

chicken-488, # 703-545-155; donkey-anti-rabbit-647, # 711-605-152; donkey-anti-mouse-647, #

715-605-150; Life Technologies, donkey-anti-mouse-555, # A31570; donkey-anti-rabbit-555, #

A31572). Cultures were counterstained with Hoechst (Bis-benzimide, 1:500, Sigma-Aldrich, #

B2261) and coverslipped with EverBrite medium (Biotium, # 23001).

        To examine Ctgf expression in the spinal cord lesion site, we processed 1 slide per animal

with ~6 equally-spaced sagittal sections (25 μm thick) throughout spinal cords from the Khankan

et al. (2016) study. Our aim was to assess if transplanted OECs (n=4 rats) and transplanted

fibroblasts (n=3 rats) express Ctgf in the injury site.

Experiments were carried out to confirm gene expression of OEC subtype markers in the

8-10-week-old olfactory system of rats. Soft tissues and the jaw were removed from fixed heads

that were decalcified in undiluted formic acid bone decalcifier (Immunocal, Decal Chemical

Corps, Tallman, NY) for 12 hrs., washed, and then infiltrated with increasing concentrations of

5% to 20% sucrose for 24 hr. Heads in 20% sucrose were heated in a 37°C oven and then placed

in a gelatin sucrose mixture for 4 hrs, embedded in warm gelatin sucrose in a plastic mold,

allowed to gel, frozen on dry ice and stored at -70°C. Heads were sectioned coronally at 40 μm

thick, and mounted in series on 16 glass slides. Several antibodies from each of the 4 OEC

clusters were chosen to determine their distribution in olfactory bulb sections. Primary antibodies

used are listed in Suppl. Table 4, together with the experimental parameters used. For most

antibodies, the olfactory bulb sections first underwent a heat-induced antigen retrieval step with

10mM citric acid at pH 6 for 5 min and then were transferred to 1.5% NDS and 0.1% Triton-X-

100 in TBS-BSA (0.1M Tris; 1.4% NaCl; 0.1% BSA) for 1 h followed by overnight incubation

in primary antibody at RT. Following multiple washes, species-appropriate secondary antibodies

with fluorescence for 488 and 555 were used to visualize the immunoreactivity.

For experiments carried out using a Tyramide Signal Amplification (TSA) kit, sections

were first incubated in 1% hydrogen peroxide and 0.1% sodium azide in phosphate-buffered

saline PBS for 30 mins followed by 10% NDS and 0.1% Triton X-100 in PBS for 1 hr. Next, the

sections were incubated with Avidin-Biotin solution (Vector laboratories; kit #SP-2001) before

an overnight incubation with the primary antibody at RT. The next day, sections were washed in

PBS followed by TNT buffer (0.1M Tris-HCl; 0.15M NaCl; 0.05% Tween) and then a 1 hr

incubation with biotinylated horse anti-goat IgG (1:1000; Vector laboratories; #BA-9500),

biotinylated donkey anti-rabbit IgG (1:1000; Jackson Immunoresearch; #711-065-152), or

biotinylated donkey anti-mouse IgG (1:1000; Jackson Immunoresearch; #715-065-150) in TSA-

specific blocking buffer (TNB; 0.1M Tris-HCl; 0.15M NaCl; 0.5% Blocking reagent;

PerkinElmer; #FP1020). Sections were then washed with TNT followed by a 1 h incubation with

streptavidin-conjugated horseradish peroxidase (1:1000; PerkinElmer; #NEL750001EA) in TNB,

and a 10-min incubation with TSA Plus Fluorescein (1:150; PerkinElmer; #NEL741001KT) or

TSA Plus Cyanine 3 (Cy3; 1:100; PerkinElmer; #NEL744001KT). Sections were imaged on an

Olympus AX70 microscope with a Zeiss AxioCam HRcRv.2 camera. Images were transferred to

Adobe Photoshop for assembly, cropping, and adjustment of brightness/contrast to create the

compiled figures.

Morphological analyses of Ki67 OEC subtypes

To determine if OEC progenitor cells marked with Ki67 immunoreactivity have a distinctive

morphology, purified and fixed OEC cultures from four rats were processed with anti-Ngfr^{p75},

anti-Ki67 and counterstained with Hoechst (Bis-benzimide, 1:500, Sigma-Aldrich, #B2261).

Images were acquired from 7-10 randomly selected fields/sample using an Olympus AX70

microscope and Zen image processing and analysis software (Carl Zeiss). We distinguished the

larger, flat 'astrocyte-like' OECs from the smaller, fusiform 'Schwann cell-like' OECs, and

recorded their expression of Ngfr^{p75} and Ki67. Cell counts from each field were averaged per rat

and then averaged into a group mean ± SEM. A Student t-test was conducted to compare the

effect of Ngfr^{p75}-labeled cell morphology and the proliferative marker Ki67. Statistical

significance was determined by $p < 0.05$.

References

Abadesco, A. D., Cilluffo, M., Yvone, G. M., Carpenter, E. M., Howell, B. W., & Phelps, P. E. (2014). Novel Disabled-1-expressing neurons identified in adult brain and spinal cord. *Eur J Neurosci*, *39*(4), 579-592. https://doi.org/10.1111/ejn.12416

Asan, E., Langenhan, T., Holtmann, B., Bock, H., Sendtner, M., & Carroll, P. (2003). Ciliary neurotrophic factor in the olfactory bulb of rats and mice. *Neuroscience*, *120*(1), 99-112. https://doi.org/10.1016/s0306-4522(03)00211-2

Barraud, P., Seferiadis, A. A., Tyson, L. D., Zwart, M. F., Szabo-Rogers, H. L., Ruhrberg, C., Liu, K. J. & Baker, C. V. H. (2010). Neural crest origin of olfactory ensheathing glia. *Proc. Natl. Acad. Sci. U.S.A.* 107, 21040-21045. https://onlinelibrary.wiley.com/doi/full/10.1002/glia.23870

Boekhoorn, K., van Dis, V., Goedknegt, E., Sobel, A., Lucassen, P. J., & Hoogenraad, C. C. (2014). The microtubule destabilizing protein stathmin controls the transition from dividing neuronal precursors to postmitotic neurons during adult hippocampal neurogenesis. *Dev Neurobiol*, *74*(12), 1226-1242. https://doi.org/10.1002/dneu.22200

Bregman, B. S. (1987) Spinal cord transplants permit the growth of serotonergic axons across the site of neonatal spinal cord transection. *Dev Brain Res 43*(2), 265-270. https//doi.org/10.1016/0165-3806(87)90214-8

Bregman, B. S., Kunkel-Bagden, E., Reier, P. J., Dai, H. N., McAtee, M., and Gao, D. (1993) Recovery of function after spinal cord injury: Mechanisms underlying transplant-mediated recovery of function differ after spinal cord injury in newborn and adult rats. *Expt Neurol 123*(1),3-16. https://doi.org/10.1006/exnr.1993.1136

Browaeys, R., Saelens, W., & Saeys, Y. (2020). NicheNet: modeling intercellular communication by linking ligands to target genes. *Nat Methods*, *17*(2), 159-162. https://doi.org/10.1038/s41592-019-0667-5

Burda, J. E., & Sofroniew, M. V. (2014). Reactive gliosis and the multicellular response to CNS damage and disease. *Neuron*, *81*(2), 229-248. https://doi.org/10.1016/j.neuron.2013.12.034

Chandler, C. E., Parsons, L. M., Hosang, M., & Shooter, E. M. (1984). A monoclonal antibody modulates the interaction of nerve growth factor with PC12 cells. *J Biol Chem*, *259*(11), 6882-6889. https://www.ncbi.nlm.nih.gov/pubmed/6327698

Chen, E. Y., Tan, C. M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G. V., Clark, N. R., & Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics*, *14*, 128. https://doi.org/10.1186/1471-2105-14-128

Christopherson, K. S., Ullian, E. M., Stokes, C. C., Mullowney, C. E., Hell, J. W., Agah, A., Lawler, J., Mosher, D. F., Bornstein, P., & Barres, B. A. (2005). Thrombospondins are astrocyte-secreted proteins that promote CNS synaptogenesis. *Cell*, *120*(3), 421-433. https://doi.org/10.1016/j.cell.2004.12.020

Chung, R. S., Woodhouse, A., Fung, S., Dickson, T. C., West, A. K., Vickers, J. C., & Chuah, M. I. (2004). Olfactory ensheathing cells promote neurite sprouting of injured axons in vitro by direct cellular contact and secretion of soluble factors. *Cell Mol Life Sci*, *61*(10), 1238-1245. https://doi.org/10.1007/s00018-004-4026-y

Conrad, S., Schluesener, H. J., Adibzahdeh, M., & Schwab, J. M. (2005). Spinal cord injury induction of lesional expression of profibrotic and angiogenic connective tissue growth

factor confined to reactive astrocytes, invading fibroblasts and endothelial cells. *J Neurosurg Spine*, *2*(3), 319-326. https://doi.org/10.3171/spi.2005.2.3.0319

Cregg, J. M., DePaul, M. A., Filous, A. R., Lang, B. T., Tran, A., & Silver, J. (2014). Functional regeneration beyond the glial scar. *Exp Neurol*, *253*, 197-207. https://doi.org/10.1016/j.expneurol.2013.12.024

D'Arcangelo, G., Miao, G. G., Chen, S. C., Soares, H. D., Morgan, J. I., & Curran, T. (1995). A protein related to extracellular matrix proteins deleted in the mouse mutant reeler. *Nature*, *374*(6524), 719-723. https://doi.org/10.1038/374719a0

D'Arcangelo, G., Miao, G. G., & Curran, T. (1996). Detection of the reelin breakpoint in reeler mice. *Brain Res Mol Brain Res*, *39*(1-2), 234-236. https://doi.org/10.1016/0169-328x(96)00046-0

Daboussi, L., Costaguta, G., Gullo, M., Jasinski, N., Pessino, V., O'Leary, B., Lettieri, K., Driscoll, S., & Pfaff, S. L. (2023). Mitf is a Schwann cell sensor of axonal integrity that drives nerve repair. *Cell Rep*, *42*(11), 113282. https://doi.org/10.1016/j.celrep.2023.113282

Dairaghi, L., Flannery, E., Giacobini, P., Saglam, A., Saadi, H., Constantin, S., Casoni, F., Howell, B. W., & Wray, S. (2018). Reelin can modulate migration of olfactory ensheathing cells and Gonadotropin releasing hormone neurons via the canonical pathway. *Front Cell Neurosci*, *12*, 228. https://doi.org/10.3389/fncel.2018.00228

Dixie, K. L. I. (2019). Transplantation of olfactory ensheathing cells combined with epidural stimulation and climb training as a long-term treatment for severe spinal cord injury in rodents. *UCLA. ProQuest ID*, *Dixie_ucla_0031D_18445. Merritt ID: ark:/13030/m5ps32vp*.

Doucette, R. (1990). Glial influences on axonal growth in the primary olfactory system. *Glia*, *3*(6), 433-449. https://doi.org/10.1002/glia.440030602

Doucette, R. (1991). PNS-CNS transitional zone of the first cranial nerve. *J Comp Neurol*, *312*(3), 451-466. https://doi.org/10.1002/cne.903120311

Durante, M. A., Kurtenbach, S., Sargi, Z. B., Harbour, J. W., Choi, R., Kurtenbach, S., Goss, G. M., Matsunami, H., & Goldstein, B. J. (2020). Single-cell analysis of olfactory neurogenesis and differentiation in adult humans. *Nat Neurosci*, *23*(3), 323-326. https://doi.org/10.1038/s41593-020-0587-9

Eroglu, C., & Barres, B. A. (2010). Regulation of synaptic connectivity by glia. *Nature*, *468*(7321), 223-231. https://doi.org/10.1038/nature09612

Eshed, Y., Feinberg, K., Poliak, S., Sabanay, H., Sarig-Nadir, O., Spiegel, I., Bermingham, J. R., Jr., & Peles, E. (2005). Gliomedin mediates Schwann cell-axon interaction and the molecular assembly of the nodes of Ranvier. *Neuron*, *47*(2), 215-229. https://doi.org/10.1016/j.neuron.2005.06.026

Forni, P. E., Taylor-Burds, C., Melvin, V. S., Williams, T., & Wray, S. (2011). Neural crest and ectodermal cells intermix in the nasal placode to give rise to GnRH-1 neurons, sensory neurons, and olfactory ensheathing cells. *J Neurosci*, *31*(18), 6915-6927. https://doi.org/10.1523/JNEUROSCI.6087-10.2011

Franceschini, I. A., & Barnett, S. C. (1996). Low-affinity NGF-receptor and E-N-CAM expression define two types of olfactory nerve ensheathing cells that share a common lineage. *Dev Biol*, *173*(1), 327-343. https://doi.org/10.1006/dbio.1996.0027

Franssen, E. H. P., de Bree, F. M., & Verhaagen, J. (2007) Olfactory ensheathing glia: Their

contribution to primary olfactory nervous system regeneration and their regenerative potential following transplantation into the injured spinal cord. *Brain Res Reviews 56,* 236-258. https://doi.org/10.1016/j.brainresrev.2007.07.013

Fu, S. Y., & Gordon, T. (1997). The cellular and molecular basis of peripheral nerve regeneration. *Mol Neurobiol*, *14*(1-2), 67-116. https://doi.org/10.1007/BF02740621

Granger, N., Blamires, H., Franklin, R. J., & Jeffery, N. D. (2012). Autologous olfactory mucosal cell transplants in clinical spinal cord injury: a randomized double-blinded trial in a canine translational model. *Brain*, *135*(Pt 11), 3227-3237. https://doi.org/10.1093/brain/aws268

Graziadei, P. P., & Monti Graziadei, G. A. (1985). Neurogenesis and plasticity of the olfactory sensory neurons. *Ann N Y Acad Sci*, *457*, 127-142. https://doi.org/10.1111/j.1749-6632.1985.tb20802.x

Gresle, M. M., Butzkueven, H., Perreau, V. M., Jonas, A., Xiao, J., Thiem, S., Holmes, F. E., Doherty, W., Soo, P. Y., Binder, M. D., Akkermann, R., Jokubaitis, V. G., Cate, H. S., Marriott, M. P., Gundlach, A. L., Wynick, D., & Kilpatrick, T. J. (2015). Galanin is an autocrine myelin and oligodendrocyte trophic signal induced by leukemia inhibitory factor. *Glia*, *63*(6), 1005-1020. https://doi.org/10.1002/glia.22798

Guerout, N., Derambure, C., Drouot, L., Bon-Mardion, N., Duclos, C., Boyer, O., & Marie, J. P. (2010). Comparative gene expression profiling of olfactory ensheathing cells from olfactory bulb and olfactory mucosa. *Glia*, *58*(13), 1570-1580. https://doi.org/10.1002/glia.21030

Hunt, D., Hossain-Ibrahim, K., Mason, M. R., Coffin, R. S., Lieberman, A. R., Winterbottom, J., & Anderson, P. N. (2004). ATF3 upregulation in glia during Wallerian degeneration: differential expression in peripheral nerves and CNS white matter. *BMC Neurosci*, *5*, 9. https://doi.org/10.1186/1471-2202-5-9

Ingram, N. T., Khankan, R. R., & Phelps, P. E. (2016). Olfactory ensheathing cells express alpha7 integrin to mediate their migration on laminin. *PLoS One*, *11*(4), e0153394. https://doi.org/10.1371/journal.pone.0153394

Jessen, K. R., Morgan, L., Stewart, H. J., & Mirsky, R. (1990). Three markers of adult non-myelin-forming Schwann cells, 217c(Ran-1), A5E3 and GFAP: development and regulation by neuron-Schwann cell interactions. *Development*, *109*(1), 91-103. https://doi.org/10.1242/dev.109.1.91

Khankan, R. R., Griffis, K. G., Haggerty-Skeans, J. R., Zhong, H., Roy, R. R., Edgerton, V. R., & Phelps, P. E. (2016). Olfactory ensheathing cell transplantation after a complete spinal cord transection mediates neuroprotective and immunomodulatory mechanisms to facilitate regeneration. *J Neurosci*, *36*(23), 6269-6286. https://doi.org/10.1523/JNEUROSCI.0085-16.2016

Khankan, R. R., Wanner, I. B., & Phelps, P. E. (2015). Olfactory ensheathing cell-neurite alignment enhances neurite outgrowth in scar-like cultures. *Exp Neurol*, *269*, 93-101. https://doi.org/10.1016/j.expneurol.2015.03.025

Krstic, D., Rodriguez, M., & Knuesel, I. (2012). Regulated proteolytic processing of Reelin through interplay of tissue plasminogen activator (tPA), ADAMTS-4, ADAMTS-5, and their modulators. *PLoS One*, *7*(10), e47793. https://doi.org/10.1371/journal.pone.0047793

Kubasak, M. D., Brooks, R., Chen, S., Villeda, S. A., & Phelps, P. E. (2004). Developmental distribution of Reelin-positive cells and their secreted product in the rodent spinal cord. *J Comp Neurol*, *468*(2), 165-178. https://doi.org/10.1002/cne.10946

Lakatos, A., Barnett, S. C., & Franklin, R. J. (2003). Olfactory ensheathing cells induce less host astrocyte response and chondroitin sulphate proteoglycan expression than Schwann cells following transplantation into adult CNS white matter. *Exp Neurol*, *184*(1), 237-246. https://doi.org/10.1016/s0014-4886(03)00270-x

Lakatos, A., Franklin, R. J., & Barnett, S. C. (2000). Olfactory ensheathing cells and Schwann cells differ in their in vitro interactions with astrocytes. *Glia*, *32*(3), 214-225. https://doi.org/10.1002/1098-1136(200012)32:3

Lambert de Rouvroit, C., de Bergeyck, V., Cortvrindt, C., Bar, I., Eeckhout, Y., & Goffinet, A. M. (1999). Reelin, the extracellular matrix protein deficient in reeler mutant mice, is processed by a metalloproteinase. *Exp Neurol*, *156*(1), 214-217. https://doi.org/10.1006/exnr.1998.7007

Lamond, R., & Barnett, S. C. (2013). Schwann cells but not olfactory ensheathing cells inhibit CNS myelination via the secretion of connective tissue growth factor. *J Neurosci*, *33*(47), 18686-18697. https://doi.org/10.1523/JNEUROSCI.3233-13.2013

Lane, M. A., Truettner, J. S., Brunschwig, J-P., Gomez, A., Bunge, M. B., Dietrich, W. D., Dziegielewska, K. M., Ek, C. J., VandeBerg, J. L., & Saunders, N. R. (2007) Age-related differences in the local cellular and molecular responses to injury in developing spinal cord of the opossum, *Monodelphis domestica*. *Eur J Neurosci 25(6),* 1725-1742. https://doi.org/10.1111/j.1460-9568.2007.05439.x

Leung, J. Y., Chapman, J. A., Harris, J. A., Hale, D., Chung, R. S., West, A. K., & Chuah, M. I. (2008). Olfactory ensheathing cells are attracted to, and can endocytose, bacteria. *Cell Mol Life Sci*, *65*(17), 2732-2739. https://doi.org/10.1007/s00018-008-8184-1

Li, Y., Li, D., & Raisman, G. (2005). Interaction of olfactory ensheathing cells with astrocytes may be the key to repair of tract injuries in the spinal cord: the 'pathway hypothesis'. *J Neurocytol*, *34*(3-5), 343-351. https://doi.org/10.1007/s11068-005-8361-1

Martini, R., & Schachner, M. (1988). Immunoelectron microscopic localization of neural cell adhesion molecules (L1, N-CAM, and myelin-associated glycoprotein) in regenerating adult mouse sciatic nerve. *J Cell Biol*, *106*(5), 1735-1746. https://doi.org/10.1083/jcb.106.5.1735

Meyer, M., Matsuoka, I., Wetmore, C., Olson, L., & Thoenen, H. (1992). Enhanced synthesis of brain-derived neurotrophic factor in the lesioned peripheral nerve: different mechanisms are responsible for the regulation of BDNF and NGF mRNA. *J Cell Biol*, *119*(1), 45-54. https://doi.org/10.1083/jcb.119.1.45

Miller, J. E., Spiteri, E., Condro, M. C., Dosumu-Johnson, R. T., Geschwind, D. H., & White, S. A. (2008). Birdsong decreases protein levels of FoxP2, a molecule required for human speech. *J Neurophysiol*, *100*(4), 2015-2025. https://doi.org/10.1152/jn.90415.2008

Misumi, Y., Misumi, Y., Miki, K., Takatsuki, A., Tamura, G., & Ikehara, Y. (1986). Novel blockade by brefeldin A of intracellular transport of secretory proteins in cultured rat hepatocytes. *J Biol Chem*, *261*(24), 11398-11403. https://www.ncbi.nlm.nih.gov/pubmed/2426273

Mokalled, M. H., Patra, C., Dickson, A. L., Endo, T., Stainier, D. Y., & Poss, K. D. (2016). Injury-induced ctgfa directs glial bridging and spinal cord regeneration in zebrafish. *Science*, *354*(6312), 630-634. https://doi.org/10.1126/science.aaf2679

Nakhjavan-Shahraki, B., Yousefifard, M., Rahimi-Movaghar, V., Baikpour, M., Nasirinezhad, F., Safari, S., Yaseri, M., Jafari, A. M., Ghelichkhani, P., Tafakhori, A. & Hosseini, M. (2018) Transplantation of olfactory ensheathing cells on functional recovery and

neuropathic pain after spinal cord injury; systematic review and meta-analysis. *Scientific*
Reports 8:325. https://doi.10.1038/s41598-017-18754-4

National Spinal Cord Statistical Center. (2024) Traumatic Spinal Cord Injury Facts and Figures
at a Glance -2024. https://sites.uab.edu/nscisc/

Nazareth, L., Lineburg, K. E., Chuah, M. I., Tello Velasquez, J., Chehrehasa, F., St John, J. A.,
& Ekberg, J. A. (2015). Olfactory ensheathing cells are the main phagocytic cells that
remove axon debris during early development of the olfactory system. *J Comp Neurol*,
523(3), 479-494. https://doi.org/10.1002/cne.23694

Nazareth, L., Shelper, T. B., Chacko, A., Basu, S., Delbaz, A., Lee, J. Y. P., Chen, M., St John, J.
A., & Ekberg, J. A. K. (2020). Key differences between olfactory ensheathing cells and
Schwann cells regarding phagocytosis of necrotic cells: implications for transplantation
therapies. *Sci Rep*, *10*(1), 18936. https://doi.org/10.1038/s41598-020-75850-8

Panteri, R., Mey, J., Zhelyaznik, N., D'Altocolle, A., Fa`, A. D., Gangitano, C. M., Marino, R.,
Lorenzetto, E., Buffelli, M., and Keller, F. (2006). Reelin is transiently expressed in the
peripheral nerve during development and is upregulated following nerve crush. *Mol. Cell.*
Neurosci., *32*, 133-142. https://doi.org/10.1016/j.mcn.2006.03.004

Radtke, C., Sasaki, M., Lankford, K. L., Gallo, V., & Kocsis, J. D. (2011). CNPase expression in
olfactory ensheathing cells. *J Biomed Biotechnol*, *2011*, 608496.
https://doi.org/10.1155/2011/608496

Ramon-Cueto, A., Cordero, M. I., Santos-Benito, F. F., & Avila, J. (2000). Functional recovery
of paraplegic rats and motor axon regeneration in their spinal cords by olfactory
ensheathing glia. *Neuron*, *25*(2), 425-435. https://doi.org/10.1016/s0896-6273(00)80905-
8

Ramon-Cueto, A., & Valverde, F. (1995). Olfactory bulb ensheathing glia: a unique cell type
with axonal growth-promoting properties. *Glia*, *14*(3), 163-173.
https://doi.org/10.1002/glia.440140302

Roet, K. C., Bossers, K., Franssen, E. H., Ruitenberg, M. J., & Verhaagen, J. (2011). A meta-
analysis of microarray-based gene expression studies of olfactory bulb-derived olfactory
ensheathing cells. *Exp Neurol*, *229*(1), 10-45.
https://doi.org/10.1016/j.expneurol.2011.03.001

Ruitenberg, M. J., Plant, G. W., Hamers, F. P., Wortel, J., Blits, B., Dijkhuizen, P. A., Gispen,
W. H., Boer, G. J., & Verhaagen, J. (2003). Ex vivo adenoviral vector-mediated
neurotrophin gene transfer to olfactory ensheathing glia: effects on rubrospinal tract
regeneration, lesion size, and functional recovery after implantation in the injured rat
spinal cord. *J Neurosci*, *23*(18), 7045-7058.
https://www.ncbi.nlm.nih.gov/pubmed/12904465

Runyan, S. A., & Phelps, P. E. (2009). Mouse olfactory ensheathing glia enhance axon
outgrowth on a myelin substrate in vitro. *Exp Neurol*, *216*(1), 95-104.
https://doi.org/10.1016/j.expneurol.2008.11.015

Sasaki, M., Black, J. A., Lankford, K. L., Tokuno, H. A., Waxman, S. G., & Kocsis, J. D. (2006).
Molecular reconstruction of nodes of Ranvier after remyelination by transplanted
olfactory ensheathing cells in the demyelinated spinal cord. *J Neurosci*, *26*(6), 1803-
1812. https://doi.org/10.1523/JNEUROSCI.3611-05.2006

Schnaufer, C., Breer, H., & Fleischer, J. (2009). Outgrowing olfactory axons contain the Reelin
receptor VLDLR and navigate through the Reelin-rich cribriform mesenchyme. *Cell*
Tissue Res, *337*(3), 393-406. https://doi.org/10.1007/s00441-009-0762-5

Schwab, J. M., Beschorner, R., Nguyen, T. D., Meyermann, R., & Schluesener, H. J. (2001). Differential cellular accumulation of connective tissue growth factor defines a subset of reactive astrocytes, invading fibroblasts, and endothelial cells following central nervous system injury in rats and humans. *J Neurotrauma*, *18*(4), 377-388. https://doi.org/10.1089/089771501750170930

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N., Schwikowski, B., & Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res*, *13*(11), 2498-2504. https://doi.org/10.1101/gr.1239303

Smithson, L. J., & Kawaja, M. D. (2010). Microglial/macrophage cells in mammalian olfactory nerve fascicles. *J Neurosci Res*, *88*(4), 858-865. https://doi.org/10.1002/jnr.22254

Street, K., Risso, D., Fletcher, R. B., Das, D., Ngai, J., Yosef, N., Purdom, E., & Dudoit, S. (2018). Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC Genomics*, *19*(1), 477. https://doi.org/10.1186/s12864-018-4772-0

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., 3rd, Hao, Y., Stoeckius, M., Smibert, P., & Satija, R. (2019). Comprehensive integration of single-cell data. *Cell*, *177*(7), 1888-1902 e1821. https://doi.org/10.1016/j.cell.2019.05.031

Su, Z., Chen, J., Qiu, Y., Yuan, Y., Zhu, F., Zhu, Y., Liu, X., Pu, Y., & He, C. (2013). Olfactory ensheathing cells: the primary innate immunocytes in the olfactory pathway to engulf apoptotic olfactory nerve debris. *Glia*, *61*(4), 490-503. https://doi.org/10.1002/glia.22450

Tabakow, P., Raisman, G., Fortuna, W., Czyz, M., Huber, J., Li, D., Szewczyk, P., Okurowski, S., Miedzybrodzki, R., Czapiga, B., Salomon, B., Halon, A., Li, Y., Lipiec, J., Kulczyk, A., & Jarmundowicz, W. (2014). Functional regeneration of supraspinal connections in a patient with transected spinal cord following transplantation of bulbar olfactory ensheathing cells with peripheral nerve bridging. *Cell Transplant*, *23*(12), 1631-1655. https://doi.org/10.3727/096368914X685131

Takeoka, A., Jindrich, D. L., Munoz-Quiles, C., Zhong, H., van den Brand, R., Pham, D. L., Ziegler, M. D., Ramon-Cueto, A., Roy, R. R., Edgerton, V. R., & Phelps, P. E. (2011). Axon regeneration can facilitate or suppress hindlimb function after olfactory ensheathing glia transplantation. *J Neurosci*, *31*(11), 4298-4310. https://doi.org/10.1523/JNEUROSCI.4967-10.2011

Tepe, B., Hill, M. C., Pekarek, B. T., Hunt, P. J., Martin, T. J., Martin, J. F., & Arenkiel, B. R. (2018). Single-cell RNA-Seq of mouse olfactory bulb reveals cellular heterogeneity and activity-dependent molecular census of adult-born neurons. *Cell Rep*, *25*(10), 2689-2703 e2683. https://doi.org/10.1016/j.celrep.2018.11.034

Thornton, M. A., Mehta, M. D., Morad, T. T., Ingraham, K. L., Khankan, R. R., Griffis, K. G., Yeung, A. K., Zhong, H., Roy, R. R., Edgerton, V. R., & Phelps, P. E. (2018). Evidence of axon connectivity across a spinal cord transection in rats treated with epidural stimulation and motor training combined with olfactory ensheathing cell transplantation. *Exp Neurol*, *309*, 119-133. https://doi.org/10.1016/j.expneurol.2018.07.015

Trupp, M., Ryden, M., Jornvall, H., Funakoshi, H., Timmusk, T., Arenas, E., & Ibanez, C. F. (1995). Peripheral expression and biological activities of GDNF, a new neurotrophic factor for avian and mammalian peripheral neurons. *J Cell Biol*, *130*(1), 137-148. https://doi.org/10.1083/jcb.130.1.137

Ubink, R., Calza, L., & Hokfelt, T. (2003). 'Neuro'-peptides in glia: focus on NPY and galanin. *Trends Neurosci*, *26*(11), 604-609. https://doi.org/10.1016/j.tins.2003.09.003

Vincent, A. J., Choi-Lundberg, D. L., Harris, J. A., West, A. K., & Chuah, M. I. (2007). Bacteria and PAMPs activate nuclear factor kappaB and Gro production in a subset of olfactory ensheathing cells and astrocytes but not in Schwann cells. *Glia*, *55*(9), 905-916. https://doi.org/10.1002/glia.20512

Vincent, A. J., West, A. K., & Chuah, M. I. (2005). Morphological and functional plasticity of olfactory ensheathing cells. *J Neurocytol*, *34*(1-2), 65-80. https://doi.org/10.1007/s11068-005-5048-6

Wanner, I. B., Anderson, M. A., Song, B., Levine, J., Fernandez, A., Gray-Thompson, Z., Ao, Y., & Sofroniew, M. V. (2013). Glial scar borders are formed by newly proliferated, elongated astrocytes that interact to corral inflammatory and fibrotic cells via STAT3-dependent mechanisms after spinal cord injury. *J Neurosci*, *33*(31), 12870-12886. https://doi.org/10.1523/JNEUROSCI.2121-13.2013

Watzlawick, R., Rind, J., Sena, E. S., Brommer, B., Zhang, T., Kopp, M. A., Dirnagi, U., Macleod, M. M., Howells, D. Q.,Schwab, J. M. (2016) Olfactory ensheathing cell transplantation in experimental spinal cord injury: Effect size and reporting bias of 62 experimental treatments:A systematic review and meta-analysis. *PLOS Biol*, 14(5): e1002468. https://doi.org/10.1371/journal.pbio.1002468

Williams, S. K., Franklin, R. J., & Barnett, S. C. (2004). Response of olfactory ensheathing cells to the degeneration and regeneration of the peripheral olfactory system and the involvement of the neuregulins. *J Comp Neurol*, *470*(1), 50-62. https://doi.org/10.1002/cne.11045

Windus, L. C., Claxton, C., Allen, C. L., Key, B., & St John, J. A. (2007). Motile membrane protrusions regulate cell-cell adhesion and migration of olfactory ensheathing glia. *Glia*, *55*(16), 1708-1719. https://doi.org/10.1002/glia.20586

Witheford, M., Westendorf, K., & Roskams, A. J. (2013). Olfactory ensheathing cells promote corticospinal axonal outgrowth by a L1 CAM-dependent mechanism. *Glia*, *61*(11), 1873-1889. https://doi.org/10.1002/glia.22564

Woodhall, E., West, A. K., & Chuah, M. I. (2001). Cultured olfactory ensheathing cells express nerve growth factor, brain-derived neurotrophic factor, glia cell line-derived neurotrophic factor and their receptors. *Brain Res Mol Brain Res*, *88*(1-2), 203-213. https://doi.org/10.1016/s0169-328x(01)00044-4

Yates, A. D., Achuthan, P., Akanni, W., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M. R., Armean, I. M., Azov, A. G., Bennett, R., Bhai, J., Billis, K., Boddu, S., Marugan, J. C., Cummins, C., Davidson, C., Dodiya, K., Fatima, R., Gall, A., . . . Flicek, P. (2020). Ensembl 2020. *Nucleic Acids Res*, *48*(D1), D682-D688. https://doi.org/10.1093/nar/gkz966

You, S., Petrov, T., Chung, P. H., & Gordon, T. (1997). The expression of the low affinity nerve growth factor receptor in long-term denervated Schwann cells. *Glia*, *20*(2), 87-100. https://doi.org/10.1002/(sici)1098-1136(199706)20:2

 Figure legends

 Figure 1. Transplanted GFP-OECs in the center of the lesion core associate with numerous

 axons. Sagittal sections show rostral and caudal glial scar borders of the injury site which are

 identified with glial fibrillary acidic protein (GFAP, blue). The GFAP-negative lesion core

 contains GFP-OECs (green) and is marked with asterisks. Arrowheads mark axons crossing into

 the lesion core. **(a-f)** Serotonergic axons (5-HT, red) are found in the rostral spinal cord stump

 and associate with OECs (green) in the lesion core. Single channels for GFAP **(c)**, OECs **(d)**, 5-

 HT **(e),** and a combination of 5-HT and OECs **(f)**. **(g-l)** Nearby injury site section from the same

 rat **(a)**. Numerous neurofilament-positive axons (white) are associated with the OECs (green) in

 the lesion core. Single channels for GFAP **(i)**, OECs **(j)**, neurofilament **(k),** and a combination of

 neurofilament and OECs **(l)**. Bridge formation across the injury site was observed in 2 of 8 OEC-

 transplanted and 0 of 11 fibroblast- or media-transplanted, spinal cord transected rats. Combined

 with the 1/5 OEC-transplanted rats with axons crossing the injury and 0/5 fibroblast controls in

 our previous study (Thornton, 2018), we observed bridges in 3/13 OEC-transplanted rats vs 0/16

 controls (p=0.042, two-sample proportion test). Scale bars: a, g = 500 µm; b, h = 100 µm.

 Reprinted from: Dixie, KL, 2019, *UCLA*. ProQuest ID: Dixie_ucla_0031D_18445.

 Figure 2. scRNA-seq results show distinct clusters of OECs and leftover cell samples.

 (a) Cells in tSNE plot colored by sample source, with cells from leftover samples in pink and

 cells from immunopurified OECs in cyan. **(b)** Clustering analysis revealed a total of 7 distinct

 cell clusters, each indicated with a different color. **(c)** A heatmap showing expression patterns of

 top marker genes of the 8 individual cell clusters. **(d)** Cell clusters showed distinct expression

 patterns for known cell type markers for fibroblasts, microglia, and OECs. Clusters 0, 1, and 5

 had high expression of fibroblast markers and are labeled as fibroblast in the Y-axis; clusters 4,

6, and 7 showed high expression of microglia markers and are labeled as microglia; clusters 2, 3,

and 7 showed high expression of OEC markers and are labeled as OEC. Known marker genes for

different cell types are on the X-axis. Feature plots and violin plots for select marker genes are in

Suppl. Figure 1. **(e)** Based on known cell type markers, cell clusters in tSNE plot were labeled

with the corresponding cell types. **(f)** The expression of cell-type specific marker genes is

depicted in a dot plot. **(g)** Genes that distinguish purified OECs vs OECs in leftover cultures are

shown. The top 3 genes were higher in purified OECs, whereas the bottom 3 genes were higher

in 'leftover' cultures.

1245

Figure 3. Well-established OEC markers are revealed by scRNA-seq and

immunofluorescence.

OEC cultures were replated from extra cells prepared for scRNA-seq. Immunolabeled OECs are

marked with arrows and all cell nuclei are stained with Hoechst (blue nuclei). tSNE maps of the

gene expression in the 5 clusters are shown next to the protein expression in a-l. **(a, b)** Cultured

OECs express Ngfr^{p75} protein and *Ngfr^{p75}* gene expression. **(c, d)** Blbp and Sox10

immunoreactive OECs with *Fabp7* gene expression. **(e, f)** S100β-labeled OECs together with

S100β expression. **(g, h)** L1cam and Sox 10 labeling next to *L1cam* expression. **(i, j)** Ncam1

protein and gene expression**. (k, l)** N-Cadherin and Sox10 markers with *Cdh2* expression. **(m-p)**

scRNA-seq data for *Sox10, Cntf, Itga7*, and *Mpz* (references in text). Scale bars: a, c, e, g, i, k =

50 μm.

1257

Figure 4. OEC subclusters, marker genes, and enriched pathways.

(a) Clustering analysis of OECs revealed five separate clusters (0-4). **(b)** Heatmap depicts the

expression patterns of known marker genes of glial cell types (x-axis) versus microglia, all OECs

and OEC subclusters (y-axis). OEC subclusters express select markers of other glial cell types.

(c) Heatmap depicts the top five marker genes (y-axis) for each OEC subcluster (x-axis). **(d)**

Pathways associated with marker genes of different OEC clusters are shown. The dashed line

indicates the false discovery rate (FDR) <0.05 in the pathway analysis. **(e)** Dot plot showing that

cluster 3 has higher potential lysosomal function based on lysosome pathway genes than the

other clusters. **(f)** Both cluster 3 and 4 express select genes involved in positive regulation of cell

migration. **(g)** Trajectory analysis reveals two trajectories, one including subclusters 2, 0, and 4,

and another involving subclusters 2, 0, 1, and 3. **(h)** NicheNet ligand-receptor network plot

demonstrates intracellular communication between OEC subtypes. Hexagons represent

subclusters 0 to 4 and circles indicate ligands secreted from subclusters. Edges (arrows) point to

clusters where receptors of the ligands are expressed. Different colors represent different OEC

subclusters.

1273

Figure 5. Confirmation of *Ccn2/Ctgf* (Connective tissue growth factor) in cultured OECs

and following OEC implantation after spinal cord injury.

(a, b) The scRNA-seq plots of the 9th (*Ccn2/Ctgf)* and 4th (*Ccn3/Nov*) highest-ranked marker

genes are strongly expressed in subcluster 0. **(c, c1, c2)** The well-known matricellular protein

Connective tissue growth factor (Ctgf) identifies OECs in cell cultures (red, c, c2), and Sox10

nuclear expression (white nuclei, c, c2) confirms they are neural crest-derived cells. **(d, d1, d2)**

A sagittal section from a rat that received a complete spinal cord transection followed by Green

Fluorescent Protein (GFP)-OEC implantation was fixed 2 weeks postinjury. Glial fibrotic acidic

protein (d, Gfap, blue) marks the edges of the borders of the glial scar. GFP-OECs (green) that

express Ctgf (red) in the injury site are outlined by the box in d. High expression of Ctgf is

detected by GFP-OECs that bridge part of the injury site in d1. The single channel of Ctgf is

shown in d2. Scale bars: c-c2 = 50 μm, d-d2 = 250 μm.

1286

Figure 6. Subcluster 2 is characterized by cell cycle and proliferative markers.

(a-f) These scRNA-seq plots show high expression of cell proliferation markers in cluster 2,

supporting their function as OEC progenitor cells. Only *Stmn1* (e, Stathmin, a microtubule

destabilizing protein) is broadly expressed across all five clusters. **(g-g2, h-h2)** Most cultured

OECs are spindle-shaped and have high Ngfr^{p75} expression (red, white arrows). Of the OEC

progenitors that express Ki67, 76% ± 8 of them display low levels of Ngfr^{p75} immunoreactivity

and a "flat" morphology (g2, h2; green nuclei, arrowheads). The remainder of Ki67-expressing

OECs express high levels of Ngfr^{p75} and are fusiform in shape (24% ± 8%, n=4 cultures,

Student-t test, p= 0.023). Hoechst marks all nuclei (g1, h1, blue nuclei, arrowheads). Scale bars:

g, h = 50 μm.

1297

Figure 7. Spatial confirmation of the defined OECs subclusters within the olfactory nerve

layer.

The protein expression of a number of top 20 genes from this scRNA-seq study of purified OEC

cultures is verified in olfactory bulb sections. In all images, the olfactory nerve layer (ONL, layer

I) is at the bottom of the image, the glomeruli (GL, layer II) next, and the remainder of the

olfactory bulb toward the top. Suppl. figure 2 illustrates the corresponding tSNE plots for each

gene in Figure 7. **(a-c)** The top gene in the largest cluster, Peripheral myelin protein 22, is highly

expressed throughout the ONL. Gliomedin, the third-ranked gene in cluster 0, is detected as

small discrete dot-like structures overlaying the ONL (b, box enlarged in c). **(d-f)** OECs in

cluster 1 are immunolabeled by antibodies against the axonal growth factor Atf3 (d) and the

intermediate filament Nestin (f). High levels of Gap43 in the ONL (e) are due to expression by

OECs and axons of immature olfactory sensory neurons. **(g-i)** Strong immunoreactivity of

Stathmin-1 in the ONL reflects labeled axons from immature olfactory sensory neurons and

OECs. Ube2c, a G2/M cell cycle regulator, is expressed in a small number of cells in the ONL,

whereas Mki67-labeled cells are widespread. **(j-l)** The immune function of this cluster is

confirmed by antibodies against Apoe (j), Anxa3 (k), and Aif1/Iba1, markers expressed by

microglia and macrophages. Scale bars: a, b, d-l = 50 µm; c, insets in j-l = 25 µm.

1315

Figure 8. Reelin-signaling pathway marker genes in OEC clusters and Reelin

immunoreactivity in embryonic olfactory system.

(a-f) These six tSNE plots show OEC gene expression levels associated with Reelin signaling.

Reln is highly expressed in most OEC clusters (a), *Disabled-1* (*Dab1*, b), and *Apolipoprotein E*

receptor 2 (*Apoer2*, c) show little expression, and *Very-low-density lipoprotein receptor* has

moderate expression (*Vldlr*, d). The serine protease tissue Plasminogen Activator (e, *tPA*) is

highly expressed in all OEC subsets and cleaves secreted Reelin only at its C-terminus.

ADAMTS-4 cleaves both the N- and C-terminals of Reelin, and is expressed at low levels by

OECs (f). **(g-g2)** Sagittal section of the olfactory epithelium (OE) and olfactory bulb from an

E16.5 *Reln*$^{+/+}$ mouse immunolabeled for Blbp (g, g1; red) and Reelin (g, g2; green). Boxed area

in g is enlarged in g1 and g2. Axons of olfactory sensory neurons (arrows) are surrounded by

peripheral OECs that express both Blbp (g1, arrowheads) and Reelin (g2, arrowheads). High

Reln expression is obvious in olfactory bulb neurons (g, g1, green) but the olfactory nerve layer

(ONL) only expresses Blbp (g, g2). **(h, h1)** No Reelin is detected in a sagittal section from a

Reln$^{-/-}$ mouse, yet Blbp expression in the ONL appears normal. Scale bars: g-h = 50 µm; g1, g2 =

50 µm.

1332

Figure 9. Reelin is expressed and secreted by olfactory ensheathing cells.

(a-c) Rat OEC cultures were treated with Brefeldin-A to inhibit protein transport and subsequent

secretion. Reelin immunoreactivity (red) in a primary culture was detected in GFP-labeled OECs

(a, arrows), but not in another cell type (arrowhead). GFP-labeled OECs which were

immunopurified with anti-Ngfr^{p75} also express Reelin (red; b, c). **(d)** Western blot confirms the

expression of Reelin in rat Olfactory Nerve Layer I and II (ONL; lane 1 of western blot). *Reln*$^{+/+}$

and *Reln*$^{-/-}$ mouse olfactory bulbs were used as positive and negative controls, respectively

(lanes: 2 and 3). Reelin synthesized by cultured OECs was found at low levels in whole cell

lysates (WCL; lanes: 4, 6, and 8), compared to Reelin secreted by cultured OECs into tissue

culture medium, termed OEC "conditioned medium" (CM; lanes: 5 and 7). GAPDH was the

loading control for tissue homogenates (lanes 1-4, 6, 8). **(e)** All three Reelin isoforms (400, 300,

and 150 kDa) were detected with a 4-15% gradient gel. *Reln*$^{+/+}$ and *Reln*$^{-/-}$ mouse cortices were

used as controls (lanes: 1 and 2). Reelin was detected in the rat ONL (lane 3) and in three rat

OEC-CM samples (lanes: 4, 5, and 6). GAPDH was the loading control for tissue homogenates

(lanes: 1-3). **(f)** Quantitation of multiple isoforms of Reelin from 4-15% gradient gels. Positive

and negative controls are *Reln*$^{+/+}$ and *Reln*$^{-/-}$ mouse cortices. Both rat tissue from the ONL (n=3)

and CM (n=9) contain significantly more 400 and 300 kDa Reelin than the *Reln*$^{-/-}$ mice. Bars

represent the standard deviation of the mean. One-sided Mann-Whitney U test was used to test

that protein expression levels in the other groups are greater than those in the *Reln*$^{-/-}$ group,

indicative of significant expression of *Reln* in the test groups. *$p < 0.05$. Scale bars: a-c = 40 µm.

Supplemental Figures

Figure S1. Marker genes were expressed specifically in different areas associated with

Figure 2.

(**a-c**) Marker genes characteristic of OECs (**a**), Microglia (**b**), and fibroblasts (**c**) are shown in

their tSNE plots (top) and violin plots (bottom).

1358

Figure S2. The tSNE plots corresponding to the OEC subclusters illustrated in Figure 7.

(**a-b**) Cluster 0 tSNE plots show that *Pmp22* is expressed in all 5 clusters with *Gldn* concentrated

in clusters 0, 2, and 4. (**d-f**) For cluster 1, tSNE plots illustrate a wide distribution of *Atf3* and

Nes, whereas *GAP43* is found primarily in cluster 1. (**g-i**) The cell cycle and proliferative

markers, *Ube2c* and *Mki67*, appear limited to cluster 2, but *Stmn1* is detected in all clusters but

most heavily concentrated in cluster 2. (**j-l**) Cluster 3 tSNE plots show high expression of *Apoe*,

Anxa3, and *Aif1*, but *Apoe* is also expressed in other clusters.

1366

1367 **Figure S3. Violin plots for quality control**

1368 (**a**) Before quality control measurement. (**b**) After quality control measurement. nFeature_RNA

1369 indicates gene counts; nCount RNA indicates unique molecular identifier (UMI) counts;

1370 percent.mt indicates mitochondria content. Samples (x-axis) are labeled by concatenation of

1371 batch (1 or 2), type (OEC or leftover), and sample ID.

1372

(a) Ngfr^p75 / Hoechst

(b) *Ngfr* (Ngfr^p75) Expression

(c) Blbp / Sox10 / Hoechst

(d) *Fabp7* (Blbp) Expression

(e) s100β / Hoechst

(f) *S100β* Expression

(g) L1cam / Sox10 / Hoechst

(h) *L1cam* Expression

(i) Ncam1 / Hoechst

(j) *Ncam1* Expression

(k) N-cadherin / Sox10 / Hoechst

(l) *Cdh2* (N-cadherin) Expression

(m) *Sox10* Expression

(n) *Cntf* Expression

(o) *Itga7* Expression

(p) *Mpz* (P0) Expression

(a) *Ccn2* Expression

(b) *Ccn3* Expression

(c) CTGF SOX10 Hoechst

(c1) SOX10

(c2) CTGF

(d) CTGF GFAP GFP-OEC

(d1) CTGF GFP-OEC

(d2) CTGF

(a) *Cdk1* Expression

(b) *Cenpf* Expression

(c) *Prc1* Expression

(d) *Ube2c* Expression

(e) *Stmn1* Expression

(f) *Mki67* Expression

(a) *Reln* Expression

(b) *Dab1* Expression

(c) *Apoer2* (*Lrp8*) expression

(d) *Vldlr* Expression

(e) *tPA* tissue Plasminogen activator

(f) *ADAMTS4* Expression

(g) *Reln* $^{+/+}$

ONL

OE

Blbp
Reelin

(h) *Reln* $^{-/-}$

ONL

OE

Blbp
Reelin

(g1)

(g2)

(h1)