Plant metabolites modulate animal social networks and lifespan

Pragya Singh[1,*], Leon Brueggemann[1], Steven Janz[1], Yasmina Saidi[1], Gaurav Baruah[2] and

Caroline Müller[1]

[1]Chemical Ecology, Bielefeld University, Universitätsstr. 25, 33615 Bielefeld, Germany

[2]Theoretical Biology, Bielefeld University, Universitätsstr. 25, 33615 Bielefeld, Germany

*Corresponding Author:

Email: pragya.singh42019@gmail.com

ORCiD: https://orcid.org/0000-0002-7411-3206

Acknowledgements

This study was funded by the German Research Foundation (DFG) as part of the SFB TRR

212 (NC³), project number 396777467 (granted to CM). We thank Karin Djendouci for her

help in conducting the experiments and Rohit Sasidharan for help with collecting sawflies.

Author Contributions

Conceptualization and experimental design: PS; data collection: YS for experiment 2, SJ for

experiment 3, PS for other experiments; chemical analysis: LB; data validation and analysis:

PS and GB for network analysis, PS for other experiments; data visualization: PS; writing

original draft: PS, CM; reviewing and editing: PS, CM, LB, GB; funding acquisition: CM.

Abstract

Social interactions are prevalent in animals, with some individuals participating in more interactions than others. However, the mechanisms underlying this heterogeneity in social interaction frequency and its potential fitness consequences remain poorly characterized. Here, we examine the effects of non-nutritive plant metabolites ('clerodanoids') on social networks and lifespan in the turnip sawfly. The adult sawflies facultatively acquire clerodanoids from non-host plant species. We found high variation in clerodanoid amounts in both wild-caught and lab-reared individuals. Furthermore, sawflies could acquire clerodanoids by nibbling on conspecifics that had prior access to these metabolites, with asymmetric clerodanoid access leading to increased agonistic interactions. Network analysis revealed more social interactions in sawfly groups where some or all individuals had prior access to clerodanoids, while groups with no prior access had fewer interactions. The number of social interactions depended on the clerodanoid status of the focal individual and also of other conspecifics. Finally, we observed a shorter lifespan in adults with prior clerodanoid access when paired with individuals without prior access, suggesting that social interactions to obtain clerodanoids have fitness costs. Our findings demonstrate that individual variation in non-nutritional sequestered plant metabolites can skew social interactions, with potential fitness implications that impact individual social niches.

Keywords

Athalia rosae, mating behaviour, aggressive behaviour, pharmacophagy, specialized plant metabolites, non-nutritional plant-insect interactions

Introduction

Social interactions are widespread in animals (Frank 2007) and lead to the emergence of diverse social networks in groups, from nominally to highly interactive. Such social interactions can range from mating to agonistic. A common aspect of animal social networks is skewness in the number of interactions individuals have, with some individuals having many interactions while others have few (Fisher *et al.* 2019; Krause *et al.* 2015). More or few social interactions may have associated benefits or costs for an individual (Koto *et al.* 2023; Ruan & Wu 2008; Snyder-Mackler *et al.* 2020). For example, increased agonistic social interactions can lead to injuries or even mortality (Archer 1988), although such costs are more widely documented in species that have teeth, claws or other armaments than in unarmed species (Guo & Dukas 2020). These social interactions can depend on various biotic and abiotic factors (Candolin & Wong 2012; Fisher *et al.* 2021). One of these biotic factors may be plant-animal associations, but their role in explaining the observed skew in social interactions is little explored.

Studies have shown that several specialised plant metabolites are taken up by certain animal species independently of nutrition, and stored (sequestered), for example, for use in mating or as own defence (Nishida 2014; Opitz & Müller 2009). This phenomenon is called pharmacophagy (Boppré 1984). Such uptake of non-nutritive plant metabolites can have direct effects on the behaviour of the individuals (Amano *et al.* 1999; Conner *et al.* 2000; Gonzalez *et al.* 1999). Remarkably, such plant metabolites, or further metabolised compounds, can also be acquired from other animals, either conspecifics (Paul *et al.* 2021; Singh *et al.* 2022) or heterospecifics (Tea *et al.* 2021). This can impact social interactions, which in turn can influence the fitness of individuals (Formica *et al.* 2012; Royle *et al.* 2012; Silk *et al.* 2003) and thus the individualised social niche (Kaiser et al. under revision).

Moreover, taking up plant metabolites may come with some costs, e.g. if the metabolites are

toxic (Kortbeek *et al.* 2019).

Interestingly, many species show intraspecific variation in the presence and quantity of

sequestered plant metabolites (Liu *et al.* 2023; Mattila *et al.* 2021; Opitz & Müller 2009;

Speed *et al.* 2012). Such variation can result from variation in the plant metabolite

concentration per se, potential costs related to metabolite acquisition by the animal or factors

such as age, sex or immunological status of the individuals (Dimarco & Fordyce 2017;

Massad *et al.* 2011; Moore *et al.* 2014; Smilanich *et al.* 2009; Zvereva & Kozlov 2015). This

leads us to the question of whether such intraspecific variation in sequestered plant

metabolites can shape social interactions in animals and whether the sequestered plant

metabolites affect fitness proxies such as lifespan of individuals, either directly or via altered

social interactions. In cases where plant metabolites are passed between individuals, we can

expect them to alter the social interactions of both the focal and the recipient individual.

However, despite the documented effect of sequestered plant metabolites on individual

behaviour (Boppré 1984; Nishida 2014), we still know relatively little about how these effects

scale up to impact social interactions.

To target these open questions, we examined the effect of intraspecific variation in

sequestered plant metabolites on social interactions and survivorship in the turnip sawfly,

Athalia rosae (Hymenoptera: Tenthredinidae). The adults of *A. rosae* feed on floral nectar,

usually from Apiaceae plants. In addition, they visit non-host plant species such as *Ajuga*

reptans (Lamiaceae), on which they exhibit pharmacophagy by taking up specialised

metabolites, neo-clerodane diterpenoids (hereafter called 'clerodanoids'), which they further

metabolize slightly (Brueggemann *et al.* 2023). These clerodanoids can serve as a resource for

A. rosae adults in multiple ways (Paul *et al.* 2022). Access to clerodanoids increases their

mating success (Amano *et al.* 1999) and also provides chemical defence against predators

(Nishida & Fukami 1990; Singh *et al.* 2022). Moreover, individuals can obtain clerodanoids

via agonistic interactions like nibbling on conspecifics that have had access to plant

clerodanoids (Paul & Müller 2021; Singh *et al.* 2022). Such agonistic interactions can

potentially be costly, as suggested by the transcriptional upregulation of metabolic pathways

in sawflies that nibbled on conspecifics (Paul *et al.* 2021). Moreover, the uptake of

clerodanoids may also directly pose fitness costs, such as a reduced lifespan (Zanchi *et al.*

2021). However, laboratory cultures of *A. rosae* are usually maintained without access to

clerodanoids, indicating that clerodanoids are not essential for individual survival.

In the present study, we first examined the clerodanoid contents of wild-caught sawflies to

investigate whether there is intraspecific variation in amounts of these metabolites in natural

A. rosae populations. Next, we tested the effect of no access, indirect access to clerodanoids

via conspecifics and direct access to clerodanoids via *A. reptans* leaves on behavioural

interactions in laboratory-reared individuals and analysed their clerodanoid contents. As we

observed intraspecific variation in clerodanoid content in both wild-caught and laboratory-

reared individuals, we, next, tested how such variation may impact social interactions.

Therefore, we set up a gradient of treatment groups with intraspecific variation in

clerodanoids, such that in a group either none, half or all individuals had access to

clerodanoids. We used network analysis to construct social interaction networks and calculate

individual and group measures of the network (Wey *et al.* 2008), allowing us to estimate the

effects of intraspecific variation in clerodanoids. Finally, we examined potential fitness costs

of clerodanoid acquisition and the resulting social interactions on lifespan, by setting up pairs

of individuals, in which either none, one or both individuals had prior access to clerodanoids.

We expected intraspecific variation in clerodanoids in wild-caught individuals. Furthermore,

we hypothesised that groups with no clerodanoid access should have the least social

interactions, while asymmetric clerodanoid acquisition should lead to increased social

interactions. We expected individuals without clerodanoid access to show differences in their

social interaction frequency depending on the clerodanoid acquisition status of other

conspecifics. Finally, we expected agonistic interactions to obtain clerodanoids from

conspecifics to be costly and to lead to a reduced lifespan.

Materials and Methods

Maintenance of study organism

Adults of *A. rosae* used in the experiments (other than wild-caught adults) were taken from a

laboratory stock population that had been established from sawflies collected in and around

Bielefeld, Germany and annually supplemented with field-caught individuals (see Singh *et*

al., 2023 for laboratory rearing details). The individuals were provided a 2% (v/v) honey-

water solution on paper that was refreshed every alternate day. The laboratory population was

maintained in multiple mesh cages (each 60 x 60 x 60 cm) at room temperature (15-25°C)

with a 16:8 h light:dark cycle and approximately 60% relative humidity.

Experiment 1. Chemical analysis for clerodanoid content of wild-caught sawflies

In total eighteen female and eight male adult *A. rosae* were collected in the wild at three

locations (SI.1). For analysis of clerodanoid amounts, the insects were individually frozen,

lyophilized, homogenized and extracted in ethyl acetate (LC-MS grade, VWR, Leuven,

Belgium), using ultrasonication. . After centrifugation, supernatants were dried and resolved

in 100 µl of methanol with mefenemic acid (Sigma-Aldrich GmbH, Taufkirchen, Germany)

as internal standard. Samples were filtered and analysed using an ultra-high performance

liquid chromatograph (UHPLC; Dionex UltiMate 3000, Thermo Fisher Scientific, San José,

CA, USA) equipped with a Kinetex XB-C18 column (1.7 µm, 150 × 2.1 mm, with guard

column, Phenomenex) kept at 45 °C, coupled to a quadrupole time of flight mass

spectrometer (QTOF-MS/MS; compact, Bruker Daltonics, Bremen, Germany) in negative

electrospray ionisation mode, following Brueggemann *et al.* (2023; for details see

Supplement).Samples were separated with a gradient from 0.1% formic acid (p.a., eluent

additive for LC-MS, ~98%, Sigma-Aldrich) in millipore water (eluent A) to 0.1% formic acid

in acetonitrile (LC-MS grade, Fisher Scientific, Loughborough, UK; eluent B) at a flow rate

of 0.5 ml min^{-1}, starting eluent B at 2%, increasing to 30% within 20 min, and then to 75%

within 9 min, followed by column cleaning and equilibration. The settings for the MS mode

were capillary voltage 3000 V, end plate offset 500 V, nebuliser (N_2) pressure 3 bar, dry gas

(N_2, 275 °C) flow 12 L min^{-1}, quadrupole ion energy 4 eV and collision energy 7 eV. Line

spectra were captured at 6 Hz, with a range of 50-1300 *m/z*. For recalibration of the *m/z* axis,

a Na(HCOO)-solution was injected prior to each sample. Blanks without animal material

were prepared in the same way and measured under the same conditions.

Mass axis recalibration and peak picking with the T-ReX 3D algorithm including spectral

background subtraction were performed in Metaboscape 2021b (Bruker Daltonics), with an

intensity threshold of 1000 counts and minimum peak length of 9 spectra. The allowed ion

types for bucket generation were: [M-H]$^-$, [M-H2O-H]$^-$, [M+Cl]$^-$, [M+HCOOH-H]$^-$,

[M+CH$_3$COOH-H]$^-$ and [2M-H]$^-$. From each bucket only the feature with the highest intensity

was used. We focussed on two metabolites which are likely taken up from *A. reptans* and

metabolised by the sawflies (Brueggemann *et al.* 2023), putative clerodanoid 1 eluting at 16.3

min (*m/z* 529.229 [M+HCOOH-H]$^-$) and putative clerodanoid 2 eluting at 18.2 min (*m/z*

527.214 [M+HCOOH-H]$^-$), with the monoisotopic masses of 484.231 Da (proposed sum

formula $C_{24}H_{36}O_{10}$) and 482.214 Da ($C_{24}H_{34}O_{10}$), respectively. Peak intensities of these target

metabolites were related to the peak intensity of mefenemic acid for semi-quantitative

analysis.

Experiment 2. Effect of direct and indirect clerodanoid acquisition on social interactions

We investigated the effects of no clerodanoid access (C-), indirect clerodanoid access from a conspecific (C+ from conspecific), or direct clerodanoid access from a plant (C+ from plant) by females on social interactions with C- males. Note that 'C+ from conspecific' has been previously referred to as AC+ (Paul *et al.* 2021; Singh *et al.* 2022). For the 'C+ from plant' treatment, freshly eclosed females were provided individually an *A. reptans* leaf disc (~1 cm^2) for 48 h, while 'C+ from conspecific' treatment females were kept with a C+ conspecific (2-4 days old female that had previously contact to *A. reptans*) for 48 h. Males (6- 10 days post-eclosion) were kept individually in Petri dishes for 48 h prior to the assays. The males were mated to a non-focal C- female (~3 days old) 24 h prior to the experiment to avoid any differences arising due to first mating (Torres-Vila & Jennions 2005). For the behavioural assays, one female was set up with one male in a clean Petri dish (5.5 cm diameter) and recorded (Sony HDR-CX410VE camcorder, AVCHD – 1920 × 1080 − 25 fps) for 25 minutes (16 replicates per treatment combination). One to two replicates of each treatment were setup simultaneously in a trial. An observer scored the number of occurrences of agonistic behaviour, including frontlimb battling, fighting and nibbling events (see Paul & Müller, 2021 for detailed behavioural descriptions), and mating behaviour (number of matings, latency until first mating) in each movie replicate using the software BORIS v7.13.9 (Friard & Gamba 2016), being blind to the treatment. After the behavioural assays, five females from each treatment were chosen randomly, frozen and analysed for their clerodanoid content as in experiment 1.

Experiment 3. Effect of intraspecific variation in clerodanoids on social networks

We found intraspecific variation in clerodanoid content in both wild-caught and lab-reared sawflies (see Results). To test the effect of clerodanoid acquisition status on social networks, we set up three group types of two males and two females each, in which either no individual

had prior access to clerodanoids (C- group), only one female and one male had prior access to

clerodanoids (Mixed group) or all individuals had prior access to clerodanoids from

A. reptans leaves (C+ group). We set up 15 replicates per group type. Two hours before the

behavioural assay, each individual was colour-marked (green, red, white, or yellow; Uni

POSCA marker pens, Japan) on its thorax and wings to enable identification during behaviour

scoring. The colours were alternated between sexes and treatments across replicates to avoid

any colour effects. In each trial, one to two replicates per group type were set up

simultaneously in Petri dishes and recorded for 25 minutes. Agonistic and mating interactions

of each individual in a movie replicate were scored by an observer as in experiment 2.

For the social network analysis, we analysed group as well as individual measures of the

network. Therefore, we constructed an undirected weighted network for the social interactions

(sum of agonistic and mating interactions) in each replicate. We calculated the network

weights as the total number of interactions between each pair of individuals (dyad). We

normalised the weights at the replicate level and at the dyadic level in each replicate, by

dividing by the maximum number of interactions across all replicates and all dyads,

respectively. At the group level, we calculated the density of social interactions in each

replicate, as following:

$$Density = \frac{sum\ of\ weights}{[number\ of\ vertices * (number\ of\ vertices - 1) * 0.5},$$

where vertices is the number of individuals in a replicate. Moreover, at the group level, we

counted the number of distinct dyads that exhibited social interactions in each replicate. At

the individual level, we calculated the strength (Barrat *et al.* 2004) and harmonic centrality

(Marchiori & Latora 2000). The strength is a measure of how strongly an individual is

directly connected to other individuals in the network. Harmonic centrality calculates the

'average' distance of an individual to the other individuals in the network. We used harmonic

centrality as we had disconnected networks in some replicates.

Experiment 4. Costs of direct and indirect clerodanoid acquisition on lifespan

To examine potential fitness costs of direct and indirect clerodanoid acquisition, we measured

the lifespan of differently treated females. We collected freshly eclosed *A. rosae* females and

assigned them to either C- or C+, offering the latter *A. reptans* leaves for 48 hours. Next, we

put together either two C- individuals (C-C- pair), a C- and a C+ individual (C-C+ pair) or

two C+ individuals (C+C+ pair). The C- individual from the C-C+ pair could acquire

clerodanoids from the C+ conspecific. Individuals in a replicate were marked with green or

white colour paint to allow for differentiation. After 48 hours, we randomly chose one

individual from each replicate of the C-C- pair and the C+C+ pair, and both individuals from

the C-C+ pair. Each individual was placed in a Petri dish and checked daily for survivorship.

Statistical analyses

All analyses were done in R version 4.2.1 (R Core Team 2022). To examine the effect of

female treatment on the number of agonistic and mating interaction occurrences in experiment

2, we used a poisson-distributed generalised linear mixed model (GLMM) with trial as

random effect (package 'lme4' version 1.1-30, package 'lmerTest' version 3.1-3; Bates *et al.*

2015; Kuznetsova *et al.* 2017). We examined the effect of female treatment on mating latency

using a linear mixed model (LMM) with trial as random effect. We tested for differences

between C-, 'C+ from conspecific' and 'C+ from plant' in clerodanoid compounds 1 and 2

using a Kruskal-Wallis test followed by Dunn test with Holm correction for post-hoc tests

('FSA' package version 0.9.4, Ogle *et al.* 2023). In experiment 3, we ran LMM for the

density and binomial GLMM for number of dyads with social interactions, with group type as

fixed effect and trial as random effect. Next, we analysed whether the group type had an

effect on the individual-level network measures of strength and harmonic centrality

(calculated using 'igraph' package version 1.5.1, Csárdi *et al.* 2023) for C- and C+ sawflies

separately. We performed poisson-distributed GLMM for strength and LMM for harmonic

centrality. The models included group type, sex and their interaction as fixed effects and

replicate ID and trial as random effect. We conducted post hoc analyses with Holm correction

using the package 'multcomp' version 1.4-20 (Hothorn *et al.* 2008) for LMM and GLMM. If

the interaction terms were non-significant, they were dropped to test the significance of the

lower-order terms. The distribution of model residuals was checked. In experiment 4, to

analyse differences in lifespan of C- (from C-C- pair), C- and C+ individual (from C-C+ pair)

and C+ (from C+C+ pair), we produced Kaplan-Meier plots and compared the curves by the

log-rank test using 'survival' package version 3.5-0 (Therneau 2023) and 'survminer' package

version 0.4.9 (Kassambara *et al.* 2021). Similarly, we tested for differences in lifespan

between C+ individuals from C+C+ pairs and C+ individuals from C-C+ pairs.

Results

Clerodanoid amounts vary in wild-caught female and male sawflies

254   Both females and males showed intraspecific variation in the amount of putative clerodanoid

255   1 and clerodanoid 2 across the three locations from which they were collected (Figure 1).

256   Some individuals had no clerodanoids, while others had relatively high quantities, suggesting

257   that natural populations show variation in clerodanoid acquisition.

Clerodanoids from plants impact behaviour and clerodanoid amounts varies in lab-reared

sawflies

There was a significant effect of female treatment on number of agonistic interaction

occurrences (χ_2^2 = 48.56, P < 0.001, Figure 2a), but not on mating occurrences (χ_2^2 = 1.76, P

= 0.41, Figure 2b). C- and 'C+ from conspecific' treatment females had significantly fewer

agonistic interactions than 'C+ from plant' treatment females (SI.2a). There was a significant

effect of female treatment on mating latency (χ_2^2 = 9.90, P = 0.007, Figure 2c) with C- females

having a significantly longer latency until mating than 'C+ from plant' females, while 'C+

from conspecific' females showed an intermediate mating latency (SI.2b).

There was a significant difference between C-, 'C+ from conspecific' and 'C+ from plant'

females in the amount of clerodanoid 1 (χ_2^2 = 8.11, P = 0.017, Figure 2d) and clerodanoid 2

(χ_2^2 = 8.10, P = 0.017, Figure 2e). No clerodanoids were detected in C- females, while there

was inter-individual variation in clerodanoids in both 'C+ from conspecific' and 'C+ from

plant' females, which did not differ significantly from each other (SI.2c, SI.2d).

Intraspecific variation in clerodanoids modulates social networks

The group types varied in their social networks, with the C- group having fewer interactions

than the Mixed and the C+ group (Figure 3a). Our results showed that there was a significant

effect of group type on density (χ_2^2 = 27.98, P < 0.001, Figure 3b) and number of dyads with

social interactions (χ_2^2 = 41.29, P < 0.001, Figure 3c). Mixed groups had a significantly

higher density and more dyads that had interacted than C- and C+ groups (SI.3a, SI.3b).

Together this suggests that access to clerodanoids increases social interactions with more

individuals networking, particularly in the Mixed group.

For C- sawflies, there was a significant effect of group type (χ_1^2 = 63.55, P < 0.001) on

strength (Figure 4a) such that C- sawflies in Mixed groups had a higher strength, i.e., had

more social interactions, than in C- groups (SI.4a). In contrast, there was a significant effect

of sex ($\chi_1^2 = 25.14$, $P < 0.001$) on strength of interactions (Figure 4b) for C+ sawflies, with

males having a lower strength than females across Mixed and C+ groups (SI.4a). For

harmonic centrality, there was a significant effect of group type ($\chi_1^2 = 6.88$, $P = 0.008$) in C-

sawflies (Figure 4c), such that those in Mixed groups had higher harmonic centrality values.

There was no significant effect of any predictor variable on harmonic centrality in C+

sawflies (Figure 4d, SI.4c).

C+ individuals from C-C+ pairs have a shorter lifespan

There was no significant difference between the lifespan of C- females from C-C- pairs, C-

females from C-C+ pairs and C+ females from C+C+ pairs ($\chi_2^2 = 2.3$, $P = 0.3$, Figure 5).

However, C+ adults from C-C+ pairs had a significantly shorter lifespan than C+ adults from

C+C+ pairs ($\chi_1^2 = 5.2$, $P = 0.02$), indicating that agonistic interactions to obtain clerodanoids

may have fitness costs.

Discussion

We observed variation in clerodanoid amounts in wild-caught *A. rosae* adults of both sexes

and from all locations. A capture release study had shown that most of the *A. rosae* adults

could be recaptured from within 100 m of their release, suggesting limited dispersal

(Nagasaka 1992). Thus, our field populations were likely separated, indicating that

intraspecific variation in sequestered compounds may be common in natural populations of

sequestering herbivorous insects. Moreover, we observed intraspecific variation in

clerodanoid amounts of lab-reared sawflies with both indirect (via nibbling on conspecifics)

and direct (via plant leaves) access to clerodanoids, similar to previous studies (Singh *et al.*

2022). Such intraspecific variation in sequestered plant chemicals in animals is widely

observed (Opitz & Müller 2009; Speed *et al.* 2012). While this variation may result from environmental stochasticity, e.g., variation in metabolite concentrations across plants, it could also have an evolutionarily basis, e.g., some individuals of an animal species may be chemically defended while others are not (Brower *et al.* 1970; Speed *et al.* 2012). This can be seen as automimicry in which undefended individuals (mimics) profit from the unpalatability of defended individuals (models). Studies have shown that clerodanoids can serve as a chemical defence against predators, like mantids (Singh *et al.* 2022) and tree sparrows (Nishida & Fukami 1990), indicating that intraspecific variation in chemical defence within wild populations can lead to automimicry.

The presence and quantity of clerodanoids can be considered as important parts of the chemical phenotype of individuals, which could play a key role in shaping the individualised niche, as has been argued for plants (Müller & Junker 2022). While effects of host plants on an individual's phenotype and behaviour are widely documented (Geiselhardt *et al.* 2012; Jarrett & Miller 2024; Müller & Müller 2017; Singh *et al.* 2020), we demonstrate here that plant chemicals sequestered from non-host plants can also modulate social interactions. Our results demonstrated the highest density and number of interacting dyads in social networks of Mixed groups, followed by C+ groups. The asymmetric access to clerodanoids in Mixed groups may lead to an increase in social interactions, potentially due to C- trying to obtain clerodanoids from C+ individuals. Moreover, C- individuals in Mixed groups could acquire clerodanoids from C+ individuals, which may then increase their social interactions, as it will change their clerodanoid acquisition status. Individuals in C- groups had the least density and number of interacting dyads, suggesting that when all individuals lack clerodanoids, they have only few intra-and inter-sex social interactions.

Remarkably, the social interactions of the individuals depended not only on their own clerodanoid acquisition status but also on that of other conspecifics in the group. For example,

C- individuals had a higher number of social interactions in Mixed than in C- groups (see

Figure 4a,4c), suggesting that even for the same clerodanoid-holding status, an individual

may have low or high frequencies of social interactions, depending on the group composition.

Similar effects of conspecifics on an individual's behaviour have been shown in the aquatic

insect, *Notonecta irrorata*, where an individual's propensity to disperse depended on the

behavioural type of conspecifics present (Kitchen & Chalcraft 2020). Given the intraspecific

variation in clerodanoid content that we observed in wild-caught individuals of *A. rosae*, we

expect social interactions in natural populations to range from low, for example, if no *A.*

reptans plants are around from which clerodanoids can be sequestered, to high, when some

insects could sequester clerodanoids, leading to a skewed distribution of social interaction

frequency.

Social interactions may accrue over time, and influence fitness of individuals through

different ways, such as injury/mortality from agonistic interactions, frequency of mating and

disease progression or spread (Alwash & Levine 2019; Dawson *et al.* 2018; Guo & Dukas

2020; Stroeymeyt *et al.* 2018). We observed a reduced lifespan in C+ females from C-C+

pairs compared to C+ females from C+C+ pairs (experiment 4). In experiment 2, we saw that

C+ females paired with C- males had the most agonistic interactions. Similarly, in Mixed

groups, we found more frequent interactions than in C- or C+ groups. Thus, we can expect

that C+ females in C-C+ pairs had more social, especially agonistic, interactions with their C-

conspecific, which may have nibbled on the C- individual to get access to clerodanoids. These

increased agonistic interactions may have led to a decreased lifespan for C+ females in C-C+

pairs. Such costs of aggression have also been shown in the fruit fly, *Drosophila*

melanogaster, where flies that were engaged in aggressive behaviour had a reduced survival

despite not showing any differences in wing damage from no-aggression treatment flies (Guo

& Dukas 2020). The authors hypothesised that such costs could arise from physiological and

metabolic changes triggered by stress and aggressive conflicts. Indeed, a previous study has shown transcriptional upregulation of genes involved in metabolic pathways in *A. rosae* sawflies that nibbled on conspecifics (Paul *et al.* 2021), which may indicate potential costs of the aggressive interactions.

Along with chemical defence against predation and increased mating success, sequestered plant metabolites could also lead to benefits for the offspring (Eisner *et al.* 2000; Eisner & Meinwald 1995; Gonzalez *et al.* 1999). However, at least under controlled laboratory conditions, no significant effects of parental clerodanoid status on offspring was found in *A. rosae* (Paul *et al.* 2022). Thus, while clerodanoid acquisition may be associated with benefits like defence against predation (Singh *et al.* 2022) and mating success (Amano *et al.* 1999; Paul & Müller 2021), there may be trade-off with costs in terms of lifespan due to agonistic interactions. This in turn may maintain intraspecific variation in clerodanoids in natural populations of *A. rosae*. Understanding the effects of variation in these sequestered plant metabolites is essential because these ecological effects can scale up from the individual level to factors that shape animal social interaction networks.

References

Alwash, N. & Levine, J.D. (2019). Network analyses reveal structure in insect social groups. *Current Opinion in Insect Science*, 35, 54–59.

Amano, T., Nishida, R., Kuwahara, Y. & Fukami, H. (1999). Pharmacophagous acquisition of clerodendrins by the turnip sawfly (*Athalia rosae* ruficornis) and their role in the mating behavior. *Chemoecology*, 9, 145–150.

Archer, J. (1988). *The behavioural biology of aggression*. Cup Archive.

Barrat, A., Barthélemy, M., Pastor-Satorras, R. & Vespignani, A. (2004). The architecture of complex weighted networks. *Proc. Natl. Acad. Sci. U.S.A.*, 101, 3747–3752.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting linear mixed-effects models Using lme4. *Journal of Statistical Software*, 67, 1–48.

Boppré, M. (1984). Redefining "pharmacophagy." *Journal of Chemical Ecology*, 10, 1151–1154.

Brower, L.P., Pough, F.H. & Meck, H. (1970). Theoretical investigations of automimicry, I. Single trial learning. *Proceedings of the National Academy of Sciences*, 66, 1059–1066.

Brueggemann, L., Tewes, L.J. & Müller, C. (2023). Characterisation and localisation of plant metabolites involved in pharmacophagy in the turnip sawfly. *PLoS ONE*, 18, e0291180.

Candolin, U. & Wong, B.B. (2012). *Behavioural responses to a changing world: mechanisms and*
consequences. Oxford University Press.
Conner, W.E., Boada, R., Schroeder, F.C., Gonzalez, A., Meinwald, J. & Eisner, T. (2000). Chemical
defense: Bestowal of a nuptial alkaloidal garment by a male moth on its mate. *Proceedings of*
the National Academy of Sciences, 97, 14406–14411.
Csárdi, G., Nepusz, T., Traag, V., Horvát, S., Zanini, F., Noom, D., *et al.* (2023). *igraph: Network*
Analysis and Visualization in R.
Dawson, E.H., Bailly, T.P.M., Dos Santos, J., Moreno, C., Devilliers, M., Maroni, B., *et al.* (2018). Social
environment mediates cancer progression in *Drosophila*. *Nat Commun*, 9, 3574.
Dimarco, R.D. & Fordyce, J.A. (2017). Not all toxic butterflies are toxic: high intra- and interspecific
variation in sequestration in subtropical swallowtails. *Ecosphere*, 8.
Eisner, T., Eisner, M., Rossini, C., Iyengar, V.K., Roach, B.L., Benedikt, E., *et al.* (2000). Chemical
defense against predation in an insect egg. *Proc. Natl. Acad. Sci. U.S.A.*, 97, 1634–1639.
Eisner, T. & Meinwald, J. (1995). The chemistry of sexual selection. *Proceedings of the National*
Academy of Sciences, 92, 50–55.
Fisher, D.N., Kilgour, R.J., Siracusa, E.R., Foote, J.R., Hobson, E.A., Montiglio, P., *et al.* (2021).
Anticipated effects of abiotic environmental change on intraspecific social interactions.
Biological Reviews, 96, 2661–2693.
Fisher, D.N., Rodríguez-Muñoz, R. & Tregenza, T. (2019). Dynamic networks of fighting and mating in
a wild cricket population. *Animal Behaviour*, 155, 179–188.
Formica, V.A., Wood, C.W., Larsen, W.B., Butterfield, R.E., Augat, M.E., Hougen, H.Y., *et al.* (2012).
Fitness consequences of social network position in a wild population of forked fungus beetles
(*Bolitotherus cornutus*). *J of Evolutionary Biology*, 25, 130–137.
Frank, S.A. (2007). All of life is social. *Current Biology*, 17, R648–R650.
Friard, O. & Gamba, M. (2016). BORIS: a free, versatile open-source event-logging software for
video/audio coding and live observations. *Methods in ecology and evolution*, 7, 1325–1330.
Geiselhardt, S., Otte, T. & Hilker, M. (2012). Looking for a similar partner: host plants shape mating
preferences of herbivorous insects by altering their contact pheromones. *Ecol Lett*, 15, 971–
977.
Gonzalez, A., Rossini, C., Eisner, M. & Eisner, T. (1999). Sexually transmitted chemical defense in a
moth (*Utetheisa ornatrix*). *Proceedings of the National Academy of Sciences*, 96, 5570–5574.
Guo, X. & Dukas, R. (2020). The cost of aggression in an animal without weapons. *Ethology*, 126, 24–
31.
Hothorn, T., Bretz, F. & Westfall, P. (2008). Simultaneous Inference in General Parametric Models.
Biometrical Journal, 50, 346–363.
Jarrett, B.J.M. & Miller, C.W. (2024). Host plant effects on sexual selection dynamics in phytophagous
insects. *Annu. Rev. Entomol.*, 69, annurev-ento-022823-020258.
Kassambara, A., Kosinski, M. & Biecek, P. (2021). *survminer: Drawing Survival Curves using "ggplot2."*
Kitchen, A.J. & Chalcraft, D.R. (2020). An individual's propensity to disperse is dependent on the
behavioral type of its peers but not its own behavioral type. *Oecologia*, 194, 403–413.
Kortbeek, R.W.J., Van Der Gragt, M. & Bleeker, P.M. (2019). Endogenous plant metabolites against
insects. *Eur J Plant Pathol*, 154, 67–90.
Koto, A., Tamura, M., Wong, P.S., Aburatani, S., Privman, E., Stoffel, C., *et al.* (2023). Social isolation
shortens lifespan through oxidative stress in ants. *Nat Commun*, 14, 5493.
Krause, J., James, R., Franks, D.W. & Croft, D.P. (2015). *Animal social networks*. Oxford University
Press, USA.
Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2017). **lmerTest** Package: Tests in Linear Mixed
Effects Models. *J. Stat. Soft.*, 82, 1–26.
Liu, Y., Heath, J.J., Zhang, S., Van Wijk, M., Wang, G., Buellesbach, J., *et al.* (2023). A mosaic of
endogenous and plant-derived courtship signals in moths. *Current Biology*, 33, 3529-3535.e4.
Marchiori, M. & Latora, V. (2000). Harmony in the small-world. *Physica A: Statistical Mechanics and*
its Applications, 285, 539–546.

Massad, T.J., Fincher, R.M., Smilanich, A.M. & Dyer, L. (2011). A quantitative evaluation of major
plant defense hypotheses, nature versus nurture, and chemistry versus ants. *Arthropod-Plant*
Interactions, 5, 125–139.

Mattila, A.L.K., Jiggins, C.D., Opedal, Ø.H., Montejo-Kovacevich, G., Pinheiro de castro, É.C., McMillan,
W.O., *et al.* (2021). Evolutionary and ecological processes influencing chemical defense
variation in an aposematic and mimetic *Heliconius* butterfly. *PeerJ*, 9, e11523.

Moore, B.D., Andrew, R.L., Külheim, C. & Foley, W.J. (2014). Explaining intraspecific diversity in plant
secondary metabolites in an ecological context. *New Phytologist*, 201, 733–750.

Mueller, T. & Mueller, C. (2017). Host plant effects on the behavioural phenotype of a Chrysomelid.
Ecological Entomology, 42, 336–344.

Müller, C. & Junker, R.R. (2022). Chemical phenotype as important and dynamic niche dimension of
plants. *New Phytologist*, 234, 1168–1174.

Nagasaka, K. (1992). Movement patterns of three *Athalia* sawflies in relation to the spatio-temporal
distributions of their habitats. *Population Ecology*, 34, 1–14.

Nishida, R. (2014). Chemical ecology of insect–plant interactions: ecological significance of plant
secondary metabolites. *Bioscience, Biotechnology, and Biochemistry*, 78, 1–13.

Nishida, R. & Fukami, H. (1990). Sequestration of distasteful compounds by some pharmacophagous
insects. *J Chem Ecol*, 16, 151–164.

Ogle, D.H., Doll, J.C., Wheeler, A.P. & Dinno, A. (2023). *FSA: Simple Fisheries Stock Assessment*
Methods.

Opitz, S.E.W. & Müller, C. (2009). Plant chemistry and insect sequestration. *Chemoecology*, 19, 117–
154.

Paul, S.C., Dennis, A.B., Tewes, L.J., Friedrichs, J. & Müller, C. (2021). *Consequences of*
pharmacophagous uptake from plants and conspecifics in a sawfly elucidated using chemical
and molecular techniques (preprint). Zoology. BioRxiv. doi: 10.1101/2021.02.09.430406.

Paul, S.C. & Müller, C. (2021). Fighting over defense chemicals disrupts mating behavior. *Behavioral*
Ecology, arab117.

Paul, S.C., Singh, P., Dennis, A.B. & Müller, C. (2022). Intergenerational effects of early life starvation
on life-history, consumption, and transcriptome of a holometabolous insect. *The American*
Naturalist, 719397.

R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. R Foundation for
Statistical Computing, Vienna, Austria.

Royle, N.J., Pike, T.W., Heeb, P., Richner, H. & Kölliker, M. (2012). Offspring social network structure
predicts fitness in families. *Proc. R. Soc. B.*, 279, 4914–4922.

Ruan, H. & Wu, C.-F. (2008). Social interaction-mediated lifespan extension of *Drosophila* Cu/Zn
superoxide dismutase mutants. *Proc. Natl. Acad. Sci. U.S.A.*, 105, 7506–7510.

Silk, J.B., Alberts, S.C. & Altmann, J. (2003). Social bonds of female baboons enhance infant survival.
Science, 302, 1231–1234.

Singh, P., van Bergen, E., Brattström, O., Osbaldeston, D., Brakefield, P.M. & Oostra, V. (2020).
Complex multi-trait responses to multivariate environmental cues in a seasonal butterfly.
Evol Ecol, 34, 713–734.

Singh, P., Grone, N., Tewes, L.J. & Müller, C. (2022). Chemical defense acquired via pharmacophagy
can lead to protection from predation for conspecifics in a sawfly. *Proc. R. Soc. B.*, 289,
20220176.

Singh, P., Wolthaus, J., Schielzeth, H. & Müller, C. (2023). State dependency of behavioural traits is a
function of the life stage in a holometabolous insect. *Animal Behaviour*, 203, 29–39.

Smilanich, A.M., Dyer, L.A., Chambers, J.Q. & Bowers, M.D. (2009). Immunological cost of chemical
defence and the evolution of herbivore diet breadth. *Ecology Letters*, 12, 612–621.

Snyder-Mackler, N., Burger, J.R., Gaydosh, L., Belsky, D.W., Noppert, G.A., Campos, F.A., *et al.* (2020).
Social determinants of health and survival in humans and other animals. *Science*, 368,
eaax9553.

Speed, M.P., Ruxton, G.D., Mappes, J. & Sherratt, T.N. (2012). Why are defensive toxins so variable?
An evolutionary perspective. *Biological Reviews*, 87, 874–884.

Stroeymeyt, N., Grasse, A.V., Crespi, A., Mersch, D.P., Cremer, S. & Keller, L. (2018). Social network
plasticity decreases disease transmission in a eusocial insect. *Science*, 362, 941–945.

Tea, Y., Soong, J.W., Beaver, E.P. & Lohman, D.J. (2021). Kleptopharmacophagy: Milkweed butterflies
scratch and imbibe from Apocynaceae-feeding caterpillars. *Ecology*.

Therneau, T.M. (2023). *A Package for Survival Analysis in R*.

Torres-Vila, L.M. & Jennions, M.D. (2005). Male mating history and female fecundity in the
Lepidoptera: do male virgins make better partners? *Behavioral Ecology and Sociobiology*, 57,
318–326.

Wey, T., Blumstein, D.T., Shen, W. & Jordán, F. (2008). Social network analysis of animal behaviour: a
promising tool for the study of sociality. *Animal Behaviour*, 75, 333–344.

Zanchi, C., Lo, L.K., R, R., Moritz, I., Kurtz, J. & Müller, C. (2021). Survival of the sawfly *Athalia rosae*
upon infection by an entomopathogenic fungus and in relation to clerodanoid uptake. *Front.*
Physiol., 12, 637617.

Zvereva, E.L. & Kozlov, M.V. (2015). The costs and effectiveness of chemical defenses in herbivorous
insects: a meta-analysis. *Ecological Monographs*, 15-0911.1.

Figures

Figure 1. Intraspecific variation in peak intensities of putative (a) clerodanoid 1 and (b)

clerodanoid 2 per individual in wild-caught *A. rosae* female (n=18) and male (n=8) sawflies

from three locations.

Figure 2. Effect of female treatment, no (C-), indirect (C+ from conspecific: prior contact to

other female that had access to *Ajuga reptans* leaf) or direct (C+ from plant: prior access to

A. reptans leaf) access to clerodanoids, on number of occurrences of (a) agonistic and (b)

mating interactions, and on (c) mating latency with C- males of *Athalia rosae*. Effect of

female treatment on peak intensity of putative (d) clerodanoid 1 and (e) clerodanoid 2 in lab-

reared female sawflies. The boxplots represent the median and 25th and 75th percentiles; the

whiskers indicate the values within 1.5 times the interquartile range. The coloured circles

represent individual data-points. The given *P*-values are the outcome of a GLMM (a,b), LMM

(c) or Kruskal-Wallis (d,e) test. Different lower-case letters in the plots represent significant

differences from post-hoc tests (*P* < 0.05).

Figure 3. (a) Networks representing the C-, Mixed and C+ groups (n=15 per group type).

Each cell in the grid corresponds to one replicate and consisted of two female (circular nodes)

and two male (square nodes) adults of *Athalia rosae*. The sawflies had either no access (C-,

red nodes) or prior access to *Ajuga reptans* (C+, blue nodes). Social interactions between a

dyad are represented by lines between the individuals and the frequency of their social

interactions is represented by the thickness of the line. The lighter colour-shade of nodes

represents those individuals that had no social interactions during the observation period.

Effect of group type on (b) density of social interactions, and (c) number of dyads with social

interactions. The boxplots represent the median and 25th and 75th percentiles; the whiskers

537   indicate the values within 1.5 times the interquartile range. The coloured circles represent

538   individual data-points. The given P-values are the outcome of a LMM (b) or GLMM (c)

539   model. Different lower-case letters in the plots represent significant differences from post-hoc

540   tests ($P < 0.05$).

Figure 4. Effect of group type (C-, Mixed or C+ groups: represented by different colours) on strength (top row) and harmonic centrality (bottom row) in C- treatment sawflies (a,c) and C+ treatment sawflies (b,d) for both sexes. The boxplots represent the median and 25th and 75th percentiles; the whiskers indicate the values within 1.5 times the interquartile range. The coloured circles represent individual data-points. The given *P*-values are the outcome of a GLMM (a,b) or LMM (c,d) model. Different lower-case letters in the plots represent significant differences from post-hoc tests (*P* < 0.05).

Figure 5. Survival of *Athalia rosae* females kept for 48 h with an individual of the same or a

different treatment (C-: no access, C+: prior access to *Ajuga reptans* leaf), represented by

Kaplan-Meier curves. The black dashed line represents the median survival of each treatment

given by the point where it intersects the treatment curve. Note that the C- individuals from

C-C+ pairs and C+ individuals from C-C+ pairs originate from the same treatment (see text

for details).