Xylosyltransferase Bump-and-hole Engineering to Chemically Manipulate

Proteoglycans in Mammalian Cells

Zhen Li[a,b], Lucia Di Vagno[b,c,#], Himanshi Chawla[a,b,#], Aisling Ni Cheallaigh[d], Meg Critcher[e,f],

Douglas Sammon[g,%], David C. Briggs[h], Nara Chung[i], Vincent Chang[i], Keira E. Mahoney[i],

Anna Cioce[a,b], Lloyd D. Murphy[j], Yen-Hsi Chen[k,&], Yoshiki Narimatsu[k], Rebecca L. Miller[k],

Lianne I. Willems[j], Stacy A. Malaker[i], Mia L. Huang[e,f], Gavin J. Miller[d], Erhard Hohenester[g],

Benjamin Schumann[a,b,*]

[a] Department of Chemistry, Imperial College London, W12 0BZ, London, United Kingdom.

[b] Chemical Glycobiology Laboratory, The Francis Crick Institute, NW1 1AT London, United

Kingdom

[c] Proteomics Science Technology Platform, The Francis Crick Institute, London NW1 1AT,

United Kingdom.

[d] Lennard-Jones Laboratory, School of Chemical & Physical Sciences, and Centre for

Glycoscience, Keele University, Keele, Staffordshire, ST5 5BG United Kingdom.

[e] Skaggs Graduate School of Chemical and Biological Sciences, Scripps Research, La Jolla,

California.

[f] Department of Chemistry, Scripps Research, La Jolla, California.

[g] Department of Life Sciences, Imperial College London, London SW7 2AZ, UK.

[h] Signalling and Structural Biology Laboratory, The Francis Crick Institute, London NW1 1AT,

United Kingdom.

[i] Ddepartment of Chemistry, Yale University, CT 06511, New Haven, United States.

[j] York Structural Biology Laboratory and York Biomedical Research Institute, Department of

Chemistry, University of York, York YO10 5DD, United Kingdom

[k] Copenhagen Center for Glycomics, Department of Cellular and Molecular Medicine,

Faculty of Health Sciences, University of Copenhagen; Copenhagen-N, Denmark.

[%] Current address: Avacta, London, United Kingdom.

[&] Current address: GlycoDisplay ApS, Copenhagen, Denmark.

[#] These authors contributed equally.

* Correspondence: b.schumann@imperial.ac.uk

Abstract

Mammalian cells orchestrate signalling through interaction events on their surfaces.

Proteoglycans are an intricate part of these interactions, carrying large glycosaminoglycan

polysaccharides that recruit signalling molecules. Despite their importance in development,

cancer and neurobiology, a relatively small number of proteoglycans have been identified. In

addition to the complexity of glycan extension, biosynthetic redundancy in the first protein

glycosylation step by two xylosyltransferase isoenzymes XT1 and XT2 complicates

annotation of proteoglycans. Here, we develop a chemical genetic strategy that manipulates

the glycan attachment site of cellular proteoglycans. By employing a tactic termed bump-

and-hole engineering, we engineer the two isoenzymes XT1 and XT2 to specifically transfer

a chemically modified xylose analogue to target proteins. The chemical modification contains

a bioorthogonal tag, allowing the ability to visualise and profile target proteins modified by

both transferases in mammalian cells. The versatility of our approach allows pinpointing

glycosylation sites by tandem mass spectrometry, and exploiting the chemical handle to

manufacture proteoglycans with defined GAG chains for cellular applications. Engineered XT

enzymes permit a view into proteoglycan biology that is orthogonal to conventional

techniques in biochemistry.

Main

Proteoglycans are large biomolecules that consist of a core protein and one or more

glycosaminoglycan (GAG) modifications. Ubiquitous on cell surfaces and within the

extracellular matrix in metazoa, proteoglycans form a myriad of interactions with cellular

receptors as well as soluble signalling molecules, and provide structural support in

connective tissues such as cartilage.[1,2] Growth factors, neurotrophic factors and chemokines

can be recruited to target cells through GAG binding sites, rendering proteoglycans

important determinants for development.[3,4] Consequently, dysfunctions in GAG biosynthesis

cause severe phenotypes from embryonic lethality to skeletal and muscular deficiencies.[5]

Binding events between proteoglycans and their receptors are impacted by the core protein

as well as the identity of GAG polysaccharides that are classified into either heparan sulfate

(HS), chondroitin sulfate (CS), dermatan sulfate (DS) or keratan sulfate.[6,7] Biochemistry and

genetic engineering have linked proteoglycan physiology to the GAG structures on particular

cell types or even on distinct subcellular locations.[8–13] Despite their relevance in physiology,

only a relatively small number of less than 50 proteoglycans is known in humans.[14,15]

Furthermore, it is challenging to dissect the role of the proteoglycan backbone in physiology

from the role of the GAG chain, necessitating strategies in chemistry to manipulate and alter

proteoglycans.[16]

An impediment for profiling proteoglycans is the large size of GAG modifications that renders

analysis by mass spectrometry (MS) challenging. While GAG-carrying glycopeptides contain

common amino acid signatures such as acidic patches and a central O-glycosylated Ser with

often flanking Gly or Ala residues, there is no consensus sequence to predict GAG

glycosylation in the Golgi.[6,9,17,18] Common strategies to identify proteoglycans feature

enzymatic digestion of GAG chains either before or after isolation of glycopeptides.[9,19–24]

While powerful, such procedures make use of complex digestion and purification protocols

and focus solely on the GAG-carrying glycopeptide, without the advantages of shotgun

(glyco-)proteomics methods that employ the full MS peptide coverage of individual proteins

for detection.

Fig. 1: Design of a xylosyltransferase bump-and-hole system. **a**, principle of the BH approach. WT-

XTs transfer Xyl to substrate proteins that can be extended to GAG chains. BH-XTs transfer a

bioorthogonal Xyl analogue for visualisation and MS profiling. **b**, structural considerations of XT1

engineering to accept UDP-6AzGlc instead of UDP-Xyl. Insert: Gatekeeper residues in the XT1

crystal structure (PDB 6EJ7) and structural trajectory of 6-azidomethyl modification in UDP-6AzGlc. **c**,

in vitro glycosylation of a fluorescently labelled bikunin substrate peptide by WT- or mutant XT1 and

different UDP-sugars. Data are means ± SD from three technical replicates from one out of two

independent experiments. **d**, Michaelis-Menten kinetics of *in vitro* peptide glycosylation by different

XT1/UDP-sugar combinations. Data are one independent replicate with three technical replicates

(top) and average ± range from two independent replicates each (bottom).

The biosynthesis of HS and CS commences via a common O-linked glycan "linker"

modification consisting of a glucuronic acid (GlcA), two galactoses (Gal) and a xylose (Xyl) in

the GlcA(β-3)Gal(β-3)Gal(β-4)Xyl(β-)Ser sequence (Fig. 1a), with optional modifications

such as phosphorylation on the core Xyl.[7,25] The first glycosylation step attaching Xyl to Ser

is subject to biosynthetic redundancy by the xylosyltransferase isoenzymes XT1 and XT2

that utilize uridine diphosphate (UDP)-Xyl as a substrate. The isoenzymes share 60% amino

acid identity but display tissue-specific expression patterns and dysfunctions are associated

with different genetic disorders - Desbuquois Dysplasia type 2 and Spondylo-Ocular

Syndrome for patient *XYLT1* and *XYLT2* mutations, respectively.[26–29] Differential roles in

physiology have been attributed to XT1 and XT2.[30,31] Although XT2 appears to be the

dominant isoenzyme in cell lines and serum,[32,33] *Xylt1* and *Xylt2* knockout (KO) mice display

differential defects in development.[30,31] Despite their importance in physiology and the

unanswered questions about substrate repertoires, it is currently not possible to directly

profile the substrate proteins or even individual glycosylation sites of XT isoenzymes in cells

or *in vivo*.

Here, we employ a chemical biology tactic termed bump-and-hole (BH) engineering to probe

the substrates of human xylosyltransferases in living cells. Based on structural

considerations, we mutate a bulky amino acid in the active site of XT1 to a smaller residue to

accept the chemically modified substrate UDP-6AzGlc that is not accepted by the wildtype

(WT) enzyme. The chemical modification contains an azide group for bioorthogonal

incorporation of fluorophores or biotin (Fig. 1a). Judicious choice of the analogue as a

derivative of UDP-glucose enables cytosolic delivery, circumventing the lack of a cellular

salvage pathway for "direct" analogues of Xyl. After in-depth biochemical characterisation,

we install the XT1 BH system in mammalian cells to directly visualise and probe

proteoglycans. Using MS glycoproteomics, we find that BH-engineered XT1 modifies the

native glycosylation site of a model proteoglycan in mammalian cells. We further show that

BH engineering can be applied to the isoenzyme XT2 simply by harnessing sequence

conservation in protein engineering, allowing differential substrate profiling of both

isoenzymes in mammalian cells. Introduction of a chemical handle at the native glycosylation

site enables attachment of a bioorthogonally tagged GAG chain, furnishing "designer"

proteoglycans to modulate cellular behaviour. By developing an XT BH system, we

chemically manipulate proteoglycans in mammalian cells for applications in MS and

functional evaluation.

Results

Design of a xylosyltransferase bump-and-hole system

Our XT BH design was prompted by biosynthetic and structural considerations (Fig. 1b). In

the absence of a functional group that allows facile chemical modification,[34,35] the most

common approach to develop bioorthogonal reporters of monosaccharides is the replacement of hydroxyl groups with azido groups.[36–38] Beahm et al. developed a 4-azido-substituted Xyl analogue that is incorporated into proteoglycans by XT1,[39] but the corresponding UDP-sugar cannot be biosynthesized in mammalian cells since there is no salvage pathway for Xyl.[40] The bump-and-hole tactic uses substrate analogues that would normally not be accepted by GTs.[41–43] We sought to employ this feature to our benefit and reprogram XTs to accept a UDP-sugar that is not accepted by WT-XT1 but can be biosynthesized in mammalian cells. WT-XT1 has been reported to use UDP-6AzGlc with approx. 20-fold lower enzymatic efficiency than UDP-Xyl.[44] We opted to develop a mutant with reversed selectivity to accept UDP-6AzGlc over UDP-Xyl. As an analogue of glucose (Glc), 6AzGlc was projected to hijack parts of the UDP-Glc salvage pathway and therefore allow cellular biosynthesis unlike Xyl analogues.[36]

We recently reported the crystal structure of XT1, revealing a two-lobe architecture containing a catalytic glycosyltransferase domain.[17] Since XT1 contains an unusually constricted UDP-Xyl binding site that prevents the use of larger UDP-sugars such as UDP-6AzGlc, we deemed it possible to generate additional space (a "hole") in the active site by mutation. XT1 harbours several bulky "gatekeeper" amino acids in close proximity to C-5 of UDP-Xyl, namely Trp392, Leu525 and Leu526 (Fig. 1b). We designed, expressed and purified from Expi293 cells a total of 16 XT1 single and double mutant variants in which these residues were replaced with smaller amino acids (Fig. 1c, Supporting Fig. 1). *In vitro* glycosylation of a well-known bikunin substrate peptide in an HPLC assay served to assess glycosylation from the sugar donors UDP-Xyl, UDP-6AzGlc and, as a substrate of intermediate size of the "bump", UDP-Glc.[17]

Fig. 2: BH engineering preserves protein substrate specificity of XT1. **a**, *in vitro* glycosylation of a

peptide substrate panel based on the bikunin peptide indicated at the top of the panel as assessed by

a luminescence assay, with every position substituted for each of the 20 amino acids. Intensity of grey

scale indicates % turnover. The WT bikunin peptide was present 12 times in the panel (copied in the

top row) and all other peptide reactions were normalised on the average of these 12 data points. Data

are from one out of three independent experiments. **b**, *in vitro* glycosylation of a membrane protein

fraction of XT2-KO CHO[KO Xyl2t] cells as assessed by streptavidin blot. Reactions contained 250 µM

UDP-6AzGlc and 100, 200 or 300 µM UDP-Xyl, respectively, and were reacted with biotin-alkyne

before blotting. Data are from one out of two independent experiments. **c**, glycosylation by BH-

XT1/UDP-6AzGlc can be prevented by pre-incubation with WT-XT1/UDP-Xyl. Reactions were

processed as in **b**. Data are from one experiment. **d**, *in vitro* glycosylation of a GAG-free preparation

of human decorin purified from pgsA-745 CHO cells as assessed by streptavidin blot. Reactions were

run with 250 µM UDP-sugars and processed as in **b** and **c**. Data are from one out of two independent

experiments. **e**, analysis of the glycosylation site on decorin introduced by BH-XT1 by MS and HCD

fragmentation. Decorin was *in vitro* glycosylated as in **c**, subjected to CuAAC with ITag-alkyne,[45]

digested and subjected to MS-glycoproteomics. Fragments are annotated on the tryptic peptide from

mature decorin. "Tag" denotes the signature ion 322.1508 *m/z* from the depicted chemically modified

sugar. Data are from one experiment.

WT-XT1 displayed exclusive activity for UDP-Xyl in our hands, with no activity towards UDP-Glc or UDP-6AzGlc.[44] Most engineered XT1 variants displayed either no activity at all or were still selective for UDP-Xyl, with some displaying activity toward UDP-Glc. Strikingly, the mutant Leu526Gly preferred UDP-6AzGlc as a substrate, with 7-8-fold higher turnover than using UDP-Xyl or UDP-Glc in an endpoint assay. Compared to the Leu526Gly mutant (henceforth termed "BH-XT1"), the construct Leu526Ala displayed no such selectivity, with equal activity on all three UDP-sugars (Fig. 1c). We determined optimal enzyme concentrations and kinetic constants for the native and BH enzyme-substrate pairs (Fig. 1d, Supporting Figs 2, 3). We found that the K_M of the BH pair (36.9 µM) was similar to the WT pair (38.6 µM), while the k_{cat} was 2.8-fold reduced in the BH pair. In contrast, BH-XT1 uses UDP-Xyl with an approx. 10-fold lower catalytic efficiency than UDP-6AzGlc, suggesting that the native substrate UDP-Xyl should not be able to outcompete UDP-6AzGlc in cellular applications. Taken together, we established a sensitive structure-activity relationship in the development of a XT1 bump-and-hole variant. BH-XT1 fulfilled the crucial pre-requisite of preferring a chemically modified substrate that is not used by the WT enzyme, with comparable kinetics to the WT enzyme-substrate pair.

Bump-and-hole engineering retains the peptide specificity of WT-XT1

To assess whether BH engineering retains substrate preference of WT-XT1 towards the proteoglycan backbone, we first tested the BH enzyme-substrate pair (BH-XT1, UDP-6AzGlc) with a panel of 240 substrate peptides in an *in vitro* glycosylation assay. The panel contained derivatives of the well-characterised bikunin XT1 substrate peptide in which each amino acid was substituted with each of the other 19 proteinogenic amino acids. We had previously used the same peptide panel to extract amino acid preferences of the native enzyme-substrate pair (WT-XT1, UDP-Xyl) in a luminescence-based assay.[17] Employing the same assay, the peptide substrate preferences were found to be remarkably conserved between WT- and BH-XT1 (Fig. 2a, for WT peptide preference see Briggs and Hohenester[17]). For instance, introducing basic Lys or Arg residues anywhere in the substrate peptide lowered enzyme activity, while acidic Glu and Asp tended to increase activity. An exception was the substitution of Glu at -4 position with Asp that led to a decrease in turnover for both WT- and BH-XT1.[17] As in WT-XT1, substitutions of glycine residues at positions -1 and +1 of the central Ser were not well-tolerated by BH-XT1. An exception was substitution of Gly at +1 position to hydrophobic amino acids Leu, Met or Phe, which led to residual activity in BH- but not WT-XT1. Since this Gly is in contact with Leu526 in WT-XT1, we reasoned that the Leu526Gly "hole" left space for substitutions to larger hydrophobic amino acids in substrate peptides. Since these were the only reproducible differences

between WT and BH-XT1 (other replicates in Supporting Fig. 4) and low in number out of a

240-member peptide library, we concluded that BH engineering exhibits conservation of

peptide substrate preference *in vitro*.

BH-XT1 glycosylates proteoglycans at GAG attachment sites *in vitro*

We next assessed whether BH-XT1 retains the activity of WT-XT1 to initiate GAG

attachment sites on proteoglycan backbones. We prepared membrane fractions from

Chinese Hamster Ovary (CHO) cells with or without KO for endogenous xylosyltransferase

genes *Xylt1* and *Xylt2*.[10] Membrane fractions were incubated with recombinant WT- or BH-

XT1 as well as synthetic UDP-Xyl and UDP-6AzGlc, followed by reaction with alkyne-biotin

under copper-catalysed azide-alkyne cycloaddtion (CuAAC) "click" conditions. Analysis by

streptavidin blot suggested labelling of lysate proteins with 6AzGlc only when BH-XT1, but

not WT-XT1 was present (Supporting Fig. 5). Since *Xylt2* is the major xylosyltransferase

gene expressed in CHO cells,[10] we used $CHO^{KO\ Xylt2}$ cells for further *in vitro* glycosylation

experiments. We established that labelling by BH-XT1 with UDP-6AzGlc could not be

outcompeted with increasing concentrations of UDP-Xyl, suggesting that BH-XT1 specifically

and potently recognises UDP-6AzGlc as a substrate (Fig. 2b). Pre-incubation of the

membrane protein fraction with WT-XT1 and UDP-Xyl abrogated incorporation of 6AzGlc by

BH-XT1, suggesting that the same glycosylation sites are introduced by both enzymes (Fig.

2c).

We next confirmed *in vitro* that BH-XT1 emulates the activity of WT-XT1 to glycosylate

proteoglycans. Human decorin has a single site of GAG attachment. Recombinant

expression in the CHO cell mutant pgsA-745 that lacks endogenous XT activity results in a

GAG-free decorin preparation.[7,8] We incubated this GAG-free decorin with either WT- or BH-

XT1 in the presence of UDP-Xyl and/or UDP-6AzGlc, followed by CuAAC with alkyne-biotin

and streptavidin blot. (Fig. 2d). While WT-XT1 activity did not lead to discernible streptavidin

signal on decorin, BH-XT1 in the presence of UDP-6-AzGlc led to intense streptavidin signal

that could be abrogated by pre-incubation of decorin with WT-XT1/UDP-Xyl. These data

indicate that the single GAG attachment site was blocked with a xylose residue by WT-XT1,

preventing BH-XT1 activity. We observed the same behaviour in GAG-free preparation of

human glypican 1 (Supporting Fig. 6), suggesting that BH-XT1 recapitulates the activity of

WT-XT1 across a range of proteoglycans.

We confirmed the glycosylation site modified by BH-XT1 on *in vitro* glycosylated

recombinant decorin by tandem MS. Two fragmentation methods are routinely employed for

O-glycopeptides. High energy collision-induced dissociation (HCD) primarily fragments the

glycosidic bond to detect glycan oxonium ions while electron transfer dissociation (ETD)

fragments the peptide backbone to allow glycan site annotation. The clickable azide tag was

essential to improve sugar identification in mass spectra, allowing incorporation of functional

groups that are beneficial to analysis. Specifically, we employed a clickable imidazolium tag

(ITag) that carries a permanent positive charge and increased the charge state of

glycopeptides, allowing direct glycosylation site annotation.[45] We first employed a standard

workflow in which HCD fragmentation led to an ITag-containing, 6AzGlc-derived signature

ion that was used to trigger ETD on the same glycopeptide.[35,41,43,45] This tandem strategy is

used because O-linked glycan modifications are usually too labile to be detected within

peptide fragments during HCD, hampering glycosylation site localisation. Surprisingly, ITag-

modified 6AzGlc was detected on peptide fragments in HCD spectra on a tryptic

glycopeptide derived from decorin, without the need for additional ETD (Fig. 2e).

Fig. 3: Biosynthesis of UDP-6AzGlc. **a**, biosynthetic pathway of UDP-Glc via hexokinase (HK),

phosphoglucomutase (PGM) and UDP-Glc pyrophosphorylase (UGP). Biosynthesis of UDP-6AzGlc

from caged phosphate **1** bypasses the HK and PGM steps. Insert: crystal structure of UGP with UDP-

Glc indicating that the 6-OH group protrudes into an open cavity (PDB 4R7P). **b**, synthesis of caged

sugar-1-phosphates **1** and **2** from 6AzGlc **3**. **c**, biosynthesis of UDP-6AzGlc in K562 cells as

assessed by ion exchange HPLC of lysates fed with compounds **1** or **2**. Data are from one out of two

independent experiment. Reagents and conditions: a) Ac_2O, pyridine, DMAP, r.t., 90%; b) AcOH,

ethylene diamine, r.t. 45-73%; c) 2 M LDA in THF, **5**, -78°C to -70°C, (**1**) 60% (**2**) 9.8%. DMAP = 4-

Dimethylaminopyridine; LDA = lithium diisopropylamide. RU = relative units.

Decorin is proteolytically processed during secretion to remove a propeptide and shorten the N-terminus.[46] HCD fragmentation allowed for direct identification of Ser34 as the attachment site of 6AzGlc by BH-XT1 on this mature form, consistent with Ser34 being the site of cellular GAG attachment (Fig. 2e).[47] Taken together, these results suggest that the XT1 bump-and-hole enzyme-substrate pair glycosylates native GAG attachment sites in proteoglycans *in vitro*. We also surmise that the 6AzGlc modification is stable towards HCD fragmentation, allowing for facile annotation of GAG attachment sites by MS.

Biosynthesis of UDP-6AzGlc in mammalian cells via a caged sugar-1-phosphate

Application of a GT BH system in living cells requires biosynthesis of the nucleotide-sugar. In general, caged, membrane-permeable monosaccharide precursors are employed with ester modifications that are deprotected in the cytosol. Free monosaccharides can then be converted to UDP-sugars before transport to the Golgi.[42,43,48] Although human cells are devoid of a salvage pathway for UDP-Xyl, our strategic use of UDP-6AzGlc provided an opportunity for cellular application by hijacking the biosynthetic pathway for UDP-Glc instead. Glc is activated in mammalian cells first by phosphorylation to Glc-6-phosphate and, subsequently, isomerization by phosphoglucomutase PGM to Glc-1-phosphate (Fig. 3a). Conversion to UDP-Glc then features the enzymes UDP-Glc pyrophosphorylase 1 or 2, UGP1/2. Since phosphorylation at the 6-position was prevented by the presence of an azido group, we sought to bypass the kinase and PGM steps and provide a sugar-1-phosphate as a direct substrate for UGP1/2. We were encouraged by analysis of the UGP1/UDP-Glc co-crystal structure in which the UDP-Glc 6-hydroxyl group is solvent-exposed, suggesting that an azido group at that position should be tolerated by the enzyme (Fig. 3a).[49] While we and others have made sugar-1-phosphates caged as labile bis-*S*-acetylthioethyl (SATE) phosphotriesters, synthesis of SATE-caged 6AzGlc-1-phosphate failed in our hands.[43,50,51] Instead, we took inspiration from the increasingly popular protide technology that has gained attention to cage phosphates in antiviral nucleotides,[52,53] and has been recently used to cage sugar-phosphates.[54–56] We synthesized phosphoramidate diester **1** as a caged sugar-1-phosphate to be deprotected by hydrolases in the cytosol of living mammalian cells (Fig. 3a).[52] The synthesis proceeded from lactol mixture **3** via the intermediate triacetate **4**. Treatment of **4** with **5** under basic conditions yielded both α-phosphate **1** (60% yield) and β-phosphate **2** (9.8% yield) (Fig. 3b).[57] Since UGP1/2 is naturally restricted to α-configured Glc-1-phosphate, preferential formation of the α-anomer was gratifying. In turn, β-phosphate **2** served as a negative control in feeding experiments. Feeding K-562 cells the α-phosphate **1** led to notable and reproducible biosynthesis of UDP-6AzGlc (Fig. 3c). In turn, β-configured phosphate **2** led to negligible UDP-6AzGlc levels, possibly arising from very small (4%)

amounts of α-configured phosphate **1** as a contaminant of **2** (Supporting Information). These

data suggest that α-phosphate **1** is a suitable precursor to deliver UDP-6AzGlc to

mammalian cells by entering the UDP-Glc biosynthetic pathway.

Fig. 4: BH-engineered xylosyltransferases label glycoproteins in mammalian cells. **a**, overview of

cellular BH engineering. UDP-6AzGlc is biosynthesized after delivery through compound **1**, and used

by BH-engineered XTs. Cell surface CuAAC detects proteoglycans. **b**, chemical tagging of

proteoglycans by BH-XT1 on pgsA-745 CHO cells as assessed by in-gel fluorescence. Cells stably

expressing WT- or BH-XT1 or non-transfected were fed with compounds in the indicated

concentrations in µM before on-cell CuAAC and in-gel fluorescence. Western Blots are from a

separate gel using the same samples. Data are from one out of two independent replicates. **c**,

structural alignment between the crystal structure of human XT1 (PDB 6EJ7, green) and the

AlphaFold structure of human XT2 (accession no. Q9H1B5, purple) with aligned gatekeeper residues

in the insert. **d**, chemical tagging of proteoglycans by BH-XT2 on pgsA-745 CHO cells as assessed by

in-gel fluorescence. Cells stably transfected with WT- or BH-XT2 were fed and treated as in **b**. Data

are from one out of two independent experiments.

Development of a cellular XT1 bump-and-hole system

With a strategy for UDP-6AzGlc delivery in hand, we established a BH system to chemically tag XT1 substrate proteins in mammalian cells (Fig. 4a). The pgsA-745 CHO cell line was employed because of the lack of competition with internal xylosyltransferases. Cells were stably transfected with plasmids encoding WT- or BH-XT1 using a transposase-based genome insertion method,[43,58] followed by feeding the 6AzGlc-1-phosphate precursor **1**. After overnight incubation, CuAAC was performed to attach clickable alkyne-CF680 to 6AzGlc on the cell surface while keeping cells alive.[35,43,48] Surplus click reagents were washed away, cells lysed and fluorophore incorporation assessed by SDS-PAGE and in-gel fluorescence (Fig. 4b). Minimal background fluorescence was observed in cells fed with DMSO or only expressing WT-XT1, even when fed with increasing concentrations of 6AzGlc-1-phosphate precursor **1**. In the presence of BH-XT1, clear fluorescent proteins were observed at 38 kDa, 90 kA and 260 kDa. With increasing feeding concentration of **1**, a dose-dependent increase of fluorescence was observed, along with labelled protein bands of lower intensity, especially between 50 and 90 kDa. In accordance with biosynthetic experiments, the β-configured 6AzGlc-1-phosphate **2** yielded a weak and diffuse labelling signal when fed to cells, indicating that UDP-6AzGlc biosynthesis is a direct prerequisite for cellular chemical tagging of glycoproteins by BH-XT1. The fluorescent bands at 90 kDa and 260 kDa in the BH-XT1/compound **1** lanes were observed weakly when compound **2** was fed, which we attribute to small residual levels of compound **1** in the preparation (*see* Fig. 3c). Ac$_4$ManNAz as a precursor to azide-tagged sialic acid yielded a strong and uniform fluorescent signal across all cell lines tested.[59] Our data indicate that BH-XT1 specifically tags cell surface proteins with bioorthogonal 6AzGlc in the secretory pathway of mammalian cells.

The mammalian genome encodes two xylosyltransferase isoenzymes that have been differentially implicated in disease and proteoglycan function.[28,30,32,60] To allow for substrate profiling of the second isoenzyme XT2, we extended the BH strategy by employing structural similarities between the two isoenzymes. A structural overlay between the XT1 crystal structure and the XT2 structure predicted by AlphaFold highlighted conservation of amino acids interacting with UDP-Xyl (Fig. 4d). We identified the gatekeeper residues Leu431, Leu432 and Trp298 in XT2 that occupied the same role as the respective residues in XT1, with Leu432 in XT2 overlaying with Leu526 in XT1. The BH-XT2 mutant L432G was thus stably expressed in pgsA-745 CHO cells. Feeding with 6AzGlc-1-phosphate precursor **1** and cell-surface CuAAC reaction with CF680-alkyne led to a similar band pattern by in-gel fluorescence, with glycoprotein bands at 90 kDa and 260 kDa identified in BH-XT2-expressing cells in a dose-dependent manner (Fig. 4d). We noted that the intensity of fluorescent bands specifically tagged by BH-XT2 was lower than by BH-XT1, indicating a

lower activity of the second isoenzyme. WT-XT2 expression did not lead to the same

fluorescent band pattern. Neither feeding DMSO nor compound **2** led to discernible signal

over background, and Ac$_4$ManNAz was included as a positive labelling control. Our data

suggest that the BH approach is applicable to both xylosyltransferase isoenzymes in

mammalian cells, allowing for the first time to assess their substrate profiles.

XT bump-and-hole engineering enables profiling of cellular proteoglycans

Xylosyltransferase BH engineering is poised to allow the identification of proteoglycans, a

feat that normally requires elaborate methods of glycopeptide enrichment and

characterisation.[9,21,23] A pre-requisite to glycoprotein enrichment and identification is

biosynthetic simplicity – ideally, 6AzGlc would replace an entire GAG chain without added

complexity from glycan elaboration. To establish an MS-glycoproteomics workflow, it was

thus important to assess whether 6AzGlc, like Xyl, was extended to a functional GAG linker

tetrasaccharide. We recently reported an enzymatic method for extension of xylosylated

glycopeptides by recombinant preparations of the glycosyltransferases B4GALT7, B3GALT6

and B3GAT3 (termed linker enzymes) in the presence of UDP-galactose (UDP-Gal) and

UDP-glucuronic acid (UDP-GlcA). Employing a fluorescently labelled bikunin-derived

peptide, we first confirmed by HPLC that attachment of either Xyl (by WT-XT1) or 6AzGlc (by

BH-XT1) led to a shift of peptide retention time (Fig. 5a). Upon addition of the linker

enzymes, the Xyl moiety was sequentially elongated to the full tetrasaccharide. In contrast, a

6AzGlc-modified peptide did not shift in retention time upon incubation with the GTs and

UDP-sugars. We concluded that 6AzGlc is a chain-terminating modification that is not

extended to functional GAG chains. We interpreted this substantial decrease in glycan

complexity as an advantage for MS and to manipulate the composition of proteoglycans (see

below).

We next determined that a functional XT1 system chemically tags a model proteoglycan in

mammalian cells by MS. FLAG-tagged human decorin was overexpressed as a secreted

proteoglycan in pgsA-745 CHO cells that expressed BH-XT1. Cells were fed with 6AzGlc-1-

phosphate precursor **1**, and decorin immunoprecipitated from conditioned supernatant. The

preparation was subjected to CuAAC with ITag-alkyne, digested and subjected to MS with

HCD fragmentation. We confirmed unambiguously that Ser34 was glycosylated by BH-XT1

inside mammalian cells, confirming the bump-and-hole approach as suitable to identify

proteoglycans including native Xyl attachment sites (Fig. 5b, c).

Our in-gel fluorescence data suggested that the two isoenzymes XT1 and XT2 may exhibit

an overlapping substrate glycoprotein profile. To test this notion, we compared the

glycoproteome tagged by either BH-XT1 or BH-XT2 in cells by MS. Chemically tagged

glycoproteins in conditioned supernatants of stably transfected pgsA-745 CHO cells were

treated with alkyne-biotin, enriched on neutravidin resin and subjected to on-bead proteolytic

digest. Peptide fractions were analysed by MS using data-independent acquisition over three

replicates each, with WT-XT1/2-expressing cells fed with **1** as control conditions. Data were

processed using the software DIA-NN to identify and quantify *Cricetulus griseus* proteins.[61]

Enriched peptide and protein fractions were analyzed with Perseus.[62] Glycosylation by BH-

XT1 led to a striking enrichment of cellular proteoglycans, including decorin, syndecan-4,

glypican-1 and CD44, compared to the supernatant of cells expressing WT-XT1 (Fig. 5d,

Supporting Fig. 7). BH-XT2 led to strong enrichment of the same glycoproteins with

exception of CD44 (Fig. 5e, Supporting Fig. 7). In both cases, we analyzed the identity of

detected peptides from these enriched proteins and found them to lie outside previously

annotated glycosylation site(s) (Supporting Table 1). These findings suggests that cellular

XT1/2 BH engineering allows unambiguous proteoglycan profiling without the need to

annotate the GAG attachment site. When a less stringent comparison was made between

cells expressing BH-XT1/2 fed with compound **1** against the same cell lines fed with DMSO

as a control, the *bona fide* proteoglycans nidogen-1, chondroitin sulfate proteoglycan 4, agrin

and aggrecan were additionally found to be enriched (Supporting Fig. 8). We conclude that

BH engineering enables straightforward detectability of proteoglycans from mammalian cells.

Chemical functionalization allows for functional annotation of the GAG modification

on proteoglycans

The BH approach replaces the GAG chain of a proteoglycan with a bioorthogonal

modification. We reasoned that this approach could be used to introduce suitably modified

GAG polysaccharides, creating neo-glycoproteins that allow for functional dissection of the

proteoglycan components in cell biology. We have previously used unnatural amino acid

introduction via amber stop codon reassignment to produce syndecan-1 (SDC1) chemically

tagged at position 37 (termed $SDC1_{37}$) with a clickable alkyne.[16] Through CuAAC, SDC_{37}

could be furnished with azide-containing, clickable heparin to generate a neo-proteoglycan

termed $SDC1_{37}$-Hep. In the presence of either fully glycosylated, HS-containing SDC1 or

$SDC1_{37}$-Hep, MDA-MB-231 cells exhibit enhanced spreading on vitronectin-coated surfaces

in an integrin $\alpha_v\beta_3$-dependent fashion that can be analyzed by microscopy. Soluble SDC1

preparations can be deposited on cells by virtue of their hexa-His tags through the use of

synthetic Ni^{2+}-displaying, membrane-anchored cholesterol anchor.[16] We used BH-XT1 to

introduce 6AzGlc into a non-glycosylated WT-SDC1 preparation *in vitro*, to generate SDC1-

6AzGlc. Heparin was furnished with an anomeric alkyne tag via hydrazide chemistry, and

introduced into 6AzGlc-SDC1 via CuAAC to furnish SDC1-6AzGlc-Hep. Both SDC1-Hep

conjugates displayed an increase in molecular weight (Supporting Fig. 9). MDA-MB-231

cells were depleted of endogenous SDC1 by siRNA-based knockdown, significantly

impacting cell spreading (Fig. 5f). Deposition of non-glycosylated SDC1$_{37}$ or SDC1-6AzGlc

enhanced cell spreading by approx. 20%, suggesting a role of the SDC backbone alone for

MDA-MB-231 spreading that has been noted before.[16] Heparin-containing SDC1$_{37}$-Hep and

SDC1-6AzGlc-Hep fully rescued cell spreading, indicating that BH engineering and

bioorthogonal chemistry restored the functional properties of a proteoglycan.

Fig. 5: Chemical functionalization by BH-XT1/2 enables detection and manipulation of proteoglycans.

a, 6AzGlc is not elongated by GAG linker enzymes. Fluorescently labelled bikunin-derived peptide

was incubated with WT-XT1/UDP-Xyl or BH-XT1/UDP-6AzGlc and subsequently with the indicated

soluble glycosyltransferases[7] and corresponding UDP-sugars. Elongation was assessed on each step

by HPLC. Data are from one experiment each with different combinations of transferases. **b**, using

BH-XT1/2 to profile proteoglycans by MS. **c**, analysis of the glycosylation site on decorin introduced

by BH-XT1 in living cells by MS and HCD fragmentation. Decorin was co-expressed with BH-XT1 in

cells fed with **1**, then subjected to CuAAC with ITag-alkyne,[45] digested and subjected to tandem MS.

Fragments are annotated on the tryptic peptide from mature decorin. Not all fragments are shown.

"Tag" denotes the signature ion 322.1508 *m/z* from chemically modified sugar. Data are from one

experiment. **d**, differential MS experiment of peptide fractions enriched in BH- vs. WT-XT1-expressing

pgsA-745 CHO cells fed with compound **1** from three independent replicates. The proteoglycans were

assigned and processed from the *Cricetulus griseus* protein database search (Uniprot) followed by

visualization of data on Perseus. **e**, differential MS experiment of peptide fractions enriched in BH- vs.

WT-XT2-expressing pgsA-745 CHO cells fed with compound **1** from three independent replicates. **f**,

analysis of cell spreading of MDA-MB-231 cells with SDC1-knockdown and rescue with clickable

heparin introduced either into alkyne-tagged $SDC1_{37}$ or azide-tagged SDC1-6AzGlc. Spreading relies on the interaction between HS/heparin-containing SDC1 and $\alpha_v\beta_3$ integrin on MDA-MB-231 cells and vitronectin on coated wells. Statistical significance was assessed by one-way ANOVA, and asterisks denote p values: **** $p < 0.0001$; ** $p < 0.01$; * $p < 0.05$. Data are means + SEM from three independent replicates with cells counted in at least six different frames per replicate.

Discussion

The importance of proteoglycans in physiology is undisputed, as the vast majority of signalling events between cells or with the extracellular matrix are modulated by the associated GAG chains. While great efforts are being made to understand the details of GAG polysaccharide sequence on biology,[10,14,22,63,64] we still lack important information on the first step of glycosylation to the protein backbone. The two human xylosyltransferases display tissue selectivity and differences in attached GAG sequences but we do not yet have fundamental detail on their individual biological functions.[28,31–33,60] Furthermore, only a relatively small number of annotated mammalian proteoglycans has been identified, with new annotations requiring considerable effort.[9] A chemical tool to dissect XT1/2 biology must accurately report on XT1/2 activity while being orthogonal to other glycosylation events in the secretory pathway and deliverable to living mammalian cells. The use of a UDP-Glc analogue matched these pre-requisites. Both catalytic efficiency and peptide substrate preference of the XT1 bump-and-hole enzyme-substrate pair were remarkably conserved, which we attribute to the careful structure-based design of the pair. Incorporation of 6AzGlc into cell surface proteins was dependent on the presence of BH-XT1, indicating that the sugar does not enter other major glycosylation pathways. We note that this finding does not exclude incorporation into glycoconjugates that naturally contain D-Glc, as most of these are either glycolipids or intracellular glycoproteins, neither of which would be visible with our assays.

The finding that 6AzGlc is not extended to the common GAG linker tetrasaccharide was expected due to the restrictive active site architecture of the follow-on enzyme B4GALT7.[65] However, we recognize this non-extension as a particular advantage for MS since the sugar 6AzGlc is structurally well-defined and does not require any further simplification by glycosidase treatment prior to analysis. In our MS experiments, annotation was further simplified by the availability of the ITag technology to facilitate MS.[45,66] Furthermore, a chain-terminating, clickable inhibitor of chain extension has the potential to be employed to study GAG biology *in vitro* or *in vivo*,[16,39,67] substantially expanding our toolbox.

Establishing a cellular XT bump-and-hole system required a biosynthetic entry point for UDP-6AzGlc. Previously used per-acetylated 6AzGlc was not a suitable precursor for glycosylation in our hands,[36] but we note that cell lines from different organisms can vary in

their biosynthetic potential.[68] Nevertheless, a protide-based caged sugar-1-phosphate was a reliable precursor for UDP-6AzGlc to fashion a cellular bump-and-hole system. We did not assess whether UDP-6AzGlc was used by other glycosyltransferases, but cell surface labelling experiments suggested that BH-XT activity was necessary to introduce the chemical modification into glycoproteins.

While XT2 appears to be the dominant isoenzyme expressed in humans, dysfunctions in both enzymes lead to severe yet differential disorders in mouse models and in patients.[30–33] After fully characterising an XT1 BH system, we designed a functional BH-XT2 mutant simply based on structural homology. BH-XT2 was directly applied in living cells without characterizing the corresponding soluble recombinant enzyme first, showcasing the reliability of the tactic as well as the importance of structural data for identification of nucleotide-sugar binding.

Proteoglycans can be perceived as a modular assembly between a protein backbone and one or more GAG chains. While biological function can be imparted by either component, it can be challenging to differentiate both. The biosynthetic details of GAG extension by either HS or CS/DS are beginning to be understood,[7] but methods to reliably swap the GAG chain on a given proteoglycan are of fundamental importance for proteoglycan biology. We have recently employed stop codon reassignment to introduce an alkyne-tagged amino acid into a recombinant proteoglycan backbone,[16] as a critical aspect of understanding the role of SDC1 for cell spreading. We applied BH engineering to underpin these findings, allowing to chemoenzymatically attach a chemical modification to the recombinant protein. We note the ability to employ this modular approach to generate "designer proteoglycans" with unnatural GAG chains to generate functional understanding.

Our work is setting the foundation to establish the fine differences between XT1 and XT2 and profile proteoglycans in a range of different model systems.

Acknowledgements

We thank Ganka Bineva-Todd and Andrea Marchesi for help with chemical synthesis and analysis, and Khadija Babiker for help with cloning similar constructs. We thank Chloë Roustan, Svend Kjaer and the Francis Crick Institute Structural Biology Science Technology Platform for help with protein expression and purification. We further thank Tania Auchynnikava and Mark Skehel for help with mass spectrometry, and the Crick Proteomics, Chemical Biology and Cell Services Science Technology Platforms for valuable support. This work was supported by the Francis Crick Institute (to B. S.) which receives its core funding from Cancer Research UK (CC2127, CC2068), the UK Medical Research Council (CC2127, CC2068) and Wellcome Trust (CC2127, CC2068), the BBSRC (BB/T01279X/1 to E. H. and B. S., BB/V008439/1 to B. S.), the EPSRC (EP/T007397/1 to G.J.M) and the MRC

(MR/T019522/1 to G.J.M). This work was supported in part by the NIH under award

R35GM142462 (to M. L. H.). We also thank the EPSRC UK National Mass Spectrometry

Facility (NMSF) at Swansea University. This work was supported by the Novo Nordisk

Foundation grant No. NNF22OC0073736 (to R. L. M.). L. I. W. gratefully acknowledges

funding from the European Research Council (ERC) under the European Union's Horizon

2020 research and innovation programme (Grant agreement No 851448). For the purpose of

Open Access, the author has applied a CC BY public copyright licence to any Author

Accepted Manuscript version arising from this submission.

1. Xu, D. & Esko, J. D. Demystifying Heparan Sulfate–Protein Interactions. *Annu Rev Biochem* **83**, 129–157 (2014).

2. Bishop, J. R., Schuksz, M. & Esko, J. D. Heparan sulphate proteoglycans fine-tune mammalian physiology. *Nature* **446**, 1030–1037 (2007).

3. Pickford, C. E. *et al.* Specific Glycosaminoglycans Modulate Neural Specification of Mouse Embryonic Stem Cells. *Stem Cells* **29**, 629–640 (2011).

4. Kreuger, J. *et al.* Fibroblast growth factors share binding sites in heparan sulphate. *Biochem J* **389**, 145–150 (2005).

5. Mizumoto, S. & Yamada, S. Congenital Disorders of Deficiency in Glycosaminoglycan Biosynthesis. *Front Genet* **12**, 717535 (2021).

6. Zhang, L. & Esko, J. D. Amino acid determinants that drive heparan sulfate assembly in a proteoglycan. *J of Biol Chem* **269**, 19295–19299 (1994).

7. Sammon, D. *et al.* Molecular mechanism of decision-making in glycosaminoglycan biosynthesis. *Nat Commun 2023* **14**, 1–13 (2023).

8. Esko, J. D., Stewart, T. E. & Taylor, W. H. Animal cell mutants defective in glycosaminoglycan biosynthesis. *Proc Natl Acad Sci U S A* **82**, 3197–3201 (1985).

9. Zhang, P. *et al.* Heparan Sulfate Organizes Neuronal Synapses through Neurexin Partnerships. *Cell* **174**, 1450-1464.e23 (2018).

10. Chen, Y. H. *et al.* The GAGOme: a cell-based library of displayed glycosaminoglycans. *Nat Methods* **15**, 881–888 (2018).

11. Qiu, H. *et al.* A mutant-cell library for systematic analysis of heparan sulfate structure–function relationships. *Nat Methods 2018* **15**, 889–899 (2018).

12. Ji, S. K. *et al.* PTPσ functions as a presynaptic receptor for the glypican-4/LRRTM4 complex and is essential for excitatory synaptic transmission. *Proc Natl Acad Sci U S A* **112**, 1874–1879 (2015).

13. Kamimura, K. *et al.* Perlecan regulates bidirectional Wnt signaling at the Drosophila neuromuscular junction. *J Cell Biol* **200**, 219–233 (2013).

14. Merry, C. L. R., Lindahl, U., Couchman, J. & Esko, J. D. Proteoglycans and Sulfated Glycosaminoglycans. *Essentials of Glycobiology* (2022)

15. Noborn, F. & Sterky, F. H. Role of neurexin heparan sulfate in the molecular assembly of synapses – expanding the neurexin code? *FEBS J* **290**, 252–265 (2023).

16. O'Leary, T. R. *et al.* Chemical editing of proteoglycan architecture. *Nat Chem Biol* **18**, 634–642 (2022).

17. Briggs, D. C. & Hohenester, E. Structural Basis for the Initiation of Glycosaminoglycan Biosynthesis by Human Xylosyltransferase 1. *Structure* **26**, 801-809.e3 (2018).

18. Zhang, L., David, G. & Esko, J. D. Repetitive Ser-Gly Sequences Enhance Heparan Sulfate Assembly in Proteoglycans. *J Biol Chem* **270**, 27127–27135 (1995).

19. Ramarajan, M. G. *et al.* Mass spectrometric analysis of chondroitin sulfate-linked peptides. *J Protein Proteomics 2022* **13**, 187–203 (2022).

20. Ly, M., Laremore, T. N. & Linhardt, R. J. Proteoglycomics: Recent progress and future challenges. *OMICS* **14**, 389–399 (2010).

21. Noborn, F., Nilsson, J. & Larson, G. Site-specific glycosylation of proteoglycans: A revisited frontier in proteoglycan research. *Matrix Biology* vol. 111 289–306 (2022).

22. Persson, A., Nikpour, M., Vorontsov, E., Nilsson, J. & Larson, G. Domain Mapping of Chondroitin/Dermatan Sulfate Glycosaminoglycans Enables Structural Characterization of Proteoglycans. *Mol Cell Proteom* **20**, 100074 (2021).

23. Noborn, F. *et al.* A Glycoproteomic Approach to Identify Novel Proteoglycans. *Meth Mol Biol* **2303**, 71–85 (2022).

24. Noborn, F. *et al.* Site-specific identification of heparan and chondroitin sulfate glycosaminoglycans in hybrid proteoglycans. *Sci Rep* **6**, 1–11 (2016).

25. Wen, J. *et al.* Xylose phosphorylation functions as a molecular switch to regulate proteoglycan biosynthesis. *Proc Natl Acad Sci U S A* **111**, 15723–15728 (2014).

26. Bui, C. *et al.* XYLT1 mutations in desbuquois dysplasia type 2. *Am J Hum Genet* **94**, 405–414 (2014).

27. Munns, C. F. *et al.* Homozygosity for frameshift mutations in XYLT2 result in a spondylo-ocular syndrome with bone fragility, cataracts, and hearing defects. *Am J Hum Genet* **96**, 971–978 (2015).

28. Wilson, I. B. H. The never-ending story of peptide O-xylosyltransferase. *Cell Mol Life Sci* **61**, 794–809 (2004).

29. Götting, C., Kuhn, J., Zahn, R., Brinkmann, T. & Kleesiek, K. Molecular cloning and expression of human UDP-D-xylose: Proteoglycan core protein β-D-xylosyltransferase and its first isoform XT-II. *J Mol Biol* **304**, 517–528 (2000).

30. Ferencz, B. *et al.* Xylosyltransferase 2 deficiency and organ homeostasis. *Glycoconj J* **37**, 755 (2020).

31. Taieb, M., Ghannoum, D., Barré, L. & Ouzzine, M. Xylosyltransferase I mediates the synthesis of proteoglycans with long glycosaminoglycan chains and controls chondrocyte hypertrophy and collagen fibers organization of in the growth plate. *Cell Death Dis 2023 14:6* **14**, 1–13 (2023).

32. Roch, C., Kuhn, J., Kleesiek, K. & Götting, C. Differences in gene expression of human xylosyltransferases and determination of acceptor specificities for various proteoglycans. *Biochem Biophys Res Commun* **391**, 685–691 (2010).

33. Kuhn, J. *et al.* Xylosyltransferase II is the predominant isoenzyme which is responsible for the steady-state level of xylosyltransferase activity in human serum. *Biochem Biophys Res Commun* **459**, 469–474 (2015).

34. Choi, J. *et al.* Engineering Orthogonal Polypeptide GalNAc-Transferase and UDP-Sugar Pairs. *J Am Chem Soc* **141**, 13442–13453 (2019).

35. Debets, M. F. *et al.* Metabolic precision labeling enables selective probing of O-linked N-acetylgalactosamine glycosylation. *Proc Natl Acad Sci U S A* **117**, 25293–25301 (2020).

36. Darabedian, N., Gao, J., Chuh, K. N., Woo, C. M. & Pratt, M. R. The Metabolic Chemical Reporter 6-Azido-6-deoxy-glucose Further Reveals the Substrate Promiscuity of O-GlcNAc Transferase and Catalyzes the Discovery of Intracellular Protein Modification by O-Glucose. *J Am Chem Soc* **140**, 7092–7100 (2018).

37. Darabedian, N. *et al.* O-Acetylated Chemical Reporters of Glycosylation Can Display Metabolism-Dependent Background Labeling of Proteins but Are Generally Reliable Tools for the Identification of Glycoproteins. *Front Chem* **8**, 529502 (2020).

38. Daughtry, J. L., Cao, W., Ye, J. & Baskin, J. M. Clickable Galactose Analogues for Imaging Glycans in Developing Zebrafish. *ACS Chem Biol* **15**, 318–324 (2020).

39. Beahm, B. J. *et al.* A visualizable chain-terminating inhibitor of glycosaminoglycan biosynthesis in developing zebrafish. *Angew Chem Int Ed Engl* **53**, 3347–3352 (2014).

40. Bakker, H. *et al.* Functional UDP-xylose transport across the endoplasmic reticulum/golgi membrane in a Chinese hamster ovary cell mutant defective in UDP-xylose synthase. *J Biol Chem* **284**, 2576–2583 (2009).

41. Cioce, A. *et al.* Cell-specific bioorthogonal tagging of glycoproteins. *Nat Commun* **13**, 1–18 (2022).

42. Cioce, A., Malaker, S. A. & Schumann, B. Generating orthogonal glycosyltransferase and nucleotide sugar pairs as next-generation glycobiology tools. *Curr Opin Chem Biol* **60**, 66–78 (2021).

43. Schumann, B. *et al.* Bump-and-Hole Engineering Identifies Specific Substrates of Glycosyltransferases in Living Cells. *Mol Cell* **78**, 824-834.e15 (2020).

44. Gao, J. *et al.* Exploration of Human Xylosyltransferase for Chemoenzymatic Synthesis of Proteoglycan Linkage Region. *Org Biomol Chem* **19**, 3374–3378 (2021)

45. Calle, B. *et al.* Benefits of Chemical Sugar Modifications Introduced by Click Chemistry for Glycoproteomic Analyses. *J Am Soc Mass Spectrom* **32**, 2366–2375 (2021).

46. von Marschall, Z. & Fisher, L. W. Decorin is processed by three isoforms of bone morphogenetic protein-1 (BMP1). *Biochem Biophys Res Commun* **391**, 1374–1378 (2010).

47. Seo, N. S., Hocking, A. M., Höök, M. & McQuillan, D. J. Decorin Core Protein Secretion Is Regulated by N-Linked Oligosaccharide and Glycosaminoglycan Additions. *J Biol Chem* **280**, 42774–42784 (2005).

48. Cioce, A. *et al.* Optimization of Metabolic Oligosaccharide Engineering with Ac4GalNAlk and Ac4GlcNAlk by an Engineered Pyrophosphorylase. *ACS Chem Biol* **16**, 1961–1967 (2021).

49. Führing, J. I. *et al.* A Quaternary Mechanism Enables the Complex Biological Functions of Octameric Human UDP-glucose Pyrophosphorylase, a Key Enzyme in Cell Metabolism. *Sci Rep* **5**, 1–11 (2015).

50. Yu, S. H. *et al.* Metabolic labeling enables selective photocrosslinking of O-GlcNAc-modified proteins to their binding partners. *Proc Natl Acad Sci U S A* **109**, 4834–4839 (2012).

51. Murphy, L. D. *et al.* Synthesis of biolabile thioalkyl-protected phosphates from an easily accessible phosphotriester precursor. *Chem Sci* **14**, 5062–5068 (2023).

52. Mehellou, Y., Rattan, H. S. & Balzarini, J. The ProTide Prodrug Technology: From the Concept to the Clinic. *J Med Chem* **61**, 2211–2226 (2018).

53. Guinan, M., Huang, N., Smith, M. & Miller, G. J. Design, chemical synthesis and antiviral evaluation of 2′-deoxy-2′-fluoro-2′-C-methyl-4′-thionucleosides. *Bioorg Med Chem Lett* **61**, 128605 (2022).

54. Morozzi, C. *et al.* Targeting GNE Myopathy: A Dual Prodrug Approach for the Delivery of N-Acetylmannosamine 6-Phosphate. *J Med Chem* **62**, 8178–8193 (2019).

55. McGuigan, C. *et al.* Phosphate prodrugs derived from N-acetylglucosamine have enhanced chondroprotective activity in explant cultures and represent a new lead in antiosteoarthritis drug discovery. *J Med Chem* **51**, 5807–5812 (2008).

56. Kufleitner, M. *et al.* Next-Generation Metabolic Glycosylation Reporters Enable Detection of Protein O−GlcNAcylation in Living Cells without S-Glyco Modification. *Angew Chem Int Ed* **63**, e202320247 (2024).

57. Ross, B. S., Ganapati Reddy, P., Zhang, H. R., Rachakonda, S. & Sofia, M. J. Synthesis of diastereomerically pure nucleotide phosphoramidates. *J Org Chem* **76**, 8311–8319 (2011).

58. Kowarz, E., Löscher, D. & Marschalek, R. Optimized Sleeping Beauty transposons rapidly generate stable transgenic cell lines. *Biotechnol J* **10**, 647–653 (2015).

59. Saxon, E. & Bertozzi, C. R. Cell surface engineering by a modified Staudinger reaction. *Science* **287**, 2007–2010 (2000).

60. Götting, C., Kuhn, J. & Kleesiek, K. Human xylosyltransferases in health and disease. *Cell Mol Life Sci* **64**, 1498–1517 (2007).

61. Demichev, V. *et al.* DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. *Nat Methods* **17**, 41–44 (2020).

62. Tyanova, S. *et al.* The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat Methods* **13** 731–740 (2016).

63. Barnett, M. W., Fisher, C. E., Perona-Wright, G. & Davies, J. A. Signalling by glial cell line-derived neurotrophic factor (GDNF) requires heparan sulphate glycosaminoglycan. *J Cell Sci* **115** 4495–4503 (2002).

64. Sterner, E., Meli, L., Kwon, S. J., Dordick, J. S. & Linhardt, R. J. FGF-FGFR signaling mediated through glycosaminoglycans in microtiter plate and cell-based microarray platforms. *Biochemistry* **52**, 9009–9019 (2013).

65. Tsutsui, Y., Ramakrishnan, B. & Qasba, P. K. Crystal Structures of β-1,4-Galactosyltransferase 7 Enzyme Reveal Conformational Changes and Substrate Binding. *J Biol Chem* **288**, 31963–31970 (2013).

66. Gonzalez-Rodriguez, E. *et al.* O-Linked Sialoglycans Modulate the Proteolysis of SARS-CoV-2 Spike and Likely Contribute to the Mutational Trajectory in Variants of Concern. *ACS Cent Sci* **9**, 393–404 (2023).

67. Maciej-Hulme, M. L. *et al.* Selective inhibition of heparan sulphate and not chondroitin sulphate biosynthesis by a small, soluble competitive inhibitor. *Int J Mol Sci* **22**, 6988 (2021).

68. Batt, A. R., Zaro, B. W., Navarro, M. X. & Pratt, M. R. Metabolic Chemical Reporters of Glycans Exhibit Cell-Type-Selective Metabolism and Glycoprotein Labeling. *ChemBioChem* **18**, 1177–1182 (2017).